Of Shared Homes and Pathways: Free-Ranging Dog Movement and

Habitat Use in a Human-Wildlife Landscape in India

Sanjana Vadakke Kuruppath*, #12, Valmiki Nagar, Thiruvanmiyur, Chennai 600041, India

Nilanjan Chatterjee*, Senckenberg Biodiversity and Climate Research Institute, Frankfurt am

Main, 60325, Germany

Ramesh Krishnamurthy, Wildlife Institute of India, Dehradun, Uttarakhand 248001, India

*Corresponding authors:

vksanjana@gmail.com

nilanjan.chatterjee@senckenberg.de

Abstract

Background: Free-ranging dogs are widely considered to potentially have negative impacts on

wildlife in shared landscapes. Understanding their use of these landscapes is therefore essential

to develop effective management strategies in wildlife-adjacent regions. In this study, we

investigated the movement patterns of free-ranging dogs in the protected area matrix of

Mudumalai Tiger Reserve (MTR) in Southern India.

Methods: Fifteen dogs were collared from two adjacent villages with different levels of human

habitation. We hypothesized that increased settlement cover would provide dogs with a greater

human shield effect, allowing them more freedom of movement. We estimated dog activity

ranges using autocorrelated kernel density estimates (AKDE) and characterized broad-scale

movement through metrics including intensity of space use and mean distance from home. We

then compared these metrics with respect to different variables to clarify underlying drivers of

movement patterns. Lastly, we analysed fine-scale habitat selection using integrated step

selection analysis (iSSA).

Results: We found that free-ranging dogs in MTR have small activity ranges, with the mean

AKDE activity range being 9.88 (median = 6.09 ha, SD = 7.69 ha, range 3.28-26.16 ha), and

primarily utilise the area within 500m of their homes. No movement metric varied significantly

with respect to any variable apart from intensity of use, which varied significantly between

villages, indicating the dogs in the village with less human settlement restricted their movement

more. iSSA revealed that dogs did not select for any specific habitat type other than human

settlements and mostly moved within human settled or human-modified spaces, moving

significantly faster in forest land than in any other habitat.

Conclusions: These findings support the human shield hypothesis and highlight the role of

human land use in shaping the movement of domestic dogs, in turn mediating potential effects of

dogs on wildlife. Our study shows that dogs show plasticity in space use even at fine scales of

5km or less, which has important implications for management strategies that aim to curb dog

movement.

Introduction

As the most abundant carnivore on earth with a long history of domestication [1], dogs are as widely distributed as humans, with the most abundant category possibly being rural or village dogs [2]. These dogs are highly likely to be free-ranging, with previous studies on rural dog demography across the world indicating that a major proportion of village dogs are free to roam without any restraint, whether alone or in the company of humans [3–6]. In areas close to natural ecosystems, free-ranging dogs can have a range of negative impacts on wildlife [7,8], potentially ranging from mild to severe depending on the local socio-ecological context [9]. Broadly, the most common negative impact of dogs on threatened wildlife is predation, estimated at 78.9%, followed by disturbance (such as barking, chasing or harassment) and disease transmission at 7.0% and 4.5% respectively, and lastly competition with native carnivores (1.5%) and hybridisation with wild canids (1.0%) [10].

In order to clarify the true extent of these effects, understanding how dogs utilise wildlife-adjacent landscapes is perhaps the most necessary avenue of research, as patterns of movement and their driving factors underpin all subsequent impacts, from disease transmission to predation. A metric as fundamental as the home range (or activity range, for short-term studies) can vary regionally, seasonally, and individually, as previous studies from Australia [11–16] Chile [17], Kenya [18], Cambodia [19] and Mexico [20,21] have shown. It follows that roaming behaviour and habitat use, aspects of interest from a wildlife conservation perspective, may be as varied and may depend on a wide range of natural as well as anthropogenic factors. For example, dogs that disrupted sea turtle nests to scavenge for eggs were fed significantly less food by their owners compared to dogs that did not scavenge [21].

The need for research on dog-movement is particularly pressing in India, where the majority of

the dog population is free ranging [22] and may often come into contact with increasingly

fragmented natural ecosystems [23] that harbour a rich diversity of wildlife [24]. Previous

studies have investigated other impacts of dogs on wildlife in India [25–27], with one study

recording reports of negative interactions between dogs and 80 wild species [28]; however, to

our knowledge, only two studies have estimated free-ranging dog movement ecology in rural or

wildlife adjacent areas, both in the state of Maharashtra [29,30]. Of these, only Vanak and

Gompper (2010), utilised telemetry to estimate home range and habitat use. As free-ranging dogs

may exhibit spatial behaviour that is highly context-dependent, fine-scale telemetry offers a

much-needed tool to document their movement ecology in matrix landscapes with joint human-

wildlife presence.

In this study, we address this research gap utilising a fine-scale telemetry study on village dogs

from a multiple-use landscape of a protected area in Southern India. As rural free-ranging dogs

are usually closely associated with human habitation [1], we investigate how dog movement is

affected by the relative presence of this habitation. We collared dogs in two different villages

inside the forest with varying levels of human population, built up area, and village-forest

boundary length. We hypothesize that a higher proportion of built up area, in conjunction with a

smaller village-forest boundary, will produce a human shield effect [31] that allows dogs more

freedom of movement in a landscape dominated by forests and wildlife. To test this hypothesis,

we designed a study with two objectives: 1. Estimate and compare activity ranges and movement

metrics between villages and 2. Evaluate how dog use of habitat types differs between villages.

Materials and Methods

Study area

The study was conducted in the villages of Masinagudi and Moyar, in Mudumalai Tiger Reserve (henceforth, MTR), Tamil Nadu, India. MTR is a protected area contiguous with multiple other wildlife sanctuaries, forming an integral part of the Nilgiri Biosphere Reserve in the highly biodiverse Western Ghats. Several villages are embedded within MTR, forming a matrix landscape where human settlements and major roads are interspersed within the forest. The study sites, Masinagudi and Moyar, are situated towards the centre of the Reserve on the Sigur Plateau, where the primary vegetation types are dry deciduous and dry thorn forest. MTR supports a variety of mammalian wildlife, including tiger (*Panthera tigris*), leopard (*Panthera pardus*), elephant (*Elephas maximus*), dhole (*Cuon alpinus*), chital (*Axis axis*), sambar (*Rusa unicolor*), blackbuck (*Antilope cervicapra*), black-naped hare (*Lepus nigricollis*), gaur (*Bos gaurus*), and sloth bear (*Melursus ursinus*) [32].

As the villages are situated within the forest, the village boundaries are routinely utilized by humans and domestic animals as well as wildlife (SVK, unpub. data). Masinagudi is a smaller village with a larger population of 1641 households (as of the 2011 Government of India census), has more contiguous built-up area, and hosts several facilities unavailable at other villages including garages, medical stores and clinics, multiple eateries, and a petrol pump. The major source of income for most residents is from tourism, by running homestays or driving safari jeeps. In contrast, Moyar hosts 486 households, with houses clustered in a few small settlements dispersed among fallow or agricultural land, which occupies the majority of the village area. Here, income is primarily derived either from agriculture or from working at the Electricity

Board, which has implemented a hydroelectric project at the Moyar dam. Masinagudi and Moyar

are situated 6.8km apart by road, separated by a straight-line distance of 4.7km. The study area is

shown in Figure 1.

Figure 1: Map of the study area (C) with Masinagudi and Moyar village boundaries outlined in

yellow. The red and blue dots show the location of collaring of all the dogs, with blue dots

indicating those retained for analysis. The inset maps show the study area location in India (A)

and in the state of Tamil Nadu (B).

Dogs are commonly kept by most people in this landscape, primarily as security against wildlife

but also as pets (SVK, unpub. data). Dogs usually accompany owners and livestock [cattle, goats

and sheep) during grazing. Dog management has been ongoing since 1997 in the form of

vaccination and sterilization programmes, carried out by local animal welfare organisations

(India Project for Animals and Nature, IPAN, and Worldwide Veterinary Service-India, WVS-I).

At present, almost 70% of dogs are sterilized and the male:female ratio is almost 1:1 (SVK,

unpub. data).

Collar deployment

GSM GPS collars (by Arcturus Inc., Bengaluru, India) were deployed on the dogs with the

permission and assistance of the owners. A total of 26 dogs were collared between September-

November, 2024. The collars were set to record location every 5 minutes, with an accuracy

(HDOP) of ten metres or less, and we aimed to place each collar for a maximum of ten days. In

practice, the number of deployment days varied as owners sometimes requested that the collar be

removed early or the collar was reported to have come off on its own, resulting in a range of

deployment durations of 3-10 days. The dogs were all photographed and several demographic

variables were recorded using the phone app Epicollect [33], including sex, sterilization status,

approximate age according to the owner, and the GPS location of the owner's home. All dogs

were adults and were fed a minimum of once and a maximum of thrice per day by the owner.

Analysis

The collated GPS data from all the collars comprised 27,850 GPS points after excluding points

collected outside deployment times. Of the initial 26 collared dogs, the data from four dogs were

completely lost due to collar error or premature collar removal by unknown persons. The data

uploaded to Movebank is therefore from the 22 remaining dogs. Outliers were removed through

the online Movebank interface in two steps, first using a speed filter (excluding points that would

require the dog to move at a speed > 5 m/s) and second manually. Points removed manually were

presumably due to poor satellite reception and were implausible for accessibility (e.g., the middle

of a fenced waterbody) or distance reasons (e.g., random single points > 500m away from

clustered preceding and subsequent points). 4193 points were removed with the filter and 186

were removed manually, resulting in a final count of 23,471 points. Subsequent data analysis

was conducted with the data downloaded from Movebank (see Data Availability) in order to

ensure conformity to a standard format. Of the 21 dogs included in this dataset, seven were

excluded during analysis as their data were excessively patchy and the number of data points

collected for each of the six was less than 300, which is approximately only one day of data [one

full day of data would comprise 288 points, as 12 points were collected every hour for 24 hours).

Three additional points were removed due to low accuracy (HDOP > 10). The final dataset

therefore comprised data from 15 dogs with robust datasets.

Movement Metrics:

To understand if the movement of the dogs differed between villages, we utilised multiple

different movement metrics. We calculated the activity range, or the area traversed by the dog

during the short-term period of deployment [19]. To do so, we utilised the package *ctmm* (v1.2.0)

[34] in R version 4.4.1 [35] to estimate the activity range for each dog using autocorrelated

kernel density estimation [AKDE], similar to Wilson-Aggarwal et. al, 2021. Six models were

fitted to each dog's spatial data, and the best fit model was selected using the *ctmm.select*

function. The models included Ornstein-Uhlenbeck (OU), OUF (Ornstein-Uhlenbeck with

foraging), and IID (Independent and Identically Distributed), which does not consider

autocorrelation. After model selection, the 95% AKDE activity range was calculated. We also

calculated two movement metrics; first, the intensity of use, or the active time spent per unit

area, defined as the ratio of the total distance moved during the sampling period to the square

root of the 100% minimum convex polygon (the maximum area used by the animal) [37,38]

using the package *amt* [39]. Second, the mean distance between the owner's home and each GPS

fix for each dog. The distance between home and the dog's location was calculated using the

package *sf* [40] for every set of GPS coordinates recorded, and subsequently averaged to provide

the mean as a measure of how far away from home each dog tended to travel during the study

period. We repeated this process with the distance from the centroid of the activity range to

assess whether there was any significant difference between the two.

The movement data were further categorised with respect to day and night for the mean distance

from home, and with respect to sex of the dog and the village where the dog was collared, for

both movement metrics as well as the activity range. Since these dogs were owned and fed

regularly, the level of human subsidy was not included as a variable in this study. As the

collaring period was relatively short, the difference between sunset and sunset times during the

study was negligible; therefore, we used 0600-1800 hours as day time and 1800-0600 hours as

night. We tested the difference between the movement metrics and activity range with respect to

these variables with the non-parametric Wilcoxon rank sum test.

Habitat Selection:

We also evaluated the selection and movement of the dogs for different habitats. We classified

the areas of each village into settlement, agriculture, fallow or scrubland [uncultivated land

beyond settlements but within village boundaries), and forest land to understand habitat-wise dog

movement (Fig. 2). Forest land included both deciduous forest and thorn forest vegetation types.

We used land use/land cover Sentinel-2 imagery at a resolution of 10 metres, classifying specific

land use types (such as fallow land) manually, given the fine scale of the study area. As an initial

coarse assay of habitat selection, we used landscape imagery of approximately 70 sq. km,

enough to maintain a buffer of 1 km in all directions from both villages, and calculated the

proportion of dog presence with respect to habitat availability of each class type.

We then utilized integrated step-selection analysis, or iSSA [41] to study the fine-scale influence

of habitat covariates on the habitat selection and movement rate of the dogs. iSSAs contrast the

environmental attributes of each animal location with a set of random locations generated from

the distribution of movement parameters (step lengths and turn angles) of the steps the dogs took

[41]. As with other habitat selection analyses, iSSAs compare covariate values at used (animal

GPS locations) and random points (locations generated using the distribution of movement

parameter) for each animal step. iSSAs are considered to be an improved approach to habitat

select analysis as the framework can simultaneously estimate coefficients of habitat selection and

movement rate, potentially increasing the accuracy of both [41,42]. We generated 20 available

steps per used steps from the observed movement parameter distribution of the dogs. As the data

was collected at a very high resolution, we resampled the movement data to 15 minutes and 30

minutes to control and test iSSA models for spatial correlation. The village of origin of the dogs

was considered as a random effect for habitat and movement to elucidate differences in

movement between the two villages. We utilized the Poisson trick [42] and the variability in

movement across villages using the random effect approach [43]. Models were fitted using the

package "glmmTMB" [44] and visualized using package ggplot2 [45]. All codes to reproduce

the analysis are publicly available [see Data Availability).

Figure 2. Map of the study area with classified habitats.

Results

Of the 15 dogs from 15 households, 8 were female and 7 male, with 8 from Masinagudi and 7

from Moyar, of which one male was unsterilized (see Supplementary Table S1). The average age

of the dogs tracked was 5 (minimum 3, maximum 8). The mean number of tracking days was

8.07 (median = 9, range = 3-10). The total number of GPS fixes recorded and used for analysis

was 22277, with an average points per dog of 1485 (range 619-2783). Post outlier filtering, the

precision of the data was high (mean HDOP = 1.66, median = 1.47, range = 0 - 9.13).

The best fit model for all individuals was OU anisotropic, often with a ΔAIC of approximately 2

with the next best model, which was OUF anisotropic (see Supplementary Table S3 for all ΔAIC

values), while IID was ranked last for all individuals. AKDE activity range estimates based on

this model varied across dogs. The lowest activity range was 3.28 ha and the highest was 26.16

ha, estimated for a dog that routinely accompanied her owner during livestock grazing. The mean

activity range was 9.88 ± 7.69 ha (median = 6.09 ha). For distance travelled from home, the

longest distance for each dog ranged from 341.9 m to 1197.1 m, the mean distance from home

across all dogs was 74.44 ± 119.53 m, and the median was 36.12m. 91.8% of all GPS points

were within 250m of the home, while 98.4% were within 500m. Lastly, the mean intensity of use

was 115.94 ± 54.50, and the median was 91.45.

We found no significant difference between mean distance from home and mean distance from

the centroid, with the distance between home and centroid sometimes varying by as little as 10

metres; therefore, all further analysis was conducted with respect to distance from home only.

When the three metrics, activity range, mean distance travelled from home for each dog, and the

intensity of use for each dog, were compared with a Wilcoxon rank sum test with respect to day

and night (for mean distance from home only), sex, and village, the only significant difference (p

value = 0.014) was between intensity of use in each village. The movement metrics for each dog

are presented in Table 1, and the p-values and U statistic for each comparison are presented in

Table 2. The intensity of use and activity ranges compared between villages are provided as a

box plot (Fig. 3).

Table 1: Individual dog movement metrics.

Dog ID number	Activity range [ha)	Activity range 5%-95% CI	Mean distance from home [m]	Intensity of use
D1V1	11.459	10.318 -12.660	76.835	87.835
D2V2	4.485	4.240 - 4.736	58.948	213.082
D3V2	4.189	3.868 - 4.522	50.858	166.233
D4V2	23.860	21.323 - 26.535	124.933	179.025
D5V2	3.389	3.220 - 3.565	41.505	146.008
D6V1	26.163	22.971 - 30.670	127.269	72.462
D7V2	3.277	3.056 - 3.550	308.419	64.976
D8V1	4.564	4.369 - 4.828	45.343	137.936
D9V1	9.876	7.842 - 12.188	52.689	54.607
D10V2	19.606	16.214 - 23.316	99.701	91.446

D11V1	3.807	3.576 - 4.046	39.944	131.479
D12V1	7.189	6.456 - 7.961	54.108	87.589
D13V1	14.398	10.318 - 19.147	79.283	62.863
D14V1	5.797	4.811 - 6.875	49.048	48.887
D15V2	6.087	5.785 - 6.397	55.570	194.622

Includes 95% activity range, mean distance from dog home and intensity of use for all dogs

included in the analysis. ID codes refer to the dog's serial number [D1) and village of origin [V1

or V2 for Masinagudi or Moyar respectively). Further details about dog demographic parameters,

data collected and duration of collaring may be found in Supplementary Table S1.

Table 2: Wilcoxon test results for movement metric comparisons, with respect to different

variables.

Metric	Variable	P-value	U statistic
Mean distance to home vs mean distance to centroid	NA	0.486	130
Mean distance to home	Day/night	0.148	148
Mean distance to home	Sex	0.867	30
Mean distance to home	Village	0.397	20

Activity range	Sex	0.336	37
Activity range	Village	0.336	37
Intensity of use	Sex	0.121	42
Intensity of use	Village	0.014*	7

Significant values (<0.05) are marked with an asterisk.

Figure 3: Histogram showing the differences in the home-range (A) and intensity of area use (B)

for collared dogs from two villages. The values on the top left show the p-value from the

Wilcoxon rank-sum test.

The initial coarse assay of habitat selection found that 85.9% of all dog locations (n=19145) were

located in settlements, followed by 7.8% of locations (n= 1742) in agricultural areas and 2.6%

(n=571) in scrub/fallow land. Together, 96.5% of all locations were in human-settled or human-

modified habitats. With 1.96% of locations (n=436) in deciduous forest and 1.23% (n=273) in

thorn forest, only 3.05% of locations were in the protected area. As the availability of settlements

in the 70 sq. km landscape was 1.85%, while agriculture was 2.76% and fallow land was 2.56%,

compared to 32.4% of deciduous forest and 59.89% of thorn forest, dogs strongly selected for

human modified habitats.

This supported the results of the integrated step-selection analysis, which evaluated the fine-scale

habitat selection of dogs from the study area. The outcome of iSSA was similar for both time

scales (15 min, 30 min), with a higher standard error for the data resampled at 30 minutes (Fig.

4.); therefore, we used the results of the 15 min resampled data for further analysis. Of all

available habitat classes, anthropogenic habitats (that is, settlements and agricultural land) were

significantly preferred, while others were not. However, the habitat-movement parameter

interaction was significant for multiple habitats. The observed step-lengths were significantly

longer than random steps in the forest habitat ($\beta = 0.0092$, p =0.043), whereas they were

marginally significant in agriculture ($\beta = 0.0077$, p =0.077) and scrubland/fallow habitat ($\beta =$

0.0079, p =0.075). Overall, although there was no significant preference or avoidance of habitat

classes apart from anthropogenic ones, the movement speed in each class was substantially

different, with dogs moving with the longest step lengths of approximately 85m-90m in the

forested areas (traversing longer distances between GPS fixes), followed by scrubland and

agricultural fields, where the step lengths were closer to 70m-75m (Fig. 5). The slowest

movement with the shortest step length of 45-50m was recorded in settlements, and there was no

significant difference when compared to random steps. Between villages, the movement trend

was similar in both, but the movement speed was marginally higher in Moyar across all habitat

classes (Fig. 5).

Figure 4: Coefficient estimates from the integrated step-selection analysis for the two different

temporal scales. The orange dots represent the model where the temporal scale for consecutive

fixes was resampled at 15 minutes, and the green dots represent the model where the temporal

scale for consecutive fixes was resampled at 30 minutes.

Figure 5: Estimates of mean step length (speed) for movement estimated for different habitats

and different villages derived using the integrated step-selection function.

Discussion

Our study investigates the movement ecology of 15 owned free-ranging domestic dogs in

Mudumalai Tiger Reserve, India. We found that dogs usually stayed close to their homes, with

the vast majority of recorded locations remaining within 500m of the owner's house, limiting

their potential for significant ecological impact. The best fit model for all individuals was OU

anisotropic, indicating directionally dependent home-ranging behaviour. The activity range for

most dogs was smaller than 10 hectares, and while it ranged from smaller than 5 ha to higher

than 25 ha across individuals, this variation was relatively low compared to the findings of

similar studies, where activity ranges as low as 0.78 ha and as high as 509 ha have been

documented [19]. The maximum distance from home travelled by a dog (1197m) was fairly short

compared to ranging distance in other areas like Cambodia (up to 4km) [19] and Australia (8-

30km) [14], and was similar to the distance travelled in Chile (0.5-1.9km) [46]. Given their fairly

restricted movement patterns, these dogs seem to fall under the class of *stay-at-home* dogs,

according to the three classes defined by Hudson et al. (2017). They display minimal roaming

and are likely to have the majority of the requirements fulfilled at or close to their home, by the

humans they depend on [13]. The range and median of activity ranges are similarly on the lower

side of activity ranges recorded elsewhere [19] (See also Supplementary Table S2).

Estimates of movement metrics (the activity range, mean distance to home, and intensity of use),

when compared based on village, sex of the dog, and day/night, showed no significant difference

apart from intensity of use, which was significantly different between villages (Table 2, Fig. 3).

We did not expect any sex-based differences with respect to any of these metrics, given that all

dogs were neutered or spayed except one male. Additionally, though neutered males have been

recorded as having significantly larger activity ranges than spayed females elsewhere [47], we

did not find that to be true here (Table 2). The lack of significant difference between day and

night mean distance from home was unexpected, as we expected that dogs would move less at

night to minimize chances of encountering wildlife that are more active at village boundaries at

night (such as elephants or leopards). The lack of difference between day and night may be a

product of the short sampling period, or it may be due to the fact that we defined 'night' as post

1800 hours, when dogs might still be quite active for a few hours post sunset (16) or well into the

night (17).

The significant difference in intensity of use between villages, however, indicates that dogs use

their activity ranges differently in each village, despite the similarity in the size of the area used.

In Moyar, the human population is small and village homes are divided into clusters of

settlements separated by large stretches of agricultural land. In Masinagudi, however, the human

population is several times larger and agricultural patches are isolated by well-connected roads

and densely constructed houses. Therefore, we interpret the higher intensity of use by dogs in

Moyar as a pattern of movement restricted to safe, well-known areas, with minimal forays to

novel areas with higher risk of wildlife encounters. This is likely due to the heightened risk

posed by wildlife to dogs, stemming from two factors: the smaller human population with

fragmented clusters of settlements, and the larger, more porous perimeter with the forest and

swathes of empty land inside the village that are conducive to wild animal movement. Indeed,

spotted deer, wild pigs, elephants, and a leopard have previously been camera trapped inside

Moyar (SVK, unpub. data). That these free-ranging dogs adapted different roaming behaviours

depending on their village of origin, even when the two study sites were situated within 5km of

each other, indicates a high degree of behavioural plasticity.

This interpretation is also supported by the habitat use and integrated step selection analysis

results. We found that dogs mostly use human-modified land types, primarily settlement and

agricultural land, with only 3.05% of all fixes located in the protected area. This is similar to

previous findings in India [30], Chad [36] and Chile [17,46]. The results of the step selection

analysis are also similar to previous studies [17,48], where anthropogenic habitats were found to

be the most preferred habitat type. Although dogs did not significantly prefer or avoid any

habitat type except settlements, which were highly preferred, the difference in step length, or

movement speed, in these habitats (Fig. 5) reflected how dogs used them. Forest areas were

traversed almost twice as fast as settlements, and about 1.2 times as fast as agricultural or

scrubland areas. This indicates that forest areas were used in a utilitarian manner, solely for

movement (perhaps exploratory forays or travel between points) while settlements were used for

rest or play. Movement in agricultural and scrubland areas may have been while accompanying

humans, as all the dogs in this study were owned. Additionally, the higher movement speed in

Moyar, where the human population was smaller and more fragmented, while not statistically

significant, highlights how human use of wildlife-adjacent land can closely influence the

movement patterns of domestic commensals like dogs. While previous studies [e.g., 17, 46] have

reported on free-ranging dog habitat usage, utilizing the robust framework of integrated step-

selection analysis to model dog habitat selection allowed us to compare free-ranging dog

movement between two sites at a fine geographical scale, presenting further support for the

human-shield hypothesis [49]. Our findings not only contribute to dog movement and

behavioural ecology, but suggest that this movement - and therefore potential impact on wildlife

- can differ significantly within villages in the same landscape, a nuance that should be taken into

consideration when designing, implementing and monitoring dog management strategies that

aim to curb roaming behaviour.

Fieldwork for this study was primarily limited by logistic and technological difficulties, namely

the unauthorized removal and subsequent loss of collars (as people were apprehensive about the

negative repercussions of their dogs being tracked) and collar malfunction errors resulting in data

loss. These issues reduced our effective sample size from 26 dogs to 15 dogs. We were also

unable to collar truly feral or unowned dogs, which are most likely to range into the forest alone,

as this would have required chemical immobilization by a veterinarian. We were therefore

unable to differentiate dog movement in the presence versus absence of humans. Though this

also constrained the diversity of our sample pool, our findings provide valuable insight into the

movement of sterilized and owned free-ranging dogs, which comprises the vast majority of the

large dog population in the Sigur Plateau region of Mudumalai Tiger Reserve (SVK, unpub.

data). They also provide a standard of comparison to evaluate the impact of sterilization

programmes on dog movement patterns. Lastly, the duration of sampling [up to ten days per dog

and approximately two months overall) was relatively short when compared to other, similar

studies [see Supplementary Table S2). This was a disadvantage, as dog movement may differ

significantly across seasons [36), while this study was limited to a single season study.

Conclusion

In summary, this study reveals that while coarse indices of movement such as dog activity range

remained limited and showed no significant difference between the two villages studied, habitat

use differed as a potential consequence of human presence and habitation providing a shelter

effect. Our findings suggest that owned free-ranging dogs, specifically those sterilized and

provisioned by their owners, may pose a relatively low risk to wildlife in Mudumalai Tiger

Reserve in terms of the size of area used and time spent in the forest. We therefore recommend

that the programmes carried out by local animal welfare NGOs be sustained in the long term

with a stronger push to sterilize and vaccinate feral dogs as well, ideally with consistent financial

and logistical support from relevant local government bodies. While the limited scale of this

study does not allow for direct extrapolation to other regions, it supports the idea that local

context is highly relevant to dog ecology [9], indicating the need for further studies from a broad

range of socio-ecological contexts. In general, research on the fundamental ecology of free-

ranging dogs at the human-wildlife interface is of vital importance in order to effectively

mitigate dog-wildlife conflict.

For future studies, we recommend collaring more broadly across dog demographics (namely

ownership, sex, age and sterilization status) as well as across human and village related

parameters (e.g., length of the village-forest perimeter, household density, agricultural cover) at

larger spatial and temporal scales. In particular, studies from non-protected areas that support

wildlife may not only be easier to carry out but may provide much-needed insight into the dog-

wildlife interface in areas with no restrictions on human movement and land use.

Declarations

Ethics approval and consent to participate

All GPS collars were fitted with the owners' consent, and often by the owners themselves.

Collars were within required weight limits (<250 grams) and caused no physical distress to dogs

during the duration of deployment.

Consent for publication

Not applicable

Availability of data and materials

The dataset generated during this study is available in the Movebank repository under the study

name "Free-ranging dogs - vksanj - Mudumalai TR - GPS collaring", study ID 5328251712, at

the following link -

Codes for data analysis are available on Github at the following link -

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by The Habitats Trust (Seed Grant, March 2024) in a grant provided to

SVK.

Authors' contributions

SVK acquired funding for the study, conducted fieldwork, participated in data analysis, and

drafted the manuscript. NC led data analysis and contributed to manuscript writing. RK provided

logistic and supervisory support, and contributed to manuscript writing. All authors read and

approved the final manuscript.

Acknowledgements

We acknowledge and appreciate the support provided by staff at the Indian Institute of Science

field station in Masinagudi, namely Mr. Sharan B. Gowda, Mr. Albert Suresh, Mr. Bharanaiah,

Mr. Arjunan, and Dr. Suresh H. S. We are grateful to Mr. Arasu for field support, as well as the

dog owners of Masinagudi and Moyar for permission to collar their dogs.

References

1. Gompper ME. The dog-human-wildlife interface: assessing the scope of the problem. In:

Free-ranging dogs and wildlife conservation. 1st ed. OUP Oxford; 2014. p. 9–54.

2. Ortolani A, Vernooij H, Coppinger R. Ethiopian village dogs: Behavioural responses to a

stranger's approach. Appl Anim Behav Sci. 2009 Jul 1;119(3):210–8;

doi:10.1016/j.applanim.2009.03.011.

3. Acosta-Jamett G, Cleaveland S, Cunningham AA, Bronsvoort BM deC. Demography of

domestic dogs in rural and urban areas of the Coquimbo region of Chile and implications for

disease transmission. Prev Vet Med. 2010 May;94(3–4):272–81;

doi:10.1016/j.prevetmed.2010.01.002.

4. Butler JRA, Bingham J. Demography and dog-human relationships of the dog population in

Zimbabwean communal lands. Vet Rec. 2000 Oct;147(16):442–6; 10.1136/vr.147.16.442.

5. Fiorello CV, Noss AJ, Deem SL. Demography, Hunting Ecology, and Pathogen Exposure of

Domestic Dogs in the Isoso of Bolivia. Conserv Biol. 2006 Jun;20(3):762–71;

doi:10.1111/j.1523-1739.2006.00466.x.

6. Kitala P, McDermott J, Kyule M, Gathuma J, Perry B, Wandeler A. Dog ecology and

demography information to support the planning of rabies control in Machakos District,

Kenya. Acta Trop. 2001 Mar 30;78(3):217–30; doi:10.1016/S0001-706X(01)00082-1.

7. Twardek WM, Peiman KS, Gallagher AJ, Cooke SJ. Fido, Fluffy, and wildlife conservation:

The environmental consequences of domesticated animals. Environ Rev. 2017

Dec;25(4):381–95; doi:10.1139/er-2016-0111.

8.   Young JK, Olson KA, Reading RP, Amgalanbaatar S, Berger J. Is Wildlife Going to the

Dogs? Impacts of Feral and Free-roaming Dogs on Wildlife Populations. BioScience. 2011

Feb 1;61(2):125–32; doi:10.1525/bio.2011.61.2.7.

9.   Gompper ME. Adding Nuance to Our Understanding of Dog–Wildlife Interactions and the

Need for Management. Integr Comp Biol. 2021 Jul 1;61(1):93–102;

doi:10.1093/icb/icab049.

10. Doherty TS, Dickman CR, Glen AS, Newsome TM, Nimmo DG, Ritchie EG, et al. The

global impacts of domestic dogs on threatened vertebrates. Biol Conserv. 2017 Jun

1;210:56–9; doi:10.1016/j.biocon.2017.04.007.

11. Dürr S, Dhand NK, Bombara C, Molloy S, Ward MP. What influences the home range size

of free-roaming domestic dogs? Epidemiol Infect. 2017 May;145(7):1339–50;

doi:10.1017/S095026881700022X.

12. Dürr S, Ward MP. Roaming behaviour and home range estimation of domestic dogs in

Aboriginal and Torres Strait Islander communities in northern Australia using four different

methods. Prev Vet Med. 2014 Nov 15;117(2):340–57; doi:10.1016/j.prevetmed.2014.07.008.

13. Hudson EG, Brookes VJ, Dürr S, Ward MP. Domestic dog roaming patterns in remote

northern Australian indigenous communities and implications for disease modelling. Prev

Vet Med. 2017 Oct 1;146:52–60; doi:10.1016/j.prevetmed.2017.07.010.

14. Meek PD. The movement, roaming behaviour and home range of free-roaming domestic

dogs, Canis lupus familiaris, in coastal New South Wales. Wildl Res. 1999;26(6):847–55;

doi:10.1071/wr97101.

15. Molloy S, Burleigh A, Dürr S, Ward M. Roaming behaviour of dogs in four remote

Aboriginal communities in the Northern Territory, Australia: preliminary investigations.

Aust Vet J. 2017;95(3):55–63; doi:10.1111/avj.12562.

16. Sparkes J, Körtner G, Ballard G, Fleming PJS. Spatial and temporal activity patterns of

owned, free-roaming dogs in coastal eastern Australia. Prev Vet Med. 2022 Jul

1;204:105641; doi:10.1016/j.prevetmed.2022.105641.

17. Schüttler E, Saavedra-Aracena L, Jiménez JE. Spatial and temporal plasticity in free-ranging

dogs in sub-Antarctic Chile. Appl Anim Behav Sci. 2022 May 1;250:105610;

doi:10.1016/j.applanim.2022.105610.

18. Muinde P, Bettridge JM, Sousa FM, Dürr S, Dohoo IR, Berezowski J, et al. Who let the dogs

out? Exploring the spatial ecology of free-roaming domestic dogs in western Kenya. Ecol

Evol. 2021;11(9):4218–31; doi:10.1002/ece3.7317.

19. Ladd R, Meek P, Eames JC, Leung LKP. Activity range and patterns of free-roaming village

dogs in a rural Cambodian village. Wildl Res. 2023 Jul;51(1); doi:10.1071/WR23024.

20. Pérez GE, Conte A, Garde EJ, Messori S, Vanderstichel R, Serpell J. Movement and home

range of owned free-roaming male dogs in Puerto Natales, Chile. Appl Anim Behav Sci.

2018 Aug 1;205:74–82; doi:10.1016/j.applanim.2018.05.022.

21. Ruiz-Izaguirre E, van Woersem A, Eilers K (c.) H a. M, van Wieren SE, Bosch G, van der

Zijpp AJ, et al. Roaming characteristics and feeding practices of village dogs scavenging

sea-turtle nests. Anim Conserv. 2015;18(2):146–56; doi:10.1111/acv.12143.

22. Belsare A, Vanak AT. Modelling the challenges of managing free-ranging dog populations.

Sci Rep. 2020 Nov 2;10(1):18874; doi:10.1038/s41598-020-75828-6.

23. Reddy CS, Sreelekshmi S, Jha CS, Dadhwal VK. National assessment of forest

fragmentation in India: Landscape indices as measures of the effects of fragmentation and

forest cover change. Ecol Eng. 2013 Nov 1;60:453–64; doi:10.1016/j.ecoleng.2013.09.064.

24. Venkataraman K, Sivaperuman C. Biodiversity Hotspots in India. In: Sivaperuman C,

Venkataraman K, editors. Indian Hotspots: Vertebrate Faunal Diversity, Conservation and

Management Volume 1. Singapore: Springer; 2018. p. 1–27. Available from:

https://doi.org/10.1007/978-981-10-6605-4_1

25. Belsare AV, Vanak AT, Gompper ME. Epidemiology of Viral Pathogens of Free-Ranging

Dogs and Indian Foxes in a Human-Dominated Landscape in Central India. Transbound

Emerg Dis. 2014;61(s1):78–86; doi:10.1111/tbed.12265.

26. Mahar N, Habib B, Hussain SA. Do we need to unfriend a few friends? Free-ranging dogs

affect wildlife and pastoralists in the Indian Trans-Himalaya. Anim Conserv. 2024

Feb;27(1):53-64; doi: 10.1111/acv.12876.

27. Vanak AT, Thaker M, Gompper ME. Experimental examination of behavioural interactions

between free-ranging wild and domestic canids. Behav Ecol Sociobiol. 2009 Dec

1;64(2):279–87; doi:10.1007/s00265-009-0845-z.

28. Home C, Bhatnagar YV, Vanak AT. Canine Conundrum: domestic dogs as an invasive

species and their impacts on wildlife in India. Anim Conserv. 2018;21(4):275–82;

doi:10.1111/acv.12389.

29. Tiwari HK, Bruce M, O'Dea M, Robertson ID. Utilising Group-Size and Home-Range
Characteristics of Free-Roaming Dogs (FRD) to Guide Mass Vaccination Campaigns against
Rabies in India. Vaccines. 2019 Dec;7(4):136; doi:10.3390/vaccines7040136.

30. Vanak AT, Gompper ME. Interference competition at the landscape level: the effect of free-
ranging dogs on a native mesocarnivore. J Appl Ecol. 2010;47(6):1225–32;
doi:10.1111/j.1365-2664.2010.01870.x.

32. Suresh HS, Dattaraja HS, Sukumar R. Tree flora of mudumalai sanctuary, Tamil Nadu,
Southern India. Indian For. 1996;122:507–19.

33. Aanensen DM, Huntley DM, Menegazzo M, Powell CI, Spratt BG. EpiCollect+: linking
smartphones to web applications for complex data collection projects. F1000Research. 2014
Aug 20;3:199; doi:10.12688/f1000research.4702.1.

34. Fleming CH, Calabrese JM. ctmm: Continuous-Time Movement Modeling. 2023. Available
from: https://github.com/ctmm-initiative/ctmm

35. R Core Team. R: a Language and Environment for Statistical Computing. (Internet]. Vienna,
Austria.: R Foundation for Statistical Computing; 2021. Available from: https://www.R-
project.org/

36. Wilson-Aggarwal JK, Goodwin CED, Moundai T, Sidouin MK, Swan GJF, Léchenne M, et
al. Spatial and temporal dynamics of space use by free-ranging domestic dogs Canis
familiaris in rural Africa. Ecol Appl. 2021;31(5):e02328; doi:10.1002/eap.2328.

37. Almeida PJAL, Vieira MV, Kajin M, Forero-Medina G, Cerqueira R. Indices of movement

behaviour: conceptual background, effects of scale and location errors. Zool Curitiba. 2010

Oct;27:674–80; doi:10.1590/S1984-46702010000500002.

38. Loretto D, Vieira MV. The Effects of Reproductive and Climatic Seasons on Movements in

the Black-Eared Opossum (Didelphis aurita Wied-Neuwied, 1826). J Mammal. 2005 Apr

15;86(2):287–93; doi:10.1644/BEH-117.1.

39. Signer J, Fieberg J, Avgar T. Animal movement tools (amt): R package for managing

tracking data and conducting habitat selection analyses. Ecol Evol. 2019;9:880–90.

40. Pebesma E. Simple Features for R: Standardized Support for Spatial Vector Data. R J.

2018;10(1):439–46.

41. Avgar T, Potts JR, Lewis MA, Boyce MS. Integrated step selection analysis: bridging the

gap between resource selection and animal movement. Methods Ecol Evol. 2016;7(5):619–

30; doi:10.1111/2041-210X.12528.

42. Muff S, Signer J, Fieberg J. Accounting for individual-specific variation in habitat-selection

studies: Efficient estimation of mixed-effects models using Bayesian or frequentist

computation. J Anim Ecol. 2020;89(1):80–92; doi:10.1111/1365-2656.13087.

43. Chatterjee N, Wolfson D, Kim D, Velez J, Freeman S, Bacheler NM, et al. Modelling

individual variability in habitat selection and movement using integrated step-selection

analysis. Methods Ecol Evol. 2024;15(6):1034–47; doi:10.1111/2041-210X.14321.

44. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.

glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized

587   Linear Mixed Modeling. R J. 2017;9(2):378–400.

588 45. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York; 2016.

589 46. Sepúlveda M, Pelican K, Cross P, Eguren A, Singer R. Fine-scale movements of rural free-

590   ranging dogs in conservation areas in the temperate rainforest of the coastal range of

591   southern Chile. Mamm Biol. 2015 Jul 1;80(4):290–7; doi:10.1016/j.mambio.2015.03.001.

592 47. Sparkes J, Körtner G, Ballard G, Fleming PJS, Brown WY. Effects of Sex and Reproductive

593   State on Interactions between Free-Roaming Domestic Dogs. PLOS ONE. 2014 Dec

594   26;9(12):e116053; doi:10.1371/journal.pone.0116053.

595 48. Cunha Silva L, Friker B, Warembourg C, Kanankege K, Wera E, Berger-González M, et al.

596   Habitat selection by free-roaming domestic dogs in rabies endemic countries in rural and

597   urban settings. Sci Rep. 2022 Dec 3;12(1):20928; doi:10.1038/s41598-022-25038-z.

598 **Supplementary files**

599 Data:

600 Table S1: Dog demographic data — Dog ID number, age, sex, sterilisation status, village of

601 origin, number of fixes and days collared for each dog.

602 Table S2: Home/activity range comparisons from global studies — expanded table adopted from

603 Ladd et al., 2023, with home/activity range estimates and sampling durations from dog telemetry

604 studies across countries.

605 Table S3: ΔAIC comparison across activity range models — ΔAIC values for all trialled models

606 during activity range estimation