Of Shared Homes and Pathways: Free-Ranging Dog Movement and

Habitat Use in a Human-Wildlife Landscape in India

Sanjana Vadakke Kuruppath*, #12, Valmiki Nagar, Thiruvanmiyur, Chennai 600041, India
Nilanjan Chatterjee*, Senckenberg Biodiversity and Climate Research Institute, Frankfurt am
Main, 60325, Germany
Ramesh Krishnamurthy, Wildlife Institute of India, Dehradun, Uttarakhand 248001, India
7
8 *Corresponding authors:
9 vksanjana@gmail.com
10 nilanjan.chatterjee@senckenberg.de
11
12

Abstract

Background: Free-ranging dogs are widely considered to have negative impacts on wildlife in

shared landscapes. Understanding their use of these landscapes is therefore essential to develop

effective management strategies in wildlife-adjacent regions. In this study, we investigated the

movement patterns of free-ranging dogs in the protected area matrix of Mudumalai Tiger

Reserve (MTR) in Southern India.

Methods: Fifteen owned free-ranging dogs (FRD) were collared from two adjacent villages with

different levels of human habitation. We hypothesized that increased settlement cover would

provide dogs with a greater human shield effect, allowing them more freedom of movement. We

estimated dog activity ranges and movement patterns using autocorrelated kernel density

estimates (AKDE). Large-scale movement was characterized through multiple metrics (intensity

of space use and mean distance from home) and compared across villages to understand

underlying drivers. Lastly, we quantified fine-scale movement and habitat selection using

integrated step selection analysis (iSSA).

Results: We found that FRD in MTR had directionally dependent home-ranging behavior and

small activity ranges, with the mean AKDE activity range being 9.88 ± 7.69 ha (median = 6.09

ha, range 3.28-26.16 ha), and primarily utilized the area within 500 m of their homes. None of

the movement metrics varied significantly between villages, except for intensity of use, which

was higher in one village than the other, suggesting that dogs in the less populated village

perceived a greater threat from surrounding wildlife and were more driven to seek refuge inside

their activity ranges. iSSA revealed that dogs selected strongly for human settled or human-

modified habitats, and moved significantly faster in forest land than any other habitat.

Conclusions: Our findings indicate that owned FRD pose a relatively lower threat to wildlife in

MTR than feral dogs may. They also support the human shield hypothesis, show that FRD

display behavioral plasticity at fine scales of <5km, and highlight the role of human land use

intensity in shaping the movement of domestic dogs. As humans can thus mediate potential

effects of dogs on wildlife, anthropogenic factors should be taken in consideration when

designing management strategies that aim to curb dog movement.

Key words: Human-wildlife conflict, movement ecology, human shield hypothesis, step-

selection analysis

Introduction

As the most abundant carnivore on earth with a long history of domestication [1], domestic dogs (*Canis familiaris*) are almost as widely distributed as humans across the world [1], with rural dogs forming a sizable category of the population in underdeveloped areas [2]. These dogs are highly likely to be free-ranging, with previous studies on rural dog demography across the world indicating that a major proportion of rural dogs are free to roam without any restraint, whether alone or in the company of humans [3–6]. In areas close to natural ecosystems, free-ranging dogs (FRD) can have a range of negative impacts on wildlife [7,8], ranging from mild to severe depending on the local socio-ecological context [9]. Broadly, the most common negative impact of domestic dogs on threatened wildlife is predation, estimated at 78.9%, followed by disturbance (such as barking, chasing or harassment) and disease transmission at 7.0% and 4.5% respectively, and lastly competition with native carnivores (1.5%) and hybridization with wild canids (1.0%) [10]. While the intensity of these impacts is unclear, there is no doubt that they are globally pervasive [10].

In order to clarify the true extent of dog-related interference with wildlife, understanding how FRD utilize wildlife-adjacent landscapes is a necessary avenue of research, as patterns of movement into wildlife habitat underpin all subsequent impacts on wild species. These patterns can vary widely across ecosystems, with a metric as fundamental as the home range (or the activity range, for short-term studies) varying regionally, seasonally, or individually, as previous studies from Australia [11–16], Chile [17], Kenya [18], Cambodia [19], and Mexico [20,21] have shown. Therefore, aspects of dog movement ecology that are of particular interest from a wildlife conservation perspective, such as ranging behavior and habitat selection, may be equally varied

and may be influenced by a range of natural as well as anthropogenic factors. Indeed, given the

highly social and commensal relationship between dogs and humans [1], even subtle

anthropogenic factors may significantly influence dog movement and behavior [10]. For

example, dogs that disrupted sea turtle nests to scavenge for eggs in Mexico were fed

significantly less food by their owners than to dogs that did not scavenge [21].

While anthropogenic factors are usually considered to be resources that humans actively provide

dogs with, such as food, there may also be passive benefits to existing in close proximity to

humans. The human shield effect, where prey species are hypothesized to seek refuge from

predators that fear humans by increasing their use of human-modified areas [22], is an example

of one such potential benefit. Although this hypothesis was developed with reference to the

behavior of wild prey species [22], a diverse range of apex carnivores have also been

documented to prey on dogs across the world [23]. There is, therefore, scope for FRD to also

benefit from the protection offered by human presence, by using settlement areas to avoid

encountering predators. As a result, different levels of habitation could affect how freely dogs

utilize landscapes shared with wildlife, providing new insight into how FRD patterns of

movement differ across anthropogenic contexts.

The need to elucidate such potential patterns underlying dog movement ecology is particularly

pressing in India, where the majority of the dog population is free ranging [24] and may often

come into contact with increasingly fragmented natural ecosystems [25] that harbor a rich

diversity of wildlife [26]. Previous studies have investigated other impacts of dogs on wildlife in

India [27–29], with one study recording reports of negative interactions between dogs and 80

wild species [30]; however, to our knowledge, only two studies have estimated FRD movement

ecology in rural or wildlife adjacent areas, both in the state of Maharashtra [31,32], of which

only Vanak and Gompper (2010), utilized VHF telemetry to estimate home range and habitat

use. While non-invasive methods such as camera trapping can be used to estimate movement

parameters, they are relatively time consuming, effort-intensive [33], and can be spatially

constrained compared to telemetry (specifically GPS telemetry), which allows for the collection

of individual-level, precise, fine-scale data around the clock across large areas [34]. When

studying FRD that may exhibit highly context-dependent spatial behavior, such data become

particularly valuable.

We address this research gap with a fine-scale GPS telemetry study on rural FRD from the

multiple-use landscape of a Tiger Reserve in Southern India, where we investigate how FRD

movement and utilization of multiple habitat types are affected by different levels of

anthropogenic land use intensity. We collared FRD in two different villages, both situated inside

the Reserve boundary and zone of influence, with varying levels of human population, built-up

area, and village-forest boundary length. We hypothesized that while FRD would primarily use

settlement areas, a higher proportion of built-up land, in conjunction with a smaller village-forest

boundary, would produce a human shield effect to allow dogs more freedom of movement in a

forested landscape harboring large carnivore species. To test this hypothesis, we designed a

study with two objectives: 1. Estimate and compare FRD activity ranges and movement metrics

between villages and 2. Evaluate how FRD use of habitat types differs between villages.

Materials and Methods

Study area

The study was conducted in the villages of Masinagudi and Moyar, in Mudumalai Tiger Reserve

(MTR), Tamil Nadu, India (Figure 1). MTR is a protected area contiguous with multiple other

wildlife sanctuaries, forming an integral part of the Nilgiri Biosphere Reserve in the highly

biodiverse Western Ghats. Several villages are embedded within MTR, forming a matrix

landscape where human settlements and major roads are interspersed within the forest. The study

sites, Masinagudi and Moyar, are situated towards the center of the Reserve on the Sigur Plateau,

where the primary vegetation types are dry deciduous and dry thorn forest. MTR supports a

variety of mammalian wildlife, including tiger (*Panthera tigris*), leopard (*Panthera pardus*),

elephant (*Elephas maximus*), dhole (*Cuon alpinus*), chital (*Axis axis*), sambar (*Rusa unicolor*),

blackbuck (*Antilope cervicapra*), gaur (*Bos gaurus*), and sloth bear (*Melursus ursinus*) [35].

Leopards in particular have been documented to prey on dogs in India since colonial times [23]

and commonly do so in MTR at present (SVK, unpub. data).

Masinagudi is smaller in area (2.29 km^2) with a larger population of 1641 households (as of the

2011 Government of India census; the current number is likely higher), has more dense and

contiguous built-up area, and has the most developed infrastructure of all villages on the Sigur

Plateau. A major source of income for most Masinagudi residents is from tourism, by running

homestays or driving safari jeeps. In contrast, Moyar is slightly larger (2.37 km^2) but hosts only

486 households, with houses clustered in a few small settlements separated by stretches of fallow

or agricultural land, which occupy the majority of the village area. Here, income is primarily

derived either from agriculture or from working at a hydroelectric project at the Moyar dam. As

the villages are situated within the forest, village-forest boundaries are routinely utilized by

humans and domestic animals as well as wildlife in both places, although several wild species

(including spotted deer (chital), sambar, wild pig, elephant, and leopard) enter Moyar more

frequently than Masinagudi (SVK, unpub. data). Masinagudi and Moyar are situated 4.7 km

apart on the Sigur Plateau, connected by a 6.8 km long road.

Figure 1: Map of the study area, showing dog collaring locations inside the villages of

Masinagudi and Moyar (A) inside MTR (B) in peninsular India (C). Collaring locations are

shown for the dogs whose data were included in this study.

Most people commonly keep dogs in this landscape, primarily as security against wildlife but

also as pets (SVK, unpub. data). Dog reproductive management has been ongoing since 1997 in

the form of vaccination and sterilization programs, carried out by local animal welfare

organizations (India Project for Animals and Nature, IPAN, and Worldwide Veterinary Service-

India, WVS-I). At present, almost 70% of dogs are vaccinated and sterilized, and the

male:female ratio is almost 1:1 (SVK, unpub. data).

Collar deployment

GSM GPS collars (by Arcturus Inc., Bengaluru, India) were deployed on owned dogs with the

permission and assistance of the owners. These collars relied on mobile 2G network to transmit

collected data to a data server. A total of 26 dogs were collared between September-November,

2024. The collars were set to record a location every 5 minutes, with an accuracy (measured by

horizontal dilution of precision, or HDOP) of ten meters or less, and we aimed to collect data for

maximum of ten days for each dog. Ten days was thought to be a reasonable duration, as other

studies have tracked dogs for shorter periods [11,18,36]. Additionally, residents were aware that

the Forest Department prohibits the presence of dogs in the forest and were afraid that allowing

their dogs to be collared would result in punitive action, making it difficult to request that collars

be kept on their dogs for longer. Indeed, in practice, the number of deployment days varied as

owners sometimes requested that the collar be removed early, the collar was removed without

our knowledge, or the collar was reported to have fallen off on its own, resulting in a range of

deployment durations of 3-10 days.

We selected adult FRD at random from households in several different localities within each

village in order to obtain a diverse sample pool, although we attempted to maintain an even sex

ratio. Collaring was accomplished after the owner gave permission and in many cases the collar

was placed by the owner under supervision of the collaring team. Most collared FRD belonged to

people living and working in the interior of the village, as dog owners living near the village-

forest boundary were especially reluctant to allow their dogs to be collared. Owners also reported

that their dogs sometimes chose to stay near the house or roam on their own, rather than

accompany them during the day. Therefore, while dogs in MTR generally go with their owners

to agricultural fields or during livestock grazing, we did not expect collared FRD to closely

mirror human movement patterns, and no collared dog was expected to routinely visit the forest

in the company of their owner. In particular, it should be noted that collared FRD presence in the

forest would not reflect associated human movement for hunting; apart from the danger posed to

humans by elephants, sloth bears and large carnivores, hunting is strictly illegal in protected

areas in India. Therefore, even if people did engage in any kind of poaching, they would be

highly unlikely to do so regularly, and would not take a collared FRD with them.

After collaring, the dogs were photographed and several demographic variables were recorded

using the mobile phone application Epicollect [37], including sex, sterilization status,

approximate age according to the owner, and the GPS location of the owner's home. All dogs

were adults and were fed a minimum of once and a maximum of three times per day by the

owner.

Analysis

We collated the GPS data that was successfully collected during deployment and uploaded them

to Movebank, a free online repository for animal movement data from across the world [38].

Outliers were removed through the online Movebank interface in two steps, first using a speed

filter (excluding points that would require the dog to move at a speed > 5 m/s) and second

manually. 5 m/s was considered a lenient upper limit, since while dogs are capable of reaching

this speed, they would be unlikely to maintain it consistently for the five-minute interval between

fixes to run for a stretch of 1.5 km. We considered that speeds below this limit would account for

any potential encounters with wildlife, whether chasing prey species or escaping from predator

species.

Points removed manually were implausible for accessibility reasons (e.g., the middle of a fenced

waterbody) or distance reasons (e.g., random single points > 500 m away from clustered

preceding and subsequent points), and were presumably due to poor satellite reception.

Subsequent data analysis was conducted with the data downloaded from Movebank (see Data

Availability) in order to ensure conformity to a standard format. Dogs with excessively patchy

data (fewer than 300 points collected) were excluded, since these constituted approximately only

one day of data and they often had gaps of multiple days in between fixes, making it unfeasible

to use them for a robust analysis. Lastly, points with an accuracy (HDOP) of less than 10 m were

removed from the dataset.

Movement Metrics:

To examine the difference in dog movement patterns between villages, we utilized multiple

different movement metrics, namely the activity range, the intensity of use, and the mean

distance between the owner's home and each GPS fix for each dog. We estimated the activity

range for each dog, or the area traversed by the dog during the short-term period of deployment

[19] with the package *ctmm* (v1.2.0) [39] in R version 4.4.1[40], using autocorrelated kernel

density estimation (AKDE), similar to Wilson-Aggarwal et. al, 2021. Six models were fitted to

each dog's spatial data, and the best fit model was selected using the *ctmm.select* function, which

ranks models according to the Aikake Information Criterion [39]. The models included Ornstein-

Uhlenbeck (OU), OUF (Ornstein-Uhlenbeck with foraging), and IID (Independent and

Identically Distributed), of which IID alone does not consider autocorrelation. After model

selection, we estimated the 95% AKDE activity range for the dog based on the best fit model

using the *akde* function [39].

The second movement metric, the intensity of use, is a measure of the active time spent by the

animal per unit area, defined as the ratio of the total distance moved during the sampling period

to the square root of the 100% minimum convex polygon (the maximum area used by the

animal) [41,42]. We estimated it with the package *amt* [43], using the *intensity_use* function. The

third movement metric was the mean distance between the owner's home and each GPS fix for

each dog. We calculated the distance between home and the dog's location using the package *sf*

[44] for every set of GPS coordinates recorded, and subsequently averaged the distances to

provide the mean as a measure of how far away from home each dog tended to travel. We

repeated this process with the distance from the centroid of the activity range to assess whether

there was any significant difference between the two.

We further categorized the movement data with respect to day and night for the mean distance

from home, and with respect to sex of the dog and the village of origin for all three movement

metrics. We hypothesized that FRD would move more during the day to avoid wild animals at

night, that a higher amount of built-up area and a smaller village-forest boundary in Masinagudi

would allow FRD more freedom of movement than in Moyar, and that males would cover larger

areas that than females despite both being sterilized, as has been recorded elsewhere [e.g. 45].

Overall, since these FRD were owned and provisioned, we expected that they would tend to

remain in the vicinity of their owners' homes for most of the time they were active. The level of

human subsidy was not included as a variable in this study since feeding was regular. As the

collaring period was relatively short, the difference between sunrise and sunset times during the

study was negligible; therefore, we used 0600-1800 hours as day time and 1800-0600 hours as

night. We tested the difference between movement metrics with respect to these variables using

the non-parametric Wilcoxon rank sum test.

We also estimated the dog daily activity pattern using the nonparametric kernel density

estimation method, expecting FRD to occupy a strongly diurnal diel niche. GPS data from

individual dogs were used to calculate movement rates as the distance between successive

locations divided by the time between them. We then converted the rates to randomly distributed

pseudo-detections within the time interval between GPS fixes, so that a higher speed translates to

more pseudo-detections, to obtain a general, continuous activity pattern [46].

Habitat Selection:

To evaluate the selection and movement of the dogs in different habitats, we first classified the

areas of each village into settlement, agriculture, fallow or scrubland (uncultivated land beyond

settlements but within village boundaries), and forest land (Fig. 2). Forest land included both

deciduous forest and thorn forest vegetation types. We used land-use/cover Sentinel-2 imagery at

a resolution of 10 meters, classifying specific habitat types (such as fallow/scrub land) manually,

given the fine scale of the study area. As an initial coarse assay of habitat selection, we used

landscape imagery of approximately 70 sq. km, enough to maintain a buffer of 1 km in all

directions from both villages, and calculated the proportion of dog presence with respect to

habitat availability for each dog.

We then utilized integrated step-selection analysis, or iSSA [47] to study the fine-scale influence

of habitat covariates on the habitat selection and movement rate of the dogs. iSSAs contrast the

environmental attributes of each animal location with a set of random locations generated from

the distribution of movement parameters (step lengths and turn angles) of the steps the dogs took

[47]. As with other habitat selection analyses, iSSAs compare covariate values at used (animal

GPS locations) and random points (locations generated using the distribution of the movement

parameter) for each animal step. iSSAs are considered to be an improved approach to habitat

select analysis as the framework can simultaneously estimate coefficients of habitat selection and

movement rate, potentially increasing the accuracy of both [47,48]. We generated 20 available

steps per used step from the observed movement parameter distribution of the dogs. As the data

was collected at a very high resolution, we resampled the movement data to 15 minutes and 30

minutes to control and test iSSA models for spatial autocorrelation. The village of origin of the

dogs was considered as a random effect for habitat and movement to elucidate differences in

movement between the two villages. We utilized the Poisson trick [48] and the variability in

movement across villages using the random effect approach [49]. Models were fitted using the

package "glmmTMB" [50] and visualized using package ggplot2 [51]. All codes to reproduce

the analysis are publicly available (see Data Availability).

Figure 2. Land use/cover classes of the study area. Black lines indicate dog movement tracks in

the vicinity of each village. One track is highlighted in grey in each village to exemplify the dogs

with the largest activity ranges from each village.

Results

Of the initial 26 collared dogs, the data from four dogs were completely lost due to collar error or

premature collar removal by unknown persons, leaving us with data from 22 dogs totaling

27,850 GPS points. Of these, 4144 points were removed as they were null points (i.e. points

recorded as 0,0 when the collar failed to acquire a GPS fix), 49 points were removed with the

Movebank speed filter, 186 were removed manually, and three points were removed due to low

accuracy, resulting in a final count of 23,468 points. Of the 22 dogs included in the Movebank

dataset, seven were excluded during analysis as their data were few in number and excessively

patchy, presumably due to satellite connection or mobile network coverage issues. The final

dataset therefore comprised GPS location data from 15 FRD with robust datasets, with the

number of points totaling 22277.

The set of 15 FRD were almost equally divided both in terms of sex as well as the village of

origin, with 8 females and 7 males, and 8 FRD from Masinagudi and 7 FRD from Moyar. Of

these, all were sterilized apart from one male dog (Table 1). The mean age of the all dogs tracked

was 5 (range = 3 - 8). The mean number of tracking days was 8.07 ± 2.40 (range = 3 - 10). The

mean number of GPS points per dog was 1485 ± 738 (range 619 - 2783) and post outlier

filtering, the precision of the data was high (mean HDOP = 1.66 m $\pm$ 0.70 m, range = 0 - 9.13

m).

The activity range best fit model for individuals from both villages was OU anisotropic, often

with a ΔAIC of approximately 2 with the next best model, which was OUF anisotropic (see

Supplementary Table S3 for all ΔAIC values), while IID was ranked last for all individuals,

indicating that collared FRD displayed directionally dependent home-ranging behavior. The

mean AKDE activity range estimate across all dogs was 9.88 ± 7.69 ha (median = 6.09 ha, range

= 3.28 ha – 26.26 ha) (Table 2). We found no significant difference between mean distance from

home and mean distance from the centroid, with the distance between home and centroid

sometimes varying by as little as 10 meters; therefore, all further analysis was conducted with

respect to distance from home only.

The mean distance from home across all dogs was 74.44 ± 119.53 m (median = 36.12 m, range =

0 m - 1197.06 m) (Table 2). 91.8% of all GPS points (n=20453 of 22277) were within 250 m of

the home, while 98.4% (n=21917) were within 500 m. The mean intensity of use was $115.94 \pm$

54.46 (range = 48.89 – 213.08, median = 91.45) (Table 2). When the three metrics, activity

range, intensity of use, and mean distance travelled from home were compared with a Wilcoxon

rank sum test with respect to sex, village, and day and night (the last used as a variable only for

mean distance from home), the only significant difference was between intensity of use in each

village, which was higher in Moyar than Masinagudi (Table 3, Fig. 3). The activity pattern

analysis produced a relatively unimodal distribution, with a broad peak from the early morning

hours (beginning at 1:00) until around noon with a dip at 6:00, followed by a steep decline until

late night (Fig. 4). The collared dogs therefore displayed a partly diurnal, partly nocturnal

activity pattern.

Table 1: Individual dog demographic parameters and tracking data. Values rounded to two

decimal points.

Dog ID number	Village of origin	Sex	Age	Sterilization status	No. of fixes	No. of days tracked
D1	Masinagudi	Male	7	Yes	1169	8
D6	Masinagudi	Female	3	Yes	906	10
D8	Masinagudi	Male	5	Yes	2174	10

D9	Masinagudi	Female	4	Yes	647	7
D11	Masinagudi	Female	5	Yes	1140	8
D12	Masinagudi	Male	3	Yes	1501	8
D13	Masinagudi	Female	5	Yes	619	3
D14	Masinagudi	Female	3	Yes	635	9
D2	Moyar	Female	Unknown	Yes	2446	9
D3	Moyar	Male	Unknown	Yes	2783	10
D4	Moyar	Male	5	No	2172	10
D5	Moyar	Female	7	Yes	1874	10
D7	Moyar	Female	8	Yes	757	3
D10	Moyar	Male	8	Yes	1207	6
D15	Moyar	Male	5	Yes	2247	10

348   Table 2: Individual dog movement metrics. Values rounded to two decimal points.

Dog ID number	Village of origin	Activity range (ha)	Activity range, 95% confidence interval	Distance from home, mean (m)	Distance from home, range (m)	Intensity of use
D1	Masinagudi	11.46	10.32 -12.66	76.84	0.24 – 667.73	87.84
D6	Masinagudi	26.16	22.97 - 30.67	127.27	0 – 606.64	72.46
D8	Masinagudi	4.56	4.37 - 4.83	45.34	0 – 859.40	137.94
D9	Masinagudi	9.88	7.84 - 12.19	52.69	1.56 – 900.99	54.61
D11	Masinagudi	3.81	3.58 - 4.05	39.94	1.09 – 341.86	131.48

D12	Masinagudi	7.19	6.46 - 7.96	54.12	0 – 923.06	87.59
D13	Masinagudi	14.40	10.32 - 19.12	79.28	1.56 – 690.90	62.86
D14	Masinagudi	5.80	4.81 - 6.88	49.05	0 – 777.84	48.89
D2	Moyar	4.49	4.24 - 4.74	58.95	1.09 – 428.27	213.08
D3	Moyar	4.19	3.87 - 4.52	50.86	0 – 913.85	166.23
D4	Moyar	23.86	21.32 - 26.54	124.93	0 – 1197.06	179.03
D5	Moyar	3.39	3.22 - 3.57	41.51	1.09 – 576.37	146.01
D7	Moyar	3.28	3.06 - 3.55	308.42	10.95 – 668.06	64.98
D10	Moyar	19.61	16.21 - 23.32	99.70	1.09 – 888.80	91.45
D15	Moyar	6.09	5.79 - 6.40	55.57	0 – 910.11	194.62

350   Table 3: Wilcoxon test results for movement metric comparisons, with respect to different

351   variables. Significant values (<0.05) are marked with an asterisk.

Metric	Variable	P-value	U statistic
Mean distance to home vs mean distance to centroid	NA	0.486	130
Mean distance to home	Day/night	0.148	148
Mean distance to home	Sex	0.867	30

Mean distance to home	Village	0.397	20
Activity range	Sex	0.336	37
Activity range	Village	0.336	37
Intensity of use	Sex	0.121	42
Intensity of use	Village	0.014*	7

Figure 3: Box plots showing the differences in the home-range (A) and intensity of area use (B)

for collared dogs from two villages (visualizing select values from Table 3). The values on the

top left show the p-value from the Wilcoxon rank-sum test (significant values marked by an

asterisk).

Figure 4: Kernel density plot showing collared FRD activity pattern across 24 hours.

The initial coarse assay of habitat selection found that 85.94% of all dog locations (n=19145)

were in settlements, followed by 7.82% of locations (n=1742) in agricultural areas and 2.56%

(n=571) in fallow/scrubland. Together, 96.32% of all locations were in human-settled or human-

modified habitats. With 1.96% of locations (n=436) in deciduous forest and 1.23% (n=273) in

thorn forest, 3.18% of all GPS locations were in the protected area. When examined by

individual, 99.5% of the points located in the forest were from four FRD (Table S2,

supplementary material), indicating that the other 11 either rarely ventured or did not venture

into the forest at all during the sampling period. As the availability of settlements in the 70 sq.

km landscape was 1.85%, while agriculture was 2.76% and fallow/scrub land was 2.56%,

compared to 32.4% of deciduous forest and 59.89% of thorn forest, dogs strongly selected for

human modified habitats over other land-use types.

This supported the results of the integrated step-selection analysis, which evaluated the fine-scale

habitat selection of dogs from the study area. The outcome of iSSA was similar for both time

scales (15 min, 30 min), with a higher standard error for the data resampled at 30 minutes (Fig.

5); therefore, we used the results of the 15 min resampled data for further analysis. Of all

available habitat classes, anthropogenic habitats (that is, settlements and agricultural land) were

significantly preferred, while others were not. However, the habitat-movement parameter

interaction was significant for multiple habitats. The observed step-lengths were significantly

longer than random steps in the forest habitat ($\beta = 0.0092$, p =0.043), whereas they were non-

significant in agriculture ($\beta = 0.0077$, p =0.077) and fallow/scrub habitat ($\beta = 0.0079$, p =0.075).

Overall, while the only significant habitat preference was for human-modified habitat classes, the

movement speed in each class was substantially different, with FRD moving with the longest

step lengths of approximately 85m-90m in the forested areas (traversing longer distances

between GPS fixes), followed by fallow/scrubland and agricultural fields, where the step lengths

were closer to 70m-75m (Fig. 6). The slowest movement with the shortest step length of 45-50m

was recorded in settlements, and there was no significant difference when compared to random

steps. Between villages, the movement trend was similar in both, but the movement speed was

marginally higher in Moyar across all habitat classes (Fig. 6).

394
Figure 5: Coefficient estimates with 95% confidence interval from the integrated step-selection

analysis for the two different temporal scales. The orange dots and lines represent the model

where the temporal scale for consecutive fixes was resampled at 15 minutes, and the blue dots

and lines represent the model where the temporal scale for consecutive fixes was resampled at 30

minutes.

Figure 6: Estimates of mean step length (speed) for movement estimated for different habitats

and different villages derived using the integrated step-selection function. The black bars

indicate the 95% confidence interval for each of the habitat classes.

Discussion

Our study investigates the movement ecology of owned free-ranging dogs in Mudumalai Tiger

Reserve, India. We found that GPS-tracked FRD primarily selected for anthropogenic habitats,

with an overwhelming preference for settlement areas, followed by agricultural land and fallow

or scrubland at village boundaries. Overall, FRD utilized human-modified habitats more than

96% of the time and the area within 500 m of their owners' homes more than 98% of the time.

This corroborates the well-documented phenomenon of dogs being closely associated with

human-modified habitats in rural areas [31,52–54], and usually staying within a few hundred

meters of their owner's home [17,21,54]. Indeed, almost all dog movement in forested land could

be attributed to only four FRD of the 15 that were tracked. Additionally, other movement metrics

were low compared to estimates from other dog telemetry studies. For instance, the median

activity range of 6.09 ha was much lower than median activity ranges from studies in Chad (54-

34 ha) [55], and Chile (19.20, 15.8-24.4) [17,56] (see also Table S2, supplementary material),

while the maximum distance from home of 1197 m was fairly short compared to ranging

distances recorded in Cambodia (up to 4km) [19] and Australia (8-30km) [14].

Given these limited movement patterns, the rural FRD in our study seem to fall into the class of

stay-at-home dogs, according to the three classes defined by Hudson et al. (2017), unlike

roamers or *explorers*, both of which travel to areas distant from owners' home with notable

frequency. As the FRD in MTR displayed minimal roaming and consistently used the area

immediately surrounding their owners' homes, they are likely to have the majority of their

requirements fulfilled by the humans they depend on [13]. This was an expected finding, since

we collared owned FRD which were subsidized by their owners, and indicates that this class of

dogs is unlikely to pose a persistent threat to wildlife. However, although our results are relevant

to local dog management in MTR, as the vast majority of the dog population in villages on the

Sigur Plateau are owned and sterilized (SVK, unpub. data), we note that we have not investigated

the movement of feral or non-owned dogs, and cannot comment on their potential ranging into

forest areas or impact on wildlife.

Our study was also constrained by a few other limitations, mainly owner apprehensions resulting

in the unauthorized removal and subsequent loss of collars, and collar malfunction resulting in

data loss, which reduced our effective sample size from 26 dogs to 15 dogs and limited the

sampling period. While several telemetry studies have been conducted on dogs with shorter

sampling periods (see Table S2, supplementary material), we acknowledge that the sampling

period should ideally be longer to more thoroughly investigate dog behavior and distinguish

between short-term and long-term movement patterns; for instance, dog home ranges have been

found to vary significantly across seasons [55]. Additionally, since we were unaware of when

dogs accompanied humans, we could not differentiate between natural movement versus

movement that reflected human activities (such as working in agricultural fields). However, we

believe this is unlikely to have significantly impacted our results, as many owners reported that

their dogs did not always accompany them but roamed freely on their own. This is supported by

the kernel density activity pattern, which showed that dogs were generally more active than

humans between 0:00-6:00 and less active than humans between 15:00-19:00.

This activity pattern also indicates why we found no significant difference between day and night

mean distance from home. While we expected that dogs would move less at night to minimize

chances of encountering wildlife, as predator species are more active at village boundaries after

sunset, the activity pattern showed that FRD moved during daylight as well as nighttime, with a

peak between early morning hours before dawn until around noon. This pattern differs from the

bimodal (morning-evening) activity pattern displayed commonly displayed by dogs elsewhere

[57] but tallies with the dog activity pattern in areas with puma presence [58]. If the observed

pattern is due to the close proximity to wildlife-adjacent areas, FRD may have adapted to

predators in MTR being more active late in the night and decreasing their activity closer to dawn,

or they may may derive sufficient protection from living in settled areas to be active at night;

however, confirming either theory would require data on wild predator activity patterns. We also

found no difference in movement metrics with respect to sex, unlike in other studies, where

neutered males have been recorded as having significantly larger activity ranges than spayed

females [45]; this may also be due to the presence of predators, as they may constrain male FRD

to similar activity ranges and distance travelled from home as female FRD.

In contrast, the significant difference in intensity of use between villages indicates that FRD used

their activity ranges differently in each village. In Moyar, the human population is smaller and

village homes are divided into settlement patches separated by large stretches of agricultural

land, while the human population in Masinagudi is several times larger and agricultural patches

are fragmented by well-connected roads and densely clustered houses. Therefore, we interpret

the higher intensity of use by Moyar FRD, indicating that they travelled longer distances or more

frequently within activity ranges similar to those of Masinagudi FRD, as dogs having to move

between safe, well-known areas inside their activity ranges in search of refuge from wildlife

encounters. Searching for resources or mates is unlikely to be a significant factor here as dogs

were fed at home by their owners and almost all dogs were sterilized. The higher intensity of use

by Moyar FRD is likely due to the heightened predation risk posed by wildlife, stemming from

two factors: the smaller human population with fragmented clusters of settlements, and the

larger, more porous perimeter with the forest and swathes of open agricultural land inside the

village that are conducive to wild animal movement. Indeed, several wild species, including

leopards, have previously been camera trapped in such land inside Moyar (SVK, unpub. data). In

Masinagudi, on the other hand, the larger human population, smaller village-forest boundary and

increased anthropogenic infrastructure potentially provide more relative safety to FRD

throughout their activity ranges, allowing FRD to carry out their daily activities with less total

movement. That the FRD of Moyar and Masinagudi adopted different roaming behaviors, even

when the two study sites were situated within 5km of each other, indicates that FRD can display

a high degree of behavioral plasticity depending on fine-scale ecological context.

The results of the integrated step selection analysis, namely the significant difference in step

length between in different habitats, also support the hypothesis that FRD rely on the human

shield and perceive higher risk in forest land. Forest areas were traversed almost two times faster

than settlements, and about 1.2 times faster agricultural or scrubland areas. This indicates that

forest areas were used in a more utilitarian manner (perhaps for travel between points) while

settlements were used for rest or play. Movement in agricultural and scrubland areas may have

been exploratory forays or may have been while accompanying owners. Additionally, the higher

movement speed across all habitat classes in the less populated and more fragmented village of

Moyar, while not statistically significant, corroborates our interpretation of the difference in

intensity of use between villages as being due to the difference in potential for wildlife

encounters. It also suggests that average dog movement speed could potentially be inversely

proportional to the settlement level in areas where FRD are preyed on by wildlife; that is, dogs

may move slower in more built-up areas and faster in less built-up areas in response the level of

perceived threat, although confirming this would require studying FRD across more villages with

a larger range of settlement levels.

Overall, we recommend that studies documenting patterns of dog movement should be

undertaken before designing and implementing dog management strategies that aim to modify

roaming behavior, as they may uncover similar unexpected nuances based on human-mediated

factors. For such studies, we suggest collaring more broadly across dog demographics (namely

ownership, sex, age and sterilization status) as well as across human and village related

parameters (e.g., length of the village-forest perimeter, household density, agricultural cover) at

larger spatial and temporal scales. In particular, studies from non-protected areas that support

wildlife may not only be easier to carry out but may provide much-needed insight into the dog-

wildlife interface in areas with no restrictions on human movement and land use.

Conclusion

Using movement metrics and the robust framework of integrated step-selection analysis to model

dog habitat selection, we compared free-ranging dog movement between two sites at a fine

geographical scale, interpreted through the lens of the human shield hypothesis [49]. Our

findings, while not generalizable across all dog-wildlife interfaces and limited by a relatively

short sampling period, contribute to research on dog movement and space use by documenting

metrics such as the activity range, and showing that FRD display high behavioral plasticity, as

movement patterns can differ between adjacent villages based on the level of protection afforded

by human settlements, putatively to avoid wild predators. They also show that owned,

provisioned and sterilized free-ranging dogs from villages in MTR spend the vast majority of

their time inside human settlement areas, making their potential impact on wildlife relatively low

compared to other dog classes, such as feral dogs. Differential access to beneficial resources – in

this case, refuge from wildlife encounters — provided by the intensity of human land use should

be taken into account when managing dog movement and behavior in wildlife adjacent areas,

given their highly commensal relationship with humans.

532
533

Declarations

Ethics approval and consent to participate

All GPS collars were fitted with the owners' consent. Collars weighed less than 250 g and were

within the required weight limit of < 5% of the dog's bodyweight, causing no physical distress to

dogs during the duration of deployment.

Consent for publication

Not applicable

Availability of data and materials

The dataset generated during this study is available in the Movebank repository under the study

name "Free-ranging dogs - vksanj - Mudumalai TR - GPS collaring", study ID 5328251712, at

the following link -

https://www.movebank.org/cms/webapp?gwt_fragment=page=studies,path=study5328251712

Codes for data analysis are available on Github at the following link -

https://github.com/nilanjanchatterjee/Dog-population-and-movement-

Analysis/blob/main/02_Movement_analysis.html

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by The Habitats Trust (Seed Grant, March 2024) in a grant provided to

SVK.

Authors' contributions

SVK acquired funding for the study, conducted fieldwork, participated in data analysis, and

drafted the manuscript. NC led data analysis and contributed to manuscript writing. RK provided

logistic and supervisory support, and contributed to manuscript writing. All authors read and

approved the final manuscript.

Acknowledgements

We acknowledge and appreciate the support provided by staff at the Indian Institute of Science

field station in Masinagudi, namely Mr. Sharan B. Gowda, Mr. Albert Suresh, Mr. Bharanaiah,

Mr. Arjunan, and Dr. Suresh H. S. We are grateful to Mr. Arasu for field support, as well as the

dog owners of Masinagudi and Moyar for permission to collar their dogs. We thank two

anonymous reviewers for their comments that helped improve this manuscript.

References

1. Gompper ME. The dog-human-wildlife interface: assessing the scope of the problem. Free-Ranging Dogs Wildl Conserv. 1st ed. OUP Oxford; 2014. p. 9–54.

2. Ortolani A, Vernooij H, Coppinger R. Ethiopian village dogs: Behavioural responses to a stranger's approach. Appl Anim Behav Sci. 2009;119:210–8.

3. Acosta-Jamett G, Cleaveland S, Cunningham AA, Bronsvoort BM deC. Demography of domestic dogs in rural and urban areas of the Coquimbo region of Chile and implications for disease transmission. Prev Vet Med. 2010;94:272–81.

4. Butler JRA, Bingham J. Demography and dog-human relationships of the dog population in Zimbabwean communal lands. Vet Rec. 2000;147:442–6.

5. Fiorello CV, Noss AJ, Deem SL. Demography, Hunting Ecology, and Pathogen Exposure of Domestic Dogs in the Isoso of Bolivia. Conserv Biol. 2006;20:762–71.

6. Kitala P, McDermott J, Kyule M, Gathuma J, Perry B, Wandeler A. Dog ecology and demography information to support the planning of rabies control in Machakos District, Kenya. Acta Trop. 2001;78:217–30.

7. Twardek WM, Peiman KS, Gallagher AJ, Cooke SJ. Fido, Fluffy, and wildlife conservation: The environmental consequences of domesticated animals. Environ Rev. 2017;25:381–95.

8. Young JK, Olson KA, Reading RP, Amgalanbaatar S, Berger J. Is Wildlife Going to the Dogs? Impacts of Feral and Free-roaming Dogs on Wildlife Populations. BioScience. 2011;61:125–32.

9. Gompper ME. Adding Nuance to Our Understanding of Dog–Wildlife Interactions and the Need for Management. Integr Comp Biol. 2021;61:93–102.

10. Doherty TS, Dickman CR, Glen AS, Newsome TM, Nimmo DG, Ritchie EG, et al. The global impacts of domestic dogs on threatened vertebrates. Biol Conserv. 2017;210:56–9.

11. Dürr S, Ward MP. Roaming behaviour and home range estimation of domestic dogs in Aboriginal and Torres Strait Islander communities in northern Australia using four different methods. Prev Vet Med. 2014;117:340–57.

12. Dürr S, Dhand NK, Bombara C, Molloy S, Ward MP. What influences the home range size of free-roaming domestic dogs? Epidemiol Infect. 2017;145:1339–50.

13. Hudson EG, Brookes VJ, Dürr S, Ward MP. Domestic dog roaming patterns in remote northern Australian indigenous communities and implications for disease modelling. Prev Vet Med. 2017;146:52–60.

14. Meek PD. The movement, roaming behaviour and home range of free-roaming domestic dogs, Canis lupus familiaris, in coastal New South Wales. Wildl Res. 1999;26:847–55.

15. Molloy S, Burleigh A, Dürr S, Ward M. Roaming behaviour of dogs in four remote Aboriginal communities in the Northern Territory, Australia: preliminary investigations. Aust Vet J. 2017;95:55–63.

16. Sparkes J, Körtner G, Ballard G, Fleming PJS. Spatial and temporal activity patterns of owned, free-
roaming dogs in coastal eastern Australia. Prev Vet Med. 2022;204:105641.

17. Schüttler E, Saavedra-Aracena L, Jiménez JE. Spatial and temporal plasticity in free-ranging dogs in
sub-Antarctic Chile. Appl Anim Behav Sci. 2022;250:105610.

18. Muinde P, Bettridge JM, Sousa FM, Dürr S, Dohoo IR, Berezowski J, et al. Who let the dogs out?
Exploring the spatial ecology of free-roaming domestic dogs in western Kenya. Ecol Evol.
2021;11:4218–31.

19. Ladd R, Meek P, Eames JC, Leung LK-P. Activity range and patterns of free-roaming village dogs in
a rural Cambodian village. Wildl Res. 2023;51.

20. Pérez GE, Conte A, Garde EJ, Messori S, Vanderstichel R, Serpell J. Movement and home range of
owned free-roaming male dogs in Puerto Natales, Chile. Appl Anim Behav Sci. 2018;205:74–82.

21. Ruiz-Izaguirre E, van Woersem A, Eilers K (C.) HAM, van Wieren SE, Bosch G, van der Zijpp AJ,
et al. Roaming characteristics and feeding practices of village dogs scavenging sea-turtle nests. Anim
Conserv. 2015;18:146–56.

22. Gaynor KM, Wooster EIF, Martinig AR, Green JR, Chhen A, Cuadros S, et al. The Human Shield
Hypothesis: Does Predator Avoidance of Humans Create Refuges for Prey? Ecol Lett. 2025;28:e70138.

23. Butler JRA, Linnell JDC, Athreya V, Lescureux N, McKeown A. Dog eat dog, cat eat dog: social-
ecological dimensions of dog predation by wild carnivores. Gompper ME Ed Free-Ranging Dogs Wildl
Conserv. Oxford, UK: Oxford University Press.; 2014. p. 117–43.

24. Belsare A, Vanak AT. Modelling the challenges of managing free-ranging dog populations. Sci Rep.
2020;10:18874.

25. Reddy CS, Sreelekshmi S, Jha CS, Dadhwal VK. National assessment of forest fragmentation in
India: Landscape indices as measures of the effects of fragmentation and forest cover change. Ecol Eng.
2013;60:453–64.

26. Venkataraman K, Sivaperuman C. Biodiversity Hotspots in India. In: Sivaperuman C, Venkataraman
K, editors. Indian Hotspots Vertebr Faunal Divers Conserv Manag Vol 1. Singapore: Springer; 2018. p.
1–27.

27. Belsare AV, Vanak AT, Gompper ME. Epidemiology of Viral Pathogens of Free-Ranging Dogs and
Indian Foxes in a Human-Dominated Landscape in Central India. Transbound Emerg Dis. 2014;61:78–
86.

28. Vanak AT, Thaker M, Gompper ME. Experimental examination of behavioural interactions between
free-ranging wild and domestic canids. Behav Ecol Sociobiol. 2009;64:279–87.

29. Mahar N, Habib B, Hussain SA. Do we need to unfriend a few friends? Free-ranging dogs affect
wildlife and pastoralists in the Indian Trans-Himalaya. Anim Conserv. 2024;27:53-64.

30. Home C, Bhatnagar YV, Vanak AT. Canine Conundrum: domestic dogs as an invasive species and
their impacts on wildlife in India. Anim Conserv. 2018;21:275–82.

31. Vanak AT, Gompper ME. Interference competition at the landscape level: the effect of free-ranging
dogs on a native mesocarnivore. J Appl Ecol. 2010;47:1225–32.

32. Tiwari HK, Bruce M, O'Dea M, Robertson ID. Utilising Group-Size and Home-Range Characteristics
of Free-Roaming Dogs (FRD) to Guide Mass Vaccination Campaigns against Rabies in India. Vaccines.
2019;7:136.

33. Rowcliffe JM, Jansen PA, Kays R, Kranstauber B, Carbone C. Wildlife speed cameras: measuring
animal travel speed and day range using camera traps. Remote Sens Ecol Conserv. 2016;2:84–94.

34. Hebblewhite M, Haydon DT. Distinguishing technology from biology: a critical review of the use of
GPS telemetry data in ecology. Philos Trans R Soc B Biol Sci. 2010;365:2303–12.

35. Suresh HS, Dattaraja HS, Sukumar R. Tree flora of Mudumalai Sanctuary, Tamil Nadu, Southern
India. Indian For. 1996;122:507–19.

36. Van Kesteren F, Mastin A, Mytynova B, Ziadinov I, Boufana B, Torgerson PR, et al. Dog ownership,
dog behaviour and transmission of Echinococcus spp. in the Alay Valley, southern Kyrgyzstan.
Parasitology. 2013;140:1674–84.

37. Aanensen DM, Huntley DM, Menegazzo M, Powell CI, Spratt BG. EpiCollect+: linking smartphones
to web applications for complex data collection projects. F1000Research. 2014;3:199.

38. Kays R, Davidson SC, Berger M, Bohrer G, Fiedler W, Flack A, et al. The Movebank system for
studying global animal movement and demography. Methods Ecol Evol. 2022;13:419–31.

39. Fleming CH, Calabrese JM. ctmm: Continuous-Time Movement Modeling [Internet]. 2023 [cited
2025 Mar 8]. Available from: https://github.com/ctmm-initiative/ctmm

40. R Core Team. R: a Language and Environment for Statistical Computing. [Internet]. Vienna, Austria.:
R Foundation for Statistical Computing; 2021. Available from: https://www.R-project.org/

41. Almeida PJAL, Vieira MV, Kajin M, Forero-Medina G, Cerqueira R. Indices of movement
behaviour: conceptual background, effects of scale and location errors. Zool Curitiba. 2010;27:674–80.

42. Loretto D, Vieira MV. The Effects of Reproductive and Climatic Seasons on Movements in the
Black-Eared Opossum (Didelphis aurita Wied-Neuwied, 1826). J Mammal. 2005;86:287–93.

43. Signer J, Fieberg J, Avgar T. Animal movement tools (amt): R package for managing tracking data
and conducting habitat selection analyses. Ecol Evol. 2019;9:880–90.

44. Pebesma E. Simple Features for R: Standardized Support for Spatial Vector Data. R J. 2018;10:439–
46.

45. Sparkes J, Körtner G, Ballard G, Fleming PJS, Brown WY. Effects of Sex and Reproductive State on
Interactions between Free-Roaming Domestic Dogs. PLOS ONE. 2014;9:e116053.

46. Lashley MA, Cove MV, Chitwood MC, Penido G, Gardner B, DePerno CS, et al. Estimating wildlife
activity curves: comparison of methods and sample size. Sci Rep. 2018;8:4173.

47. Avgar T, Potts JR, Lewis MA, Boyce MS. Integrated step selection analysis: bridging the gap
between resource selection and animal movement. Methods Ecol Evol. 2016;7:619–30.

48. Muff S, Signer J, Fieberg J. Accounting for individual-specific variation in habitat-selection studies:
Efficient estimation of mixed-effects models using Bayesian or frequentist computation. J Anim Ecol.
2020;89:80–92.

49. Chatterjee N, Wolfson D, Kim D, Velez J, Freeman S, Bacheler NM, et al. Modelling individual
variability in habitat selection and movement using integrated step-selection analysis. Methods Ecol Evol.
2024;15:1034–47.

50. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al. glmmTMB
Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.
R J. 2017;9:378–400.

51. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York; 2016.

52. Silva-Rodríguez EA, Ortega-Solís GR, Jiménez JE. Conservation and ecological implications of the
use of space by chilla foxes and free-ranging dogs in a human-dominated landscape in southern Chile.
Austral Ecol. 2010;35:765–77.

53. Cunha Silva L, Friker B, Warembourg C, Kanankege K, Wera E, Berger-González M, et al. Habitat
selection by free-roaming domestic dogs in rabies endemic countries in rural and urban settings. Sci Rep.
2022;12:20928.

54. Sepúlveda M, Pelican K, Cross P, Eguren A, Singer R. Fine-scale movements of rural free-ranging
dogs in conservation areas in the temperate rainforest of the coastal range of southern Chile. Mamm Biol.
2015;80:290–7.

55. Wilson-Aggarwal JK, Goodwin CED, Moundai T, Sidouin MK, Swan GJF, Léchenne M, et al.
Spatial and temporal dynamics of space use by free-ranging domestic dogs Canis familiaris in rural
Africa. Ecol Appl. 2021;31:e02328.

56. Saavedra-Aracena L, Grimm-Seyfarth A, Schüttler E. Do dog-human bonds influence movements of
free-ranging dogs in wilderness? Appl Anim Behav Sci. 2021;241:105358.

57. Griss S, Riemer S, Warembourg C, Sousa FM, Wera E, Berger-Gonzalez M, et al. If they could
choose: How would dogs spend their days? Activity patterns in four populations of domestic dogs. Appl
Anim Behav Sci. 2021;243:105449.

58. de Cassia Bianchi R, Olifiers N, Riski LL, Gouvea JA, Cesário CS, Fornitano L, et al. Dog activity in
protected areas: behavioral effects on mesocarnivores and the impacts of a top predator. Eur J Wildl Res.
2020;66:36.

Additional files

Additional file 1, format: PDF

Data:

Table S1: Number of GPS points for each individual dog in each habitat type.

Table S2: Home/activity range comparisons from global studies — expanded table adopted from

Ladd et al., 2023, with home/activity range estimates and sampling durations from dog telemetry

studies across countries.

Table S3: ΔAIC comparison across activity range models — ΔAIC values for all trialed models

during activity range estimation.

Figure S1. Time plots for GPS locations in the forest, illustrating individual temporal variation in

dogs' use of forest land.

Figure S2. Compilation of photos of collared dogs.

Additional file 2: Mapped kernel density estimates for all dogs. Please note that this includes

data from all 22 dogs uploaded to Movebank, and dogs are referred to by names and not ID

numbers.