Optimized chemogenetic ablation and regeneration of enteric nervous system neurons in

zebrafish

Mujahid Ali Shah[1], Katherine K. Moran[1], Helen Rueckert[1,2], Abigail V. Sharrock[3], David F.

Ackerley[3], Jeff S. Mumm[4], Julia Ganz[1] *

[1]Department of Integrative Biology, Michigan State University, East Lansing, MI, USA,

[2]Department of Cell Biology, Developmental and Stem Cell Biology Program, Duke University

School of Medicine, Durham, NC, USA, [3]School of Biological Sciences, Victoria University of

Wellington, Wellington, New Zealand, [4] Department of Ophthalmology, Wilmer Eye Institute,

Johns Hopkins University, Baltimore, MD, USA.

*Correspondence, ganz@msu.edu

Abstract

The enteric nervous system (ENS) is the intrinsic nervous system of the gut and regulates

essential gut functions, including motility, digestion, and immune response, ensuring gut

homeostasis. ENS dysfunction or loss is associated with gastrointestinal disorders such as

Hirschsprung disease (HSCR). Currently, surgery is the only treatment for HSCR, but it often

has lifelong, severe complications. Restoring missing ENS neurons by stimulating endogenous

neuronal regeneration presents a promising therapeutic approach for ENS disease. To reveal

the cellular-molecular mechanisms regulating neuronal regeneration we study a species

capable of robust ENS restoration, the zebrafish. For this, we developed a chemogenetic

ablation model in zebrafish using the Gal4/UAS NTR 2.0 system for targeted ENS neuron

ablation. Spatially and temporally controlled neuronal death was confirmed by morphological

changes, quantification of neuronal loss, and TUNEL assays. We observed an acute immune

response that normalizes at 1 day of treatment. Quantification of regenerated neurons

demonstrated complete restoration of ENS neuron numbers to control levels by 9 days post

treatment, with recovery of gut motility. Among the regenerated neurons, nitrergic, cholinergic

and VIPergic subtypes showed full recovery, whereas serotonergic neurons only displayed

partial recovery, indicating subtype-specific differences in regenerative capacity and/or timing of

cell replacement. Our study establishes a robust platform for dissecting the cellular-molecular

mechanisms of ENS regeneration to develop potential treatment approaches for ENS-related

diseases.

Keywords: Enteric neurons, chemogenetic ablation, functional recovery, gut motility, ENS

diseases, immune response

Introduction

The enteric nervous system (ENS), the largest subdivision of the peripheral nervous system, consists of an intricate network of neurons and glial cells derived from neural crest progenitors that migrate into the developing gastrointestinal tract (Ganz, 2018; Hao & Young, 2009; Taylor et al., 2016). This complex network plays a pivotal role in regulating essential gastrointestinal functions such as gut motility, nutrient absorption, epithelial secretion, and blood flow (Kuil et al., 2021; Kunze & Furness, 1999; Niesler et al., 2021). Dysfunction or loss of the ENS has been linked to various gastrointestinal disorders such as Hirschsprung disease (HSCR), achalasia, and autism spectrum disorder (Niesler et al., 2021). HSCR, which is characterized by variable degrees ENS neuron loss in the distal gut. HSCR one of the most common congenital anomalies of the ENS, leading to severe intestinal obstruction in neonates, with an estimated incidence of 1 in 5,000 live births (Butler Tjaden & Trainor, 2013). To date, ENS disorders can only be treated symptomatically or by surgical removal of the area with ENS neuron deficits. The current standard treatment for HSCR is surgical resection of the affected gut segment. However, patient outcomes vary widely, and most continue to experience lifelong gastrointestinal defects, highlighting the need for alternative treatments. One of the therapeutic approaches for HSCR is the stimulation of resident stem cells to regenerate missing ENS cells with promising results but varying efficacy depending on the age and the specific gut region (Burns et al., 2016; Joseph et al., 2011; Soret et al., 2020). The successful translation of these approaches requires a comprehensive understanding of the cellular and molecular mechanisms that underlie effective ENS regeneration. Elucidating these mechanisms necessitates the development of a reliable animal model that exhibits robust and reproducible regeneration of ENS neurons following injury.

So far, only a few studies - mostly in mammalian models - have investigated ENS regeneration using chemical, mechanical, or chemical-genetic (chemogenetic) injury. Although these approaches have provided valuable insights, mammalian models show limited and incomplete neuronal regeneration in the ENS (Rueckert & Ganz, 2022; Stavely et al., 2024). In contrast, the zebrafish (*Danio rerio*), a non-mammalian animal model system, is a robust model for studying organ regeneration including various parts of the nervous system such as the brain (Kizil et al., 2012) retina (Hanovice et al., 2019), ENS (El-Nachef & Bronner, 2020; Ohno et al., 2021), but also other organs, such as kidney (Diep et al., 2011), heart (Cao & Poss, 2018), and fin (Romero et al., 2018). Zebrafish also offer numerous experimental advantages, including rapid development, large numbers of offspring, optical transparency that enables live imaging, and a high degree of genetic homology with humans (Howe et al., 2013). Despite these experimental advantages (Kuil et al., 2021; Ganz, 2018), so far, only two studies have studied ENS regeneration in zebrafish using a

laser ablation approach (El-Nachef & Bronner, 2020; Ohno et al., 2021). Those studies provided the

first evidence that the ENS can regenerate in zebrafish following focal neuronal loss. In particular,

Ohno et al., showed that neural crest-derived cells migrate into the ablated region, proliferate, and

give rise to new neurons, partially restoring ENS neuron numbers. However, several important

issues remain to demonstrate that the ENS fully regenerates in zebrafish: First, the number of

regenerated neurons did not reach control levels in the studied time frame. Second, it remains to be

shown whether different ENS neuronal subtypes are able to regenerate. Third, it remains unclear

whether the regenerated neurons functionally integrated into existing ENS circuits to re-establish

normal gut function. In addition, focal laser ablation has several technical limitations, including

collateral damage to surrounding tissues, labor-intensive protocols, and restricted ablation of just a

few cells within a small gut region, hindering downstream analysis for functional recovery

assessment (Ohno et al., 2021).

To overcome these challenges, we leveraged an established approach for spatiotemporally

controlled cell ablation: the chemogenetic bacterial nitroreductase (NTR) system. This method

utilizes the NTR enzyme, which converts prodrugs, e.g., Metronidazole (MTZ), into cytotoxic agents,

thereby selectively killing cells that express NTR (Medico et al., 2001). In zebrafish, this

chemogenetic method has been successfully applied in regenerative studies by ablating specific cell

populations in heart, pancreas, liver, retina, and other organs (Ariga et al., 2010; Curado et al.,

2007, Sharrock et al., 2022; White et al., 2017). However, this method has not yet been tested for

the ablation of ENS neurons.

In this study, we took advantage of the precise spatial-temporal control of the *Gal4/UAS* system

(Asakawa et al., 2008; Zhang et al., 2019) for the chemogenetic ablation of ENS neurons in

postembryonic zebrafish larvae (**Figure 1A**). We additionally employed the most advanced iteration

of the nitroreductase technology, NTR 2.0, which provides effective ablation of target cells at ~10- to

100-fold lower concentrations of MTZ than the original NTR system (Sharrock et al., 2022). Using

this approach, we establish a robust experimental paradigm of ENS neuron regeneration by ablating

only NTR-expressing neurons through MTZ treatment. Morphological and TUNEL analysis

confirmed that the ablated cells underwent cell death with a concomitant acute immune response

that normalizes within 1 day post treatment (dpt). By 9 dpt, neuronal recovery was complete, with full

recovery of most neuronal subtypes, including nitrergic, cholinergic and VIPergic neurons. Our

results demonstrate, for the first time, functional recovery of ENS neuron numbers including

restoration of gut motility. These findings pave the way for understanding the cellular and molecular

mechanisms of ENS regeneration, a first step for developing new therapeutic approaches for
gastrointestinal disorders associated with ENS dysfunction in the future.

Material and methods
Animals
All experiments were carried out in accordance with animal welfare laws, guidelines, and
policies and were approved by the Michigan State University Institutional Animal Care and Use
Committee. Zebrafish lines were maintained in a laboratory breeding colony according to
established protocols (Westerfield, 2000). Adult zebrafish were bred naturally in system water,
and fertilized eggs were collected and transferred to 100 mm petri dishes containing embryo
medium. Embryos were allowed to develop at 28.5°C and staged by days post fertilization (dpf)
as described earlier (Kimmel et al., 1995; Webb et al., 2009). The zebrafish lines utilized in this
study are listed in **Table 1**. Double transgenic larvae were established to co-express YFP and
NTR 2.0 in the ENS neurons. For this, the *Tg(5xUAS:GAP-TagYFP-2A-NTR2.0)jh513Tg*
reporter/effector line was maintained together with the *Et(2xNRSE-cfos KALT4)gmc617;*
Tg(5xUAS:GAP-ECFP,he1.1:GAP-ECFP)gmc1913 transgenic fish line. The
Et(2xNRSE:cfos:KalTA4)gmc617 line expresses KalTA4, a zebrafish-optimized Gal4/VP16
fusion protein with neuronal specificity (Xie et al., 2012; Sharrock et al., 2022). We confirmed
that the *Tg(SAGFF(LF)217B)* drives reporter expression in ENS neurons (described in the
results section), as an incross of *Tg(5xUAS:GAP-TagYFP-2A-NTR2.0)*; *Et(2xNRSE-cfos*
KALT4)gmc617; Tg(5xUAS:GAP-ECFP,he1.1:GAP-ECFP)gmc1913 did not show YFP
expression in ENS neurons. The *Tg(5xUAS:GAP-ECFP,he1.1:GAP-ECFP)gmc1913* line
expresses cyan fluorescent protein (CFP), our analysis was focused on YFP expression driven
by the Gal4/UAS system. Genotyping of Gal4 driver lines was performed with the following
primers for Gal4 (GAL4F: 5'-CGCTACTCTCCCAAAACCAAAAGG-3'; GAL4R: 5'-
TCTCTTCCGATGATGATGTCGCAC -3' using the PCR protocol: 95° x 3 min - [95° x 30s - 60° x
30s - 72° x 30s] x 35 - 72° x 5' 12° ∞.

Table 1: Transgenic lines used in this study

Full name	Abbreviation	Source
Tg(SAGFF(LF)217B)	*217B:Gal4*	Asakawa et al., 2008
Tg(SAGFF(LF)234A)	*234A:Gal4*	Asakawa et al., 2008
Tg(gSAIzGFFM495A)	*495A:Gal4*	Asakawa et al., 2008
Tg(5xUAS:EGFP)nkuasgfp	*UAS:GFP*	Kawakami et al., 2010
Tg(UAS-E1b:NfsB-mCherry)c264	*UAS:NTR-mCherry*	Davison et al., 2007

Full name	Abbreviation	Source
Tg(5xUAS:GAP-TagYFP-2A-NTR2.0)jh513Tg; Et(2xNRSE-cfos KALT4)gmc617; Tg(5xUAS:GAP-ECFP, he1.1:GAP-ECFP)gmc1913	*UAS:YFP-NTR 2.0**	Sharrock et al., 2022; Xie et al., 2012

129   *The *Tg(5xUAS:GAP-TagYFP-2A-NTR2.0)jh513Tg* fish were maintained together with the *Et(2xNRSE-cfos*
130   *KALT4)gmc617; Tg(5xUAS:GAP-ECFP,he1.1:GAP-ECFP)gmc1913* lines.

Chemogenetic ablation using Metronidazole

The *217B:Gal4* driver line was crossed with the *UAS:YFP-NTR 2.0* responder line. Offspring from this cross were sorted based on YFP+ cells in the gut into YFP+ and YFP− groups at 5 dpf using a Zeiss Axio ZoomV16 fluorescent stereo microscope. The sorted larvae were then treated at 5 dpf or for the later time points transferred to the nursery until the day of the assay. Prior to MTZ treatment, larvae - except for the 5 dpf groups - were starved for 24 hours to clear the gut of autofluorescent matter. Approximately 50 larvae were transferred to a petri dish (100 mm × 15 mm) containing 40 mL of 400 µM MTZ (M1547 Sigma Aldrich) (Sharrock et al., 2022) in zebrafish system water (diluted from a 5 mM stock solution). The following experimental groups were used: 1) YFP+ MTZ (YFP+ larvae treated with MTZ; 2) YFP+ control (YFP+ larvae, not treated with MTZ); and 3) YFP- MTZ (YFP- larvae, treated with MTZ). Larvae were incubated in the dark for 24 hours to ensure maximal drug exposure as MTZ is light sensitive. After 24 hours, the effects of MTZ treatment on YFP+ cell ablation were assessed using a Zeiss Axio Zoom.V16 fluorescent stereo microscope. After confirmation of cell loss, the MTZ solution was discarded and larvae were returned to the nursery, placed in 2.8-liter tanks (~40 larvae per tank), and raised until the 21 dpt / 33 dpf.

ENS neuron regeneration analysis

The regeneration of MTZ-treated YFP+ cells in the gut was assessed at 12 dpf (0 dpt, before ablation) and subsequently at three-day intervals from 13 dpf (1 dpt) to 27 dpf (15 dpt) and compared to YFP+ untreated controls. At each regeneration time point, both control and treated larvae were checked briefly using a Zeiss Axio Zoom.V16 fluorescent stereo microscope, then fixed and immunoassayed (as described below) by confocal microscopy imaging for data acquisition.

Sample fixation

To ensure optimal imaging conditions, larvae were fasted for 24 hours prior to sampling at the required stages to allow for gut clearance of autofluorescent matter. Then, larvae were

euthanized with an overdose of Tricaine Methanesulfonate (MS-222). **For immunolabeling**,

euthanized samples were fixed in 4% paraformaldehyde (PFA) prepared in 1x sweet buffer (8%

sucrose, 0.2 M CaCl$_2$, 0.2 M PO$_4$ buffer, pH 7.3) for 2 hours at room temperature. Then, they

were washed subsequently 2 times in 0.5% Phosphate buffer saline-Triton™ X-100 (PBSTX) for

15 mins at room temperature, and stored in PBSTX at 4°C. **For TUNEL assay**, euthanized

samples were fixed overnight in 10% Natural buffer saline (NBF) at 4°C, then washed three

times in PBS for 5 minutes at room temperature and stored in PBS at 4°C. **For Hybridization**

Chain Reaction (HCR), euthanized larvae were rinsed thoroughly in RNAase free water, and

fixed in RNAase free 4% PFA in PBS at 4°C for ~16 hours, washed three times in PBS for 5

minutes at room temperature, dehydrated in a graded methanol series (50, 70, 90, 95 and 100

% methanol in nuclease free water), and stored in 100% methanol at −20°C for a minimum of

16 hours or up to six months (Ibarra-García-Padilla et al., 2021).

TUNEL assay

TUNEL assays were performed using the Click-iT™ Plus TUNEL Assay Kits (C10618, Thermo

Fisher Scientific) for In Situ Apoptosis Detection. For this, PBS was removed from fixed samples

and permeabilization was carried out using Proteinase K (20 µg/mL) for 50 minutes at room

temperature, followed by three PBS washes and a deionized water rinse. Then samples were

re-fixed in 10% NBF in PBS for 5 minutes at 37°C, followed by three additional PBS washes and

another deionized water rinse. To detect DNA strand breaks (i.e., for the positive control), the

larvae were incubated in 1 unit of DNase I (Cat. No. 18068-015) diluted into 1× DNase I

Reaction Buffer for 30 minutes at room temperature (**Figure S1**). After that, larvae were

preincubated in 150 µL of Terminal deoxynucleotidyl transferase (TdT) buffer for 10 minutes at

37°C, incubated in the 100 µL TdT reaction mixture {94 µL TdT Reaction Buffer, 2 µL 5-ethynyl-

2'-deoxyuridine-5'-triphosphate (EdUTP), and 4 µL TdT enzyme} for 30 minutes at 37°C, rinsed

in deionized water, washed twice in 3 % bovine serum albumin (BSA) in 0.1% PBS-Triton™ X-

100 for 5 minutes each, and one time in PBS. Then the larvae were incubated with the reaction

cocktail (90 µL TUNEL Supermix and 10 µL 10× Click-iT Plus TUNEL Reaction Buffer Additive)

for 30 minutes at 37°C in the dark, followed by two 3% BSA in PBS washes and two final PBS

washes. TUNEL staining was assessed using a Zeiss Axio Zoom.V16 fluorescence

stereomicroscope. Immunostaining was subsequently performed under dark conditions, as

described below, and larvae were stored in PBS until imaging.

Immunostaining

Whole-mount immunostaining was performed as previously described (Bandla et al., 2022;
Uyttebroek et al., 2010). The following primary antibodies: 5-HT (1:5,000, catalog number
20080), HuC/D (1:1,000, catalog number A-21271), mouse GFP (1:300, catalog number
A11120), rabbit GFP (1:1,000, catalog number A11122), nNOS (1:1,000, catalog number
GTX133407) and L-plastin, Lcp1 (1:100, catalog number GTX124420) were utilized to perform
immunostaining. The primary antibodies were detected using Alexa Fluor 488 and/or 568
secondary antibodies, followed by washes with 0.3% PBSTX. The staining was assessed using
Zeiss Axio Zoom.V16 fluorescent stereo microscope and samples were stored in 0.3% PBSTX
at 4°C until imaging.
203

Hybridization Chain Reaction
The (HCR) kit, probes, *vip* (Accession number: NM_001114553.3) and *slc18a3a* (Accession
number: NM_001077550.1), and hairpins were from Molecular Instruments, Inc.
(https://www.molecularinstruments.com) and the HCR protocol was performed as previously
described (Ibarra-García-Padilla et al., 2021) with minor modifications. The HCR protocol was
followed by immunostaining. The fixed samples stored in 100% methanol were rehydrated in
graded methanol series, permeabilized with pre-chilled acetone at −20°C for 25 min, treated
with 20 µg of Proteinase K for 1 h at room temperature, refixed in 4% PFA in PBS for 20 min,
and washed with PBS-Tween before hybridization. For HCR hybridization, samples were pre-
hybridized at 37°C, incubated with a probe mixture (4 µL of each 1 µM probe set stock) for 12–
16 hours, washed to remove excess probes, and amplified for 12–16 hours, followed by
sequential washes (Ibarra-García-Padilla et al., 2021). For antibody staining, samples were
blocked in 5% goat serum, incubated with primary anti-mouse GFP (1:300, catalog number
A11120) antibodies overnight at 4°C, washed extensively, and incubated with secondary
antibodies overnight (Ibarra-García-Padilla et al., 2021). Throughout the protocol, samples were
protected from light to prevent photobleaching, ensuring optimal visualization of hybridization
and staining results. Afterward, stained samples were stored in PBST at 4°C until imaging.
221

Imaging, data acquisition, and quantification
For analysis of colocalization and quantification, intestines were dissected with forceps and
mounted on glass slides in 0.3% PBSTX or Fluoromount-G® mounting media (Southern Biotech,
catalog number 0100-01) and covered with a coverslip. For the analysis of Gal4-based reporter
co-expression with the pan-neuronal marker HuC/D, mounted intestines were imaged on a
Zeiss Axio Imager microscope operated through the ZEN software to determine colocalization.

Confocal z-stacks images were captured using the 3i spinning disk confocal microscope with a
10, 20, or 60× objectives, operated through Slidebook6 software (Version 6.0.13). For cell
quantification, z-stacks were generated from confocal stacks in Slidebook6, which were used for
manual cell counts and colocalization analysis. For analysis within gut regions, the images of
the intestine were divided into three sections - anterior, middle, and posterior - as previously
described (Jianlong Li et al., 2020; Wallace et al., 2005). Cell counts were normalized to 200 μm
to account for varying lengths of each of gut regions between larvae and to 1000 μm for the
whole gut.
236

237 **Analysis of neuron morphology**
238 To analyze morphological changes in neurons after MTZ exposure, control and MTZ-treated
239 larvae were euthanized at the timepoint indicated, their intestines were dissected, mounted on
240 glass slide in PBS and imaged on the 3i spinning disk confocal microscope with a 60× objective,
241 operated through Slidebook6 software (Version 6.0.13).
242

Gut anterograde movements analysis
Larvae from control and treated groups were fasted 24 hours prior to analysis, anesthetized with
MS-222, and embedded in 0.8% low-melting-point agarose. Videos of the gut region from
midgut to posterior were recorded on the Zeiss Axio Zoom.V16 fluorescent stereo microscope
with ZEN Imaging Software (ZEN 2.3 blue edition). All time lapse videos were recorded using
the same conditions, including exposure time (206s), frame rate (100 frames / 206 sec), and
pixels (2752 x 2208). The total number of anterograde, peristaltic wave movements was
analyzed from the recorded videos. The Peristaltic Wave Interval (PWI) time was calculated as:
$PWI = T/N$ where PWI represents the average time interval (in seconds) between two
consecutive anterograde peristaltic movements. The total duration of each video (T, in seconds)
and the total number of peristaltic movements observed (N) were recorded.
254

Statistical Analysis
For multiple comparisons, one-way and two-way ANOVA were applied to normally distributed
data, while the non-parametric Kruskal–Wallis test was used for non-normally distributed data.
For pairwise comparisons, unpaired t-tests were used for normally distributed data, and the
Mann–Whitney test was used for non-normally distributed data. To evaluate temporal changes
across multiple time points in regeneration studies, multiple unpaired t-tests were performed
between MTZ-treated and control larvae. For fractional analysis of neuronal colocalization

patterns, descriptive statistics were applied. Fold-change analysis was calculated based on the

average number of cells in each group. All statistical analyses were performed using GraphPad

Prism 9.1 (GraphPad Software, San Diego, CA, USA). A significance threshold of $p < 0.05$ was

used to determine statistically significant differences.

Results

Identification of Gal4 driver line for specific ablation of ENS neurons

To identify a Gal4 driver line for maximizing the breadth and selectivity of ENS neuron ablation,

we crossed three Gal4 gene-trap driver lines (*495A:Gal4; 234A:Gal4; 217B:Gal4*) with a *UAS*

reporter line to analyze the extent and specificity of reporter gene expression in the ENS and

surrounding tissues. To determine if reporter gene expression within the ENS is primarily

restricted to neurons, thereby minimizing off-target ablation, we investigated reporter gene co-

expression with the pan-neuronal marker HuC/D in the ENS at 5 dpf (*495A:Gal4xUAS:GFP*,

234A:Gal4xUAS:NTR:DsRed) and at 5, 6 and 7 dpf (*217B:Gal4xUAS:GFP*) (**Figure 1**).

Our analysis revealed notable differences in the co-expression profiles among the driver lines.

The *495A:Gal4* line has previously been reported to show largely gut-restricted expression, but

which gut cells express the reporter had not been determined (Kawakami et al., 2010). We

found that the intestines of the *495A:Gal4* line displayed only about half HuC/D+/Reporter+ cells

with many reporter+ cells being HuC/D- and HuC/D+/Reporter- cells (**Figure 1B**). The

expression of *234A:Gal4* has been reported to be expressed in the ENS, but also broadly

outside the gut (Kawakami et al., 2010; Heanue et al., 2016). In contrast, the *217B:Gal4* had

been reported to have reporter expression in the gut with minimal expression outside the gut

compared to the *234A:Gal4* line. Consistent with these reports, we found that both *234A:Gal4*

and *217B:Gal4* driver lines exhibited higher number of neurons co-expressing HuC/D+ and

Reporter with only a small population of HuC/D-/Reporter+ and HuC/D+/Reporter- cells (**Figure**

1B-C).

Comparatively, both the *234A:Gal4* and *217B:Gal4* lines had sufficient Reporter+ cells in the

gut. However, *234A:Gal4* drives extensive reporter gene expression outside the ENS and in

ENS progenitor cells (Heanue et al., 2016), which would lead to ablation of additional, non-

targeted cells. Therefore, we chose the *217B:Gal4* line to maximize the extent and specificity of

ENS neuron ablation (**Figure 1C**).

Figure 1. Identification of a Gal4 driver line for efficient ENS neuron ablation using a chemogenetic system. (A) Pictorial representation of chemogenetic ablation of ENS neurons. A carrier of the Gal4 driver line, *Tg*(*Gal4)*, which drives expression in ENS neurons was crossed with a carrier of the responder/effector line *Tg(UAS:YFP-NTR 2.0)*, where the Gal4 protein binds to Upstream Activating Sequence (*UAS*) element, initiating expression of both *yellow fluorescence protein* (YFP, reporter protein) and *nitroreductase* (NTR 2.0). Left side: Schematic representation of the untreated control intestine, where enteric neurons co-expressing the *Gal4* and *UAS* transgenes also express both YFP and NTR 2.0 in ENS neurons (green). Right side: Schematic representation of intestine following treatment with 400 µM metronidazole (MTZ). The prodrug MTZ is enzymatically converted by NTR 2.0 into a cytotoxic compound, leading to the selective ablation of YFP-NTR 2.0-expressing enteric neurons. The bar graphs show colocalization percentage of the pan-neuronal marker HuC/D with Reporter (Rep) expression within the gut for Gal4 driver. **(B)** For the *234A:Gal4* line, most Rep+ cells are HuC/D+ at 5 dpf (81.3% HuC/D+/Rep+). The remaining cells consist of 12.7% HuC/D+/ Rep- and 6.0% HuC/D-/Rep+ (n = 12 larvae per group, 2 experiments). For the *495A:Gal4* line, only 55% cells are HuC/D+/Rep+ at 5 dpf, the remaining cells are 23.5% HuC/D+/ Rep- and 21.2% HuC/D-/Rep+ (n = 12 larvae per group, 2 experiments). **(C)** For the *217B:Gal4* line, there is a high degree of overlap between Rep and HuC/D expression [(85.3% (5 dpf), 85.7% (6 dpf), and 90.2% (7 dpf)]. The remaining cells consist of HuC/D+/ Rep- [14% (5 dpf) 12.3% (6 dpf) and 9.0% (7 dpf)] and HuC/D-/Rep+ [0.7% (5 dpf), 2.0% (6 dpf), and 0.8% (7 dpf)], n = 13 (5 dpf), 15 (6 dpf), and 14 (7 dpf) larvae per group, 2 experiments per time point. dpf = days post fertilization.

Establishing a robust chemogenetic ablation assay for ENS neuron
To establish a robust ENS neuron ablation assay, we used offspring from a cross between the
217B:Gal4 and *UAS:YFP-NTR 2.0* transgenic lines that co-express YFP and NTR 2.0 in cells in
the gut (YFP+). To ablate YFP-NTR 2.0 expressing cells, we treated YFP+ larvae with 400 µM
MTZ for 24 hours (Sharrock et al., 2022) starting at 5 or 12 dpf. To assess the extent of ENS
cell loss, we quantified YFP+ cells at 6 dpf and 13 dpf (1 dpt for MTZ-treated larvae). In
untreated controls, no significant difference in YFP+ cells was observed between 5 and 6 dpf, or
12 and 13 dpf. Conversely, MTZ-treated groups showed a near-complete loss of YFP+ cell
numbers at 1 dpt compared to 0 dpt (before treatment) and untreated control (**Figure 2A-C**).
As 24 h of MTZ treatment showed robust YFP+ cell ablation at both developmental stages, we
decided to move forward with MTZ treatment at 12 dpf for further analyses. Embryonic

neurogenesis is still very active at 5 dpf (Olden et al., 2008; Roy-Carson et al., 2017). Therefore,

ablating neurons at 5 dpf might include processes of developmental neurogenesis rather than of

neuronal regeneration, which we aimed to avoid.

To confirm that YFP+ cells are neurons at 12 dpf, we analyzed co-expression of YFP and

HuC/D in *217B:Gal4*; *UAS:YFP-NTR 2.0* fish. Quantification revealed that, on average, 44.4

YFP+ neurons were observed among a total of 204.6 HuC/D+ neurons. We found that 96.3% of

YFP+ cells were HuC/D+ indicating that the vast majority of YFP+ cells are ENS neurons while

only 3.7% YFP+ cells are HuC/D-. Therefore, we will hereafter refer to YFP+ cells in these lines

as YFP+ neurons (**Figure 2D-F**).

Figure 2. Establishment of a robust ENS neuron ablation assay. (A) The timeline of dpf
and dpt illustrates the experimental procedure for MTZ treatments initiated at either 5 or 12
dpf. **(B)** Whole-mount side views of live larval guts at the stage indicated. Left panel: YFP+

neurons (white dots) are present before treatment (0 dpt) at 5 or 12 dpf in YFP+ larvae. Right panel: after 24 hours treatment with 400 µM Metronidazole (MTZ), YFP+ cells are completely absent in MTZ-treated fish (YFP+ MTZ) but present in untreated controls (YFP+ control). **(C)** Quantification of YFP+ neurons in control and MTZ-treated groups. Data are shown as mean ± SD (n = 10 larvae per group, 2 experiments); statistical significance ****p < 0.0001; ns = non-significant. **(D)** At 12 dpf, YFP+ cells [(anti-GFP immunostaining (top, green)] and the pan-neuronal marker HuC/D+ cells (middle, magenta) highly colocalize (bottom, overlay). **(E)** Quantification of GFP+/HuC/D+, GFP+/HuC/D- and GFP-/HuC/D+ cells. Data are shown as mean (n= 10 larvae per group, 2 experiments). **(F)** Percentage of YFP+ cells that are HuC/D+ and HuC/D- cells in **E**. (D) Maximum projection confocal images of dissected intestine at 12 dpf; dpf = days post fertilization. dpf = days post fertilization; dpt = days post treatment; SD = standard deviation; Scale bar = 100 µm.

YFP+ neurons undergo DNA-damage induced cell death after MTZ treatment

To determine the timeline of neuronal ablation, we quantified number of YFP+ neurons at 6, 12 and 24 hpt. We find that YFP+ neurons are essentially absent at 24 hpt indicating that MTZ-induced ablation of YFP+ neurons is completed within 24 hours post-treatment (**Figure S2**). To test if YFP+ neurons undergo cell death after MTZ treatment, we evaluated changes in neuronal morphology and performed TUNEL assays to detect cells undergoing cell death. YFP+ neurons in untreated control larvae displayed normal morphology with no visible signs of morphological changes (**Figure 3A**). In contrast, YFP+ neurons in MTZ-treated samples exhibited significant morphological changes, including axonal and cell body fragmentation and retraction of neuronal processes, consistent with neurons undergoing cell death [**Figure 3**, (Cusack et al., 2013; Ham et al., 2010)]. Additionally, we conducted TUNEL assays combined with immunostaining for YFP to assess DNA damage in control and MTZ-treated fish 6 at hpt. As expected, we found GFP+/TUNEL+ cells in both treated and control groups since apoptotic cell death is a natural ongoing process at 12 dpf (Lee et al., 2023). However, significantly higher numbers of GFP+/TUNEL+ cells were observed in the MTZ-treated group compared to YFP+ untreated control (**Figure 3B, C**). In addition, GFP+/TUNEL+ neurons exhibited clear morphological changes indicating that YFP+ neurons were undergoing cell death (**Figure 3D**). These results indicate that MTZ treatment induces DNA damage which leads to cell death with pronounced structural alterations attending this process, consistent with prior studies using this system for selective cell death (Kulkarni et al., 2018; White & Mumm, 2013).

Figure 3. MTZ treatment induces DNA-damage cell death in YFP+ neurons. (A) In 12 dpf larvae, YFP+ neurons displays normal neuronal morphology in the untreated control group, while MTZ treated YFP+ neurons show fragmented and retracted neuronal processes at 1 hpt (white arrowheads; n=6 larvae per group, 2 experiments). **(B)** At 6 hpt, there is an increase in TUNEL staining (magenta) in YFP+ neurons (anti-GFP immunostaining) in MTZ treated larvae compared to untreated controls. In YFP+ control group, GFP+ neurons have normal morphology. In contrast, GFP+ neurons in the MTZ-treated group show fragmented morphology indicative of cell death. **(C)** Quantification of GFP+/TUNEL+ cells shows significant increase in YFP+ MTZ-treated larvae compared to YFP+ untreated control. Data are shown as mean ± SD (n= 9 larvae per group, 2

experiments); statistical significance ***$p < 0.001$. **(D)** Magnified images of GFP+/TUNEL+ cells (white arrowhead) in the YFP+ MTZ-treated group. **(A-D)** Maximum projection confocal images of dissected intestines at 1 hpt **(A)** and 6 hpt **(B, D)**. dpf = days post fertilization; hpt = hours post treatment; Scale bars = 20 μm in A; 100 μm in B; 40 μm in D.

MTZ-induced YFP+ cell ablation elicits acute immune cell activation

Immune cell activation plays a crucial role in neuronal regeneration (Kyritsis et al., 2012; White

et al., 2017). To determine whether MTZ-induced cell ablation stimulates an immune response,

we analyzed immune cells dynamics using the pan-leukocyte marker L-plastin (Hiroto

Shinomiya, 2012). In YFP- MTZ-treated (non-ablated) and YFP+ MTZ-treated (ablated) larvae,

we quantified L-plastin+ cell numbers in gut across the time points of neuronal cell death (6 hpt,

12 hpt, and 1 dpt; see **Figure S2**). At 6 hpt, YFP+ MTZ-treated larvae exhibited a significant

increase in L-plastin+ cells compared with YFP- MTZ-treated larvae. By 12 hpt and 1 dpt, no

significant differences were detected (**Figure 4A, B**). Consistently, fold-change analysis

revealed a peak in L-plastin+ cell numbers at 6 hpt, which steadily declined back to normal

levels by 1 dpt (**Figure 4C**). These findings indicate that MTZ-induced ablation of YFP+ cells

trigger a transient immune response in the gut, characterized by an acute accumulation of L-

plastin+ cells that resolves within 24 hours.

Figure 4. L-plastin+ immune cells increase rapidly after YFP+ cell death in the gut.
(A) L-plastin+ immune cells (magenta) in YFP- MTZ (non-ablated) and YFP+ MTZ (ablated) larvae at 6 hpt, 12 hpt, and 1 dpt. **(B)** Quantification of L-plastin+ cells at indicated time points. At 6 hpt, YFP+ MTZ groups shows a significant increase in L-plastin+ cells in the gut compared with YFP- MTZ. No significant difference (ns) was detected at 12 hpt and 1 dpt. **(C)** Fold-change analysis of L-plastin+ cell numbers in YFP+ MTZ larvae relative to YFP- MTZ larvae across time points shows a peak at 6 hpt followed by a decline to baseline levels by 1 dpt. **(A)** Maximum projection confocal images of dissected intestines at time point indicated. Data are shown as mean ± SD (n= 10 larvae per group, 2 experiments); Statistical significance: **p < 0.01; ns = no significant difference; hpt = hours post treatment; dpt = day post treatment; Scale bar = 50 μm.

YFP+ neurons regenerate following targeted ablation

Next, we investigated whether YFP+ neuron numbers were restored to control levels, which would indicate that they regenerate after MTZ treatment. We quantified YFP+ neurons along the whole gut at different time points before and after treatment in YFP+ MTZ and YFP+ control larvae. YFP+ control larvae exhibited an obvious neuronal network from 0 to 15 dpt (**Figure 5A-B**). In contrast, YFP+ neuronal numbers were reduced in the YFP+ MTZ larvae at 1 dpt, confirming successful ablation of YFP+ neurons (**Figure 5A-B**). By 2 dpt, we find a small number of YFP+ neurons along with developing neurites that bridge across the gut (**Figure S3**). YFP+ neuron numbers continue to increase, but through 6 dpt there was a significant difference in YFP+ numbers compared to the control group. By 9 dpt, regenerated YFP+ neuron numbers reached levels comparable to the controls and remained at control levels through 12 and 15 dpt (**Figure 5A-B**). We noticed that approximately half of the YFP+ MTZ larvae exhibited abnormal body posture after 9 dpt. This phenotype might be caused by off-target ablation of reporter-positive cell in the spinal cord area (**Figure S4**). To investigate whether this affected YFP+ neuron regeneration, we compared regenerated YFP+ neuron counts in larvae with normal posture and those with defective posture at 12 dpt but found no significant difference (**Figure S5**). This indicates that postural defects did not impact neuron regeneration. To determine if there were differences in the timeline of neuron regeneration within discrete gut regions compared to the whole gut, we quantified YFP+ neurons in three main gut regions - anterior, midgut, and posterior (Jianlong Li et al., 2020; Wallace et al., 2005). We found that the timeline of YFP+ neuron regeneration was the same in the three gut regions as in the whole gut (**Figure 5C, B-F**).

As shown above, YFP+ neurons colocalized with a small subset of HuC/D+ neurons, indicating that only a fraction of ENS neurons were ablated (**Figure 2D-F**). To assess if ablation of YFP+ neurons has any effects on the YFP- ENS neuronal population, we quantified HuC/D+ neurons in YFP+ MTZ and YFP+ control groups at 0 dpt (before treatment) and 1 and 3 dpt. We found no significant difference in HuC/D+ neuron numbers between treated and control groups after ablation of YFP+ neurons (**Figure S6A-D**). This lack of significance may be due to the relatively small proportion of YFP+ neurons within the total HuC/D+ population—on average, 44.4 out of 204.6 neurons (**Figure 2F**). Additionally, MTZ treatment alone in YFP- MTZ treated larvae did not significantly decrease overall HuC/D numbers (**Figure S6E**).

Figure 5. Timeline of YFP+ neuron regeneration in the whole intestine and defined gut regions.

(A) YFP+ neurons (labelled with anti-GFP antibody) and the neuronal network in YFP+ control and YFP+ MTZ zebrafish larvae at 0 (before treatment) and 1, 3, 6, 9, 12, and 15 dpt.
(B) YFP+ control larvae exhibited consistent GFP+ neuron numbers till 6 dpt with a slight decrease by 9 through 15 dpt. In contrast, the YFP+ MTZ group showed a dramatic decrease in YFP+ neurons, confirming complete ablation of YFP+ neurons ($p < 0.001$) at 1 dpt. A small number of YFP+ neurons appeared by 3 and 6 dpt, but YFP+ neuron numbers were still

significantly decreased compared to the control. By 9 dpt, neuronal number fully recovered with no significant difference between treated and controls at 9, 12 and 15 dpt. **(C)** Schematic representation of the zebrafish gut regions shows the three analyzed gut regions (magenta = anterior, green = midgut, blue = posterior) at 0–9 dpt and 12–18 dpt based on Jianlong Li et al. (2020) and Wallace et al. (2005). Anterior **(D)**, mid **(E)** and posterior **(F)** gut regions show the same timeline of YFP+ neuron regeneration as the whole gut **(B)**. **(A)** Maximum projection confocal images of dissected intestines at time point indicated; Data are presented as mean ± SD (10 larvae per group, 2 experiments); Statistical significance: ***$p < 0.001$, **$p < 0.01$ and *$p < 0.05$; dpt = days post treatment; Scale bar = 100 µm.

Specific co-expressed neuronal subtypes regenerate following ablation

The ENS is composed of various neuron subtypes which control the different functions of the gut including gut motility (Holmberg et al., 2006; Roach et al., 2013; Uyttebroek et al., 2010). The subtypes can be grouped into three main classes which include nitrergic/VIPergic inhibitory motor neurons [neuronal nitric oxide synthase+ (nNOS+); *vasoactive intestinal peptide*+ (*vip*+)], cholinergic excitatory motor neurons [*solute carrier family 18 member 3a*+ slc18a3a+)], and serotonergic neurons [5-hydroxytryptamine+, 5-HT+)] (Kuil et al., 2023; Li et al., 2025; Uyttebroek et al., 2010). To determine the neuronal subtype composition within the YFP+ neuronal population, we analyzed co-expression of these subtype markers with YFP at 12 dpf, prior to MTZ treatment. At 0 dpt, we found that 57.54%, 12.26%, 13.33%, and 6.58% of YFP+ neurons co-expressed nNOS, 5-HT, *vip*, and *slc18a3a*, respectively, indicating the presence of distinct YFP+ neuronal subtypes prior to ablation, which fall into the three major classes of ENS neurons [(Kuil et al., 2023; Li et al., 2025; Uyttebroek et al., 2010), **Figure S7**].

To examine if all YFP+ expressing subtypes regenerated, we analyzed the number of YFP+/subtype+ neurons in the MTZ-treated group at 9 dpt and compared them to untreated control and baseline at 0 dpt control. Notably, the number of YFP+/nNOS+ neurons remained unchanged between 0 control and 9 dpt control, and there was no significant difference between MTZ-treated and control group, suggesting robust stability and regenerative capacity for YFP+/nNOS+ neurons (**Figure 6A, B**). Analysis of YFP+/5-HT+ neurons also revealed no significant difference between 0 dpt control and the 9 dpt control group. However, in the MTZ-treated group, the number of YFP+/5-HT+ neurons at 9 dpt remained significantly lower compared to control, indicating a limited regenerative capacity or delayed regenerative kinetics for this neuronal subtype (**Figure 6C, D**).

Using HCR to detect *vip* and *slc18a3a* expression, we found that the number of YFP+/*vip*+

neurons significantly decreased within the control group between 0 and 9 dpt. However, no

significant difference was observed between the MTZ-treated and control groups at 9 dpt,

suggesting that YFP+/*vip*+ neurons had successfully regenerated by this time point (**Figure 6E,**

F). Lastly, analysis of the cholinergic marker *slc18a3a* showed no significant differences across

groups, indicating that YFP+/*slc18a3a*+ neurons remained stable in the control group and were

successfully restored following regeneration (**Figure 6G, H**).

Overall, these findings suggest that the majority of YFP+ cells are nitrergic neurons before

ablation. After regeneration, YFP+/nNOS+, YFP+/*vip*+ and YFP+/*slc18a3a*+ neurons are

successfully restored. In contrast the YFP+/5-HT+ neurons exhibited lower and/or slower

regenerative capacity compared to the other subtypes.

Figure 6: YFP+/nNOS+, YFP+/*vip*+ and YFP+/*slc18a3a*+ but not YFP+/5-HT+ neurons regenerate following ablation.

(A) Example of nNOS+ (magenta) and YFP+ (labeled with anti-GFP antibody, green) neurons that co-localize (overlay) at 0 dpt control (before treatment) and at 9 dpt in YFP+ control and YFP+ MTZ groups (white arrowheads). (B) Quantification of GFP+/nNOS+ neurons at 0 dpt control and 9 dpt in YFP+ control and YFP+ MTZ shows no significant difference. (C) Example of 5-HT (magenta) and YFP+ (labeled with anti-GFP antibody, green) neurons that co-localize (overlay) at 0 dpt control and 9 dpt in YFP+ control and YFP+ MTZ groups (white arrowheads). (D) Quantification of GFP+/5-HT+ at 0 dpt control and 9 dpt in YFP+ control and YFP+ MTZ. There is no significant difference between 0 dpt and 9 dpt in the YFP+ control group, but GFP+/5-HT+ neurons are significantly reduced in in the MTZ-treated group compared to the control group at 9 dpt. (E) Example of *vip* (magenta) and YFP+ (labeled with anti-GFP antibody, green) neurons that co-localize (overlay) at 0 dpt control (before treatment), and 9 dpt in YFP+ control and YFP+ MTZ (white arrowheads). (F) Quantification of GFP+/*vip*+ neuron at 0 dpt control and 9 dpt in both YFP+ control and YFP+ MTZ groups. A significant difference is observed between 0 dpt and 9 dpt control groups. However, there is no significant difference between 9 dpt control and MTZ-treated groups. (G) Example of *slc18a3a* (magenta) and YFP+ (labeled with anti-GFP antibody, green) neurons that co-localize (overlay) at 0 dpt control and 9 dpt of YFP+ control and YFP+ MTZ (white arrowheads). (H) Quantification of GFP+/*slc18a3a*+ neurons shows no significant difference between the groups. (A, C, E, G) Maximum projection confocal images of dissected intestines at stage indicated. Data are presented as mean ± SD (n = 10 larvae per group, 2 experiments). Statistical significance: ***$p < 0.001$ and *$p < 0.05$; ns = non-significant difference; dpt = days post treatment; Scale bars = 10 µm.

430 **Decreased gut wave movements recover to control levels after neuronal regeneration**

431 ENS neurons play a critical role in regulating gut motility. Excitatory motor neurons mediate muscle

432 contractions and inhibitory motor neurons induce muscle relaxation (Furness, 2012; Kunze &

433 Furness, 1999). We find that major portion of YFP+ neurons co-express inhibitory and excitatory

434 motor neurons markers (**Figure S6**). Thus, we aimed to determine i) whether the loss of YFP+

435 neurons affects gut motility and ii) whether the regenerated neurons restore gut function after

436 ablation.

437 To assess this, we examined one measure of gut motility, the time of the Peristaltic Wave Interval

438 (PWI) of anterograde movements (Holmberg et al., 2006). We measured the PWI at three different

439 time points at 1 dpt, 3 dpt and 9 dpt in YFP+ and YFP- MTZ treated fish, and YFP+ untreated

controls. We included the YFP- MTZ treated group to determine if MTZ treatment alone impacted

PWI. At 1 dpt, we found no significant difference in the PWI time between control and treated groups

(**Figure 7A, Supplementary Videos S1-3**). However, at 3 dpt, the PWI time in the YFP+ MTZ

treated group increased significantly compared to both control groups, suggesting slower peristaltic

movement in the YFP+ MTZ group (**Figure 7B, supplementary videos S4-6**). At 9 dpt, the PWI

time in the YFP+ MTZ treated group was back to control levels (**Figure 7C, supplementary videos**

S7-9). Overall, our results suggest that gut motility function is affected after YFP+ neuronal loss and

then restored after regeneration, indicating functional recovery.

Figure 7. Peristaltic Interval Wave time recovers at 9 dpt after YFP+ neuron ablation. (A) At
1dpt, there is no significant difference in gut PWI time across all experimental groups, YFP+ MTZ
treated, YFP+ untreated control, YFP- MTZ treated. **(B)** At 3dpt, YFP+MTZ treated larvae showed
significantly slower gut PWI time compared to both control groups. **(C)** At 9dpt, there was no
significant difference in PWI across all experimental groups. Data are presented as mean ± SD (n
= 6 larvae per group, 2 experiments); Statistical significance: **$p < 0.01$; ns = non-significant; dpt
= days post treatment.

Discussion

450   The topic of reparative regeneration has gained significant attention over the past decade due to its

451   potential therapeutic implications, for example to treat ENS diseases with loss of ENS neurons. Yet,

452   to date, no study has demonstrated functional ENS neuron regeneration either in mammalian

453   models or in zebrafish. In the present study, we show targeted chemogenetic ablation and functional

regeneration of ENS neurons for the first time in zebrafish (**Figure 8**). Our results determined the spatial and temporal cellular dynamics of targeted neuron ablation as demonstrated by increased TUNEL labeling at 6hpt. Additionally, we observed an acute L-plastin+ immune cell response at 6 hpt that resolves within 1 dpt (**Figure 8**). We find full recovery of ablated neuron numbers by 9 dpt for the majority on ENS neuron subtypes investigated. Specifically, among the regenerated neurons, we found full recovery of nitrergic, cholinergic, and VIPergic subtypes, but only partial or delayed regeneration of serotonergic neurons. Importantly, we also showed functional impacts of neuronal ablation on a gut motility parameter and its functional recovery suggesting functional regeneration in our system (**Figure 8**).

Figure 8. Timeline of events after targeted ablation and regeneration of ENS neurons in zebrafish. Timeline from 0 dpt / 12 dpf to 9 dpt / 21 dpf depicts the different events: i) Neuronal subtype composition of YFP+ neuron population at 0 dpt; ii) Chemogenetic ablation approach activates cell death of YFP+ neurons to completely ablate YFP+ neurons; iii) neuronal regeneration at 3 and 6 dpt which affects gut peristalsis at 3dpt; iv) complete regeneration of YFP+ cells and functional recovery at 9 dpt. **Panel 1** (0dpt / 12dpf): YFP+ cells co-localize with the pan-neuronal marker HuC/D and various neuronal subtypes (nNOS, 5-HT, *vip*, and *slc18a31a*). **Panel 2** (1dpt / 13dpf): Set-up of Gal4-UAS:NTR 2.0 based ablation system. MTZ treatment induces DNA-damage cell death in YFP+ neurons as

confirmed by TUNEL assay at 6hpt. Additionally, there is an acute accumulation of L-plastin+ immune cells at 6hpt that resolves within 24 hours. **Panel 3** (3dpt / 15 dpf): Neuronal ablation resulted in increased peristaltic wave interval time, reflecting reduced gut motility. **Panel 4** (9dpt / 21dpf): Complete recovery of YFP+ neurons and most neuronal subtypes (nNOS+, *vip+*, *slc18a31+*), except 5-HT+ neurons and full functional recovery of gut motility. dpf = days post fertilization, dpt = days post treatment.

Extensive ENS neuron ablation and their complete regeneration in zebrafish

In zebrafish, the only two studies to study ENS neurons regeneration used focal laser ablation to
target a few ENS neurons in a small gut region. Both studies reported generation of new neurons
following ablation (El-Nachef & Bronner, 2020, Ohno et al., 2021). However, quantification of
regenerated neurons showed that they did not return to control levels in the analyzed time frame,
nor were impacts on gut function investigated (Ohno et al., 2021). In our study, using chemogenetic
ablation of ~44.4 ENS neurons, we find robust regeneration of ablated cells to control levels by 9 dpt
which stay at control levels through 15 dpt, the last time point that we analyzed. We find that
different neuronal subtypes within the three main ENS neuron classes are able to regenerate but
identify interesting subtype-specific differences in regenerative capacity or timing (**Figure 8**). Our
work also shows for the first time recovery of a parameter of gut function - gut wave propagation -
indicative of successful functional regeneration (**Figure 8**). In addition, our approach has several
experimental advantages compared to laser ablation. First, laser ablation is less efficient, as it can
only ablate a limited number of neurons within specific gut regions (El-Nachef & Bronner, 2020,
Ohno et al., 2021). For instance, one study only ablated ~16 neurons within one-somite-width of the
posterior gut at 10-15 dpf (Ohno et al., 2021). In contrast, our chemogenetic ablation system
successfully ablated on average 44 neurons throughout the whole gut, significantly improving
efficiency. Second, laser ablation is time-consuming as it requires individual larval preparation and
precise manual targeting (Ohno et al., 2021). In contrast, our methods enable the simultaneous
ablation of ENS neurons in several hundred of larvae. Third, laser irradiation can cause unintended
damage to non-neuronal, surrounding tissues, including smooth muscle and epithelial cells, which
may impact regeneration and complicate downstream analyses such as gut motility. In contrast, our
approach is spatiotemporally controlled ablation that precisely targets only specific ENS neurons
and had no deleterious effects on the overall neuronal population in the gut (**Figure 8**).

Zebrafish are a powerful model system to study ENS regeneration

So far, ENS regeneration research has mostly focused on mammalian model systems with only limited work in non-mammalian models such as zebrafish (Corvetti et al., 2001; El-Nachef & Bronner, 2020; Gabella & Trigg, 1984; Galligan et al., 1989; Hanani et al., 2003; Jew et al., 1989; Joseph et al., 2011; Karaosmanoğlu et al., 1996; Laranjeira et al., 2011; Luck et al., 1993; Ohno et al., 2021; Stavely et al., 2024; Tokui et al., 1994). Previous studies in mammalian models involving surgical (Galligan et al., 1989; Karaosmanoğlu et al., 1996; Tokui et al., 1994), stenosis (Corvetti et al., 2001; Gabella & Trigg, 1984; Jew et al., 1989; Joseph et al., 2011), and chemical (Hanani et al., 2003; Laranjeira et al., 2011; Luck et al., 1993) ablation methods consistently report incomplete neuronal regeneration, and gut region- or species-specific differences in regenerative outcomes of ENS neurons (Rueckert and Ganz, 2022). Notably, all previous approaches also reported structural abnormalities at the injury site by co-lateral damage of surrounding tissues. To address this, recent works in adult mice reported focal ablation specifically of ENS neurons using a diphtheria toxin-based chemogenetic ablation approach (Bhave et al., 2019; Stavely et al., 2024). Even though this study reported that mature, post-mitotic enteric neurons in adult mice can regenerate axons and reinnervate the gut after injury, there was still limited regeneration of neurons even three months after ablation (Stavely et al., 2024). This indicates that mammals have generally limited ENS regenerative abilities. In contrast, the present study demonstrates that zebrafish are a powerful model to study the signals and cellular-molecular processes underlying functional ENS regeneration. Our spatio-temporally controlled approach ablates approximately one fifth of all ENS neurons by inducing cell death with subsequent full recovery after only 9 days. Ablation along the entire gut will enable the analysis of cellular responses and gene regulatory programs active during ENS regeneration as whole guts can be analyzed. We find an acute immune response during the time of cell death which normalizes at 1 dpt. Studies on adult zebrafish brain traumatic injury and selective rod photoreceptor ablation in the retina have demonstrated that immune responses are crucial components of regenerative processes (Kyritsis et al., 2012; White et al., 2017). In our study, selective ablation of enteric neurons (**Figure 8**) led to a marked accumulation of L-plastin+ cells as early as 6 hours after MTZ treatment. Given that L-plastin is strongly expressed in leukocytes, including macrophages, monocytes, and neutrophils (Hiroto Shinomiya, 2012), this finding suggests that selective neuronal death in the ENS triggers an acute immune cell recruitment, similar to what has been described in models of traumatic brain injury and retinal regeneration (Kyritsis et al., 2012; White et al., 2017). Future studies will be needed to provide mechanistic insights into how immune–neural interactions (Graves et al., 2021) regulate ENS regeneration. Our approach thus provides a

rapid experimental setup to identify the signals and cellular responses important for neuronal

regeneration in the ENS in the future (**Figure 8**).

Regeneration of neurons in the zebrafish CNS *vs* ENS

The zebrafish central nervous system has been extensively studied with regards to its regenerative

capacity (Kaslin et al., 2009, 2013; März et al., 2010) while little is known about ENS regeneration

(Ohno et al., 2021). In the central nervous system, brain regions show different capacities to

regenerate neuronal subtypes. In the zebrafish telencephalon, all ablated neuron types regenerate

(Barbosa et al., 2015; Kroehne et al., 2011; Reimer et al., 2008), whereas the cerebellum is capable

of regenerating many, but not all neuron types (Kaslin et al., 2017). This variable regenerative

capacity of the cerebellum has been attributed to varying ability of cerebellar stem and progenitor

cells to generate neurons and glia throughout the lifespan (Kaslin et al., 2009, 2013). We also find

differences in regeneration of neuron types in the ENS where some subtypes (nitrergic, cholinergic

and VIPergic) regenerate fully, whereas serotonergic neurons do not reach control levels (**Figure 8**).

There could be three possibilities for this difference in regeneration: (1) serotonergic neurons take

longer to regenerate, (2) the progenitor cells or cellular mechanisms to regenerate serotonergic

neurons are not activated at this stage of post-embryonic development, and (3) the low level of

serotonergic cell loss incurred is insufficient to trigger serotonergic cell replacement. Further work

will determine which cellular responses are triggered after cell ablation and the progenitor

population(s) that generate new ENS neurons and neuronal subtypes in zebrafish.

Supplementary Materials

The following supporting information can be downloaded at: https:

Author Contributions

Conceptualization: M.A.S, K.M., H.R., J.G. Methodology, Data curation, Validation, Formal

Analysis and Investigation: M.A.S, K.M., H.R. Visualization: M.A.S. Writing—Original Draft

Preparation: M.A.S. Writing—Review and Editing: M.A.S, K.M., H.R., A.V.S., D.F.A., J.S.M., and

J.G. Supervision, Project Administration, Funding Acquisition and Recourses: D.F.A., J.S.M.,

J.G. All authors have read and agreed to the published version of the manuscript.

Funding:

This work was supported by NSF CAREER #2143267 to J.G., NIH R21NS123629 to J.G., and a Research Grant from the American Neurogastroenterology and Motility Society to J.G. NIH R01OD020376 to J.S.M. and D.F.A., supported development of the improved NTR 2.0 enzyme.

Institutional Review Board Statement:

The animal study protocol was approved by the Michigan State University Institutional Animal Care and Use Committee.

Acknowledgments

We want to thank Rachel Alcorn, Taylor Lawrence, Giovanna Haddock, Kristen Lounsbury, Anthony Turner, Samantha Lampman, and Sharon An for excellent fish care.

Conflicts of Interest

The authors declare no competing or financial interests

List of Abbreviations

ENS = Enteric nervous system

MTZ = Metronidazole

HSCR = Hirschsprung Disease

DPF = Days post fertilization

DPT = Days post treatment

HPT = Hours post treatment

NTR = Nitroreductase

YFP = Yellow fluorescent protein

GFP = Green fluorescent protein

UAS = Upstream activating sequence

PWI = Peristaltic Wave Interval

TdT = Terminal deoxynucleotidyl Transferase

BSA = Bovin serum albumin

PBS = Phosphate-Buffered Saline

HCR = Hybridization Chain Reaction

SD = standard deviation

References

Ariga, J., Walker, S. L., & Mumm, J. S. (2010). Multicolor time-lapse imaging of transgenic zebrafish: Visualizing retinal stem cells activated by targeted neuronal cell ablation. *Journal of Visualized Experiments*, *43*. https://doi.org/https://doi.org/10.3791/2093

Asakawa, K., Suster, M. L., Mizusawa, K., Nagayoshi, S., Kotani, T., Urasaki, A., Kishimoto, Y., Hibi, M., & Kawakami, K. (2008). Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4 gene and enhancer trapping in zebrafish. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(4), 1255–1260. https://doi.org/10.1073/pnas.0704963105

Bandla, A., Melancon, E., Taylor, C. R., Davidson, A. E., Eisen, J. S., & Ganz, J. (2022). A New Transgenic Tool to Study the Ret Signaling Pathway in the Enteric Nervous System. *International Journal of Molecular Sciences*, *23*(24), 15667. https://doi.org/10.3390/ijms232415667

Barbosa, J. S., Sanchez-Gonzalez, R., Di Giaimo, R., Baumgart, E. V., Theis, F. J., Götz, M., & Ninkovic, J. (2015). Live imaging of adult neural stem cell behavior in the intact and injured zebrafish brain. *Science*, *348*(6236), 789–793. https://doi.org/10.1126/science.aaa2729

Bhave, S., Arciero, E., Baker, C., Ho, W. L., Stavely, R., Goldstein, A. M., & Hotta, R. (2019). Enteric neuronal cell therapy reverses architectural changes in a novel diphtheria toxin-mediated model of colonic aganglionosis. *Scientific Reports*, *9*(1), 1–12. https://doi.org/10.1038/s41598-019-55128-4

Burns, A. J., Goldstein, A. M., Newgreen, D. F., Stamp, L., Metzger, M., Hotta, R., Young, H. M., Andrews, P. W., Belkind-gerson, J., Bondurand, N., Bornstein, J. C., Yee, W., Cheah, K., Gershon, M. D., Heuckeroth, R. O., Robert, M. W., Just, L., Kapur, R. P., King, S. K., … Urology, P. (2016). *White paper on guidelines concerning enteric nervous system stem cell therapy for enteric neuropathies⋆*. *417*(2), 229–251. https://doi.org/10.1016/j.ydbio.2016.04.001

Butler Tjaden, N. E., & Trainor, P. A. (2013). The developmental etiology and pathogenesis of Hirschsprung disease. *Translational Research*, *162*(1), 1–15. https://doi.org/10.1016/j.trsl.2013.03.001

Cao, J., & Poss, K. D. (2018). The epicardium as a hub for heart regeneration. *Nature Reviews Cardiology*, *15*(10), 631–647. https://doi.org/10.1038/s41569-018-0046-4

Corvetti, G., Fornaro, M., Geuna, S., Poncino, A., & Giacobini-Robecchi, M. G. (2001). Unscheduled DNA synthesis in rat adult myenteric neurons: An immunohistochemical study. *NeuroReport*, *12*(10), 2165–2168. https://doi.org/10.1097/00001756-200107200-00024

Cusack, C. L., Swahari, V., Henley, W. H., Ramsey, J. M., & Deshmukh, M. (2013). Distinct Pathways Mediate Axon Degeneration during Apoptosis and Axon-Specific Pruning. *Nature Communications*, *4*(1876). https://doi.org/doi: 10.1038/ncomms2910

Davison, J. M., Akitake, C. M., Goll, M. G., Rhee, J. M., Gosse, N., Baier, H., Halpern, M. E., Leach, S. D., & Parsons, M. J. (2007). Transactivation from Gal4-VP16 transgenic insertions for tissue-specific cell labeling and ablation in zebrafish. *Developmental Biology*, *304*(2), 811–824. https://doi.org/10.1016/j.ydbio.2007.01.033

Diep, C. Q., Ma, D., Deo, R. C., Holm, T. M., Naylor, R. W., Arora, N., Wingert, R. A., Bollig, F., Djordjevic, G., Lichman, B., Zhu, H., Ikenaga, T., Ono, F., Englert, C., Cowan, C. A., Hukriede, N. A., Handin, R. I., & Davidson, A. J. (2011). Identification of adult nephron progenitors capable of kidney regeneration in zebrafish. *Nature*, *470*, 95–101. https://doi.org/10.1038/nature09669

El-Nachef, W. N., & Bronner, M. E. (2020). De novo enteric neurogenesis in post-embryonic zebrafish from Schwann cell precursors rather than resident cell types. *Development (Cambridge)*, *147*(13). https://doi.org/10.1242/dev.186619

Furness, J. B. (2012). The enteric nervous system and neurogastroenterology. *Nature Reviews Gastroenterology and Hepatology*, *9*(5), 286–294. https://doi.org/10.1038/nrgastro.2012.32

Gabella, G., & Trigg, P. (1984). Size of neurons and glial cells in the enteric ganglia of mice, guinea-pigs, rabbits and sheep. *Journal of Neurocytology*, *13*(1), 49–71. https://doi.org/10.1007/BF01148318

Galligan, J. J., Furness, J. B., & Costa, M. (1989). Migration of the myoelectric complex after interruption of the myenteric plexus: Intestinal transection and regeneration of enteric nerves in the guinea pig. *Gastroenterology*, *97*(5), 1135–1146. https://doi.org/10.1016/0016-5085(89)91683-1

Ganz, J. (2018). Gut feelings: Studying enteric nervous system development, function, and disease in the zebrafish model system. *Developmental Dynamics*, *247*(2), 268–278. https://doi.org/10.1002/dvdy.24597

Graves, C. L., Chen, A., Kwon, V., & Shiau, C. E. (2021). Zebrafish harbor diverse intestinal macrophage populations including a subset intimately associated with enteric neural processes. *IScience*, *24*(6), 102496. https://doi.org/10.1016/j.isci.2021.102496

Ham, T. J., Mapes, J., Kokel, D., & Peterson, R. T. (2010). Live imaging of apoptotic cells in zebrafish. *The FASEB Journal*, *24*(11), 4336–4342. https://doi.org/10.1096/fj.10-161018

Hanani, M., Ledder, O., Yutkin, V., Abu-Dalu, R., Huang, T. Y., Härtig, W., Vannucchi, M. G., & Faussone-Pellegrini, M. S. (2003). Regeneration of myenteric plexus in the mouse colon after experimental denervation with benzalkonium chloride. *Journal of Comparative Neurology*, *462*(3), 315–327. https://doi.org/10.1002/cne.10721

Hanovice, N. J., Leach, L. L., Slater, K., Gabriel, A. E., Romanovicz, D., Shao, E., Collery, R., Burton, E. A., Lathrop, K. L., Link, B. A., & Gross, J. M. (2019). Regeneration of the zebrafish retinal pigment epithelium after widespread genetic ablation. In *PLoS Genetics* (Vol. 15, Issue 1). https://doi.org/10.1371/journal.pgen.1007939

Hao, M. M., & Young, H. M. (2009). Development of enteric neuron diversity. *Journal of Cellular and Molecular Medicine*, *13*(7), 1193–1210. https://doi.org/10.1111/j.1582-4934.2009.00813.x

Heanue, T. A., Boesmans, W., Bell, D. M., Kawakami, K., Berghe, P. Vanden, & Pachnis, V. (2016). *A Novel Zebrafish ret Heterozygous Model of Hirschsprung Disease Identifies a Functional Role for mapk10 as a Modifier of Enteric Nervous System Phenotype Severity*. *12*(11), 1–23. https://doi.org/doi:10.1371/journal.pgen.1006439

Hiroto Shinomiya. (2012). Plastin Family of Actin-Bundling Proteins: Its Functions in
Leukocytes, Neurons, Intestines, and Cancer. *Nternational Journal of Cell Biology*,
11(1), 1–8. https://doi.org/10.1155/2012/213492

Holmberg, A., Olsson, C., & Holmgren, S. (2006). The effects of endogenous and
exogenous nitric oxide on gut motility in zebrafish Danio rerio embryos and larvae.
Journal of Experimental Biology, *209*(13), 2472–2479.
https://doi.org/10.1242/jeb.02272

Howe, K., Clark, M. D., Torroja, C. F., Torrance, J., Berthelot, C., & Muffato, M. (2013).
The zebrafish reference genome sequence and its relationship to the human
genome. *Nature*, *496*(7446), 598–503. https://doi.org/10.1038/nature12111.The

Ibarra-García-Padilla, R., Howard, A. G. A., Singleton, E. W., & Uribe, R. A. (2021). A
protocol for whole-mount immuno-coupled hybridization chain reaction (WICHCR)
in zebrafish embryos and larvae. *STAR Protocols*, *2*(3).
https://doi.org/10.1016/j.xpro.2021.100709

Jew, J. Y., Zhang, M. Q., Williams, T. H., & Gabella, G. (1989). The Intestine as a Model
for Neuronal Plasticity. *Archives of Histology and Cytology*, *52*(Suppl.), 167–180.
https://doi.org/10.1679/aohc.52.Suppl_167

Jianlong Li, Prochaska, M., Maney, L., & Wallace, K. N. (2020). Development and
organization of the zebrafish intestinal epithelial stem cell niche. *Development
Dynamics*, *249*(1), 76–87.

Joseph, N. M., He, S., Quintana, E., Kim, Y. G., Núñez, G., & Morrison, S. J. (2011).
Enteric glia are multipotent in culture but primarily form glia in the adult rodent gut.
Journal of Clinical Investigation, *121*(9), 3398–3411.
https://doi.org/10.1172/JCI58186

Karaosmanoğlu, T., Müftüoğlu, S., Dağdeviren, A., Durgun, B., Aygün, B., & Örs, Ü.
(1996). Morphological changes in the myenteric plexus of rat ileum after transection
and end-to-end anastomosis. *Journal of Anatomy*, *188*(2), 323–331.

Kaslin, J., Ganz, J., Geffarth, M., Grandel, H., Hans, S., & Brand, M. (2009). Stem cells
in the adult zebrafish cerebellum: Initiation and maintenance of a novel stem cell
niche. *Journal of Neuroscience*, *29*(19), 6142–6153.
https://doi.org/10.1523/JNEUROSCI.0072-09.2009

Kaslin, J., Kroehne, V., Benato, F., Argenton, F., & Brand, M. (2013). Development and
specification of cerebellar stem and progenitor cells in zebrafish: From embryo to
adult. *Neural Development*, *8*(1). https://doi.org/10.1186/1749-8104-8-9

Kaslin, J., Kroehne, V., Ganz, J., Hans, S., Brand, M., Kaslin, J., Kroehne, V., Ganz, J.,
Hans, S., & Brand, M. (2017). Correction to: Distinct roles of neuroepithelial-like
and radial glia-like progenitor cells in cerebellar regeneration (Development
(Cambridge), (2017) 144, (1462-1471), 10.1242/dev.144907). *Development
(Cambridge)*, *144*(18), 3388. https://doi.org/10.1242/dev.158600

Kawakami, K., Abe, G., Asada, T., Asakawa, K., Fukuda, R., Ito, A., Lal, P., Mouri, N.,
Muto, A., Suster, M. L., Takakubo, H., Urasaki, A., Wada, H., & Yoshida, M. (2010).
ZTrap: Zebrafish gene trap and enhancer trap database. *BMC Developmental
Biology*, *10*. https://doi.org/10.1186/1471-213X-10-105

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., & F., B. U. homas. (1995). Stages of
embryonic development of the zebrafish. *Developmental Dynamics*, *203*, 253–310.
https://doi.org/https://doi.org/10.1002/aja.1002030302

Kizil, C., Kaslin, J., Kroehne, V., & Brand, M. (2012). Adult neurogenesis and brain regeneration in zebrafish. *Developmental Neurobiology*, *72*(3), 429–461. https://doi.org/10.1002/dneu.20918

Kroehne, V., Freudenreich, D., Hans, S., Kaslin, J., & Brand, M. (2011). Regeneration of the adult zebrafish brain from neurogenic radial glia-type progenitors. *Development*, *138*(22), 4831–4841. https://doi.org/10.1242/dev.072587

Kuil, L. E., Chauhan, R. K., Cheng, W. W., Hofstra, R. M. W., & Alves, M. M. (2021). Zebrafish: A Model Organism for Studying Enteric Nervous System Development and Disease. *Frontiers in Cell and Developmental Biology*, *8*(January), 1–15. https://doi.org/10.3389/fcell.2020.629073

Kuil, L. E., Kakiailatu, N. J. M., Windster, J. D., Bindels, E., Zink, J. T. M., van der Zee, G., Hofstra, R. M. W., Shepherd, I. T., Melotte, V., & Alves, M. M. (2023). Unbiased characterization of the larval zebrafish enteric nervous system at a single cell transcriptomic level. *IScience*, *26*(7), 107070. https://doi.org/10.1016/j.isci.2023.107070

Kulkarni, A. A., Conteh, A. M., Sorrell, C. A., Mirmira, A., Tersey, S. A., Mirmira, R. G., Linnemann, A. K., & Anderson, R. M. (2018). An in vivo zebrafish model for interrogating ROS-mediated pancreatic β-cell injury, response, and prevention. *Oxidative Medicine and Cellular Longevity*, 1–8. https://doi.org/10.1155/2018/1324739

Kunze, W. A. A., & Furness, J. B. (1999). The enteric nervous system and regulation of intestinal motility. *Annual Review of Physiology*, *61*, 117–142. https://doi.org/10.1146/annurev.physiol.61.1.117

Kyritsis, N., Kizil, C., Zocher, S., Kroehne, V., Kaslin, J., Freudenreich, D., Iltzsche, A., & Brand, M. (2012). Acute inflammation initiates the regenerative response in the adult zebrafish brain. *Science*, *338*(6112), 1353–1356. https://doi.org/10.1126/science.1228773

Laranjeira, C., Sandgren, K., Kessaris, N., Richardson, W., Potocnik, A., Vanden Berghe, P., & Pachnis, V. (2011). Glial cells in the mouse enteric nervous system can undergo neurogenesis in response to injury. *Journal of Clinical Investigation*, *121*(9), 3412–3424. https://doi.org/10.1172/JCI58200

Lee, E., Song, C. H., Bae, S. J., Ha, K. T., & Karki, R. (2023). Regulated cell death pathways and their roles in homeostasis, infection, inflammation, and tumorigenesis. *Experimental and Molecular Medicine*, *55*(8), 1632–1643. https://doi.org/10.1038/s12276-023-01069-y

Li, C., Gehring, J., & Bronner, M. E. (2025). Spatiotemporal dynamics of the developing zebrafish enteric nervous system at the whole-organ level. *Developmental Cell*, *60*(4), 613-629.e6. https://doi.org/10.1016/j.devcel.2024.11.006

Luck, M. S., Dahl, J. L., Boyeson, M. G., & Bass, P. (1993). Neuroplasticity in the smooth muscle of the myenterically and extrinsically denervated rat jejunum. *Cell and Tissue Research*, *271*(2), 363–374. https://doi.org/10.1007/BF00318623

März, M., Chapouton, P., Diotel, N., Vaillant, C., Birgit Hesl, M. T., Lam, C. S., Kah, O., Bally-Cuif, L., & Strähle, U. (2010). Heterogeneity in Progenitor Cell Subtypes in the Ventricular Zone of the Zebrafish Adult Telencephalon. *Glia*, *58*(7), 870–888. https://doi.org/10.1002/glia.20971.

Medico, E., Gambarotta, G., Gentile, A., Comoglio, P. M., & Soriano, P. (2001). A gene trap vector system for identifying transcriptionally responsive genes. *Nature Biotechnology*, *19*(6), 579–582. https://doi.org/10.1038/89343

Niesler, B., Kuerten, S., Demir, I. E., & Schäfer, K. H. (2021). Disorders of the enteric nervous system — a holistic view. *Nature Reviews Gastroenterology and Hepatology*, *18*(6), 393–410. https://doi.org/10.1038/s41575-020-00385-2

Ohno, M., Nikaido, M., Horiuchi, N., Kawakami, K., & Hatta, K. (2021). The enteric nervous system in zebrafish larvae can regenerate via migration into the ablated area and proliferation of neural crest-derived cells. *Development*, *148*(2). https://doi.org/10.1242/dev.195339

Olden, T., Akhtar, T., Beckman, S. A., & Wallace, K. N. (2008). Differentiation of the Zebrafish Enteric Nervous System and Intestinal Smooth Muscle. *Genesis*, *46*(9), 484–498. https://doi.org/10.1002/dvg.20429.

Reimer, M. M., Sörensen, I., Kuscha, V., Frank, R. E., Liu, C., Becker, C. G., & Becker, T. (2008). Motor neuron regeneration in adult zebrafish. *Journal of Neuroscience*, *28*(34), 8510–8516. https://doi.org/10.1523/JNEUROSCI.1189-08.2008

Roach, G., Heath Wallace, R., Cameron, A., Emrah Ozel, R., Hongay, C. F., Baral, R., Andreescu, S., & Wallace, K. N. (2013). Loss of ascl1a prevents secretory cell differentiation within the zebrafish intestinal epithelium resulting in a loss of distal intestinal motility. *Developmental Biology*, *376*(2), 171–186. https://doi.org/10.1016/j.ydbio.2013.01.013

Romero, M. M. G., McCathie, G., Jankun, P., & Roehl, H. H. (2018). Damage-induced reactive oxygen species enable zebrafish tail regeneration by repositioning of Hedgehog expressing cells. *Nature Communications*, *9*(1). https://doi.org/10.1038/s41467-018-06460-2

Roy-Carson, S., Natukunda, K., Chou, H. chao, Pal, N., Farris, C., Schneider, S. Q., & Kuhlman, J. A. (2017). Defining the transcriptomic landscape of the developing enteric nervous system and its cellular environment. *BMC Genomics*, *18*(1), 1–24. https://doi.org/10.1186/s12864-017-3653-2

Rueckert, H., & Ganz, J. (2022). How to Heal the Gut's Brain: Regeneration of the Enteric Nervous System. *International Journal of Molecular Sciences*, *23*(9). https://doi.org/10.3390/ijms23094799

Sharrock, A. V., Mulligan, T. S., Hall, K. R., Williams, E. M., White, D. T., Zhang, L., Emmerich, K., Matthews, F., Nimmagadda, S., Washington, S., Le, K. D., Meir-Levi, D., Cox, O. L., Saxena, M. T., Calof, A. L., Lopez-Burks, M. E., Lander, A. D., Ding, D., Ji, H., … Mumm, J. S. (2022). NTR 2.0: a rationally engineered prodrug-converting enzyme with substantially enhanced efficacy for targeted cell ablation. *Nature Methods*, *19*(2), 205–215. https://doi.org/10.1038/s41592-021-01364-4

Soret, R., Schneider, S., Bernas, G., Christophers, B., Souchkova, O., Charrier, B., Righini-Grunder, F., Aspirot, A., Landry, M., Kembel, S. W., Faure, C., Heuckeroth, R. O., & Pilon, N. (2020). Glial Cell-Derived Neurotrophic Factor Induces Enteric Neurogenesis and Improves Colon Structure and Function in Mouse Models of Hirschsprung Disease. *Gastroenterology*, *159*(5), 1824-1838.e17. https://doi.org/10.1053/j.gastro.2020.07.018

Stavely, R., Rahman, A. A., Mueller, J. L., Leavitt, A. R., Han, C. Y., Pan, W., Kaiser, K. N., Ott, L. C., Ohkura, T., Guyer, R. A., Burns, A. J., Koppes, A. N., Hotta, R., &

Goldstein, A. M. (2024). Mature enteric neurons have the capacity to reinnervate the intestine with glial cells as their guide. *Neuron*, *112*(18), 3143-3160.e6. https://doi.org/10.1016/j.neuron.2024.06.018

Taylor, C. R., Montagne, W. A., Eisen, J. S., & Ganz, J. (2016). Molecular fingerprinting delineates progenitor populations in the developing zebrafish enteric nervous system. *Developmental Dynamics, 245*(11), 1081–1096. https://doi.org/10.1002/dvdy.24438

Tokui, K., Sakanaka, M., & Kimura, S. (1994). Progressive reorganization of the myenteric plexus during one year following reanastomosis of the ileum of the guinea pig. *Cell & Tissue Research*, *277*(2), 259–272. https://doi.org/10.1007/BF00327773

Uyttebroek, L., Shepherd, I. T., Harrisson, F., Hubens, G., Blust, R., Timmermans, J.-P., & Nassauw, L. Van. (2010). Neurochemical Coding of Enteric Neurons in Adult and Embryonic Zebrafish (Danio rerio). *J Comp Neuro*, *518*(21), 4419–4438. https://doi.org/10.1002/cne.22464.Neurochemical

Wallace, K. N., Akhter, S., Smith, E. M., Lorent, K., & Pack, M. (2005). Intestinal growth and differentiation in zebrafish. *Mechanisms of Development*, *122*(2), 157–173. https://doi.org/10.1016/j.mod.2004.10.009

Webb, J. F., Shirey, J. E., Desvignes, T., Carey, A., Postlethwait, J. H., Kimmel, S. R., & Ullmann, B. (2009). *Normal table of postembryonic zebra!sh development: Staging by externally visible anatomy of the living fish*. *238*(12), 2975–3015. https://doi.org/10.1002/dvdy.22113

Westerfield, M. (2000). *A Guide for the Laboratory Use of Zebrafish Danio (Brachydanio) rerio* (4th, Ed.). Institute of Neuroscience, University of Oregon, Eugene. Published by the University of Oregon Press.

White, D. T., & Mumm, J. S. (2013). The nitroreductase system of inducible targeted ablation facilitates cell-specific regenerative studies in zebrafish. *Methods*, *62*(3), 232–240. https://doi.org/10.1016/j.ymeth.2013.03.017

White, D. T., Sengupta, S., Saxena, M. T., Xu, Q., Hanes, J., Ding, D., Ji, H., & Mumm, J. S. (2017). Immunomodulation-accelerated neuronal regeneration following selective rod photoreceptor cell ablation in the zebrafish retina. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(18), E3719–E3728. https://doi.org/10.1073/pnas.1617721114

Zhang, Y., Ouyang, J., Qie, J., Zhang, G., Liu, L., & Yang, P. (2019). Optimization of the Gal4/UAS transgenic tools in zebrafish. *Applied Microbiology and Biotechnology*, *103*(4), 1789–1799. https://doi.org/10.1007/s00253-018-09591-0

863   **Supplementary figures**

Figure S1: Inducing cell death shows an expected increase in TUNEL+ cells.
Inducing cell death as a positive control results in a high number of TUNEL-positive cells
(magenta, middle panel) in YFP+ control (green) larvae at 6 hpt (n= 9 larvae, 2
experiments). Maximum projection confocal images of dissected intestine at stage
indicated. Scale bar = 100 μm; hpt = hours post treatment.

865
866
867

Figure S2. Metronidazole (MTZ)-induced ablation of YFP+ neurons is completed within 24 hours post-treatment. (A) YFP+ neurons (labeled with anti-GFP, green) are present at 0 dpt (before treatment of MTZ), but YFP+ neuron numbers continuously decrease after treatment of MTZ at 6 hpt and 12 hpt, and are completely gone at 1 dpt. **(B)** Quantification of YFP+ neurons at time points indicated in YFP+ control (red line) and YFP+ MTZ groups (blue line). The YFP+ control group maintains stable cell numbers

across all time points. In the YFP+ MTZ-treated group YFP+ neuron numbers rapidly decline at 6 hpt, with near-complete ablation by 1 dpt. **(A-C)** Maximum projection confocal images of dissected intestines at time point indicated. Data presented as mean (n = 10 larvae per group, 2 experiments); hpt = hours post-treatment, dpt = hours/days post-treatment; Scale bar = 100 μm.

Figure S3. Regenerating neurites in YFP+ neurons following ablation. At 2 dpt, newly generated YFP+ neurons form neurites as the beginning step of neuron regeneration. Upper panel: at 1 dpt there is complete ablation of YFP+ neurons. Lower panel: at 2 dpt developing neurites that form a bridge (arrow) between the newly generated neurons (n = 6 larvae per group. 2 experiments). Maximum projection of confocal image of dissected intestine at time point indicated. dpt = days post treatment; Scale bar = 100 μm.

Figure S4. Off-target ablation of YFP+ cells from the spinal cord region. (A) In YFP+ control larvae, the intestine contains a network of YFP+ ENS neurons along the gut and in cells within the area of the spinal cord and in the brain. **(B)** Magnified view shows YFP+ cells in the spinal cord area. **(C)** In YFP+ MTZ treated larvae, YFP+ ENS neurons are essentially absent. **(D)** Magnified view shows that YFP+ cells in spinal cord area are also reduced after MTZ treatment, demonstrating off-target NTR-mediated cell ablation in the spinal cord. **(A-C)** Whole-mount side views of live images of YFP+ control (untreated) and YFP+ MTZ (treated) at stages indicated. Arrowheads indicate reporter-positive cells in the gut region. Dashed box outlines the region for magnified view. dpt = days post treatments; (A, C) scale bar = 100 μm; (B, D) scale bar = 20 μm.

Figure S5. Abnormalities in body posture do not affect YFP+ neuron regeneration. (A-B) Whole-mount side views of larvae at stage indicated from treated (YFP-/+ MTZ treated) and Control (YFP+ untreated) groups. Only half of YFP+MTZ treated shows the body posture defect after 9 dpt. **(C)** The bar graph shows the number of regenerated YFP+ neurons in normally developed larvae compared to those with defective body posture at 12 days post treatment. Data presented as mean ± SD (n = 8 larvae per group, 2 experiments); ns = no statistically significant differences; Scale bar = 100 μm.

Figure S6. YFP+ neuron loss and MTZ treatment of YFP- larvae does not affect neuron numbers. (A) At 12 dpf (before treatment, 0 dpt), the pan-neuronal marker HuC/D (magenta) and YFP+ cells (labeled with anti-GFP, green) show high colocalization (overlay). **(B)** Following MTZ treatment at 13dpf (1dpt), GFP-HuC/D+ cells are present, whereas GFP+ cells are completely absent due to ablation of YFP+ neurons. **(C)** 15dpf (3dpt) shows GFP+ cells and exons (indicated by while arrow) regeneration process. **(D)** Quantification of HuC/D-positive neuron counts before (0 dpt) and after MTZ treatment (1 and 3dpt). There was no significant difference in total HuC/D+ cells between control and treatment group (n = 10 larvae

per group, 2 experiments). **(E)** Quantification of HuC/D+ neurons in the gut in YFP- MTZ treated and YFP+ untreated controls at 12dpf (before MTZ treatment) and 13dpf (after MTZ treatment). Bar graph shows the number of HuC/D+ cells. No significant difference was observed between treatment groups (n = 10 larvae, 2 experiments). Data presented as mean ± standard deviation; ns = not significant, dpt = days post treatment. **(A-C)** Maximum projection confocal images of dissected intestines. Scale bar = 100 μm; Data are presented as mean ± SD, ns = not significant; dpf = days post fertilization; dpt = days post treatment.

Figure S7. The YFP+ neuronal population consists of different neuronal subtypes at 12 days post fertilization. Quantification of YFP and neuronal subtype colocalization. Bar graph shows the percentage of co-expression of each subtype marker with YFP at 12 days post fertilization / 0 day post treatment. Data presented as mean (n = 10 larvae per group, 2 experiments).

878 **Supplementary videos have been uploaded to the designated repository.**

Groups	Video description
1dpt	Video S1 = YFP- MTZ
	Video S2 = YFP+ control
	Video S3 = YFP+ MTZ
3dpt	Video S4 = YFP- MTZ

	Video S5 = YFP+ control
	Video S6 = YFP+ MTZ
9dpt	Video S7 = YFP- MTZ
	Video S8 = YFP+ control
	Video S9 = YFP+ MTZ
dpt = days post treatment	

879