Gamete release in *Ciona robusta*: roles of gonadotropin-releasing hormone and the

photoreception system

Tomohiro Osugi[1*], Shin Matsubara[1], Akira Shiraishi[1], Azumi Wada[1], Yuki Miyamoto[2], Issei S.

Shimada[3], Yasunori Sasakura[4], Takehiro G. Kusakabe[2], Honoo Satake[1].

[1]Bioorganic Research Institute, Suntory Foundation for Life Sciences, Soraku-gun, Kyoto, Japan.

[2]Department of Biology, Faculty of Science and Engineering and Institute for Integrative

Neurobiology, Konan University, Kobe, Japan.

[3]Department of Cell Biology, Graduate School of Medical Sciences, Nagoya City University, Japan.

[4]Shimoda Marine Research Center, University of Tsukuba, Shimoda, Shizuoka, Japan.

*Correspondence to: Tomohiro Osugi

E-mail: osugi@sunbor.or.jp

Abstract

Gamete release, an essential event for animals, is regulated by various factors including environmental stimuli, neural circuit, and endocrine molecules. In this study, we investigated the mechanisms regulating gamete release in the ascidian *Ciona robusta*, part of the sister group of vertebrates. *Ciona* is a hermaphrodite, releasing sperm first from orange-pigmented organ (OPO) at the end of the spermiduct, followed by eggs from the oviduct beneath the spermiduct knob-like structure. In this study, behavioral and morphological analyses revealed that this sequential release occurs because the knob-like structure compresses the oviduct opening at the onset of gamete release. Observations of transgenic *Ciona* expressing Kaede under the gonadotropin-releasing hormone 2 (*Gnrh2*) promoter showed that *Gnrh2*-expressing neurons and fibers accumulate in the epithelium around the spermiduct openings. In contrast, *Gnrh1*-expressing neurons are localized in the cerebral ganglion and project toward the ovary, suggesting distinct roles of GnRH1 and GnRH2 in reproductive regulation. RNA-seq and real-time PCR analyses revealed that *Opsin2*, *Opsin3*, beta-carotene-15,15'-monooxygenase (*Bco*), and several ion channel genes are specifically expressed in the OPO, along with *Gnrh2*. *In situ* hybridization showed that these genes are localized in the innermost OPO layer, suggesting that the OPO functions as a photoreceptive organ. Although, *Gnrh2* expression has been considered low in adult *Ciona*, our study revealed strong OPO-specific expression with photoreceptive genes. Collectively, these findings suggest that GnRH2 plays a central role in gamete release regulation, potentially in coordination with the photoreceptive system, while GnRH1 may regulate ovarian functions. These results indicate that *C. robusta* employs distinct GnRH systems in a tissue-specific manner to regulate reproduction.

Introduction

Gamete release, an essential event for animals, is regulated by environmental stimuli and various biological factors including neural circuit and endocrine molecules. The mode and timing of gamete release is important especially for aquatic animals because the site of fertilization is in water. Light is one of the common environmental stimuli inducing gamete release. In jelly fish, light directly stimulates gonadal photosensory-neurosecretory cells which induce gamete maturation and release.[1] In fish, the timing of spawning is tightly connected with the lunar and diurnal cycles during the spawning season in pufferfish and the molecular mechanisms regulating spawning are well established.[2-5] In ascidians, the sister group of vertebrates, the relationship between light and gamete release was previously studied in *Ciona intestinalis* while interestingly, the timing of spawning is still controversial. Several papers reported that spawning generally takes place at sunrise.[6-8] On the other hand, it was reported that spawning took place during the evening and night-time hours.[9] Furthermore, the other paper reported that *Ciona* may spawn and settle at any time of the day.[10] In contrast to the controversy, continuous illumination prevents uncontrolled spawning,[11,12] suggesting that light is an essential factor inhibiting spawning behavior in *Ciona*.

Ciona is a hermaphroditic animal that releases both sperm and eggs. Sperm is released first from the distal end of sperm duct, where orange-colored cells accumulate to form a structure known as orange-pigmented organ (OPO).[13] Egg release occurs from the oviduct beneath the spermiduct knob-like structure at the OPO after a period of sperm release. Interestingly, orange-colored cells are typically observed in the OPO of *Ciona intestinalis* type A, which is distributed in the Mediterranean, northeast Atlantic, and Pacific regions, whereas in *Ciona intestinalis* type B, found in the north Atlantic region, pigmentation is restricted to the spermiduct.[14] Based on geographical distribution, morphological characteristics, and genetic information, *Ciona intestinalis* is now be reclassified into two distinct species: *Ciona robusta* (formerly *Ciona intestinalis* type B) and *Ciona intestinalis* (formerly *Ciona intestinalis* type A).[14,15]

Neuropeptides are important regulators of reproduction in animals. Among them, gonadotropin-releasing hormone (GnRH) plays a central role in the hypothalamus-pituitary-gonadal (HPG) axis in vertebrates. In tetrapods, GnRH is generally believed to promote gonadal maturation by stimulating the secretion of pituitary gonadotropins: follicle-stimulating hormone (FSH) and luteinizing hormone (LH).[16-18] In fish, GnRH has also been reported to regulate spawning behavior.[19,20] More recently, a "dual GnRH model" has been proposed in fish for the regulation of FSH and LH secretion, in which GnRH stimulates LH secretion, while cholecystokinin stimulates FSH secretion.[21] The origin of GnRH can be traced back to a common ancestor of bilaterians, forming a superfamily that includes GnRH, adipokinetic hormone, and corazonin, and acts as a multifunctional peptide in both protostomes and deuterostomes.[22-24] In the deuterostome lineage,

GnRH genes have been well conserved. In *Ciona*, two GnRH genes (*Gnrh1* and *Gnrh2*) and their encoded GnRH peptides, and four GnRH receptor genes have been identified.[25-29] These GnRH and receptor genes are thought to have emerged by lineage-specific duplication events.[30] GnRH plays essential roles during the development of *Ciona*, which are independent of the regulation of reproduction.[31-33] In adult *Ciona*, morphological analyses have shown that GnRH immunoreactive neurons and nerve fibers directly reach the ovary, suggesting that GnRH is involved in reproductive regulation.[34] Interestingly, GnRH injection into gonoduct induced gamete release, while the molecular mechanisms underlying this process remain to be clarified.[26,35] In addition, the relationship between the GnRH system and above-mentioned light/dark conditions that induce gamete release remains enigmatic.

In this study, we focused on *Ciona robusta* (hereafter referred to as *C. robusta*) and investigated the mechanisms regulating gamete release. Behavioral observations and histological analyses revealed that the sequential release of sperm and eggs occurs in this order because the knob-like structure on the spermiduct physically compresses the oviduct opening during the initial stage of gamete release. This structural characteristic suggests that *C. robusta* actively regulates sperm release but not egg release. Anatomical observations of transgenic *C. robusta* expressing Kaede under the control of the *Gnrh1* or *Gnrh2* promoter further revealed distinct morphological differences between GnRH1 and GnRH2 nervous systems: GnRH1 neurons in the cerebral ganglion project to the ovary, whereas GnRH2 neurons in the OPO project to the cerebral ganglion. Gene expression and histological analyses of the OPO further revealed that not only *Gnrh2* but also photoreception-related genes, including *Opsin2, Opsin3*, and *beta-carotene 15,15'-monooxygenase* (*Bco*), and several ion channel genes are specifically expressed in the OPO. These findings suggest that GnRH2 and the photoreceptive system may represent a missing link in the regulation of gamete release in *C. robusta*.

Materials and Methods

Animals

Wild adult *C. robusta* was used for behavioral and morphological observations, gene expression analyses, and histochemical studies. *C. robusta* was also used to generate transgenic lines. The CiGnRH1>*Kaede* lines and CiGnRH2>*Kaede* lines of *C. robusta*, Tg[MiCignrh1K]2, Tg[MiCignrh2K]1 and Tg[MiCignrh2K]2[36,37], which were created by Minos transposon-mediated transgenesis, were used for anatomical observations of the GnRH1 and GnRH2 nervous systems. No significant differences were observed between the Tg[MiCignrh2K]1 and Tg[MiCignrh2K]2 lines. The transgenic lines were cultured using an inland culture system as described previously[38].

Observation of spawning behavior

Wild *C. robusta* were placed in a 30-cm cubic tank and maintained under dark conditions for 30 minutes to 1 hour by covering the tank with a box draped with a blackout curtain. Spawning behavior was monitored under the dark conditions or after removing the box, and video recordings were captured using a digital camera (PENTAX K-70; RICOH IMAGING COMPANY, LTD., Tokyo, Japan), and an action camera (GoPro HERO13 Black; GoPro, Inc., San Mateo, CA, USA) with close-up lens (AOI UCL-03; Fisheye Co.,Ltd., Tokyo, Japan). For observations under a stereomicroscope, wild *C. robusta* were anesthetized using L-menthol according to a previously reported method[39,40]. Briefly, a 0.56% (weight/volume) L-menthol solution in ethanol was prepared and used as a stock solution. The stock solution was further diluted with 1% (volume/volume) artificial seawater before use. Animals were soaked in the diluted L-menthol solution for 10 minutes. After anesthetization, parts of the tunic, and body-wall muscle were excised, and the OPO was exposure under a stereomicroscope (ZEISS SteREO Discovery.V8; Carl Zeiss, Oberkochen, Germany). Gamete release was observed, and video recordings were captured using the stereomicroscope equipped with a USB digital camera (3R-DKMC04; 3R SOLUTION CORP., Fukuoka, Japan) and Anyty microscope adapter software ver. 9.0 (3R SOLUTION CORP.).

Whole-mount tissue preparation

Transgenic animals were anesthetized as described above. Parts of the tunic, body-wall muscle, and pharynx were excised under a fluorescence stereomicroscope (Leica M205 FA; Leica Microsystems, Wetzlar, Germany), and whole animals were fixed in 4% paraformaldehyde in phosphate buffer (PB) at 4 °C overnight. The fixed specimens were then soaked in 2 mg/ml glycine in PBS to quench the paraformaldehyde. Subsequently, the specimens were washed three times with PBS and used for morphological analyses. Anatomical analyses were performed under the stereomicroscope (Leica M205 FA; Leica Microsystems) using precision tweezers (Outils Rubis SA, Stabio, Switzerland) and microscissors (Inami & Co., Ltd., Tokyo, Japan).

Whole-mount observation

After removal of the tunic, fixed tissues were placed either in a glass dish with a thick silicone resin sheet or in a slide glass chamber (AGC TECHNO GLASS, Tokyo, Japan) filled with PBS and secured with needles as appropriate. The fluorescence stereomicroscope (Leica M205 FA; Leica Microsystems) and image acquisition software (Leica AF6000E; Leica Microsystems) were used for low-magnification observations.

Histological analysis of tissue sections of the OPO

The OPO tissues were fixed with Bouin's fixative solution at 4 °C overnight. The fixed tissues were

then soaked in 30% sucrose in PBS at 4 °C overnight. The tissues were embedded in Tissue-Tek

Cryomold #4565 (Sakura Finetek Japan Co., Ltd., Tokyo, Japan) using Super Cryoembedding

Medium-L1 (Leica Microsystems Japan, Tokyo, Japan). The tissues were subsequently frozen in

n☐hexane cooled to −196 °C with liquid nitrogen and stored at −80 °C until sectioning. Sections of

10 μm thickness were prepared using a CryoStar NX70 cryostat (Thermo Fisher Scientific, Inc.,

Waltham, MA) at −20 °C. Hematoxylin–eosin staining was performed following a modified version

of a previously reported method[40]. Briefly, the sections were washed with distilled water for

5 minutes and immersed in Mayer's hemalum solution for 10 seconds. They were then washed with

PBS for 2 minutes and distilled water for 1 minutes, followed by immersed in eosin solution for

2 minutes. After rinsing with distilled water, the sections were dehydrated through a graded ethanol

series (70%, 80%, 95%, and 100%) and xylene. Finally, the sections were mounted with a cover

glass (Matsunami Glass Ind., Ltd.) and MOUNT-QUICK (Daido Sangyo Co., Ltd., Tokyo, Japan),

and observed under a light microscope equipped with 20× or 40× objectives (Axio Imager 2; Carl

Zeiss, Oberkochen, Germany) in phase-contrast mode. OPO tissues of Tg[MiCignrh1K]2 line were

fixed in 4% PFA and sections of 10 μm thickness were prepared as described above. Sections were

washed three times with PBS for 5 minutes and mounted with a cover glass (Matsunami Glass Ind.,

Ltd.) and Fluoromount (Diagnostic BioSystems, Pleasanton, CA, USA). The sections were observed

under a fluorescent microscope with with 20× or 40× objectives (Axio Imager 2; Carl Zeiss), or a

confocal microscope equipped with 100× objectives (FLUOVIEW FV3000; Olympus, Tokyo,

Japan).

RNA-sequence analysis

Animals were anesthetized as described above. OPO, spermiduct, oviduct, and ovary were dissected

under a stereo microscope (ZEISS SteREO Discovery.V8; Carl Zeiss, Oberkochen, Germany), and

the tissues were snap-frozen in liquid nitrogen. Total RNA was extracted from each tissue using

Sepasol-RNA I (Nacalai Tesque, Kyoto, Japan). The extracted RNA was further purified and treated

with TURBO DNase as previously described.[41,42] RNA quality was assessed using a bioanalyzer

(Agilent 2100; Agilent, Santa Clara, CA, USA), confirming that the RNA integrity number (RIN) was 10.

RNA sequencing was conducted by Novogene Co., Ltd as contract research *via* NIPPON Genetics

Co, Ltd, and raw reads from each tissue were obtained. The resulting reads were aligned to *C.robusta*

genes (HT version, KY21 gene models) using the Hisat2 algorithm. Gene models were downloaded

as "HT.KY21Gene.2.fasta" from the Ghost Database

(http://ghost.zool.kyoto-u.ac.jp/default_ht.html). Gene expression levels were calculated as

transcripts per million (TPM). Genes that were expressed at levels more than tenfold higher in the

OPO compared to other tissues were identified, and their expression levels were further analyzed by
Real-time PCR.
200
Real-time PCR analysis
Total RNA was extracted from the cerebral ganglion, OPO, spermiduct, oviduct, and ovary as
described above. A 500-ng aliquot of DNase-treated total RNA was used for first-strand cDNA
synthesis. Real-time PCR was performed using a CFX96 Touch Real-Time PCR Detection System
(Bio-Rad Laboratories, Inc., Hercules, CA) as previously described[43] to validate the RNA-seq data.
The primers used for real-time PCR analyses are listed in Table 1. Gene expression levels were
normalized to the expression of the ubiquitin-associated domain containing 1 gene (*CiUbac1*,
KY21.Chr12.26), which was constitutively expressed among tissues according to the RNA-seq
analysis.
210
In situ **hybridization**
The RNA probes for *Opsin2* (XM_002119927), *Opsin3* (NM_001032468), and *Bco*
(NM_001078422) were designed to span exon-intron junctions. Approximately 300-bp fragments of
these genes were inserted into the pCR4-TOPO vector (Thermo Fisher Scientific Inc., Waltham, MA,
USA). The plasmids were linearized and used for in vitro transcription with a digoxigenin-labeled
RNA labeling kit (Roche Applied Science, Penzberg, Germany). OPO tissues were dissected and
fixed in 4% paraformaldehyde in 0.1 M phosphate buffer at 4 °C overnight. The fixed tissues were
then soaked in a refrigerated sucrose solution (30% sucrose in PBS) until they sank. They were
embedded in Super Cryoembedding Medium-L1 (Leica Microsystems Japan, Tokyo, Japan) and
sectioned at 10 µm thickness using a CryoStar NX70 cryostat (Thermo Fisher Scientific Inc.,
Waltham, MA, USA) at −20 °C. The sections were mounted onto MAS-coated slides (Matsunami
Glass Ind., Ltd., Osaka, Japan). Hybridization, washing, and detection were performed as previously
described.[43] No positive signals were observed when sense probes were used, confirming the
specificity of the hybridization. The primer sets used for probe preparation are listed in Table 1. In
tissue sections of the OPO, orange-pigmented cells were observed prior to *in situ* hybridization.
However, the pigmentation disappeared following the hybridization procedure. To accurately
determine the localization of *in situ* hybridization signals relative to the original pigmentation, we
captured microscopic images of the OPO both before and after *in situ* hybridization and overlaid the
images using Adobe Photoshop Elements 13 (Adobe Systems, San Jose, CA, USA).
230
231

Results

Observation of spawning behavior

Since continuous light is known to inhibit spawning in *C. robusta*, we hypothesize that dark

conditions would stimulate spawning behavior. To test this hypothesis, we placed *C. robusta* under

dark conditions and monitored spawning behavior. Spawning began approximately 30 minutes after

the onset of darkness (Movie 1). Another video recording revealed that *C. robusta* initiates spawning

by releasing sperm, followed by egg release (Movie 2). To examine this sequence more closely, we

conducted further observations under a stereomicroscope, focusing on the OPO. These observations

confirmed that sperm release precedes egg release (Movie 3). Notably, at the onset of spawning, a

knob-like structure on the spermiduct was observed to suppress the oviduct opening. As sperm

release progressed, the size of this knob gradually decreased, allowing eggs to begin moving toward

release (Movie 3).

Whole mount observations of transgenic animals

We examined the distribution of *Gnrh1*- and *Gnrh2*-promoter-driven Kaede-positive cells throughout

the *C. robusta* body. Figure 1 provides an overview of the distribution of Kaede-positive cells and

nerve fibers in the cerebral ganglion, dorsal strand plexus (DSP), OPO, anus, and ovary. In

Gnrh1>Kaede transgenic animals, Kaede-positive cells were observed in the cerebral ganglion and

DSP, with both cells and nerve fibers extending toward the ovary (Fig. 1A-D). In contrast, no

projections were observed toward the OPO or anus. These findings are consistent with previous

immunohistochemical studies of GnRH peptides[26,34]. In *Gnrh2>Kaede* transgenic animals,

Kaede-positive cells were not detected in the cerebral ganglion, but were present in the OPO and

anus (Fig. 1 E-G). Nerve fibers appear to project from the OPO to the cerebral ganglion (Fig. 1 A).

In *Gnrh2>Kaede* transgenic animals, nerve fibers were not observed projecting to the ovary. Weak

nerve projections were detected in the oral siphon in both *Gnrh1>Kaede* and *Gnrh2>Kaede* animals,

whereas nerve fibers projected to the atrial siphon only in *Gnrh2>Kaede* animals (Supplemental Fig.

1). Whole-mount observations revealed distinct patterns of cellular and fiber distribution between

Gnrh1 and *Gnrh2* transgenic animals. Interestingly, while *Gnrh2* expression in adult *C. robusta* has

been considered to be very low[44], the present observations suggest specific expression of *Gnrh2* in

the OPO and anus, which have not been examined in previous studies.

Histological analysis of tissue sections of the OPO

Hematoxylin-eosin staining revealed the basic structures of the OPO, spermiduct, and oviduct. The

OPO is composed of several papillae, which serve as outlet ports for sperm. These papillae are filled

with orange-pigmented cells and are entirely covered by an epithelium layer as reported previously[45]

(Fig. 2). The spermiduct terminates at the OPO, and sperm accumulate in a structure referred to as

the spermiduct knob-like structure in this study. The oviduct lies adjacent to the spermiduct, and its

opening seems to be pressed down by the spermiduct knob-like structure (Fig. 2). Orange-pigmented

cells were observed beneath the epithelial cells, in a region where hemocytes circulate. As shown in

Figure 1F, Kaede-positive cells were observed in the OPO of *Gnrh2>Kaede* animals. To further

investigate their localization, we examined tissue sections for the distribution of Kaede-positive cells.

Fluorescence microscopy revealed that Kaede-positive cells are located within the epithelial layer

surrounding the outlet ports of the OPO (Fig. 3). Confocal microscopy further demonstrated that

these Kaede-positive cells are neurons, with observable nerve fiber projections (Fig. 3D).

RNA sequencing and real-time PCR

To investigate gene expression in the OPO, we performed RNA sequencing (RNA-seq) on four

tissues: OPO, spermiduct, oviduct, and ovary. RNA-seq yielded approximately 40 to 50 million

paired-end reads per tissue. For each dataset, approximately 50,000 transcripts were successfully

mapped. The resulting FASTQ files were deposited in the Sequence Read Archive (SRA) under the

following accession number: PRJNA1279780. Genes expressed at levels more than tenfold higher in

the OPO than in the other tissues were identified and listed. Notably, *Gnrh2* was among the top 100

OPO-specific genes (Supplemental Table 1), consistent with our histological findings based on

Gnrh2 promoter-driven Kaede fluorescence in the OPO of transgenic animals. Interestingly, several

photoreception-related genes including *Opsin2*, *Opsin3*, and beta-carotene-15,15'-monooxygenase

(*Bco*), were specifically expressed in the OPO (Supplemental Table 1). In addition, multiple ion

channel-related genes were expressed abundantly in the OPO, including potassium/sodium

hyperpolarization-activated cyclic nucleotide-gated channel 2 (*Hcn2*)-like, sodium channel protein

type 4 subunit alpha B (*Scn4b*)-like, potassium voltage-gated channel subfamily H member 8

(*Kcnh8*)-like, and voltage-dependent T-type calcium channel subunit alpha-1G (*Cacna1*)

(Supplemental Table 1). We further examined the expression of these genes by quantitative real-time

PCR (qRT-PCR) in the cerebral ganglion, OPO, spermiduct, oviduct, and ovary and found that their

expression was almost exclusive in the OPO (Fig. 4). We also analyzed the expression of GnRH

receptor genes, including *Gnrhr1* through *Gnrhr4*. In contrast to the OPO-specific genes, the GnRH

receptor genes showed relatively high expression in the cerebral ganglion and oviduct, rather than in

the OPO (Fig. 4).

***In situ* hybridization**

To examine the localization of OPO-specific genes, we performed *in situ* hybridization on OPO

tissue sections. As shown in Figure 2, orange-pigmented cells were observed beneath the epithelial

layer. *In situ* hybridization signals were detected in a restricted region of the epithelial cells, where

those cells are adjacent to the sperm in the papilla of the OPO (Fig. 5). No signal was observed at the

distal tip of the OPO opening. Interestingly, all examined genes, including photoreception-related

genes and ion channels examined were localized to the same epithelial region (Fig. 5). No

hybridization signal was detected using the sense probe, confirming the specificity of the observed

signals. Some blue staining was observed in the outer region of the epithelium, which was

considered nonspecific (Fig. 5).

Discussion

In ascidians, *C. intestinalis* and *C. robusta*, which were previously considered to be the same species,

have both been used to study spawning behavior. Interestingly, several studies have reported

differing light/dark conditions that stimulate spawning in *Ciona*. Some have shown that spawning

generally occurs around sunrise.[6-8] Conversely, other studies have reported spawning during the

evening and night-time hours.[9] Furthermore, one report suggested *Ciona* may spawn and settle at

any time of the day.[10] In our observations, *C. robusta* began spawning approximately 30 minutes

after the onset of darkness under laboratory conditions. These results support the view that spawning

occurs in *Ciona* occurs during night-time hours.[9] Given that many earlier studies did not distinguish

between *C. intestinalis* and *C. robusta*, this lack of taxonomic resolution may have contributed to the

discrepancies in previous findings. In light of our results, it may be necessary to re-evaluate the

relationship between light–dark stimuli and spawning behavior, particularly in *C. intestinalis*. Until

now, the sequence of sperm and egg release during spawning in *C. robusta* has not been thoroughly

investigated. In this study, we demonstrated that egg release is physically delayed due to the

anatomical arrangement of the spermiduct and oviduct openings, resulting in the release of sperm

occurring first. These findings suggest that sperm release in *C. robusta* is actively regulated. In

contrast, egg release appears to be passively controlled, suggesting that *Ciona* employs a dexterous

strategy which can control the timing of sperm and egg release strictly with the minimum biological

cost of active regulation.

In *Ciona*, two gonadotropin-releasing hormone homolog genes, *Gnrh1* and *Gnrh2* have been

identified.[26] Both genes are expressed during embryonic development and in the larval stage.[26,44] In

adult animals, *Gnrh1* is expressed in the neural complex, including the cerebral ganglion, whereas

Gnrh2 is not.[44] Immunohistochemical studies revealed that GnRH-positive cells are detected in the

cerebral ganglion and they project fibers, along with the DSP, to the ovary,[26,34] suggesting that the

GnRH system plays a regulatory role in ovarian functions. Our whole-mount observations of

transgenic animals further revealed that the GnRH1-expressing cells are localized in the cerebral

ganglion and project nerve fibers to the ovary, supporting the idea that the GnRH1 nervous system

may regulate ovarian functions. Interestingly, our observations also showed that GnRH2-expressing

cells are localized in the OPO. *Gnrh2*-positive neurons are situated near the outlet ports of the OPO

and extend projections to the cerebral ganglion, suggesting that GnRH2 nervous system may be

involved in transmitting sperm-release signals to the cerebral ganglion, or in suppressing sperm

release under light conditions. To further investigate the role of *Gnrh2* in spawning behavior, we are

currently conducting gene knockout experiments.

The present study found key molecules associated with spawning behavior and light/dark

conditions. Two opsin genes, *Opsin2* and *Opsin3*, were specifically expressed in the OPO. In a

previous phylogenetic analysis, *Opsin2* was grouped within the clade of vertebrate pigments and

related opsins, while *Opsin3* was clustered with the human retinal G protein-coupled receptor (RGR),

a non-visual opsin that can bind and photoisomerize all-trans-retinal to 11-cis-retinal.[46] In addition,

Bco was also expressed in the OPO. All these photoreception-related genes were localized in the

innermost region of the OPO, where they are situated adjacent to the orange-pigmented cells. Since

BCO and *RGR* are expressed in the reginal pigment epithelium (RPE) of mammalian eyes,[47,48] the

innermost region of the OPO may be similar to the RPE in mammals, although it does not form a

single-cell layer as in the mammalian RPE.[49] It is particularly interesting that *Opsin2*, a

photoreceptor gene, was expressed in the same region as *Bco* and *Opsin3*, given that in mammalian

eyes, RPE cells are functionally or physically separated from photoreceptor cells.[48] Although *Ciona*

is positioned as the sister group of vertebrates and its organs are less differentiated, these findings

suggests that the OPO functions as a photoreceptive organ involved in spawning behavior in

response to light/dark cues. In *Ciona* larvae, *Opsin1* a paralog of *Opsin2*, is expressed in the outer

segment of photoreceptor cells, whereas *Opsin2* is not found in the ocellus photoreceptor cells but

instead is expressed in the brain vesicle and oral siphon rudiment.[46] *Opsin3* is localized in both the

ocellus photoreceptor cells and surrounding non-photoreceptor cells in the brain vesicle.[46,50] These

photoreceptive systems in *Ciona* larvae are important for the regulation of swimming behavior,[51,52]

and likely contribute to the regulation of spawning behavior in adult *Ciona*, although the

developmental transition of this system remains to be elucidated.

In this study, we also focused on the ion channels in the RNA-seq data, as they are known to

play important roles in visual function in vertebrates.[53,54] Several putative ion channel genes,

including *Hcn2*-like, *Scn4b*-like, *Kcnh8*-like, and *Cacna1*, were specifically expressed in the same

region of the photoreceptive site in the OPO. Although we have not yet conducted double-staining

analyses, photoreceptor-related genes and ion channel genes appear to be co-expressed in the same

cells. In mammals, phototransduction pathway involved in non-image-forming vision depends on

HCN channels.[54] In retinoblastoma cells, *CACNA1G* is expressed but is downregulated after

differentiation.[55] These findings suggest that *Hcn2*-like and *Cacna1* may be involved in

phototransduction in the OPO in a manner similar to their roles in mammals. In contrast, to the best

of our knowledge, there is no evidence that *Scn4b* and *Kcnh8* participate in vertebrate photoreceptor

systems. These two ion channels may therefore have *Ciona*-specific roles within the photoreceptive

system in the OPO.

In summary, this study revealed that spawning behavior in *C. robusta* is regulated by

light/dark conditions, with spawning initiated after the onset of darkness. Morphological, gene

expression, and histological analyses indicate that the OPO functions as a photoreceptive organ, and

that the GnRH2, rather than GnRH1, neural system participates in the regulation of spawning

behavior. Further investigations will be necessary to elucidate potential interactions among GnRH2,

photoreception-related genes, and ion channels in the context of gamete release regulation in the

OPO. Functional analyses of the identified genes and neural circuits are currently in progress and

will be essential for fully understanding the mechanism of light-mediated spawning in ascidians.

Acknowledgments

We gratefully acknowledge Prof. Fumihiko Sato for his fruitful comments on the manuscript. We

thank the National Bio-Resource Project (NBRP) for providing ascidians. This study was supported

by JSPS Grants-in-Aid for Scientific Research (KAKENHI); Grant Numbers 22K06327 and

25K09712 to T.O., and 23K27185 to T.G.K.

Author contributions

T.O. designed the study; T.O., S.M., A.S., A.W., Y.M., I.S.S., Y.S., T.G.K., and H.S. performed the

research; T.O., S.M., A.S., Y.S., T.G.K., and H.S. wrote, reviewed, and edited the paper. Y.S.

produced and provided transgenic animals *via* NBRP.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to T.O.

References

1. Quiroga Artigas, G., Lapebie, P., Leclere, L., Takeda, N., Deguchi, R., Jekely, G., Momose, T., and Houliston, E. (2018). A gonad-expressed opsin mediates light-induced spawning in the jellyfish Clytia. Elife *7*. 10.7554/eLife.29555.

2. Motohashi, E., Hamabata, T., and Ando, H. (2008). Structure of neurohypophysial hormone genes and changes in the levels of expression during spawning season in grass puffer (Takifugu niphobles). Gen. Comp. Endocrinol. *155*, 456-463. 10.1016/j.ygcen.2007.07.009.

3. Ando, H., Shahjahan, M., and Hattori, A. (2013). Molecular neuroendocrine basis of lunar-related spawning in grass puffer. Gen. Comp. Endocrinol. *181*, 211-214. 10.1016/j.ygcen.2012.07.027.

4. Ando, H., Shahjahan, M., and Kitahashi, T. (2018). Periodic regulation of expression of genes for kisspeptin, gonadotropin-inhibitory hormone and their receptors in the grass puffer: Implications in seasonal, daily and lunar rhythms of reproduction. Gen. Comp. Endocrinol. *265*, 149-153. 10.1016/j.ygcen.2018.04.006.

5. Chen, J., Katada, Y., Okimura, K., Yamaguchi, T., Guh, Y.J., Nakayama, T., Maruyama, M., Furukawa, Y., Nakane, Y., Yamamoto, N., et al. (2022). Prostaglandin E(2) synchronizes lunar-regulated beach spawning in grass puffers. Curr. Biol. *32*, 4881-4889 e4885. 10.1016/j.cub.2022.09.062.

6. Thorson, G. (1964). Light as an ecological factor in the dispersal and settlement of larvae of marine bottom invertebrates. Ophelia *1*, 167-208. 10.1080/00785326.1964.10416277.

7. Millar, R.H. (1971). The Biology of Ascidians. Adv. Mar. Biol. *9*, 1-100.

8. Svane, I., and Young, C.M. (1989). The ecology and behaviour of ascidian larvae. Oceanography and Marine Biology *27*, 45-90.

9. Yamaguchi, M. (1971). Spawning periodicity and settling time in ascidians, Ciona intestinalis and Sryela plicata. Rec. Oceanogr. Wks. *10*, 147-155.

10. Svane, I., and Havenhand, J.N. (1993). Spawning and Dispersal in Ciona intestinalis (L.). Marine Ecology *14*, 53-66. https://doi.org/10.1111/j.1439-0485.1993.tb00364.x.

11. Lambert, C.C., and Brandt, C.L. (1967). The Effect of Light on the Spawning of Ciona Intestinalis. Biol. Bull. *132*, 222-228. 10.2307/1539890.

12. Yamaguchi, A., Saito, T., Yamada, L., Taniguchi, H., Harada, Y., and Sawada, H. (2011). Identification and localization of the sperm CRISP family protein CiUrabin involved in gamete interaction in the ascidian Ciona intestinalis. Mol. Reprod. Dev. *78*, 488-497. 10.1002/mrd.21329.

13.   Ohta, N., Horie, T., Satoh, N., and Sasakura, Y. (2010). Transposon-mediated enhancer
detection reveals the location, morphology and development of the cupular organs,
which are putative hydrodynamic sensors, in the ascidian Ciona intestinalis. Zoolog. Sci.
27, 842-850. 10.2108/zsj.27.842.

14.   Caputi, L., Andreakis, N., Mastrototaro, F., Cirino, P., Vassillo, M., and Sordino, P.
(2007). Cryptic speciation in a model invertebrate chordate. Proc. Natl. Acad. Sci. U. S.
A. *104*, 9364-9369. 10.1073/pnas.0610158104.

15.   Brunetti, R., Gissi, C., Pennati, R., Caicci, F., Gasparini, F., and Manni, L. (2015).
Morphological evidence that the molecularly determined *Ciona intestinalis* type A and
type B are different species: *Ciona robusta* and *Ciona intestinalis*. Journal of Zoological
Systematics and Evolutionary Research *53*, 186-193. https://doi.org/10.1111/jzs.12101.

16.   Coss, D. (2018). Regulation of reproduction via tight control of gonadotropin hormone
levels. Mol. Cell. Endocrinol. *463*, 116-130. 10.1016/j.mce.2017.03.022.

17.   Tsutsumi, R., and Webster, N.J. (2009). GnRH pulsatility, the pituitary response and
reproductive dysfunction. Endocr. J. *56*, 729-737. 10.1507/endocrj.k09e-185.

18.   Hasunuma, I. (2024). Central regulation of reproduction in amphibians. J Exp Zool A
Ecol Integr Physiol *341*, 219-229. 10.1002/jez.2769.

19.   Ando, H., and Urano, A. (2005). Molecular regulation of gonadotropin secretion by
gonadotropin-releasing hormone in salmonid fishes. Zoolog. Sci. *22*, 379-389.
10.2108/zsj.22.379.

20.   Zohar, Y., Zmora, N., Trudeau, V.L., Munoz-Cueto, J.A., and Golan, M. (2022). A half
century of fish gonadotropin-releasing hormones: Breaking paradigms. J.
Neuroendocrinol. *34*, e13069. 10.1111/jne.13069.

21.   Uehara, S.K., Nishiike, Y., Maeda, K., Karigo, T., Kuraku, S., Okubo, K., and Kanda, S.
(2024). Identification of the FSH-RH as the other gonadotropin-releasing hormone. Nat
Commun *15*, 5342. 10.1038/s41467-024-49564-8.

22.   Fodor, I., Osugi, T., Matsubara, S., Shiraishi, A., Kawada, T., Pirger, Z., and Satake, H.
(2024). Characterization of corazonin signaling in a molluscan model species, Lymnaea
stagnalis. Gen. Comp. Endocrinol. *357*, 114594. 10.1016/j.ygcen.2024.114594.

23.   Zandawala, M., Tian, S., and Elphick, M.R. (2018). The evolution and nomenclature of
GnRH-type and corazonin-type neuropeptide signaling systems. Gen. Comp.
Endocrinol. *264*, 64-77. 10.1016/j.ygcen.2017.06.007.

24.   Sakai, T., Shiraishi, A., Kawada, T., Matsubara, S., Aoyama, M., and Satake, H. (2017).
Invertebrate Gonadotropin-Releasing Hormone-Related Peptides and Their Receptors:
An Update. Front. Endocrinol. (Lausanne) *8*, 217. 10.3389/fendo.2017.00217.

25.   Powell, J.F., Reska-Skinner, S.M., Prakash, M.O., Fischer, W.H., Park, M., Rivier, J.E.,
Craig, A.G., Mackie, G.O., and Sherwood, N.M. (1996). Two new forms of
gonadotropin-releasing hormone in a protochordate and the evolutionary implications.
Proc. Natl. Acad. Sci. U. S. A. *93*, 10461-10464. 10.1073/pnas.93.19.10461.

26.   Adams, B.A., Tello, J.A., Erchegyi, J., Warby, C., Hong, D.J., Akinsanya, K.O., Mackie,
G.O., Vale, W., Rivier, J.E., and Sherwood, N.M. (2003). Six novel
gonadotropin-releasing hormones are encoded as triplets on each of two genes in the
protochordate, Ciona intestinalis. Endocrinology *144*, 1907-1919.
10.1210/en.2002-0216.

27.   Kusakabe, T., Mishima, S., Shimada, I., Kitajima, Y., and Tsuda, M. (2003). Structure,
expression, and cluster organization of genes encoding gonadotropin-releasing hormone
receptors found in the neural complex of the ascidian Ciona intestinalis. Gene *322*,
77-84. 10.1016/j.gene.2003.08.013.

28.   Tello, J.A., Rivier, J.E., and Sherwood, N.M. (2005). Tunicate gonadotropin-releasing
hormone (GnRH) peptides selectively activate Ciona intestinalis GnRH receptors and
the green monkey type II GnRH receptor. Endocrinology *146*, 4061-4073.
10.1210/en.2004-1558.

29.   Kawada, T., Ogasawara, M., Sekiguchi, T., Aoyama, M., Hotta, K., Oka, K., and Satake,
H. (2011). Peptidomic analysis of the central nervous system of the protochordate,
Ciona intestinalis: homologs and prototypes of vertebrate peptides and novel peptides.
Endocrinology *152*, 2416-2427. 10.1210/en.2010-1348.

30.   Kawada, T., Aoyama, M., Sakai, T., and Satake, H. (2013). Structure, Function, and
Evolutionary Aspects of Invertebrate GnRHs and their Receptors. In
Gonadotropin-releasing hormone (GnRH), S.S. Eric, ed. (Nova Science Publishers, Inc),
pp. 1-15.

31.   Kusakabe, T.G., Sakai, T., Aoyama, M., Kitajima, Y., Miyamoto, Y., Takigawa, T.,
Daido, Y., Fujiwara, K., Terashima, Y., Sugiuchi, Y., et al. (2012). A conserved
non-reproductive GnRH system in chordates. PLoS One *7*, e41955.
10.1371/journal.pone.0041955.

32.   Kamiya, C., Ohta, N., Ogura, Y., Yoshida, K., Horie, T., Kusakabe, T.G., Satake, H., and
Sasakura, Y. (2014). Nonreproductive role of gonadotropin-releasing hormone in the
control of ascidian metamorphosis. Dev. Dyn. *243*, 1524-1535. 10.1002/dvdy.24176.

33.   Hozumi, A., Matsunobu, S., Mita, K., Treen, N., Sugihara, T., Horie, T., Sakuma, T.,
Yamamoto, T., Shiraishi, A., Hamada, M., et al. (2020). GABA-induced GnRH release
triggers chordate metamorphosis. Curr. Biol. *30*, 1555-1561 e1554.
10.1016/j.cub.2020.02.003.

34.   Tsutsui, H., Yamamoto, N., Ito, H., and Oka, Y. (1998). GnRH-immunoreactive
neuronal system in the presumptive ancestral chordate, Ciona intestinalis (Ascidian).
Gen. Comp. Endocrinol. *112*, 426-432. 10.1006/gcen.1998.7160.

35.   Terakado, K. (2001). Induction of gamete release by gonadotropin-releasing hormone in
a protochordate, Ciona intestinalis. Gen. Comp. Endocrinol. *124*, 277-284.
10.1006/gcen.2001.7728.

36.   Sasakura, Y., Awazu, S., Chiba, S., and Satoh, N. (2003). Germ-line transgenesis of the
Tc1/mariner superfamily transposon Minos in *Ciona intestinalis*. Proc. Natl. Acad. Sci.
U. S. A. *100*, 7726-7730. 10.1073/pnas.1230736100.

37.   Matsuoka, T., Awazu, S., Shoguchi, E., Satoh, N., and Sasakura, Y. (2005). Germline
transgenesis of the ascidian *Ciona intestinalis* by electroporation. Genesis *41*, 67-72.
10.1002/gene.20096.

38.   Joly, J.S., Kano, S., Matsuoka, T., Auger, H., Hirayama, K., Satoh, N., Awazu, S.,
Legendre, L., and Sasakura, Y. (2007). Culture of *Ciona intestinalis* in closed systems.
Dev. Dyn. *236*, 1832-1840. 10.1002/dvdy.21124.

39.   Osugi, T., Sasakura, Y., and Satake, H. (2017). The nervous system of the adult ascidian
Ciona intestinalis Type A (*Ciona robusta*): Insights from transgenic animal models.
PLoS One *12*, e0180227. 10.1371/journal.pone.0180227.

40.   Osugi, T., Sasakura, Y., and Satake, H. (2020). The ventral peptidergic system of the
adult ascidian *Ciona robusta* (*Ciona intestinalis* Type A) insights from a transgenic
animal model. Sci. Rep. *10*, 1892. 10.1038/s41598-020-58884-w.

41.   Matsubara, S., Shiraishi, A., Osugi, T., Kawada, T., and Satake, H. (2019). The
regulation of oocyte maturation and ovulation in the closest sister group of vertebrates.
Elife *8*. 10.7554/eLife.49062.

42.   Matsubara, S., Osugi, T., Shiraishi, A., Wada, A., and Satake, H. (2021). Comparative
analysis of transcriptomic profiles among ascidians, zebrafish, and mice: Insights from
tissue-specific gene expression. PLoS One *16*, e0254308.
10.1371/journal.pone.0254308.

43.   Osugi, T., Miyasaka, N., Shiraishi, A., Matsubara, S., and Satake, H. (2021). Cionin, a
vertebrate cholecystokinin/gastrin homolog, induces ovulation in the ascidian *Ciona*
intestinalis type A. Sci. Rep. *11*, 10911. 10.1038/s41598-021-90295-3.

44.   Kawada, T., Aoyama, M., Okada, I., Sakai, T., Sekiguchi, T., Ogasawara, M., and
Satake, H. (2009). A novel inhibitory gonadotropin-releasing hormone-related
neuropeptide in the ascidian, Ciona intestinalis. Peptides *30*, 2200-2205.
10.1016/j.peptides.2009.08.014.

45.   Tajima, Y., Hozumi, A., Yoshida, K., Treen, N., Sakuma, T., Yamamoto, T., and
Sasakura, Y. (2020). Hox13 is essential for formation of a sensory organ at the terminal
end of the sperm duct in Ciona. Dev. Biol. *458*, 120-131. 10.1016/j.ydbio.2019.10.028.
46.   Kusakabe, T., and Tsuda, M. (2007). Photoreceptive systems in ascidians. Photochem.
Photobiol. *83*, 248-252. 10.1562/2006-07-11-IR-965.
47.   Redmond, T.M., Gentleman, S., Duncan, T., Yu, S., Wiggert, B., Gantt, E., and
Cunningham, F.X., Jr. (2001). Identification, expression, and substrate specificity of a
mammalian beta-carotene 15,15'-dioxygenase. J. Biol. Chem. *276*, 6560-6565.
10.1074/jbc.M009030200.
48.   Radu, R.A., Hu, J., Peng, J., Bok, D., Mata, N.L., and Travis, G.H. (2008). Retinal
pigment epithelium-retinal G protein receptor-opsin mediates light-dependent
translocation of all-trans-retinyl esters for synthesis of visual chromophore in retinal
pigment epithelial cells. J. Biol. Chem. *283*, 19730-19738. 10.1074/jbc.M801288200.
49.   Yang, S., Zhou, J., and Li, D. (2021). Functions and Diseases of the Retinal Pigment
Epithelium. Front. Pharmacol. *12*, 727870. 10.3389/fphar.2021.727870.
50.   Tsuda, M., Kusakabe, T., Iwamoto, H., Horie, T., Nakashima, Y., Nakagawa, M., and
Okunou, K. (2003). Origin of the vertebrate visual cycle: II. Visual cycle proteins are
localized in whole brain including photoreceptor cells of a primitive chordate. Vision
Res. *43*, 3045-3053. 10.1016/j.visres.2003.09.012.
51.   Tsuda, M., Kawakami, I., and Shiraishi, S. (2003). Sensitization and habituation of the
swimming behavior in ascidian larvae to light. Zoolog. Sci. *20*, 13-22.
10.2108/zsj.20.13.
52.   Salas, P., Vinaithirthan, V., Newman-Smith, E., Kourakis, M.J., and Smith, W.C. (2018).
Photoreceptor specialization and the visuomotor repertoire of the primitive chordate
Ciona. J. Exp. Biol. *221*. 10.1242/jeb.177972.
53.   Rashwan, R., Hunt, D.M., and Carvalho, L.S. (2021). The role of voltage-gated ion
channels in visual function and disease in mammalian photoreceptors. Pflugers Arch.
473, 1455-1468. 10.1007/s00424-021-02595-2.
54.   Jiang, Z., Yue, W.W.S., Chen, L., Sheng, Y., and Yau, K.W. (2018). Cyclic-Nucleotide-
and HCN-Channel-Mediated Phototransduction in Intrinsically Photosensitive Retinal
Ganglion Cells. Cell *175*, 652-664 e612. 10.1016/j.cell.2018.08.055.
55.   Hirooka, K., Bertolesi, G.E., Kelly, M.E., Denovan-Wright, E.M., Sun, X., Hamid, J.,
Zamponi, G.W., Juhasz, A.E., Haynes, L.W., and Barnes, S. (2002). T-Type calcium
channel alpha1G and alpha1H subunits in human retinoblastoma cells and their loss
after differentiation. J. Neurophysiol. *88*, 196-205. 10.1152/jn.2002.88.1.196.
584

Figure legends

Figure 1. Whole-mount observations of transgenic *C. robusta* expressing Kaede under the

control of either the *Gnrh1* or *Gnrh2* promoter. Images were taken using a fluorescence stereo

microscope. The names of transgenic lines are indicated at the top-right corner of each panel. (A, B)

Cerebral ganglion of *Gnrh1* and *Gnrh2* transgenic lines. (C, D) Dorsal strand plexus (DSP) and

orange-pigmented organ (OPO) of *Gnrh1* and *Gnrh2* transgenic lines. (E, F) Anus of *Gnrh1* and

Gnrh2 transgenic lines. (G, H) Ovary and DSP of *Gnrh1* and *Gnrh2* transgenic lines. Scale bars: 500

µm in A, C, E, F, G, and H; 250 µm in B and D.

Figure 2. HE staining of the OPO tissue section. The spermiduct terminates at the OPO, which

serves as the outlet port for sperm. The OPO consists of multiple papillae filled with

orange-pigmented cells. The oviduct terminates beneath the knob-like structure of the spermiduct,

and its openings are compressed by this structure (arrowheads). Scale bar: 200 µm.

Figure 3. Morphological observations of the OPO section of *Gnrh2* transgenic line

(Tg[MiCignrh2K]2). (A) Bright-field image of papillae in the OPO. (B) Dark-field image of the

OPO showing *Gnrh2*-positive cells. (C) Merged image of the bright-field and dark-field images.

Gnrh2-positive cells are observed around the outlet ports of the OPO. (D) Confocal microscopic

image showing *Gnrh2*-positive cells located in the epithelium (EP), adjacent to the

orange-pigmented cells (OP). Nerve fibers are also visible. Scale bars: 50 µm in A, B, and C; 20 µm

in D.

Figurer 4. Real-time PCR analysis in the cerebral ganglion, OPO, spermiduct, oviduct, and

ovary. Relative expression levels of target genes are shown for each tissue. Data are presented as

means ± SEM (n = 3).

Figure 5. *In situ* hybridization of OPO-specific genes. The upper panels show antisense probe

signals for each gene, and lower panels show the corresponding sense controls. mRNA expression is

indicated by purple staining. (A, B) *Opsin2*; (C, D) *Opsin3*; (E, F) *Bco*; (G, H) *Kcnh8*-like isoform

X1; (I, J) *Hcn2*-like; (K, L) *Cac1g* isoform X2; (M, N) *Scn4ab*-like isoform X4. Scale bars: 50 µm.

Supplemental Figure 1. Whole-mount observations of transgenic *C. robusta* expressing Kaede

under the control of either the *Gnrh1* or *Gnrh2* promoter. Images were taken using a

fluorescence stereo microscope. The names of transgenic lines are indicated at the top-right corner of

each panel. The right panels show dark-field images, and the left panels show merged images of bright-field and dark-field views. (Aa, Ab) Oral siphon of *Gnrh1* transgenic line. (Ac, Ad) Oral siphon of *Gnrh2* transgenic line. (Be, Bf) Atrial siphon of *Gnrh1* transgenic line. (Bg, Bh) Atrial siphon of *Gnrh2* transgenic line. Scale bars: 1 mm.

Movies

Movie 1, 2, and 3 are available in the Zenodo repository under the following doi number: 10.5281/zenodo.15734177.

632 **Table 1. Primer sets used in this study.**

Gene name	Accession no.	Forward	Reverse
Real time PCR			
GnRH2	NM_001032628	GGGAGCAGAAGAAAGCGGAT	TTGTAGCGACCGAAGTCACC
Opsin2	XM_002119927	TTGGTGCCGTGCCAGTATTT	CAACTGTCGTTACCCAGCCC
Opsin3	NM_001032468	TCGCTGTTCGTGGAGAAGAC	AAGCAAACTGTAGCCGGTGA
Bco	NM_001078422	GGTTCAGATTTTGCTGAATAGGCA	GTAGCAAGCGCAGTCAAGTG
GnRHR1	NM_001033825	TTCAGTTCGACACGCTTCAC	CTGCCCCTTAGTGACCAGAG
GnRHR2	NM_001033824	CTTCATGGTTGTGGTGATCG	GCAGCAAGAACCATGCAGTA
GnRHR3	NM_001171881	CTGAGAAGTTCGGGTCTTGC	TTATAAAGGGGACGCTGGTG
GnRHR4	NM_001171880	CTCCAATCTTCTGGCTTTCG	CGAATGTGATGGTGATGAGC
Kcnh8	XP_009859607.1	ACCATGGTTTGTCATGCTCA	TTCAACTTCCGAGTGGTTCC
Hcn2	XM_009863621.3	GTCCACCAGCCATATGCTTT	GGTAGTTTGCGGAATTGCAT
Scn4ab	XP_026692361.1	ACGTTTTCACCGCCATCTAC	GAGCGAACGATGGCTTTTAG
Cacna1g	XP_018667819.1	CCACTTTATTTGCCCTGGAA	GCTGGAGAGCTGGCATAAAC
Ubac1	XM_018815256	CCACCTAGTGTTGGACAACG	TCATGCGAGTCGTCCACCTT
Probes for *in situ* hybridization			
Opsin2	XM_002119927	GGTTTGGTCGGCGGTTTCAT	CAGCTTGTCCCAAGTCCTTC
Opsin3	NM_001032468	CTGCTCTCTCTTGCAACCAG	CAGTTTCGAATCCATATCGC
Bco	NM_001078422	CTGAATGGTAGCTGGTACCG	GGCGAACAGATGATCACCGA
Kcnh8	XP_009859607.1	CAATGGACACACTTAACCCAC	CTTCGTCTTCTACGCCTGCT
Hcn2	XM_009863621.3	GTCCACCAGCCATATGCTTT	AGATCCGGTGACGTAAGCCA
Scn4ab	XP_026692361.1	CTATTCGGCCAGGAAAGTCG	CCAGCATTGTGGAGCGAACG
Cacna1g	XP_018667819.1	ATCATAATCAGCCCACAGGG	CCACGTTGTCCATAGTTTTCA

Outlet port of sperm

Epithelium

Orange-pigmented cells

Spermiduct

Spermiduct knob-like structurue

Oviduct

Egg

200 µm

Opsin2

Opsin3

Bco

Kcnh8-like isoform X1

Hcn2-like

Cac1g isoform X2

Scn4ab-like isoform X4