Selection inference in a complex genomic landscape: the role of inversions in

the spruce bark beetle

Julia Morales-García[1,2], Wiesław Babik[1], Piotr Zieliński[1], Krystyna Nadachowska-Brzyska[1*]

[1]Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University,

Gronostajowa 7, 30-387, Kraków, Poland

[2]Doctoral School of Exact and Natural Sciences, Jagiellonian University, ul. Łojasiewicza 11,

30-348, Kraków, Poland

[*]Corresponding author: Krystyna Nadachowska-Brzyska, krystyna.nadachowska@uj.edu.pl

Abstract

Understanding the genetic basis of adaptation is a key objective in evolutionary biology. Although advances in genomic selection scans have greatly improved our ability to detect signatures of adaptation, distinguishing true signals from false positives remains challenging. This task is particularly difficult in regions of reduced recombination, such as polymorphic inversions. In this study, we examine the genome-wide landscape of selection in the European spruce bark beetle (*Ips typographus*), Europe's most destructive forest pest, which has one of the most complex inversion-associated recombination landscapes. Using whole-genome resequencing data from 312 individuals across 23 populations, we applied two complementary selection scan methods to assess how structural genomic variants, such as inversions, affect the detection of adaptive signals. Our findings reveal that large polymorphic inversions are significantly enriched for selection signals, thereby reinforcing their pivotal role in adaptation. However, these same inversions can complicate the interpretation of selection signals within their boundaries, where recombination suppression can obscure the true targets of selection. In contrast, the detection of selection in collinear regions is largely unaffected by the presence of inversions. To overcome the confounding effects within inversions, we demonstrate that focusing on homozygote genotypes enables the identification of candidate regions and haplotype-specific signals. Our findings highlight the contribution of inversions to adaptation, emphasising the importance of understanding species-specific genomic architecture when interpreting selection scan results.

Key words: inversion-rich genomes, spruce bark beetle, selection scans, recombination suppression

Introduction

Elucidating the genetic basis of adaptation is one of the major goals of evolutionary biology (Bomblies & Peichel 2022). The long-standing debate over whether selection acts primarily on many loci of small effect or few loci of large effect, and whether de novo mutations or standing genetic variation are more important in adaptation (Barghi et al. 2020, Barrett & Schluter 2008, Bomblies & Peichel 2022, Herrmisson & Pennings 2017, Höllinger et al. 2019, Rockman 2012, Sella & Barton 2017, Stephan 2019) has led to a large body of empirical adaptation research (e.g. Bomblies & Peichel 2022, Enbody et al. 2023, Roberts Kingman et al. 2021). The progress in this field has been facilitated by a growing number of increasingly

sophisticated methods for genomic selection scans (Cheng & DeGiorgio 2022, DeGiorgio &

Szpiech 2022, Kern & Schrider 2018). Despite many methodological advances and the fact

that selection leaves characteristic signatures around its targets, there are still serious

limitations in distinguishing true selection signals from false positives. In particular, we still

have a limited understanding of how regions of reduced recombination (e.g. polymorphic

inversions) affect our ability to identify true selection signals and the general importance of

such regions in adaptation (Lotterhos 2019, Schaal et al. 2022).

A growing body of theoretical (Guerrero et al. 2012, Fuller et al. 2018, Kirkpatrick & Barton

2006, Schaal et al. 2022, Yeaman 2013, Yeaman et al. 2016) and empirical research identifies

polymorphic inversions as key drivers of adaptation (Berdan et al. 2023, Harringmeyer &

Hoekstra 2022, Kapun & Flatt 2019, Rane at al. 2015, Thompson & Jiggins 2014). Inversions

are chromosomal mutations that reverse the order of the sequence in the affected region and

suppress recombination between two inversion haplotypes in heterozygous individuals. The

overall magnitude of recombination reduction depends on the frequency of heterozygous

individuals within the population/species (Seich Al Basatena et al. 2013) and as suggested by

recent studies, the effect of suppressed recombination may extend beyond the inverted region

itself (Adrion et al. 2020, Crown et al. 2018, Stevison et al. 2011). Recombination suppression

between the inverted and non-inverted arrangement is a critical mechanism that contributes to

keeping coadapted genes together, facilitating local adaptation and allowing specific allelic

combinations to remain sheltered from the homogenising effects of gene flow (Hoffmann &

Rieseberg 2008, Kirkpatrick 2010).

Accumulating evidence suggests that genomes harbouring multiple polymorphic inversions,

and thus characterised by complex recombination landscapes, may be more common than

previously thought (so called inversion-rich genomes; Harringmeyer & Hoekstra 2022, Reeve

et al. 2024). Emerging and still largely unexplored questions include: Does positive selection

contribute to shaping patterns of inversion polymorphisms in species with inversion-rich

genomes? Are we able to identify causal variants underlying adaptation within inversions using

population genomic approaches? Simulation studies have shown that genome-wide studies can

distinguish between neutral and adaptive inversions (Berdan et al. 2023, Lotterhos, 2019,

Schaal et al. 2022), and many studies use either inversion frequency and/or phenotypic

information to associate specific adaptive traits with inversion genotypes/haplotypes (Faria et

al. 2019, Harringmeyer & Hoekstra 2022, Merot et al. 2021). However, empirical studies using

classical selection scan approaches, that consider the effect of inversion landscapes on the

selection inference are still scarce. Such an approach may provide valuable insights into how

inversions contribute to adaptation and how powerful selection scans are in the context of

complex genomic recombination landscapes.

The European spruce bark beetle (*Ips typographus*, hereafter the spruce bark beetle) is found

in Norway spruce forests throughout Eurasia. It is often considered to be the most destructive

forest pest, which under certain conditions can cause mass mortalities of spruce stands (Hlásny

et al. in 2021, Seidl et al. in 2016). In recent decades, exacerbated by climate change, the

frequency and severity of outbreaks have dramatically increased, making the spruce bark beetle

a major economic and environmental concern in European forests (Dobor et al. 2024, Gohli et

al. 2024, Kautz et al. 2024). Extensive research on species ecology and biology, as well as, on

causes and consequences of the spruce bark beetle outbreaks, has been recently complemented

by population genomic studies (Mykhailenko et al. 2024, Zieliński et al. 2025). Importantly,

these studies identified 27 large polymorphic inversions covering approximately 30% of the

species genome (Mykhailenko et. al 2024). Most of the identified inversions are polymorphic

across species range but can significantly vary in frequency among populations which makes

the spruce bark beetle an excellent model to study natural selection signatures in the context of

inversion rich genome.

Here, we performed genome-wide selection scans in the spruce bark beetle, with a particular

focus on the multiple inversion polymorphisms present across its genome. We investigated how

inversion-rich genome landscape affects performance of two selection scans methods that

differ in their assumptions and explored solutions to minimise the effects of inversions on

selection inference. Finally, we revealed candidate loci under selection within the spruce bark

beetle genome.

Materials and Methods

Sampling, sequencing, and data preparation

We used data presented in Mykhailenko et al. (2024) and Zieliński et al. (2025). The data

included 312 individuals from 23 populations (13-14 individuals per population). Samples

covered most of the species' range in Europe and ranged from northern Italy to northern

Fennoscandia (Figure 2; Table S1). The details of library preparation, sequencing and

processing are given in Mykhailenko et al (2024) and Zieliński et al. (2025). In short, DNA

was extracted from the whole beetles using the Wizard Genomic DNA Purification Kit
(Promega). Genomic libraries were subject to 2×150 bp paired-end sequencing using the
NovaSeq technology (Illumina Inc.). Raw data was subject to quality control with FastQC
(Andrews 2010). Low quality reads and adaptors were trimmed with Trimmomatic (Bolger et
al. 2014) and mapped to the spruce bark beetle genome (Powell et al. 2021) with Bowtie2
(Langmead & Salzberg 2012). Duplicate reads were removed with PicardMarkDulicates
(https://broadinstitute.github.io/picard/). Variant calling, genotyping, base quality score
recalibration, and filtering were performed with GATK, following the GATK best practice
guideline. Filtering included: removing five sites around indels and sites within repeat-masked
regions of the genome, and variants exhibiting: low-quality (mapping quality below 30), total
depth higher than mean + 1SD, excessive heterozygosity. Individual genotypes with coverage
lower than 8 and genotype quality lower than 20 were treated as missing. In downstream
analysis we used only biallelic SNPs with more than 50% of genotypes called. The data for all
individuals was phased using default parameters of Beagle 5.2 (Browning et al. 2021). All
analyses were performed using contigs larger than 1 >Mb (IpsContigs 1-36, 78% of assembly
length). Moreover, IpsContig33 which had high similarity to mtDNA and potentially sex linked
contigs were excluded (IpsContig9, IpsContig27, IpsContig31 and IpsContig35,
Nadachowska-Brzyska in preparation).

The effect of inversions on selection scans - general overview and predictions

There are two important aspects of inversions that can affect selection scans: low
recombination rate and genetic subdivision within the inverted regions. Inversion regions are
characterised by lower recombination due to suppressed recombination in heterozygous
individuals. Thus, the recombination landscape in these regions is shaped by the frequency of
inversion haplotypes - the more balanced the frequencies, the higher the number of
heterozygous individuals in the population and the lower the population recombination rate in
the inverted region. Importantly, haplotype-based selection statistics tend to be biased toward
low-recombination regions - selection signals are more frequently found in such regions
(O'Reilly et al. 2024) and, at least in some cases, this may be a consequence of a higher false-
positive rate in such regions (Ferrer-Admetlla et al. 2014). Additionally, a direct consequence
of suppressed recombination between alternative inversion haplotypes is independent
accumulation of mutations on both arrangements. This creates a genetic subdivision within the

inversion region - a situation that is akin to a demographic scenario of two isolated populations.

In addition, gene flux (double-crossover and conversion events) that can happen between

inversion haplotypes mimics a scenario of two isolated populations with some migration

between them. This is important since panmixia is a common assumption of selection scan

methods. Inference methods are often evaluated for their robustness to population subdivision

(or other complex demographic scenarios; DeGiorgio & Szpiech 2022, Ferrer-Admetlla et al.

2014) but almost never consider inversions as a source of potential bias (Novo et al. 2023).

To test the effect of inversions on selection inference in the spruce bark beetle genome, we used

two different selection scan methods: the number of segregating sites by length (nSL, Ferrer-

Admetlla et al. 2014) and Λ from the saltiLASSI maximum likelihood approach (DeGiorgio &

Szpiech 2022). We focused on methods that do not require recombination rate information, as

this is still lacking for many of the species for which genomic data are being generated. Both

methods are designed to search for hard and soft sweep signatures of ongoing or recent

selection but differ in the way they identify and summarise selection signals and can be

differently affected by reduced recombination and genetic subdivision within inversion

regions. While nSL has been shown to be quite robust to variation in recombination rate, Λ

tends to be negatively correlated with recombination rate, and strong Λ signals in low-

recombination regions have to be interpreted with extra caution (DeGiorgio & Szpiech 2022,

Ferrer-Admetlla et al. 2014). Due to within inversions genetic subdivision nSL may be affected

by substructure created by inversions – the stronger the isolation between isolated haplotypes

the stronger the bias (Ferrer-Admetlla et al. 2014 tested several different demographic

scenarios including population subdivision). We may expect a similar effect on Λ, but

population subdivision has not been evaluated in saltiLASSI. In addition, both methods may

be differentially affected by the presence of multiple inversions within the investigated

genome. The saltiLASSI method is based on a maximum likelihood framework that

incorporates spatial patterns of haplotype frequency distortion into a statistical model of a

sweep. This is achieved by defining a genome-wide background haplotype frequency spectrum

representing neutral expectations and haplotype frequency spectra representing distorted

frequency spectra consistent with a model of one or more sweeping haplotypes. Selection

targets are identified by looking for extreme distortions in haplotype frequency spectra relative

to genome-wide (neutral) expectations. Since the neutral expectations are based on whole-

genome data, in the case of inversion-rich genomes, they will include multiple (and often large)

regions that will shape neutral expectation while they can clearly have different haplotype

spectra patterns than collinear parts of the genome. In contrast, nSL does not consider genome-wide expectations but its value depends on allele frequency and nSL values are normalised in frequency bins using whole genome variation information. Inversions, especially if old and maintained as polymorphic by, e.g., balancing selection, may harbour multiple fixed differences (resulting in intermediate allele frequency when both haplotypes are considered together), influencing the assignment of the nSL values into the frequency bins and the normalisation process. Finally, both methods are capable of detecting hard and soft sweeps, and it is possible that multiple high-frequency inversion haplotypes can mimic soft sweep signatures. Given the different approaches and assumptions used by these methods, predicting their behaviour in inversion-rich genomes is challenging, requires empirical assessment and careful interpretation of the results.

nSL was calculated in selscan 2.0.2 (Szpiech 2024) using default parameters and phased data (it is not possible to use unphased data if it contains missing genotypes - a very common situation with whole-genome resequencing data). The nSL values for each SNP were normalised against genome-wide distribution using 'norm' function and default settings. Because the sign of nSL depends on the unknown ancestral versus derived state of the allele, we followed standard practice and used the absolute values to capture both types of signatures. The top 1% of nSL absolute values were treated as outliers. Λ was inferred using LASSI-Plus1.1.2 (DeGiorgio & Szpiech 2022). In contrast to the nSL approach, which provides statistics for each analysed SNP, Λ is given for windows consisting of a predefined number of SNPs (hereafter called SNP-based window). In addition, different window sizes may be more suitable for detecting recent or more ancient sweeps (DeGiorgio & Szpiech 2022). We tested different window sizes and based on our tests and previous studies (DeGiorgio & Szpiech 2022, Yang et al. 2023), we decided to use two window sizes: 52 (sliding window step 12) and 117 SNPs (sliding window step 12). The top 1% of Λ values were considered outliers.

To be able to compare selection signals between statistics, nSL results were translated into SNP-based Λ windows and mean values of nSL were calculated in each window. The top 1% mean (window) nSL absolute values were considered outliers. In order to be able to compare different datasets that contained different individuals and thus different numbers of SNPs and different numbers of SNP-based windows, we further standardised the results by summarising the statistics (both nSL and Λ) in non-overlapping 50 kb genomic windows. For each 50 kb window, we calculated the mean value of each statistic of all SNP-based windows that

overlapped it. This allowed a consistent window-based comparison across datasets and
statistics.

Finally, to test if selection signals tend to cluster within inversions, we plotted outlier windows
(SNP or 50kb windows according to each case) along the genome, investigated the position of
each outlier window in comparison to inversion coordinates and calculated the sizes of clusters
composed of adjacent outlier windows. Additionally, to correlate selection statistics with the
genetic differentiation between inversion haplotypes we summarised Fst in SNP-based
windows (in the same way as we did for nSL results) using vcftools (Danecek et al. 2011).

Selection scans - datasets, tested hypotheses and specific predictions

Multiple datasets used in the study are presented in Figure 1. Individuals were grouped
according to the previously described genetic structuring into the southern and northern clade
(Mykhailenko et al. 2024, Zielinski et al. 2025; Figure 2, Table S1) creating three datasets: 1)
all individuals from southern and northern region (SN, 258 individuals), 2) only individuals
from the southern populations (S, 138 individuals), 3) only individuals from the northern
populations (N, 120 individuals). These datasets did not include Polish populations that are
highly admixed. Each of these datasets was analysed in two versions, differing in the included
parts of the genome: inversion regions were either included (SN, S, N) or excluded (SNcol,
Scol, Ncol; only included collinear genome). To investigate the effects of selection in inverted
homozygotes, all individuals were also divided into datasets based on their inversion genotype,
forming three datasets per inversion: 1) individuals carrying major inversion haplotype (e.g.
MJAInv2); 2) individuals carrying minor inversion haplotype (e.g. MNAInv2); 3) individuals
carrying all possible inversion genotypes (e.g. SNInv2). Polish individuals were included in
this analysis to ensure sufficient sample sizes, and because this analysis focused on differences
between inversion genotypes rather than population-level structure. Inversion datasets
consisted of inversion in question and collinear part of the genome (thus, excluding other
inversions apart from the one in question).

The analysis of the datasets with and without inversions (SN, S and N, and their respective
SNcol, Scol and Ncol) allowed us to investigate whether all or some inversions appear as
selection outliers, to test whether the presence of multiple inversions affects the detection of
selection signals outside of inversions (in collinear part of the genome), and to compare the
behaviour of the two methods in these respects. Specifically, we predicted: 1) to find more

selection signals in inversions than in collinear parts due to the effect of reduced recombination
and inversions' genetic subdivision, 2) that balanced inversions with intermediate frequency of
both inversion haplotypes will show stronger selection signals than inversions with one allele
rarer (assuming that rare allele is younger and have low within haplotype polymorphism), 3)
that outlier detection outside inversions is affected by inversions present within the genome
due to their influence on haplotype frequency spectra (Λ) and the assignment of the statistics
values into the frequency bins (nSL) . To test whether selection signals are more often found
in inversion regions than in collinear parts of the genome we calculated the proportion of
outliers in inversions and performed a permutation test for each of the analysed dataset (1000
permutations, permuting SNP-based window labels - inverted and collinear). To check the
effect of inversion polymorphism frequency on selection scan performance, we correlated the
inversion haplotype frequency and the percentage of outliers within the inversion. To test
whether inversions affect selection inference in the collinear part of the genome we compared
datasets with and without inversions. We counted the number of outlier windows and
investigated whether the same regions were classified as outliers in both datasets and whether
new selection signals appear if we excluded inversions from the analysis. We compared both
methods by checking the number and distribution along the genome of the outlier windows
identified in both methods and across all datasets.

To be able to identify selection signals specific for the particular genetically differentiated
groups identified across the species range, selection scans described above were performed
separately for southern and northern regions/clades (S and N; Scol and Ncol; Figure 1-2). To
test if selection signals correlate with Fst between the southern and northern group we
correlated Fst with nSL and Λ values (all calculated in SNP-based windows, significance of
the correlation was calculated using permutations).

The inversion datasets (MJAInv, MNAInv and SNInv) allowed us to test whether: 1) selection
signals present in inversions when analysing dataset with all individuals (SNInv; all possible
inversion genotypes) disappear when the selection scans were performed using only
homozygous individuals (MJAInv or MNAInv); 2) There is a positive association between
inversion polymorphism and signal disappearance (more false positive selection signals
expected in high frequency of heterozygotes); 3) Selection signals disappear only in one
homozygotes revealing haplotype which carries alleles under selection. We expect that the
behaviour of the scans depends on the frequency of the inversion haplotype (produced by lower
recombination within haplotypes and higher genetic subdivision between them) and that the

most pronounced changes in selection signal occur when the inversion frequency is the highest

(around 0.5). We also predict that in some inversions false positive signals would disappear,

leaving only inversion haplotype-specific selection signals (if they exist).

In total, twenty-four inversions were analysed. In each of the inversion dataset all the other

inversions (apart from the inversion in question) were excluded before performing selection

scans. We calculated the percentage of outliers in each inversion in all datasets and in both of

the selection scans. Since 1% of the selection scans results are always treated as outliers (even

if the selection landscape is flat) we corrected the observed percentage of outliers in each

inversion for the expected number of outliers given the inversion length and analysed genome

size (given in number of SNP windows). Since several observed percentages equalled zero, we

did not normalize them by simply dividing observed by expected values. Instead, we

transformed the observed data as (observed percentage - expected percentage)/(observed

percentage + expected percentage)/2. This produced values that were easy to interpret: positive

values indicated regions enriched in the selection signal and negative values regions that were

depleted in the selection signals. These transformed percentages were used in all the

downstream tests.

To test for a difference between the observed and the expected percentage of outliers within

inversions and to test whether inversions are enriched in (Observed – Expected > 0) or depleted

(Observed – Expected < 0) from selection signals we used Wilcoxon signed-rank test. To test

whether the magnitude of selection signals depends on the frequency of the inversion

haplotype, we fitted logistic regression models for each dataset (SNInv, MJAInv, MNAInv).

We transformed the percentage of outliers into binary categories—"Depleted" (negative values)

and "Enriched" (positive values). These categories were used as response variables, and the

frequency of the major allele and its square (to capture a potential nonlinear relationships) were

included as explanatory variables. Finally, to test if selection signals disappear only in one

homozygote (major or minor) revealing haplotype which carries loci under selection we

classified selection signals into two signal categories: 1) selection signal gets much stronger in

one of the homozygous dataset (percentage of outliers in one MJAInv (or MNAInv) dataset is

higher than 50% of the percentage observed in the SNInv dataset (all genotypes) and the

percentage of outliers in the other homozygote dataset is lower than 50% of the percentage

observed in the SNInv dataset); 2) percentage of outliers is different from the first category.

After categorization we used binomial regression model with signal behaviour as response

variable and frequency of the inversion major haplotype as explanatory variable. The
significance of the variables was tested using a Wald test.

Candidate genes under selection and gene ontology enrichment analysis

We identified genes harbouring or closest to outlier windows from the top 1% of nSL or Λ
values. We used gene annotations from Powell at. al (2021) and tested for overrepresentation
of GO using R package topGO (Alexa et al. 2006). We applied Fisher's exact test and
"weight01" algorithm (Alexa & Rahnenführer 2009) to deal with the GO graph structure and
only GO categories with at least 10 members among the SNP associated genes were considered.
GO analysis were performed for datasets including: 1) selection outliers outside inversions and
specific for northern group; 2) selection outliers outside inversions and specific for southern
group; 3) selection outliers outside inversions common for both groups; 4) selection outliers
specific for particular inversion haplotype or present in both inversion haplotypes (the
inversion signals that we consider true signals).

Results

We used whole-genome sequence data from hundreds of individuals and performed nSL and Λ
selection scans across the inversion-rich spruce bark beetle genome. The list of all datasets and
their characteristics (e.g. whether they contained inversions or not) are given in Figure 1. In
short, the overall results of the selection scans are presented in Figure 3 and Tables 1-2 (nSL
and Λ for 52 SNPs window size), Figures S1 and Table S4 (nSL and Λ for 117 SNPs window)
and Figure S2 (nSL results before translating to SNP-based windows). The distribution of
selection signals in collinear regions is shown in Figure 4 and Table 3 (nSL and Λ for 50kb
windows), and the selection scan results in homozygous datasets are shown in Figure 5, Table
S5 and Figure S4 (nSL and Λ for 52 SNPs window size).

Selection landscape in inversion-rich genome

1. Inversion regions are enriched in selection signals

Most of the outliers were identified within inversion regions (Figure 3, Figure S1, Table 1),
many more than expected by chance (permutation tests; $p < 0.001$; Supplementary Information

1.1). This pattern was consistent across both statistics, window sizes and all datasets (SN, S,

N), except for Λ scan and SN dataset and 52 SNPs window size, where more outliers than

expected were found in the collinear part of the genome (Table 1). In the latter case the signal

was mainly driven by one region in IpsContig3 (39% of all outliers and 55% of collinear

outliers in IpsContig3). Only 7 inversions (out of 24) did not show any selection signals in nSL

or Λ analysis (Table S2). Inversions not only included more outliers in total but also harboured

longer clusters (in particular in Λ scans; permutation test, $p < 0.001$; Table S3; Supplementary

Information 1.8). This was also true for the Λ scan with 52 SNPs window size that had a smaller

percentage of outliers in inversions compared to the collinear part of the genome, even

considering that inversions represent only ~30% of the genome (Figure 3, Table S3). The

largest inversion outlier clusters were found in inversion Inv16 (0.324 Mb, 3% of the inversion,

in S dataset) and Inv5 (0.253 Mb, 6% of the inversion, in N dataset, Figure 3, Table S3). The

largest cluster outside inversion was found in IpsContig3 (ranging from 0.317 to 0.369 Mb in

different datasets). We did not find a significant correlation between frequency of inversion s

and amount of selection signals (Supplementary Information 1.3). In summary, we found that

inversions are enriched in selection signals and that these signals form large blocks of outlier

regions.

2.   Selection landscape identified by both methods in collinear parts is minimally affected
      by inversions present in the genome

Both nSL and Λ scans conducted using the dataset with inversions, revealed multiple signals

of selection outside inversion regions (Table 1-2, Figure 3-4, Figure S1-S2). The proportion of

outliers detected depended on the dataset and selection scan method used. Figure 4 and Table

3 compare the selection landscape in the collinear part of the genome in datasets that included

inversions when the selection scans were performed (SN, S, N, inversions were kept or

removed for the outlier threshold calculation) and datasets in which inversions were excluded

before selection scan analysis (SNcol, Ncol, Scol; both in 50kb widows that allowed us to

compare datasets with different number of SNP windows). There was a significant correlation

between the results obtained in the collinear part of the genome using datasets that included

and excluded inversions (r^2 ranged from 0.76-0.95, permutation test; p-value < 0.001;

Supplementary Information 1.6). In summary, the general selection landscape in collinear

genome was almost the same between datasets that included and excluded inversions indicating

minimal influence of inversions on the selection scan results in the collinear part of the genome.

3. Both methods identify a similar selection landscape

Both selection scan methods produced a fairly similar selection landscape across all datasets

(r^2 ranged from 0.30 to 0.45, permutation test; $p < 0.001$; Supplementary Information 1.4).

Only the selection landscape obtained from Λ scans with window size 117 SNPs was visually

different and dominated by two large outlier clusters in inversions Inv5 and Inv7 (Figure S1,

Table S4). Despite this difference, many regions identified in the analysis that used 52 SNPs

window size (both nSL and Λ) were still visible as distinct peaks when using window size of

117 SNPs but were not classified as outliers in this dataset due to the extreme values of the two

inversions mentioned above. In general, the genome-wide selection landscapes were similar

across different statistics (as indicated by a high correlation of values in 50kb widows), but the

actual overlap between outlier SNP-based windows in nSL and Λ was rather small due to the

difference in values for particular windows - the same regions appear as outliers but particular

small SNP windows can differ in the magnitude of selection signals (range from 11% to 20%

of shared outliers in Λ and nSL for 52 SNPs window size in different datasets; Table 1), but

this is still more overlap then expected by chance (permutation test, $p < 0.001$; Supplementary

Information 1.4). In summary, both methods identify similar selection landscapes despite the

observed difference in magnitude of selection signal between nSL and Λ.

4. Northern and southern regions have similar selection landscapes with few region-

specific selection signals

Selection landscapes inferred for S and N datasets were highly correlated with each other (r^2

0.76 for Λ, 0.81 for nSL; $p < 0.001$; Figure 3, Figures S1–S2, Tables 1-2, Supplementary

Information 1.5). This overall similarity was consistent across both nSL and Λ and reflected

the largely shared genomic landscape of selection between geographic regions. However, a

region-specific outlier windows were also detected (Figure S3, Table 2). In addition, both nSL

and Λ values were positively correlated with Fst values calculated across the genome and

between the N and S datasets (r^2 0.0004–0.0634 across datasets; $p < 0.001$; Supplementary

Information 1.2), indicating that regions of high genetic differentiation may be enriched for

signals of selection and at least some of them may be involved in local (regional) adaptation.

Selection scans in inversion homozygotes - detecting false positive and haplotype-specific signals

To minimise the effect of the recombination suppression and genetic subdivision between inversion haplotypes, both selection scans were performed in individuals carrying particular inversions only in homozygous states (Figure 1, MJAInv and MNAInv datasets). Each inversion was evaluated separately for both inversion homozygotes and excluding the genomic regions that contained inversions (e.g. MJAInv2 and MNAInv2; Figure 5) and compared to the results obtained using a dataset that included all genotypes in inversion in question (dataset e.g. SNInv2). We tested several hypothesis: 1) Selection signals present in inversions when analysing a dataset with SNInv individuals disappear when selection scans are performed using only homozygous individuals (MJAInv or MNAInv); 2) Selection signals disappear only in inversions with the highest frequency of heterozygotes (false positive signals would be more often in such cases); 3) Selection signals disappear only in one homozygote, revealing a haplotype that carries alleles under selection.

5. *Selection scans in homozygote datasets reveal haplotype-specific and false positive signals*

nSL selection signal was detected in ten inversions in both homozygotes, eight inversions in only one homozygote (four of which were minor alleles), and in three inversions selection signals disappeared completely in the homozygous datasets (Table S5). In the Λ scans, three inversions showed selection signals in both homozygotes, and five inversions showed signals in only one homozygote. None of the inversions lost their selection signals in both homozygotes. Figure 5 shows the large variation in the difference between the observed and the expected percentages of outliers among inversions and datasets. For example, Inv2 and Inv16 show a much higher proportion of observed outliers in SNInv and one of the homozygote datasets—dozens observed compared to only a few percent expected. In contrast, Inv13 exhibits similar observed and expected percentages in both SNInv and MJNInv, and notably fewer observed outliers than expected in MJAInv (Figure 5a). Overall, analysis of homozygote datasets revealed that the majority of inversions harbour selection signals in either both or one inversion haplotype (in the latter case revealing haplotype under selection). However, in few cases selection signals in homozygote datasets disappeared completely indicating false positive signals obtained in SNInv datasets.

 6. Selection signal tends to be reduced in homozygote datasets

While analyses using SN dataset (all individuals and all inversions), showed that inversion

regions are overall enriched for selection signals compared to collinear regions, we observed

that the strength of these signals tends to be reduced in homozygote datasets. Our tests indicated

a depletion of the signal in homozygotes (MJAInv and MNAInv datasets) compared to the

SVInv dataset (Wilcoxon tests, $p < 0.02$ for both nSL and Λ scans; see

Supplementary_Information_Selection_model.html). However, the magnitude of the selection

signal in inversion regions did not correlate with inversion haplotype frequency. Similarly,

inversion haplotype frequency did not explain the magnitude of selection signals within

inversions or indicate any clear patterns regarding what happens to selection signals in

inversions (the disappearance of the signal is not dependent on inversion frequency; see

Supplementary_Information_Selection_model.html).

In summary, although selection signal tends to be reduced in homozygous datasets compared

to SNInv, there is a considerable heterogeneity in this respect among inversions and among

datasets (Figure 5, Figure S4, Table S5). The strength of the selection signals and its behaviour

in different datasets did not depend on the frequency of inversions.

Gene ontology terms enriched in selection targets

To explore the functions associated with regions under selection, we performed Gene Ontology

(GO) enrichment analysis using outlier windows that (1) were located outside inversions and

specific to N population, (2) were located outside inversions and specific to S population, (3)

were shared between S, N, and SN datasets, or (4) remained significant after excluding

heterozygotes in the analysis performed with only homozygotes. In total, we identified 24

significantly enriched GO terms (Table S6). Of these, four terms were shared between both

nSL and Λ statistics in the only homozygote datasets, while five terms were exclusive to Λ and

three to nSL.

In the parts outside of the inversions from SN, S, and N datasets, two GO terms were shared.

Only one term was enriched specifically for S and also only one for N populations. Enriched

functions included terms such as "signalling receptor activity" and "response to stimulus,"

potentially related to odorant receptor function. A complete list of enriched GO terms, including

fold enrichment and significance levels, is provided in Table S6.

Discussion

In this study we used whole-genome resequencing data from 312 individuals of the spruce bark beetle to investigate how selection signals are distributed throughout its genome and how chromosomal inversions influence these patterns and our ability to detect them.

As predicted, we observed a significant enrichment of selection signals within inversion regions using both nSL and Λ statistics. Notably, even in the homozygotes, most inversions exhibit some selection signals in at least one homozygote dataset. These findings reinforce the view that most inversions contribute to adaptation (Christmas et al. 2019, Nadachowska-Brzyska et al. 2019, Pegueroles et al. 2010, Rane et al. 2015) and are unlikely to be selectively neutral (Berden et al. 2023), particularly in the case of large polymorphic inversions (Connallon & Olito, 2022) — the kind identified in the spruce bark beetle genome (Mykhailenko et al. 2024).

On the other hand, we observed substantial variation in both the magnitude and behaviour (enrichment or depletion) of selection signals across inversions and datasets, in particular when all genotypes were contrasted with homozygotes. In general, selection signals were weaker in the homozygote datasets, suggesting that some signals detected in the entire dataset may be inflated by the presence of heterozygous individuals. By restricting analyses to individuals carrying only one inversion haplotype, we reduced the likelihood of false positives and were able to detect haplotype-specific selection signals. This approach also helped refine candidate selection regions and distinguish genuine signals from spurious ones. Importantly, the lack of a clear relationship between the extent of inversion polymorphism and the behaviour of selection signals across datasets underscores a key challenge: although inversions are enriched for selection signals, identifying the specific targets of selection within these regions remains difficult.

Contrary to our expectations, we found that the presence of inversions had only a minimal impact on the shape of the selection landscape   and our ability of detect selection signals in the collinear genome. Importantly, this funding suggests that identifying loci under selection outside of inversions remains feasible, even in genomes characterized by extensive inversion polymorphism. This result aligns with our earlier findings on demographic inference (Zieliński et al. 2025), where we showed that overlooking chromosomal inversions had little effect on model selection and introduced only minor biases in the estimation of demographic parameters.

Additionally, the spruce bark beetle selection scans supported the necessity of using multiple complementary approaches to selection scans. While the overall patterns of selection inferred by Λ and nSL were similar, differences were observed at a local scale. There was a high correlation of selection signals at 50 kb windows, but only a small overlap among outlier windows at 52 SNPs, which demonstrates the general difficulty of selection scans in pinpointing the focal gene under selection (Vitti et al. 2013). Moreover, the selection of window sizes in the Λ scans can influence which regions are considered candidates for selection. The differences observed in the spruce bark beetle between 52 and 117 SNP windows can, to some extent, reflect differences in the ability to detect more recent and more ancient sweep signatures: smaller window sizes are more effective at detecting ancient sweeps (DeGiorgio & Szpiech 2022).

Our study revealed a broadly similar selection landscape between the southern and northern populations of the spruce bark beetle, suggesting that these populations are likely subject to comparable selective pressures and/or that their divergence is relatively recent, with the observed signals possibly reflecting selection in a shared ancestral population. According to our recent demographic modelling, the divergence between the southern and northern clades occurred approximately 80,000 years ago (Zieliński et al. 2025). Signals of selection in the site frequency spectrum are expected to persist for roughly 0.1Ne generations, while haplotype-based signals fade more rapidly, typically within ~0.01Ne generations (Pennings & Hermisson 2006, Przeworski 2002). Assuming an ancestral Ne of 250,000 (Zieliński et al. 2025), haplotype-based signatures could persist for around 2,500 generations—well after the estimated divergence time—supporting the idea that shared selection pressures may be acting across both populations. The inferred Ne for the northern and southern populations after the split was 180,000 and 830,000, respectively, and even under the higher Ne scenario, the timescale of detectable selection signals remains post-divergence. These findings suggest that spruce bark beetle populations across Europe may be subject to similar selective pressure. Nonetheless, we also detected evidence of regional differentiation: one of the strongest selection signals outside of inversions (on IpsContig3) overlapped with a pronounced Fst peak between the northern and southern groups, indicating that divergent selection may also contribute to shaping the selection landscape in this species.

Finally, our GO enrichment analysis identified terms such as "signalling receptor activity" and "response to stimulus," which are potentially linked to odorant receptor function. These findings are of particular interest considering recent studies showing that inversions are

enriched in genes encoding odorant receptors (Mykhailenko et al. 2024), and that these
inversions may contain receptor variants that respond differently to different components of
pheromones (Johny et al. 2025).

Conclusions and future directions

Our study provides new insights into the selection landscape across the spruce bark beetle
genome, identifying candidate genes that may guide future research into the genomic basis of
adaptation in this ecologically and economically important species. We show that most
polymorphic inversions in the genome are enriched for selection signals, supporting their role
as key contributors to adaptation, while also highlighting the importance of accounting for
structural variation — particularly inversions — when performing selection inference. As
illustrated by the changing selection patterns in the homozygote datasets, inversions can affect
the detection of selection within their regions. Nevertheless, their presence does not obscure
the identification of selection signals outside inversions, emphasizing the feasibility of
conducting genome-wide selection scans even in inversion-rich genomes. More broadly, our
results underscore the need to systematically evaluate the performance and robustness of
evolutionary inference methods in the presence of structural variation — a factor still largely
overlooked in standard methodological assessments, with notable exceptions such as the recent
evaluation of GONE, which used simulations to assess how balanced inversion polymorphisms
affect effective population size estimates (Novo et al. 2023). However, such efforts remain rare,
and further research is needed to understand how inversions influence evolutionary inference,
particularly given that their effects likely vary across species with differing numbers and
evolutionary histories of large polymorphic inversions.

Acknowledgments

We thank members of the Genomics and Experimental Evolution Group at Jagiellonian
University for their help in improving this manuscript. This work was funded by a Polish
National Science Center 2018/30/E/NZ8/00105 grant to K.N.B. We gratefully acknowledge
Polish high-performance computing infrastructure PLGrid (HPC Centers: ACK Cyfronet
AGH) for providing computer facilities and support within computational grant no.
PLG/2024/017278 to K.N.B.

Data Availability

All DNA sequences have been deposited in the National Centre for Biotechnology Information Sequence Read Archive under BioProject ID PRJNA1013983. Scripts used for data processing, generation of intermediate files, and analyse, have been archived and are publicly available in the Dryad Digital Repository entry [will be added as soon as possible].

Author contributions

J.M.G., K.N.B. conceived the study and wrote the first draft of the manuscript. J.M.G, performed all the main analyses. P.Z processed and supervised the raw data processing. W.B. performed the GO analysis. W.B. and P.Z helped in data interpretation and provided feedback on all manuscript versions. All authors read and approved the final manuscript.

References

Adrion, J. R., et al (2020). Predicting the landscape of recombination using deep learning. *Molecular Biology and Evolution*, vol. 37(6):1790–808. https://doi.org/10.1093/molbev/msaa038.

Alexa, A., et al. (2006). Improved scoring of functional groups from gene expression data by decorrelating GO graph structure'. *Bioinformatics*, vol. 22(13):1600–16007. https://doi.org/10.1093/bioinformatics/btl140.

Alexa, A., & Rahnenführer, J. (2009). Gene set enrichment analysis with topGO. *Bioconductor Improv*, 27(1-26): 776.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. Braham Bioinformatics.

Barrett, R., & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends in Ecology & Evolution*, vol. 23(1):38–44. https://doi.org/10.1016/j.tree.2007.09.008.

Berdan, E. L., et al. (2023) How chromosomal inversions reorient the evolutionary process. *Journal of Evolutionary Biology*, vol. 36(12):1761–1782. https://doi.org/10.1111/jeb.14242.

Bolger, A. M., et al. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, vol. 30(15):2114–2120. https://doi.org/10.1093/bioinformatics/btu170.

Bomblies, K., and Peichel, C. L. (2022). Genetics of adaptation. *Proceedings of the National Academy of Sciences*, vol. 119(30):1-10. https://doi.org/10.1073/pnas.2122152119.

Browning, B. L., et al. (2021) Fast two-stage phasing of large-scale sequence data. *The American Journal of Human Genetics*, vol. 108(10):1880–1890. https://doi.org/10.1016/j.ajhg.2021.08.005.

Cheng, X., and DeGiorgio, M. (2022). BalLeRMix +: mixture model approaches for robust

joint identification of both positive selection and long-term balancing selection.

Bioinformatics, vol. 38(3):861–63. https://doi.org/10.1093/bioinformatics/btab720.

Christmas, M. J., et al. (2019). Chromosomal inversions associated with environmental

adaptation in honeybees. *Molecular Ecology*, vol. 28(6):1358–1374.

https://doi.org/10.1111/mec.14944.

Connallon, T., and Olito, C. (2022). Natural selection and the distribution of chromosomal

inversion lengths. *Molecular Ecology*, vol. 31(13):3627–3641.

https://doi.org/10.1111/mec.16091.

Crown, K. N., et al. (2018). Local inversion heterozygosity alters recombination throughout

the genome. *Current Biology*, vol. 28(18):2984-2990.

https://doi.org/10.1016/j.cub.2018.07.004.

Danecek, P., et al. (2011). The Variant Call Format and VCFtools'. *Bioinformatics*, vol.

27(15):2156–2158. https://doi.org/10.1093/bioinformatics/btr330.

DeGiorgio, M., and Szpiech. Z. A. (2022). A spatially aware likelihood test to detect sweeps

from haplotype distributions. *PLOS Genetics*, vol. 18(4):1-37

https://doi.org/10.1371/journal.pgen.1010134.

Dobor, L., et al. (2024). The interacting effect of climate change and herbivory can trigger

large-scale transformations of European temperate forests. *Global Change Biology*,

vol. 30(2):1-20. https://doi.org/10.1111/gcb.17194.

Enbody, E. D., et al. (2023). Community-wide genome sequencing reveals 30 years of

Darwin's finch evolution. *Science*, vol. 381(6665):1-9.

https://doi.org/10.1126/science.adf6218.

Faria, R., et al. (2019). Evolving inversions. *Trends in Ecology & Evolution*, vol. 34(3):239–

48. https://doi.org/10.1016/j.tree.2018.12.005.

Ferrer-Admetlla, A., et al. (2014). On detecting incomplete soft or hard selective sweeps

using haplotype structure. *Molecular Biology and Evolution*, vol. 31(5):1275–1291.

https://doi.org/10.1093/molbev/msu077.

Fuller, Z. L., et al. (2018). Ancestral polymorphisms explain the role of chromosomal

inversions in speciation. *PLOS Genetics*, vol. 14(7):1-26.

https://doi.org/10.1371/journal.pgen.1007526.

Gohli, J., et al. (2024). Climatic and management-related drivers of endemic European spruce

bark beetle populations in Boreal forests. *Journal of Applied Ecology*, vol. 61(4):809–

820. https://doi.org/10.1111/1365-2664.14606.

Guerrero, R. F., et al. (2012). Coalescent patterns for chromosomal inversions in divergent

populations. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

vol. 367(1587):430–438. https://doi.org/10.1098/rstb.2011.0246.

Harringmeyer, O. S., and Hoekstra, H. E. (2022). Chromosomal inversion polymorphisms

shape the genomic landscape of deer mice. *Nature Ecology & Evolution*, vol.

6(12):1965–1979. https://doi.org/10.1038/s41559-022-01890-0.

Hlásny, T., et al. (2021). Bark beetle outbreaks in Europe: state of knowledge and ways

forward for management. *Current Forestry Reports*, vol. 7(3):138–165.

https://doi.org/10.1007/s40725-021-00142-x.

Hoffmann, A. A., and Rieseberg, L. H. (2008). Revisiting the impact of inversions in

evolution: from population genetic markers to drivers of adaptive shifts and

speciation? *Annual Review of Ecology, Evolution, and Systematics*, vol. 39(1):21–42.

https://doi.org/10.1146/annurev.ecolsys.39.110707.173532.

Höllinger, I., et al. (2019). Polygenic adaptation: from sweeps to subtle frequency shifts.

PLOS Genetics, vol. 15(3):1-26. https://doi.org/10.1371/journal.pgen.1008035.

Johny, J., et al. (2025). Not all bark beetles smell the same: population-level functional

olfactory polymorphisms in *Ips Typographus* pheromone receptor ItypOR33.

Molecular Ecology, vol. 34(6):1-19. https://doi.org/10.1111/mec.17693.

Kapun, M., et al. (2023). An ancestral balanced inversion polymorphism confers global

adaptation. *Molecular Biology and Evolution*, vol. 40(6):1-28.

https://doi.org/10.1093/molbev/msad118.

Kautz, M., et al. (2024). Early detection of bark beetle (*Ips typographus*) infestations by

remote sensing – a critical review of recent research. *Forest Ecology and*

Management, vol. 556:1-17. https://doi.org/10.1016/j.foreco.2023.121595.

Kern, A. D., and Schrider, D. R. (2018). diploS/HIC: an updated approach to classifying

selective sweeps. *G3 Genes|Genomes|Genetics*, vol. 8(6):1959–1970.

https://doi.org/10.1534/g3.118.200262.

Kirkpatrick, M. (2010). How and why chromosome inversions evolve. *PLoS Biology*, vol.

8(9):1-5. https://doi.org/10.1371/journal.pbio.1000501.

Kirkpatrick, M., and Barton, N. (2006). Chromosome inversions, local adaptation and

speciation. *Genetics*, vol. 173(1):419–434.

https://doi.org/10.1534/genetics.105.047985.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, vol. 9(4):357–359. https://doi.org/10.1038/nmeth.1923.

Lotterhos, K. E. (2019). The effect of neutral recombination variation on genome scans for

selection. *G3 Genes|Genomes|Genetics*, vol. 9(6):1851–1867.

https://doi.org/10.1534/g3.119.400088.

Mérot, C., et al. (2021). Locally adaptive inversions modulate genetic variation at different

geographic scales in a seaweed fly. *Molecular Biology and Evolution*, vol.

38(9):3953–3971. https://doi.org/10.1093/molbev/msab143.

Mykhailenko, A., et al. (2024). Complex genomic landscape of inversion polymorphism in

Europe's most destructive forest pest. *Genome Biology and Evolution*, vol. 16(12):1-

23. https://doi.org/10.1093/gbe/evae263.

Nadachowska-Brzyska, K., et al. (2019). Footprints of adaptive evolution revealed by whole

Z chromosomes haplotypes in flycatchers. *Molecular Ecology*, vol. 28(9):2290–2304.

https://doi.org/10.1111/mec.15021.

Novo, I., et al. (2023). Impact of population structure in the estimation of recent historical

effective population size by the software GONE. *Genetics Selection Evolution*, vol.

55(1):1-15. https://doi.org/10.1186/s12711-023-00859-2.

O'Reilly, G. D., et al. (2024). A new method for ecologists to estimate heterozygote excess

and deficit for multi-locus gene families. *Ecology and Evolution*, vol. 14(7):1-13.

https://doi.org/10.1002/ece3.11561.

Pegueroles, C., et al. (2010). Recombination and selection in the maintenance of the adaptive

value of inversions. *Journal of Evolutionary Biology*, vol. 23(12):2709-2717.

https://doi.org/10.1111/j.1420-9101.2010.02136.x.

Pennings, P. S., and Hermisson, J. (2006). Soft sweeps II—molecular population genetics of

adaptation from recurrent mutation or migration. *Molecular Biology and Evolution*,

vol. 23(5):1076-1084. https://doi.org/10.1093/molbev/msj117.

Powell, D., et al. (2021). A highly contiguous genome assembly of the Eurasian spruce bark

beetle, *Ips typographus*, provides insight into a major forest pest. *Communications*

Biology, vol. 4(1):1-9. https://doi.org/10.1038/s42003-021-02602-3.

Przeworski, M. (2002). The signature of positive selection at randomly chosen loci. *Genetics*,

vol. 160(3):1179–1189. https://doi.org/10.1093/genetics/160.3.1179.

Rane, R. V., et al. (2015). Genomic evidence for role of inversion 3RP of *Drosophila*

Melanogaster in facilitating climate change adaptation. *Molecular Ecology*, vol.

24(10):2423–2432. https://doi.org/10.1111/mec.13161.

Reeve, J., et al. (2024). Chromosomal inversion polymorphisms are widespread across the

species ranges of rough periwinkles (*Littorina Saxatilis* and *L. arcana*). *Molecular*

Ecology, vol. 33(24):1-18. https://doi.org/10.1111/mec.17160.

Roberts Kingman, G. A., et al. (2021). Predicting future from past: the genomic basis of

recurrent and rapid stickleback evolution. *Science Advances*, vol. 7(25):1-12.

https://doi.org/10.1126/sciadv.abg5285.

Rockman, M. V. (2012). The QTN program and the alleles that matter for evolution: all that's

gold does not glitter. *Evolution*, vol. 66(1):1–17. https://doi.org/10.1111/j.1558-

5646.2011.01486.x.

Schaal, S. M., et al. (2022). Inversion invasions: when the genetic basis of local adaptation is

concentrated within inversions in the face of gene flow. *Philosophical Transactions of*

the Royal Society B: Biological Sciences, vol. 377(1856):1-14.

https://doi.org/10.1098/rstb.2021.0200.

Seich Al Basatena, N. K., et al. (2013). The effect of genomic inversions on estimation of

population genetic parameters from SNP data. *Genetics*, vol. 193(1):243-253.

https://doi.org/10.1534/genetics.112.145599.

Seidl, R., et al. (2016). Small beetle, large-scale drivers: how regional and landscape factors

affect outbreaks of the European spruce bark beetle. *Journal of Applied Ecology*, vol.

53(2):530-540. https://doi.org/10.1111/1365-2664.12540.

Sella, G., and Barton, N. H. (2019). Thinking about the evolution of complex traits in the era

of genome-wide association studies. *Annual Review of Genomics and Human*

Genetics, vol. 20(1):461-493. https://doi.org/10.1146/annurev-genom-083115-

022316.

Stephan, W. (2019). Selective sweeps. *Genetics*, vol. 211(1):5–13.

https://doi.org/10.1534/genetics.118.301319.

Stevison, L. S., et al. (2011). Effects of inversions on within- and between-species

recombination and divergence. *Genome Biology and Evolution*, vol. 3:830-841.

https://doi.org/10.1093/gbe/evr081.

Szpiech, Z. A., et al. (2021). Application of a novel haplotype-based scan for local adaptation

to study high-altitude adaptation in rhesus macaques. *Evolution Letters*, vol. 5(4):408-

421. https://doi.org/10.1002/evl3.232.

Szpiech, Z. A. (2024). Selscan 2.0: scanning for sweeps in unphased data. *Bioinformatics*,

vol. 40(1):1-6. https://doi.org/10.1093/bioinformatics/btae006.

Thompson, M. J., and Jiggins, C. D. (2014). Supergenes and their role in evolution. *Heredity*,

vol. 113(1):1–8. https://doi.org/10.1038/hdy.2014.20.

Vitti, J. J., et al. (2013). Detecting natural selection in genomic data. Annual Review of

Genetics, vol. 47(1):97-120. https://doi.org/10.1146/annurev-genet-111212-133526.

Yang, X., et al. (2022). Lineage-specific targets of positive selection in three leaf beetles with

different defence capacity against a parasitic wasp. *Authorea*.

https://doi.org/10.22541/au.166754711.16782535/v1.

Yeaman, S., et al. (2016). Convergent local adaptation to climate in distantly related conifers.

Science, vol. 353(6306):1431–1433. https://doi.org/10.1126/science.aaf7812.

Yeaman, S. (2013). Genomic rearrangements and the evolution of clusters of locally adaptive

loci. *Proceedings of the National Academy of Sciences*, vol. 110(19):1743-1751.

https://doi.org/10.1073/pnas.1219381110.

Zieliński, P., et al. (2025). Demographic history inferred from an inversion-rich spruce bark

beetle genome. Submitted.

Figures

<u>Figure 1</u>. A graphical summary of the datasets used for selection scans in the spruce bark beetle.
nSL and Λ selection scans were performed with two window sizes: 52 and 117 SNPs.
Individual beetles were assigned to different groups based on the following two factors: 1)
geographic origin: southern populations (S and Scol ("col" abbreviation of "collinear" to refers
to datasets that included only collinear parts of the genome, excluding inversions), red),
northern populations (N and Ncol, blue), or all individuals (SN and SNcol, southern and
northern region combined, yellow), and 2) inversion genotype: major allele homozygotes
(MJAInv, orange), minor allele homozygotes (MNAInv, green), and groups including all

possible genotypes (SNInv, yellow). Each dataset included versions that included (e.g., SN) or
excluded (e.g., SNcol) inversions from the analysis.

Figure 2. The geographic distribution, admixture, and genetic differentiation of *Ips*
typographus populations as inferred in Zielinski et al. 2025. Top left: map of sampled
populations, with admixture proportions illustrated by coloured circles. The colours represent
the admixture proportions that were identified (top right). Admixture plot showing individuals
admixture proportions for K=2. The populations are separated by dashed lines and ordered
according to longitude. Bottom plot: pairwise Fst calculated between the two genetic clusters
across the genome. Grey and black represent different contigs.

Figure 3. Results of the selection scans for the 52 SNP window sizes and SN, S, and N datasets for both selection scans (a) Λ and (b) nSL (absolute values). Results using both collinear and inversion regions are shown in black and the results of the same analysis performed excluding inversion regions are overlaid (SNcol – yellow, Scol – red, and Ncol – blue). The dashed lines indicate the outlier threshold (1% of all values). In all plots, grey dotted vertical lines represent the end of each contig.

Figure 4. Comparison of the selection landscape in a collinear genome across different datasets, including or excluding inversion regions. (a) nSL selection scans results (absolute values); (b) Λ selection scans results. Only collinear regions are shown. The dashed lines represent the outlier thresholds calculated as 1% of the values in each respective dataset. SN, S, and N correspond to datasets in which selection scans were performed using the entire genome (including collinear and inverted regions). These datasets have two outlier thresholds: one including (grey) and one excluding (orange) inversions in the threshold calculation. SNcol, Scol, and Ncol represent datasets in which only collinear regions were included in the selection scans and the outlier threshold is shown in pink. Outliers values are colour-coded as follows: violet: outliers in collinear regions identified based on datasets that included inversions, and inversions were not removed before identifying the outliers; purple: outliers in collinear regions

identified based on datasets that included inversions and inversions were removed before

identifying the outliers; and pink: outliers in collinear regions identified based on datasets that

included only collinear regions. The visual comparison of coloured dots indicates that

inversions have negligible effect on the identification of outliers in the collinear genome.

Figure 5. Comparison of the expected and the observed percentages of outliers detected in

inversions. The expected values were calculated based on the length of the inversion and a 1%

outlier threshold. The results are shown for nSL (a) and Λ (b). All selection scans were

performed using 52 SNP windows and three datasets: datasets including individuals carrying

any of the possible inversion genotypes (SNInv, yellow); datasets including only individuals

that were homozygous for the major allele for the inversion in question (MJAInv, orange); and

datasets including only individuals that were homozygous for the minor allele for the inversion

in question (MNAInv, green). Each dataset included one analysed inversion and a collinear

genome. The inversions are ordered according to the major allele frequency (from lowest to

highest, top to bottom).

Tables

Table 1. The comparison between Λ and nSL selection scans using different datasets and a 52-SNP window size. Calculations from nSL analysis are based on the absolute values of the statistic. Genomic region: part of the genome that is included in each dataset. Inversion-associated outliers: the percentage of outlier windows located within inversion regions, calculated from the total number of outliers. Collinear-region outliers: percentage of outliers located outside inversion regions calculated from the total number of outliers. Shared outliers: percentage of outliers that are shared between Λ and nSL results. Shared outliers in inversions: the percentage of outliers shared between Λ and nSL results that are in inversion regions. Shared outliers in collinear: the percentage of outliers shared between Λ and nSL results that are in the collinear genome.

Dataset	Genomic region	Statistic	Inversion associated outliers	Collinear region outliers	Shared outliers	Shared outliers in inversions	Shared outliers in collinear
SN	Inversions + collinear	Λ	29%	71%	15%	12%	3%
		nSL	86%	14%			
SNcol	Only collinear	Λ		100%	11%		11%
		nSL		100%			
S	Inversions + collinear	Λ	70%	30%	20%	18%	2%
		nSL	87%	13%			
Scol	Only collinear	Λ		100%	16%		16%
		nSL		100%			
N	Inversions + collinear	Λ	56%	44%	17%	15%	2%
		nSL	91%	9%			
Ncol	Only collinear	Λ		100%	19%		19%
		nSL		100%			

Compared datasets	Genomic region	Statistic	Shared outliers	Shared outliers in inversions	Shared outliers in collinear
SN, N	Inversions + collinear	Λ	23%	5%	18%
		nSL	33%	31%	2%
SNcol, Ncol	Only collinear	Λ	19%		19%
		nSL	32%		32%
SN, S	Inversions + collinear	Λ	30%	6%	24%
		nSL	48%	39%	9%
SNcol, Scol	Only collinear	Λ	44%		44%
		nSL	44%		44%
S, N	Inversions + collinear	Λ	9%	2%	7%
		nSL	20%	18%	2%
Scol, Ncol	Only collinear	Λ	24%		24%
		nSL	20%		20%

Table 3. Changes in percentage of outliers found in collinear parts of the genome when

including or excluding inversions. Results are shown for three datasets (SN, N, and S) analysed

using both selection scan methods (Λ and nSL) with results summarised in 50kb windows.

Calculations from nSL analysis are based on the absolute values of the statistic. Outliers in

collinear regions before excluding inversions: the percentage of outliers in collinear regions

identified based on datasets that included inversions, inversions were not removed before

identifying the outliers. Lost outliers in collinear parts after excluding inversions: the

percentage of outlier windows in collinear parts that disappear after excluding inversions and

recalculating the outlier threshold. New outliers in collinear parts after excluding inversions:

the percentage of outlier windows in collinear parts that appear after excluding inversions and

recalculating the outlier threshold.

Dataset	Statistic	Outliers in collinear regions before excluding inversions	Lost outliers in collinear parts after excluding inversions	New outliers in collinear parts after excluding inversions
SN	Λ	83%	9%	0%
	nSL	30%	0%	42%
S	Λ	48%	0%	26%
	nSL	36%	0%	36%
N	Λ	60%	0%	14%
	nSL	21%	0%	53%

Norway spruce native range

Norway spruce introduced

Fst

Genomic position