Title: Florigen Activation Complex Dynamics and SVP-Mediated Repression
Orchestrate Temperature-Regulated Flowering in Saffron

Authors: Diksha Kalia[1, 2], Joel Jose-Santhi[1, 2], Firdous Rasool Sheikh[1, 2], and Rajesh
Kumar Singh[1, 2, *]

Affiliations:
1. Plant Adaptation and Developmental Biology Lab, Department of Biotechnology,
CSIR-Institute of Himalayan Bioresource Technology, Palampur, H.P., India-
176061
2. Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, 201002,
India

Author Details:
Diksha Kalia, email: dikshakalia47@gmail.com,
Joel Jose -Santhi, email: joeljosesanthi@gmail.com,
Firdous Rasool Sheikh, email: sheikhfirdous191@gmail.com,
Rajesh Kumar Singh: *Corresponding Author, email: rajeshsingh@ihbt.res.in, ORCID
Id: 0000-0001-7831-7946

Abstract
Saffron, a high-value spice cultivated worldwide for its therapeutic and culinary uses,
is a sterile triploid species, rendering conventional breeding approaches ineffective.
This limitation underscores the need for molecular and biotechnological strategies for
its genetic improvement. Flowering, a key determinant of saffron yield, is strongly
influenced by temperature; however, the genetic regulatory networks underlying this
process remain poorly understood. Our study identifies key regulators of saffron's
flowering, focusing on the Florigen Activation Complex (FAC) components:
FLOWERING LOCUS T (FT), bZIP transcription factor FD, and *TERMINAL FLOWER-*
1 (TFL-1), and demonstrate their temperature-dependent roles in floral regulation.
Spatiotemporal expression analyses suggested that *CsatFT3* and *CsatFD2*,
expressed in the floral meristem promote floral induction, while *CsatTFL1*-3 acts as a
floral repressor. Protein interaction studies showed that CsatFT3 and CsatTFL1-3
compete for binding to CsatFD2, and their balance modulates floral induction.
Functional validation in Arabidopsis and Saffron confirmed these findings.
Furthermore, we identified *CsatSVP2*, an ortholog of *SHORT VEGETATIVE PHASE*
(SVP), as a low temperature-responsive repressor that directly binds the CsatFT3

promoter to inhibit its expression. Together, these findings enhance our understanding
of temperature mediated floral induction in saffron and provide insights and lay the
groundwork for genetic interventions to enhance yield under variable temperature
conditions.

Keywords: Florigen activation complex (FAC), Flowering locus T(FT), bZIP
transcription factor-FD, *TFL1*, Geophytes, Thermoresponsive, Thermoperiodic, Floral
induction, Phosphatidyl Ethanolamine Binding Protein (PEBP), VIGS,

Introduction

Saffron (*Crocus sativus* L.) is a sterile, clonally propagated, autotriploid geophytic
monocot cultivated for its highly valued dried stigmas, which constitute the world's
most expensive spice. While the stigmas are the primary edible and culinary
component, other parts of the saffron plant also have diverse applications, including
uses in agriculture, traditional medicine, and as a natural colouring agent, making the
entire flower economically valuable (Ahrazem et al., 2015). Saffron has been
predominantly cultivated for centuries in select regions such as Iran, India, Greece,
Italy, Afghanistan, Morocco, Spain, and various Mediterranean basins (Cardone et al.,
2020). Flowering is the key determinant of saffron's crop productivity, and this critical
phase is increasingly influenced by environmental changes. As a thermoperiodic plant,
saffron's flowering is primarily regulated by temperature. During the warm summer
months, the transition from vegetative to reproductive growth occurs underground,
with floral bud emergence only triggered by exposure to cooler temperatures (Molina
et al., 2005a). This biphasic thermal requirement consists of an induction phase at
warm temperatures lasting 50–150 days, followed by an emergence phase under
cooler conditions, a pattern that has been well documented (Jose-Santhi et al., 2023;
Molina et al., 2005a). The inappropriate temperature during the flowering transition
leads to flower atrophy or no flower, causing yield loss (Wang et al., 2021). Saffron
corms can sense the temperature change and modify the response accordingly
(Molina et al., 2005b). Thus, there exists a thermoresponsive regulation of flowering
in saffron which is still not well studied at the molecular level.
Flowering is a key transition in the angiosperm life cycle, marking the shift from
vegetative to reproductive growth and representing a tightly regulated process vital for
survival. This process is orchestrated by a complex network of environmental cues,

internal signalling pathways, hormonal regulation, and genetic programs that together
ensure flowering occurs under optimal conditions (Zik and Irish, 2003). Key
environmental pathways influencing flowering include photoperiod, ambient
temperature, light quality, vernalization, and gibberellin signalling. These pathways
operate in coordination with endogenous factors along with plant age and nutritional
status, allowing plants to synchronize reproduction with favourable environmental
windows (Andrés and Coupland, 2012; Freytes et al., 2021; Pyo et al., 2014; Srikanth
and Schmid, 2011). Extensive studies in model organisms such as Arabidopsis
thaliana have elucidated major regulatory components of flowering (Kinoshita and
Richter, 2020). Central to this regulation are floral integrator genes, particularly
FLOWERING LOCUS T (FT) a member of the phosphatidylethanolamine-binding
protein (PEBP) family. FT functions to promote flowering by forming the Florigen
Activation Complex (FAC), which includes FD-like bZIP transcription factors and the
florigen receptor 14-3-3 protein (Abe et al., 2005; Taoka et al., 2013). This complex
activates floral meristem identity genes such as *LEAFY* and *APETALA1*, initiating floral
development (Putterill et al., 2004; Simon et al., 1996; Wigge et al., 2005). In contrast,
TERMINAL FLOWER 1 (TFL1), another PEBP family member, acts antagonistically
to repress flowering (Wickland and Hanzawa, 2015). Members of the FD family are
key interaction partners of the PEBP family, playing roles in both promoting and
repressing meristem differentiation, depending on their specific interacting partners
(Zhu et al., 2020). Concurrently, the flowering program suppresses negative
regulators, including *FLOWERING LOCUS C* (FLC), and *SHORT VEGETATIVE*
PHASE (SVP), which are crucial for ensuring a proper transition from the vegetative
to the reproductive phase mainly regulated by temperature (Bowman et al., 2012;
Turck et al., 2008). Temperature-dependent flowering has been widely studied, with
FT-like genes playing a central role (Capovilla et al., 2015) . Upstream regulators
include MADS-domain transcription factors such as SVP and FLC, which modulate
flowering by forming complexes and repressing FT expression via direct binding to its
promoter (Balasubramanian et al., 2006; Gu et al., 2013; Lee et al., 2007). However,
the thermoresponsive pathway governing floral induction in saffron remains largely
unexplored.
In contrast to annuals, the regulation of flowering in perennial geophytes involves
modified or additional mechanisms to accommodate their extended life cycles, periods
of dormancy and dual reproduction (Khosa et al., 2021) . This has been observed in

species like *Narcissus tazetta* (Noy-Porat et al., 2013), tulip (Tulipa spp.) (Leeggangers et al., 2017b), and saffron (Molina et al., 2005a), where floral induction occurs even in the absence of leaves. In these species, flowering is thought to be mediated directly at the meristem level, independent of traditional photoperiod or vernalization cues. Instead, ambient temperature plays a dominant role in determining flowering time. These observations highlight the need to identify the specific genes and mechanisms that govern flowering in geophytic perennials. Over evolutionary time, the molecular mechanisms controlling flowering have grown increasingly complex, involving intricate interactions between floral regulator genes and their transcription factor partners, which form regulatory complexes (Wickland and Hanzawa, 2015) . Gene duplication and neofunctionalization have contributed to the diversification of these regulatory genes. Both *FT* and *TFL1,* the central regulators of flowering, have undergone gene duplication events in various lineages, leading to paralogs with distinct functions that promote or inhibit flowering (Jin et al., 2021) . This evolutionary diversification is evident across many species. In geophytes, for instance, members of the PEBP gene family have extended their functions beyond flowering to include the formation of underground storage organs, contributing to both sexual and asexual reproduction (Khosa et al., 2021). In species like potato, tulip, and others, gene duplication has led to distinct roles in flowering, vegetative growth, and tuber or bulb formation (Bellinazzo et al., 2025; Jing et al., 2023; Navarro et al., 2011). Similar divergence has been observed in onion (Allium cepa), sugar beet (Beta vulgaris), and hybrid aspen, demonstrating the complexity of flowering regulation in perennials (Hsu et al., 2011; Lee et al., 2013b; Pin et al., 2010). Beyond the PEBP gene family, functional diversification is seen in related pathways. In hybrid aspen, two FD homologs, *FDL1* and *FDL2*, regulate seasonal growth differently (Tylewicz et al., 2015). In rice, *OsFD2* contributes to leaf development while *OsFD1* regulates flowering (Tsuji et al., 2013). FD-like transcription factors can have dual roles, promoting or repressing flowering depending on their partners. For instance, *CmFDa* in Chrysanthemum morifolium represses flowering through epigenetic regulation (Xue et al., 2025). These examples highlight the evolutionary plasticity of flowering networks and the importance of species-specific regulatory mechanisms.

Saffron, due to its sterile nature and extremely limited genetic diversity, is not amenable to conventional breeding approaches, which hinders efforts to develop climate-resilient, high-yielding, and stress-tolerant varieties. As a result, there have

been no significant advancements in improving saffron's adaptability to climate change, making its cultivation increasingly vulnerable to environmental fluctuations. Consequently, only biotechnological interventions can be employed to introduce desirable traits. Despite its economic and agricultural importance, research on the molecular regulation of flowering in saffron remains limited. Most existing studies have focused on temperature effects on flowering physiology (Molina et al., 2005a; Wang et al., 2021), and while several flowering-related genes have been identified through transcriptomic and genomic analyses (Hu et al., 2020; Kalia et al., 2022; Renau-Morata et al., 2021; Singh et al., 2023; Tsaftaris et al., 2013), these remain largely descriptive and lack functional validation. Expression profiling has indicated the involvement of multiple *FT* and *TFL1* homologs in saffron flowering (Kalia et al., 2022; Renau-Morata et al., 2021; Tsaftaris et al., 2013), yet no functional characterization of these genes has been reported. To fill this gap, we present a comprehensive molecular framework for temperature-dependent floral induction in saffron—a geophytic monocot with a unique flowering phenotype. By integrating gene expression profiling, heterologous functional assays, protein–protein interaction studies, promoter binding analyses, and virus-induced gene silencing, we identify key regulatory components that mediate the floral transition in response to ambient temperature. Overall, these findings advance our understanding of temperature-mediated flowering in saffron and provide a foundation for genetic interventions aimed at improving yield under fluctuating environmental conditions

Methods

Plant materials and growth conditions

Saffron corms *(Crocus sativus* L.) were grown under controlled conditions at CSIR-IHBT, and samples were collected monthly from January to September, covering both the vegetative growth and flowering phases (Jose-Santhi et al., 2023; Kalia et al., 2022). To analyze gene expression patterns, apical buds, axillary buds, and corm tissues were harvested across this period, with three biological replicates per time point, each comprising pooled tissue from ten individual corms. For RT-qPCR analysis, apical buds, axillary bud, corm tissue collected during the vegetative and flowering stages were used, with detailed methods and results provided in the Supplementary Information.

ViGS experiments were conducted using corms grown under the same controlled
conditions as described earlier. Corms with high flowering competency (weighing >10
g) were selected for the silencing of *CsatFT3* and *CsatFD2*. To investigate the role of
CsatSVP2, two experimental conditions were used: (i) standard growth conditions, and
(ii) a temperature-sensitive treatment in which corms were incubated at 8 °C during
the flowering phase to assess temperature-dependent gene regulation. For silencing
of *CsatTFL1-3/CEN1*, corms with lower flowering competency (weighing 7–9 g) were
utilized.
Arabidopsis thaliana (Col-0) was used as the wild-type background for transformation
experiments, and the ft1 mutant in the Ler background was used for functional
complementation of CsatFT3. *Nicotiana benthamiana* was used for transient
expression assays. Arabidopsis seeds were surface-sterilized with 4% sodium
hypochlorite and germinated on half-strength Murashige and Skoog (MS) medium. To
synchronize germination, seeds were stratified at 4°C for two days before being
transferred to a soil mixture of peat, vermiculite, and perlite (2:1:1). All plants were
cultivated in controlled growth chambers under long-day conditions (16 h light/8 h
dark) at 22°C for Arabidopsis and 25°C for *Nicotiana benthamiana*.
186
RNA isolation and expression analysis by RT-qPCR
Total RNA was isolated from saffron apical buds, axillary buds, corm tissues, and
Arabidopsis using the Plant Total RNA Isolation Kit (Sigma-Aldrich), according to the
manufacturer's protocol. For each sample, 1 μg of total RNA was used to synthesize
cDNA. Reverse transcription was performed using the RevertAid cDNA Synthesis Kit.
The transcript level of each gene was quantified by quantitative real-time PCR (q-PCR)
using gene-specific primers (Supplementary Table 1), generated by SnapGene
Viewer 5.3.1 software (version 0.4.0) using the System (BioRad CFX Opus Real time
PCR). Each expression analysis included three biological replicates, with each
replicate derived from pooled tissue of ten individual corms. RT-qPCR results are
presented as the mean ± standard error (SEM) of three technical replicates per
biological sample.
199
200 **Sequence alignments and phylogenetic analysis**
201 Protein sequences were aligned using the ClustalW algorithm with default parameters
202 to generate high-quality multiple sequence alignments. Visualization of the alignments

was performed using ESPript 3.x (Robert & Gouet, 2014) to highlight conserved regions and structural features. The aligned sequences were exported in Newick (.nwk) format for phylogenetic analysis. Phylogenetic tree construction was carried out using the "build" function of ETE3 v3.1.3 (Huerta-Cepas et al., 2016), implemented via the GenomeNet platform (https://www.genome.jp/tools/ete/). FastTree v2.1.8 (Price et al., 2010) was used with default settings to infer approximately maximum-likelihood phylogenetic trees.

Plant transformation

To generate Arabidopsis transgenic lines, full-length coding sequences (CDS) of the target genes were amplified and first cloned into the pENTR entry vector using Gateway® cloning technology. Subsequently, LR recombination was performed to transfer the CDS into the destination vectors pK2GW7 (for overexpression analysis), under the control of the Cauliflower Mosaic Virus (CaMV) 35S promoter. The resulting constructs included 35S::*CsatFD1*, 35S::*CsatFD2*, 35S::*CsatFD3*, 35S::*CsatFT3*, 35S::*CsatTFL1-1*, 35S::*CsatTFL1-2*, 35S::*CsatTFL1-3*, 35S::*CsatSVP1*, and 35S::*CsatSVP2*. Primer sequences used for vector construction are provided in the supplementary table 1 file. The final recombinant plasmids were introduced into *Arabidopsis thaliana* (Col-0) via Agrobacterium tumefaciens strain GV3101 using the floral dip transformation method. Transgenic seeds were harvested individually and screened based on the appropriate selection marker. Homozygous T3 lines were identified and used for further phenotypic characterization. Flowering time was recorded as the number of days from germination to the emergence of the first floral bud.

Yeast one hybrid (Y1H) assay

To examine the binding of CsatSVP2 to the CsatFT3 promoter, yeast one-hybrid (Y1H) assays were conducted using the Matchmaker® Gold Yeast One-Hybrid System Kit (Clontech). Promoter fragments of CsatFT3 containing the core CArG-box motif were cloned into the pAbAi vector. The resulting recombinant pAbAi-promoter plasmids were linearized with BstBI (NEB) and integrated into the genome of the Y1HGold yeast strain. Transformants were selected on synthetic dextrose (SD) medium lacking uracil (SD/-Ura). The coding sequence of CsatSVP2 was amplified and inserted into the pGADT7-AD vector. The recombinant pGADT7-AD constructs were then introduced

into the Y1HGold strains containing the integrated CsatFT3 promoter. Protein–DNA
interaction was assessed by culturing the transformed yeast on SD medium lacking
leucine (SD/-Leu) and supplemented with Aureobasidin A (AbA) at a minimal inhibitory
concentration of 150 ng/mL for four days. Primer sequences used for cloning are
provided in Supplementary Table 1.

Histochemical localization of GUS activity
GUS activity was performed using the histochemical staining protocol described by
Jefferson(Jefferson et al., 1987). Transgenic *Arabidopsis thaliana* lines with
ProCsatFT3::GUS construct were incubated in a staining solution composed of 50 mM
sodium phosphate buffer (pH 7.2), 2.0 mM potassium ferricyanide [$K_3Fe(CN)_6$], 2.0
mM potassium ferrocyanide [$K_4Fe(CN)_6$], 0.1% (v/v) and 1.0 mg/mL X-Gluc (5-bromo-
4-chloro-3-indolyl-β-D-glucuronide). Samples were incubated at 37 °C for 5 hours to
allow GUS expression to develop. Following staining, tissues were cleared with 70%
(v/v) ethanol to remove chlorophyll and enhance contrast, then examined under a
stereo microscope (ZEISS Stemi 508). To validate GUS expression patterns, at least
ten independent transgenic lines were screened for the construct. Plants from three
confirmed GUS-positive lines were selected for detailed expression analysis at various
developmental stages, including 7, 14, and 21 days after germination (DAG),
inflorescence emergence, flowering, and silique formation. For the flowering and
silique stages, staining was performed on aerial tissues from soil-grown plants

Yeast two-hybrid (Y2H) assays
Yeast two-hybrid assays were conducted following the manufacturer's protocol using
the Matchmaker™ Two-Hybrid System (Clontech). The coding sequences of
CsatFD1, CsatFD2, and CsatFD3 were cloned into the bait vector pGBKT7, whereas
those of CsatFT3, CsatTFL1-1, CsatTFL1-2, and CsatTFL1-3 were inserted into the
prey vector pGADT7. The primers used for amplification are provided below. Bait and
prey plasmids were co-transformed into the Y2HGold yeast strain and cultured on
synthetic dextrose (SD) medium lacking tryptophan (SD/-Trp) and leucine (SD/-Leu)
for 3–5 days at 30°C. Protein-protein interactions were assessed by selecting colonies
on SD medium lacking Trp, Leu, and histidine (SD/-Trp/-Leu/-His).

Bimolecular fluorescence complementation (BiFC)

The coding sequences of *CsatFD1*, *CsatFD2*, *CsatFD3* Cloned into YFP C-terminal
(YFPC) and *CsatFT3*, *CsatTFL1-1*, *CsatTFL1-2*, *CsatTFL1-3* were cloned into the
YFP N-terminal (YFPN) vectors. Primer sequences used for vector construction are
provided in the supplementary table 1. These constructs were then introduced into
Agrobacterium tumefaciens strain GV3101 for transient expression in 5-week-old
Nicotiana benthamiana leaves. After co-infiltration for 48 hours, fluorescence was
imaged using a confocal laser scanning microscope (Leica TCS SP5, Germany).YFP
fluorescence was excited at 514 nm, with emission detected between 522 and 560
nm.
280

Firefly Complementation Assay
The full-length coding sequences (CDSs) of CsatFT3, CsatTFL1-3, and CsatFD2 were
cloned into the pCAMBIA1300-nLUC and pCAMBIA1300-cLUC vectors, respectively,
and introduced into Agrobacterium tumefaciens strain GV3101. Various combinations
were co-infiltrated into Nicotiana benthamiana leaves. The infiltrated plants were
incubated at 22 °C for two days. Prior to LUC activity detection, the leaves were
sprayed with 1 mM D-luciferin (Potassium Salt, BioVision) and incubated in the dark
for 8 minutes to eliminate background fluorescence. Fluorescence was detected using
a cooled CCD camera system (Bio-Rad Chemi Doc). Relative LUC intensity was
quantified using ImageJ software. Statistical analysis was performed using one-way
ANOVA, and significance was assessed using the Compact Letter Display (CLD)
method to indicate differences at a significant p-value level. Primer sequences used
in these assays are provided in Supplementary Table 1.
294

Genome walker library for promoter amplification
The 5′ upstream region of the CsatFT3 gene from Crocus sativus was obtained using
the Genome Walker™ kit (Clontech, USA). Genomic DNA was extracted from leaf
tissue following the protocol provided with the QIAGEN DNA isolation kit, then
digested with two blunt-end restriction enzymes, DraI and EcoRV, and further ligated
with adapters, to generate Genome Walker libraries. Gene-specific primers (GSPs),
designed based on the CsatFT3 cDNA sequence, were used alongside adaptor
primers from the kit to carry out primary and nested PCR amplifications as listed in
Supplementary Table 1. The amplified PCR products were resolved on a 1% (w/v)
agarose gel, purified, and cloned into the pGEM-T Easy vector (Promega, USA) for

sequencing. The resulting CsatFT3 promoter sequence was analyzed using the PlantCARE database (Lescot et al., 2002) to identify potential cis-regulatory elements involved in transcriptional regulation.

Virus-induced gene silencing (VIGS)

Virus-induced gene silencing (VIGS) was carried out following the protocol established in lab (Kalia et al., 2024). Gene-specific fragments were amplified for *CsatFT3* (300 bp, flanked by EcoRI sites), *CsatFD2* (292 bp, flanked by BamHI sites), *CsatTFL1-3* (302 bp, containing both EcoRI and BamHI sites), and *CsatSVP2* (318 bp, with EcoR1and BamH1 sites) restriction sites at both termini. These fragments were cloned into the pTRV2 vector to generate the constructs. Agrobacterium-mediated infiltration was performed as previously described (Kalia et al., 2024). To confirm gene silencing, transcript levels of *CsatFT3*, *CsatFD2*, *CsatTFL1-3*, and *CsatSVP2* were quantified by RT-qPCR in apical buds collected at the floral induction stage. Phenotypic analysis of flower development was conducted when floral buds reached the floral emergence stage. Primer sequences used for the VIGS assay are listed in Supplementary table 1.

Luciferase assay

To investigate the regulatory effect of *CsatSVP2* on the *CsatFT3* promoter, a luciferase reporter assay was conducted in *Nicotiana benthamiana* leaves. Promoter sequences of *CsatFT3* were cloned into the Gateway-compatible vector pGWB435 (Invitrogen) upstream of the luciferase (Luc) reporter gene. Three promoter constructs were generated: (i) the full-length promoter (FL), (ii) a truncated fragment containing the CArG motif(T1), and (iii) a truncated fragment lacking the CArG motif (T2). The *CsatSVP2* coding sequence was separately cloned into pGWB402 under the control of the CaMV35S promoter to serve as an effector. Control treatments included empty vectors (pGWB435 and pGWB402) as a negative control, and *ProCsatFT3*::Luc alone as a positive control. Constructs were introduced into the *Agrobacterium tumefaciens* strain GV3101 and co-infiltrated into *Nicotiana benthamiana* leaves. Each leaf was divided into four quadrants to enable side-by-side comparisons of different treatments within the same biological context. Luciferase activity was visualized and quantified using ImageJ software. The experiment was repeated independently at least three times in separate leaves to ensure reproducibility. One-way ANOVA was used for

339 statistical analysis, and differences among treatments were determined using
340 Compact Letter Display (CLD) to indicate statistically significant differences (p < 0.05).
341

Statistical analyses
All statistical analyses were conducted using GraphPad Prism version 10 for Windows
(GraphPad Software, Boston, MA, USA; www.graphpad.com). One-way ANOVA
followed by Tukey's Honest Significant Difference (HSD) test was applied for multiple
comparisons. Statistically significant differences between groups at p < 0.05 are
indicated by different letters, as determined by Compact Letter Display (CLD).Two-
way ANOVA to assess the effects of temperature, month, and their interaction,
followed by Sidak's multiple comparisons test and Student's t-test for one-to-one
comparison. (*: p<0.05; **: p<0.01***: p<0.001) (****: p<0.0001).
351

Results:
CsatFT3 functions as a temperature-sensitive regulator of floral induction in
saffron
Our previous work identified several PEBP family genes implicated in flowering
regulation in saffron, with *CsatFT3* emerging as a potential key regulator of floral
induction (Kalia et al., 2023). However, that study was limited to spatial and temporal
expression profiling during the reproductive phase transition. To further elucidate the
functional role of *CsatFT3*, we investigated its expression dynamics across vegetative
and reproductive phases in saffron apical buds. We found that *CsatFT3* is specifically
expressed during the reproductive phase, with transcript accumulation coinciding with
the floral induction stage (Figure 1A, Supplementary Figure 1). In contrast, its
expression was barely detectable in apices of newly developing corms and during the
vegetative phase in winter. Given that low temperatures are known to suppress floral
induction in saffron (Jose-Santhi et al., 2023), we next assessed *CsatFT3* expression
in apical buds of corms stored under low (8°C) and ambient (25°C) temperature
conditions following the vegetative phase. Consistent with a temperature-responsive
role, *CsatFT3* transcript levels were markedly reduced in corms stored at 8°C
compared to those maintained at 25°C (Figure 1B).
To further investigate tissue- and temperature-specific expression of the *CsatFT3*
gene, we isolated a 1.2 kb fragment of the *CsatFT3* promoter (Supplementary figure
2) that contains several core, distal, and proximal promoter elements and constructed

a promoter-GUS fusion construct. GUS-driven transcription reporter lines of *CsatFT3*
displayed meristem and flower-specific expression in Arabidopsis, consistent with our
spatiotemporal expression profiling in saffron (Figure 1C and Supplementary figure 3).
Furthermore, these *ProCsatFT3::GUS* fusion lines demonstrated temperature-
sensitive regulation, with expression gradually decreasing as the temperature was
lowered, in line with our previous findings that low temperatures suppress the
expression of *CsatFT3* (Figure 1D). At low temperatures (4°C), GUS activity was
nearly undetectable, indicating a significant suppression of expression under cold
conditions. These findings reinforce the role of *CsatFT3* in temperature-sensitive
flowering regulation in saffron and suggest that its expression is activated under the
relatively high-temperature conditions that promote floral induction.
To functionally validate the role of *CsatFT3* in floral regulation, we ectopically
expressed *CsatFT3* in Arabidopsis Col-0 and complemented the late-flowering ft-10
mutant (Ler-0). Overexpression in Col-0 resulted in a significant early-flowering
phenotype (Figure 1E–G), while ft-10 plants transformed with *CsatFT3* exhibited
restored or even earlier flowering compared to wild-type (Figure 1H–J). These results
confirm that *CsatFT3* is functionally conserved and capable of promoting flowering in
a heterologous system. Following confirmation of CsatFT3's involvement in floral
induction and flowering, finally, to evaluate the role of *CsatFT3* in its native we
performed Virus-Induced Gene Silencing (VIGS) of *CsatFT3* in saffron corms and
monitored flowering. Silencing of *CsatFT3* led to a significant reduction in its transcript
levels in apical buds (Figure 1K) and was accompanied by a marked decrease in
flowering percentage compared to TRV2 mock-infected controls (Figure 1L–M). No
significant decrease of *CsatFT1* and *CsatFT2* was observed in TRV2-silenced
CsatFT3 corms (Supplementary Figure 4). Together, these findings establish CsatFT3
as a temperature-sensitive floral inducer in saffron, functioning as a key regulatory
component during the transition from vegetative to reproductive growth.
400

Identification of florigen activation complex component FD, involved in
flowering induction in saffron
FT like genes requires interaction with bZIP transcription factor-*FD* to form FT-FD
complex that directly activates the floral meristem identity genes. Thus, to identify the
FD like genes involved in floral induction of saffron we have cloned full length cDNAs
for three FD-like genes (*CsatFD1*, *CsatFD2* and *CsatFD3*), which encodes for 197,

215 and 245 aa of proteins respectively. All the three FDs are similar to FD proteins from other plants and contains the conserved basic region, Leucine zipper region and the functionally important conserved threonine (T)/SAP motif at the C terminus (Supplementary Figure 5A). Phylogeny aligns CsatFD's with monocot FDs with Asparagus officinalis as the nearest family member (Supplementary Figure 5B).

We next investigated the expression profiles of the three saffron *FD* genes during the vegetative and flowering induction stages. Interestingly, the expression of *CsatFD2* coincided with the floral induction stage, whereas *CsatFD3* showed higher expression at later stages of sprouting, specifically in July, when floral differentiation and vegetative growth begin (Figure 2 A). In contrast, *CsatFD1* was expressed at relatively stable levels throughout flower initiation and the vegetative phase, but its expression was generally lower compared to *CsatFD2* and *CsatFD3*. In saffron, the apical bud gives rise to the flower, while the axillary buds contribute only to vegetative growth, thereby distinguishing the reproductive meristem from the vegetative meristems (Kalia et al., 2022). Notably, *CsatFD2* was predominantly expressed in apical buds and flower tissues, indicating a primary role in flowering (Figure 2B, C). Additionally, *CsatFD2* showed comparatively higher expression in flower tissues compared to the other two FD genes (Figure 2C). We also examined the temperature-sensitive expression of these genes in corms stored at low and ambient high temperatures. Our results showed that only *CsatFD2* expression was significantly downregulated at low temperatures (Figure 2D), while the expression of *CsatFD1* and *CsatFD3* did not exhibit a similar pattern (Supplementary Figure 6A). Interestingly, *CsatFD3* was highly expressed in corms stored at low temperatures (Supplementary Figure 6B).

Next, we examined the effect of ectopic expression of the saffron *FD* genes in Arabidopsis. Ectopic expression of *CsatFD2* can only induce early flowering in Arabidopsis, whereas there was no significant phenotypic change observed in *CsatFD1* and *CsatFD3* expressing Arabidopsis plants (Figure 2E). These findings were further supported by measurements of days to flowering and the number of rosette leaves in the Arabidopsis transformants (Figures 2F-K). These results further suggest that *CsatFD2* may be the primary gene involved in flowering regulation in saffron. To confirm the role of *CsatFD2* during floral induction, we silenced the gene using Virus-Induced Gene Silencing (VIGS) in flower-competent saffron corms. Compared to the TRV2 control, silencing of *CsatFD2* resulted in a reduction in flower formation and overall flowering percentage (Figure 2L). VIGS silencing of CsatFD2

also led to a significant decrease in its transcript levels (Figure 2M), but did not affect

the expression of *CsatFD1* and *CsatFD3* (Supplementary Figure 7 A and B). These

results suggest that *CsatFD2* is a key component of the Floral Activator Complex

(FAC) involved in floral induction in saffron.

TFL1-3/CEN1 is a newly identified negative regulator of flowering induction in

saffron

In our previous study (Kalia et al., 2023), we identified two homologs of *TFL-1/CEN*,

named Csat*TFL1-1* and Csat*TFL1-2*, and through spatio-temporal expression

profiling, we proposed that TFL1-2 may play a role in determining flowering

competency in saffron (Supplementary figure 8A and B). However, neither of these

two *TFL-1* genes exhibited a temperature-sensitive response (Supplementary Figure

9A and B). Furthermore, ectopic expression of *CsatTFL1-1* and *CsatTFL1-2* in

Arabidopsis led to delayed bolting in only *CsatTFL1-2* plants compared to wild-type

(WT) under long-day (LD) conditions, whereas *CsatTFL1-1* transformants bolted

normally, similar to WT plants (Supplementary figure 10A-F). Additionally, interaction

studies, including BiFC and Y2H assays, did not reveal any positive interactions with

CsatFD2, our identified positive regulator of flowering (Supplementary Figure 11A and

B). These findings prompted us to investigate further, hypothesizing that another

homolog of TFL1/CEN might be involved in floral induction in saffron. Through corm

development studies in our lab (Jose-Santhi et al., 2023), we identified a third *TFL1*

homolog, which exhibited higher expression during the vegetative phase. This gene

shares similarities with *CsatTFL1-1* and *CsatTFL1-2*, but phylogenetic analysis

clustered it within the CEN subgroup. Based on this, we named it *CsatTFL1-3/CEN1*

(Supplementary Figure 12A and B). Like other TFL1/CEN family members, TFL1-

3/CEN1 contains the conserved domains characteristic of this gene family.

Quantitative PCR (qPCR) expression profiling revealed that *CsatTFL1-3/CEN1*

expression is elevated during the vegetative phase, with a significant decrease during

the reproductive phase (Figure 3A). Tissue-specific expression analysis confirmed

that *CsatTFL1-3/CEN1* is predominantly expressed in the axillary bud tissue, further

supporting its potential role in negatively regulating flowering (Figure 3 C and D).

Next, we examined the expression of *CsatTFL1-3/CEN1* in corms stored at low (8°C)

and ambient (25°C) temperatures. Interestingly, there was no reduction in *TFL1-*

3/CEN1 expression in corms stored at low temperatures compared to those stored at

ambient high temperatures, suggesting that it may be regulated by temperature and
could play a role in temperature-mediated floral induction in saffron (Figure 3B). We
then performed ectopic expression of *CsatTFL1-3/CEN1* in *Arabidopsis thaliana* (Col-
0) under the control of the constitutive 35S promoter. Transgenic lines exhibited a
significant delay in flowering compared to wild-type plants, as evidenced by increased
days to bolting and a higher number of rosette leaves at flowering (Figure 3F-G).
These results confirm the ability of TFL1-3/CEN1 to act as a floral repressor in
Arabidopsis, consistent with its proposed function in saffron. Notably, ectopic
expression of TFL1-3/CEN1 also led to a hyper-vegetative shoot phenotype,
characterized by the development of an inflorescence composed of leaf primordia
surrounded by small, serrated leaves—a phenotype not observed in TFL1-1 or TFL1-
2 expressing Arabidopsis lines (Figure 3E). Finally, we performed Virus-Induced Gene
Silencing (VIGS) of *CsatTFL1-3/CEN1* in low-flower-competent saffron corms. In
comparison to TRV2 control plants, silencing of *CsatTFL1-3/CEN1* led to the formation
of floral structures in 7-9 g corms (classified as low-flower-competent), and these
flowers appeared earlier than in TRV2 controls (Figure 3G-I). Importantly, VIGS of
CsatTFL1-3/CEN1 did not affect the transcript levels of the other two *TFL1* homologs,
TFL1-1 and TFL1-2 (Supplementary Figure 13 A and B), suggesting TFL1-3/CEN1
and not TFL1-1 or TFL1-2 could play a role in flowering in saffron. This observation
suggests that TFL1-3/CEN1 acts as a negative regulator of floral induction, and its
silencing promotes floral transition in otherwise low-flowering competence corms.
496
The FT-FD-TFL complex involved in floral regulation
Our earlier analyses identified *CsatFT3*, *CsatFD2*, and *CsatTFL1-3/CEN1* as potential
core components of a Floral Activator Complex (FAC) in saffron. To investigate
whether these proteins physically interact to form functional regulatory complexes,
we employed two complementary approaches: yeast two-hybrid (Y2H) assays and
bimolecular fluorescence complementation (BiFC) in *Nicotiana benthamiana*. In the
Y2H assays, the coding sequences (CDS) of the three *CsatFD* genes were cloned into
the pGADT7 prey vector, and co-transformed into yeast with either *CsatFT3* or
CsatTFL1-3/CEN1 fused to the pGBKT7 bait vector. Interaction screening revealed
that both CsatFT3 and TFL1-3/CEN1 specifically interacted with CsatFD2, but not with
CsatFD1 or CsatFD3 (Figure 4 A and B). These results suggest that CsatFD2 may

serve as a central interacting hub for both activator (FT3) and repressor (TFL1-3)
proteins within the FAC.

We further validated these interactions by BiFC assays in *Nicotiana benthamiana*
leaves. YFP fluorescence was observed in nuclei of cells co-expressing the C-terminal
half of YFP fused to CsatFD2 (cYFP–CsatFD2) with the N-terminal half of YFP fused
to either CsatFT3 (nYFP–CsatFT3) or CsatTFL1-3 (nYFP–CsatTFL1-3) (Figure 4 C
and D). No YFP fluorescence was detected in negative controls, including co-
expression of cYFP–CsatFD2 with nYFP alone or cYFP alone with nYFP–CsatFT3 or
CsatTFL1-3. Furthermore, consistent with the Y2H results, no BiFC signal was
observed when CsatFT3 or TFL1-3/CEN1 were co-expressed with CsatFD1 or
CsatFD3 (Supplementary Figure). Together, these findings strongly support a model
in which CsatFD2 specifically interacts with both CsatFT3 and TFL1-3/CEN1,
suggesting that competitive complex formation may modulate floral transition in
saffron.

Given the antagonistic roles of CsatFT3 and CsatTFL1-3 in the regulation of flowering,
and their respective interactions with the transcription factor CsatFD2, it was
hypothesized that CsatFT3 and CsatTFL1-3 may compete for binding to CsatFD2. To
investigate this, a luciferase complementation imaging (LCI) assay was conducted in
Nicotiana benthamiana leaves to evaluate the interaction strength between CsatFD2-
nLUC and cLUC-CsatTFL1-3 in the presence or absence of SK-CsatFT3.Strong
luciferase (LUC) activity was observed when CsatFD2-nLUC and cLUC-CsatTFL1-3
were coexpressed, indicating a robust interaction. However, coexpression with SK-
CsatFT3 significantly reduced LUC activity, suggesting that CsatFT3 interferes with
the interaction between CsatTFL1-3 and CsatFD2. Moreover, this suppression was
dose-dependent, with increasing amounts of CsatFT3 leading to a progressive
decrease in LUC signal (Figure 4E). Biological replicates supporting these
observations are presented in Supplementary figure 15 A and B. These results
strongly suggest that CsatFT3 competes with CsatTFL1-3 for interaction with
CsatFD2, which interferes with CsatTFL1-3's ability to interact with CsatFD2.

CsatSVP2 Acts Upstream of FT3 in Temperature-Mediated Floral Repression
To identify upstream genes potentially involved in temperature-dependent floral
induction in saffron, we screened transcriptome data representing the suppression of
floral induction under low temperatures (Jose-Santhi et al., 2023) and identified two

partial SVP-like MADS-box genes that were upregulated in non-flowering samples.
SVP-like genes are known to regulate flowering in a temperature-dependent manner
(Lee et al., 2007). Using long-read and other in-house transcriptome data, and based
on sequence similarity, we obtained full-length sequences of these two SVP-like
genes. We isolated and cloned them from saffron, naming them *CsatSVP1* and
CsatSVP2. Sequence analysis indicated that while these genes are highly similar, they
code for different amino acid sequences. Multiple sequence alignment revealed that
both CsatSVP1 and CsatSVP2 contain the conserved MADS-box domain, and
phylogenetic analysis grouped them with SVP-like genes from other geophytes, such
as Narcissus, Zingiber, and Lilium (Supplementary Figure 16 A and B).

To further characterize their role in flowering, we performed expression analysis of
CsatSVP1 and *CsatSVP2* during vegetative and floral induction stages. Both genes
showed a significant reduction in expression during the floral induction stage (June);
however, the reduction in *CsatSVP2* was more pronounced and correlated closely with
floral induction and subsequent stages (Figure 5A). We then evaluated their
expression in corms stored under floral inductive (25°C) and non-inductive (8°C)
conditions. *CsatSVP2* transcript levels remained high at both temperatures,
suggesting it may contribute to the floral repression observed at low temperatures
(Figure 5C). In contrast, *CsatSVP1* expression decreased under both conditions,
showing no temperature-dependent pattern (Figure 5B). Additionally, *CsatSVP2*
expression was comparatively higher in floral tissues than *CsatSVP1*, further
supporting its potential regulatory role (Supplementary Figure 17).

Next, to functionally validate these genes, we ectopically expressed both *CsatSVP1*
and *CsatSVP2* in Arabidopsis under a constitutive promoter. Ectopic expression of
both genes resulted in delayed flowering, with a more pronounced delay in the
CsatSVP2-transformed lines compared to both the control and *CsatSVP1* lines (Figure
5D-I). Arabidopsis plants expressing *CsatSVP2* displayed various phenotypic changes
in flower bud and flower morphology, producing smaller, more compact flowers and
shorter siliques (Supplementary Figure 18). To investigate the role of *SVP* in
temperature-mediated floral suppression, we silenced *CsatSVP2* in saffron corms
using Virus-Induced Gene Silencing (VIGS). Csat*SVP2* was selected due to its
expression pattern and phenotype in Arabidopsis, which correlated with its expected
function. Since low temperatures during storage or dormancy are known to suppress
flowering, and CsatSVP2 may be involved in this process, we stored the *CsatSVP2-*

silenced corms at low temperatures, alongside a mock control (TRV2). Silencing of
SVP2 resulted in flowering in corms at low temperatures, whereas the TRV2-treated
corms stored at low temperatures did not flower (Figure 5J). In parallel, corms stored
at 25°C—an inductive condition—showed flowering in both control and SVP2-silenced
groups, confirming that *CsatSVP2* acts specifically in the floral suppression
mechanism under cold conditions (Figure 5K). Gene expression analysis further
revealed that *CsatSVP2* silencing prevented the typical low-temperature-mediated
suppression of *CsatFT3* expression, thereby facilitating flowering (Figure 5P and Q).
These findings suggest that CsatSVP2 and CsatFT3 act within the same regulatory
pathway, mediating temperature-responsive floral induction in saffron.
586
SVP2 Directly Binds to the CsatFT3 Promoter and Represses Its Expression
Our previous results indicated that CsatSVP2 and CsatFT3 function in the same
temperature-sensitive regulatory pathway: low temperature maintains or upregulates
CsatSVP2 expression, which in turn suppresses *CsatFT3*, thereby repressing floral
induction. Notably, *CsatFT3* transcript levels were restored in *CsatSVP2*-silenced
corms, suggesting that CsatSVP2 functions upstream of *CsatFT3*. To further explore
the mechanism of this regulation, we analyzed the CsatFT3 promoter sequence to
identify potential binding motifs for MADS-box transcription factors. In silico analysis
of the 1.2 kb FT3 promoter, previously shown to drive temperature-responsive
expression, revealed a conserved CArG-box motif (CCATTTAAGG) located
approximately 827 bp upstream of the ATG start codon that is established for binding
of MADS boxes genes including SVP orthologs (Figure 6A). To experimentally test
further whether CsatSVP2 directly binds to the *CsatFT3* promoter, we performed a
yeast one-hybrid (Y1H) assay. The Y1H assay confirmed that CsatSVP2 specifically
binds to the *CsatFT3* promoter, supporting the hypothesis of direct regulation (Figure
6B). To further validate the functional specificity of this interaction, we employed a
luciferase reporter assay using *Nicotiana benthamiana* transient expression (Figure
6C) and conducted a luciferase reporter assay using three different FT3 promoter
fragments: (1) the full-length promoter (~1.2kb), (2) a truncated version containing the
conserved CArG motif (T1), and (3) a truncated version lacking the motif (T2) (Figure
6D). Co-expression of CsatSVP2 with these constructs in *Nicotiana benthamiana*
revealed that luciferase activity was significantly reduced in both the full-length and T1
constructs (Figure6 D and F). In contrast, no repression was observed with the T2

construct, confirming that SVP2 specifically suppresses FT3 expression through direct
binding to the CArG motif (Figure 6F). Together, these results establish a mechanistic
model in which CsatSVP2 acts as a temperature-responsive floral repressor that
directly binds to the FT3 promoter via a conserved CArG motif, thereby repressing
FT3 expression and floral induction under low-temperature conditions.
615

Discussion
Saffron is a high-value crop cultivated in limited regions due to its strict temperature
requirements, which distinctly regulate its vegetative and reproductive growth. High
temperatures induce flowering, while subsequent cooling triggers flower formation. In
contrast, low temperatures suppress floral induction and support vegetative growth,
enabling overwintering and daughter corm development (Molina et al., 2005a). This
temperature sensitivity restricts saffron cultivation to specific climates. Although
environmental control of flowering has enabled cultivation in soilless and controlled
settings, the molecular mechanisms underlying temperature-regulated flowering
remain poorly understood. As a sterile crop, saffron cannot be improved through
conventional breeding, making biotechnological interventions essential. However,
limited genomic and genetic resources, challenges in genetic transformation, and a
scarcity of molecular studies have hindered progress in functional genomics—creating
a critical knowledge gap that must be addressed.
In this study, we have identified a temperature-sensitive transcriptional network that
regulates flowering in saffron, uncovering both conserved and saffron-specific
components involved in floral induction (Figure 7). Our data establish a regulatory
module involving CsatFT3, CsatFD2, CsatTFL1-3/CEN1, and CsatSVP2 as key
components in mediating floral transition in response to temperature cues. Although
the core elements of the Florigen Activation Complex (FAC)—FT, FD, and TFL1—are
conserved across angiosperms, their expression patterns, protein interactions, and
functional dynamics in saffron suggest neofunctionalization in line with its geophytic
lifecycle. For instance, Notably, CsatFT3 and CsatFD2 are upregulated at higher
temperatures and co-expressed in the floral meristem, promoting flowering. In
contrast, other FT and FD homologs show no such expression correlation (Figure 2;
(Kalia et al., 2022), indicating their limited or no involvement in floral induction.
Similarly, CsatTFL1-3 functions as a floral repressor by competitively binding to
CsatFD2, whereas the other two TFL1 homologs neither interact with CsatFD2 nor

delay flowering in Arabidopsis, suggesting a lack of functional redundancy and

functional specificity. Functional validation in both Arabidopsis and saffron confirms

that floral induction is governed by the dynamic interplay between CsatFT3, CsatFD2,

and CsatTFL1-3.Given the roles of gene duplication, neofunctionalization, and

lineage-specific gene expansion in driving the specialization of reproductive and

vegetative pathways in geophytes (Khosa et al., 2021), our identification of distinct

regulatory factors and their molecular interactors from multigene families in saffron

represents a significant advance.

A distinguishing feature of floral induction observed in saffron is the spatial restriction

of flowering-related gene expression predominantly to the meristem. All the genes

identified in this study were specifically expressed in the apical (flowering) meristem.

During floral induction, saffron corms are either underground or stored in the dark and

remain leafless, indicating that the molecular signals initiating flowering are localized

within the apical buds. Such meristem-localized control may reflect an adaptation to

the geophytic habit of saffron, which undergoes floral induction in the absence of

foliage, similar to what has been observed in other geophytes such as tulips

(Leeggangers et al., 2017b). This localization supports the hypothesis that

temperature cues are sensed and interpreted locally within the corm, rather than

systemically via leaves, as observed in many other species, including some geophytes

(Lee et al., 2013b; Navarro et al., 2011; Wigge, 2011; Wu et al., 2022; Yan et al.,

2021).

Temperature plays a pivotal role in regulating flowering time across plant species,

often through conserved thermosensory pathways. In Arabidopsis, warmer

temperatures promote flowering via *FT* activation (Balasubramanian et al., 2006).

while in crops like strawberry and onion, cold exposure induces flowering by triggering

FT-like genes (Heide, 1977; Lee et al., 2013b). Interestingly, the effect of temperature

on flowering is not always straightforward. Temperatures above 25°C seem to have

both inductive and repressive effects on flowering, depending on the species and

environmental conditions (Nakano et al., 2013; Noy-Porat et al., 2013; Oda et al.,

2012). This temperature sensitivity is closely linked to the regulation of FT, a key gene

that integrates various environmental signals and controls the timing of flowering. In

Narcissus tazetta (daffodils), floral induction is triggered by high ambient temperatures

(25°C) through the activation of *NtFT*, while in tulips, *TgTFL1* downregulation and

TgFT2 upregulation contribute to temperature-induced floral induction (Leeggangers

et al., 2017a; Noy-Porat et al., 2013). In contrast, in species like onion and lily, which
flower in spring, vernalization (low-temperature treatment) is required to activate the
expression of *AcFT2* and *LiFT*, respectively (Lee et al., 2013b). In our study, we also
found that *CsatFT3*, a key flowering-time gene in saffron, is temperature-regulated,
with its expression being induced by high ambient temperatures and suppressed by
low temperatures (Figure 1B). Similarly, *CsatFD2*, a partner of *CsatFT3* in the
flowering regulatory network, follows a similar expression pattern, suggesting their
involvement in the temperature-mediated floral induction pathway in saffron.
Interestingly, we observed that *CsatTFL1-3*, a gene that maintains the vegetative state
and prevents early flowering, exhibits a reciprocal expression pattern. In Arabidopsis
low and ambient temperature regulation of flowering involves the gene *TFL1* (Strasser
et al., 2009). Higher expression of *CsatTFL1-3* is observed during the vegetative
phase, which coincides with the low winter temperatures during saffron's growth cycle
(Figure 3A). This expression pattern is crucial for maintaining the vegetative phase
and delaying flowering, as confirmed by the hyper-vegetative shoot phenotype in
TFL1-3 overexpressing Arabidopsis plants (Figure 3) similar to what was   observed
when Arabidopsis *TFL1* is overexpressed (Lee et al., 2019).   In contrast, as
temperatures rise in the summer, the expression of CsatFT3 increases (Figure 1A),
likely triggering the activation of floral pathway genes and the transition to reproductive
development. Overall, a balance between CsatFT3, CsatTFL1-3, and their interaction
with CsatFD2 governs floral induction in a temperature-dependent manner. This
delicate interplay between temperature and gene expression is vital for ensuring
proper timing of flowering in saffron, aligning the plant's reproductive phase with
favourable environmental conditions.

We also identify CsatSVP2, a MADS-box transcription factor, as a critical floral
repressor acting upstream of FT–FD–TFL core module. MADS-box genes are key
regulators of flowering time, with SVP-like genes playing central roles in ambient
temperature-mediated floral control (Jin and Ahn, 2021). CsatSVP2 directly binds a
conserved CArG-box in the CsatFT3 promoter, repressing its expression under low
temperatures. Silencing of SVP2 alleviated low-temperature-induced flowering
suppression and restored *CsatFT3* expression, thereby establishing a direct
mechanistic link between ambient temperature perception and floral induction (Figure
5). In *Arabidopsis thaliana*, a comparable regulatory pathway has been described,
where SVP negatively regulates FT expression by directly binding to a CArG motif in

its promoter (Lee et al., 2007). Additionally, SVP levels are known to increase under

low temperatures and are degraded at higher temperatures, facilitating early flowering

as SVP abundance declines (Blázquez et al., 2002; Lee et al., 2013a; Sureshkumar

et al., 2016). Interestingly, individual members of the SVP gene family have been

shown to play distinct roles in processes such as bud dormancy, flowering initiation,

and floral development reflecting adaptations of developmental timing to local

environmental conditions and seasonal temperature fluctuations. In tulips, gene

expression analyses have similarly identified *TgMADS15/16* and *TgMADS9* as low-

temperature-responsive SVP-like genes involved in floral induction (Lu et al., 2023),

reinforcing the conserved role of this gene family in temperature-mediated flowering

regulation across geophytic species. SVP homologs in Lilium and *Narcissus tazetta*

have also been reported as flowering repressors, though their temperature

responsiveness remains unexplored (Li et al., 2015; Wu et al., 2024). Notably, ectopic

expression of *CsatSVP2* in Arabidopsis caused floral morphology defects, reminiscent

of phenotypes observed with Lilium SVP expression (Wu et al., 2024), suggesting that

these homologs may also contribute to flower development in addition to regulating

flowering time.

Conclusion

Collectively, our findings establish a detailed molecular framework for temperature-

regulated flowering in saffron, centered on the dynamic balance between floral

activators (CsatFT3, CsatFD2) and repressors (CsatTFL1-3, CsatSVP2). The

spatiotemporal expression patterns and protein interactions among these components

provide key insights into how flowering is precisely modulated in response to

environmental temperature fluctuations. By uncovering saffron-specific adaptations

within otherwise conserved gene networks, this study advances our understanding of

floral regulation in geophytes and sterile crops. Importantly, these results lay a

foundation for the development of genetic and biotechnological strategies to improve

saffron yield, synchronize flowering, and enhance resilience to climate variability. Our

work highlights a saffron-specific model of thermoresponsive flowering that builds

upon conserved floral regulatory pathways, yet exhibits functional divergence shaped

by the geophytic lifecycle. Beyond saffron, these findings offer a valuable framework

for translational research in other temperature-sensitive geophytes such as tulip,

Narcissus, Hyacinthus, and Iris, where molecular insights remain scarce (Khodorova

and Boitel-Conti, 2013). By identifying novel regulatory targets, this study opens new avenues for improving flowering and productivity in saffron and related high-value geophytic species.

Acknowledgments

We gratefully acknowledge the financial support from the Council of Scientific and Industrial Research (CSIR) under the Genome Editing Mission for Crop Improvement (MMP25301), the intramural CSIR grant to IHBT (MLP201), the SERB-Start-up Research Grant (SRG/GAP0288), and the Department of Biotechnology (DBT/GAP0307) to Rajesh Kumar Singh. Senior Research Fellowships provided to Diksha Kalia (UGC), Joel Jose-Santhi (CSIR), and Firdous Rasool Sheikh (CSIR) are also gratefully acknowledged. We sincerely thank Dr. Rishikesh Bhalerao, Umeå Plant Science Centre, Swedish University of Agricultural Sciences, Sweden, for his careful reading of the manuscript and for providing constructive feedback, which significantly improved the clarity and quality of the work. Our sincere thanks go to Dr. Rimpy Diman, Technical Officer at CSIR-IHBT, for her valuable assistance with confocal microscopy. This manuscript represents the CSIR-IHBT communication number "5819".

Author contribution statement

R.K.S. conceptualized the research idea and designed the work plan. D.K., J.J.S., and F.R.S. conducted the experiments and collected the data. D.K., J.J.S., and R.K.S. analyzed the data and contributed to writing the manuscript. All authors read and approved the final version of the manuscript.

Conflicts of Interest

The authors declare no conflicts of interest.

Figure Legends

Figure 1. *CsatFT3* is involved in temperature mediated floral induction in saffron. **(A)** CsatFT3 transcript levels in apical buds across developmental stages show specific expression during the reproductive phase. **(B)** Relative expression of CsatFT3 in apical buds of saffron corms stored at ambient (25°C) and low (8°C) temperatures. Data are means ± SEM from 3 biological replications, where n=10 for each replicate. **(C)** Histochemical GUS assay of the *CsatFT3* promoter (proCsatFT3) in transgenic

Arabidopsis in long-day conditions (LD) during various developmental stages. 1.2kb

sequence upstream of the CsatFT3 translation start sites (proCsatFT3) was fused with

uidA (GUS) reporter gene. Different developmental stages include Arabidopsis plants

after 7 Days After Germination (DAG), 14DAG, 21DAG, and 30DAG. FT3 expression

is confined to meristem and inflorescence **(D)** Temperature sensitive GUS activity at

3 different temperatures. The experiment was repeated three times independently with

similar results. **(E-G)** Ectopic expression of CsatFT3 in Arabidopsis resulted in early

flowering phenotype. Statistical significance was analyzed by Student's t-test (*P <

0.05). **(H-J)** Functional complementation of the ft-10 mutant with CsatFT3 restores

flowering. Representative line expressing number of rosette leaves and number of

days to flower. Total leaf number was calculated by combining total rosette leaves.

Data represent a minimum of 10 plants scored for each line. Statistical significance

was analyzed by Student's t-test (*P < 0.05). **(K)** Representative flowering vs. non-

flowering phenotypes in control and CsatFT3-silenced saffron corms. The scale bar

represents 2 cm. **(L)** qRT-PCR validation of CsatFT3 transcript knockdown in saffron

corm apices following VIGS (n=3 technical replicates, 10 independent corms). Error

bars are shown as Standard Error of Mean (SEM) of three technical repeats. **(M)**

Percentage of flowering in VIGS-silenced vs. TRV2 control corms. The data presented

includes the mean percentage of flowering, with error bars representing the Standard

Error of Mean (SEM) of three technical repeats (n=3) and for each replicate, 10 corms

were used. All data in this figure are shown as mean ± SEM. Statistical Significance

for Data in panel A, I, & J are analyzed using one-way ANOVA followed by Tukey's

Honest Significant Difference (HSD) test for multiple comparisons. Different letters

indicate statistically significant differences between groups at p < 0.05, according to

Compact Letter Display (CLD). Data in panels B are analyzed using two-way ANOVA

to assess the effects of temperature, month, and their interaction, followed by Sidak's

multiple comparisons test. Asterisks indicate significant differences between 8°C and

25°C within each month (**** p < 0.0001), F, **G**, **L**, and **M** are analyzed using the

student's t-test (*: p<0.05; **: p<0.01).

Figure 2: Functional characterization of FD-like genes in saffron (A) Expression

profiles of CsatFD1, CsatFD2, and CsatFD3 during vegetative and floral induction

stages in the apical bud by RT-qPCR. Data are means ± SE of 3 independent

replicates. **(B)** Expression patterns of CsatFD1, CsatFD2, and CsatFD3 in the Main

bud (MB), axillary bud (AB), and corm tissue (CT) during the floral induction stage. Values are means ± SE of 3 independent replicates. **(C)** Tissue specific expression of FDs in different parts of saffron plant. **(D)** Expression analyses of *CsatFD* genes in apical buds of corms stored at 25∘ C and 8 during months of May and June **(E)** Phenotypic comparison of early flowering in CsatFD2-expressing Arabidopsis compared to control. **(F-K)** Measurements of days to flowering and rosette leaf number in CsatFD1, CsatFD2, and CsatFD3 expressing Arabidopsis plants. Values are means ± SE (n = 10). Asterisks indicate a significant difference between the transgenic Arabidopsis lines and wild-type plants (Col-0; Student's t-test, $P < 0.05$). **(L)** Phenotypes of CsatFD2-TRV2 silenced lines and TRV2 lines. The scale bar represents 2 cm. **(M)** qRT-PCR validation of CsatFD2 knockdown in saffron corm apices after VIGS (n=3). Error bars are shown as SD of three technical repeats **(N)** Flowering percentage in TRV2 control and CsatFD2 VIGS-silenced saffron corms. The data presented includes the mean percentage of flowering, with error bars representing the standard error of mean (SEM) of three technical repeats (n=3) and for each replicate, 10 corms were used. All data in this figure are shown as mean ± SEM. Statistical Significance for data in panel **A** & **B** are analyzed using two-way ANOVA followed by Tukey's Honest Significant Difference (HSD) test for multiple comparisons. Different letters indicate statistically significant differences between groups at $p < 0.05$, according to Compact Letter Display (CLD). Data in panel **D** are analyzed using two-way ANOVA to assess the effects of temperature, month, and their interaction, followed by Sidak's multiple comparisons test. Asterisks indicate significant differences between 8°C and 25°C within each month (**** $p < 0.0001$), , **F**,**G**,**H**,**I**,**J**,**K**,**M** and **N** are analyzed using the Student's t-test (*: $p<0.05$; **: $p<0.01$).

Figure 3: CsatTFL1-3 is negative regulator of floral induction in saffron (A) Relative expression of CsatTFL1-3 during vegetative and reproductive phase in the apical bud by RT-qPCR. Data are means ± SE of 3 independent replicates. **(B)** Expression analyses of CsatTFL1-3 gene in apical buds of corms stored at 25°C and 8 °C during floral induction time **(C)** Corm tissue specific expression of TFL1-3 during floral induction. **(D)** Tissue Specific Expression of CsatTFL1-3. **(E)**Phenotypic analysis of Arabidopsis Col-0 plants ectopically expressing TFL1-3/CEN1, Hyper vegetive phase phenotype observed in CsatTFL1-3 transgenic lines of Arabidopsis. **(F and G)** Rosette leaves number and number of days to flower of 35S::TFL1-3 transgenic and

control plants respectively. Values are means ± SEM (n = 10). Asterisks indicate a

significant difference between the transgenic Arabidopsis lines and wild-type plants

(Col-0; Student's t-test, P < 0.05). **(H)** Virus-Induced Gene Silencing (VIGS) of TFL1-

3/CEN1 in low-flower-competent saffron corms. The scale bar represents 2 cm. **(I)** The

expression of *CsatTFL1-3* in *CsatTFL1-3*-TRV2 silenced lines and TRV2 lines (n=3).

(J) Comparison of flowering percentages in VIGS-silenced versus control saffron

corms. The data presented includes the mean percentage of flowering of three

technical repeats (n=3) and for each replicate, 10 corms were used. All data in this

figure are shown as mean ± SEM. Statistical Significance for Data in panel **A**,**C** & **D**

are analyzed using one-way ANOVA followed by Tukey's Honest Significant

Difference (HSD) test for multiple comparisons. Different letters indicate statistically

significant differences between groups at $p < 0.05$, according to Compact Letter

Display (CLD). Data in panel B are analyzed using two-way ANOVA to assess the

effects of temperature, month, and their interaction, followed by Sidak's multiple

comparisons test. Asterisks indicate significant differences between 8°C and 25°C

within each month (**** $p < 0.0001$), , F,G,I, & J are analyzed using the Student's t-

test (*: $p<0.05$; **: $p<0.01$).

Figure 4. Interaction analysis of CsatFT3, CsatTFL1-3, and CsatFD2.

(A, B) Yeast two-hybrid (Y2H) assays showing physical interactions between

CsatFT3/CsatTFL1-3 and CsatFD2. The coding sequences of *CsatFT3* and

CsatTFL1-3 were fused to the activation domain in the pGADT7 vector, and *CsatFD2*

was fused to the DNA-binding domain in the pGBKT7 vector. Yeast transformants

were grown on double dropout (DDO) medium lacking tryptophan and leucine

(selection for transformation) and quadruple dropout (QDO) medium lacking

tryptophan, leucine, histidine, and adenine (selection for interaction). pGBKT7-53 and

pGADT7-T were used as positive control; pGBKT7-lam and pGADT7-T interactions

served as the negative control. **(C-D)** Bimolecular fluorescence complementation

(BiFC) assay showing nuclear localization of the interactions between CsatFT3 or

CsatTFL1-3 with CsatFD2 in *Nicotiana benthamiana* leaf epidermal cells. CsatFD2

was fused to the C-terminal half of YFP (cYFP), and CsatFT3 or CsatTFL1-3 were

fused to the N-terminal half of YFP (nYFP). YFP signal indicates interaction. Bars =

20 µm. **(E)** Competitive luciferase complementation assay demonstrating the

competition between CsatFT3 and CsatTFL1-3 for binding to CsatFD2 in *N.*

benthamiana leaves. CsatTFL1 was fused to the N-terminal half of luciferase (nLUC), and CsatFD2 to the C-terminal half (cLUC). Increasing amounts of CsatFT3 (non-tagged, driven by 35S promoter, in SK vector) were co-infiltrated at different ratios. Images represents three independent replications. **(F)** Relative luminescence intensities indicate competition for FD2 binding with TFL1-3 and FT3. Data represented is mean of three replicates. Statistical Significance are analyzed using one-way ANOVA followed by Tukey's Honest Significant Difference (HSD) test for multiple comparisons. Different letters indicate statistically significant differences between groups at $p < 0.05$, according to Compact Letter Display (CLD).

Figure 5: Functional characterization of CsatSVP1 and CsatSVP2 in temperature-dependent floral regulation (A) Expression profiling of CsatSVP1 and CsatSVP2 in apical bud during different vegetative and flowering stages. Data are means ± SE from 3 replications. **(B and C)** Expression of SVP1 and SVP2 genes in corms stored at inductive (25°C) and non-inductive (8°C) temperatures respectively. **(D)** Flowering phenotypes of Arabidopsis lines ectopically expressing CsatSVP1. **(E and F)** Rosette leaves number and number of days to flower of CsatSVP1 transgenic and control plants respectively **(G)** Phenotypic changes of the floral organs in 35S::CsatSVP2 transgenic Arabidopsis plants. **(H-I)** Rosette leaves number and number of days to flower of CsatSVP2 transgenic and control plants respectively. The total leaf number was calculated by combining total rosette leaves. **(J-K)** Silencing of CsatSVP2 by VIGS rescues flowering under low-temperature conditions in saffron. The scale bar represents 2 cm. **(L-M)** qRT-PCR validation of CsatSVP2 knockdown in saffron corm apices after VIGS stored at different temperatures (n=3). Error bars are shown as SD of three technical repeats. **(N-O)** Flowering percentage in TRV2 control and CsatSVP2 VIGS-silenced saffron corms stored at low (8°C) and ambient (23°C) temperatures. The data presented includes the mean percentage of flowering, with error bars representing the standard deviation (SD) of three technical repeats (n=3) and for each replicate, 10 corms were used. **(P-Q)** Relative expression of CsatFT3 in SVP2-silenced versus control corms under different temperature regimes. All data in this figure are shown as mean ± SEM. Statistical Significance for data in panel A are analyzed using two-way ANOVA followed by Tukey's Honest Significant Difference (HSD) test for multiple comparisons. Different letters indicate statistically significant differences between groups at $p < 0.05$, according to Compact Letter Display (CLD).

Data in panel B & C are analyzed using two-way ANOVA to assess the effects of
temperature, month, and their interaction, followed by Sidak's multiple comparisons
test. Asterisks indicate significant differences between 8°C and 25°C within each
month (**** $p < 0.0001$), E, F, H, I, L, M, N, O, P & Q are analyzed using the student's
t-test (*: $p<0.05$; **: $p<0.01$).

Figure 6. CsatSVP2 directly binds to the FT3 promoter via the CArG motif and
represses its transcription (A) Schematic diagram of the 1.2 kb CsatFT3 promoter
showing the location of a conserved CArG-box motif (CCATTTAAGG) approximately
827 bp upstream of the ATG start codon. **(B)** Yeast one-hybrid assay showing
interaction between CsatSVP2 and the FT3 promoter region containing the CArG
motif. Growth on selective media confirms binding specificity. **(C)** Schematic diagram
of the effectors and reporters used in dual-luciferase reporter. **(D)** Schematic of the
dual-luciferase constructs used for transient expression in *Nicotiana benthamiana*: FL
(Full length promoter) T1 (with CArG motif) and T2 (without). **(E)** Relative luciferase
activity in leaves co-expressing CsatSVP2 with the full promoter and T1 fragment
construct of *CsatFT3* gene. **(F)** Relative luciferase activity in leaves co-expressing
CsatSVP2 and CsatFT3 full promoter and with T2 fragment constructs. Significant
repression is observed only in the T1 construct, indicating that CsatSVP2 represses
CsarFT3 transcription through the CArG motif. **(G and H)** Quantification of luciferase
activity. Bars represent mean ± SEM (n = 3). All data in figure are shown as mean ±
SEM. Statistical Significance for Data in panel **F & H** are are analyzed using one-way
ANOVA followed by Tukey's Honest Significant Difference (HSD) test for multiple
comparisons. Different letters indicate statistically significant differences between
groups at $p < 0.05$, according to Compact Letter Display (CLD)

Figure 7. Proposed model of temperature-mediated flowering regulation in
saffron. This schematic model illustrates the temperature-regulated floral induction
pathway in saffron, emphasizing the key roles of CsatFT3, CsatFD2, TFL1-3/CEN1,
and CsatSVP2 in transitioning from vegetative to reproductive growth.   At low
temperatures (e.g., below 10°C), *CsatFT3* expression remains low, preventing the
formation of the Floral Activator Complex (FAC). TFL1-3/CEN1 binds to CsatFD2,
maintaining the shoot apical meristem in a vegetative state and inhibiting flowering.
CsatFD2 expression is minimal, suppressing floral induction. At low temperatures,

CsatSVP2 is upregulated and directly binds to the FT3 promoter at a conserved CArG motif, repressing its expression and preventing FAC formation. As temperatures rise above the threshold (e.g., 25°C), *CsatFT3* expression increases, leading to the formation of the FAC by binding to CsatFD2. This complex activates the expression of floral meristem identity genes and promotes floral development. At the same time, *TFL1-3/CEN1* expression decreases, allowing the activation of CsatFD2 and the initiation of flowering.

Figure 1

994   **Figure 2**

1004   **Figure 3**

Figure 4

Figure 5

1033 **Figure 6**

1034

1035

1036

1037

1038

1039

1040

1041 **Figure 7**

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

References

Abe, M., Kobayashi, Y., Yamamoto, S., Daimon, Y., Yamaguchi, A., Ikeda, Y., Ichinoki, H., Notaguchi, M., Goto, K. and Araki, T. (2005) FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* **309**, 1052-1056.

Ahrazem, O., Rubio-Moraga, A., Nebauer, S.G., Molina, R.V. and Gomez-Gomez, L. (2015) Saffron: Its Phytochemistry, Developmental Processes, and Biotechnological Prospects. *J Agric Food Chem* **63**, 8751-8764.

Andrés, F. and Coupland, G. (2012) The genetic basis of flowering responses to seasonal cues. *Nature Reviews Genetics* **13**, 627-639.

Balasubramanian, S., Sureshkumar, S., Lempe, J. and Weigel, D. (2006) Potent induction of Arabidopsis thaliana flowering by elevated growth temperature. *PLoS genetics* **2**, e106.

Bellinazzo, F., Manders, I., Heidemann, B., Bolanos, M.A., Stouten, E., Busscher, J., Abarca, D., van der Wal, F., Dornelas, M.C. and Angenent, G.C. (2025) Differential growth and flowering capacity of tulip bulbs and the potential involvement of PHOSPHATIDYLETHANOLAMINE-BINDING PROTEINS (PEBPs). *Biology Direct* **20**, 29.

Blázquez, M.A., Trénor, M. and Weigel, D. (2002) Independent control of gibberellin biosynthesis and flowering time by the circadian clock in Arabidopsis. *Plant Physiology* **130**, 1770-1775.

Bowman, J.L., Smyth, D.R. and Meyerowitz, E.M. (2012) The ABC model of flower development: then and now. *Development* **139**, 4095-4098.

Capovilla, G., Schmid, M. and Posé, D. (2015) Control of flowering by ambient temperature. *Journal of Experimental Botany* **66**, 59-69.

Cardone, L., Castronuovo, D., Perniola, M., Cicco, N. and Candido, V. (2020) Saffron (Crocus sativus L.), the king of spices: An overview. *Scientia Horticulturae* **272**, 109560.

Freytes, S.N., Canelo, M. and Cerdán, P.D. (2021) Regulation of flowering time: when and where? *Current Opinion in Plant Biology* **63**, 102049.

Gu, X., Le, C., Wang, Y., Li, Z., Jiang, D., Wang, Y. and He, Y. (2013) Arabidopsis FLC clade members form flowering-repressor complexes coordinating responses to endogenous and environmental cues. *Nature Communications* **4**, 1947.

Heide, O.M. (1977) Photoperiod and temperature interactions in growth and flowering of strawberry. *Physiologia Plantarum* **40**, 21-26.

Hsu, C.-Y., Adams, J.P., Kim, H., No, K., Ma, C., Strauss, S.H., Drnevich, J., Vandervelde, L., Ellis, J.D. and Rice, B.M. (2011) FLOWERING LOCUS T duplication coordinates reproductive and vegetative growth in perennial poplar. *Proceedings of the National Academy of Sciences* **108**, 10756-10761.

Hu, J., Liu, Y., Tang, X., Rao, H., Ren, C., Chen, J., Wu, Q., Jiang, Y., Geng, F. and Pei, J. (2020) Transcriptome profiling of the flowering transition in saffron (Crocus sativus L.). *Scientific Reports* **10**, 9680.

Jefferson, R.A., Kavanagh, T.A. and Bevan, M.W. (1987) GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *Embo j* **6**, 3901-3907.

Jin, S. and Ahn, J.H. (2021) Regulation of flowering time by ambient temperature: repressing the repressors and activating the activators. *New Phytologist* **230**, 938-942.

Jin, S., Nasim, Z., Susila, H. and Ahn, J.H. (2021) Evolution and functional diversification of FLOWERING LOCUS T/TERMINAL FLOWER 1 family genes in plants. *Seminars in Cell & Developmental Biology* **109**, 20-30.

Jing, S., Jiang, P., Sun, X., Yu, L., Wang, E., Qin, J., Zhang, F., Prat, S. and Song, B. (2023) Long-distance control of potato storage organ formation by SELF PRUNING 3D and FLOWERING LOCUS T-like 1. *Plant communications* **4**.

Jose-Santhi, J., Sheikh, F.R., Kalia, D. and Singh, R.K. (2023) Sugar metabolism mediates temperature-dependent flowering induction in saffron (Crocus sativus L.). *Environmental and Experimental Botany* **206**, 105150.

Kalia, D., Jose-Santhi, J., Kumar, R. and Singh, R.K. (2022) Analysis of PEBP Genes in Saffron Identifies a Flowering Locus T Homologue Involved in Flowering Regulation. *Journal of Plant Growth Regulation*, 1-20.

Kalia, D., Jose-Santhi, J., Kumar, R. and Singh, R.K. (2023) Analysis of PEBP genes in saffron identifies a flowering locus T homologue involved in flowering regulation. *Journal of Plant Growth Regulation* **42**, 2486-2505.

Kalia, D., Jose-Santhi, J., Sheikh, F.R., Singh, D. and Singh, R.K. (2024) Tobacco rattle virus-based virus-induced gene silencing (VIGS) as an aid for functional genomics in Saffron (Crocus sativus L.). *Physiol Mol Biol Plants* **30**, 749-755.

Khodorova, N.V. and Boitel-Conti, M. (2013) The role of temperature in the growth and flowering of geophytes. *Plants* **2**, 699-711.

Khosa, J., Bellinazzo, F., Kamenetsky Goldstein, R., Macknight, R. and Immink, R.G.H. (2021) PHOSPHATIDYLETHANOLAMINE-BINDING PROTEINS: the conductors of dual reproduction in plants with vegetative storage organs. *Journal of Experimental Botany* **72**, 2845-2856.

Kinoshita, A. and Richter, R. (2020) Genetic and molecular basis of floral induction in Arabidopsis thaliana. *Journal of Experimental Botany* **71**, 2490-2504.

Lee, C., Kim, S.J., Jin, S., Susila, H., Youn, G., Nasim, Z., Alavilli, H., Chung, K.S., Yoo, S.J. and Ahn, J.H. (2019) Genetic interactions reveal the antagonistic roles of FT/TSF and TFL1 in the determination of inflorescence meristem identity in Arabidopsis. *The Plant Journal* **99**, 452-464.

Lee, J.H., Ryu, H.-S., Chung, K.S., Posé, D., Kim, S., Schmid, M. and Ahn, J.H. (2013a) Regulation of Temperature-Responsive Flowering by MADS-Box Transcription Factor Repressors. *Science* **342**, 628-632.

Lee, J.H., Yoo, S.J., Park, S.H., Hwang, I., Lee, J.S. and Ahn, J.H. (2007) Role of SVP in the control of flowering time by ambient temperature in Arabidopsis. *Genes & development* **21**, 397-402.

Lee, R., Baldwin, S., Kenel, F., McCallum, J. and Macknight, R. (2013b) FLOWERING LOCUS T genes control onion bulb formation and flowering. *Nat Commun* **4**, 2884.

Leeggangers, H.A., Nijveen, H., Bigas, J.N., Hilhorst, H.W. and Immink, R.G. (2017a) Molecular regulation of temperature-dependent floral induction in Tulipa gesneriana. *Plant physiology* **173**, 1904-1919.

Leeggangers, H.A., Nijveen, H., Bigas, J.N., Hilhorst, H.W. and Immink, R.G. (2017b) Molecular Regulation of Temperature-Dependent Floral Induction in Tulipa gesneriana. *Plant Physiol* **173**, 1904-1919.

Li, X.-F., Wu, W.-T., Zhang, X.-P., Qiu, Y., Zhang, W., Li, R., Xu, J., Sun, Y., Wang, Y. and Xu, L. (2015) Narcissus tazetta SVP-like gene NSVP1 affects flower development in Arabidopsis. *Journal of Plant Physiology* **173**, 89-96.

Lu, J., Qu, L., Xing, G., Liu, Z., Lu, X. and Han, X. (2023) Genome-Wide Identification and Expression Analysis of the MADS Gene Family in Tulips (Tulipa gesneriana). *Genes (Basel)* **14**.

Molina, R., Valero, M., Navarro, Y., Guardiola, J. and Garcia-Luis, A. (2005a) Temperature effects on flower formation in saffron (Crocus sativus L.). *Scientia horticulturae* **103**, 361-379.

Molina, R.V., M., V., Y., N., A., G.-L. and and Guardiola, J.L. (2005b) Low temperature storage of corms extends the flowering season of saffron (Crocus sativus L.). *The Journal of Horticultural Science and Biotechnology* **80**, 319-326.

Nakano, Y., Higuchi, Y., Sumitomo, K. and Hisamatsu, T. (2013) Flowering retardation by high temperature in chrysanthemums: involvement of FLOWERING LOCUS T-like 3 gene repression. *Journal of Experimental Botany* **64**, 909-920.

Navarro, C., Abelenda, J.A., Cruz-Oró, E., Cuéllar, C.A., Tamaki, S., Silva, J., Shimamoto, K. and Prat, S. (2011) Control of flowering and storage organ formation in potato by FLOWERING LOCUS T. *Nature* **478**, 119-122.

Noy-Porat, T., Cohen, D., Mathew, D., Eshel, A., Kamenetsky, R. and Flaishman, M.A. (2013) Turned on by heat: differential expression of FT and LFY-like genes in Narcissus tazetta during floral transition. *Journal of experimental botany* **64**, 3273-3284.

Oda, A., Narumi, T., Li, T., Kando, T., Higuchi, Y., Sumitomo, K., Fukai, S. and Hisamatsu, T. (2012) CsFTL3, a chrysanthemum FLOWERING LOCUS T-like gene, is a key regulator of photoperiodic flowering in chrysanthemums. *Journal of Experimental Botany* **63**, 1461-1477.

Pin, P.A., Benlloch, R., Bonnet, D., Wremerth-Weich, E., Kraft, T., Gielen, J.J. and Nilsson, O. (2010) An antagonistic pair of FT homologs mediates the control of flowering time in sugar beet. *Science* **330**, 1397-1400.

Putterill, J., Laurie, R. and Macknight, R. (2004) It's time to flower: the genetic control of flowering time. *BioEssays* **26**, 363-373.

Pyo, Y., Park, S., Xi, Y. and Sung, S. (2014) Regulation of flowering by vernalisation in Arabidopsis. In: *Advances in botanical research* pp. 29-61. Elsevier.

Renau-Morata, B., Nebauer, S.G., García-Carpintero, V., Cañizares, J., Gómez Minguet, E., de los Mozos, M. and Molina, R.V. (2021) Flower induction and development in saffron: Timing and hormone signalling pathways. *Industrial Crops and Products* **164**, 113370.

Simon, R., Igeño, M.I. and Coupland, G. (1996) Activation of floral meristem identity genes in Arabidopsis. *Nature* **384**, 59-62.

Singh, D., Sharma, S., Jose-Santhi, J., Kalia, D. and Singh, R.K. (2023) Hormones regulate the flowering process in saffron differently depending on the developmental stage. *Front Plant Sci* **14**, 1107172.

Srikanth, A. and Schmid, M. (2011) Regulation of flowering time: all roads lead to Rome. *Cellular and molecular life sciences* **68**, 2013-2037.

Strasser, B., Alvarez, M.J., Califano, A. and Cerdán, P.D. (2009) A complementary role for ELF3 and TFL1 in the regulation of flowering time by ambient temperature. *The Plant Journal* **58**, 629-640.

Sureshkumar, S., Dent, C., Seleznev, A., Tasset, C. and Balasubramanian, S. (2016) Nonsense-mediated mRNA decay modulates FLM-dependent thermosensory flowering response in Arabidopsis. *Nat Plants* **2**, 16055.

Taoka, K.-i., Ohki, I., Tsuji, H., Kojima, C. and Shimamoto, K. (2013) Structure and function of florigen and the receptor complex. *Trends in plant science* **18**, 287-294.

Tsaftaris, A., Pasentsis, K. and Argiriou, A. (2013) Cloning and Characterization of FLOWERING LOCUS T-Like Genes from the Perennial Geophyte Saffron Crocus (Crocus sativus). *Plant Mol. Biol. Report.* **31**, 1558–1568.

Tsuji, H., Nakamura, H., Taoka, K. and Shimamoto, K. (2013) Functional diversification of FD transcription factors in rice, components of florigen activation complexes. *Plant Cell Physiol* **54**, 385-397.

Turck, F., Fornara, F. and Coupland, G. (2008) Regulation and identity of florigen: FLOWERING LOCUS T moves center stage. *Annu. Rev. Plant Biol.* **59**, 573-594.

Tylewicz, S., Tsuji, H., Miskolczi, P., Petterle, A., Azeez, A., Jonsson, K., Shimamoto, K. and Bhalerao, R.P. (2015) Dual role of tree florigen activation complex component FD in photoperiodic growth control and adaptive response pathways. *Proceedings of the National Academy of Sciences* **112**, 3140-3145.

Wang, Z., Li, X., Xu, J., Yang, Z. and Zhang, Y. (2021) Effects of ambient temperature on flower initiation and flowering in saffron (Crocus sativus L.). *Scientia Horticulturae* **279**, 109859.

Wickland, D.P. and Hanzawa, Y. (2015) The FLOWERING LOCUS T/TERMINAL FLOWER 1 gene family: functional evolution and molecular mechanisms. *Molecular plant* **8**, 983-997.

Wigge, Philip A. (2011) FT, A Mobile Developmental Signal in Plants. *Current Biology* **21**, R374-R378.

Wigge, P.A., Kim, M.C., Jaeger, K.E., Busch, W., Schmid, M., Lohmann, J.U. and Weigel, D. (2005) Integration of spatial and temporal information during floral induction in Arabidopsis. *Science* **309**, 1056-1059.

Wu, X., Ling, W., Pan, Y., Yang, Z., Ma, J., Yang, Y., Xiang, W., Zhou, L., Sun, M., Chen, J., Chen, H., Zheng,
S., Zeng, J. and Li, Y. (2024) Functional analysis of a lily SHORT VEGETATIVE PHASE ortholog in
flowering transition and floral development. *Plant Physiology and Biochemistry* **206**, 108287.
Wu, Y.-M., Ma, Y.-J., Wang, M., Zhou, H., Gan, Z.-M., Zeng, R.-F., Ye, L.-X., Zhou, J.-J., Zhang, J.-Z. and
Hu, C.-G. (2022) Mobility of FLOWERING LOCUS T protein as a systemic signal in trifoliate
orange and its low accumulation in grafted juvenile scions. *Horticulture Research* **9**.
Xue, W., Shi, J., Li, Z., Zhang, Y., Tang, M., Du, X., An, Z., Chen, H., Yang, D. and Li, X. (2025) CmFDa-
mediated epigenetic regulation of flowering in chrysanthemum. *Plant Biotechnology Journal*.
Yan, X., Cao, Q.-Z., He, H.-B., Wang, L.-J. and Jia, G.-X. (2021) Functional analysis and expression
patterns of members of the FLOWERING LOCUS T (FT) gene family in Lilium. *Plant Physiology
and Biochemistry* **163**, 250-260.
Zhu, Y., Klasfeld, S., Jeong, C.W., Jin, R., Goto, K., Yamaguchi, N. and Wagner, D. (2020) TERMINAL
FLOWER 1-FD complex target genes and competition with FLOWERING LOCUS T. *Nature
communications* **11**, 5118.
Zik, M. and Irish, V.F. (2003) Global identification of target genes regulated by APETALA3 and
PISTILLATA floral homeotic gene action. *The Plant Cell* **15**, 207-222.

1220