Title: Sensory and palatability coding of taste stimuli in cortex involves dynamic and asymmetric

cortico-amygdalar interactions

Alternative title: Both sensory and palatability coding of taste stimuli in cortex involve a back-and-

forth conversation between amygdala and cortex

Abbreviated title: The amygdala-cortical loop processes taste

Authors: Mahmood, A.[1,2], Steindler, J.R.[3], Katz, D.B.[1,2]

Department of Psychology[1], Volen Center for Complex Systems[2], Brandeis University, Waltham, MA

Lewis Katz School of Medicine[3], Temple University, Philadelphia, PA

Corresponding Author: Donald B. Katz, dbkatz@brandeis.edu

Conflicts of Interest: None declared

Keywords: taste, reciprocal connectivity, cortico-limbic interaction, asymmetric influence,

response dynamics, palatability

Contents

Abstract

Gustatory cortical (GC) and basolateral amygdalar (BLA) taste responses consist of an inter-regionally coherent 3-part state sequence. This coherence suggests that reciprocal BLA-GC connectivity is important for taste processing, but it remains unknown: 1) whether BLA-GC coherence actually reflects a reciprocal "conversation" (as opposed to one region simply driving the other); and 2) whether such a "conversation" has anything to do with the taste processing observed within GC response dynamics. Here, we address these questions using network and single-neuron analysis of simultaneously-recorded GC and BLA taste responses in awake rats. We find asymmetric, reciprocal μ-frequency influences that reflect taste processing dynamics: BLA→GC influence dominates between 300 and 1000msec (the epoch in which BLA codes palatability); afterward, when GC responses become palatability-related and GC has been shown to release a behavior-relevant signal, the direction of influence reverses, becoming GC→BLA. Follow-up analyses demonstrate that this "turn-taking" exists alongside effectively synchronous amygdala-cortical coupling—the two regions functioning as a unified structure. Finally, to assess the implications of these interactions for single-neuron responses, we tested the response properties of GC neurons categorized by their inferred connectivity with BLA: GC neurons influenced by BLA produce stronger taste-specific and palatability-related responses than other GC neurons, and the strongest taste encoding is specifically found in GC neurons that both influence and receive influence from BLA—those most deeply embedded in the reciprocal circuit. These results, consistent with findings in multiple systems, support the novel conclusion that taste processing and decision-making is a function of the amygdala-cortical loop.

New & Noteworthy

Conventionally, taste circuitry is considered feedforward, travelling up from the brainstem, with each additional node containing more sophisticated information. We challenge this convention by demonstrating that amygdala and cortex instead influence each other bidirectionally, yet in a direction-specific and asymmetric manner. These influences appear to drive different parts of the taste response, with distinct patterns for decision-making and behavioral output; furthermore, involvement in cortex-amygdala functional connectivity determines the strength of encoding in cortical single neurons.

Introduction

Stimulus-evoked activity in gustatory cortical (GC) neurons transitions through a sequence of population states, with each successive state containing more highly processed information (1,2). Across the 1-1.5 sec following taste delivery to the tongue, these states reflect, in turn, the fact that the taste is on the tongue, then taste identification, and finally the decision to either consume or reject the taste on the basis of its perceived palatability (3,4). Such ensemble neural dynamics are predicted to arise in systems with dense recurrent connectivity (5,6), which has been shown, in studies on visual processing, to generate dynamic neural responses observed across multiple regions, as inter-region influence cascades in forward and reverse directions across the visual stream (7–9).

Another expected consequence of reciprocal inter-region connectivity is coupling of activity across regions. In line with this expectation, recent work has shown both correlations in the latencies of the above-described firing-rate transitions and significant cross-coherence between GC and Basolateral Amygdalar (BLA) taste responses (10), both which have strong reciprocal connections (11–15). these findings dovetail nicely with work demonstrating cross-coherence between reciprocally connected structures in other sensory systems (16,17), together suggesting that the reciprocal connectivity between BLA and GC may be central to processing of the taste itself.

Current literature stops well short of testing that hypothesis, however. Previous studies on inter-regional interactions in taste processing have largely focused on the impact of perturbing one region on processing in the other (4,18–22), and while a few studies have investigated multi-region recordings, they have done so using cross-coherence measures (10), usually collapsed across time (19,23). Since these techniques are agnostic to which connections cause "coupling," they cannot distinguish unidirectional from bidirectional influences. That is, it is unknown whether the evolution of the neural processing of tastes reflects moment-to-moment changes in which region's activity is "influencing" that in the other.

The current study answers these questions by bringing advanced analyses to bear on simultaneously-recorded BLA and GC taste-evoked activity (in a paradigm that avoids confounds related to GC's involvement in interoception, see Methods). We show that the BLA-GC interaction is a conversation in which the two regions take turns influencing one another, and that this asymmetry of influence is epoch-specific: BLA primarily influences GC during the 300-1000ms post-stimulus period, which is the epoch during which BLA responses are palatability-related (10,23,24), and just before GC responses become so (2,3); GC influence on BLA then peaks briefly at approximately the time of the transition into the next epoch—the point at which GC has been shown to emit behavior-related signals (25). We perform follow-up analyses to show that this "turn taking" asymmetry exists within the framework of nearly synchronous, coordinated dynamics—as expected for a system in which function relies on reciprocal interactions.

We then move on to characterizing the functional properties of single neurons underlying these
interactions, testing the hypothesis that embeddedness in the amygdala-cortical loop determines
involvement in taste processing. We show that roughly half of our GC single-neuron sample can be
inferred to either receive input from or send output to BLA. These overlapping functional "input" and
"output" populations are both primarily made up of putative pyramidal neurons, and yet are distinct
with regard to their roles in taste processing: GC neurons receiving influence from BLA show
significantly higher levels of both taste specificity and palatability-relatedness in their responses
than those that only influence BLA; furthermore, GC neurons that both influence and receive
influence from BLA produce the strongest taste-related responses.

Collectively, these results shed light onto the forces underlying taste-related processing by
demonstrating that dynamic recurrent interactions between BLA and GC play a role in the emergence
of GC neural codes related to the generation of the behavioral response, and that the degree to which
a GC neuron is embedded in the reciprocal circuit largely determines its involvement in taste
processing. These results highlight the importance of inter-region interactions in understanding
perceptual function.

Results

BLA-GC interaction dynamics are asymmetric but reciprocal

Insight into the involvement of reciprocal amygdala-cortical coupling in taste processing can only be gained through an examination of the dynamics of directional influence—of the moment-to-moment degree to which activity in each area impacts activity in the other—and of the degree to which this interplay reflects the known taste processing dynamics. We began this examination by applying Spectral Granger Causality (GrCa) to local field potentials (LFPs) simultaneously recorded from BLA and GC. Rather than simply assessing similarity between timeseries (as cross-coherence analyses do), GrCa allows us to disentangle directional dependencies by making use of time-lagged regressions: if future GC activity is better predicted using the histories of both GC and BLA, compared to just using the history of GC, then BLA can be thought to "causally influence" the future of GC (26–30). Significance of influence in GrCa can then be assessed for each time and frequency bin, and thus we can test whether GC and BLA are mutually influencing one another (and whether those directional influences have different time-courses). We note that GrCa infers functional connectivity between regions, and neither has information regarding, nor makes claims about, the physical connectivity between the sources of its data; hence, we use the term "influence" here to describe only a functional connection without claiming any specific properties of a neural projection (excitatory or inhibitory).

Fig. 1 shows the basic (collapsed across time) output of this analysis brought to bear on simultaneously recorded GC and BLA LFPs. The influence of each region on the other is almost entirely contained within the 1-60Hz range (**Fig. 1A**; note that the low-pass filter used when collecting the LFPs allows for the detection of influence at much higher frequencies, had they existed, see Methods). More to the point of our investigation, the GC→BLA influence proved quite distinct from the BLA→GC influence within that range (**Fig. 1B**): GC→BLA influence, while above zero across this entire range of frequencies, dominates in the 20 to 60Hz range (β and γ frequencies; see Ref# 31); BLA→GC influence, meanwhile, is essentially non-existent (and non-significant) across most of the β/γ range, but is the dominant direction of influence in the θ and μ frequency ranges (2-12Hz), the latter of which is sometimes referred to as α (31,32). This latter frequency range is known to be both particularly prominent in GC evoked activity and modulated in a manner related to the dynamics of GC taste responses (31–33). In summary, BLA influences GC and GC influences BLA during taste processing, and these two influences are distinct from one another.

We note that it is unlikely that this relatively clean separation in directional frequencies is an artifact of the methodology, as GrCa can appropriately recognize influences in identical or overlapping frequency bands, as has been demonstrated before (26,34–36). Furthermore, the distinctness between the frequency bands of the BLA→GC and GC→BLA influences aligns with previous work showing that pharmacological inhibition of taste thalamus (VPMpc) reduces GC LFP power in the 20-

162   60Hz band, but increases it in the 2-12Hz band (19)—a finding that corroborates the idea of the 2-

163   12Hz and 20-60Hz bands as being distinct communication "channels".

Fig. 1) GC-BLA interaction is rich with processing-related asymmetries and dynamics. A) Mean (bold lines) and individual sessions' (thin lines) significance of Spectral Granger Causality for 0-2000ms post-stimulus activity across datasets for BLA→GC (black) and GC→BLA (red) influences. **B)** Directional differences in mean (bold red line) and individual subjects' (thin black lines) influence across frequency bands for data shown in A). Above-zero differences mean stronger BLA→GC influence, and below-zero differences mean stronger GC→BLA influence. BLA→GC is stronger below 12Hz, and GC→BLA dominates between 20 and 60Hz; ** = p<0.05, *** = p<0.005. **C)** The data from Fig. 1A unpacked into significant time-frequency bins for **i)** BLA→GC and **ii)** GC→BLA influence. Brighter color indicates that a larger fraction of the recording sessions displayed significant Granger Causality at that time-frequency bin (heatmap legend to right). In both panels the difference between higher and lower frequency bands can be seen, and across panels the asymmetry in influence can be seen; furthermore, dynamics of influence are apparent in the lower frequency band. **D)** The data shown in C collapsed across frequencies within each identified bandwidth (frequency band limits are given above each panel). Dynamics of influence are restricted to the <12Hz bandwidth; shaded regions indicate periods of strongest significance determined using cluster-permutation testing. See text and Methods for more details.

The distinction between lower and higher frequency ranges is also easily seen when direction-specific influence is examined across time (**Fig. 1C**; compare the upper two thirds to the lower third in each panel), as is the difference between GC→BLA and BLA→GC influence (compare the left and right panels). This presentation further suggests that, in general, the β/γ influence may be slightly higher spontaneously than during taste processing. But most centrally with regard to our hypothesis, the differences between BLA→GC (**Fig. 1Ci**) and GC→BLA (**Fig. 1Cii**) influences within the θ/μ frequency range can be seen to change repeatedly across taste processing time, and this "turn-taking" reflects the oft-described GC dynamics.

To rigorously assess the significance of this latter phenomenon, we plotted the average proportion of sessions in which power in each frequency-time bin is significantly above zero (the "significant fraction") in each direction of influence, collapsed across frequencies within each band. This analysis reveals near-tonic influences in the higher frequency band (**Fig. 1Di** and **ii**; the average GC→BLA magnitude of influence is of course higher than that of BLA→GC influence), although there is a slight trend toward a reduction of influence following taste delivery.

The μ/α band results, meanwhile, do in fact reflect the known GC taste-response dynamics (**Fig. 1Diii** and **iv**). In each panel, the shaded regions represent the result of cluster-based permutation testing, identifying the longest continuous periods in which this result is significantly higher than that of randomized data (note that there are no such periods for β/γ influence; **Fig. 1Di/ii**). With regard to BLA→GC influence in the lower frequency band (**Fig. 1Diii**), this period is a good match to the GC identity epoch (approx. 300-1000ms post-stimulus; see **Fig. 3A**). Given that BLA responses are palatability-related during this period (24), and that GC responses become palatability-related only at the end of this period, this result also accords well with classic studies on the role of amygdala in stimulus valuation, which have led both human and rodent researchers to theorize that amygdala "passes" emotional valence to cortical regions (4,18,37,38) see also Discussion).

GC→BLA influence in this lower frequency range, meanwhile (**Fig. 1Div**), peaks only briefly (note the thin string of yellow bins between 1.0 and 1.5sec in the bottom rows of **Fig. 1Cii**), with that peak coming immediately following the BLA→GC push. This influence "turn taking" further corroborates the above interpretation, suggesting that GC, after being influenced by BLA, emits decision-related activity capable of influencing behavior, consistent with work demonstrating that the transition into palatability-related firing in GC (which typically occurs around this time) is causally linked to the driving of behavior (3,25). This result establishes bi-directional interactions between BLA and GC during taste processing, with dynamic asymmetries of amygdala-cortical coupling that directly reflect what is known of GC taste processing.

Asymmetric BLA-GC interaction dynamics are nonetheless synchronized

Again, the fact that this amygdala-cortical "conversation" is asymmetric was predicted (see Discussion section "The division of function(s) between BLA-GC directional influences"). And while it might seem at first blush that they contradict the previously-described coherence of activity in the two structures (10), further analysis revealed that even these asymmetries of influence reflect the fact that the two regions process tastes according to the same "clock"—that changes in influence

occur simultaneously. This can easily be seen when Principal Component Regression is performed
on each time-series shown in **Fig. 1C**. In **Fig. 2A**, the trajectories for the data in the two Fig. 1C panels
obtained from this regression are shown overlain on one another. The dynamics of the two time-
series, which are represented as changes in the trajectories' directions through PCA space, are
clearly very similar.

We quantified this similarity by assessing the mean-squared error (MSE) of attempts to linearly
predict one time-series from the other (**Fig. 2B**, red dashed line) compared to that produced by the
same linear regression of the two time-series after shuffling them temporally (linear, inter, shuffled),
and also by linear and non-linear (i.e., using multi-layered artificial neural network) auto-regression
(i.e., predictions of each time series from itself; see Methods), which give us a lower bound for
prediction error. The results of these analyses make it clear that the linear prediction of the BLA→GC
time-series from the GC→BLA time-series (in blue) was almost as accurate as the non-linear auto-
regression (orange) and better than the linear auto-regression (green; this was not surprising given
the nonlinearities in the BLA-GC interaction dynamics). Simply put, the inter-series prediction nears
the limit of prediction accuracy, a fact that serves as novel evidence for the suggestion that BLA and
GC form, upon receiving taste input, a single distributed dyad with time-locked, coupled dynamics—
albeit dynamics within which GC→BLA and BLA→GC influences are asymmetric.

Fig. 2) Dynamics of BLA→GC and GC→BLA influences are time-locked. A) Aligned principal components of spectra shown in **Fig. 1C**, indicating that the dynamics of BLA→GC and GC→BLA influence are strongly coordinated, in that the trajectories change direction together through time. Elapsing post-stimulus time is shown with changing colors (legend to left). **B)** Quantification of alignment shown in A). Prediction of one time-series (GC→BLA) from the other (BLA→GC) using linear regression (Linear, Inter) is better than both shuffled linear regression (Linear, Inter, Shuffled), and linear autoregression (Linear, Auto) of timeseries, and almost as good as non-linear autoregression (Non-linear, Auto).

As a convergent measure, we tested the specific timing of the coordination between the ensemble
firing-rate state changes (**Fig. 3A**) that have been repeatedly identified in GC and BLA (10,24). While
our previous results show the latencies of GC and BLA state changes to be correlated, this leaves
open the question of whether there are significant lags (or offsets) between ensemble firing-rate
transitions in either region (see **Fig. 3B**), which would suggest a general direction of "influence flow."
We therefore directly assessed these lags (see **Fig. 3C** for a schematic of the analysis).

The between-region lags for each individual session (**Fig. 3Di**, presented as a heat map) appear to be
highly reliable across the 18 sessions (from 6 rats), appearances that are confirmed by the tightness
of the across-session distributions of mean lags (**Fig 3Dii**). Analysis of these latter distributions
reveals that, for the earliest transition, the sudden transition in BLA ensemble firing rates tends to
lead that in GC (permutation test, 95%CI=[57, 183], p<0.001), but fail to show significant between-
region deviation from simultaneity (zero-lag) of later transitions (Transition 2: 95%CI=[-74.5, 51.5],
p=0.78; Transition 3: 95%CI=[-86, 39.5], p=0.43). The distributions of lags for transitions 2 and 3 were
both significantly different from that for transition 1 (1 vs. 2: p=0.001, 1 vs. 3: p=0.015, Wilcoxon
signed-rank test), but not from one another (Wilcoxon signed-rank test, p=0.747). This finding agrees
with previous results which showed that the first transition is not significantly correlated between
BLA and GC, but the 2nd and 3rd transitions are (10).

These results support the hypothesis that BLA and GC are not a joint processing network "at rest,"
but that they coalesce into a dyad—not just working together, but becoming essentially a single
processing object—after producing uncoordinated initial transients in their taste responses (see
Discussion). The lag present in the first transition suggests that BLA and GC taste responses are
initially driven independently (and provides proof that the analysis is capable of detecting even small
reliable lags), although there are other possible explanations. Regardless, this initially uncoupled
activation gives way to the two regions cycling through the later transitions of their taste responses
as a single processing unit, such that the relatively long axons connecting the two regions do not
induce detectable delays in the firing-rate transitions that characterize taste processing (see
Discussion).

Fig 3. BLA initially leads GC in state transition but both regions subsequently show zero-lag coordination. A) Summary schematic of the timing and "coding content" previously demonstrated to characterize state sequences in GC and BLA ensemble spiking activity. Overlain are the numbers used in this study to label these transitions. **B)** State transitions can be strongly coordinated but may still show significant lags. **C)** A schematic illustrating how transition lags between GC and BLA were calculated and how we define the sign of the lags. **Di)** Distributions of transition lags, presented as heat maps (legend at right shows the # of trials with a specific transition lag) for 18 sessions (y-axis) across 6 animals for each of the 3 transitions. **Dii)** Summary distributions of lags for each transition, with red arrows indicating the grand-average lag. The significance of the difference between that grand-average lag and 0 was evaluated using bootstrapped 95% confidence intervals of the distribution mean (see text and Methods).

Modeling of single-neuron functional connectivity reveals overlapping groups of GC neurons differentiated on the basis of whether they influence or are influenced by BLA neuron activity

As noted above, the finding that different patterns of influence appear to "live" in different frequency bands is not without precedent—bottom-up and top-down influences in visual pathways have also been suggested to segregate into separate frequency bands (35,36,39). Meanwhile, the fact that BLA→GC influence is focused in the 2-12Hz range provides further motivation for our central hypothesis that the involvement of GC neurons in taste processing (which can be quantified in terms of the magnitude of GC responses' taste-specificity and palatability-relatedness) might be a function of their degree of coupling with BLA, given that previous studies have shown that GC taste-processing dynamics are also prominently visible in GC μ activity (19,31–33).

To set up the test of our central hypothesis, we trained Poisson Generalized Linear Models (pGLM, see Methods for important differences between pGLM and the more commonly-used General Linear Model) to infer connectivity and influence between individual GC-BLA neuron pairs—an aim that cannot be achieved with GrCa (see Methods), and that allows us to categorize individual GC neurons as putatively influencing and/or being influenced by activity in BLA (and vice-versa). We fit pGLMs to each single neuron in our dataset, using spike trains from all other simultaneously recorded neurons as input (40–42).

An advantage of this analytic approach to inferring functional connectivity is that, while it cannot capture the dynamics of influence, it is inherently more conservative than correlational methods such as spike-train cross-coherence (43). Standard correlational approaches are significantly more prone to false positives due to the shared latent dynamics across neurons within regions and across regions (41). In pGLMs stimulus input and the history of spiking in the current neuron "compete" against one another, and also against the recent spiking of other neurons (**Fig. 4A**), for significance during pGLM inference. The evoked activity of a neuron can potentially be explained as a function of any of these variables, and only explanatory value above and beyond what is predicted by all other variables is deemed evidence of a functional inter-regional influence—influence that can be attributed to other sources is essentially "partialled out."

In total, we fit models to 151 and 160 well-isolated GC and BLA single neurons, respectively (mean±SD per session: 9.4±4.14, GC; 10.0±4.61, BLA). Before using model output to test any hypotheses about taste processing, we tested the quality of the fit to the data provided by the models. **Fig. 4B,** 2 representative examples of actual PSTHs (left) and PSTHs predicted by the pGLMs (right), visually illustrates this fit. The apparent similarity between the two is quantified in **Fig. 4C**, which shows that the model consistently produces significantly better matches to the data than models fit on spike-trains shuffled between trials (Wilcoxon signed-rank test, p < 0.001), a control that eliminates only the within-trial coordination of spiking between simultaneously recorded neurons, preserving the average statistics of the spike-trains used. This result highlights the fact that,

despite significant trial-to-trial variability in spiking, the pGLM learns within-trial relationships that
are meaningful.

As an additional test of the pGLM fits, we also compared correlations between actual PSTHs and: 1)
trial-matched predictions; 2) cross-unit predictions; and 3) predictions on circularly shuffled data
(**Fig. 4D**, see Methods for details). We found, as expected, that trial-matched predictions (blue
histogram and summary box plot) perform best. Control correlations with unit-shuffled/cross-unit
predictions (in orange) were significantly worse (Wilcoxon signed-rank test, $p < 0.001$) but also
significantly above zero, probably because many GC neuron pairs exhibit broadly similar neural
dynamics. Worst, and essentially zero, were correlations with circularly shuffled (temporally garbled)
data (in green). Together, these tests allow us to conclude that the pGLM generates reasonable
models of the mechanics (functional circuitry) underlying the observed activity.

Fig. 4) Poisson Generalized Linear Modelling of single-neuron interactions between BLA and GC provides good fits to the data. A) pGLM structure. Every neuron model depends on: 1) stimulus timing, 2) the neuron's spiking history, and 3) input from (or spiking history of) all other simultaneously recorded neurons. **B)** Actual and Predicted PSTHs of two representative units across 4 delivered tastants (different colors), showing that fitted models (right column) accurately recapitulate single-neuron activity (left column). **C)** The vast majority of models built on real data showed higher cross-validated likelihood than models trained on shuffled data (i.e., the bulk of the distribution is to the right of the vertical red dashed line at equal predictability). This confirms that the models are learning single-trial relevant features. The white arrow indicates the median of the distribution. **D)** The distributions of correlations (Spearman's Rho) between: 1) actual and predicted PSTHs (blue); 2) unit-shuffled data and prediction (orange); and 3) temporally scrambled data and prediction (Circ shuffle: green).

These results confirmed that we can confidently use the model to evaluate whether the activity of
one simultaneously recorded neuron influenced that of another (i.e., we could identify significant
coupling filters). In what follows, we for simplicity's sake refer to neurons with responses that are
concluded to exert influence on activity in the other region as "Output" neurons, and neurons with
responses that are influenced by activity in the other region as "Input" neurons; note that in neither
case are we proposing to have identified direct anatomical connections—the terms refer purely to
functional connectivity.

Fig. 5A shows the distribution of neurons so identified: clearly, our BLA sample contains more Output
neurons (40.0%) than Input neurons (6.3%), with an additional subset (9.4%) categorized as both;
the Output and Input populations in GC were more balanced (17.2% and 22.5%, respectively), with
an additional 11.9% characterized as doing both. Another subset of neurons in GC could not be
described as either influencing or being influenced by activity in BLA (and vice-versa). No claim is
made about whether these neurons might output to or take input from any structures from which we
didn't record, and therefore no further analysis is done on the (probably highly diverse set of) neurons
that were categorized as neither Input nor Output.

Given the relatively small overlap between sets of functional Input and Output neurons shown in **Fig.**
5A, it was reasonable to ask whether the two sets were distinct—that is, whether a neuron (or neuron
type) was specifically likely to be either an Input or Output neuron, but not both; as has been
suggested for projection neurons in the visual stream (17,44). Two separate analyses provided
evidence against this possibility. First, we performed a direct, Bayesian assessment of the overlap
between Output and Input sub-groups in both regions, comparing the data to a null distribution in
which labels for connection types were assigned randomly so that we could ask whether, given the
likelihood of a neuron being categorized as Input or Output, the conditional probability (i.e., the
likelihood of it being categorized as Input given that it was categorized as Output, and vice-versa) was
lower or higher than chance. **Fig. 5B** shows that the overlap we observe in the real data falls squarely
within the chance probability of overlap for this null-distribution for both regions, therefore providing
no evidence that Output and Input populations should be thought of as segregated groups.

We also assessed the likelihood that the separation of neurons into Output and Input groups aligns
with anatomically-defined neuron categories. Specifically, we asked whether putative pyramidal
cells, identified on the basis of spike width, were more likely to be Output neurons than Input
neurons. **Fig. 5C** demonstrates that our sample of isolated single neurons broke, as expected, neatly
into two groups on this basis, with a 0.45ms trough-to-peak time providing a clean cutoff between
fast-spiking putative interneurons (Int) and slower-spiking putative pyramidal neurons (Pyr; see
(4,45–47). Slightly under 10% of our sample fell into the Int category, a number that is not wildly
inconsistent with expectation based on an extensive previous literature using chronic extracellular
electrophysiology. Given the small fraction of interneurons, it can be concluded that putative
pyramidal neurons are approximately equally likely of being Input and Output neurons. In conclusion,
we could find no evidence suggesting that Output and Input neurons are distinct anatomical types
(at least at the level of gross subdivision).

Fig. 5) pGLM brought to bear on amygdala-cortical ensembles identifies functionally (but not structurally) distinct subpopulations of neurons. A) Inference of Input and Output population in the recorded BLA-GC populations (number outside colored circles indicates percentage of neurons with no identified between-region connections). **B)** The amount of overlap between Input and Output populations is not significantly different from that expected by chance in either brain region. **C)** Inferred neuron-types based on trough-to-peak times of waveforms (Inset: solid line = GC, crosses = BLA). Most neurons in the dataset (expectedly) are putative pyramidal neurons, which means that the vast majority (and similar percentage) of each functional neural subtype as shown in panel A is pyramidal.

Taste processing in GC is a function of embeddedness in the amygdala-cortical loop

While our analyses do not suggest that Output and Input neurons are separate neuron types, our ultimate hypothesis was that the "embeddedness" of GC neurons within the reciprocally-connected network—whether they influence BLA (Output neurons), are influenced by BLA (Input neurons), both, or neither—determines their involvement in taste processing. Having conservatively labeled each neuron's functional role, we tested this possibility; specifically, we tested whether we can discern a relationship between pGLM label and the magnitude of the stimulus-specificity (i.e., more distinctive responsiveness to different tastes) and palatability-relatedness (i.e., correlation with the hedonic value of the tastants) of the neuron's taste response—the basic variables that describe involvement in taste processing (1–4,10,23–25,48,49).

Our analyses did in fact reveal significant differences in both taste-coding variables as a function of subpopulation (**Fig. 6A**): GC Input neurons (again, these are those shown by pGLM to be influenced by BLA, with no anatomical specifics implied) showed higher magnitudes of both taste specificity and palatability relatedness than did GC Output neurons. Despite the status of GC as a primary sensory cortex, the recipient of ascending taste input from thalamus (14,19,50,51), involvement in taste processing (as measured by taste specificity and palatability relatedness of taste responses) appears to depend on receiving "feedback" input from BLA.

As taste discriminability and palatability are (necessarily) non-independent, correlated variables (**Fig. 6B**, Spearman's Rho=0.73, p<0.001), it was useful to aggregate and summarize the results shown in **Fig. 6A** into a single measure. We used for this summary the vector norm (the magnitude of an arrow drawn to a point in 2D-space, see Methods) of normalized discriminability and palatability coding, an aggregate measure of taste-processing involvement that we here refer to as "tastiness." An analysis of neuron tastiness (**Fig. 6C)** confirms the findings of **Fig. 6A**: GC Input neurons show significantly higher tastiness than GC Output neurons (note that this difference was not an artifact of lower firing rates in the Output neurons—we observed no significant link between firing rates and the metrics tested above; Spearman's Rho: Firing Rate vs. Discriminability = 0.1, Firing Rate vs. Palatability = 0.0).

But the most rigorous test of our central hypothesis (that involvement in taste processing depends upon being "embedded" in the amygdala-cortical loop) involves an analysis of the specific set of neurons that are most deeply so embedded—that is, the GC neurons that are categorized as both influencing and receiving influence from BLA (i.e., "Input/Output" neurons). If this hypothesis is correct, we should expect that these neurons will exhibit the strongest tastiness—even stronger than that of GC Input neurons. Indeed, this proved to be the case: the tastiness of GC Input/Output neurons is significantly higher than that of GC Input neurons (Student's t-test p=0.01).

Note that the division of neurons into functional Input and Output types was made by a model that was blind to taste coding in those responses, and therefore our results provide a conservative, risky test of the importance of BLA feedback in taste responsiveness. Thus they confirm our prediction that the degree of "embedding" in the reciprocal circuit determines its degree of involvement in taste

434   processing (**Fig. 6C**), a result that is particularly striking given the weak "tastiness" of GC Output

435   neuron responses.

Fig. 6) GC neurons that are embedded in the amygdala-cortical loop are the ones that most strongly process tastes. A) Both taste discriminability **(i)** and palatability-relatedness **(ii)** of GC Input neurons (those influenced by BLA activity), measured in the standard manner (see Methods), are significantly stronger than that in GC Output neurons. ** = p<0.005. **B)** Discriminability and Palatability are significantly correlated for GC neurons (Spearman's Rho=0.73, p<0.001) and hence can be combined into a measure ("Tastiness") that represents a single measure of involvement in taste processing. **C)** Comparison of tastiness among subpopulations confirms the results of A (higher tastiness in Input neurons than Output neurons), but also shows that the small population of GC Input/Output neurons produces responses with significantly higher tastiness than even the Input population (2-sample Student's T-test, ** = p<0.005, * = p<0.05)

Discussion

This study reveals GC taste processing to centrally involve dynamical activity of the amygdala-cortical loop. Following an initial transient in which they fire independently of one another, the flow of influence between BLA and GC becomes one of coupled, asymmetric "turn-taking." Influence in the BLA→GC direction appears dominant during the middle epoch, when BLA taste responses are palatability-related and GC taste responses are not; GC→BLA influence, meanwhile, spikes immediately afterward, at the time that GC emits a signal reflecting the reaching of a consumption decision (almost certainly based on the prior input from BLA). At the single-neuron level, any individual GC neuron's involvement in taste processing—measured in terms of the taste-specificity and palatability-relatedness of its taste responses—is shown to be a function of its embeddedness in this reciprocal circuit.

Thus, we conclude that the BLA-GC dyad, formed soon after taste delivery, processes tastes on the basis of direct recurrent connectivity between the two regions, collectively transitioning through quasi-stable states. The following sections discuss this proposal in further detail, specifically considering: 1) how stimulus-evoked metastable dynamics seen in the amygdala-cortical circuit are well-described as a distributed attractor network; 2) whether this conceptualization can be generalized to describe work from other sensory systems; and 3) the fact that the "flow of influence" in tightly connected dynamical systems is likely to be asymmetric, with distinct dynamics of influence acting in concert with one another, And 4) a brief consideration of output pathways and possible role of this loop in other proposed functions of GC.

Taste processing evolves through the functioning of an amygdala-cortical joint attractor network

The metastable population dynamics that make up GC taste responses have already been successfully modeled as an attractor network (52–55). Our current results reveal this attractor network to be made up not just of GC, but of the dyad of GC and BLA. In such a joint attractor network, there is no "source" of transitions, which are triggered by coordination of the noise in the system. As predicted by such a model, all but the initial transitions (which are "externally" driven by the stimulus) in taste-evoked activity happen, on average, with lags that are distributed around zero, despite the fact that the regions are connected by relatively long axons—a result that is difficult to explain using hierarchical/feedforward models which, by construction, contain sources and sinks of influence (56–58). Thus, we argue that taste input causes the BLA-GC system to switch from resembling a feedforward system to resembling a recurrent attractor network for purposes of taste processing.

This characterization of BLA and GC as a joint network is unlikely to be an isolated case. Our previous work has shown that at the single-neuron level, neurons in both the lateral hypothalamus (48) and the parabrancial nucleus of the pons (59) also show states in the taste-evoked responses. Indeed, as noted below, such multi-state responses are ubiquitous, and hence, suggest that BLA and GC are likely undergoing these state-transitions as part of the larger taste circuit, especially given the dense recurrent long-range connectivity between most regions in the circuit (14,15). The fact that both GC

and BLA are also emotional, interoceptive, and multisensory "hubs" (60,61) makes it reasonable to
speculate that they couple with the olfactory circuitry, as well as frontal cortices, in appropriate
functional contexts as well.

Given these considerations, we also expect that changes in internal state of the animal alters BLA-
GC coupling. Indeed, GC taste responses change with the rat's level of engagement in the task (31),
a change accompanied by dramatic changes in the GC's LFP profile suggesting a change in the
functional connectivity of the larger network. It is likely that such a change may be driven by, or in
concert with, the amygdalar (or frontal cortical) neurons, firing of which is sensitive to both internal
state and task structures (62,63).

Reciprocal interactions underlie processing in other sensory systems

Evidence supporting our proposal that reciprocal connectivity causes regions to "coalesce" into a
single perceptual processing unit abounds in work on other sensory systems. Global Workplace
Theory (GWT) (8,64), for instance, claims that sensory information is first transmitted in a
feedforward manner to specific brain regions; activation of these regions then engages a distributed
network via reciprocal projections, which essentially binds the regions together. Further, these
different states reflect different stages of perceptual processing and transitions between stages are
determined by different mechanisms. Importantly, their 2^{nd} state is driven by establishment of
recurrent interactions, is internally driven (as opposed to being stimulus-bound), and reflects
"conscious" stimulus processing. This mirrors taste research showing that processing in the 2^{nd} state
in GC taste-evoked activity is used to make consumption decisions (3,55,65–68). And finally, that
perturbation of activity in this 2^{nd} state most strongly affects stimulus-related decision making in
both the GWT-related visual blocking task (8) and taste responses (25). Similar reports of circuits
undergoing a state transition from an initial feedforward to a recurrent (core processing) state can be
found for more visual processing studies (7,9), as well as somatosensory (69), auditory (70), and
olfactory processing (71).

The division of function(s) between BLA-GC directional influences

While the fact that BLA and GC progress through taste processing in lock-step is itself evidence of
the importance of the reciprocal connectivity between them, more straightforward evidence is found
in the asymmetries in how each region influences the other. It is actually not unusual (quite the
opposite) for recurrently interacting systems to contain such asymmetric influences. Examples of
dynamical systems in which feedforward and feedback interactions have different strengths include
half-center oscillators (72), Lotka-Volterra (prey-predator) models (73), and reservoir networks (74).
Furthermore, researchers have reported asymmetric interactions analogous to our own in data
collected from visual system recordings (17,36,75).

The "turn-taking" dynamics of the amygdala-cortical interaction we have observed in this study
accord well with classic stimulus-valuation studies, including those performed by Damasio (37) and
Schoenbaum (38), which have been interpreted as suggesting that amygdala, after playing a primary
role in determining the value of a stimulus, provides this information to cortex which then uses the

information to generate a behavioral decision. Our GrCa analysis provides the first representation of
this "signal" being emitted by GC, but it is worth noting that Forseth et al. (76) observed a similar
transient "signal" emanating from somatomotor regions approximately 250ms prior to the onset of
response articulation in the above-discussed visual-naming task.

In the reverse direction, it is tempting to speculate that the tonic GC→BLA influence in the γ
frequency range powers the processing attributed to BLA, perhaps by sending taste quality
information to BLA while BLA returns palatability information, with the two regions co-evolving in this
manner to converge to a decision (77). While GC neurons that influence BLA activity without
themselves being influenced by BLA activity (i.e., GC Output neurons) showed weak taste coding,
the above conclusion can nonetheless be easily reconciled with previous work (20) showing that a
subset of GC→BLA projection neurons have more discriminatory taste-responses compared to the
average of non-projecting neurons, in response to aversive tastants vs palatable tastants. This
matches well with our data if we remember that the bidirectional GC Input/Output neurons will only
appear as GC→BLA projection neurons if bidirectional projecctions are not considered.

Output pathways and other GC functions

In the context of previous work (3,4,10,25,37,38,78), the current findings can be interpreted to
demonstrate that amygdala-cortical interactions culminate in the generation of consumption
decisions. As to what happens next—how exactly the consumption decision is behaviorally
enacted—our work is agnostic. Basic consumption-related behaviors are actuated by a central
pattern generator in the brainstem, the Reticular Formation (RF; (79–81)), and thus it is reasonable to
speculate that information flow from GC to RF is capable of driving these behaviors. A direct pathway
connects GC and RF (82,83), but there are in addition multiple indirect feedback pathways linking
the two structures (84,85). A recent proposal suggests that palatable and aversive taste behaviors
may be instigated by separate feedback pathways leading through different amygdalar structures
(86,87). While the connection between this work and ours is currently unclear (the proposal of
separate behavioral pathways relies on the suggestion that spatially separated populations of GC
neurons code the different tastes—a finding that most labs do not observe; (1,68,88–91), future work
will illuminate that connection.

Another subject of future work is the involvement of the amygdala-cortical loop in other GC
functions. GC responses have been shown in multiple contexts to represent post-ingestive factors;
relatedly, recent work has highlighted the role of GC in interoceptive processes (60,92,93). While for
the current study we expressly used an experimental protocol that minimized the influence of these
variables (delivering small aliquots of fluid such that rats neither satiated nor ingested substantial
numbers of calories, and randomizing presentation order and timing such that rats could not
anticipate any delivery) such that taste responses were stable across entire sessions (94), it is
entirely likely that the kind of amygdala-cortical dynamics exposed here are differentially invoked
when a rat is adjusting its consumption on the basis of post-ingestive need. Even more generally,
while the BLA-GC connection is likely stable during our roughly hour-long recording sessions, it will
inevitably be modulated by long-term brain-states given that neural networks display multi-scale,

context-dependent dynamics and interactions (95–97), making this an exciting avenue for future
research.

Conclusion

This study furthers our understanding of the role of inter-region interactions in sensory processing by
showing that cortico-amygdalar interactions subserve taste processing and decision-making. We
speculate that these dynamics are potentially a function of the larger taste circuit including the
brainstem (59). This taste network is initialized in a feedforward state but coalesces into a
reciprocally interacting network that evolves toward a point shown previously to generate the
behavioral output (3,25). Our investigation of the makeup of GC Input and Output neurons suggests
that network interactions are strong determinants of neural encoding and response properties . All
results presented herein align with multiple previous studies across sensory systems as well as
investigations of GC-BLA interactions during learning, allowing us to conceptualize these previous
results in a unified theoretical framework.

Methods

Experimental Design and Statistical Analyses

Subjects

Adult, female Long-Evans rats (n = 6; 300–350g at time of electrode implantation, Charles River Laboratories) served as subjects in our study (we have observed no sex differences in the basic cortical dynamics of taste responses between male and female rats, and therefore use female Long-Evans rats because they are, in our hands, calmer than males). The rats were housed in individual cages in a temperature- and humidity-controlled environment under a 12:12 hr light:dark cycle, given ad libitum access to food and water prior to the start of experimentation, and weighed daily following surgery to ensure that they never dropped below 80% of their pre-surgery weight. All experimental methods were in compliance with National Institutes of Health guidelines and were approved in advance by the Brandeis University Institutional Animal Care and Use Committee.

Electrode and intra-oral cannula construction

Custom microwire bundle drives, optimized for chronic (i.e., multi-day and multi-week) recording quality, were made with either 16 or 32 electrodes per recording site (design and construction details available at https://katzlab.squarespace.com/technology). Intra-oral cannulae—flexible tubing with a flanged tip and washer to ensure stability, connected to a plastic top complete with a locking mechanism—were built to allow the delivery of tastants directly onto the tongue (94).

Surgery

Rats were anaesthetized with an intraperitoneal injection of ketamine/xylazine cocktail (100mg/kg and 5.2 mg/kg respectively) and mounted in a stereotaxic instrument (David Kopf Instruments; Tujunga, CA) with blunt (atraumatic) ear bars. A midline incision exposed the skull and trephine holes (~2 mm diameter) were drilled above BLA and GC. Microwire bundles were implanted 0.5mm above GC (coordinates: AP +1.4 mm, ML -5.0 mm, DV −4.4mm from dura) and BLA (coordinates : AP -3.0mm, ML -5.0mm, DV -6.8mm from dura). Once in place, electrode bundles were cemented to the skull. Once electrode bundles were secured, an intra-oral cannula (IOC) was threaded through the masseter muscle (inside the zygomatic arch) to the space between the lip and gums, and the top of the cannula was cemented to the rest of the assembly with dental acrylic (94). The rat's body temperature was monitored and maintained at ~37°C by a heating pad throughout the duration of the surgery.

Acquisition of electrophysiological data

Electrophysiological signals from the micro-electrodes were sampled at 30 kHz using 32-channel analog-to-digital converter chips (RHD2132) from Intan Technologies, digitized online at the head stage and sampled jointly, along with signals from actuators marking tastant delivery, using an Intan RHD USB interface board (Part #C3100), which routed records to the hard drive of a PC for saving.

610 The experimental chamber was ensconced in a Faraday cage that shielded recordings from external
611 electromagnetic influences.

Habituation and passive taste administration

Following their recovery from surgery, we habituated rats to the experimental chamber for 2 days, to
the IOC/electrode harness for the next 2 days, and to passive water deliveries for the following 2 days,
before beginning data collection. Starting with the second habituation day, we also placed rats on a
mild water restriction schedule—20mL of water (not including the ~4mL delivered during habituation
sessions themselves) per day. This water restriction schedule was maintained till the end of the
experiment. For the 2 final habituation sessions, we attached the rats to the taste delivery apparatus,
and infused 120 pulses of distilled water (~30μL per pulse; 20s inter-pulse interval) into the animal's
oral cavity through the IOC, and drove electrode bundles deeper (by 250 μm) into target structures.

By the end of this procedure, the tips of the electrodes lay within GC and BLA. We then recorded taste
responses during 3-4 days of taste delivery sessions, between each of which the microwire bundle
was driven down approximately 60 μm. During these sessions, Sucrose (0.3M), Sodium Chloride
(0.1M), Citric Acid (0.1M), and Quinine (1mM), dissolved in ultra-pure water (~30μL per pulse; 20s
inter-pulse interval, 30 trials/tastant; ~3.6mL total volume delivered) were delivered to passive rats
(i.e., no behavior was required to elicit delivery). These concentrations were chosen to represent a
range of hedonic values, and because they are known to evoke robust responses in both GC and BLA
(2,24). We note that this volume has been shown to evoke stable neural responses in roughly hour-
long sessions, arguing against evocation of post-ingestive effects during the recording (31).

Histology

In preparation for histology, rats were deeply anesthetized with an overdose of the ketamine/xylazine
mixture. We perfused the rats through the heart with 0.9% saline followed by 10% formalin and
harvested the brain. The brain tissue was incubated in a fixing mixture of 30% sucrose and 10%
formalin for 7 days before being sectioned into 50μm coronal slices on a sliding microtome (Leica
SM2010R, Leica Microsystems). Sections containing the electrode implant sites around GC and BLA
were imaged at 2x.

Data and statistical analyses

638 The analysis of data and statistical tests were performed using custom written software in Python as
639 described below.

Local Field Potential processing and analysis

Filtering / power and phase extractions

LFPs were extracted from broadband digitized signals using a 2nd order bandpass Butterworth filter
(1-300Hz), to de-emphasize spiking and emphasize frequencies typically of interest in such data. In
order to avoid contamination from noise/artifacts on noisy/broken channels, only channels
containing isolable single neurons (see below) were used for analyses. Signal was down sampled to
1000Hz to reduce computational load for downstream analyses.

We note that this filtering process is sufficient for analysis up to the maximum frequency shown in
the Results section (that is, 100Hz), and is unlikely to be the reason we don't observe effects above
60Hz. First, a maximum frequency of 300Hz is sufficient for a Nyquist frequency for accurately
detecting signals up to 150Hz. Secondly, we use a Butterworth filter, which is "maximally flat" (see
scipy.signal.butter, Virtanen et al., 2020) within the pass band and is even permissive of frequencies
higher than 300Hz. Hence, signals with frequencies 100Hz or lower are unlikely to be lost due to
filtering.

Spectral Granger Causality

Spectral Granger Causality was assessed between pairs of electrodes in GC and BLA using the
Spectral Connectivity package (30). We selected for this analysis the electrodes from which activity
was most similar to the mean activity for the region (smallest mean-squared error relative to the
mean phase across all channels for each region; this selection of channels was constant for all trials
in a single analyzed session, and the same set of channels was used for all frequencies), to ensure
reliable representation of each.

To satisfy the normality assumption for use of Granger Causality, signals from both electrodes were
first preprocessed according to previously established criteria to remove outliers and "detrend" the
data (99). Briefly:

1. Any trials with data points exceeding 3 Median Absolute Deviations calculated using all the
trials, were removed
2. Linear detrending was performed on a single-trial basis (scipy.signal.detrend)
3. The temporal (single-trial) mean was subtracted from the data, and all data divided by the
temporal standard deviation
4. Trial-averaged mean was subtracted, and all data divided by the trial-averaged standard
deviation

Following preprocessing, the Augmented Dickey-Fuller test was run on each channel to confirm
stationarity (normality) of the signal. All preprocessed data passed the test and hence was included
in further analyses.

Using the preprocessed data, spectral granger causality was calculated using multitaper methods
implemented in the Spectral Connectivity package. The default set of tapers (Slepian) was used.
Specific parameters included:

• "sampling_frequency" = 1000 (Hz),
• "time_halfbandwitch_product" = 1
• "time_window_duration" = 0.3 (sec)
• "time_window_step" = 0.05 (sec)

Significance of the granger causality for each frequency-time bin on a single-session basis was
obtained using a shuffling procedure. Granger Causality was recalculated for 500 sets of data where
trial labels were shuffled (mismatched) between BLA and GC to generate a null distribution of "non-
specific" Granger Causality between the two regions. Deviation of the value in each frequency-time

bin from its respective null distribution was taken to indicate a significant interaction in that bin.
Multiple comparisons were addressed using the Bonferroni correction with a base alpha of 0.05 with
each frequency-time bin treated as a comparison. We further supplemented this "bin-level" testing
as below.

Cluster Permutation Testing

690 To perform statistical testing on the Granger Causality results while minimizing issues of multiple
691 comparisons, we opted to use a cluster-based permutation test (100,101) to detect the largest
692 statistically-significant deviation in the time-series for the 2-12Hz and 20-60Hz frequency bands.
693 Briefly, the procedure involves:

1. Z-scoring the time-series on which testing needs to be done (in our case, this is the mean significance of Granger Causality for a frequency band)
2. Assessing the summed value of the largest cluster (continuous chunk) above a specific threshold for the absolute of the z-scored time-series (we used a threshold of Z = 2). This value is used as the test statistic.
3. This summed value of the largest cluster is then recalculated from shuffled data to obtain a null-distribution for the test statistic.
4. If the actual value of the statistic significantly deviates from the null-distribution, the null hypothesis is rejected (alpha=0.05).

Granger Causality Similarity Analysis

To analyze similarities in the dynamics of the directional Granger Causality timeseries, we performed
Principal Component Analysis on the Granger Causality significance timeseries for 0-100Hz,
reducing each multivariate timeseries down to 3 components to mitigate overfitting in the following
regression analyses. All comparisons were then performed on these dimensionally reduced
timeseries. We next performed 1) Linear Regression between the two timeseries, 2) Linear regression
between circularly (temporally) shuffled versions of the two timeseries (as a negative control /
context for poor prediction), 3) Linear Autoregression, and 4) non-Linear Autoregression
(Multilayered perceptron regressor, (102). For both forms of autoregression, we used the previous 3
timepoints to predict the current one. To obtain an estimate of the uncertainty of prediction, we
performed bootstrapping of the data with 500 reruns of model fitting and prediction.

Analysis of Spiking Activity

Single Unit Isolation

Spikes from electrophysiological recordings were sorted and analyzed off-line using an in-house
Python-based pipeline (103). Putative single-neuron waveforms with >5:1 signal-to-noise ratio
(median absolute deviation) were sorted using a semi-supervised algorithm: recorded voltage data
were filtered between 300-3000Hz, grouped into potential clusters by a Gaussian Mixture Models
(GMM) fit to multiple waveform features; clusters were then labeled and/or refined manually (to
increase conservatism) by the experimenters.

 Single Unit Evoked Response Characterization

Evaluating single-neuron response taste specificity. To statistically determine the degree to which a
single neuron's response contained taste-specific information, firing rates were estimated by
convolving spike-trains with a rectangular window (length: 200ms). A one-way ANOVA (between
tastes, across trials) was then run on each window to identify if the response of a single neuron to
one taste was different from its responses to any other tastes at that timepoint. The (taste) specificity
of a single neuron was defined as the average of the ANOVA effect sizes for 2000ms of post-stimulus
activity.

Evaluating single-neuron response palatability-relatedness. To statistically determine the degree to
which a single neuron's response reflected the hedonic value of the stimuli delivered, we estimated
firing rates as described above, and calculated the Pearson's correlation coefficient between the
evoked firing rates and the palatability ranks of the tastants at each timepoint. Palatability ranks —
sucrose (1) > NaCl (2) > citric acid (3) > quinine (4) — directly reflected consumption of (earlier-run
squads of) rats in a Brief Access Task (see (2)). This ordering is canonical, and has been replicated in
many studies and with multiple measures of stimulus appreciation (94,104,105). The palatability of
a single neuron was defined as the average of the absolute correlation value (as the correlation can
also be strongly negative) for 2000ms of post-stimulus activity.

Calculating "Tastiness"

To combine taste specificity and palatability into a single metric, we first log-transformed specificity
to make the scales of specific and palatability more similar. This is because palatability as an
absolute value of a correlation is bounded between [0,1] whereas specificity as an effect size has the
range [0, ∞). Log-Specificity and Palatability were then Min-Max normalized following which the L2-
norm of the vector generated by the normalized metrics (normalized log-specificity, normalized
palatability) was calculated on a single-neuron basis and defined as "tastiness".

Determination of putative pyramidal vs inter-neurons

From the average of filtered spikes for each putative single-unit, we determined the time taken from
the trough to the next peak. As established previously, inter-neurons and pyramidal neurons can be
reliably separated using this "rise time" (4,45–47). The specific threshold varies depending on the
filtering frequencies used. We used a value of 0.45ms based on previous work using the same
bandpass filter frequencies as we have used here (4,47)

Changepoint Modeling of Population Activity

Model Fitting

Changepoint modelling of neural activity was performed as described previously (106) using the
pytau library (https://github.com/abuzarmahmood/pytau) written in Python. Briefly, spike-trains were
binned into 50ms bins. Changepoint models with independent Poisson emissions were fit to these
timeseries of binned spike-counts for BLA and GC populations separately using automatic
differentiation variational inference (ADVI) functionalities implemented in the pymc3 probabilistic
programming library (107). Once fit, the procedure returns: 1) the estimate of the posterior

distribution for single-trial population changepoint locations, and 2) the emission (firing) rates for
each neuron during every state. Since ADVI utilizes spherical Gaussians as the surrogate distribution,
we use the mode of the distribution for individual parameters as the "best estimate" of the parameter
for downstream analyses. Models with 4 states were fit to 2000ms of post-stimulus activity. Please
refer to (106) for more details on the process of model/state-number selection.

Calculation of Transition Lags

Lags between BLA and GC transitions were calculated on a single-trial basis by subtracting the "best
estimate" (see above) of the matched transition (that is, transition 1 in GC was compared to transition
1 in BLA for any given trial) between models fit to each region. This resulted in a distribution of lags
for each recording session (120 values, 1 for each trial in a session). The means of each session's
distribution were used to generate a distribution of average lags for each transition across all the
recording sessions. Finally, a grand average of each transition's distribution of average lags was
calculated. The significance of the deviation of this grand-average lag from zero was calculated by
bootstrapping at the level of trials for each session with a total of 5000 permutations. The 95%
confidence interval of each grand average was calculated using these bootstrapped distributions,
and deviation of zero from this 95% interval was taken as a significant difference.

Generalized Linear Modelling

As has been shown previously, Granger Causality methods perform poorly on spike-train data due to
the strong violation of the stationarity assumption used by the autoregressive processes on which
Granger Causality is based (108,109). This issue is addressed by General<u>ized</u> Linear Models (GLM;
distinct from <u>General</u> Linear Models which also follow the stationarity/normality assumption) which
can model Poisson-distributed data, and furthermore can model temporal dependencies explicitly
over a large range of timescales (40,57,110,111).

Model Specification and Model Fitting

The GLM included 3 sets of filters for each neuron modelled (see **Fig. 4A** for schematic):

• <u>Spike History Filter</u>: Captures how a neuron's past spiking activity influences its current firing
rate. This can model refractoriness (reduced probability of firing immediately after a spike)
and bursting behavior.
• <u>Stimulus Filter</u>: Models how external stimuli drive neural activity.
• <u>Coupling Filter</u>: Captures how the activity of other neurons influences the target neuron,
modeling functional connectivity.

The duration of the spike history and coupling filters was set to 100ms, while the duration of the
stimulus filter was set to 300ms, based on previous modelling work (40). To reduce number of fit
parameters and mitigate over-fitting of the models, raised cosine-basis functions were used to
parameterize the filters (40,41,110): a basis of 10 functions was used for every filter, with peaks
distributed logarithmically to model fast and slow timescales with appropriate temporal resolutions.

The conditional intensity of the neural activity was hence defined as:

$\lambda(t) = \exp(\beta_0 + \beta\_hist * h(t) + \beta\_stim * s(t) + \beta\_coup * c(t))$

• $\lambda(t)$ is the firing rate at time t
• β_0 is the baseline firing rate
•   h(t) represents the neuron's own spike history
•   s(t) represents the stimulus
•   c(t) represents the coupling with other neurons
•   β terms are the corresponding weights

The Python "statsmodels" library was used for model fitting (112).

Quality of fit was assessed in two ways. First, by comparing the log-likelihood of models fit to actual
data with models fit to data where the trial labels were shuffled for neurons in the ensemble being
fit, which would only allow the model to learn an "average" coupling filter between the neurons in the
model being fit. Second, we compared Spearman's Rho between predicted peri-stimulus time
histograms (PSTH) of a given neuron with PSTHs predicted for 1) the neuron the model was fit to
(actual data), 2) a different neuron (Unit Shuffled), and 3) temporally scrambled data for the same
neuron (Circular Shuffle).

Once models were fit, putative inter-region connections were determined using the coupling filters.
The fitting process returns p-values for each coefficient in the coupling filter (which, in our case due
to the parametrization of the filter using basis functions corresponded to the scaling factor of each
function in the basis). We asserted that a coupling filter indicated a statistically significant interaction
if a p-value in the filter was below alpha=0.05 following a Bonferroni correction (corrected p-value =
0.05 / 10 = 0.005).

Assessment of overlap between input and output populations
We assess whether the overlap observed between subpopulations labelled as "Input" and "Output"
in GC and BLA was different from that expected by chance using Monte Carlo simulations.
Essentially, keeping the number of Input, Output, and Total number of neurons in each region
constant, we label neurons as Input or Output randomly and assess the number of neurons with both
labels. We perform 10,000 simulations for each region to generate a null distribution for this overlap
as expected by random assignment. Deviation of the observed value of overlap from the 95%
confidence interval of this distribution would indicate that the observed value is significantly
different from that expected by chance.

Data Availability

We have structured our electrophysiology datasets in a hierarchical data format (HDF5) and are
hosting the files on a university-wide network share managed by Library and Technology Services
(LTS) at Brandeis University. These HDF5 files contain our electrophysiology recordings, sorted
spikes, single-neuron and population-level analyses (and associated plots and results). These files
are prohibitively large to be hosted on a general-purpose fileshare platform - we request anyone
interested in our datasets to contact the corresponding author, Donald Katz (dbkatz@brandeis.edu)

who can put them in touch with LTS in order to create a guest account at Brandeis through which they
can securely access our datasets (hosted on the katz-lab share at files.brandeis.edu).

References

1. Katz DB, Simon SA, Nicolelis MAL. Dynamic and Multimodal Responses of Gustatory Cortical Neurons in Awake Rats. J Neurosci. 2001 June 15;21(12):4478–89.

2. Sadacca BF, Rothwax JT, Katz DB. Sodium Concentration Coding Gives Way to Evaluative Coding in Cortex and Amygdala. J Neurosci. 2012 July 18;32(29):9999–10011.

3. Sadacca BF, Mukherjee N, Vladusich T, Li JX, Katz DB, Miller P. The Behavioral Relevance of Cortical Neural Ensemble Responses Emerges Suddenly. J Neurosci. 2016 Jan 20;36(3):655–69.

4. Lin JY, Mukherjee N, Bernstein MJ, Katz DB. Perturbation of amygdala-cortical projections reduces ensemble coherence of palatability coding in gustatory cortex. Schoenbaum G, Wassum KM, editors. eLife. 2021 May 21;10:e65766.

5. van Bergen RS, Kriegeskorte N. Going in circles is the way forward: the role of recurrence in visual inference. Current Opinion in Neurobiology. 2020 Dec 1;65:176–93.

6. Kay LM, Freeman WJ. Bidirectional Processing in the Olfactory-Limbic Axis During Olfactory Behavior. 2008;13.

7. Brincat SL, Connor CE. Dynamic Shape Synthesis in Posterior Inferotemporal Cortex. Neuron. 2006 Jan 5;49(1):17–24.

8. Gaillard R, Dehaene S, Adam C, Clémenceau S, Hasboun D, Baulac M, et al. Converging Intracranial Markers of Conscious Access. PLOS Biology. 2009 Mar 17;7(3):e1000061.

9. Kietzmann TC, Spoerer CJ, Sörensen LKA, Cichy RM, Hauk O, Kriegeskorte N. Recurrence is required to capture the representational dynamics of the human visual system. Proceedings of the National Academy of Sciences. 2019 Oct 22;116(43):21854–63.

10. Mahmood A, Steindler J, Germaine H, Miller P, Katz DB. Coupled Dynamics of Stimulus-Evoked Gustatory Cortical and Basolateral Amygdalar Activity. J Neurosci. 2023 Jan 18;43(3):386–404.

11. Maffei A, Haley M, Fontanini A. Neural processing of gustatory information in insular circuits. Current Opinion in Neurobiology. 2012 Aug 1;22(4):709–16.

12. Haley MS, Fontanini A, Maffei A. Laminar- and Target-Specific Amygdalar Inputs in Rat Primary Gustatory Cortex. J Neurosci. 2016 Mar 2;36(9):2623–37.

13. Stone ME, Maffei A, Fontanini A. Amygdala Stimulation Evokes Time-Varying Synaptic Responses in the Gustatory Cortex of Anesthetized Rats. Front Integr Neurosci [Internet]. 2011 [cited 2019 May 8];5. Available from: https://www.frontiersin.org/articles/10.3389/fnint.2011.00003/full

14. Shi CJ, Cassell MD. Cortical, thalamic, and amygdaloid connections of the anterior and posterior insular cortices. The Journal of Comparative Neurology. 1998 Oct 5;399(4):440–68.

15.   McDonald AJ. Cortical pathways to the mammalian amygdala. Progress in Neurobiology. 1998
June 27;55(3):257–332.

16.   Arce-McShane FI, Ross CF, Takahashi K, Sessle BJ, Hatsopoulos NG. Primary motor and
sensory cortical areas communicate via spatiotemporally coordinated networks at multiple
frequencies. Proceedings of the National Academy of Sciences. 2016 May 3;113(18):5083–8.

17.   Mejias JF, Murray JD, Kennedy H, Wang XJ. Feedforward and feedback frequency-dependent
interactions in a large-scale laminar network of the primate cortex. Science Advances. 2016
Nov 1;2(11):e1601335.

18.   Piette CE, Baez-Santiago MA, Reid EE, Katz DB, Moran A. Inactivation of Basolateral Amygdala
Specifically Eliminates Palatability-Related Information in Cortical Sensory Responses. J
Neurosci. 2012 July 18;32(29):9981–91.

19.   Samuelsen CL, Gardner MPH, Fontanini A. Thalamic Contribution to Cortical Processing of
Taste and Expectation. J Neurosci. 2013 Jan 30;33(5):1815–27.

20.   Lavi K, Jacobson GA, Rosenblum K, Lüthi A. Encoding of Conditioned Taste Aversion in
Cortico-Amygdala Circuits. Cell Reports. 2018 July 10;24(2):278–83.

21.   Lin JY, Arthurs J, Reilly S. The effects of amygdala and cortical inactivation on taste neophobia.
Neurobiology of Learning and Memory. 2018 Nov 1;155:322–9.

22.   Arieli E, Gerbi R, Shein-Idelson M, Moran A. Temporally-precise basolateral amygdala
activation is required for the formation of taste memories in gustatory cortex. The Journal of
Physiology. 2020;598(23):5505–22.

23.   Grossman SE, Fontanini A, Wieskopf JS, Katz DB. Learning-Related Plasticity of Temporal
Coding in Simultaneously Recorded Amygdala–Cortical Ensembles. J Neurosci. 2008 Mar
12;28(11):2864–73.

24.   Fontanini A, Grossman SE, Figueroa JA, Katz DB. Distinct Subtypes of Basolateral Amygdala
Taste Neurons Reflect Palatability and Reward. J Neurosci. 2009 Feb 25;29(8):2486–95.

25.   Mukherjee N, Wachutka J, Katz DB. Impact of precisely-timed inhibition of gustatory cortex on
taste behavior depends on single-trial ensemble dynamics. Colgin LL, Maffei A, Maffei A,
Ghazanfar AA, Verhagen J, Mazzucato L, editors. eLife. 2019 June 24;8:e45968.

26.   Ding M, Chen Y, Bressler SL. Granger Causality: Basic Theory and Application to Neuroscience
[Internet]. arXiv; 2006 [cited 2023 Apr 26]. Available from: http://arxiv.org/abs/q-bio/0608035

27.   Seth AK, Barrett AB, Barnett L. Granger causality analysis in neuroscience and neuroimaging. J
Neurosci. 2015 Feb 25;35(8):3293–7.

28.   Geweke J. Measurement of Linear Dependence and Feedback Between Multiple Time Series.
Journal of the American Statistical Association. 1982;77(378):304–13.

29.   Chicharro D. On the spectral formulation of Granger causality. Biol Cybern. 2011 Dec;105(5–
906   6):331–47.

907   30.   Denovellis EL, Myroshnychenko M, Sarmashghi M, Stephen EP. Spectral Connectivity: a
908   python package for computing multitaper spectral estimates and frequency-domain brain
909   connectivity measures on the CPU and GPU. Journal of Open Source Software. 2022 Dec
910   13;7(80):4840.

911   31.   Fontanini A, Katz DB. 7 to 12 Hz Activity in Rat Gustatory Cortex Reflects Disengagement From
912   a Fluid Self-Administration Task. Journal of Neurophysiology. 2005 May 1;93(5):2832–40.

913   32.   Tort ABL, Fontanini A, Kramer MA, Jones-Lush LM, Kopell NJ, Katz DB. Cortical Networks
914   Produce Three Distinct 7–12 Hz Rhythms during Single Sensory Responses in the Awake Rat. J
915   Neurosci. 2010 Mar 24;30(12):4315–24.

916   33.   Pavão R, Piette CE, Lopes-dos-Santos V, Katz DB, Tort ABL. Local Field Potentials in the
917   Gustatory Cortex Carry Taste Information. J Neurosci. 2014 June 25;34(26):8778–87.

918   34.   Baccalá LA, Sameshima K. Partial directed coherence: a new concept in neural structure
919   determination. Biol Cybern. 2001 May 1;84(6):463–74.

920   35.   van Kerkoerle T, Self MW, Dagnino B, Gariel-Mathis MA, Poort J, van der Togt C, et al. Alpha and
921   gamma oscillations characterize feedback and feedforward processing in monkey visual
922   cortex. Proc Natl Acad Sci U S A. 2014 Oct 7;111(40):14332–41.

923   36.   Bastos AM, Vezoli J, Bosman CA, Schoffelen JM, Oostenveld R, Dowdall JR, et al. Visual Areas
924   Exert Feedforward and Feedback Influences through Distinct Frequency Channels. Neuron.
925   2015 Jan 21;85(2):390–401.

926   37.   Bechara A, Damasio H, Damasio AR, Lee GP. Different Contributions of the Human Amygdala
927   and Ventromedial Prefrontal Cortex to Decision-Making. J Neurosci. 1999 July 1;19(13):5473–
928   81.

929   38.   Schoenbaum G, Chiba AA, Gallagher M. Neural Encoding in Orbitofrontal Cortex and
930   Basolateral Amygdala during Olfactory Discrimination Learning. J Neurosci. 1999 Mar
931   1;19(5):1876–84.

932   39.   Michalareas G, Vezoli J, van Pelt S, Schoffelen JM, Kennedy H, Fries P. Alpha-Beta and Gamma
933   Rhythms Subserve Feedback and Feedforward Influences among Human Visual Cortical
934   Areas. Neuron. 2016 Jan 20;89(2):384–97.

935   40.   Pillow JW, Shlens J, Paninski L, Sher A, Litke AM, Chichilnisky EJ, et al. Spatio-temporal
936   correlations and visual signalling in a complete neuronal population. Nature. 2008
937   Aug;454(7207):995–9.

938   41.   Stevenson IH, Rebesco JM, Miller LE, Körding KP. Inferring functional connections between
939   neurons. Curr Opin Neurobiol. 2008 Dec;18(6):582–8.

940   42.   Magrans de Abril I, Yoshimoto J, Doya K. Connectivity inference from neural recording data:
941   Challenges, mathematical bases and research directions. Neural Networks. 2018 June
942   1;102:120–37.

943   43.   Kobayashi R, Kurita S, Kurth A, Kitano K, Mizuseki K, Diesmann M, et al. Reconstructing
944   neuronal circuitry from parallel spike trains. Nat Commun. 2019 Oct 2;10(1):4468.

945   44.   Markov NT, Vezoli J, Chameau P, Falchier A, Quilodran R, Huissoud C, et al. Anatomy of
946   hierarchy: Feedforward and feedback pathways in macaque visual cortex. Journal of
947   Comparative Neurology. 2014;522(1):225–59.

948   45.   Sirota A, Montgomery S, Fujisawa S, Isomura Y, Zugaro M, Buzsáki G. Entrainment of
949   Neocortical Neurons and Gamma Oscillations by the Hippocampal Theta Rhythm. Neuron.
950   2008 Nov 26;60(4):683–97.

951   46.   Quirk MC, Sosulski DL, Feierstein CE, Uchida N, Mainen ZF. A defined network of fast-spiking
952   interneurons in orbitofrontal cortex: responses to behavioral contingencies and ketamine
953   administration. Front Syst Neurosci [Internet]. 2009 Nov 3 [cited 2025 May 8];3. Available
954   from: https://www.frontiersin.orghttps://www.frontiersin.org/journals/systems-
955   neuroscience/articles/10.3389/neuro.06.013.2009/full

956   47.   Herzog LE, Pascual LM, Scott SJ, Mathieson ER, Katz DB, Jadhav SP. Interaction of Taste and
957   Place Coding in the Hippocampus. J Neurosci. 2019 Apr 17;39(16):3057–69.

958   48.   Li JX, Yoshida T, Monk KJ, Katz DB. Lateral Hypothalamus Contains Two Types of Palatability-
959   Related Taste Responses with Distinct Dynamics. J Neurosci. 2013 May 29;33(22):9462–73.

960   49.   Moran A, Katz DB. Sensory Cortical Population Dynamics Uniquely Track Behavior across
961   Learning and Extinction. J Neurosci. 2014 Jan 22;34(4):1248–57.

962   50.   Kosar E, Grill HJ, Norgren R. Gustatory cortex in the rat. II. Thalamocortical projections. Brain
963   Research. 1986 Aug 6;379(2):342–52.

964   51.   Carleton A, Accolla R, Simon SA. Coding in the mammalian gustatory system. Trends in
965   Neurosciences. 2010 July 1;33(7):326–34.

966   52.   Miller P, Katz DB. Stochastic Transitions between Neural States in Taste Processing and
967   Decision-Making. J Neurosci. 2010 Feb 17;30(7):2559–70.

968   53.   Miller P, Katz DB. Accuracy and response-time distributions for decision-making: linear
969   perfect integrators versus nonlinear attractor-based neural circuits. J Comput Neurosci. 2013
970   Dec 1;35(3):261–94.

971   54.   Mazzucato L, Fontanini A, Camera GL. Dynamics of Multistable States during Ongoing and
972   Evoked Cortical Activity. J Neurosci. 2015 May 27;35(21):8214–31.

973   55.   Lang L, Camera GL, Fontanini A. Temporal progression along discrete coding states during
974   decision-making in the mouse gustatory cortex. PLOS Computational Biology. 2023 Feb
975   7;19(2):e1010865.

976   56.   Parras GG, Nieto-Diego J, Carbajal GV, Valdés-Baizabal C, Escera C, Malmierca MS. Neurons
977   along the auditory pathway exhibit a hierarchical organization of prediction error. Nat
978   Commun. 2017 Dec 15;8(1):2148.

979   57.   Glaser JI, Perich MG, Ramkumar P, Miller LE, Kording KP. Population coding of conditional
980   probability distributions in dorsal premotor cortex. Nature Communications. 2018 May
981   3;9(1):1–14.

982   58.   Heidari-Gorji H, Ebrahimpour R, Zabbah S. A temporal hierarchical feedforward model
983   explains both the time and the accuracy of object recognition. Sci Rep. 2021 Mar
984   11;11(1):5640.

985   59.   Baez-Santiago MA, Reid EE, Moran A, Maier JX, Marrero-Garcia Y, Katz DB. Dynamic taste
986   responses of parabrachial pontine neurons in awake rats. Journal of Neurophysiology. 2016
987   Jan 20;115(3):1314–23.

988   60.   Livneh Y, Sugden AU, Madara JC, Essner RA, Flores VI, Sugden LA, et al. Estimation of Current
989   and Future Physiological States in Insular Cortex. Neuron. 2020 Mar 18;105(6):1094-1111.e10.

990   61.   Fonseca E, de Lafuente V, Simon SA, Gutierrez R. Sucrose intensity coding and decision-
991   making in rat gustatory cortices. Schoenbaum G, Frank MJ, Schoenbaum G, Wesson DW,
992   Livneh Y, editors. eLife. 2018 Nov 19;7:e41152.

993   62.   Alvarez-Crespo M, Skibicka KP, Farkas I, Molnár CS, Egecioglu E, Hrabovszky E, et al. The
994   Amygdala as a Neurobiological Target for Ghrelin in Rats: Neuroanatomical,
995   Electrophysiological and Behavioral Evidence. PLOS ONE. 2012 Oct 10;7(10):e46321.

996   63.   Gründemann J, Bitterman Y, Lu T, Krabbe S, Grewe BF, Schnitzer MJ, et al. Amygdala
997   ensembles encode behavioral states. Science. 2019 Apr 19;364(6437):eaav8736.

998   64.   Dehaene S, Sergent C, Changeux JP. A neuronal network model linking subjective reports and
999   objective physiological data during conscious perception. Proceedings of the National
1000   Academy of Sciences. 2003 July 8;100(14):8520–5.

1001   65.   Rogers PJ. Why a palatability construct is needed. Appetite. 1990 June;14(3):167–70;
1002   discussion 180.

1003   66.   Araujo JA, Studzinski CM, Larson BT, Milgram NW. Comparison of the cognitive palatability
1004   assessment protocol and the two-pan test for use in assessing palatability of two similar
1005   foods in dogs. 2004 Nov 1 [cited 2025 May 6]; Available from:
1006   https://avmajournals.avma.org/view/journals/ajvr/65/11/ajvr.65.11.1490.xml

1007   67.   Breslin PAS. The Sense of Taste Encompasses Two Roles: Conscious Taste Perception and
1008   Subconscious Metabolic Responses. In: Think Tank [Internet]. Yale University Press; 2018
1009   [cited 2025 May 6]. p. 110–8. Available from:
1010   https://www.degruyterbrill.com/document/doi/10.12987/9780300235470-
1011   017/html?srsltid=AfmBOopo1-t48Heuq2k_xlyWiTnwcA6QP4aYMaFChOt_KAeaiSt3OwPX

68. Vincis R, Chen K, Czarnecki L, Chen J, Fontanini A. Dynamic Representation of Taste-Related
Decisions in the Gustatory Insular Cortex of Mice. Current Biology. 2020 May 18;30(10):1834-
1844.e5.

69. Supp GG, Siegel M, Hipp JF, Engel AK. Cortical Hypersynchrony Predicts Breakdown of
Sensory Processing during Loss of Consciousness. Current Biology. 2011 Dec 6;21(23):1988–
93.

70. Luo TZ, Kim TD, Gupta D, Bondy AG, Kopec CD, Elliot VA, et al. Transitions in dynamical regime
and neural mode underlie perceptual decision-making [Internet]. bioRxiv; 2024 [cited 2024
Nov 18]. p. 2023.10.15.562427. Available from:
https://www.biorxiv.org/content/10.1101/2023.10.15.562427v3

71. Frederick DE, Brown A, Brim E, Mehta N, Vujovic M, Kay LM. Gamma and Beta Oscillations
Define a Sequence of Neurocognitive Modes Present in Odor Processing. J Neurosci. 2016 July
20;36(29):7750–67.

72. Skinner FK, Gramoll S, Calabrese RL, Kopell N, Marder E. Frequency Control in Biological Half-
Center Oscillators. In: Eeckman FH, editor. Computation in Neurons and Neural Systems
[Internet]. Boston, MA: Springer US; 1994 [cited 2025 May 6]. p. 223–8. Available from:
https://doi.org/10.1007/978-1-4615-2714-5_36

73. Diz-Pita É, Otero-Espinar MV. Predator–Prey Models: A Review of Some Recent Advances.
Mathematics. 2021 Jan;9(15):1783.

74. Vrugt M te. An introduction to reservoir computing [Internet]. arXiv; 2024 [cited 2025 May 6].
Available from: http://arxiv.org/abs/2412.13212

75. Arce-McShane FI, Ross CF, Takahashi K, Sessle BJ, Hatsopoulos NG. Primary motor and
sensory cortical areas communicate via spatiotemporally coordinated networks at multiple
frequencies. PNAS. 2016 Apr 15;201600788.

76. Forseth KJ, Pitkow X, Fischer-Baum S, Tandon N. What The Brain Does As We Speak [Internet].
bioRxiv; 2021 [cited 2025 May 6]. p. 2021.02.05.429841. Available from:
https://www.biorxiv.org/content/10.1101/2021.02.05.429841v1

77. Ahissar E, Assa E. Perception as a closed-loop convergence process. Kleinfeld D, editor. eLife.
2016 May 9;5:e12830.

78. Li JX, Maier JX, Reid EE, Katz DB. Sensory Cortical Activity Is Related to the Selection of a
Rhythmic Motor Action Pattern. J Neurosci. 2016 May 18;36(20):5596–607.

79. Moriyama Y. Rhythmical jaw movements and lateral ponto-medullary reticular neurons in rats.
Comparative Biochemistry and Physiology Part A: Physiology. 1987 Jan 1;86(1):7–14.

80. Schafe GE, Bernstein IL. Forebrain contribution to the induction of a brainstem correlate of
conditioned taste aversion: II. Insular (gustatory) cortex. Brain Research. 1998 July
27;800(1):40–7.

81. Travers JB, DiNardo LA, Karimnamazi H. Medullary reticular formation activity during ingestion
and rejection in the awake rat. Exp Brain Res. 2000 Jan;130(1):78–92.

82. Gehrlach DA, Weiand C, Gaitanos TN, Cho E, Klein AS, Hennrich AA, et al. A whole-brain
connectivity map of mouse insular cortex. Wassum KM, Livneh Y, editors. eLife. 2020 Sept
17;9:e55585.

83. Ponserre M, Peters C, Fermani F, Conzelmann KK, Klein R. The Insula Cortex Contacts Distinct
Output Streams of the Central Amygdala. J Neurosci. 2020 Nov 11;40(46):8870–82.

84. Shammah-Lagnado SJ, Costa MSMO, Ricardo⚔ JA. Afferent connections of the parvocellular
reticular formation: A horseradish peroxidase study in the rat. Neuroscience. 1992 Sept
1;50(2):403–25.

85. Karimnamazi H, Travers JB. Differential projections from gustatory responsive regions of the
parabrachial nucleus to the medulla and forebrain. Brain Research. 1998 Dec 7;813(2):283–
302.

86. Wang L, Gillis-Smith S, Peng Y, Zhang J, Chen X, Salzman CD, et al. The coding of valence and
identity in the mammalian taste system. Nature. 2018 June;558(7708):127–31.

87. Jin H, Fishman ZH, Ye M, Wang L, Zuker CS. Top-Down Control of Sweet and Bitter Taste in the
Mammalian Brain. Cell. 2021 Jan 7;184(1):257-271.e16.

88. Accolla R, Bathellier B, Petersen CCH, Carleton A. Differential Spatial Representation of Taste
Modalities in the Rat Gustatory Cortex. J Neurosci. 2007 Feb 7;27(6):1396–404.

89. Fletcher ML, Ogg MC, Lu L, Ogg RJ, Boughter JD. Overlapping representation of primary tastes
in a defined region of the gustatory cortex. J Neurosci. 2017 July 3;0649–17.

90. Levitan D, Lin JY, Wachutka J, Mukherjee N, Nelson SB, Katz DB. Single and population coding
of taste in the gustatory cortex of awake mice. J Neurophysiol. 2019 Oct 1;122(4):1342–56.

91. Chen K, Kogan JF, Fontanini A. Spatially Distributed Representation of Taste Quality in the
Gustatory Insular Cortex of Behaving Mice. Current Biology. 2021 Jan 25;31(2):247-256.e4.

92. de Araujo IE, Gutierrez R, Oliveira-Maia AJ, Pereira A, Nicolelis MAL, Simon SA. Neural
Ensemble Coding of Satiety States. Neuron. 2006 Aug 17;51(4):483–94.

93. Stone BT, Lin JY, Mahmood A, Sanford AJ, Katz DB. LiCl-induced sickness modulates rat
gustatory cortical responses. PLOS Biology. 2022 July 25;20(7):e3001537.

94. Fontanini A, Katz DB. State-Dependent Modulation of Time-Varying Gustatory Responses.
Journal of Neurophysiology. 2006 Dec 1;96(6):3183–93.

95. Engel AK, Gerloff C, Hilgetag CC, Nolte G. Intrinsic Coupling Modes: Multiscale Interactions in
Ongoing Brain Activity. Neuron. 2013 Nov 20;80(4):867–86.

96. Kato S, Kaplan HS, Schrödel T, Skora S, Lindsay TH, Yemini E, et al. Global Brain Dynamics
Embed the Motor Command Sequence of Caenorhabditis elegans. Cell. 2015 Oct
22;163(3):656–69.

97. Raut RV, Snyder AZ, Raichle ME. Hierarchical dynamics as a macroscopic organizing principle
of the human brain. Proceedings of the National Academy of Sciences. 2020 Aug
25;117(34):20890–7.

98. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0:
fundamental algorithms for scientific computing in Python. Nat Methods. 2020 Mar;17(3):261–
72.

99. Ding M, Bressler SL, Yang W, Liang H. Short-window spectral analysis of cortical event-related
potentials by adaptive multivariate autoregressive modeling: data preprocessing, model
validation, and variability assessment. Biol Cybern. 2000 June 1;83(1):35–45.

100. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. Journal of
Neuroscience Methods. 2007 Aug 15;164(1):177–90.

101. Candia-Rivera D, Valenza G. Cluster permutation analysis for EEG series based on non-
parametric Wilcoxon–Mann–Whitney statistical tests. SoftwareX. 2022 July 1;19:101170.

102. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. J Mach Learn Res. 2011 Nov 1;12(null):2825–30.

103. Mahmood A, Mukherjee N, Stone B, Raymond M, Germaine H, Lin JY, et al.
katzlabbrandeis/blech_clust: v1.1.0 [Internet]. Zenodo; 2025 [cited 2025 May 15]. Available
from: https://zenodo.org/records/15175273

104. Clarke SNDA, Ossenkopp KP. Taste reactivity responses in rats: influence of sex and the
estrous cycle. American Journal of Physiology-Regulatory, Integrative and Comparative
Physiology. 1998 Mar;274(3):R718–24.

105. Travers JB, Norgren R. Electromyographic analysis of the ingestion and rejection of sapid
stimuli in the rat. Behavioral Neuroscience. 1986;100:544–55.

106. Mahmood A, Steindler J, Germaine H, Miller P, Katz DB. Coupled Dynamics of Stimulus-
Evoked Gustatory Cortical and Basolateral Amygdalar Activity. J Neurosci. 2023 Jan
18;43(3):386–404.

107. Salvatier J, Wiecki TV, Fonnesbeck C. Probabilistic programming in Python using PyMC3. PeerJ
Comput Sci. 2016 Apr 6;2:e55.

108. Kispersky T, Gutierrez GJ, Marder E. Functional connectivity in a rhythmic inhibitory circuit
using Granger causality. Neural Syst Circuits. 2011 May 25;1:9.

109. Gerhard F, Kispersky T, Gutierrez GJ, Marder E, Kramer M, Eden U. Successful Reconstruction
of a Physiological Circuit with Known Connectivity from Spiking Activity Alone. PLOS
Computational Biology. 2013 July 11;9(7):e1003138.

110. Stevenson IH, Rebesco JM, Hatsopoulos NG, Haga Z, Miller LE, Kording KP. Bayesian Inference
of Functional Connectivity and Network Structure From Spikes. IEEE Transactions on Neural
Systems and Rehabilitation Engineering. 2009 June;17(3):203–13.

111. Keeley SL, Zoltowski DM, Aoi MC, Pillow JW. Modeling statistical dependencies in multi-region
spike train data. Current Opinion in Neurobiology. 2020 Dec 1;65:194–202.

112. Seabold S, Perktold J. Statsmodels: Econometric and Statistical Modeling with Python. In
Austin, Texas; 2010 [cited 2025 May 16]. p. 92–6. Available from:
https://doi.curvenote.com/10.25080/Majora-92bf1922-011

1125