Morning Elevation in Insulin Enhances Afternoon Postprandial Insulin Action and

Glucose Effectiveness

Hannah L. Waterman[1], Marta S. Smith[1], Ben Farmer[1], Kalisha Yankey[1], Tristan Howard[1],

Guillaume Kraft[1], Alan D. Cherrington[1], Dale S. Edgerton[1]

[1]Department of Molecular Physiology & Biophysics, Vanderbilt University School of Medicine

Corresponding author information:

Name: Hannah Waterman

Current Affiliation: Vanderbilt University School of Medicine

Phone: (360)-536-1498

Email: Hannah.L.Waterman@Vanderbilt.Edu

Running title: AM Insulin Enhances PM Insulin Action and Glucose Effectiveness

Keywords: hepatic glucose uptake, glucose metabolism, liver metabolism, glycogen, glucagon,

insulin, meal composition, breakfast, lunch, second meal phenomenon, second meal effect,

Staub-Traugott effect, insulin action, glucose effectiveness, postprandial, insulin signaling,

glucose

Abstract

The second meal phenomenon refers to the improved glycemic response observed after consuming a second identical meal. It is well known that postprandial hepatic glucose uptake (HGU) is determined by the combined effects of three regulatory factors: insulin action (hyperinsulinemia; HI), glucose effectiveness (hyperglycemia; GE), and glucose delivery into the hepatoportal circulation (portal glucose signal; PGS). While our previous studies demonstrated the importance of morning (AM) hyperinsulinemia in priming the liver for increased HGU later in the day, the question remains as to which aspect of the afternoon (PM) response is enhanced by morning insulin. Thus, to provide insight into the underlying mechanism of the 2^{nd} meal effect, we sought to determine the extent to which PM HI, GE, and the PGS are impacted by AM hyperinsulinemia. Dogs underwent an AM clamp with either a 4h hyperinsulinemic prime (Prime, *n=8*) or basal insulin delivery (No Prime, *n=8*). After a 1.5h non-clamp period, both groups were challenged with a PM hyperglycemic clamp (with the PGS) in the presence of basal insulin. During the PM clamp, net HGU was significantly elevated in Prime vs. No Prime group (mean of 2.2±0.3 vs. 0.1±0.3 mg/kg/min, respectively, P<0.005), indicating an enhancement of GE and/or the PGS. There was no difference in PM non-HGU (4.5±0.3 vs. 4.0±0.3 mg/kg/min). In previous experiments, when all three factors (HI, GE, PGS) were present in the PM, net HGU was 6.3±1.7 vs. 2.4±1.1 mg/kg/min in groups receiving an AM insulin prime vs. basal insulin, respectively. Therefore, the AM insulin prime enhanced the effects of HI, GE, and the PGS on PM HGU by 3.9 mg/kg/min, whereas it enhanced the effects of glucose (GE and the PGS), in the absence of a rise in insulin, by 2.1 mg/kg/min. Taken together, the data show that AM HI enhanced PM insulin action and GE (including the PGS) to an equal extent. This suggests that AM insulin priming enhances cellular targets common to both insulin and glucose signaling in the PM.

Introduction

Postprandial glucose regulation is essential for maintaining metabolic homeostasis, as impaired glucose disposal is a hallmark of insulin resistance and type 2 diabetes (1; 2). A well-characterized but not fully understood phenomenon in glucose metabolism is the 2nd meal effect, where the glycemic response to a second, identical meal is improved compared to the first (3-5). This adaptive response is linked to changes in liver glucose metabolism, specifically hepatic glucose uptake (HGU), which plays a crucial role in regulating postprandial glucose excursions (6). While the general improvement in HGU following the second meal is recognized, the precise mechanisms underlying how prior insulin exposure enhances subsequent HGU remain to be elucidated.

The regulation of postprandial HGU is influenced by the interaction of three primary factors: insulin action (hyperinsulinemia; HI), glucose effectiveness (hyperglycemia; GE), and the portal glucose signal (PGS) (7; 8). In this context, hyperinsulinemia refers to the physiological rise in insulin levels following a meal, which promotes insulin-dependent glucose uptake in the liver and peripheral tissues (9). Insulin's action on the liver helps regulate both glucose production and storage, and its effects on HGU are key in postprandial glucose clearance, even equaling muscle (10). Glucose effectiveness, on the other hand, describes the ability of glucose itself to enhance its own uptake and storage, independent of insulin (11). This insulin-independent effect contributes significantly to glucose homeostasis, especially when insulin action is compromised (12; 13). The portal glucose signal is the third factor, referring to the neural response that occurs when glucose is absorbed or delivered directly into the hepatoportal circulation (7). This signal plays a crucial role in coordinating the liver's response to changes in glucose levels, ensuring efficient glucose utilization (14-16).

In our previous work, we demonstrated that morning (AM) hyperinsulinemia primes the

liver for an enhanced HGU response to a hyperinsulinemic-hyperglycemic (HIHG) clamp later in

the day (17). However, the specific mechanisms driving the augmented HGU observed in the

afternoon (PM) response, particularly in the context of the 2nd meal phenomenon, are not fully

understood. While AM hyperinsulinemia has been shown to enhance the liver's response to the

full combination of factors that regulate HGU (HI, GE, and PGS), it is unclear which of these

factors is responsible for the enhanced HGU observed during a second meal.

We hypothesize that AM hyperinsulinemia will enhance PM glucose uptake through both

insulin-dependent and insulin-independent mechanisms, indicating that hepatic metabolic

pathways common to both insulin action and glucose effectiveness are primed by prior insulin

exposure for subsequent HGU. By dissecting the contributions of insulin action, glucose

effectiveness, and the portal glucose signal, this study will provide new mechanistic insights into

the temporal dynamics of hepatic glucose metabolism. Ultimately, this research will expand our

understanding of how metabolic conditions established in the morning influences subsequent

postprandial responses and could have significant therapeutic implications for improving

glucose control, particularly in individuals with glucose intolerance or type 2 diabetes.

Research Design and Methods

Animal care and surgical procedures

16 adult mongrel dogs (8 males, 8 females) with an average body weight of 27.6 ± 1.0

kg were used in these experiments. The animals were obtained from a USDA-approved vendor

and housed according to the standards set by the American Association for the Accreditation of

Laboratory Animal Care. All procedures were reviewed and approved by the Vanderbilt

Institutional Animal Care and Use Committee. Two weeks before the experimental procedures,

the animals underwent a laparotomy under general anesthesia for the surgical placement of

blood flow probes around the hepatic artery and hepatic portal vein. Additionally, catheters were

implanted for blood sampling in the hepatic vein, hepatic portal vein, and femoral artery. Infusion

catheters were positioned in the splenic and jejunal veins, which supply the hepatic portal vein,

as well as the inferior vena cava. These catheters were secured subcutaneously until the day of

the experiment, when they were exteriorized. The dogs were fed a controlled diet consisting of

chow and meat, with a macronutrient composition of 46% carbohydrates, 34% protein, 14.5%

fat, and 5.5% fiber. Before the experiment, the animals underwent an 18-hour fasting period.

Assessments were conducted to ensure the dogs were healthy before the experiment, requiring

that the dogs had consumed at least 75% of their most recent meal, maintained a leukocyte

count below 18,000/mm³, and had a hematocrit level above 34%. Blood collection was limited to

no more than 20% of the animal's total blood volume over the course of a day.

Experimental Design

The experimental protocol lasted 8 hours and included two glucose clamp phases,

separated by a rest period. These clamps were designed to simulate postprandial glucose

dynamics at times corresponding to morning and midday meals. The first phase, referred to as

the AM clamp, occurred from 0 to 240 minutes, followed by a break before the PM clamp, which

took place from 330 to 480 minutes **(Fig. 1).** Blood samples were collected every 15 to 30

minutes from the femoral artery, hepatic portal vein, and hepatic vein to measure hormone and

metabolite levels. Arterial plasma glucose was assessed every five minutes, and exogenous

glucose infusion into a peripheral vein was adjusted as needed to maintain target glucose

concentrations.

Morning (AM) Clamp Period (0-240 min) and Non-Clamp Period (240 – 330 min)

To suppress endogenous pancreatic insulin and glucagon secretion, somatostatin

(Bachem, Torrance, CA) was infused into the inferior vena cava (0.8 µg/kg/min) at the onset of

the AM clamp. Glucagon (GlucaGen, Boehringer Ingelheim, Ridgefield, CT) was replaced

intraportally at basal levels (0.57 ng/kg/min). One of the experimental groups, Prime, received

morning insulin (Novolin R; Novo Nordisk, Basværd, Denmark) infusion designed to replicate

the physiologic insulin secretion pattern previously observed following a morning duodenal

glucose infusion (replaced intraportally at 2.1 mU/kg/min from 0-30 min, 2.4 mU/kg/min from 30-

60 min, and 1.5 mU/kg/min from 60-240 min) (17). The other group, No Prime, received

intraportal basal insulin replacement in the morning (0.25 mU/kg/min). Prior research has shown

that the enhancement of HGU in the afternoon is driven by an increase in insulin during the

morning, rather than a rise in morning glucose levels (17). Therefore, glucose was clamped at

euglycemia (~100 mg/dL plasma glucose) for the entirety of the AM clamp. To allow for tracer-

determined HGU measurements in the afternoon period, a peripheral infusion of [3-^{3}H] glucose

(Revvitty, Waltham, MA) was initiated at 180 min in both groups, with a priming dose of 38 µCi,

followed by a continuous infusion at 0.38 µCi/min. After the AM clamp (240 min), all infusions,

except for the tracer, were discontinued for 1.5h (non-clamp period). To characterize the hepatic

metabolic state prior to the initiation of the PM clamp, blood samples were obtained from the

femoral artery, portal vein, and hepatic vein at 300, 315, and 330 minutes **(Fig. 1).**

Afternoon (PM) Clamp Period (330 – 480 min)

Both the Prime and No Prime groups underwent an identical 2.5-h hyperglycemic clamp

with basal insulin and glucagon during the afternoon phase **(Fig. 1).** This protocol was designed

to isolate the effects of the morning insulin prime on PM insulin action vs. glucose and the portal

glucose signal. Thus, basal insulin levels were maintained throughout the PM clamp. To achieve

this, somatostatin was continuously infused via a peripheral vein (0.8 µg/kg/min), while basal

glucagon (0.57 ng/kg/min) and fourfold basal insulin (1.2 mU/kg/min) were delivered

intraportally. To stimulate the portal glucose signal, glucose was directly infused into the hepatic

portal vein at a rate of 4 mg/kg/min. Additionally, a primed, continuous glucose infusion into a

peripheral vein was administered to rapidly elevate plasma glucose levels to ~200 mg/dL,

maintaining hyperglycemia for the duration of the PM clamp. These conditions were intended to

mimic the postprandial metabolic state seen following carbohydrate ingestion (7). To further

investigate the liver's metabolic response, all dogs were anesthetized with infusions still

ongoing. At the end of the PM clamp (480 min) hepatic tissue was rapidly excised, flash-frozen

in liquid nitrogen, and stored at -80°C for subsequent molecular analyses, following which the

animals were euthanized.

Analyses

Biochemical and Molecular Methods

To evaluate hormone and substrate flux across the liver, whole blood samples were

collected and analyzed thereby permitting the use of arteriovenous balance techniques. Plasma

glucose levels were measured in quadruplicate with an Analox GM9 glucose analyzer

throughout the experiment. The concentrations of insulin (PI-12K, MilliporeSigma, Burlington,

MA), glucagon (GL-32K, MilliporeSigma), and cortisol (VUMC Analytical Services Core in-house

primary antibody with I^{125} cortisol from MP Biomedicals, Santa Ana, CA) were quantified using

radioimmunoassay (18). Key metabolic intermediates, including lactate, glycerol, alanine, and

non-esterified fatty acids (NEFA), were assessed via enzymatic spectrophotometric assays. To

measure the specific activity of [3-^{3}H]-glucose, plasma samples were deproteinized using the

Somogyi method, followed by liquid scintillation counting (19; 20). The terminal hepatic glycogen

content was quantified using the Keppler and Decker amyloglucosidase method (21).

At the end of the PM clamp, hepatic tissue was collected for qPCR analysis, which has

previously been optimized for canine-specific targets using validated primers (18). Gene

expression levels were normalized to GAPDH. Additionally, baseline liver samples from five

dogs that were fasted overnight were included for comparison.

Calculations

Arterio-venous difference calculations were used to determine net hepatic glucose

balance (NHGB) using the following equation: NHGB = [HGL$_{out}$] - [HGL$_{in}$]. Glucose output from

the liver (HGL$_{out}$) was calculated as [HGL$_{out}$] = [BF$_h$ x G$_h$], while glucose coming into the liver

was calculated as [HGL$_{in}$] = [(BF$_a$ × G$_a$) + (BF$_p$ × G$_p$)]. In these equations, G represents the

concentration of blood glucose, BF indicates measured blood flow, and A, P, and H correspond

to the hepatic artery, hepatic portal vein, and hepatic vein, respectively. Negative values of

NHGB indicate that hepatic glucose uptake exceeds glucose production, suggesting net glucose

uptake. This approach was also applied to assess net hepatic balance for other substrates and

hormones. To calculate hepatic fractional glucose extraction, NHGB was divided by HGLin.

Unidirectional hepatic glucose uptake (HGU) was determined by multiplying HGLin by the

fractional glucose extraction of [3-^{3}H]-glucose. Non-HGU, which primarily represents glucose

uptake by skeletal muscle, was calculated by subtracting tracer-derived HGU from the glucose

infusion rate (GIR) and adjusting for changes in glucose mass over time. Direct hepatic

glycogen synthesis was estimated by dividing the total radiolabeled glycogen stored in the liver

by the specific activity of the glucose precursor pool. This was measured in disintegrations per

minute (dpm) of labeled glucose incorporated into glycogen. The specific activity of the glucose

precursor pool was determined by the ratio of labeled to unlabeled glucose (dpm/mg) in the

inflowing blood, considering contributions from both the hepatic artery and portal vein. To

calculate net hepatic carbon retention (NHCR), which reflects total glycogen synthesis over

time, the equation NHCR = NHGU + NHLU + 2(NHAU) + NHGlyU − HGO was used. Here,

NHLU, NHAU, and NHGlyU represent net hepatic uptake of lactate, alanine, and glycerol,

respectively, and HGO refers to hepatic glucose oxidation. NHAU was doubled to account for

net hepatic uptake of all amino acids, as alanine accounts for approximately 50% of

gluconeogenic amino acid precursor uptake (22; 23). Previous studies have shown that hepatic

glucose oxidation is minimal, approximately 0.2 mg/kg/min under hyperinsulinemic-

hyperglycemic clamp conditions, thus contributing little to overall hepatic glucose flux (24; 25).

Glycolytic flux was estimated by adding HGO and the net hepatic output of lactate, alanine, and

glycerol. Finally, plasma concentrations of insulin and glucagon in the hepatic sinusoidal space

were derived based on the relative contributions of the hepatic artery and portal vein, as

previously established.

Statistics

Results are expressed as mean ± SEM. Group differences and temporal changes in flux

analyses were evaluated using a two-way repeated-measures ANOVA, followed by Tukey's

post hoc test for multiple comparisons. Areas under the curve were compared using an

unpaired two-tailed t-test, while molecular outcomes were analyzed via one-way ANOVA with

Tukey's post hoc correction. Data normality was confirmed before statistical testing. A threshold

of $P < 0.05$ was used to define statistical significance. All analyses were performed using

GraphPad Prism software.

Results

AM Clamp Glucose and Hormone Data

Plasma glucose levels were clamped at euglycemia (105-115 mg/dL) in both groups

during the morning clamp **(Fig. 2A)**. Arterial plasma insulin was maintained at basal levels in the

No Prime group (mean of 6 ± 1 μU/mL), whereas in the Prime group, insulin levels were

elevated as intended (mean of 32 ± 3 μU/mL) **(Fig. 2B, C).** This pattern was reflected in the

hepatic sinusoidal plasma insulin levels, with insulin levels in the Prime group being nearly 6-

fold of that in the No Prime group **(Fig. 2D, E, Table 1).** Arterial and hepatic sinusoidal plasma

glucagon levels were matched between the two groups **(Table 1).** The exogenous glucose

delivered to adequately clamp plasma glucose levels was significantly greater in the Prime

group compared to the No Prime group, as expected **(Fig. 2F, G, Table 1).** In summary, the

Prime group received elevated amounts of insulin (and subsequently glucose to maintain

euglycemia) in the hepatic sinusoids and arterial circulation during the AM clamp, representative

of a morning insulin prime, whereas the No Prime group received basal levels of insulin,

representative of an extended fast.

PM Clamp Glucose, Hormone, and Metabolite Data

Both groups were subject to identical afternoon conditions, which consisted of clamping

glucose levels at hyperglycemia (230-240 mg/dL, **Fig. 3A, B)**, activating the portal glucose

signal by delivering some glucose into the hepatic portal vein to establish a negative arterial-to-

portal glucose gradient (**Fig. 3C**) and maintaining arterial and hepatic sinusoidal plasma insulin

at basal levels **(Table 1, Fig. 3D).** Plasma glucagon levels were replaced at basal levels **(Table**

1). Despite having matched parameters during the PM clamp, the Prime group required 37%

more glucose to maintain hyperglycemia (mean GIR of 7.0 ± 0.4 vs. 4.6 ± 0.4 mg/kg/min in

Prime vs. No Prime, respectively, **Fig. 4A, B**). This increase in the GIR in the Prime group was

not attributed to non-hepatic glucose uptake, which was not different between the two groups

(mean of 4.5 ± 0.3 mg/kg/min in Prime vs. mean of 4.0 ± 0.3 mg/kg/min in No Prime, **Fig. 4C,**

D). Rather, it was due to a substantial increase in PM net hepatic glucose uptake in the Prime

group (mean of 2.2 ± 0.3 vs. 0.1 ± 0.3 mg/kg/min in Prime vs. No Prime, respectively, **Fig. 4E,**

F). The enhancement in net hepatic glucose uptake in the Prime group compared to the No

Prime group was attributed to an increase in net glycolytic flux **(Fig. 4G)** and net glycogenesis

(Fig. 4H). Thus, these data show that a morning insulin prime has lasting effects on afternoon

hepatic glucose handling, resulting in increased afternoon net hepatic glucose uptake despite

the presence of basal insulin during the PM clamp.

254   When comparing plasma non-esterified fatty acid (NEFA) levels, arterial NEFA levels

255 were significantly greater in the No Prime group vs. the Prime group, although net hepatic NEFA

256 uptake was similar between the two groups **(Fig. 5A, B)**. It is important to note that arterial

257 NEFA levels were not at the same level at the onset of the PM clamp, likely due to the lingering

258 effects of the morning insulin prime on suppressing lipolysis in the Prime group. Additionally,

259 arterial blood glycerol and net hepatic glycerol uptake were not significantly different between

260 groups **(Fig. 5C, D)** suggesting that, despite the elevated arterial NEFA levels in the No Prime

261 group, hepatic processing of NEFAs and glycerol was not impaired. This may indicate that the

262 excess circulating NEFAs were diverted toward re-esterification in peripheral tissues rather than

263 being taken up and oxidized by the liver, although this pathway likely contributes only minimally.

264 As for alanine, while arterial blood levels were not different between groups, the Prime group

265 had reduced net hepatic alanine uptake compared to the No Prime group **(Fig. 5E, F).** Lastly,

266 arterial blood lactate levels were similar between groups, but the Prime group had 5-fold greater

267 net hepatic lactate output than the No Prime group, which is consistent with the Prime group

268 exhibiting net glycolysis **(Fig. 5G, H).**

270 ***Glycogen Data and Terminal Hepatic Tissue Analyses***

271   The amount of tritiated glucose incorporated into hepatic glycogen stores during the PM

272 clamp, denoted as direct glycogen synthesis, was comparable between groups **(Fig. 6A)**.

273 However, terminal hepatic glycogen was greater in the Prime group compared to the No Prime

274 group (mean of 43 ± 3 vs. 36 ± 1 mg glycogen/g liver in Prime vs. No Prime, respectively, **Fig.**

275 **6B)**. Upon assessing hepatic gene expression at the end of the PM clamp, compared to

276 baseline, transcript levels of phosphoenolpyruvate carboxykinase 1 (PCK1; **Fig. 6C**), glucose-6-

277 phosphatase catalytic subunit 1 (G6PC1, **Fig. 6D**), and liver-specific glycogen phosphorylase

278 (PYGL) mRNA **(Fig. 6E)** were reduced similarly between the two groups. On the other hand,

279 glucokinase (GCK, **Fig. 6F**), sterol regulatory element binding protein-1c (SREBP-1c, **Fig. 6G**),

and fatty acids synthase (FAS) gene expression (**Fig. 6H)** at the end of the PM clamp were

significantly elevated in the Prime group compared to the No Prime group. Acetyl-coA

carboxylase (ACC) mRNA was not significantly different between groups (data not shown).

These data further support the notion that morning insulin exposure has a lasting effect on the

transcriptional landscape of the liver, allowing for enhanced PM glucose handling.

Discussion

The second meal phenomenon, characterized by an improved glycemic response to a

second, identical meal, has recently gained increasing attention due to its potential therapeutic

implication in managing postprandial glucose levels, particularly in individuals with insulin

resistance and type 2 diabetes (3; 5; 26-30). Our previous research established that a morning

elevation in insulin primes the liver for enhanced HGU in response to a subsequent glucose

load (17; 31). However, the precise mechanisms underlying this improvement during PM

hepatic glucose handling have remained unclear. In the present study, we identify the key

mechanisms responsible for the augmented HGU observed in the afternoon following AM

insulin priming, uncovering both insulin-dependent and insulin-independent components that

contribute to enhanced PM glucose metabolism. Specifically, we demonstrate that AM

hyperinsulinemia amplifies not only insulin action in the afternoon, but also glucose

effectiveness and/or the portal glucose signal, leading to a coordinated increase in net HGU.

These findings highlight the complex interplay between these signals in modulating postprandial

HGU and, for the first time, provide novel mechanistic insight into how a morning insulin prime

elicits its effects on the liver to improve glucose handling with subsequent glucose loads.

To assess the impact of AM insulin priming on PM net hepatic glucose uptake, we

previously performed a series of control experiments in which dogs underwent a PM HIHG

clamp following either morning basal insulin or elevated insulin delivery **(Fig. 7A).** The priming

effect of AM insulin was quantified as the difference in PM NHGU between these two conditions,

which was 3.8 mg/kg/min. In the current study, dogs underwent a PM hyperglycemic clamp with

basal insulin to isolate the effects of glucose (including both glucose effectiveness and the

activation of the portal glucose signal) from the effects of insulin **(Fig. 7B).** Consistent with prior

findings, the AM insulin prime significantly enhanced PM NHGU, with the difference between

the No Prime and Prime groups, amounting to 2.1 mg/kg/min. This represents the contribution

of glucose effectiveness and the portal glucose signal to the priming effect, independent of

elevated PM insulin levels. Comparing the findings from both studies, AM insulin priming

enhanced PM NHGU by 3.8 mg/kg/min when PM insulin was elevated versus by 2.1 mg/kg/min

when insulin remained at basal levels, indicating that approximately 45% of the priming effect

was attributable to increased PM insulin action **(Fig. 7C).** Insulin-primed animals exhibited

significantly greater PM net HGU, indicating increased hepatic sensitivity to glucose itself. Thus,

AM insulin priming had a similar impact on PM insulin action and glucose action, suggesting that

priming influences cellular mechanisms common to both insulin and glucose signaling. These

findings provide strong evidence that prior insulin exposure causes the liver to be conditioned to

respond more effectively to both insulin and glucose in the PM, optimizing subsequent hepatic

glucose disposal.

To understand how AM insulin priming enhances hepatic responsiveness to glucose

independently of insulin, we examined hepatic intermediary metabolism under basal PM insulin

conditions. Beyond improved PM net HGU, AM insulin priming altered downstream metabolism,

with primed animals showing elevated PM net glycolytic flux and higher terminal hepatic

glycogen content, despite similar rates of direct glycogen synthesis from tritiated glucose. This

suggests enhanced glucose cycling in the Prime group, likely via increased indirect glycogen

synthesis or accelerated glucose-6-phosphate turnover, facilitating more efficient glucose

storage. These results demonstrate that prior insulin exposure metabolically conditioned the

liver to utilize glucose more effectively in the PM without requiring elevated insulin. This flexible

metabolic phenotype is further supported by a fivefold increase in net hepatic lactate output in

the Prime group, consistent with increased glycolytic throughput and possibly enhanced

pyruvate cycling. While elevated lactate in the Prime group may reflect glycolytic flux surpassing

immediate oxidative demands, the concurrent rises in glycogen content and glucokinase mRNA

expression suggest coordinated activation of multiple glucose disposal pathways during the PM

clamp.

To explore the molecular basis for the enhancement of HGU following AM insulin

priming, we assessed hepatic gene expression at the end of the PM clamp. Compared to

animals that received basal insulin in the morning (No Prime), those that received an AM insulin

prime (Prime) exhibited elevated expression of GCK, SREBP-1c, and FAS, consistent with

increased activation of insulin-responsive and lipogenic transcriptional programs (32; 33).

Interestingly, ACC expression did not differ between groups, suggesting a selective induction of

SREBP-1c targets or potential post-transcriptional regulation limiting ACC expression despite

transcriptional priming (34). Meanwhile, gluconeogenic and glycogenolytic genes such as PCK1

and PYGL were similarly suppressed in both groups when compared to basal, indicating that

the presence of elevated glucose alone is sufficient to downregulate these genes and that this

effect is independent of prior insulin exposure. These findings support the concept that AM

insulin primes the liver by selectively enhancing glucose uptake and lipid synthesis pathways

while suppressing opposing catabolic processes. This coordinated shift in gene expression

establishes a permissive transcriptional environment that poises the liver to more effectively

dispose of glucose during the postprandial period.

In addition to changes in glucose metabolism, the PM clamp data revealed important

differences in hepatic and systemic substrate handling that further characterize the metabolic

adaptations of enhanced PM glucose action induced by AM insulin priming. During the PM

clamp, arterial NEFA levels were significantly elevated in the No Prime group compared to the

Prime group, yet net hepatic NEFA uptake remained similar between groups. This dissociation

indicates that hepatic fatty acid metabolism was not a primary factor driving the enhanced

glucose disposal observed in the primed liver. Similarly, glycerol fluxes were comparable

between groups, suggesting hepatic lipid processing was largely unaffected. Notably, the No

Prime group exhibited increased net hepatic alanine uptake despite similar arterial alanine

concentrations, potentially reflecting a compensatory increase in amino acid catabolism due to

impaired glucose handling, although gluconeogenesis was not different between groups. These

substrate flux patterns reinforce the concept that AM insulin priming shifts hepatic metabolism to

preferentially utilize glucose, promoting more efficient energy partitioning and metabolic

flexibility during postprandial nutrient influx.

Collectively, these findings support the concept of hepatic metabolic memory, whereby

early hormonal cues elicit lasting molecular and functional adaptations that sensitize the liver to

subsequent nutrient challenges. This memory likely involves coordinated transcriptional

reprogramming, enzyme activity modulation, substrate availability shifts, and potentially

epigenetic modifications (35). Understanding the cellular and molecular underpinnings of this

memory will be critical to harnessing it therapeutically. Clinically, these results have important

implications for individuals with type 2 diabetes or impaired early-phase insulin secretion, where

inadequate morning insulin surges may impair hepatic priming and exacerbate postprandial

hyperglycemia throughout the day (36). Therapeutic strategies aimed at restoring or mimicking

hepatic insulin priming, whether via hepato-preferential insulin delivery, incretin-based agents,

or meal timing optimization, could enhance hepatic glucose handling. Furthermore, insulin

priming may interact with hepatic circadian rhythms, as the liver's metabolic programs are

regulated by an endogenous clock anticipating feeding-fasting cycles (37). Disruption of this

synchrony, common in shift work or metabolic syndrome, may blunt the priming effect and

worsen glucose intolerance (38-41). Future research should explore the crosstalk between

insulin signaling and circadian regulators to develop interventions that restore metabolic

homeostasis.

In conclusion, a physiological rise in morning insulin initiates a coordinated hepatic

program that enhances glucose uptake, utilization, and storage later in the day. This priming

effect engages both insulin-dependent and insulin-independent mechanisms, including hepatic

glucose effectiveness and the portal glucose signal. Collectively, the data demonstrate that

morning hyperinsulinemia equally enhances afternoon insulin and glucose action, suggesting

that insulin priming likely targets shared cellular pathways. These findings provide a mechanistic

basis for the second meal phenomenon and highlight the liver as a dynamic integrator of prior

hormonal cues that shape its metabolic responsiveness. By revealing how early insulin

exposure primes hepatic glucose metabolism, this work offers new opportunities to improve

glycemic control in metabolic disease.

Acknowledgments

Author Contributions. H.W., D.S.E., and A.D.C. participated in the design of experiments;

H.W. directed all experiments and collected data; B.F., K.Y., and T.H. participated in the

experiments; B.F. and K.Y. were responsible for surgical preparation and oversight of animal

care; H.W. and M.S.S. carried out biochemical and tissue analysis; H.W., D.S.E., and A.D.C.

interpreted experimental results; H.W. prepared figures and drafted the manuscript; H.W.,

D.S.E., G.K., and A.D.C. edited and revised the manuscript; All authors approved the final

version of the manuscript.

Guarantor Statement. A.D.C. is the guarantor of this work and, as such, has full access to all

the data in the study and takes responsibility for the integrity of the data and the accuracy of the

data analysis.

Conflict of Interest. A.D.C. has research contracts with Alnylam, Abvance, Fluidics, and Novo

Nordisk, as well as grants from the NIH and the Helmsley Charitable Trust. A.D.C. is a

consultant to Novo Nordisk, Paratus, Portal Insulin, Fractyl, and Thetis. No other individuals

have conflicts of interest relevant to this article.

Funding. This work was supported by National Institutes of Health (NIH) grant R01DK131082.

H.W. was supported by NIH grant 1F31DK142297 and by the Vanderbilt University Training

Program in Molecular Endocrinology NIH grant 5T32DK007563. Hormone analysis was

completed by Vanderbilt's Hormone Assay and Analytical Services Core, supported by NIH

grants DK020593 and DK059637. Vanderbilt's Large Animal Core provided surgical expertise,

supported by the NIH Grant DK020593.

Prior Presentation. This work was orally presented at the American Diabetes Association 2025

meeting. The abstract can be viewed at doi.org/10.2337/db25-362-OR.

References

1. Basu A, Dalla Man C, Basu R, Toffolo G, Cobelli C, Rizza RA: Effects of type 2

diabetes on insulin secretion, insulin action, glucose effectiveness, and postprandial

glucose metabolism. Diabetes Care 2009;32:866-872

2. Taylor R: Insulin resistance and type 2 diabetes. Diabetes 2012;61:778-779

3. Jovanovic A, Gerrard J, Taylor R: The second-meal phenomenon in type 2 diabetes.

Diabetes Care 2009;32:1199-1201

4. Bonuccelli S, Muscelli E, Gastaldelli A, Barsotti E, Astiarraga BD, Holst JJ, Mari A,

Ferrannini E: Improved tolerance to sequential glucose loading (Staub-Traugott effect):

size and mechanisms. Am J Physiol Endocrinol Metab 2009;297:E532-537

5. Wajngot A, Grill V, Efendić S, Cerasi E: The Staub-Traugott effect. Evidence for

multifactorial regulation of a physiological function. Scand J Clin Lab Invest

1982;42:307-313

6. Moore MC, Smith MS, Farmer B, Kraft G, Shiota M, Williams PE, Cherrington AD:

Priming Effect of a Morning Meal on Hepatic Glucose Disposition Later in the Day.

Diabetes 2017;66:1136-1145

7. Moore MC, Coate KC, Winnick JJ, An Z, Cherrington AD: Regulation of hepatic

glucose uptake and storage in vivo. Adv Nutr 2012;3:286-294

8. Cherrington AD: Banting Lecture 1997. Control of glucose uptake and release by the

liver in vivo. Diabetes 1999;48:1198-1214

9. Rahman MS, Hossain KS, Das S, Kundu S, Adegoke EO, Rahman MA, Hannan MA,

Uddin MJ, Pang MG: Role of Insulin in Health and Disease: An Update. Int J Mol Sci

2021;22

10. Najjar SM, Perdomo G: Hepatic Insulin Clearance: Mechanism and Physiology.

Physiology 2019;34:198-215

11. Dube S, Errazuriz-Cruzat I, Basu A, Basu R: The forgotten role of glucose

effectiveness in the regulation of glucose tolerance. Curr Diab Rep 2015;15:605

12. Bergman RN: Lilly lecture 1989. Toward physiological understanding of glucose

tolerance. Minimal-model approach. Diabetes 1989;38:1512-1527

13. Bergman RN, Ider YZ, Bowden CR, Cobelli C: Quantitative estimation of insulin

sensitivity. Am J Physiol 1979;236:E667-677

14. Galassetti P, Chu CA, Neal DW, Reed GW, Wasserman DH, Cherrington AD: A

negative arterial-portal venous glucose gradient increases net hepatic glucose uptake in

euglycemic dogs. Am J Physiol 1999;277:E126-134

15. Hsieh PS, Moore MC, Neal DW, Cherrington AD: Importance of the hepatic arterial

glucose level in generation of the portal signal in conscious dogs. Am J Physiol

Endocrinol Metab 2000;279:E284-292

16. Myers SR, McGuinness OP, Neal DW, Cherrington AD: Intraportal glucose delivery

alters the relationship between net hepatic glucose uptake and the insulin

concentration. J Clin Invest 1991;87:930-939

17. Moore MC, Smith MS, Farmer B, Coate KC, Kraft G, Shiota M, Williams PE,

Cherrington AD: Morning Hyperinsulinemia Primes the Liver for Glucose Uptake and

Glycogen Storage Later in the Day. Diabetes 2018;67:1237-1245

18. Waterman HL, Moore MC, Smith M, Farmer B, Scott M, Edgerton DS, Cherrington

AD: Duration of Morning Hyperinsulinemia is Key to the Enhancement of Hepatic

Glucose Uptake and Glycogen Storage Later in the Day. bioRxiv

2024:2024.2005.2010.593551

19. Sindelar DK, Balcom JH, Chu CA, Neal DW, Cherrington AD: A Comparison of the

Effects of Selective Increases in Peripheral or Portal insulin on Hepatic Glucose

Production in the Conscious Dog. Diabetes 1996;45:1594-1604

20. Nelson N: A PHOTOMETRIC ADAPTATION OF THE SOMOGYI METHOD FOR

THE DETERMINATION OF GLUCOSE. Journal of Biological Chemistry 1944;153:375-

380

21. Keppler D, Decker K: Methods of Enzymatic Analysis, Edited by Bergmeyer, HU.

Academic Press, New York and London, 1974

22. Shah AM, Wondisford FE: Tracking the carbons supplying gluconeogenesis. J Biol

Chem 2020;295:14419-14429

23. Edgerton DS, Cardin S, Emshwiller M, Neal D, Chandramouli V, Schumann WC,

Landau BR, Rossetti L, Cherrington AD: Small Increases in Insulin Inhibit Hepatic

Glucose Production Solely Caused by an Effect on Glycogen Metabolism. Diabetes

2001;50:1872-1882

24. Moore MC, Satake S, Lautz M, Soleimanpour SA, Neal DW, Smith M, Cherrington

AD: Nonesterified fatty acids and hepatic glucose metabolism in the conscious dog.

Diabetes 2004;53:32-40

25. Satake S, Moore MC, Igawa K, Converse M, Farmer B, Neal DW, Cherrington AD:

Direct and indirect effects of insulin on glucose uptake and storage by the liver.

Diabetes 2002;51:1663-1671

26. Abraira C, Buchanan B, Hodges L: Modification of glycogen deposition by priming

glucose loads: the second-meal phenomenon. Am J Clin Nutr 1987;45:952-957

27. Chen MJ, Jovanovic A, Taylor R: Utilizing the second-meal effect in type 2 diabetes:

practical use of a soya-yogurt snack. Diabetes Care 2010;33:2552-2554

28. Clark CA, Gardiner J, McBurney MI, Anderson S, Weatherspoon LJ, Henry DN,

Hord NG: Effects of breakfast meal composition on second meal metabolic responses in

adults with Type 2 diabetes mellitus. Eur J Clin Nutr 2006;60:1122-1129

29. Jenkins DJ, Wolever TM, Taylor RH, Griffiths C, Krzeminska K, Lawrie JA, Bennett

CM, Goff DV, Sarson DL, Bloom SR: Slow release dietary carbohydrate improves

second meal tolerance. Am J Clin Nutr 1982;35:1339-1346

30. Lee SH, Tura A, Mari A, Ko SH, Kwon HS, Song KH, Yoon KH, Lee KW, Ahn YB:

Potentiation of the early-phase insulin response by a prior meal contributes to the

second-meal phenomenon in type 2 diabetes. Am J Physiol Endocrinol Metab

2011;301:E984-990

31. Waterman HL, Moore MC, Smith MS, Farmer B, Yankey K, Scott M, Edgerton DS,

Cherrington AD: Improved Afternoon Hepatic Glucose Disposal and Storage Requires

Morning Engagement of Hepatic Insulin Receptors. Diabetes 2024;

32. Foretz M, Guichard C, Ferré P, Foufelle F: Sterol regulatory element binding

protein-1c is a major mediator of insulin action on the hepatic expression of glucokinase

and lipogenesis-related genes. Proc Natl Acad Sci U S A 1999;96:12737-12742

33. Griffin MJ, Sul HS: Insulin regulation of fatty acid synthase gene transcription: roles

of USF and SREBP-1c. IUBMB Life 2004;56:595-600

34. Dentin R, Pégorier JP, Benhamed F, Foufelle F, Ferré P, Fauveau V, Magnuson

MA, Girard J, Postic C: Hepatic glucokinase is required for the synergistic action of

ChREBP and SREBP-1c on glycolytic and lipogenic gene expression. J Biol Chem

2004;279:20314-20326

35. Chen SS, Otero YF, Mulligan KX, Lundblad TM, Williams PE, McGuinness OP:

Liver, but not muscle, has an entrainable metabolic memory. PLoS One 2014;9:e86164

36. Di Giuseppe G, Ciccarelli G, Soldovieri L, Capece U, Cefalo CMA, Moffa S, Nista

EC, Brunetti M, Cinti F, Gasbarrini A, Pontecorvi A, Giaccari A, Mezza T: First-phase

insulin secretion: can its evaluation direct therapeutic approaches? Trends Endocrinol

Metab 2023;34:216-230

37. Fougeray T, Polizzi A, Régnier M, Fougerat A, Ellero-Simatos S, Lippi Y, Smati S,

Lasserre F, Tramunt B, Huillet M, Dopavogui L, Salvi J, Nédélec E, Gigot V, Smith L,

Naylies C, Sommer C, Haas JT, Wahli W, Duez H, Gourdy P, Gamet-Payrastre L,

Benani A, Burnol AF, Loiseau N, Postic C, Montagner A, Guillou H: The hepatocyte

insulin receptor is required to program the liver clock and rhythmic gene expression.

Cell Rep 2022;39:110674

38. Honma K, Hikosaka M, Mochizuki K, Goda T: Loss of circadian rhythm of circulating

insulin concentration induced by high-fat diet intake is associated with disrupted

rhythmic expression of circadian clock genes in the liver. Metabolism 2016;65:482-491

39. Mason IC, Qian J, Adler GK, Scheer F: Impact of circadian disruption on glucose

metabolism: implications for type 2 diabetes. Diabetologia 2020;63:462-472

40. Zimmet P, Alberti K, Stern N, Bilu C, El-Osta A, Einat H, Kronfeld-Schor N: The

Circadian Syndrome: is the Metabolic Syndrome and much more! J Intern Med

2019;286:181-191

41. Boivin DB, Boudreau P, Kosmadopoulos A: Disturbance of the Circadian System in

Shift Work and Its Health Impact. J Biol Rhythms 2022;37:3-28

Table 1 – Mean hormone and substrate clamp data

	Experimental Group	
AM Clamp Parameter (mean 30-240 min)	No Prime (n=8)	Prime (n=8)
Hepatic Sinusoidal Plasma Insulin (µU/mL)	18 ± 1	106 ± 6
Glucose Infusion Rate (mg/kg/min)	0.4 ± 0.3	8.8 ± 0.8
Arterial Plasma Glucagon (pg/mL)	22 ± 2	21 ± 2
Hepatic Sinusoidal Plasma Glucagon (pg/mL)	29 ± 2	29 ± 2
PM Clamp Parameter (mean 360-480 min)		
Arterial Plasma Insulin (µU/mL)	6 ± 1	6 ± 1
Hepatic Sinusoidal Plasma Insulin (µU/mL)	20 ± 1	24 ± 2
Arterial Plasma Glucagon (pg/mL)	21 ± 1	20 ± 1
Hepatic Sinusoidal Plasma Glucagon (pg/mL)	29 ± 1	27 ± 2

Figure Legends

1. **Figure 1: Experimental protocol.** Canines (*n*=8/group) underwent a 4h euglycemic clamp in the morning (0 to 240 min) with either basal insulin (No Prime) or hyperinsulinemia (Prime). Near the end of the AM clamp (180 min), tracer infusion began to allow enough time for equilibration before the start of the PM clamp. Following the AM clamp, there was a 1.5h rest period (240-330 min) where all infusions were halted. The dogs then underwent a 2.5h PM hyperglycemic clamp with portal glucose delivery and basal insulin (330 to 480 min). Tissue collection occurred at the end of the PM clamp. IV; intravenous infusion.

2. **Figure 2: Morning (AM) clamp glucose and insulin flux data.** Arterial plasma glucose (A) and insulin (B), hepatic sinusoidal plasma insulin (D), and the amount of exogenous glucose required to maintain euglycemia (F) are shown for the No Prime and Prime groups; *n*=8/group. Bar graphs indicate the area under the curve (AUC) for arterial insulin concentrations (C), hepatic sinusoidal insulin concentrations (E), and glucose infusion rates (G) during the 4h AM period. Data are expressed as mean ± SEM. ****$P<0.0001$ between groups. ns = non-significant.

3. **Figure 3: Afternoon (PM) clamp glucose and insulin data.** A vertical line at 330 min separates the resting period from the onset of the PM clamp. Arterial plasma glucose (A), hepatic glucose load (B), the difference between the artery and the portal vein plasma glucose levels (C), and hepatic sinusoidal plasma insulin (D) are shown for the 2.5h euinsulinemic-hyperglycemic PM clamp period in the No Prime and Prime groups; *n*=8/group. There were no differences between the two groups for any of the parameters measured. Data are expressed as mean ± SEM. ns = non-significant.

4. **Figure 4: Glucose uptake and storage during the PM euinsulinemic-hyperglycemic**

clamp. A vertical line at 330 min separates the resting period from the onset of the PM

clamp. Glucose infusion rate (A), non-hepatic glucose uptake (C), net hepatic glucose

uptake (E), net glycolytic/gluconeogenic flux (G), and net glycogenesis/glycogenolysis

flux (H) are shown over time for the No Prime and Prime groups, n=8/group. The

respective AUCs for the glucose infusion rate, non-hepatic glucose uptake, and net

hepatic glucose uptake are shown for both groups (B, D, F). Data are expressed as

mean $\pm$ SEM. *P<0.05, **P<0.01, ***P<0.001 between groups. ns = non-significant.

5. **Figure 5: PM clamp fatty acid and metabolite flux data.** A vertical line at 330 min

separates the resting period from the onset of the PM clamp. Arterial plasma non-

esterified fatty acids (NEFA) (A), net hepatic NEFA uptake (B), arterial blood glycerol

(C), net hepatic glycerol uptake (D), arterial blood alanine (E), net hepatic alanine uptake

(F), arterial blood lactate (G), and net hepatic lactate output (H) are shown for the PM

euinsulinemic-hyperglycemic clamp for the No Prime and Prime groups, n=8/group. Data

are expressed as mean $\pm$ SEM. **P<0.01 between groups. ns = non-significant.

6. **Figure 6: Glycogen and terminal hepatic tissue gene expression data.** Tracer-

determined direct glycogen synthesis during the PM clamp (A), terminal hepatic

glycogen reserves (B), phosphoenolpyruvate carboxykinase 1 (PCK1) mRNA (C),

glucose-6-phosphatase catalytic subunit 1 (G6PC1) mRNA (D), and liver-specific

glycogen phosphorylase (PYGL) mRNA (E), glucokinase (GCK) mRNA (F), sterol

regulatory element-binding protein 1c (SREBP-1c) mRNA (G), and fatty acid synthase

(FAS) mRNA (H) levels are shown for the No Prime and Prime groups, n=8/group, as

well as basal tissue when applicable (n=5). Data are expressed as mean $\pm$ SEM. *P<0.5,

**P<0.01, ***P<0.001, ****P<0.0001 between groups. ns = non-significant.

7. **Figure 7: Summary of PM net hepatic glucose uptake (NHGU) data. A)** Mean PM

NHGU data from control experiments previously completed. One of these groups (grey

bar) received basal insulin in the AM with a PM HIHG clamp. The other (blue bar)

received hyperinsulinemia in the AM with the same PM HIHG clamp. The size of the

priming effect of AM insulin on PM NHGU is the difference in PM NHGU between these

two groups, which is 3.8 mg/kg/min (shown in red). **B)** Mean PM NHGU data from the

current experiments. The No Prime group (orange) received basal insulin in the AM with

a PM hyperglycemic clamp with basal insulin delivery. The Prime group (green) received

AM hyperinsulinemia with the same euinsulinemic-hyperglycemic clamp in the PM. The

only difference between these two groups and the control groups presented in (A) is that

they underwent a PM hyperglycemic clamp with **basal** insulin rather than having an

elevation in insulin. The impact of the AM insulin prime on the PM effects of glucose

(glucose effectiveness (GE) and the portal glucose signal (PGS) is equal to the

difference in PM NHGU between the No Prime and Prime groups, which is equal to 2.1

mg/kg/min (shown in purple). **C)** Of the 3.8 mg/kg/min enhancement in PM NHGU

following an AM insulin prime, 45% of that (1.7 mg/kg/min) was due to an enhancement

in PM insulin action, whereas the other 55% (2.1 mg/kg/min) was due to the

enhancement in PM glucose action.

AM CLAMP INSULIN RATES (mU/kg/min):

No Prime (Basal insulin, *n=8*):
0-240 min: 0.25

Prime (Hyperinsulinemia, *n=8*):
0-30 min: 2.1
30-60 min: 2.4
60-240 min: 1.5

Figure 1: Experimental protocol. Canines (*n*=8/group) underwent a 4h euglycemic clamp in the morning (0 to 240 min) with either basal insulin (No Prime) or hyperinsulinemia (Prime). Near the end of the AM clamp (180 min), tracer infusion began to allow enough time for equilibration before the start of the PM clamp. Following the AM clamp, there was a 1.5h rest period (240-330 min) where all infusions were halted. The dogs then underwent a 2.5h PM hyperglycemic clamp with portal glucose delivery and **basal** insulin (330 to 480 min). Tissue collection occurred at the end of the PM clamp. IV; intravenous infusion.

Figure 2: Morning (AM) clamp glucose and insulin flux data. Arterial plasma glucose (A) and insulin (B), hepatic sinusoidal plasma insulin (D), and the amount of exogenous glucose required to maintain euglycemia (F) are shown for the No Prime and Prime groups; *n*=8/group. Bar graphs indicate the area under the curve (AUC) for arterial insulin concentrations (C), hepatic sinusoidal insulin concentrations (E), and glucose infusion rates (G) during the 4h AM period. Data are expressed as mean ± SEM. ****P<0.0001 between groups. ns = non-significant.

Figure 3: Afternoon (PM) clamp glucose and insulin data. A vertical line at 330 min separates the resting period from the onset of the PM clamp. Arterial plasma glucose (A), hepatic glucose load (B), the difference between the artery and the portal vein plasma glucose levels (C), and hepatic sinusoidal plasma insulin (D) are shown for the 2.5h euinsulinemic-hyperglycemic PM clamp period in the No Prime and Prime groups; *n*=8/group. There were no differences between the two groups for any of the parameters measured. Data are expressed as mean ± SEM. ns = non-significant.

Figure 4: Glucose uptake and storage during the PM euinsulinemic-hyperglycemic clamp. A vertical line at 330 min separates the resting period from the onset of the PM clamp. Glucose infusion rate (A), non-hepatic glucose uptake (C), net hepatic glucose uptake (E), net glycolytic/gluconeogenic flux (G), and net glycogenesis/glycogenolysis flux (H) are shown over time for the No Prime and Prime groups, *n*=8/group. The respective AUCs for the glucose infusion rate, non-hepatic glucose uptake, and net hepatic glucose uptake are shown for both groups (B, D, F). Data are expressed as mean ± SEM. *P<0.05, **P<0.01, ***P<0.001 between groups. ns = non-significant.

Figure 5: PM clamp fatty acid and metabolite flux data. A vertical line at 330 min separates the resting period from the onset of the PM clamp. Arterial plasma non-esterified fatty acids (NEFA) (A), net hepatic NEFA uptake (B), arterial blood glycerol (C), net hepatic glycerol uptake (D), arterial blood alanine (E), net hepatic alanine uptake (F), arterial blood lactate (G), and net hepatic lactate output (H) are shown for the PM euinsulinemic-hyperglycemic clamp for the No Prime and Prime groups, *n*=8/group. Data are expressed as mean ± SEM. **P<0.01 between groups. ns = non-significant.

Figure 6: Glycogen and terminal hepatic tissue gene expression data. Tracer-determined direct glycogen synthesis during the PM clamp (A), terminal hepatic glycogen reserves (B), phosphoenolpyruvate carboxykinase 1 (PCK1) mRNA (C), glucose-6-phosphatase catalytic subunit 1 (G6PC1) mRNA (D), and liver-specific glycogen phosphorylase (PYGL) mRNA (E), glucokinase (GCK) mRNA (F), sterol regulatory element-binding protein 1c (SREBP-1c) mRNA (G), and fatty acid synthase (FAS) mRNA (H) levels are shown for the No Prime and Prime groups, *n*=8/group, as well as basal tissue when applicable (*n*=5). Data are expressed as mean ± SEM. *P<0.5, **P<0.01, ***P<0.001, ****P<0.0001 between groups. ns = non-significant.

Figure 7: Summary of PM net hepatic glucose uptake (NHGU) data. A) Mean PM NHGU data from control experiments previously completed. One of these groups (grey bar) received basal insulin in the AM with a PM HIHG clamp. The other (blue bar) received hyperinsulinemia in the AM with the same PM HIHG clamp. The size of the priming effect of AM insulin on PM NHGU is the difference in PM NHGU between these two groups, which is 3.8 mg/kg/min (shown in red). **B)** Mean PM NHGU data from the current experiments. The No Prime group (orange) received basal insulin in the AM with a PM hyperglycemic clamp with basal insulin delivery. The Prime group (green) received AM hyperinsulinemia with the same euinsulinemic-hyperglycemic clamp in the PM. The only difference between these two groups and the control groups presented in (A) is that they underwent a PM hyperglycemic clamp with **basal** insulin rather than having an elevation in insulin. The impact of the AM insulin prime on the PM effects of glucose (glucose effectiveness (GE) and the portal glucose signal (PGS)) is equal to the difference in PM NHGU between the No Prime and Prime groups, which is equal to 2.1 mg/kg/min (shown in purple). **C)** Of the 3.8 mg/kg/min enhancement in PM NHGU following an AM insulin prime, 45% of that (1.7 mg/kg/min) was due to an enhancement in PM insulin action, whereas the other 55% (2.1 mg/kg/min) was due to the enhancement in PM glucose action.