Morning Elevation in Insulin Enhances Afternoon Hepatic Postprandial Glucose Disposal in the Dog by Increasing Both Insulin Signaling and Glucose Action

Hannah L. Waterman[1], Marta S. Smith[1], Ben Farmer[1], Kalisha Yankey[1], Tristan Howard[1], Guillaume Kraft[1], Alan D. Cherrington[1], Dale S. Edgerton[1]

[1]Department of Molecular Physiology & Biophysics, Vanderbilt University School of Medicine

Corresponding author information:

Name: Hannah Waterman

Current Affiliation: Vanderbilt University School of Medicine

Phone: (360)-536-1498

Email: Hannah.L.Waterman@Vanderbilt.Edu

Running title: AM Insulin Enhances PM Glucose Action and Insulin Action

Keywords: hepatic glucose uptake, glucose metabolism, liver metabolism, glycogen, glucagon, insulin, meal composition, breakfast, lunch, second meal phenomenon, second meal effect, Staub-Traugott effect, insulin action, glucose effectiveness, postprandial, insulin signaling, glucose

Abstract

The second-meal phenomenon refers to the improved glycemic response to a

subsequent identical meal. Postprandial net hepatic glucose uptake (NHGU) is

governed by the combined effects of three regulatory factors: insulin action (IA), initiated

by hyperinsulinemia; glucose effectiveness (GE), driven by hyperglycemia; and the

portal glucose signal (PGS), activated by glucose delivery into the hepatoportal

circulation. Previous studies demonstrated that morning (AM) hyperinsulinemia primes

the liver, causing substantially enhanced NHGU later in the day; however, it remained

unclear which component of the afternoon (PM) response is augmented. To address

this, we assessed how AM insulin elevation influences PM IA, GE, and the PGS. Dogs

underwent an AM clamp with either a 4h hyperinsulinemic prime (Prime, *n=8*) or basal

insulin delivery (No Prime, *n=8*). After a 1.5-hour rest, both groups underwent a PM

hyperglycemic clamp (with portal glucose delivery) under basal insulin conditions.

During the PM clamp, NHGU was significantly greater in the Prime versus No Prime

group (2.2 ± 0.3 vs. 0.1 ± 0.3 mg/kg/min, $P<0.005$), indicating priming enhanced GE

and/or PGS effects. In prior experiments with all three stimuli present in the PM (IA, GE,

and PGS), AM insulin priming increased PM NHGU by 3.8 mg/kg/min. Thus, while AM

insulin priming alone enhanced GE and/or PGS, the full effect requires an elevation in

PM insulin, suggesting that morning insulin exposure primes the liver by augmenting

both afternoon insulin action and glucose action.

Introduction

Effective postprandial glucose disposal is essential for maintaining metabolic

health, and impairments in this process are a defining feature of insulin resistance and

type 2 diabetes (1-3). An incompletely understood aspect of glucose metabolism is the

second meal phenomenon, in which the glycemic response to a second, identical meal

is attenuated compared with its initial response to the meal (4-6), suggesting that prior

metabolic or hormonal cues influence later glucose handling. Consistent with this

concept, we previously demonstrated in dogs that a rise in morning (AM) insulin

enhanced net hepatic glucose uptake (NHGU) during an afternoon (PM)

hyperinsulinemic-hyperglycemic clamp (7; 8). These findings indicate that early insulin

exposure can prime the liver for more efficient glucose disposal later in the day, exerting

effects that persist beyond the period of elevated circulating insulin.

Hepatic glucose uptake, storage, and release strongly influence postprandial

glucose excursions, highlighting the liver's central role in glucose homeostasis (9).

Notably, under physiologic conditions the liver can dispose of as much postprandial

glucose as muscle (10). NHGU is regulated by the integrated actions of three primary

signals: insulin action (IA), triggered by hyperinsulinemia; glucose effectiveness (GE),

driven by hyperglycemia; and the portal glucose signal (PGS), activated by elevated

glucose concentrations in the hepatoportal circulation, such as occurs with absorption of

glucose from the gut (11; 12). Insulin promotes NHGU by suppressing endogenous

glucose production and stimulating glycogen synthesis through well-characterized

signaling pathways (13; 14). GE reflects the ability of glucose per se to suppress

hepatic glucose output and enhance NHGU in the presence of basal insulin (15-17).

The PGS arises from neural sensing of glucose delivery through the portal vein,

enabling the liver to anticipate nutrient influx and adjust glucose uptake and storage

accordingly, via mechanisms that are less well defined (11; 18-20). Together, these

factors enable the liver to function as an active integrator of postprandial glucose

signals (12).

Despite evidence that prior insulin exposure enhances NHGU later in the day, it

remains unclear which afternoon regulatory signals (IA, GE, and/or the PGS) are

augmented by morning insulin exposure. This distinction is particularly relevant because

both insulin action and glucose action are impaired in insulin-resistant and diabetic

states, contributing to defective postprandial glucose disposal and exaggerated

glycemic excursions (21; 22). Determining whether AM insulin exposure enhances the

PM hepatic response to glucose, insulin, or their combined actions is therefore essential

for defining how early hormonal cues regulate hepatic glucose metabolism and for

identifying mechanisms that may be leveraged to improve postprandial glucose control

in metabolic disease. To address this gap, we assessed how an elevation in morning

insulin influences the effects of glucose (GE and the PGS) vs. insulin (IA) during a PM

euinsulinemic-hyperglycemic clamp.

Research Design and Methods

Animal care and surgical procedures

16 adult mongrel dogs (8 males, 8 females) with an average body weight of 27.6

± 1.0 kg were used in these experiments. The animals were obtained from a USDA-

approved vendor and housed according to the standards set by the American

Association for the Accreditation of Laboratory Animal Care. All procedures were

reviewed and approved by the Vanderbilt Institutional Animal Care and Use Committee.

Two weeks before the experimental procedures, the animals underwent a laparotomy

under general anesthesia for the surgical placement of blood flow probes around the

hepatic artery and hepatic portal vein. Additionally, catheters were implanted for blood

sampling in the hepatic vein, hepatic portal vein, and femoral artery. Infusion catheters

were positioned in the inferior vena cava and the splenic and jejunal veins, which supply

the hepatic portal vein. These catheters were secured subcutaneously until the day of

the experiment, when they were exteriorized under local anesthesia. The dogs were fed

a controlled diet consisting of chow and meat, with a macronutrient composition of 46%

carbohydrates, 34% protein, 14.5% fat, and 5.5% fiber. Before the experiment, the

animals underwent an 18-hour fasting period. Assessments were conducted to ensure

the dogs were healthy before the experiment, requiring that the dogs had consumed at

least 75% of their most recent meal, maintained a leukocyte count below 18,000/mm³,

and had a hematocrit level above 34%. Blood collection was limited to no more than

20% of the animal's total blood volume over the course of a day.

Experimental Design

The experimental protocol lasted 8 hours and included two glucose clamp

phases, separated by a rest period. These clamps were designed to simulate

postprandial glucose dynamics at times corresponding to morning and midday meals.

The first phase, referred to as the AM clamp, occurred from 0 to 240 min, followed by a

90 min break before the PM clamp, which took place from 330 to 480 min **(Fig. 1).**

Blood samples were collected every 15 to 30 minutes from the femoral artery, hepatic

portal vein, and the left major hepatic vein to measure hormone and metabolite levels.

Arterial plasma glucose was assessed every five minutes, and exogenous glucose

infusion into a peripheral vein was adjusted as needed to maintain target glucose

concentrations.

Morning (AM) Clamp Period (0-240 min) and Non-Clamp Period (240 – 330 min)

To suppress endogenous pancreatic insulin and glucagon secretion,

somatostatin (Bachem, Torrance, CA) was infused into the inferior vena cava (0.8

µg/kg/min) at the onset of the AM clamp. Glucagon (GlucaGen, Boehringer Ingelheim,

Ridgefield, CT) was replaced intraportally to maintain basal levels (0.57 ng/kg/min). One

of the experimental groups, Prime, received morning insulin (Novolin R; Novo Nordisk,

Basværd, Denmark) infusion designed to replicate the physiologic insulin secretion

pattern previously observed following a morning duodenal glucose infusion (insulin was

replaced intraportally at 2.1 mU/kg/min from 0-30 min, 2.4 mU/kg/min from 30-60 min,

and 1.5 mU/kg/min from 60-240 min) (7). The other group, No Prime, received

intraportal basal insulin replacement in the morning (0.25 mU/kg/min). Prior research

from our lab has shown that the enhancement of HGU in the afternoon is driven by an

increase in AM insulin levels, rather than a rise in morning glucose levels (7). Therefore,

arterial plasma glucose was clamped at euglycemia (~110 mg/dL) for the entirety of the

AM clamp. To allow for tracer-determined direct glycogen synthesis measurements in

the afternoon, a peripheral infusion of [3-³H] glucose (Revvitty, Waltham, MA) was

initiated at 180 min in both groups, with a priming dose of 38 µCi, followed by a

continuous infusion at 0.38 µCi/min. After the AM clamp (240 min), all infusions, except

for the tracer, were discontinued for 1.5h (non-clamp period). To characterize the

hepatic metabolic state prior to the initiation of the PM clamp, blood samples were

obtained from the femoral artery, portal vein, and hepatic vein at 300, 315, and 330

minutes **(Fig. 1).**

Afternoon (PM) Clamp Period (330 – 480 min)

Both the Prime and No Prime groups underwent an identical 2.5-h hyperglycemic

clamp in the PM with insulin and glucagon kept basal **(Fig. 1).** This protocol was

designed to isolate the effects of the morning insulin prime on PM insulin action vs.

glucose and the portal glucose signal. Thus, basal insulin levels were maintained

throughout the PM clamp. To achieve this, somatostatin was continuously infused via a

peripheral vein (0.8 µg/kg/min), while basal glucagon (0.57 ng/kg/min) and basal insulin

(0.25 mU/kg/min) were delivered intraportally. To stimulate the portal glucose signal,

glucose was directly infused into the hepatic portal vein at a rate of 4 mg/kg/min.

Additionally, a primed, continuous glucose infusion into a peripheral vein was

administered to rapidly elevate plasma glucose levels to ~200 mg/dL, maintaining

hyperglycemia for the duration of the PM clamp. These conditions were intended to

mimic the postprandial metabolic state seen following carbohydrate ingestion (11). To

further investigate the liver's metabolic response, all dogs were anesthetized with

infusions ongoing at the end of the PM clamp (480 min). Hepatic tissue was then rapidly

excised, flash-frozen in liquid nitrogen, and stored at -80°C for subsequent molecular

analyses. The animals were then euthanized.

Analyses

Biochemical and Molecular Methods

Whole blood samples were collected to assess hormone and substrate flux

across the liver using arteriovenous balance techniques (23). Plasma glucose levels

were measured in quadruplicate throughout the experiment using an Analox GM9

glucose analyzer. Plasma insulin (#PI-12K, MilliporeSigma, Burlington, MA), glucagon

(#GL-32K, MilliporeSigma), and cortisol (VUMC Analytical Services in-house primary

antibody with I125 cortisol from MP Biomedicals, Santa Ana, CA) were quantified by

radioimmunoassay (23). Glucagon values were background corrected to account for

non-specific binding, ensuring that measured concentrations accurately reflected

circulating hormone levels.

Key metabolic intermediates, including lactate, glycerol, alanine, and non-

esterified fatty acids (NEFA), were measured using enzymatic spectrophotometric

assays (23). The specific activity of [3-^{3}H]-glucose in plasma samples was determined

following deproteinization using the Somogyi method, with quantification by liquid

scintillation counting (24). Terminal hepatic glycogen content was measured using the

Keppler and Decker amyloglucosidase method (25).

At the end of the PM clamp, hepatic tissue was collected for molecular analyses,

including qPCR, Western blotting, and enzymatic activity assays. A group of overnight-

fasted dogs were used as basal controls, as applicable (*n=5*). Gene expression was

normalized to GAPDH, and protein levels were standardized to total protein using

Ponceau S staining (23). All assays were optimized for canine-specific targets using

validated primers and antibodies (23). Glucokinase activity was measured using an

assay adapted from Shiota et al. (26). Detailed protocols, including reagents, primers,

antibodies, sample preparation, and calculations, are provided in the supplemental

materials.

Calculations

Arterio-venous difference calculations were used to determine net hepatic

glucose balance (NHGB) using the following equation: $NHGB = [HGL_{out}] - [HGL_{in}]$.

Glucose output from the liver (HGL_{out}) was calculated as $[HGL_{out}] = [BF_h \times G_h]$, while

glucose coming into the liver was calculated as $[HGL_{in}] = [(BF_a \times G_a) + (BF_p \times G_p)]$. In

these equations, G represents the concentration of blood glucose, BF indicates

measured blood flow, and A, P, and H correspond to the hepatic artery, hepatic portal

vein, and hepatic vein, respectively. Measurements obtained in plasma were converted

to whole blood values using a conversion factor of 0.75, as previously described (23).

Negative values of NHGB indicate net glucose uptake, meaning that hepatic glucose

uptake exceeds glucose production. This approach was also used to assess net hepatic

balance for other substrates and hormones. To calculate hepatic fractional glucose

extraction, NHGB was divided by HGLin. Unidirectional hepatic glucose uptake (HGU)

was determined by multiplying HGLin by the fractional glucose extraction of [3-³H]-

glucose. Non-HGU, which primarily represents glucose uptake by skeletal muscle, was

calculated by subtracting tracer-derived HGU from the glucose infusion rate (GIR) and

correcting for changes in glucose mass over time. Direct hepatic glycogen synthesis

was estimated by dividing the total radiolabeled glycogen stored in the liver by the

specific activity of the glucose precursor pool during the PM clamp. This was measured

in disintegrations per minute (dpm) of labeled glucose incorporated into glycogen from

330-480 min. The specific activity of the glucose precursor pool was determined by the

ratio of labeled to unlabeled glucose (dpm/mg) in the inflowing blood, weighted by

contributions from both the hepatic artery and portal vein. To calculate net hepatic

carbon retention (NHCR), which reflects total glycogen synthesis over time, the

equation NHCR = NHGU + NHLU + 2(NHAU) + NHGlyU − HGO was used. Here,

NHLU, NHAU, and NHGlyU represent net hepatic uptake of lactate, alanine, and

glycerol, respectively, and HGO refers to hepatic glucose oxidation. NHAU was doubled

to account for net hepatic uptake of all amino acids, as alanine accounts for

approximately 50% of gluconeogenic amino acid precursor uptake (27; 28). Previous

studies have shown that hepatic glucose oxidation is minimal, approximately 0.2

mg/kg/min under the hyperinsulinemic-hyperglycemic clamp conditions seen in these

dogs, contributing little to overall hepatic glucose flux (29; 30). Glycolytic flux was

estimated by adding HGO to net hepatic lactate output. Finally, plasma concentrations

of insulin and glucagon in the hepatic sinusoidal space were derived based on the

relative contributions of the hepatic artery and portal vein to hepatic blood flow, following

established methodology.

Statistics

Results are expressed as mean ± SEM. Group differences and temporal

changes in flux analyses were evaluated using a two-way repeated-measures ANOVA,

followed by Tukey's post hoc test for multiple comparisons. Areas under the curve were

compared using an unpaired two-tailed t-test, and molecular outcomes were analyzed

using a one-way ANOVA with Tukey's post hoc correction. Normality of all datasets was

confirmed using the Shapiro-Wilk test, and all data were found to follow a normal

distribution. A threshold of $P < 0.05$ was used to define statistical significance. All

analyses were performed using GraphPad Prism software.

Results

AM Clamp Glucose and Hormone Data

During the AM clamp, plasma glucose was maintained at euglycemia (105-115

mg/dL) in both groups **(Fig. 2A).** Arterial plasma insulin was maintained at basal levels

in the No Prime group (6 ± 1 μU/mL), whereas insulin was elevated in the Prime group

according to the experimental design (32 ± 3 μU/mL; **Fig. 2B, C**). As a result, hepatic

sinusoidal insulin concentrations were approximately sixfold higher in the Prime group

than in the No Prime group (**Fig. 2D, E; Table 1**). Arterial and hepatic sinusoidal

glucagon concentrations remained basal and did not differ between groups **(Table 1).**

Consistent with the higher insulin exposure, the Prime group required significantly more

exogenous glucose to maintain euglycemia during the AM clamp than the No Prime

group **(Fig. 2F, G; Table 1).** Thus, the AM clamp successfully established sustained

hyperinsulinemia in the Prime group without altering glucose or glucagon levels.

PM Clamp Glucose, Hormone, and Metabolite Data

During the PM clamp, experimental conditions were identical between groups.

Arterial plasma glucose was clamped at hyperglycemia (230-240 mg/dL; **Fig. 3A, B**),

the portal glucose signal was activated by intraportal glucose delivery resulting in a

negative arterial-to-portal glucose gradient **(Fig. 3C),** and arterial and hepatic sinusoidal

insulin concentrations were maintained at basal levels **(Fig. 3D; Table 1).** Despite

matched PM clamp conditions, the Prime group required 52% more glucose to maintain

hyperglycemia than the No Prime group (glucose infusion rate of 7.0 ± 0.4 vs. 4.6 ± 0.4

mg/kg/min, respectively; **Fig. 4A, B).** This difference was not attributable to changes in

non-hepatic glucose uptake, which did not differ between groups **(Fig. 4C, D).** Instead,

the increased glucose requirement was due to a marked increase in net hepatic glucose

uptake (NHGU) in the Prime group (2.2 ± 0.3 vs. 0.1 ± 0.3 mg/kg/min in Prime vs. No

Prime; **Fig. 4E, F).** The enhancement in NHGU in the Prime group reflected both

increased net glycolytic flux **(Fig. 4G)** and increased net glycogenesis **(Fig. 4H).** Thus,

morning insulin priming resulted in a sustained amplification of PM hepatic glucose

uptake even if insulin did not rise during the PM clamp.

Arterial non-esterified fatty acid (NEFA) concentrations were significantly higher

in the No Prime group than in the Prime group during the PM clamp, whereas net

hepatic NEFA uptake did not differ between groups **(Fig. 5A, B).** Arterial glycerol

concentrations and net hepatic glycerol uptake were similar between groups **(Fig. 5C,**

D). Arterial alanine concentrations were also comparable; however, net hepatic alanine

uptake was reduced modestly in the Prime group relative to the No Prime group **(Fig.**

5E, F). Arterial lactate concentrations did not differ between groups, but net hepatic

lactate output was approximately fivefold greater in the Prime group than in the No

Prime group **(Fig. 5G, H),** consistent with increased hepatic glucose uptake and

glycolytic flux.

Glycogen Content and Hepatic Gene and Protein Analyses

Terminal hepatic glycogen content was significantly greater in the Prime group

compared with the No Prime group (43 ± 3 vs. 36 ± 1 mg glycogen/g liver; **Fig. 6A**).

Direct glycogen synthesis during the PM clamp, assessed by incorporation of tritiated

glucose into hepatic glycogen, tended to be greater in the Prime group but this change

did not reach statistical significance **(Fig. 6B).** Phosphorylated Akt relative to total Akt

was similar in both the Prime and No Prime groups compared with basal controls,

consistent with basal insulin conditions during the PM clamp **(Fig. 6C).** *Glucokinase*

(*GCK*) mRNA abundance was significantly greater in the Prime group than in the No

Prime and basal groups at the end of the PM clamp **(Fig. 6D);** however, GCK protein

abundance and enzymatic activity did not differ between groups **(Fig. 6E, F).** Glycogen

synthase protein was similarly dephosphorylated and activated in both experimental

groups relative to basal **(Fig. 6G),** while phosphorylated glycogen phosphorylase

protein did not differ among groups **(Fig. 6H).** Transcript levels of *phosphoenolpyruvate*

carboxykinase 1 (*PCK1*) and *glucose-6-phosphatase catalytic subunit 1* (*G6PC1*) were

similarly reduced in both Prime and No Prime groups compared with basal

(Supplemental Fig. 1A, 1B). In contrast, *sterol regulatory element-binding protein-1c*

(SREBP-1c) and *fatty acid synthase (FAS)* mRNA levels were significantly elevated in

the Prime group relative to the No Prime group at the end of the PM clamp

(Supplemental Fig. 1C, 1D).

Discussion

The second meal phenomenon, in which the glycemic response to a subsequent

identical meal is improved, reflects the body's capacity to integrate prior hormonal and

nutrient exposures to optimize later postprandial glucose handling (6; 31; 32). This

adaptation provides a potential therapeutic avenue which is relevant for disease states

where postprandial glucose disposal is reduced due to impaired insulin action and/or

glucose effectiveness, such as type 1 and 2 diabetes (4; 6; 32-35). Previous studies in

dogs demonstrated that it is the physiological rise in morning insulin which is the

component of the first meal that primes the liver for enhanced NHGU and glycogen

storage later in the day and that this effect requires activation of hepatic insulin

receptors during the morning period (7; 8). Building on this foundation, the present study

was designed to define which components of the afternoon hepatic response are

enhanced by morning insulin exposure. This was done by isolating glucose-mediated

regulation of NHGU (GE and the PGS), under basal insulin conditions in the PM,

thereby distinguishing these effects from those of insulin action (11).

To address this question, we used a reductive experimental approach to identify

which regulatory signals are enhanced by a morning insulin prime in controlling

afternoon NHGU. Specifically, we compared the effect of morning insulin priming on

afternoon hepatic glucose metabolism during a hyperinsulinemic-hyperglycemic clamp

(36), either when all three regulatory factors were engaged, or when just GE and the

PGS were elevated under basal insulin conditions. When afternoon IA, GE, and the

PGS were all engaged, morning insulin priming increased afternoon NHGU by 3.8

mg/kg/min **(Fig. 7A).** In the present study, with matching hyperglycemia and portal

glucose delivery in the two groups, but in the presence of basal insulin, morning priming

increased NHGU by 2.1 mg/kg/min **(Fig. 7B).** This pattern suggests that a substantial

portion of the priming effect (about half) is mediated through enhanced responsiveness

to glucose-mediated signals, whereas the remaining component likely reflects

augmented afternoon insulin action (1.7 mg/kg/min) **(Fig. 7C).** However, this calculation

relies on cross-study comparisons and assumes additivity based on historical

experiments (37; 38). In reality, these pathways likely interact and may coordinate

synergistically within integrated hepatic metabolic networks. The key point is not the

precise quantitative partitioning of afternoon NHGU, but that morning insulin exposure

amplifies the liver's sensitivity to both hormonal and glucose-mediated cues later in the

day, enhancing postprandial glucose disposal. Thus, this study demonstrates that

morning insulin augments both the glucose-mediated (GE/the PGS) and insulin-

mediated components of afternoon NHGU.

Afternoon clamp conditions were carefully matched to control for confounding

influences. Plasma glucose, portal glucose delivery, and basal insulin and glucagon

replacement were tightly controlled in both groups with no statistical differences. Minor

differences included a slightly greater relative arterial-portal glucose gradient in the

Prime group (4 mg/dL) and a minimally higher hepatic sinusoidal insulin concentration in

the Prime group (4 µU/mL). These differences were small in magnitude, however, and

are unlikely to have meaningfully alter hepatic glucose metabolism.

Interestingly, primed animals showed significantly higher hepatic expression of

the lipogenic transcription factor *SREBP-1c*, a well-established insulin-responsive

transcription factor (39), and its downstream target, *FAS*, at the end of the PM

hyperglycemic clamp. Notably, this occurred under conditions in which the morning

clamp produced a transient rise in insulin, followed later by hyperglycemia during the

PM clamp, rather than the simultaneous high insulin and high glucose previously

reported as necessary for hepatic activation (39; 40). Since gene expression was

assessed only at the terminal time point, we cannot exclude the possibility that insulin

exposure during the morning clamp initiated or partially activated these transcriptional

programs. However, the persistence or amplification of *SREBP-1c* and *FAS* expression

into the afternoon hyperglycemic period suggests that prior insulin exposure establishes

a hepatic state that alters the transcriptional response to subsequent glucose

availability. These findings point to a potentially novel mechanism in which temporal

separation of hormonal and nutrient signals can coordinate carbohydrate and lipid

metabolism. The coordinated activation of *SREBP-1c* and *FAS*, along with enhanced

glycolytic flux and hepatic glycogen storage, reflects a hepatic memory of earlier insulin

exposure and a shift toward integrated carbohydrate and lipid metabolism. This primed

state provides a mechanistic context for the enhanced NHGU observed in primed

animals, linking early insulin exposure to subsequent regulation of both glucose and

lipid pathways. Despite this, circulating lipid availability did not parallel hepatic

transcriptional responses. Although arterial NEFA concentrations were higher in the No

Prime group, net hepatic NEFA uptake did not differ between groups, and afternoon

lipolysis was not suppressed in either condition.

The enhancement of NHGU within the first 30 minutes of the PM clamp further

indicates that the effects of morning insulin were already present at the time of glucose

exposure rather than developing progressively during the clamp. Molecular

measurements obtained at the terminal time point therefore provide mechanistic context

but cannot fully account for the dynamic flux changes observed during the clamp itself.

At the end of the PM clamp, primed animals exhibited elevated hepatic *GCK* mRNA,

whereas total GCK protein abundance, maximal GCK enzymatic activity, and the

phosphorylation states of glycogen synthase and glycogen phosphorylase were similar

between groups. The increase in *GCK* mRNA is consistent with insulin-dependent

transcription initiated in the morning and relatively slow mRNA turnover during the

afternoon hyperglycemic clamp and supports the concept that morning insulin

establishes a molecular program that enhances glucose phosphorylation capacity

during subsequent hyperglycemia (41-43).

Prior work from our group demonstrated that a physiological rise in morning

insulin results in increased hepatic *GCK* mRNA and protein, elevations that persist

through the intervening rest period (330 min) following the AM clamp (8). In the present

study, the absence of elevated GCK protein at the terminal time point (480 min), despite

a clear enhancement of NHGU in the Prime group (evident from 360-480 min), was

unexpected. However, this discrepancy does not preclude a central role for glucokinase

in mediating the priming effect. *In vivo*, GCK activity is regulated not only by total

enzyme abundance but also by dynamic processes such as subcellular

compartmentalization and protein-protein interactions. These include binding to

glucokinase regulatory protein in the nucleus (44), as well as the bifunctional enzyme

PFK-2/FBPase-2 (45) and the pro-apoptotic mitochondrial protein BAD (46). Through

these interactions and others, GCK can be sequestered or released in response to

glucose availability, thereby modulating its cytosolic activity depending on the specific

physiological context (47). Importantly, these complex regulatory mechanisms are not

captured by terminal protein measurements or *ex vivo* activity assays. It is plausible that

prior insulin exposure may sensitize these processes, although this remains to be

demonstrated and represents an important avenue for future investigation. Furthermore,

the rapid onset and sustained enhancement of NHGU argue against accelerated

degradation or loss of functional GCK during the afternoon, particularly given its long

hepatic half-life (48). Glucokinase remains the most plausible molecular mediator of

morning insulin priming, as it catalyzes the rate-limiting step of hepatic glucose uptake

under hyperglycemic conditions and uniquely integrates insulin-dependent transcription,

a long protein half-life, and rapid activation by glucose (42; 49-51). Other hepatic

proteins are less likely contributors, as they are not simultaneously insulin-inducible,

long-lived, and acutely responsive to glucose availability later in the day (50).

In summary, these studies demonstrate that a physiological rise in morning

insulin primes the liver to respond more effectively to a subsequent nutrient load, even

without a rise in afternoon insulin. This finding is notable because it suggests that

hepatic priming may confer metabolic benefit even in settings of limited or impaired

insulin availability, such as in individuals with type 1 diabetes or insulin resistance / type

2 diabetes. Whether this priming mechanism is preserved or altered in metabolically

dysfunctional states, however, remains to be determined. Under these conditions, the

priming enhances glucose-mediated regulatory mechanisms, but full second-meal

enhancement of NHGU, still requires a concurrent rise in afternoon insulin, indicating

that morning insulin also potentiates later insulin action. Glucokinase is the most

plausible molecular mediator, consistent with its insulin-dependent regulation and

central role in hepatic glucose phosphorylation. Together, these findings define a

hepatic mechanism by which early-day insulin exposure shapes later postprandial

glucose handling and suggest that strategically timed insulin or insulinotropic

interventions could improve postprandial glycemic control in insulin-resistant and

diabetic states by restoring defects in glucose effectiveness and insulin action.

Acknowledgments

Author Contributions. H.W., D.S.E., and A.D.C. participated in the design of

experiments; H.W. directed all experiments and collected data; B.F., K.Y., and T.H.

participated in the experiments; B.F. and K.Y. were responsible for surgical preparation

and oversight of animal care; H.W. and M.S.S. carried out biochemical and tissue

analysis; H.W., D.S.E., and A.D.C. interpreted experimental results; H.W. prepared

figures and drafted the manuscript; H.W., D.S.E., G.K., and A.D.C. edited and revised

the manuscript; All authors approved the final version of the manuscript.

Data availability. Values for all data points in graphs are reported in the Supporting

Data Values file. Additionally, these data and supplemental materials are publicly

available: doi.org/10.6084/m9.figshare.31049578. All other data generated and

analyzed during the current study are available from the corresponding author upon

reasonable request.

Guarantor Statement. A.D.C. is the guarantor of this work and, as such, has full

access to all the data in the study and takes responsibility for the integrity of the data

and the accuracy of the data analysis.

Conflict of Interest. A.D.C. has research contracts with Alnylam and Abvance, as well

as grants from the NIH. A.D.C. is a consultant to Novo Nordisk, Paratus, Portal Insulin,

Fractyl, and Thetis. No other individuals have conflicts of interest relevant to this article.

Funding. This work was supported by National Institutes of Health (NIH) grant

R01DK131082. H.W. was supported by NIH grant F31DK142297 and by the Vanderbilt

University Training Program in Molecular Endocrinology NIH grant T32DK007563.

Hormone analysis was completed by the Vanderbilt University Medical Center Analytical

Services Core, supported by the NIH Diabetes Research Center grant DK020593.

Vanderbilt's Large Animal Core provided surgical expertise, supported by the NIH Grant

DK020593.

Prior Presentation. This work was orally presented at the American Diabetes

Association 2025 meeting. The abstract can be viewed at doi.org/10.2337/db25-362-

OR.

451   References

1. Basu A, Dalla Man C, Basu R, Toffolo G, Cobelli C, Rizza RA: Effects of type 2

diabetes on insulin secretion, insulin action, glucose effectiveness, and postprandial

glucose metabolism. Diabetes Care 2009;32:866-872

2. Taylor R: Insulin resistance and type 2 diabetes. Diabetes 2012;61:778-779

3. Blaak EE, Antoine JM, Benton D, Björck I, Bozzetto L, Brouns F, Diamant M, Dye L,

Hulshof T, Holst JJ, Lamport DJ, Laville M, Lawton CL, Meheust A, Nilson A, Normand

S, Rivellese AA, Theis S, Torekov SS, Vinoy S: Impact of postprandial glycaemia on

health and prevention of disease. Obes Rev 2012;13:923-984

4. Jovanovic A, Gerrard J, Taylor R: The second-meal phenomenon in type 2 diabetes.

Diabetes Care 2009;32:1199-1201

5. Bonuccelli S, Muscelli E, Gastaldelli A, Barsotti E, Astiarraga BD, Holst JJ, Mari A,

Ferrannini E: Improved tolerance to sequential glucose loading (Staub-Traugott effect):

size and mechanisms. Am J Physiol Endocrinol Metab 2009;297:E532-537

6. Wajngot A, Grill V, Efendić S, Cerasi E: The Staub-Traugott effect. Evidence for

multifactorial regulation of a physiological function. Scand J Clin Lab Invest

1982;42:307-313

7. Moore MC, Smith MS, Farmer B, Coate KC, Kraft G, Shiota M, Williams PE,

Cherrington AD: Morning Hyperinsulinemia Primes the Liver for Glucose Uptake and

Glycogen Storage Later in the Day. Diabetes 2018;67:1237-1245

8. Waterman HL, Moore MC, Smith MS, Farmer B, Yankey K, Scott M, Edgerton DS,

Cherrington AD: Improved Afternoon Hepatic Glucose Disposal and Storage Requires

Morning Engagement of Hepatic Insulin Receptors. Diabetes 2024;

9. Moore MC, Smith MS, Farmer B, Kraft G, Shiota M, Williams PE, Cherrington AD:

Priming Effect of a Morning Meal on Hepatic Glucose Disposition Later in the Day.

Diabetes 2017;66:1136-1145

10. Edgerton DS, Scott M, Farmer B, Williams PE, Madsen P, Kjeldsen T, Brand CL,

Fledelius C, Nishimura E, Cherrington AD: Targeting insulin to the liver corrects defects

in glucose metabolism caused by peripheral insulin delivery. JCI Insight 2019;5

11. Moore MC, Coate KC, Winnick JJ, An Z, Cherrington AD: Regulation of hepatic

glucose uptake and storage in vivo. Adv Nutr 2012;3:286-294

12. Cherrington AD: Banting Lecture 1997. Control of glucose uptake and release by

the liver in vivo. Diabetes 1999;48:1198-1214

13. Rahman MS, Hossain KS, Das S, Kundu S, Adegoke EO, Rahman MA, Hannan

MA, Uddin MJ, Pang MG: Role of Insulin in Health and Disease: An Update. Int J Mol

Sci 2021;22

14. Najjar SM, Perdomo G: Hepatic Insulin Clearance: Mechanism and Physiology.

Physiology 2019;34:198-215

15. Dube S, Errazuriz-Cruzat I, Basu A, Basu R: The forgotten role of glucose

effectiveness in the regulation of glucose tolerance. Curr Diab Rep 2015;15:605

16. Bergman RN: Lilly lecture 1989. Toward physiological understanding of glucose

tolerance. Minimal-model approach. Diabetes 1989;38:1512-1527

17. Bergman RN, Ider YZ, Bowden CR, Cobelli C: Quantitative estimation of insulin

sensitivity. Am J Physiol 1979;236:E667-677

18. Galassetti P, Chu CA, Neal DW, Reed GW, Wasserman DH, Cherrington AD: A

negative arterial-portal venous glucose gradient increases net hepatic glucose uptake in

euglycemic dogs. Am J Physiol 1999;277:E126-134

19. Hsieh PS, Moore MC, Neal DW, Cherrington AD: Importance of the hepatic arterial

glucose level in generation of the portal signal in conscious dogs. Am J Physiol

Endocrinol Metab 2000;279:E284-292

20. Myers SR, McGuinness OP, Neal DW, Cherrington AD: Intraportal glucose delivery

alters the relationship between net hepatic glucose uptake and the insulin

concentration. J Clin Invest 1991;87:930-939

21. Ahrén B, Pacini G: Glucose effectiveness: Lessons from studies on insulin-

independent glucose clearance in mice. J Diabetes Investig 2021;12:675-685

22. Najjar SM, Caprio S, Gastaldelli A: Insulin Clearance in Health and Disease. Annu

Rev Physiol 2023;85:363-381

23. Waterman HL, Moore MC, Smith MS, Farmer B, Scott M, Edgerton DS, Cherrington

AD: Duration of morning hyperinsulinemia determines hepatic glucose uptake and

glycogen storage later in the day. Am J Physiol Endocrinol Metab 2024;327:E655-e667

24. Nelson N: A PHOTOMETRIC ADAPTATION OF THE SOMOGYI METHOD FOR

THE DETERMINATION OF GLUCOSE. Journal of Biological Chemistry 1944;153:375-

380

25. Keppler D, Decker K: Methods of Enzymatic Analysis, Edited by Bergmeyer, HU.

Academic Press, New York and London, 1974

26. Fujimoto Y, Torres TP, Donahue EP, Shiota M: Glucose Toxicity Is Responsible for

the Development of Impaired Regulation of Endogenous Glucose Production and

Hepatic Glucokinase in Zucker Diabetic Fatty Rats. Diabetes 2006;55:2479-2490

27. Shah AM, Wondisford FE: Tracking the carbons supplying gluconeogenesis. J Biol

Chem 2020;295:14419-14429

28. Edgerton DS, Cardin S, Emshwiller M, Neal D, Chandramouli V, Schumann WC,

Landau BR, Rossetti L, Cherrington AD: Small Increases in Insulin Inhibit Hepatic

Glucose Production Solely Caused by an Effect on Glycogen Metabolism. Diabetes

2001;50:1872-1882

29. Moore MC, Satake S, Lautz M, Soleimanpour SA, Neal DW, Smith M, Cherrington

AD: Nonesterified fatty acids and hepatic glucose metabolism in the conscious dog.

Diabetes 2004;53:32-40

30. Satake S, Moore MC, Igawa K, Converse M, Farmer B, Neal DW, Cherrington AD:

Direct and indirect effects of insulin on glucose uptake and storage by the liver.

Diabetes 2002;51:1663-1671

31. Abraira C, Buchanan B, Hodges L: Modification of glycogen deposition by priming

glucose loads: the second-meal phenomenon. Am J Clin Nutr 1987;45:952-957

32. Jenkins DJ, Wolever TM, Taylor RH, Griffiths C, Krzeminska K, Lawrie JA, Bennett

CM, Goff DV, Sarson DL, Bloom SR: Slow release dietary carbohydrate improves

second meal tolerance. Am J Clin Nutr 1982;35:1339-1346

33. Chen MJ, Jovanovic A, Taylor R: Utilizing the second-meal effect in type 2 diabetes:

practical use of a soya-yogurt snack. Diabetes Care 2010;33:2552-2554

34. Clark CA, Gardiner J, McBurney MI, Anderson S, Weatherspoon LJ, Henry DN,

Hord NG: Effects of breakfast meal composition on second meal metabolic responses in

adults with Type 2 diabetes mellitus. Eur J Clin Nutr 2006;60:1122-1129

35. Lee SH, Tura A, Mari A, Ko SH, Kwon HS, Song KH, Yoon KH, Lee KW, Ahn YB:

Potentiation of the early-phase insulin response by a prior meal contributes to the

second-meal phenomenon in type 2 diabetes. Am J Physiol Endocrinol Metab

2011;301:E984-990

36. H. Waterman MS, K. Yankey, B. Farmer, M. Schleh, K. Coate,, T. Howard DE, A.

Cherrington: Metabolic memory induced by morning insulin exposure enhances

liver responsiveness to a subsequent glucose load later in the day. Diabetologia

2025;68:S325

37. Pagliassotti MJ, et al.: Comparison of the Time Courses of Insulin and the Portal

Signal on Hepatic Glucose and Glycogen Metabolism in the Conscious Dog. 1996;

38. Pagliassotti MJ, Myers SR, Moore MC, Neal DW, Cherrington AD: Magnitude of

negative arterial-portal glucose gradient alters net hepatic glucose balance in conscious

dogs. Diabetes 1991;40:1659-1668

39. Foretz M, Guichard C, Ferré P, Foufelle F: Sterol regulatory element binding

protein-1c is a major mediator of insulin action on the hepatic expression of glucokinase

and lipogenesis-related genes. Proc Natl Acad Sci U S A 1999;96:12737-12742

40. Girard J, Ferré P, Foufelle F: Mechanisms by which carbohydrates regulate

expression of genes for glycolytic and lipogenic enzymes. Annu Rev Nutr 1997;17:325-

352

41. Iynedjian PB: Molecular physiology of mammalian glucokinase. Cell Mol Life Sci

2009;66:27-42

42. Iynedjian PB, Jotterand D, Nouspikel T, Asfari M, Pilot PR: Transcriptional induction

of glucokinase gene by insulin in cultured liver cells and its repression by the glucagon-

cAMP system. J Biol Chem 1989;264:21824-21829

43. Agius L: Glucokinase and molecular aspects of liver glycogen metabolism.

Biochemical Journal 2008;414:1-18

44. de la Iglesia N, Mukhtar M, Seoane J, Guinovart JJ, Agius L: The role of the

regulatory protein of glucokinase in the glucose sensory mechanism of the hepatocyte.

J Biol Chem 2000;275:10597-10603

45. Payne VA, Arden C, Wu C, Lange AJ, Agius L: Dual role of phosphofructokinase-

2/fructose bisphosphatase-2 in regulating the compartmentation and expression of

glucokinase in hepatocytes. Diabetes 2005;54:1949-1957

46. Danial NN, Gramm CF, Scorrano L, Zhang CY, Krauss S, Ranger AM, Datta SR,

Greenberg ME, Licklider LJ, Lowell BB, Gygi SP, Korsmeyer SJ: BAD and glucokinase

reside in a mitochondrial complex that integrates glycolysis and apoptosis. Nature

2003;424:952-956

47. Agius L: Targeting hepatic glucokinase in type 2 diabetes: weighing the benefits and

risks. Diabetes 2009;58:18-20

48. Niemeyer H, Ureta T, Clark-Turri L: Adaptive character of liver glucokinase. Mol Cell

Biochem 1975;6:109-126

49. Rui L: Energy metabolism in the liver. Compr Physiol 2014;4:177-197

50. Petersen MC, Vatner DF, Shulman GI: Regulation of hepatic glucose metabolism in

health and disease. Nat Rev Endocrinol 2017;13:572-587

51. Bossard P, Decaux JF, Juanes M, Girard J: Initial expression of glucokinase gene in

cultured hepatocytes from suckling rats is linked to the synthesis of an insulin-

dependent protein. Eur J Biochem 1994;223:371-380

Table 1 – Mean hormone and substrate clamp data

	Experimental Group	
AM Clamp Parameter (mean 30-240 min)	**No Prime (n=8)**	**Prime (n=8)**
Hepatic Sinusoidal Plasma Insulin (µU/mL)	18 ± 1	106 ± 6****
Glucose Infusion Rate (mg/kg/min)	0.4 ± 0.3	8.8 ± 0.8****
Arterial Plasma Glucagon (pg/mL)	22 ± 2	21 ± 2
Hepatic Sinusoidal Plasma Glucagon (pg/mL)	29 ± 2	29 ± 2
PM Clamp Parameter (mean 360-480 min)		
Arterial Plasma Insulin (µU/mL)	6 ± 1	6 ± 1
Hepatic Sinusoidal Plasma Insulin (µU/mL)	20 ± 1	24 ± 2
Arterial Plasma Glucagon (pg/mL)	21 ± 1	20 ± 1
Hepatic Sinusoidal Plasma Glucagon (pg/mL)	29 ± 1	27 ± 2

Figure Legends

1. **Figure 1: Experimental protocol.** Canines (n=8/group) underwent a 4h euglycemic clamp in the morning (0 to 240 min) with either basal insulin (No Prime) or hyperinsulinemia (Prime). Near the end of the AM clamp (180 min), tracer infusion began to allow enough time for equilibration before the start of the PM clamp. Following the AM clamp, there was a 1.5h rest period (240-330 min) where all infusions were halted. The dogs then underwent a 2.5h PM hyperglycemic clamp with portal glucose delivery and basal insulin (330 to 480 min). Tissue collection occurred at the end of the PM clamp. IV; intravenous infusion.

2. **Figure 2: Morning (AM) clamp glucose and insulin flux data.** Arterial plasma glucose (A) and insulin (B), hepatic sinusoidal plasma insulin (D), and the amount of exogenous glucose required to maintain euglycemia (F) are shown for the No Prime and Prime groups; n=8/group. Bar graphs indicate the area under the curve (AUC) for arterial insulin concentrations (C), hepatic sinusoidal insulin concentrations (E), and glucose infusion rates (G) during the 4h AM period. Data are expressed as mean $\pm$ SEM. ****P<0.0001 between groups. ns = non-significant.

3. **Figure 3: Afternoon (PM) clamp glucose and insulin data.** A vertical line at 330 min separates the resting period from the onset of the PM clamp. Arterial

plasma glucose (A), hepatic glucose load (B), the difference between the artery

and the portal vein plasma glucose levels (C), and hepatic sinusoidal plasma

insulin (D) are shown for the 2.5h euinsulinemic-hyperglycemic PM clamp period

in the No Prime and Prime groups; n=8/group. There were no differences

between the two groups for any of the parameters measured. Data are

expressed as mean $\pm$ SEM. ns = non-significant.

4. Figure 4: Glucose uptake and storage during the PM euinsulinemic-

hyperglycemic clamp. A vertical line at 330 min separates the resting period

from the onset of the PM clamp. Glucose infusion rate (A), non-hepatic glucose

uptake (C), net hepatic glucose uptake (E), net glycolytic/gluconeogenic flux (G),

and net glycogenesis/glycogenolysis flux (H) are shown over time for the No

Prime and Prime groups, n=8/group. The respective delta AUCs for the glucose

infusion rate, non-hepatic glucose uptake, and net hepatic glucose uptake are

shown for both groups (B, D, F). Data are expressed as mean $\pm$ SEM. *P<0.05,

**P<0.01, ***P<0.001 between groups. ns = non-significant.

5. **Figure 5: PM clamp fatty acid and metabolite flux data.** A vertical line at 330

min separates the resting period from the onset of the PM clamp. Arterial plasma

non-esterified fatty acids (NEFA) (A), net hepatic NEFA uptake (B), arterial blood

glycerol (C), net hepatic glycerol uptake (D), arterial blood alanine (E), net

hepatic alanine uptake (F), arterial blood lactate (G), and net hepatic lactate

output (H) are shown for the PM euinsulinemic-hyperglycemic clamp for the No

Prime and Prime groups, n=8/group. Data are expressed as mean $\pm$ SEM.

**P<0.01 between groups. ns = non-significant.

6. Figure 6. Glycogen content and hepatic gene/protein expression at the end

of the PM clamp. Terminal hepatic glycogen content is shown in (A), and tracer-

determined direct glycogen synthesis during the PM clamp is shown in (B).

Protein and gene expression are shown as follows: Akt protein (phospho/total) in

(C), glucokinase (GCK) mRNA in (D), GCK protein (normalized to Ponceau S

stain) in (E), glucokinase (GCK) activity at 0.5 Vmax in (F; n = 5 or 6/group),

glycogen synthase (GS) protein (phospho/total) in (G), and glycogen

phosphorylase (GP) protein (phospho/total) in (H). Protein data were obtained by

western blot and normalized to Ponceau S stain; representative images are

provided in Supplemental Figure 1. Data are presented for the No Prime and

Prime groups (n = 8/group), with basal tissue values included where applicable (n

= 5). All values are expressed as mean ± SEM. *P < 0.05, **P < 0.01, ***P <

0.001, ****P < 0.0001 between groups; ns = not significant.

7. **Figure 7: Summary of PM net hepatic glucose uptake (NHGU) data.** Mean

PM NHGU data from control experiments previously completed are shown in (A).

One of these groups (grey bar) received basal insulin in the AM with a PM HIHG

clamp. The other (blue bar) received hyperinsulinemia in the AM with the same

PM HIHG clamp. The size of the priming effect of AM insulin on PM NHGU is the

difference in PM NHGU between these two groups, which is 3.8 mg/kg/min

(shown in red). Mean PM NHGU data from the current experiments are shown in

(B). The No Prime group (orange) received basal insulin in the AM with a PM

hyperglycemic clamp with basal insulin delivery. The Prime group (green)

received AM hyperinsulinemia with the same euinsulinemic-hyperglycemic clamp

in the PM. The only difference between these two groups and the control groups

presented in (A) is that they underwent a PM hyperglycemic clamp with basal

insulin rather than having an elevation in insulin. The impact of the AM insulin

prime on the PM effects of glucose (glucose effectiveness (GE) and the portal

glucose signal (PGS) is equal to the difference in PM NHGU between the No

Prime and Prime groups, which is equal to 2.1 mg/kg/min (shown in purple). Of

the 3.8 mg/kg/min enhancement in PM NHGU following an AM insulin prime,

45% of that (1.7 mg/kg/min) was due to an enhancement in PM insulin action,

whereas the other 55% (2.1 mg/kg/min) was due to the enhancement in PM

glucose action (C).

AM CLAMP INSULIN RATES (mU/kg/min):

**No Prime (Basal insulin, *n=8):*
0-240 min: 0.25

**Prime (Elevated insulin, *n=8):*
0-30 min: 2.1
30-60 min: 2.4
60-240 min: 1.5

Figure 1: Experimental protocol. Canines (*n*=8/group) underwent a 4h euglycemic clamp in the morning (0 to 240 min) with either basal insulin (No Prime) or hyperinsulinemia (Prime). Near the end of the AM clamp (180 min), tracer infusion began to allow enough time for equilibration before the start of the PM clamp. Following the AM clamp, there was a 1.5h rest period (240-330 min) where all infusions were halted. The dogs then underwent a 2.5h PM hyperglycemic clamp with portal glucose delivery and **basal** insulin (330 to 480 min). Tissue collection occurred at the end of the PM clamp. IV; intravenous infusion.

Figure 2: Morning (AM) clamp glucose and insulin flux data. Arterial plasma glucose (A) and insulin (B), hepatic sinusoidal plasma insulin (D), and the amount of exogenous glucose required to maintain euglycemia (F) are shown for the No Prime and Prime groups; *n*=8/group. Bar graphs indicate the area under the curve (AUC) for arterial insulin concentrations (C), hepatic sinusoidal insulin concentrations (E), and glucose infusion rates (G) during the 4h AM period. Data are expressed as mean ± SEM. ****P<0.0001 between groups. ns = non-significant.

Figure 3: Afternoon (PM) clamp glucose and insulin data. A vertical line at 330 min separates the resting period from the onset of the PM clamp. Arterial plasma glucose (A), hepatic glucose load (B), the difference between the artery and the portal vein plasma glucose levels (C), and hepatic sinusoidal plasma insulin (D) are shown for the 2.5h euinsulinemic-hyperglycemic PM clamp period in the No Prime and Prime groups; *n*=8/group. There were no differences between the two groups for any of the parameters measured. Data are expressed as mean ± SEM. ns = non-significant.

Figure 4: Glucose uptake and storage during the PM euinsulinemic-hyperglycemic clamp. A vertical line at 330 min separates the resting period from the onset of the PM clamp. Glucose infusion rate (A), non-hepatic glucose uptake (C), net hepatic glucose uptake (E), net glycolytic/gluconeogenic flux (G), and net glycogenesis/glycogenolysis flux (H) are shown over time for the No Prime and Prime groups, *n*=8/group. The respective delta AUCs for the glucose infusion rate, non-hepatic glucose uptake, and net hepatic glucose uptake are shown for both groups (B, D, F). Data are expressed as mean ± SEM. *P<0.05, **P<0.01, ***P<0.001 between groups. ns = non-significant.

Figure 5: PM clamp fatty acid and metabolite flux data. A vertical line at 330 min separates the resting period from the onset of the PM clamp. Arterial plasma non-esterified fatty acids (NEFA) (A), net hepatic NEFA uptake (B), arterial blood glycerol (C), net hepatic glycerol uptake (D), arterial blood alanine (E), net hepatic alanine uptake (F), arterial blood lactate (G), and net hepatic lactate output (H) are shown for the PM euinsulinemic-hyperglycemic clamp for the No Prime and Prime groups, *n*=8/group. Data are expressed as mean ± SEM. **P<0.01 between groups. ns = non-significant.

Figure 6: Glycogen content and hepatic gene/protein expression at the end of the PM clamp. Terminal hepatic glycogen content is shown in (A), and tracer-determined direct glycogen synthesis during the PM clamp is shown in (B). Protein and gene expression are shown as follows: Akt protein (phospho/total) in (C), glucokinase (GCK) mRNA in (D), GCK protein (normalized to Ponceau S stain) in (E), glucokinase (GCK) activity at 0.5 Vmax in (F; n = 5 or 6/group), glycogen synthase (GS) protein (phospho/total) in (G), and glycogen phosphorylase (GP) protein (phospho/total) in (H). Protein data were obtained by western blot and normalized to Ponceau S stain; representative images are provided in Supplemental Figure 1. Data are presented for the No Prime and Prime groups (n = 8/group), with basal tissue values included where applicable (n = 5). All values are expressed as mean ± SEM. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001 between groups; ns = not significant.

Figure 7: Summary of PM net hepatic glucose uptake (NHGU) data. Mean PM NHGU data from control experiments previously completed are shown in (A). One of these groups (grey bar) received basal insulin in the AM with a PM HIHG clamp. The other (blue bar) received hyperinsulinemia in the AM with the same PM HIHG clamp. The size of the priming effect of AM insulin on PM NHGU is the difference in PM NHGU between these two groups, which is 3.8 mg/kg/min (shown in red). Mean PM NHGU data from the current experiments are shown in (B). The No Prime group (orange) received basal insulin in the AM with a PM hyperglycemic clamp with basal insulin delivery. The Prime group (green) received AM hyperinsulinemia with the same euinsulinemic-hyperglycemic clamp in the PM. The only difference between these two groups and the control groups presented in (A) is that they underwent a PM hyperglycemic clamp with **basal** insulin rather than having an elevation in insulin. The impact of the AM insulin prime on the PM effects of glucose (glucose effectiveness (GE) and the portal glucose signal (PGS) is equal to the difference in PM NHGU between the No Prime and Prime groups, which is equal to 2.1 mg/kg/min (shown in purple). Of the 3.8 mg/kg/min enhancement in PM NHGU following an AM insulin prime, 45% of that (1.7 mg/kg/min) was due to an enhancement in PM insulin action, whereas the other 55% (2.1 mg/kg/min) was due to the enhancement in PM glucose action (C).