Functional connectivity, structural connectivity, and inter-individual variability in *Drosophila melanogaster*

Takuto Okuno[a,b*], Alexander Woodward[c], Hideyuki Okano[b,d], Junichi Hata[a,b]

[a] *Graduate School of Human Health Sciences, Tokyo Metropolitan University, 7-2-10 Higashiogu, Arakawa-ku, Tokyo, 116-8551, Japan*

[b] *Laboratory for Marmoset Models of Brain Diseases, RIKEN Center for Brain Science, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan*

[c] *Intelligent Vision Systems Lab, The University of Auckland, 34 Princes Street, Auckland Central, Auckland 1010, New Zealand*

[d] *Keio University Regenerative Medicine Research Center, Kawasaki 210-0821, Kanagawa, Japan.*

* Corresponding author

Abstract

Clarifying the relationship between structure and function is important for understanding the brain. In *Drosophila melanogaster*, whole-brain and hemibrain electron microscopy connectome data, as well as whole-brain calcium imaging data are available. We applied pre-processing methods from fMRI to whole-brain calcium imaging data and comprehensively investigated the optimal parameters. Then, we found that the FC-SC (functional and structural connectivity) correlation decreased linearly with ROI count, and this trend was the same in flies and humans. We also developed a new, more robust method to quantify the degree of pre- and post-synaptic segregation and investigated this in the fly whole-brain. This revealed that many neurons have unsegregated synapses. We extracted highly unsegregated synapses and compared them with random-extracted null SC matrices. Their FC-SC correlation was significantly higher, indicating that these synapses contribute to FC well. Conversely, highly segregated-synapses showed significantly lower FC-SC correlation and contribute less to FC. Therefore, neurons with unsegregated synapses like non-spiking neurons are spread throughout the whole brain, and they are thought to have a significant influence on FC.

Introduction (Main text – up to 5,000 words)

Functional connectivity (FC) describes the relationship between functional activity in one brain region

and that in another. In previous approaches, two spike trains were analyzed by a cross-correlation

histogram [1, 2]; then, this method was applied to positron emission tomography and functional magnetic

resonance imaging (fMRI) [3]. Subsequently, the time-series correlation between two regions has become

widely used as a measure of undirected FC [4]. Underlying such functional connectivity, a structural

connectivity (SC) substrate exists, and its relationship with FC has been investigated using various

modalities [5, 6, 7, 8, 9, 10, 11]. In humans, FC-SC correlations between resting-state (rs-) fMRI and

diffusion tensor imaging have been explored, and studies have shown high relationships of r=0.66 for 66

regions and r=0.54 for 998 regions [6, 7]. Other studies have examined the relationship between rs-fMRI

and SC based on retrograde tracing from 145 injection sites in the left cortex of marmosets (r=0.379 for

9862 voxels, Ext.Data.Fig.2-1c) [10] and the relationship between calcium imaging and SC based on

electron microscopy (EM) in the hemibrain of *Drosophila melanogaster (D. melanogaster)* [12] (r=0.74

for 37 regions) [9]. Because of the disparity in species, number of ROIs, and modalities, a direct

comparison across them becomes difficult. Therefore, in this study, we used whole-brain [13] and

hemibrain [12] EM connectome data, which are currently considered to be the most detailed, and whole-

brain calcium imaging data [14] of *D. melanogaster* to comprehensively investigate the relationship

between FC and SC across various ROI shapes and numbers. Our results of the FC-SC correlations in the

fly hemibrain compared to the human cortex at various ROIs showed roughly 1.08–1.24 times higher FC-

SC correlations in the fly (Fig. 2d), and no large divergence was observed between the two species.

Because of the strong relationship between FC and SC, they can be used to investigate inter-individual

variability in *D. melanogaster*. In particular, the FlyWire whole-brain and FlyEM hemibrain connectome

data have been compared [15, 16], and widely different results are observed with cell typing [16].

However, these studies have yet to clarify whether these differences in connectome between the two

individuals are attributed to inter-individual variability or some other reasons. Therefore, we compared

these two EM connectomes based on the averaged FC of eight flies. FlyEM was scanned at a resolution of

8×8×8 nm, and the connectome data has a "confidence" threshold that indicates the degree of synaptic

detection. Conversely, FlyWire was scanned at a resolution of 4×4×40 nm, and it had a "CleftScore" [17]

threshold that shows the degree of synaptic detection. In fact, the thresholds initially employed by

FlyWire and FlyEM caused large differences in post-synapse numbers; as such, we adjusted for the

thresholds and compared the EM connectome data in each fly. We found that FlyEM clearly had higher

values (more accuracy) than FlyWire in its FC-SC correlation and detection. In our judgment, this

difference was not caused by inter-individual variability; instead, it was attributed to differences in the

accuracy of detection and annotation of dendrites, axons, and synapses because of the voxel anisotropy.

We explored how functional activity contributed to the FC. In textbook terms, post-synapses on dendrites

receive neurotransmitters and conduct an action potential along nerve fibers if they are excitatory; and

pre-synapses on axons release neurotransmitters [18, 19]. In calcium imaging, the pre-synaptic activity in

each region is detected as calcium signals, which represent functional activity. Then the FC is detected as

the synchronization of functional activities. However, the synaptic arrangement on the neurons revealed

in the EM connectome shows pre- and post-synapses that are highly intermingled in many neurons (i.e.

input and output locations may be intermingled across axon and dendritic arbors); thus, signal

transmission may be complex [19, 15, 13, 20]. Nerve cells include non-spiking neurons such as visual

neurons, CT1 amacrine cells [21, 22, 23, 24, 25], interneurons in the antennal lobe [26], and APL cells in

the mushroom body [27, 28, 29]. In *D. melanogaster*. these cells show localized functional activity taking

place on a portion of the dendrite. In CT1 and APL cells, the pre- and post-synapses are highly

intermingled (Fig.4a, Ext.Data.Fig.4-2). Therefore, we wanted to investigate the presence of these

unsegregated synapse-type neurons in the brain. We defined the pre- and post-synapse segregation index

(PPSSI) and found that only some cells have a clear synaptic segregation, and many neurons have

unsegregated synapses. When we extracted synapses with a particularly low segregation (high

intermingling) and compared them between SC and FC, we found a significantly higher FC-SC

correlation. Conversely, neurons with a high segregation had significantly lower FC-SC correlations.

Many *Drosophila* brain neurons are interconnected [30]. As such, we similarly investigated reciprocal

synapses [31, 32, 33, 34, 35], whose function is clearer than that of other unsegregated synapse types.

When we extracted such reciprocal synapses, the FC-SC correlation was significantly higher. Our results

indicated that unsegregated synapses contributed more to the FC. In other words, local activity due to

unsegregated synapses could be considered the more essential functional activity. Unsegregated synapses

are found not only in flies but also in mammals [31, 32, 18, 36, 37, 38], including humans

(Ext.Data.Fig.1-2 from [39]). Localized functional activity from pre-synaptic population dynamics can be

observed by calcium imaging or detected through the neurovascular coupling with astrocytes [40] via

fMRI. Although modalities vary, the FC-SC correlation between humans and flies does not considerably

differ, as shown earlier. Since various fMRI pre-processing and analysis methods can be directly applied

to pre-synaptic population dynamics in this study, functional activity is locally compartmentalized in the

whole brain, which is common among various species.

Results

Investigation of the optimal nuisance factor removal method for calcium imaging

First, the optimal pre-processing of the calcium imaging data of *D. melanogaster* was investigated. The

registration method used in fMRI was applied to the calcium image (Ext.Data.Fig.1-1) to register to the

FDA template [14] (Fig. 1b). Then, various nuisance factor removal methods (details in Methods) were

applied to the ROI time-series to generate the FC matrices of eight flies (Fig. 1a). Three types of ROIs

were used in the analysis: ROIs based on the anatomical neuropil, commonly used in *Drosophila*

(hemibrain ROI), ROIs based on k-means clustering of the SC matrix of 146,604 voxels in the hemibrain

(SCKm), and ROIs based on distance-based clustering of voxels in the hemibrain (DistKm). SCKm can

provide ROIs close to the shape of the anatomical neuropil across various ROI counts (Ext.Data.Fig.2-1a),

while DistKm can provide ROIs that are completely unrelated to the anatomical neuropil. Next, a

correlation analysis and ROC curve analysis were performed using the SC matrices based on FlyEM and

the FC matrices with these ROI types (Fig. 1a). Fig. 1d-f shows the FC-SC correlation results, FC-SC

detection results, and the normalized total score of FC-SC detection and correlation (details in Methods),

respectively. After the comparison of the raw (leftmost) and removal methods for various ROI types, the

combination of polynomial detrending [41] & tCompCor [42] was found to yield the optimal results. The

optimal removal methods were found to differ between fMRI [10] and calcium imaging data, and these

results are further described in the Discussion section.

Comparison of the FC-SC correlation between humans and flies

Spatial smoothing is useful for absorbing inter-individual variability and conducting second-level group

analysis [10, 43]. Here, we explored the effect of spatial smoothing (of the calcium imaging data) on

various ROI counts in DistKm (Fig. 2a) and SCKm (Ext.Data.Fig.2-1a) ROIs. In DistKm, smoothing was

highly effective, and the optimal kernel radius of the Gaussian filter was around 23 voxels (52.44 μm) for

various ROI counts (Fig. 2b). In SCKm, smoothing was ineffective for ROI count below 100, while the

kernel radius above 16 voxels (36.48 μm) was effective for ROI count above 100 (Ext.Data.Fig.2-1b).

SCKm has a larger variation in volume for each ROI count than DistKm (Fig. 1c, Fig. 2e); as such, the

effect of spatial smoothing is likely unstable. Next, we investigated whether the combination of spatial

smoothing and Poly-tCompCor was effective. We found that this combination was significantly more

effective than smoothing alone for SCKm and DistKm (Fig. 2c, Mann–Whitney U test p<1.67e-4

Bonferroni corrected). Conversely, the combination of a high-pass filter and nuisance factor removal

overlapped for the purpose of noise removal, and the effect was degraded (Ext.Data.Fig.2-1d), as found in

fMRI [10].

The optimal FC-SC correlation for various ROI counts was determined. Results showed that the

correlation decreased proportionally to the logarithm of the ROI count (Fig. 2d). This was also compared

between flies and mammals. At approximately 50 ROIs, the difference was nearly 1.08-fold to 1.22-fold

(r=0.66 for humans and r=0.71-0.81 for flies [SCKm, DistKm]); by contrast, at approximately 1,000 ROIs,

the difference was roughly 1.12-fold to 1.24-fold (r=0.54 for humans and r=0.6-0.67 for flies), which was

not a considerable difference. The difference was slightly larger (1.21 to 1.5-fold: r=0.378 for marmosets

and r=0.46-0.57 for flies) than the marmoset results at around 10,000 ROIs (Ext.Data.Fig.2-1c). Although

SCs for flies and humans were obtained from single individuals, the tracer data is a composite from 52

marmosets [44], which might contribute to the difference with the fly's FC-SC correlation.

Comparison between FlyEM and FlyWire connectome data

We used the FlyEM's SC to compare the averaged FC of eight flies. In this section, we also used the

FlyWire connectome data [45] to compare the FC and each SC, and possibly reveal the inter-individual

variability of the two EM connectome data. As mentioned earlier, FlyEM has a synaptic "confidence"

threshold and contains 21.4 million post-synapses in the hemibrain region at its initial value of 0.

Conversely, FlyWire has a 'CleftScore' [17] threshold and 18.8 million post-synapses at its initial value

of 50. Since the number of post-synapses differs at each threshold, we set the confidence threshold of 0.8

as high as possible, thereby increasing the FC-SC correlation (Fig. 3d, ExtFig.3-1b). Then we adjusted the

CleftScore to 140, which has a post-synapse number of approximately 11.4 million (Fig. 3a). These

thresholds were used for all analyses in this study. When we displayed the maximum Z-projection of the

synapse point cloud of post-synapse counts in each voxel, we found that the trends in synapse detection

completely varied (Fig. 3b). In each region, FlyEM had more synapses in the mushroom bodies (aL, bL,

etc.) and in the major neuropils of the central nervous system (CNS; EB, FB, NO, and PB). In comparison,

FlyWire had more synapses in the visual system (LO and ME) and in the periesophageal neuropils (CAN

and FLA; Fig. 3c). In the FC-SC correlation of each region, FlyEM had higher values in most regions at

58:5 (Fig. 3e, ExtFig.3-1a). As a result, FlyEM r=0.73 differed from FlyWire r=0.68 at the matrix level

(Fig. 3d). Both datasets were from genotype Canton-S-G1×w1118 adult females; FlyEM was five-day-old

[12], and FlyWire was seven-day-old [13]. Although brain development may differ depending on the

rearing environment, the tendency to detect synapses substantially varied. We attribute this to differences

in detection accuracy resulting from the resolution of EM scanning, but not to inter-individual variability.

We next used the FlyWire connectome data to investigate the ratios of neurotransmitter input (ExtFig.3-

2a) and output (Fig. 3f) in each region. The highest percentage was acetylcholine (62.8%±14.2%),

followed by glutamine (15.5%±9.2%), then GABA (11.4%±8.2%) in neurotransmitter output. The

analysis of the relationship between this ratio and FC-SC correlation in each neurotransmitter revealed

significant correlations for acetylcholine (r=0.39, p=0.0013) and GABA (r=-0.25, p=0.046). That is, the

higher the percentage of excitatory connections, the higher the FC-SC correlation; conversely, the higher

the percentage of inhibitory connections, the lower the FC-SC correlation. In humans, FC-SC correlations

are lower in the evolutionarily expanded Brodmann area 39,45 and the temporal lobe [8]; therefore, these

areas may have more long range inhibitory connections [46, 47, 48].

Unsegregated synapses contribute considerably to functional connectivity

We compared FC-SC in the macroscopic regions, and we also investigated the relationship between the

synapses, the source of functional activity (pre-synaptic population dynamics), and FC. First, we

examined the synaptic segregation in whole-brain neurons because many neurons showed pre- and post-

synaptic intermingling. Then, we determined how unsegregated and segregated synapses affected the FC.

To quantify synaptic segregation, we developed the PPSSI (Fig. 4d). Using DBSCAN [49] with a radius

of 3 μm (details on Methods), we clustered synapses in each neuron and calculated the degree of

segregation for each cluster (cPPSSI). We then multiplied the cPPSSI by the fraction of the synapse

number and obtained the sum to determine the PPSSI of each neuron. Fig. 4b shows the result of applying

PPSSI to 139,255 whole-brain neurons of FlyWire. The result showed a histogram resembling a normal

distribution. When we created unsegregated ranking (details in Methods) by considering the number of

synapses, APL and CT1 cells occupied the top 1–4 positions, indicating that we efficiently extracted

unsegregated neurons (Fig. 4a, Ext.Data.Fig.4-3a). APL was also at the top for FlyEM (Ext.Data.Fig.4-

1a). Similarly, we established segregated ranking and successfully extracted neurons with a high degree

of segregation (Fig. 4c, Ext.Data.Fig.4-3; Ext.Data.Fig.4-1c and Ext.Data.Fig.4-4 for FlyEM).

Next, synapses with a cPPSSI between 0 and 0.1 were extracted, and the SC matrices (unsegregated

synapses) of neurons and synapses were created. Synapses were then randomly sampled to generate 99

null SC matrices (random synapses), centered around the total number of connections (2,373,100) in the

SC matrix (unsegregated synapses), forming a normal distribution (Fig. 4e, p=0.832, details in Methods).

The FC-SC correlations of the SC matrices (unsegregated synapses) of neurons and synapses were

significantly higher than those of the randomly extracted samples (Fig. 4e, p=0, 6.5e-43; Ext.Data.Fig.4-

1d for FlyEM). Therefore, unsegregated synapses might strongly contribute to FC. A similar analysis was

performed on synapses with a cPPSSI of 0.9–1, which surprisingly showed a significantly lower FC-SC

correlation (Fig. 4f, p=9.2e-14, 0; Ext.Data.Fig.4-1e for FlyEM). Therefore, segregated synapses may not

be able to contribute to strongly to the FC, and these findings are further explored in the Discussion

section.

Among unsegregated synapses, reciprocal synapses have been well studied [31, 32, 33, 34, 35] and their

function is considered to be clear (Fig. 4g, Ext.Data.Fig.4-1f). The shortest distance of such reciprocal

synapses was extracted from whole-brain neurons, shown in Fig. 4h (Ext.Data.Fig.4-1g for FlyEM). They

are particularly abundant within 2 μm (Ext.Data.Fig.4-2a,c for CT1 and APL cells), and acetylcholine and

GABA constitute the majority (Fig. 4h). The comparison of the SC matrix based on reciprocal synapses

(<2 μm) with the null SC matrices showed that the FC-SC correlation was significantly higher (Fig. 4I,

p=0, 2.7e-32; Ext.Data.Fig.4-1h for FlyEM). Therefore, unsegregated synapses and reciprocal synapses

(<2 μm) strongly contributed to FC. Furthermore, since the FC was the average of eight flies, such

synapses are likely functionally common among individuals.

Region suppression analysis during fly walking

The whole-brain calcium imaging dataset from Brezovec et al. [14] provides fly walking data. Since

fMRI pre-processing methods can be applied to calcium imaging data, we investigated whether voxel-

level analysis using a general linear model (GLM) [50] could be applied to the fly walking task (Fig. 5a,

details in Methods). Fig. 5b shows the GLM analysis result. Similar to the results of Brezovec et al., our

findings revealed that LAL, FB, and IPS neuropils showed strong activity; the GNG neuropil, which is

highly connected to leg neuropils in the ventral nerve code [51], also exhibited strong activity. Although

Brezovec et al. detected voxels that are strongly correlated with velocity, the GLM extracted voxels

associated with walking behavior. Conversely, we found inactivity or strong suppression in the Wedge

and AVLP neuropils. We searched for a GABA neuron that would fit with this phenomenon and managed

to find one that extended its dendrites widely from the Wedge to AVLP neuropils (Fig. 5e, h). The

synapse point cloud could also fit in this region (Fig. 5c). We called this neuron the Wedge-AVLP GABA

neuron (WAGN). The Wedge neuropil receives mechanosensory (wind, sound) input and connects to

WPNb neurons (Ext.Data.Fig.5-1a,b), which then connect to ER1 ring neurons that act as the fly's

compass [52, 53]. Interestingly, the left WAGN was strongly interconnected with WPNb tier 2/3 [54] and

received input from 22 ascending neurons (ANs; Fig. 5d). Therefore, it may be a pathway that provides

walking behavior feedback to the compass; however, wind caused by walking was absent in the

experimental setup (the fly was fixed and walked on a treadmill), and may relate to the strong regional

suppression that was observed. As such, further experimentation is needed to pursue this topic.

The left WAGN resembles a hub neuron; however, like CT1 and APL cells, it has highly unsegregated

synapses (PPSSI=0.133). Synapses on the medial side were highly segregated, but the lateral side was

well unsegregated (Fig. 5h). It received input from ANs (212 synapses) mainly on the ventral side, and

outputs to WPNb (65 synapses), widely from the medial to the lateral side (Fig. 5e). A histogram of the

shortest straight-line distances between pre-synapses to WPNb and post-synapses from ANs is shown in

Fig. 5f, and the closest distance was 852.3 nm (Fig. 5g). Many acetylcholine inputs (disynaptic inhibition

[25]) were closer than this value (Fig. 5i), and many GABA inputs from ANs (Fig. 5f) further suppress

such disynaptic inhibition. AVLP ventral suppression (Fig. 5b white arrows) may be the suppression from

these ANs. Thus, a highly delicate and localized control is possibly exercised from WAGN to WPNb tier

2/3.

Discussion

Although nuisance factor removal methods such as global mean (GM) and aCompCor [42] are optimal for fMRI [10], calcium imaging showed different results. GM is the average of all voxels, and the regression of this signal is useful in fMRI because noise from the equipment and coils affects the entire voxel image. Conversely, in two-photon microscopy, noise does not affect all voxels because of the excitation of specific molecules by laser absorption, and GM does not show a positive effect. aCompCor extracts the component time-series from the voxel time-series of white matter and cerebrospinal fluid (CSF) via PCA and then performs a regression. In flies, the CSF had no ROI mask, but fibers had ROIs; therefore, fibers were used instead of the white matter. However, many synapses may exist around these fibers, and aCompCor was not as effective as in humans. Conversely, polynomial detrending [41] was highly effective (Fig. 1d-f), likely because of linear thermal noise and other factors caused by the continuous excitation of molecules (for 30 min). tCompCor [42] is the component time-series extracted by PCA from voxels with the top 1000 standard deviations among all voxels. It potentially captures minute fluctuations caused by fly movements and effectively removes them.

The SC matrix from a cPPSSI of 0.9–1 synapses resulted in a significantly lower FC-SC correlation than the SC matrix from randomly selected synapses (Fig. 4f for FlyWire, Ext.Data.Fig.4-1e for FlyEM). This result was surprising, but we propose several possible reasons. Neurons with highly segregated synapses include ascending neurons (ANs) and descending neurons (DNs), (Ext.Data.Fig.4-3b). They also have a small number of unsegregated synapses (i.e. Sep rank9: PPSSI=0.959, Sep rank11: PPSSI=0.964). They are thought to be specialized for input or output in the CNS regions and may show large differences between SC and FC. Another reason is that these neurons may have widespread logical OR inputs and outputs. In Fig. 4c, an action potential at one point in the input synapse (blue) is likely transmitted to all red synapses. Such an OR circuit-like disproportionate input–output possibly causes a mismatch with the FC, because the functional activity in the output region originates from activity at a single location. These neurons are likely those whose FC are difficult to capture. Conversely, neurons with highly unsegregated

synapses may capture the inputs of neighboring synapses, such as forming a logical AND circuit,

correlating well with the output regions.

The FC-SC correlation was highly dependent on ROI count in both flies and humans (Fig. 2d). Through

optimal nuisance factor removal and smoothing, we achieved r=0.87 at 20 ROIs. Furthermore, by

applying Spearman correlation instead of Pearson, we obtained r=0.928. Thus, for a reasonably small ROI

count the correlation was very high; however, for 10,000 ROIs, the correlation decreased to r=0.57. This

phenomenon could be attributed to the presence of highly segregated synapses. Such neurons are

widespread in the CNS (Fig. 4b, Ext.Data.Fig.4-2f) and are thought to provide logical OR circuits. When

ROI volume was large (when the ROI count was small), a single ROI could cover the distributed input

synapses. However, when a ROI was divided, input synapses were also divided across many ROIs, and

the mismatch between SC and FC would increase. Other possibilities include the presence of gap

junctions and FC due to neuropeptides [55], but they are outside of the scope of the present study and

their relationship with ROI count is unclear.

In this study, we applied fMRI techniques directly to calcium imaging in flies and found many similarities

between humans and flies (Fig. 2d). The calcium signal represents pre-synaptic population dynamics and

likely includes the localized functional activity of unsegregated synapses [25, 31, 32, 33, 34], such as

non-spiking [21, 22, 23, 24, 25, 26] and sub-threshold neuronal activity [18]. For fMRI, the signal

captures localized functional activity as blood-oxygenation-level-dependent signal via neuro vascular

coupling [40]. These two signals can be pre-processed (Fig. 1, Fig. 2) and analyzed (Fig. 5a) in

surprisingly similar ways. This is unlikely a coincidence, and it suggests that similar design concepts

underlie neural circuits in flies and mammals. Using *D. melanogaster* as a model animal, we were able to

elucidate detailed local functional activities around unsegregated synapses and compare these results to

the brain activities of mammals, including humans.

Materials and Methods

Preprocessing of *D. melanogaster* calcium imaging data

Whole-brain calcium imaging data of *D. melanogaster* were acquired by Brezovec et al. [14] and

downloaded from DANDI (https://dandiarchive.org/dandiset/000727/0.240106.0043). 4D NIfTI image

data (256×128×49 voxels, 3384 frames) were extracted from a NWB file by an in-house MATLAB script.

The FDA template, made from in-vivo calcium imaging, was resized (256×128×49 voxels,

2.45×2.28×3.72μm for a voxel) and used as a template because the original thresholded FDA template

[56] was extremely large (1652×768×479 voxels), and the calcium imaging file was large (more than 40

GB); therefore, a smaller template with the imaging data was preferably used. Preprocessing and image

registration were performed using Statistical Parametric Mapping (SPM12) [57] and ANTs [58]

(Ext.Data.Fig.1-1a). Motion and slice timing correction were performed using SPM12, and the averaged

NIfTI image was registered by ANTs to the resized FDA template. Lastly, 4D NIfTI images were

transformed using the transform information. The data contained n=9 flies, but 8 flies were used for the

ROI-based analysis because of missing time-series in some ROIs in one fly. Nine flies were used in the

GLM analysis.

Combinations of the following nuisance factor removal methods were investigated: global mean (average

signal across all voxels), global signal (average signal across all brain voxels), mean white matter

(average signal across fiber voxels), 6HMP (head motion parameters), 24HMP, aCompCor, tCompCor

[42], and polynomial detrending (Poly) [41]. These methods were implemented by an in-house MATLAB

script [59]. A full width at half maximum (FWHM) from 1 voxel (2.28 μm) to 30 voxels (68.4 μm), at a 1

voxel step, was investigated to determine the optimal spatial smoothing size for calcium imaging data.

Temporal high-pass filters were also examined from 0.1 Hz to 0.001 Hz.

Transformation of EM connectome data into the resized FDA template

In the FlyWire case, connectome data, such as neurons and synapses, were downloaded from FlyWire

Codex (https://codex.flywire.ai/) and Zenodo (https://zenodo.org/records/10676866) [13, 60, 45]. Detailed

synapse data were saved in an Apache feather format, which was then converted to a CSV file. Synapse

points were transformed from the FlyWire space to a JRC2018 female template [61] by the Python navis-

flybrains package [61]. Since FlyWire coordinates are left and right reversed [13], they were flipped on

the JRC2018 female template, and these points were transformed to the resized FDA template by ANTs.

The use of these two templates facilitated precise data alignment across different modalities.

In the FlyEM case, the latest connectome data were downloaded from Janelia's DVID website

(https://dvid.io/blog/release-v1.2/). Like those in the FlyWire case, synapse points were transformed from

the FlyEM space to a JRC2018 female template by the Python navis-flybrains package. These points were

transformed to a resized FDA template by ANTs.

Generation of regions of interest for the hemibrain

The hemibrain neuropil atlas (114 ROIs, including primary and sub-regions, aligned with the FlyEM

space) was downloaded from Virtual Fly Brain (https://www.virtualflybrain.org/). ROI files were

transformed from the FlyEM space to a JRC2018 female template by Java template-building scripts

(https://github.com/saalfeldlab/template-building) [61]. They were transformed to a resized FDA template

by ANTs. In Fig. 3, 63 primary ROIs mentioned above were used. The 52 ROIs in Fig. 1 were selected

from the primary ROIs, and approximately 50 ROIs (those used by Turner et al. [9]) were selected. ROIs

were segmented by SCKm (Fig. 1 and Fig. 2) via the k-means clustering of the SC matrix of 146,604

voxels in the hemibrain. They were also segmented by DistKm (Fig. 1 and Fig. 2) via the k-means

clustering of the distance of 146,604 voxels in the hemibrain.

The SC matrix was calculated according to synapse locations and ROIs. Since synapse data formats differ

between FlyEM and FlyWire, they were standardized by using post-synapses within ROIs. The input and

output neurons at each ROI were extracted from the paired pre- and post-synapse data. Then the input and

output neurons and their synapses that were common between ROI pairs were counted to create SC

matrices for neurons and synapses. The FC matrix was calculated using the pre-processed voxel time-

series and ROIs. The intensity of the voxels in the ROI was averaged to obtain the average intensity time-

series of the ROI. Then, Pearson correlation was calculated between ROI pairs to generate the FC matrix.

The correlation coefficients were z-transformed, calculated for eight flies, and then averaged to obtain the

final FC matrix.

FC-SC detection and correlation

FC-SC correlation was calculated by taking the Pearson correlation between the logarithm of the SC

matrix and the FC matrix, as done in other studies [6, 7, 10]. In the SC matrix, elements with zero

connections become *-Inf* when logarithmically converted, so we substituted zero for them. Also, the

diagonal elements of the FC matrix become *Inf* when z-transformed, so we replaced them with the

maximum value of the matrix. For one FC matrix, we calculated its correlation with both the neuron SC

matrix and the synapse SC matrix. FC-SC detection was calculated by ROC (Receiver Operating

Characteristic) curve analysis. This gives the detection accuracy of the FC matrix when the SC matrix is

used as ground truth. Thresholds were taken from 0% to 99% by 1% steps of all elements of the SC

matrix, and 100 AUCs (Area Under the Curve) were calculated. The average of these AUCs is called the

FC-SC detection. FC-SC correlation indicates the similarity of the two matrices, while FC-SC detection

indicates the detection accuracy. Although the digit number of connections differs between neurons and

synapses, FC-SC detection can absorb such differences. The FC-SC detection & correlation score was

calculated as follows:

$$FC\text{-}SC_{det\&corr\ score} = |FC\text{-}SC_{correlation}| + 2 \times |FC\text{-}SC_{detection} - 0.5|\#(1)$$

338   The FC-SC detection & correlation score is in the range of [0 2], where the larger the score, the higher the

339   similarity and detection accuracy.

PPSSI

The PPSSI (Fig. 4d) was used in DBSCAN [49] with a radius of 3 μm to generate pre- and post-synaptic

clusters. The radius was empirically defined by several criteria. Firstly, CT1 and APL cells have the

shortest distance of reciprocal synapses, within 2 μm (Ext.Data.Fig.4-2a,c). In zebrafish, reciprocal

synapses were defined within 2.5 μm [35]. Schneider-Mizell et al. [19] defined a twig as a branch within

3 μm from the backbone of the cytoskeleton, and 80% of synaptic inputs in motor neurons are in the twig.

In addition, 97% of pre-synaptic sites are located at less than 3 μm from the mitochondria. Therefore, 3

μm was deemed appropriate for the synapse cluster radius. Once the synapse clusters were determined by

DBSCAN, the degree of synaptic segregation for each cluster was calculated using the following formula:

$$cPPSSI_i = \left(\frac{pre_i - post_i}{pre_i + post_i}\right)^2, \#(2)$$

349   where pre_i and $post_i$ are the pre- and post-synapse counts in cluster i, respectively. The cPPSSI of each

350   cluster (each synapse) is in the range of [0 1]. The PPSSI of each neuron was calculated as follows:

$$PPSSI = \sum_{i=1}^{N} \frac{(pre_i + post_i)}{W} \cdot cPPSSI_i \; \#(3)$$

351   where N is the cluster number, and W is the total synapse count for a neuron.

352   The unsegregated ranking was calculated based on the PPSSI and total synapse count of neuron k:

$$unsegregated \; score_k = log_{10}(W_k) \cdot (1 - PPSSI_k)\#(4)$$

353   where $PPSSI_k$ is the PPSSI of neuron k and W_k is the total synapse count of neuron k. This allows us to

354   extract neurons with many inputs and outputs and highly unsegregated synapses. The segregated ranking

355   was also calculated in the same way, using the following score:

$$segregated \; score_k = log_{10}(W_k) \cdot PPSSI_k\#(5)$$

A similar index to our PPSSI, namely, the segregation index, has been defined in the literature [19], but

its use poses several problems. For neurons such as ANs and DNs, that have only input or output

synapses, the segregation index results in calculations of $0 \times \infty$ or division by 0; consequently,

calculation becomes impossible. Despite the [0 1] range of indices, the histogram is heavily biased toward

0 because of the use of the logarithm, and the mean is much closer to 0 than 0.5 (Ext.Data.Fig.4-1i, j).

Therefore, in the present study, we developed the more robust PPSSI because it is less biased and its

mean is close to 0.5 (Fig. 4b, Ext.Data.Fig.4-1b).

GLM Analysis for the fly walking task

GLM analysis (Fig. 5a) was performed to investigate the whole-brain functional activity during the fly

walking task. Movement velocity data were extracted from an NWB file by an in-house MATLAB script.

Walking was defined as 1σ (34%) or more of the movement velocity and divided into walking and other.

Less than 10% of movement velocity was zero cut. The canonical calcium response function (CRF) was

created by the gamma probability density function (a=1.7, b=0.25) in MATLAB based on the GCaMP6f

response [62]. The walking-task time-series was transformed into a calcium response time-series by

convolution with the CRF. GLM design matrices were created using the calcium response time-series and

nuisance factors (polynomial detrending and tCompCor). In whole-brain voxels, smoothing was applied

with a kernel radius of 8 voxels (18.24 μm). Then, a Tukey taper (taper size = 8) was used for GLM pre-

whitening [63]. A mixed-effects model [43] was used for group analysis, and the t-value of each voxel

was calculated via the second-level analysis of OLS regression with a Tukey taper. These analyses were

implemented using in-house MATLAB scripts [64].

Statistical Analysis

A Mann–Whitney U test was used to examine whether the combination of spatial smoothing and nuisance

factor removal was effective. Statistical significance was set at $p<0.05$, and Bonferroni correction was

applied to correct the familywise error (FWE) rate. A permutation test was performed to analyze whether

specific synapses (segregated, unsegregated, or reciprocal synapses) affect SC-FC correlation. The SC

matrix (specific synapses) in the 63 ROIs in the hemibrain was generated, and its SC-FC correlation was

calculated. Then, 99 null SC matrices (random synapses) were generated so that they would be normally

distributed around the total number of connections in the SC matrix (specific synapses). Since the SC-FC

correlations of the null SC matrices (random synapses) also roughly followed a normal distribution, the

T-value of the SC-FC correlation of the SC matrix (specific synapses) was calculated using the

cumulative distribution function of the normal distribution. For GLM analysis, a mixed-effects model was

used for group analysis.

Acknowledgements

389   The authors wish to thank Dr. Clandinin for providing valuable calcium imaging data and answering our

390   question.

391   This research was supported by the program for Brain Mapping by Integrated Neurotechnologies for

392   Disease Studies (Brain/MINDS 2.0) from the Japan Agency for Medical Research and Development,

393   AMED. Grant number: JP24wm0625408.

References

[1] G. L. Gerstein and D. H. Perkel, "Simultaneously recorded trains of action potentials: Analysis and functional interpretation.," *Science,* vol. 164, pp. 828-830, 1969.

[2] G. L. Gerstein, P. Bedenbaugh and A. M. Aertsen, "Neuronal assemblies," *IEEE Trans. Biomed. Eng.,* vol. 36, pp. 4-14, 1989.

[3] K. J. Friston, C. D. Frith, P. F. Liddle and R. S. Franckowiak, "Functional Connectivity: The Principal-Component Analysis of Large (PET) Data Sets," *J. Cereb. Blood Flow Metab.,* vol. 13, pp. 5-14, 1993.

[4] K. Friston, R. Moran and A. K. Seth, "Analysing connectivity with Granger causality and dynamic causal modelling," *Curr. Op. in Neuro.,* vol. 23, no. 2, pp. 172-178, 2013.

[5] P. Hagmann, L. Cammoun, X. Gigandet, R. Meuli, C. J. Honey, V. J. Wedeen and O. Sporns, "Mapping the Structural Core of Human Cerebral Cortex," *PLoS Biol,* vol. 6, no. 7, p. e159, 2008.

[6] C. J. Honey, O. Sporns, L. Cammoun, X. Gigandet, J. P. Thiran, R. Meuli and P. Hagmann, "Predicting human resting-state functional connectivity from structural connectivity," *Proc. Natl. Acad. Sci. USA,* vol. 106, no. 6, pp. 2035-2040, 2009.

[7] C. J. Honey, J.-P. Thivierge and O. Sporns, "Can structure predict function in the human brain?," *NeuroImage,* vol. 52, no. 3, pp. 766-776, 2010.

[8] G. L. Baum, Z. Cui, D. R. Roalf, R. Ciric, R. F. Betzel, B. Larsen, M. Cieslak, P. A. Cook, C. H. Xia, T. M. Moore, K. Ruparel, D. J. Oathes, A. F. Alexander-Bloch, R. T. Shinohara, A. Raznahan, R. E. Gur, R. C. Gur, D. S. Bassett and T. D. Satterthwaite, "Development of structure–function coupling in human brain networks during youth," *PNAS,* vol. 117, no. 1, pp. 771-778, 2020.

[9] M. H. Turner, K. Mann and T. R. Clandinin, "The connectome predicts resting-state functional connectivity across the Drosophila brain," *Current Biology,* vol. 31, p. 2386–2394, 2021.

[10] T. Okuno, J. Hata, C. Kawai, H. Okano and A. Woodward, "A Novel Directed Seed-Based Connectivity Analysis Toolbox Applied to Human and Marmoset Resting-State fMRI," *J. Neurosci.,* vol. 44, no. 45, p. e0389242024, 2024.

[11] R. Hira, L. B. Townsend, I. T. Smith, C.-H. Yu, J. N. Stirman, Y. Yu and S. L. Smith, "Mesoscale functional architecture in medial posterior parietal cortex," *eLife,* p. 105213v1, 2025.

[12] L. K. Scheffer, C. S. Xu, M. Januszewski, Z. Lu, S. Takemura, K. J. Hayworth, G. B. Huang, K. Shinomiya, J. Maitlin-Shepard, S. Berg, J. Clements, P. M. Hubbard, W. T. Katz, L. Umayam, T. Zhao, D. Ackerman, T. Blakely, J. Bogovic, T. Dolafi, D. Kainmueller, T. Kawase, K. A. Khairy, L. Leavitt, P. H. Li, L. Lindsey, N. Neubarth, D. J. Olbris, H. Otsuna, E. T. Trautman, M. Ito, A. S. Bates, J. Goldammer, T. Wolff, R. Svirskas, P. Schlegel, E. Neace, C. J. Knecht, C. X. Alvarado, D. A. Bailey, S. Ballinger, J. Z. Borycz, B. S. Canino, N. Cheatham, M. Cook, M. Dreher, O. Duclos, B. Eubanks, K. Fairbanks, S. Finley, N. Forknall, A. Francis, G. P. Hopkins, E. M. Joyce, S. Kim, N. A. Kirk, J. Kovalyak, S. A. Lauchie, A. Lohff, C. Maldonado, E. A. Manley, S. McLin, C. Mooney, M. Ndama, O. Ogundeyi, N. Okeoma, C. Ordish, N. Padilla, C. M. Patrick, T. Paterson, E. E. Phillips, E. M. Phillips, N. Rampally, C. Ribeiro, M. K. Robertson, J. M. Rymer, S. M. Ryan, M. Sammons, A. K. Scott, A. L. Scott, A. Shinomiya, C. Smith, K. Smith, N. L. Smith, M. A. Sobeski, A. Suleiman, J. Swift, S. Takemura, I. Talebi, D. Tarnogorska, E. Tenshaw, T. Tokhi, J. J. Walsh, T. Yang, J. A. Horne, F. Li, R. Parekh, P. K. Rivlin, V. Jayaraman, M. Costa, G. S. Jefferis, K. Ito, S. Saalfeld, R. George, I. A. Meinertzhagen, G. M. Rubin, H. F. Hess, V. Jain and S. M. Plaza, "A connectome and analysis of the adult Drosophila central brain," *eLife,* vol. 9, p. e57443, 2020.

[13] S. Dorkenwald, A. Matsliah, A. R. Sterling, P. Schlegel, S. Yu, C. E. McKellar, A. Lin, M. Costa, K. Eichler, Y. Yin, W. Silversmith, C. Schneider-Mizell, C. S. Jordan, D. Brittain, A. Halageri, K. Kuehner, O. Ogedengbe, R. Morey, J. Gager, K. Kruk, E. Perlman, R. Yang, D. Deutsch, D. Bland, M. Sorek, R. Lu, T. Macrina, K. Lee, J. A. Bae, S. Mu, B. Nehoran, E.

Mitchell, S. Popovych, J. Wu, Z. Jia, M. A. Castro, N. Kemnitz, D. Ih, A. S. Bates, N. Eckstein, J. Funke, F. Collman, D. D. Bock, G. S. Jefferis, H. S. Seung, M. Murthy and The FlyWire Consortium, "Neuronal wiring diagram of an adult brain," *Nature,* vol. 634, p. 124–138, 2024.

[14] B. E. Brezovec, A. B. Berger, Y. A. Hao, F. Chen, S. Druckmann and T. R. Clandinin, "Mapping the neural dynamics of locomotion across the Drosophila brain," *Current Biology,* vol. 34, p. 710–726, 2024.

[15] P. Schlegel, A. S. Bates, T. Stürner, S. R. Jagannathan, N. Drummond, J. Hsu, L. S. Capdevila, A. Javier, E. C. Marin, A. Barth-Maron, I. F. Tamimi, F. Li, G. M. Rubin, S. M. Plaza, M. Costa and G. S. Jefferis, "Information flow, cell types and stereotypy in a full olfactory connectome," *eLife,* vol. 10, p. e66018, 2021.

[16] P. Schlegel, Y. Yin, A. S. Bates, S. Dorkenwald, K. Eichler, P. Brooks, D. S. Han, M. Gkantia, M. D. Santos, E. J. Munnelly, G. Badalamente, L. S. Capdevila, V. A. Sane, A. M. Fragniere, L. Kiassat, M. W. Pleijzier, T. Stürner, I. F. Tamimi, C. R. Dunne, I. Salgarella, A. Javier, S. Fang, E. Perlman, T. Kazimiers, S. R. Jagannathan, A. Matsliah, A. R. Sterling, S. Yu, C. E. McKellar, FlyWire Consortium, M. Costa, H. S. Seung, M. Murthy, V. Hartenstein, D. D. Bock and G. S. Jefferis, "Whole-brain annotation and multi-connectome cell typing of Drosophila," *Nature,* vol. 634, pp. 139-152, 2024.

[17] L. Heinrich, J. Funke, C. Pape, J. Nunez-lglesias and S. Saalfeld, "Synaptic Cleft Segmentation in Non-Isotropic Volume Electron Microscopy of the Complete Drosophila Brain," *arXiv,* p. 1805.02718, 2018.

[18] T. Branco and M. Häusser, "The single dendritic branch as a fundamental functional unit in the nervous system," *Current Opinion in Neurobiology,* vol. 20, pp. 494-502, 2010.

[19] C. M. Schneider-Mizell, S. Gerhard, M. Longair, T. Kazimiers, F. Li, M. F. Zwart, A. Champion, F. M. Midgley, R. D. Fetter, S. Saalfeld and A. Cardona, "Quantitative neuroanatomy for connectomics in Drosophila," *eLife,* vol. 5, p. e12059, 2016.

[20] O. T. Shafer, G. J. Gutierrez, K. Li, A. Mildenhall, D. Spire, J. Marty, A. A. Lazar and M. D. Fernandez, "Connectomic analysis of the Drosophila lateral neuron clock cells reveals the synaptic basis of functional pacemaker classes," *eLife,* vol. 11, p. e79139, 2022.

[21] M. Meier and A. Borst, "Extreme Compartmentalization in a Drosophila Amacrine Cell," *Current Biology,* vol. 29, pp. 1545-1550, 2019.

[22] R. Behnia, D. A. Clark, A. G. Carter, T. R. Clandinin and C. Desplan, "Processing properties of ON and OFF pathways for Drosophila motion detection," *Nature,* vol. 512, pp.

427-430, 2014.

[23] E. Gruntman, P. Reimers, S. Romani and M. B. Reiser, "Non-preferred contrast responses in the Drosophila motion pathways reveal a receptive field structure that explains a common visual illusion," *Current Biology,* vol. 31, pp. 5286-5298, 2021.

[24] L. N. Groschner, J. G. Malis, B. Zuidinga and A. Borst, "A biophysical account of multiplication by a single neuron," *Nature,* vol. 603, pp. 119-123, 2022.

[25] A. Braun, A. Borst and M. Meier, "Disynaptic inhibition shapes tuning of OFF-motion detectors in Drosophila," *Current Biology,* vol. 33, pp. 2260-2269, 2023.

[26] J. E. Schenk and Q. Gaudry, "Nonspiking Interneurons in the Drosophila Antennal Lobe Exhibit Spatially Restricted Activity," *eNeuro,* vol. 10, no. 1, pp. ENEURO.0109-22.2022, 2023.

[27] K. Inada, Y. Tsuchimoto and H. Kazama, "Origins of Cell-Type-Specific Olfactory Processing in the Drosophila Mushroom Body Circuit," *Neuron,* vol. 95, pp. 357-367, 2017.

[28] H. Amin, A. A. Apostolopoulou, R. Suárez-Grimalt, E. Vrontou and A. C. Lin, "Localized inhibition in the Drosophila mushroom body," *eLife,* vol. 9, p. e56954, 2020.

[29] X. Liu and R. L. Davis, "The GABAergic anterior paired lateral neuron suppresses and is suppressed by olfactory learning," *Net. Neurosci.,* vol. 12, pp. 53-59, 2009.

[30] A. Lin, R. Yang, S. Dorkenwald, A. Matsliah, A. R. Sterling, P. Schlegel, S. Yu, C. E. McKellar, M. Costa, K. Eichler, A. S. Bates, N. Eckstein, J. Funke, G. S. Jefferis and M. Murthy, "Network statistics of the whole-brain connectome of Drosophila," *Nature,* vol. 634, pp. 153-165, 2024.

[31] W. N. Grimes, W. Li, A. E. Chávez and J. S. Diamond, "BK channels modulate pre- and postsynaptic signaling at reciprocal synapses in retina," *Nat. Neurosci.,* vol. 12, no. 5, pp. 585-592, 2009.

[32] W. N. Grimes, J. Zhang, C. W. Graydon, B. Kachar and J. S. Diamond, "Retinal Parallel Processors: More than 100 Independent Microcircuits Operate within a Single Interneuron," *Neuron,* vol. 65, pp. 873-885, 2010.

[33] I. Cervantes-Sandoval, A. Phan, M. Chakraborty and R. L. Davis, "Reciprocal synapses between mushroom body and dopamine neurons form a positive feedback loop required for

learning," *eLife,* vol. 6, p. e23789, 2017.

[34] S. Takumura, Y. Aso, T. Hige, A. Wong, Z. Lu, C. S. Xu, P. K. Rivlin, H. Hess, T. Zhao, T. Parag, S. Berg, G. Huang, W. Katz, D. J. Olbris, S. Plaza, L. Umayam, R. Aniceto, L. Chang, S. Lauchie, O. Ogundeyi, C. Ordish, A. Shinomiya, C. Sigmund, S. Takemura, J. Tran, G. C. Turner, G. M. Rubin and L. K. Scheffer, "A connectome of a learning and memory center in the adult Drosophila brain," *eLife,* vol. 6, p. e23789, 2017.

[35] A. A. Wanner, C. Genoud, T. Masudi, L. Siksou and R. W. Friedrich, "Dense EM-based reconstruction of the interglomerular projectome in the zebrafish olfactory bulb," *Nat. Neurosci.,* vol. 19, pp. 816-825, 2016.

[36] T. Euler, P. B. Detwiler and W. Denk, "Directionally selective calcium signals in dendrites of starburst amacrine cells," *Nature,* vol. 418, pp. 845-852, 2002.

[37] J. C. Patel, P. Witkovsky, M. V. Avshalumov and M. E. Rice, "Mobilization of Calcium from Intracellular Stores Facilitates Somatodendritic Dopamine Release," *J. Neurosci.,* vol. 29, no. 20, pp. 6568-6579, 2009.

[38] C. Grienberger, X. Chen and A. Konnerth, "Dendritic function in vivo," *Trends Neurosci.,* vol. 38, no. 1, pp. 45-54, 2015.

[39] A. Shapson-Coe, M. Januszewski, D. R. Berger, A. Pope, Y. Wu, T. Blakely, R. L. Schalek, P. H. Li, S. Wang, J. Maitin-Shepard, N. Karlupia, S. Dorkenwald, E. Sjostedt, L. Leavitt, D. Lee, J. Troidl, F. Collman, L. Bailey, A. Fitzmaurice, R. Kar, B. Field, H. Wu, J. Wagner-Carena, D. Aley, J. Lau, Z. Lin, D. Wei, H. Pfister, A. Peleg, V. Jain and J. W. Lichtman, "A petavoxel fragment of human cerebral cortex reconstructed at nanoscale resolution," *Science,* vol. 384, p. eadk4858, 2024.

[40] C. Iadecola, "The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease," *Neuron,* vol. 1, no. 27, pp. 17-42, 2017.

[41] O. Friman, M. Borga, P. Lundberg and H. Knutsson, "Detection and detrending in fMRI data analysis," *NeuroImage,* vol. 22, pp. 645-655, 2004.

[42] Y. Behzadi, K. Restom, J. Liau and T. T. Liu, "A component based noise correction method (CompCor) for BOLD and perfusion based fMRI," *NeuroImage,* vol. 37, pp. 90-101, 2007.

[43] A. P. Holmes and K. J. Friston, "Generalisability, Random Effects & Population Inference," *NeuroImage,* vol. 7, no. 4, p. S754, 1998.

[44] P. Majka, S. Bai, S. Bakola, S. Bednarek, J. M. Chen, N. Jermakow, L. Passarelli, D. H. Reser, P. Theodoni, K. H. Worthy, X. Wang, D. K. Wojcik, P. P. Mitra and M. G. P. Rosa, "Open access resource for cellular-resolution analyses of corticocortical connectivity in the marmoset monkey," *Nat. Comm.,* vol. 11, p. 1133, 2020.

[45] N. Eckstein, A. S. Bates, A. Champion, M. Du, Y. Yin, P. Schlegel, A. K.-Y. Lu, T. Rymer, S. Finley-May, T. Paterson, R. Parekh, S. Dorkenwald, A. Matsliah, S.-C. Yu, C. McKellar, A. Sterling, K. Eichler, M. Costa, S. Seung, M. Murthy, V. Hartenstein, G. S. Jefferis and J. Funke, "Neurotransmitter classification from electron microscopy images at synaptic sites in Drosophila melanogaster," *Cell,* vol. 187, no. 10, pp. 2574-2594.e23, 2024.

[46] R. Tomioka, K. Okamoto, T. Furuta, F. Fujiyama, T. Iwasato, Y. Yanagawa, K. Obata, T. Kaneko and N. Tamamaki, "Demonstration of long-range GABAergic connections distributed throughout the mouse neocortex," *Eur. J. Neurosci.,* vol. 21, pp. 1587-1600, 2005.

[47] S. Higo, K. Akashi, K. Sakimura and N. Tamamaki, "Subtypes of GABAergic neurons project axons in the neocortex," *Front. Neuroanat.,* vol. 3, p. 25, 2009.

[48] C. Mazo, A. Nissant, S. Saha, E. Peroni, P.-M. Lledo and G. Lepousez, "Long-range GABAergic projections contribute to cortical feedback control of sensory processing," *Nat. Comm.,* vol. 13, p. 6879, 2022.

[49] M. Ester, H.-P. Kriegel, J. Sander and X. Xu, "A density-based algorithm for discovering clusters in large spatial databases with noise.," *KDD'96: Proceedings of the Second International Conference on Knowledge Discovery and Data Mini,* pp. 226-231, 1996.

[50] R. L. Buckner, J. R. Andrews-Hanna and D. L. Schacter, "The Brain's Default Network Anatomy, Function, and Relevance to Disease," *Ann. N.Y. Acad. Sci.,* vol. 1124, pp. 1-38, 2008.

[51] S. Namiki, M. H. Dickinson, A. M. Wong, W. Korff and G. M. Card, "The functional organization of descending sensory-motor pathways in Drosophila," *eLife,* vol. 7, p. e34272, 2018.

[52] T. S. Okubo, P. Patella, I. D'Alessandro and R. I. Wilson, "A Neural Network for Wind-Guided Compass Navigation," *Neuron,* vol. 107, pp. 924-940, 2020.

[53] B. El Jundi and M. Dacke, "Insect Orientation: The Drosophila Wind Compass Pathway," *Current Biology,* vol. 31, no. 2, pp. PR83-R85, 2021.

[54] K. E. Coates, S. A. Calle-Schuler, L. M. Helmick, V. L. Knotts, B. N. Martik, F. Salman, L. T. Warner, S. V. Valla, D. D. Bock and A. M. Dacks, "The Wiring Logic of an Identified Serotonergic Neuron That Spans Sensory Networks," *J. Neurosci.,* vol. 40, no. 33, pp. 6309-6327, 2020.

[55] F. Randi, A. K. Sharma, S. Dvali and A. M. Leifer, "Neural signal propagation atlas of Caenorhabditis elegans," *Nature,* vol. 623, pp. 406-414, 2023.

[56] B. E. Brezovec, A. B. Berger, Y. A. Hao, A. Lin, O. M. Ahmed, D. A. Pacheco, S. Y. Thiberge, M. Murthy and T. R. Clandinin, "BIFROST: A method for registering diverse imaging datasets of the Drosophila brain," *Proc Natl Acad Sci USA,* vol. 121, no. 47, p. e2322687121, 2024.

[57] W. Penny, K. Friston, J. Ashburner, S. Kiebel and T. Nichols, Statistical Parametric Mapping: The Analysis of Functional Brain Images, London: Elsevier, 2007.

[58] B. B. Avants, N. J. Tustison, G. Song, P. A. Cook, A. Klein and J. C. Gee, "A reproducible evaluation of ANTs similarity metric performance in brain image registration," *NeuroImage,* vol. 54, no. 3, pp. 2033-2044, 2011.

[59] T. Okuno and A. Woodward, "Vector Auto-Regressive Deep Neural Network: A Data-Driven Deep Learning-Based Directed Functional Connectivity Estimation Toolbox," *Front. Neurosci.,* vol. 15, p. 764796, 2021.

[60] J. Buhmann, A. Sheridan, C. Malin-Mayor, P. Schlegel, S. Gerhard, T. Kazimiers, R. Krause, T. M. Nguyen, L. Heinrich, W.-C. A. Lee, R. Wilson, S. Saalfeld, G. S. Jefferis, D. D. Bock, S. C. Turaga, M. Cook and J. Funke, "Automatic detection of synaptic partners in a whole-brain Drosophila electron microscopy data set," *Nature Methods,* vol. 18, pp. 771-774, 2021.

[61] J. A. Bogovic, H. Otsuna, L. Heinrich, M. Ito, J. Jeter, G. Meissner, A. Nern, J. Colonell, O. Malkesman, K. Ito and S. Saalfeld, "An unbiased template of the Drosophila brain and ventral nerve cord," *PLoS ONE,* vol. 15, no. 12, p. e0236495, 2020.

[62] T.-W. Chen, T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, A. Baohan, E. R. Schreiter, R. A. Kerr, M. B. Orger, V. Jayaraman, L. L. Looger, K. Svoboda and D. S. Kim, "Ultrasensitive fluorescent proteins for imaging neuronal activity," *Nature,* vol. 499, pp. 295-300, 2013.

[63] M. W. Woolrich, B. D. Ripley, M. Brady and S. M. Smith, "Temporal Autocorrelation in

Univariate Linear Modeling of FMRI Data," *NeuroImage,* vol. 14, pp. 1370-1386, 2001.

[64] T. Okuno, N. Ichinohe and A. Woodward, "A reappraisal of the default mode and frontoparietal networks in the common marmoset brain," *Front. Neuroimaging,* vol. 2, p. 1345643, 2024.

Figures

Fig. 1 Comparison between SC and FC, and optimal nuisance factor removal methods. a, Schematic image of the comparison between SC matrices and FC matrices. (left) Three ROI types: Hemibrain anatomical 52 ROIs, SCKm 50 ROIs, and DistKm 50 ROIs. (center) SC matrices (neuron) and SC matrices (synapse) for each ROI type. (right) FC matrices for an average of 8 flies for each ROI type. **b,** Schematic image of calcium imaging data registration. **c,** Swarm plot of ROI volume for each ROI type. **d,** Results of nuisance factor removal method investigation. The vertical axis shows the correlation coefficient, and the horizontal axis shows the FC-SC correlation for each method. Results are plotted for the three ROI types. **e,** The vertical axis shows the averaged AUC of 100 thresholds, and the horizontal axis is the same as **d**. **f,** The vertical axis shows the FC-SC detection & correlation score, and the horizontal axis is the same as **d**.

Fig. 2 Investigating spatial smoothing size, and comparison of FC-SC correlation between mammals and flies. a, (top) Example z-slice of the reduced FDA template with DistKm ROIs from 20 to 1000 ROIs. (middle) SC matrices (neuron) of DistKm ROIs from 20 to 1000 ROIs. (bottom) FC matrices of DistKm ROIs from 20 to 1000 ROIs. **b,** From left to right, FC-SC correlation, FC-SC detection and FC-SC detection & correlation score, respectively. The solid line shows the FC vs. SC matrix (neuron), and the dashed line shows FC vs. SC matrix (synapse) at each ROI count. The vertical axis shows the correlation coefficient, averaged AUC and FC-SC detection & correlation score, respectively. The horizontal axis shows spatial smoothing size (voxels). **c,** Comparison between raw & smoothing (0 to 30 voxels) vs. Poly-tCompCor & smoothing (0 to 30 voxels) at each ROI count. Non-parametric Mann-Whitney U test was performed and Bonferroni correction was applied to correct for the familywise error rate (*** p<1.67e-4). **d,** Relation between FC-SC correlation and ROI count. (left vertical axis) Solid line shows FC-SC correlation (neuron) of SCKm and DistKm ROI type at each ROI count. (right vertical axis) The dashed line shows FC-SC detection (neuron) of SCKm and DistKm at each ROI count. **e,** Swarm plot of ROI volume of SCKm and DistKm at each ROI count.

Fig. 3 Comparison between FlyEM and FlyWire connectome data. a, Bar graph of post-synapse count in hemibrain region from FlyEM and FlyWire connectome data. The vertical axis shows post-synapse count, and the horizontal axis shows confidence thresholds for FlyEM and Cleft score thresholds for FlyWire. **b,** (top) Maximum Z-projection of synapse point cloud of FlyEM (confidence thresholds=0.8 [hereafter, cf0.8]) (bottom) Maximum Z-projection of synapse point cloud of FlyWire (Cleft score threshold=140 [hereafter, sc140]) **c,** Bar graph of post-synapse count in hemibrain primary ROIs. Blue bar shows FlyEM (cf0.8), red bar shows FlyWire (sc140). The vertical axis shows post-synapse count, and the horizontal axis shows hemibrain primary ROIs. **d,** Bar graphs of FC-SC correlation, FC-SC detection, and FC-SC detection & correlation score, respectively. The horizontal axis shows confidence thresholds for FlyEM and Cleft score thresholds for FlyWire. **e,** Bar graph of FC-SC correlation in hemibrain primary ROIs. Blue bars show FlyEM (cf0.8) vs FC (neuron), red bars show FlyWire (sc140) vs FC (neuron). The vertical axis shows FC-SC correlation, and the horizontal axis shows hemibrain primary ROIs. **f,** Bar graph of neurotransmitter rate of output neurons in hemibrain primary ROIs based on FlyWire (sc140) connectome data **[45]**. The horizontal axis shows hemibrain primary ROIs. (DA: dopamine, SER: serotonin, GABA, GLUT: glutamine, ACH: acetylcholine, OCT: octopamine) **g,** Scatter plots of neurotransmitter rate of output neurons vs. FC-SC correlation (neuron) in hemibrain primary ROIs (FlyWire sc140). Each of the six neurotransmitters was compared.

Fig. 4 Quantification of pre- and post-synapse segregation, and the relationship between FC and synapses. a, Unsegregated first (APL_L cell) and second (CT1_R cell) ranking in FlyWire (sc140) connectome data. A red dot shows pre-synapse, a blue dot shows post-synapse. **b,** The PPSSI histogram of all neurons (139,255) in FlyWire (sc140) connectome data. The vertical axis shows neuron count, and the horizontal axis shows PPSSI [0 1]. **c,** Segregated first (OA-AL2b2 cell) and second (OA-AL2b2 cell) ranking in FlyWire (sc140) connectome data. **d,** Schematic image of pre- and post-synapse clustering based on DBSCAN, and calculation of cPPSSI and PPSSI. **e,** Histogram of null SC matrices (blue bar) and extracted SC matrix with PPSSI (0-0.1) (red solid line). Black dotted line shows cumulative distribution function of the normal distribution, brown solid line shows cumulative distribution function of null & extracted SC matrices, and red dotted line shows Bonferroni-corrected p<0.05 threshold. Top shows SC matrix (synapse) total, middle shows FC-SC correlation (neuron), and bottom shows FC-SC correlation (synapse). **f,** Histogram of null SC matrices and extracted SC matrix with PPSSI (0.9-1). **g,** Example of a reciprocal-synapse in the EM image (left) and in 3D (right). **h,** Histogram of reciprocal-synaptic minimum distances in FlyWire (sc140) connectome data. The vertical axis shows reciprocal-synapse pair count, and the horizontal axis shows distance [μm]. **i,** Histogram of null SC matrices and extracted SC matrix for reciprocal-synapses ($\leq 2\mu m$).

Fig. 5 General Linear Model analysis result and Wedge-AVLP GABA neuron. a, Schematic image of GLM analysis. From the top, experimental image and walking trajectory, calcium response function, thresholded velocity, and GLM design matrix, respectively. **b,** 2nd level (group) GLM analysis result from 9 flies. Top shows Z=17 slice of the resized FDA template, and the bottom shows the Z=21 slice. The white arrow indicates the suppressed region during the fly walking task. **c,** Synapse point cloud of the left WAGN in FlyWire (sc140) connectome data. Top shows Z=17 slice of the resized FDA template, and bottom shows the Z=21 slice. **d,** Network graph of the left WAGN, ANs, WPNb tier 1 and tier 2/3 neurons from FlyWire codex (sc50, connection threshold>5). Blue is acetylcholine, ocher is GABA neuron. The small number on the edges shows synapse count. **e,** 3D image of the left WAGN, ANs, WPNb tier 1, tier 2/3 and left ER1 neurons, frontal view. Inside the black square shows the left WAGN, ANs and WPNb tier 2/3 from the back side. **f,** Histogram of closest distance from pre-synapse (to WPNb) to post-synapse (from ANs) in the left WAGN. Horizontal axis shows distance [μm]. Neurotransmitter indicates input ANs to post-synapses. **g,** Example 3D image of pre-synapse (to WPNb) to post-synapse (from ANs). **h,** All pre- and post-synapses of the left WAGN from the front. Color indicates cPPSSI in each synapse. **i,** Histogram of closest distance from pre-synapse to post-synapse in the left WAGN. The horizontal axis shows distance [μm]. Neurotransmitter indicates input neurons to post-synapses.

Data Availability

Whole-brain calcium imaging data and walking behavior data in *Drosophila Melanogaster* can be

downloaded from DANDI web site (https://dandiarchive.org/dandiset/000727/0.240106.0043). FlyWire

connectome data can be downloaded from flywire codex (https://codex.flywire.ai/) and Zenodo web site

(https://zenodo.org/records/10676866). FlyEM connectome data can be downloaded from Janeria's DVID

web site (https://dvid.io/blog/release-v1.2/).

Code Availability

Software and code used in the current study is open source and publicly available from

https://github.com/takuto-okuno-riken/flywalk

Authors' Contributions

T.O. conceived of the presented idea. T.O. developed the theory, performed the computations. T.O.

downloaded and processed fly data. T.O., A.W., H.O. and J.H. discussed the results and contributed to the

final manuscript.

Ethics declarations

All experimental procedures were approved by the Experimental Animal Committee of RIKEN or by the

Experimental Animal Committee of the National Center of Neurology and Psychiatry. The marmosets

were handled by the "Guiding Principles of the Care and Use of Animals in the Field of Physiological

Science" formulated by the Japanese Physiological Society.

Competing interests

The authors declare that they have no competing interests.

Extended Data Figures

Extended Data Fig.1-1 Schematic image of registration process and connectome data transformation in *Drosophila melanogaster.* a, Schematic image of calcium image registration to the resized FDA template. Motion and slice timing correction were conducted using SPM12 [57], and the averaged NIfTI image was registered by ANTs [58] to the resized FDA template. Lastly, 4D NIfTI images were transformed using the transformation information. **b,** Schematic image of synapse points transformation from FlyWire space to the resized FDA template. **c,** Schematic image of synapse points transformation from FlyEM space to resized FDA template.

Extended Data Fig.1-2 Pre- and Post-synapses of pyramidal cell in human temporal cortex layer 3-4. a, (left) Input to (post-synapses) a pyramidal cell in the human temporal cortex, from EM connectome data [39]. (center insert) A slice of human temporal cortex and target pyramidal cell. (right) Output from (pre-synapses) a pyramidal cell in the human temporal cortex, from EM connectome data. Blue dots show post-synaptic sites, Red dots show pre-synaptic sites.

Extended Data Fig.2-1 Investigating spatial smoothing size, and high-pass filtering. a, (top) Example z-slice of reduced FDA template with SCKm ROIs from 20 to 1000 ROIs. (middle) SC matrices (neuron) of SCKm ROIs from 20 to 1000 ROIs. (bottom) FC matrices of SCKm ROIs from 20 to 1000 ROIs. **b,** From left to right, FC-SC correlation, FC-SC detection, and FC-SC detection & correlation score, respectively. Solid lines shows FC vs. SC matrix (neuron), and dashed lines shows FC vs. SC matrix (synapse) for each ROI count. The vertical axis shows the correlation coefficient, averaged AUC, and FC-SC detection & correlation score, respectively. The horizontal axis shows spatial smoothing size (voxels). **c,** Scatter plot of FC-SC correlation in the marmoset left cortex [10] (145 injection × 9862 voxels, r=0.379). **d,** Result of high-pass filtering and nuisance factor removal combination of DistKm ROIs. From left to right, FC-SC correlation, FC-SC detection, and FC-SC detection & correlation score, respectively. The high-pass filter degraded the FC-SC relationship.

Extended Data Fig.3-1 Comparison between FlyEM and FlyWire connectome data. a, (left) A scatter plot of regional post-synapse count between FlyEM (cf0.8) and FlyWire (sc140) in the hemibrain primary 63 ROIs. The Pearson correlation is not so similar, r=0.778, but the Spearman correlation is much more similar, r=0.9. (right) A scatter plot of log10 of the regional post-synapse count between FlyEM (cf0.8) and FlyWire (sc140) in the hemibrain primary ROIs. Pearson correlation shows r=0.919, a higher similarity. **b,** From left to right, FC-SC correlation, FC-SC detection, and FC-SC detection & correlation score, respectively. Solid lines shows FC vs. SC matrix (neuron), and dashed lines shows FC vs. SC matrix (synapse) of FlyEM (cf0.5-0.9) and FlyWire (sc50-140) corresponding Fig.3d. The vertical axis shows the correlation coefficient, averaged AUC, and FC-SC detection & correlation score, respectively. The horizontal axis shows spatial smoothing & nuisance factor removal combinations. **c,** From left to right, SC matrix (neuron) (FlyEM cf0.8), SC matrix (neuron) (FlyWire sc140) and FC matrix (Poly-tCompCor, no smoothing) in the hemibrain primary 63 ROIs. **d,** (right) SC matrix (neuron) result of FlyEM (cf0.8) minus FlyWire (sc140) in hemibrain primary ROIs. (left) SC matrix (synapse) result of FlyEM (cf0.8) minus FlyWire (sc140). FlyEM shows a higher number of synaptic connections.

Extended Data Fig.3-2 In-neuron neurotransmitters and comparison between FlyEM and FlyWire connectome data. a, Bar graph of neurotransmitter rate of input neurons in hemibrain primary 63 ROIs based on FlyWire (sc140) connectome data [44]. The horizontal axis shows hemibrain primary ROIs. (DA: dopamine, SER: serotonin, GABA, GLUT: glutamine, ACH: acetylcholine, OCT: octopamine) **b,** Scatter plots of neurotransmitter rate of input neurons vs. FC-SC correlation (neuron) in hemibrain primary ROIs (FlyWire sc140). Each of the six neurotransmitters was compared. **c,** Comparison between FlyEM and FlyWire APL right neuron (synapse point cloud mask). Red voxels are FlyEM, green voxels are FlyWire, and yellow voxels are both. Sørensen-Dice index (0.716) shows similarity, so neuropil shape is similar between two. **d,** (left) A scatter plot of post-synapse count between FlyEM and FlyWire APL right neuron. Pearson correlation shows r=0.104, so not similar at all. (center) Same as left, but synapse point cloud was spatially smoothed by gaussian kernel (2 voxels). Similarity was increased (r=0.379). (right) Same as center, but smoothed by gaussian kernel (8 voxels). Similarity was increased (r=0.66). These results support that spatial smoothing can absorb inter-individual variability.

Extended Data Fig.4-1 Quantification of pre- and post-synapse segregation and relationship between FC and synapses (FlyEM). a, Unsegregated first (APL_L cell) and second (DPM_R cell) ranked cells in the FlyEM (cf0.8) connectome data. Red dots shows pre-synapses, blue dot shows post-synapses. **b,** The PPSSI histogram of hemibrain neurons (99,644) in FlyEM (cf0.8)

connectome data. The vertical axis shows neuron count, and the horizontal axis shows PPSSI [0 1]. **c,** Segregated first (AOTU019_R cell) and second (DNa02_R cell) ranked cells in the FlyEM (cf0.8) connectome data. **d,** Histogram of null SC matrices (blue bar) and extracted SC matrix with PPSSI (0-0.1) (red solid line). The black dotted line shows the cumulative distribution function of the normal distribution, the brown solid line shows the cumulative distribution function of null &

extracted SC matrices, and the red dotted line shows the Bonferroni-corrected p<0.05 threshold. Top shows the SC matrix (synapse) total, middle shows FC-SC correlation (neuron), and bottom shows FC-SC correlation (synapse). **e,** Histogram of null SC matrices and extracted SC matrix with PPSSI (0.9-1). **f,** Example of a reciprocal-synapse in the EM image (left) and 3D image (right). g, Histogram of reciprocal-synaptic minimum distances in the FlyEM (cf0.8) connectome data. The vertical axis

shows reciprocal-synapse pair count, and the horizontal axis shows distance [μm]. **h,** Histogram of null SC matrices and extracted SC matrix with reciprocal-synapses (≤2μm). **i,** The segregation index histogram of all neurons (139,255) in FlyWire (sc140) connectome data. The horizontal axis shows segregation index [0 1]. **j,** The segregation index histogram of hemibrain neurons (99,644) in the FlyEM (cf0.8) connectome data. The horizontal axis shows the segregation index [0 1].

Extended Data Fig.4-2 Reciprocal synapses of CT1 and APL neurons, and synapse point cloud for several conditions. a, Histogram of reciprocal-synaptic minimum distances of the CT1-R neuron in the FlyWire (sc140) connectome data. The vertical axis shows reciprocal-synapse pair count, and the horizontal axis shows distance [μm]. **b,** Skeleton view of the CT1-R neuron. Red dots are pre-synapses and blue dots are post-synapses of reciprocal synapses (≤ 2μm). **c,** Histogram of reciprocal-synaptic

minimum distances of the APL-R neuron in the FlyWire (sc140) connectome data. **d,** Skeleton view of the APL-R neuron, the same way as b. **e,** (top) Maximum Z-projection of pre & post synapse point cloud within PPSSI (0-0.1) in the FlyEM data (cf0.8). (bottom) Maximum Z-projection of pre & post synapse point cloud within PPSSI (0-0.1) in the FlyWire data (sc140). **f,** The same as in e, but with the condition PPSSI in 0.9-1. **g,** The same as in e, but with the condition of only reciprocal synapses (≤ 2μm).

Extended Data Fig.4-3 Unsegregated and segregated cell ranking examples (FlyWire sc140). a, Unsegregated cell ranking from 3[rd] to 11th in the FlyWire (sc140) connectome data. Red dots shows pre-synapses, blue dots shows post-synapses. **b,** Segregated cell ranking from 3[rd] to 11th.

Extended Data Fig.4-4 Unsegregated and segregated cell ranking (FlyEM cf0.8). a, Unsegregated cell ranking from 3[rd] to 11th in the FlyEM (cf0.8) connectome data. Red dots shows pre-synapse, blue dots show post-synapses. **b,** Segregated cell ranking from 3[rd] to 11th.

Extended Data Fig.5-1 Mechanosensory pathway from Wedge to Ellipsoid Body. a, 3D image of WL-L right, WED016 left, WPNb tier 1, and ER1 left neurons, frontal view. Black square insert shows WL-L right, WED016 left, WPNb tier 1 and ER1 left neurons from the left lateral view. **b,** Network graph of WL-L right, WED016 left, WPNb tier 1 and ER1 left neurons from the FlyWire codex (sc50, connection threshold≥5). Blue is for acetylcholine, ocher for GABA neurons. The small number on the edges shows synapse count.

a Unsegregated rank 1
APL_L (GABA)
PPSSI=0.012

Unsegregated rank 2
CT1_R (GABA)
PPSSI=0.036

● Pre-synapse (output)
● Post-synapse (input)

b Fly EM cf0.8   0.471

Legend: OCT, ACH, GLUT, GABA, SER, DA, Unknown

PPSSI [0 1]

c Segregated rank 1
OA-AL2b2 (ACH)
PPSSI=0.973

Segregated rank 2
OA-AL2b2 (ACH)
PPSSI=0.975

d DBSCAN $\varepsilon = 3\mu m$
Cluster 1
$cPPSSI=0$

$$cPPSSI = \left(\frac{pre - post}{pre + post}\right)^2$$

Cluster 2
$cPPSSI=0.25$

$$(neural)\ PPSSI = \frac{8}{12} \cdot 0 + \frac{4}{12} \cdot 0.25 = 0.083$$

e SC matrix (synapse) total
2373100   *p=0.832*

FC-SC correlation (neuron)
p=0

FC-SC correlation (synapse)
p=6.5e-43

Extracting pre & post synapses with PPSSI (0-0.1)

f SC matrix (synapse) total
p=0.64   28954900

FC-SC correlation (neuron)
p=9.2e-15

FC-SC correlation (synapse)
p=0

Extracting pre & post synapses with PPSSI (0.9-1)

g (90348, 62216, 2280)
102 nm
300 nm

AVLP.28 (GABA)
720575940634854554
AVLP.1861 (ACH)
720575940603595366

h Reciprocal synapse pair count
×10⁶

Legend: OCT, ACH, GLUT, GABA, SER, DA, Unknown

Distance [μm]

i SC matrix (synapse) total
1630010   *p=0.85*

FC-SC correlation (neuron)
p=0

FC-SC correlation (synapse)
p=2.7e-32

Extracting pre & post synapses with reciprocal synapse ($\leq 2\mu m$)

a Walking behavior

Calcium response function

Threshold velocity * [plot 0.04 / 0.02 / 0 over 0–5 sec]

Thresholded velocity
Behavior related calcium response

GLM Design matrix

Calcium response
Nuisance factor (Poly & tcompcor)

b 2nd level GLM result

Z-score

AVLP   AL   GNG   Z17

AVLP   LAL   EB   wedge   saddle   GNG   Z21

c WAGN left synapse point cloud

Synapse count

AVLP   AL   GNG   Z17   WAGN left

AVLP   LAL   EB   wedge   saddle   GNG   Z21   WAGN left

d

ACH   GABA

WED016 (L)   WPNb tier 1

ANs

WAGN left

WPNb tier 2/3

e From the front side

WPNb tier1, tier 2/3

WPNb tier 2/3   WAGN left

ER1s left

WAGN left   ANs   ANs

Pre (to WPNb) : 65
Post (from WPNb) : 250
Pre (to ANs) : 13
Post (from ANs) : 212

From bottom back side

f Closest Distance from pre (to WPNb) to post (from ANs) synapses on WAGN

Synapse count / Distance [μm]

OCT, ACH, GLUT, GABA, SER, DA, Unknown

g From ANs   852.3nm   To WPNb

h WAGN left PPSSI = 0.133

Pre:post = 5577:6511

cPPSSI

i Closest Distance from pre to post synapses on WAGN left

Synapse count / Distance [μm]

OCT, ACH, GLUT, GABA, SER, DA, Unknown

a

Original 4D image

256 × 128 × 49 × 3384 frames (30min)
Around 42 GB, 4D NIfTI image file

SPM12 (MATLAB)

- Motion correction
- Slice timing correction
  (slice order z: 49 to 1)

Motion corrected 4D image

MATLAB script

Make averaged image with sigmoid contrast

100 um

256 × 128 × 49 3D NIfTI image file

Resized FDA template

256 × 128 × 49 3D NIfTI image file

ANTs

Registration to FDA template

Generating GenericAffine.mat & Warp file

ANTs

Apply transform to 4D image

registered 4D NIfTI image file

b

Transform synapse points from FlyWire space to
JRC2018_FEMALE template by **Python** with **navis-flybrains**

flybrains

Flip left and right on template on JRC2018_FEMALE
template

Maximum Z projection

100 um

FlyWire synapse point cloud on JRC2018_FEMALE template

Maximum Z projection

D
R — L
V

100 um

FlyWire synapse point cloud on resized FDA template

c

Transform synapse points from FlyEM space to
JRC2018_FEMALE template by **Python** with **navis-flybrains**

flybrains

Maximum Z projection

100 um

FlyEM synapse point cloud on JRC2018_FEMALE template

Maximum Z projection

Synapse count

FlyEM synapse point cloud on resized FDA template

a

Input (post-synapses)

Output (pre-synapses)

L1

L2

L3

L4

L5

L6

1mm

● Pre-synapse
● Post-synapse

a FlyWire sc140 in-neuron neurotransmitters

Hemibrain primary 63 ROIs

Legend: OCT, ACH, GLUT, GABA, SER, DA, Unknown

b

DA: r=0.12 (p=0.34)

SER: r=-0.14 (p=0.287)

GABA: r=-0.21 (p=0.089)

GLUT: r=-0.25 (p=0.044)

ACH: r=0.41 (p=0.0007)

OCT: r=-0.39 (p=0.0015)

FlyWire sc140 out-neuron neurotransmitter rate

FC-SC correlation [r]

c Sørensen-Dice index=0.716

Maximum Z projection

Z=23/49

Z=44/49

FlyEM

FlyWire

100 um

d

Raw, synapses FlyEM vs. FlyWire r=0.104

Smoothing 2 voxels r=0.379

Smoothing 8 voxels r=0.66

FlyWire synapse count

FlyEM synapse count

a

From the front side

WPNb tier 1

ER1s left

WL-L right

WED016 left

From left side

R · D · V · P · A

D · V · L · M

b

ACH

GABA

720575940636020122

WPNb tier 1

WL-L right

720575940628443127

ER1 (L)

LAL203 (L)

LHPV6q1 (L)

LHPV6q1 (R)

LAL156b (R)

WED016 (L)