Title

Loss of the s^2U tRNA modification induces antibiotic tolerance and is linked to changes in ribosomal protein expression.

Author list

Katherine L Cotten[1], Abigail McShane[2], Peter C. Dedon[2], Thomas J. Begley[3], Kimberly M. Davis[1,4*]

Author affiliations and footnotes

[1] W. Harry Feinstone Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA
[2] Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA
[3] RNA institute and Department of Biological Sciences, University at Albany, State University of New York, Albany, NY, USA
[4] Lead contact

Contact information for corresponding authors

*Correspondence: kdavi140@jhu.edu

Abstract

Stress promotes phenotypic changes in bacteria that allow them to survive antibiotic treatment. This phenomenon, termed antibiotic tolerance, can cause treatment failure, highlighting a need to define bacterial pathways that promote survival. Previously, we found *Yersinia pseudotuberculosis* downregulates *tusB*, a gene involved in modifying tRNAs with s^2U, in response to doxycycline. Here we find that deletion of *tusB* results in loss of s^2U and induces antibiotic tolerance. Using a combination of sequencing-based approaches and analysis of gene codon usage, our data show that loss of s^2U decreases translation of ribosomal proteins. Ribosomal proteins are highly enriched in codons that require s^2U-modified tRNAs for efficient translation, and loss of s^2U results in ribosome pausing at these codons. Our results highlight a previously unknown mechanism of antibiotic tolerance where reduction in ribosomal protein abundance can globally reduce translation, and describes a novel strategy bacteria use to slow growth by modulating s^2U levels.

Introduction

Antibiotic treatment failure is a concerning and multifaceted problem. Major focus has been on antibiotic resistance, which occurs when bacteria acquire a genetic mutation or a genetic element that allows bacteria to grow in the presence of antibiotics[1, 2]. However, bacteria can also develop phenotypic changes in response to stress that allows them to survive antibiotic treatment[1, 2]. This phenomenon is known as antibiotic tolerance when the entire bacterial population responds this way, or antibiotic

persistence when only a subpopulation is recalcitrant to antibiotics[3]. These phenotypic
changes are transient and typically induce a slowed growth response that renders
antibiotics less effective[2]. Once the stress is removed, bacteria resume growth, which is
a major concern for relapsing infection[4]. Growing evidence also suggests that antibiotic
tolerance and persistence contribute to the development of antibiotic resistance[5-8]. We
currently lack strategies to target antibiotic tolerant bacteria, making it important to
understand what pathways promote bacterial survival, so we can leverage this
information to improve antibiotic efficacy.
51
Yersinia pseudotuberculosis (Yptb) is a Gram-negative bacterium in the class of
Gammaproteobacteria that is a natural pathogen of humans and rodents. Yptb has been
used as a model to study multiple aspects of pathogenesis of enteric pathogens[9-13] and
recently has been studied in a mouse model of antibiotic persistence[14]. In a previous
study we investigated the phenotypic response of Yptb to doxycycline treatment and
found expression of the gene *tusB* was significantly downregulated in response to
doxycycline[15]. TusB (2-<u>t</u>hio<u>u</u>ridine <u>s</u>ubunit <u>B</u>) is involved in a sulfur relay system that
transfers a sulfur group onto the wobble position of tRNAs encoding for glutamate (Glu),
glutamine (Gln), and lysine (Lys)[16]. TusB complexes with itself and other Tus proteins
(TusC and TusD homodimers) forming a heterohexamer that mediates the transfer of
sulfur from TusA to TusE[16, 17]. TusE then transfers sulfur to MnmA, which catalyzes the
final 2-thiolation reaction, adding the sulfur group onto the second carbon of the uridine
(U34) in the tRNA anticodon[16]. This modification restricts binding of the tRNA to purines
(A and G), rather than pyrimidines (C or U), thereby increasing translation fidelity and
efficiency.
67
Translation efficiency is generally defined as the rate of protein production per mRNA
molecule[18]. Initiation is primarily thought to be the rate limiting step of translation, but
elongation can be a limiting step as well[19]. During elongation, the speed at which
proteins are translated is impacted by the available pool of tRNAs and competition
between cognate and non-cognate tRNAs[20]. tRNA modifications that stabilize the
interaction between the anticodon and its cognate codon increase translation efficiency
by promoting more rapid decoding. tRNA modifications can thus impact protein
translation depending on which codons are decoded by the modified tRNA, and how
enriched those codons are in a given transcript[21-23]. In yeast, absence of s^2U has been
shown to impact the metabolic state of cells[24] and led to proteotoxic stress due to
improper protein folding during translation[25], but has been postulated to not have a
major impact on translation efficiency, as measured by correlating transcriptional
changes with ribosome footprint changes[24]. In contrast, loss of s^2U seems to impact
translation in *Plasmodium falciparum*, by biasing the translation of certain proteins
depending on their enrichment of lysine codons, impacting artemisinin resistance[26]. In
E. coli, it has been shown that s^2U mutants have delayed expression of specific
proteins, RpoS, Fis, and FtsZ, and impaired cell division[27, 28]. However, it is not known
whether s^2U has a global impact on translation or if this tRNA modification impacts
antibiotic susceptibility in bacteria.
87

Here, we find that deletion of *tusB* and loss of the s^2U modification renders *Yersinia*
tolerant to multiple ribosome-targeting antibiotics, and the RNA polymerase-targeting
antibiotic, rifampicin. We hypothesized that loss of the modification slows translation at
Glu, Gln, and Lys codons. In line with this, we find that ribosomes pause more
frequently at the specific codons that rely on the tRNA modification for proper decoding.
We find that ribosomal proteins themselves are highly enriched in these codons and
downregulated at the protein level, but not the transcript level, indicating they may be
translated less efficiently in the absence of s^2U. Our results support the model that a
reduction in ribosomal proteins is globally reducing translation and is responsible for
slowing bacterial growth and inducing antibiotic tolerance.
98
99 **Results**
100
101 **Deletion of *tusB* results in a complete loss of mnm^{5}s^{2}U and induces changes in**
102 **other modifications.**
103
To understand what impact *tusB* expression has on antibiotic susceptibility, we created
a *tusB* deletion strain (Δ*tusB*). First, to confirm that deletion of *tusB* results in loss of the
s^2U modification, we performed chromatography-coupled tandem quadrupole mass
spectrometric (LC-MS) analysis to detect relative tRNA modification levels in Δ*tusB*
compared to our wild-type (WT) strain (**Figure 1**). We also compared relative tRNA
modification levels in WT +/- doxycycline (dox) treatment to see if downregulation of
tusB would be sufficient to lower tRNA modification levels. We chose to look at tRNA
modification levels in the early log phase of growth to match our RNA-seq timepoint
where we saw downregulation of *tusB* under WT+dox conditions[15]. Additionally, early to
mid-log phase is when the s^2U modification should be highest in WT cells, prior to
nutrient (particularly sulfur) limitation[29].
115
Hierarchal clustering analysis shows that the *tusB* deletion strain clusters together
regardless of doxycycline treatment and WT + dox samples also cluster together
(**Figure 1**). Importantly, as expected, we found a complete loss of s^2U in Δ*tusB* (**Figure**
1, blue box). We saw minimal downregulation of mnm^{5}s^{2}U (average 0.9-fold change
WT dox vs. WT untreated) and cmnm^{5}s^{2}U (average 0.87-fold change WT dox vs. WT
untreated) in response to doxycycline.
122
Interestingly we detected other changes in relative tRNA modification levels in Δ*tusB*
compared to WT, and in response to doxycycline treatment. The sulfur modifications,
s^2C and ms^{2}i6A, were downregulated in response to doxycycline (both in WT and
Δ*tusB*) and somewhat in response to *tusB* deletion alone (**Figure 1**), which could
indicate low sulfur availability[29]. Additionally, we also detected upregulation of many
modifications commonly found on ribosomal RNA (m^{5}C, m^{1}A, m^{3}U, m^{2}G, and m^{66}A).
Our RNA isolation enriches for tRNA based on size and can include smaller ribosomal
RNA fragments if these are abundant.
131

Figure 1. Deletion of *tusB* results in a complete loss of s2U and changes in other tRNA modifications. Log phase bacterial cultures were grown for 2h at 37°C (untreated or treated with 1μg/mL of doxycycline). Small RNA was purified and digested into nucleosides. Changes in relative quantities of modified nucleosides were measured by LC-MS/MS. Each sample is normalized to the average peak area of four biological replicates for WT untreated (WT1-4), see Supplemental Data 1.

Loss of 2-thiolation tRNA modification induces tolerance to specific ribosome and RNA-polymerase targeting antibiotics.

We tested the growth and antibiotic susceptibility of Δ*tusB* and found that it was significantly slower growing and more tolerant to multiple antibiotics including the ribosome-targeting antibiotics, doxycycline and gentamicin, and the RNA polymerase-targeting antibiotic, rifampicin (**Figure 2**). We detected a slight decrease in antibiotic susceptibility to chloramphenicol, although this was not statistically significant. These phenotypes were rescued by integrating the *tusB* gene back into the genome (*tusB* rescue, **Figure 2**). Additionally, to confirm that phenotypes were specific to loss of the tRNA s[2]U modification, we complemented the deletion strain with the *Bacillus subtilis* pathway for 2-thiolation. *B. subtilis* possess an abbreviated version of the sulfur relay system where the cysteine desulfurase, YrvO, transfers sulfur groups directly to MnmA, which catalyzes the final 2-thiolation reaction[30]. As expected, expression of *yrvO-mnmA* restored growth and antibiotic susceptibility of Δ*tusB*, although this was not statistically

significant for doxycycline (**Figure 2**). Chloramphenicol susceptibility was similar to
Δ*tusB* when *yrvO-mnmA* was expressed, indicating that loss of the modification does
not greatly impact chloramphenicol susceptibility. Importantly, phenotypes were only
complemented when a functional copy of *yrvO* was expressed and not when the active
cysteine residue (C325A) of YrvO was mutated or with an empty vector control
(**Extended Data Figure 2**)[30].
161

162
163
Figure 2. Deletion of *tusB* induces slowed growth and antibiotic tolerance to ribosome
and RNA-polymerase targeting antibiotics due to loss of s^{2}U tRNA modification. Log
phase bacterial cultures were grown at 37C and sampled every 2h to enumerate CFUs (colony
forming units). Complemented strains carry a plasmid (pDS145)[30] with the *yrvO-mnmA* open
reading frames under the control of the P$_{ara}$ promoter, and were treated with 0.2% arabinose
throughout the experiment to induce expression of the construct. Cultures were treated with
either 1µg/mL doxycycline, 25µg/mL chloramphenicol, 2µg/mL gentamicin, 2µg/mL rifampicin,
0.1µg/mL ciprofloxacin, or left untreated. Fold change in CFUs are shown by normalizing each
timepoint to the 0h timepoint. Two-way ANOVA with Tukey's multiple comparisons. Asterisks
denote significant p-values as follows: **** $p <0.0001$, *** $p <0.001$, ** $p < 0.01$, and * $p < 0.050$.
Data points represent the mean of three biological replicates, error bars represent standard
deviation (SD). See Supplemental Data 2.
176
Overall, these results indicate that loss of the s^{2}U tRNA modification induces tolerance
to the ribosome and RNA-polymerase targeting antibiotics that were tested.
Interestingly, we observed no difference in susceptibility of Δ*tusB* to the DNA-gyrase
targeting antibiotic, ciprofloxacin, (**Figure 2**) indicating that this tolerance phenotype
may be specific to antibiotics that target translation and transcription. Since transcription

and translation are generally coupled in bacteria, we hypothesized that this might
indicate translation is impacted by loss of s^2U.
184
185 **ΔtusB has a starvation-like metabolic signature and decreased levels of**
186 **ribosomal proteins.**
187
To further understand the impact of loss of s^2U, we performed RNA-seq and mass
spectrometry to define the transcriptome and proteome, respectively, of Δ*tusB*
compared to WT. 259 genes were significantly downregulated, and 182 genes were
significantly upregulated at the transcript level ($p_{adj} < 0.05$ and >2-fold change). At the
protein level, 166 genes were significantly downregulated, and 197 genes were
significantly upregulated (p-value < 0.05 and >2-fold change in protein abundance).
194
KEGG Pathway Enrichment analysis of differentially regulated genes from our RNA-seq
dataset shows that Δ*tusB* upregulates multiple amino acid biosynthesis pathways
(**Figure 3A**). Many of these are also upregulated at the protein level in our mass
spectrometry dataset (**Figure 3B**). We hypothesized this could indicate the stringent
response is being induced by loss of the tRNA modification. *In vitro* studies have shown
the s^2U modification improves tRNA synthetase binding[31] making it plausible that loss of
the modification could result in uncharged tRNAs binding to the ribosome, inducing the
stringent response. However, deleting the alarmone synthetase, *relA*, in the *tusB*
deletion background did not rescue the slowed growth or antibiotic tolerance phenotype
(**Extended Data Figure 3**), indicating the Δ*tusB* phenotype is not linked to the stringent
response.
206
RNA-seq and proteomic analyses generally revealed that Δ*tusB* has a starvation-like
metabolic signature. Δ*tusB* significantly downregulated oxidative phosphorylation
(**Figure 3A, B)** as indicated by downregulation of several genes in the TCA cycle and
multiple subunits of the electron transport chain. This type of metabolic shift is often
associated with antibiotic tolerance and persistence and is consistent with the slowed
growth phenotype of Δ*tusB*.
213
Surprisingly, our proteomic analysis revealed that 37 of the 54 ribosomal protein
subunits were significantly downregulated in Δ*tusB* (**Figure 3B**). Most of these
ribosomal subunits did not show up as hits in our RNA-seq dataset and the few that did
(p<0.05) were not strongly downregulated or upregulated at the transcript level (less
than 2-fold change) (**Extended Data Table 1**). This also indicates that the stringent
response is likely not be responsible for the expression changes in Δ*tusB*, since the
ppGpp alarmone has been shown to repress transcription of ribosomal protein genes[32].
We hypothesized that this difference between our RNA-seq and proteomics datasets
could indicate that the ability to translate these proteins is impaired in Δ*tusB*.
223

224
225
Figure 3. Δ*tusB* has a starvation like metabolic profile and reduced ribosomal protein
levels. A. Log phase bacterial cultures were grown for 2h at 37°C before isolating RNA.
Significant changes in transcript levels between WT and Δ*tusB* were evaluated by DESeq2.
Enriched KEGG Pathways (p-value < 0.05) are shown for downregulated and upregulated
genes. The number of genes associated with each pathway is represented by circle size. Three
biological replicates. **B.** Protein was isolated from bacterial cultures after 2h of growth at 37°C.
Significant changes in protein abundance between WT and Δ*tusB* were evaluated by One-way
ANOVA. The number of significantly regulated genes (p-value < 0.05) associated with KEGG
pathways are shown. Three biological replicates. See Supplemental Data 3.
235
236 **Ribosomes pause more frequently at Glu, Gln, and Lys codons in Δ*tusB***
237

To understand what impact *tusB* deletion has on translation, we performed ribosome
profiling to determine if loss of the s^2U tRNA modification impacts ribosome speed. We
adopted a ribosome profiling approach developed by Mohammad et al.[33, 34], allowing us
to identify ribosome abundance and positioning at single codon resolution. Ribosome
pause scores are calculated by taking the average ribosome density at a single codon
normalized to the average ribosome density across a given gene and averaging this for
all genes in the genome. As expected, the relative ribosome codon occupancy (Δ*tusB*
pause score normalized to WT) was moderately increased at Glu, (GAG and GAA), Gln,
(CAA), and Lys (AAA and AAG), indicating that ribosomes paused more frequently at
these codons in Δ*tusB* (**Figure 4**). No change in ribosome pausing at CAG was
expected since Yptb have one Gln tRNA that is not modified with s^2U (tRNA^{CUG}) that is
likely sufficient for decoding CAG[35]. In contrast, Yptb seem to more heavily rely on
modified tRNAs for decoding Lys (4 tRNA^{UUU} and only one tRNA^{CUU})[35] and Glu (5
tRNA^{UUC} and no tRNA^{CUC})[35], which explains why we see ribosome pausing at both
isoacceptor codons for Lys and Glu.
253

254
Figure 4. Ribosomes pause more frequently at Glutamate, Glutamine, and Lysine codons.
Cultures of bacteria were grown for 5h at 37°C and then flash frozen in liquid nitrogen to stop
translation. Ribosome protected fragments were isolated and sequenced to measure the
abundance of ribosomes at each codon. Pause scores are calculated by taking the average
ribosome density at a codon normalized to the average ribosome density across a given gene
for all genes in the genome. Relative ribosome occupancy (pause score of Δ*tusB* normalized to
the pause score of WT) is shown for each codon. Data points represent the mean of two
biological replicates, error bars represent standard deviation (SD). See Supplemental Data 4.
263
In addition to indicating slower translation, ribosome pausing can also be a sign that
proteins are being mistranslated and/or not properly folded[25]. Since s^2U is important for
stabilizing the interaction between the tRNA anticodon and its cognate codon, we briefly
investigated if the absence of s^2U impacts translation fidelity, with the idea that loss of
s^2U would increase the chances that a near cognate tRNA could bind and incorporate
the wrong amino acid. We detected potential misincorporation of amino acids in a

preliminary LC-MS/MS run, considering the following changes as modifications in
peptides: glutamate to aspartate, glutamine to histidine, and lysine to asparagine. To
understand if these misincorporations occur more frequently due to loss of s^2U, we
measured the relative abundance of modified to unmodified peptides in $\Delta tusB$ vs. WT
(**Extended Data Figure 4A**, **Supplemental Data 8**). Interestingly we detected on
average higher frequency of misincorporation at Glutamine CAA versus Glutamine CAG
codons in $\Delta tusB$ compared to WT (**Extended Data Figure 4B**), consistent with our
finding that ribosomes pause at CAA and not CAG (**Figure 4**). However, with a small
sample size we cannot conclusively say whether misincorporation of amino acids is
more frequent due to loss of s^2U. Additionally, it does not seem that proteins are being
widely mistranslated in $\Delta tusB$, given that we saw no indication of proteotoxic stress in
our differential expression analysis (**Figure 3A, B**).
282
Overall, our results indicate that absence of s^2U impacts ribosome speed. The ribosome
pausing we detected at Glu, Gln, and Lys codons (**Figure 4**) is comparable to what has
been reported in yeast[25, 36], although we detected relatively higher pause scores at Gln
CAA and Lys AAG codons. These values are considered a moderate increase in
ribosome occupancy, indicating that the ribosome is transiently pausing at these sites
rather than stalling. Although transient pausing at one site may not have a great impact
on protein output, we hypothesized that if transcripts were highly enriched in these
codons, this would significantly slow down the ribosome and decrease translation of a
given transcript.
292
293 **Switching Gln codons from CAA to CAG slightly improves translation of GFP**
294
To test whether enrichment of codons with ribosome pausing is sufficient to slow
translation and impact protein output, we created a codon-optimized GFP reporter
construct where all CAA codons (6/8 total Gln codons) are switched to CAG (**Figure
5A**). We focused on Gln codons since we saw a clear increase in ribosome pausing at
CAA, but not CAG (**Figure 4**). As expected, $\Delta tusB$ grew slower than WT (**Figure 5B**).
However, $\Delta tusB$ growth was slightly improved when expressing the CAG construct
(**Figure 5B**), consistent with others have reports that bacterial growth is impaired by
expressing *gfp* with codons that slow the ribosome[20]. Oddly, we saw lower GFP
fluorescence from the CAG construct in the WT strain compared to native GFP, which
was unexpected since CAA and CAG are used equally in protein coding sequences in
Yptb and thus switching these codons was not predicted to bias translation in the WT
strain (**Figure 5C**). However, GFP fluorescence from the CAG construct was not lower
than native GFP in $\Delta tusB$, instead it was slightly higher (**Figure 5C**). As a proxy for GFP
translation, we quantified GFP fluorescence relative to *gfp* transcript levels at 4h (**Figure
5D**). Importantly we can see GFP fluorescence per relative *gfp* transcript level is
trending lower in $\Delta tusB$ vs. WT (p-value = 0.0552) for native GFP, suggesting GFP
translation is lower in $\Delta tusB$. GFP fluorescence per relative *gfp* transcript level was
slightly improved in $\Delta tusB$ with the CAG construct, suggesting that replacing CAA
codons with CAG slightly improves GFP translation, although this was not statistically
significant.
315

316
Figure 5. Switching Glutamine codons from CAA to CAG slightly improves GFP
translation. A. Graphical depiction of model. **B, C, D.** Log phase bacterial cultures were grown
for 8h at 37°C. At 0h cultures were treated with 1mM IPTG to induce expression of *gfp* or left
untreated as a control. **B.** Fold change in OD600 is shown by normalizing each timepoint to the
0h timepoint. **C.** GFP fluorescence is normalized to optical density (OD600) to account for
differences in bacterial density between samples. Values are normalized to the 0h timepoint to
show fold change in GFP fluorescence overtime from the point of IPTG induction. **D.** RNA was
isolated at 4h during the experiment and *gfp* transcript levels were quantified by qPCR. GFP
fluorescence is normalized to *gfp* transcript level as a proxy for GFP protein translation. GFP
fluorescence is normalized to the untreated (uninduced) control to account for background
autofluorescence of the bacteria. *gfp* transcript levels were also normalized to the untreated
(uninduced) control to account for any leaky expression of the construct. See Supplemental
Data 5.

330
331 Overall, these results indicate that translation is lower when s^2U is absent and is
332 partially improved by replacing CAA codons with CAG. We expect that GFP translation
333 was not significantly improved by switching only Glutamine codons since the other
334 codons, Glutamate and Lysine, likely also contribute to the lower translation of GFP in
335 $\Delta tusB$

336
Since GFP protein levels can also be impacted by protein decay and degradation, we
measured GFP decay using a ssrA tag to target the GFP protein to the ClpXP
proteosome[37, 38]. Unstable GFP (*gfp-ssrA*) had a half-life of 57 minutes in WT. In
contrast, unstable GFP (*gfp-ssrA*) had a half-life of 187 minutes in $\Delta tusB$, indicating that
protein degradation is impaired (**Extended Data Figure 5**). Additionally, stable GFP also

had a longer half-life in Δ*tusB* (277 minutes) compared to WT (165 minutes) (**Extended**
Data Figure 5). These findings may be explained by our RNA-seq and proteomic
analysis which indicates that Δ*tusB* downregulates the ClpXP proteosome (*clpX* is
downregulated 1.79-fold by RNA-seq (padj =1x10^8) and 3-fold in protein abundance (p
= 0.0292); *clpP* is downregulated 1.69-fold by RNA-seq (padj = 1.49x10^8) and 3.77-
fold in protein abundance (p=0.00508), see **Supplemental Data 9**). It is unclear why
Δ*tusB* downregulates the ClpXP proteasome, but this suggests that Δ*tusB* is impaired in
protein degradation, allowing us to conclude that the lower GFP fluorescence of Δ*tusB*
(**Figure 5C, D**) is not due to increased degradation of GFP proteins.
351
352 **Lys AAG codon enrichment is highly associated with subset of downregulated**
353 **ribosomal proteins**
354
To look globally at the impact of loss of the s^2U tRNA modification on biasing translation,
we analyzed codon usage for all protein-coding genes in the *Yersinia*
pseudotuberculosis IP2666 genome, with the idea that enrichment of Glu, Gln, and Lys
codons would be highly associated with downregulated proteins if these codons were
biasing translation. Codon Z-scores, detailing relative codon over- or under-use in a
gene (relative to genome averages), were calculated for each codon in each gene by
subtracting the average global codon frequency from the gene codon frequency and
normalizing to the standard deviation of that codon's usage throughout the genome[39].
As a result, codon Z-scores represent how many standard deviations above or below
the global frequency a specific codon is used in a given gene. We performed principal
component (PC) analysis using the codon Z-scores for all genes that were differentially
regulated at the protein level (2-fold change in protein abundance, Δ*tusB* vs. WT, p-
value < 0.05). PC scores are calculated for each gene based on their codon usage and
plotted in **Figure 6A**, revealing a cluster of downregulated proteins along PC1. We
performed principal component regression analysis using the log2fold change in protein
abundance (Δ*tusB* vs. WT) as the dependent variable and found several codons that
significantly contribute to the variance in protein expression. Codons of interest, Glu
GAA (p-value=0.0345), Lys AAG (p-value <0.0001) and Lys AAA (although not
significant, p-value = 0.1757) corresponded with the cluster of downregulated proteins
along PC1 in the lower right quadrant of the loadings plot (**Figure 6B, C**). The codon Z-
scores are particularly high for Lys AAG amongst the downregulated proteins in this
subset. Interestingly 18 of the 21 downregulated proteins are ribosomal proteins,
indicating that enrichment of Lys AAG codons may explain why these proteins are less
abundant even though none were significantly downregulated at the transcript level
(**Figure 3B**). Although we were also expecting Gln CAA codons to bias translation of
proteins, principal component regression analysis indicated they did not significantly
contribute to the variance.
382

383
Figure 6. A combination of Glu, Gln, and Lys codons are highly associated with
ribosomal proteins. Codon Z-scores were calculated for each gene by taking the frequency of
a given codon in that gene and normalizing it the global frequency of that codon and the
standard deviation of that codon. **A.** A data matrix of codon Z-scores for all genes that were
significantly regulated at the protein level in Δ*tusB* vs. WT (2-fold change, p<0.05) were
analyzed by principal component analysis. PC scores are plotted for each gene based on their
codon usage. PC1 explains 11.3% of the variance and PC2 explains 6.4% of the variance. **B.**
Biplot shows the loadings plot, linear association of each codon, with the PC scores of each
gene. **C.** Biplot of genes associated with Lys AAG, Lys AAA, and Glu GAA (PC score >2 and <-
0.5) are graphed separately for clarity. Boxplots of codon Z-scores amongst upregulated (red)
and downregulated (blue) genes in this subset are shown for Glu GAA, Lys AAA and Lys AAG.

One-way ANOVA with Dunnett's T3 multiple comparisons (* denotes $p < 0.05$). **D.** Cumulative
pause predictor scores were calculated by taking the sum of Glu (GAA), Glu (GAG), Gln (CAA),
Lys (AAA), Lys (AAG), codon Z-scores weighted by their ribosome pause score. Cumulative
pause predictor scores are higher on average amongst downregulated proteins (blue) versus
upregulated proteins (red). Mann Whitney test, ** $p < 0.01$. **E.** Ribosomal proteins that are
significantly downregulated at the protein level (purple) have on average higher cumulative
pause predictor scores compared to all other downregulated proteins (blue). Mann Whitney test,
*** $p < 0.0001$. **F.** The log2fold change in protein abundance in $\Delta tusB$ vs. WT (x-axis) is plotted
against the cumulative pause predictor score for all 54 ribosomal proteins. Simple linear
regression shows a negative slope (-3.215) and significant deviation from zero (p-value =
0.0008). See Supplemental Data 6.
406
407 **A combination of Glu, Gln and Lys enrichment likely contributes to decreased**
408 **ribosomal protein levels**
409
Finally, we also looked at the combined impact of all five codons that increased
ribosome pausing, with the idea that enrichment of a combination of Glu, Gln (CAA) and
Lys codons would have more impact than enrichment of a single codon, and that
codons with higher ribosome pause scores would have a greater impact on the
translation of a given protein. We took the sum of the codon Z-scores for Glu (GAA), Glu
(GAG), Gln (CAA), Lys (AAA), and Lys (AAG) for each gene and weighted (multiplied)
each codon Z-score by the average ribosome pause score for that respective codon.
We refer to this calculation as the "cumulative pause predictor score", with the idea that
a higher score will predict a gene will be translated less efficiently due to increased
ribosome pausing along the transcript. Consistent with this idea, we saw significantly
increased cumulative pause predictor scores among proteins that were downregulated
in $\Delta tusB$ compared to upregulated proteins (**Figure 6D**). Ribosomal proteins in
particular had significantly higher cumulative pause predictor scores than all other
downregulated proteins (**Figure 6E**), suggesting that not only enrichment of Lys AAG,
but likely enrichment of a combination of all five of these codons is slowing down the
translation of ribosomal proteins when s^2U is absent. Importantly, we do not see this
same trend in five other codons selected as a control that did not have changes in
ribosome pausing (**Extended Data Figure 6**). Additionally, for the 54 ribosomal proteins,
we see that higher cumulative pause predictor scores correlate with decreased protein
abundance (**Figure 6F**).
430
Overall, these results suggest that translation of ribosomal proteins is significantly
reduced when s^2U is lost due to an enrichment of codons that slow down the ribosome.

Discussion

This work provides evidence for a novel mechanism by which loss of s^2U induces
antibiotic tolerance. It also adds to the growing evidence that modulation of tRNA
modifications can widely impact expression of genes through codon biases in
translation[21, 23, 40]. In other reports, s^2U has been assumed to not have a drastic impact
on protein translation output when only a moderate increase in ribosome pausing is
seen[18,19,25]. However, here we see ribosome pausing highly correlates with genes that
are significantly downregulated at the protein level, but not at the transcript level.

Although pausing at one site alone is likely not sufficient to significantly slow down
translation, our results suggest that enrichment of ribosome pausing sites can lead to
reduced protein levels. Ribosomal proteins seem to be the most impacted by this codon
bias in translation as indicated by their high enrichment in Glu, Gln, and Lys codons and
the significant reduction in ribosomal protein levels. We expect that this reduction in
ribosomal proteins is globally reducing the translational capacity of the cell and is
responsible for inducing tolerance to ribosome and RNA polymerase-targeting
antibiotics.

The s^2U tRNA modification is well conserved among bacteria[16] and ribosomal protein
sequences are also generally conserved across bacterial species, including prominent
ESKAPE pathogens that are highly resistant to antibiotics (**Extended Data Figure 7**).
Therefore, it is plausible that this could be an antibiotic tolerance mechanism used
broadly by bacteria to better survive antibiotics, although this would depend on the
codon usage and tRNA availability for each species.

Future work will be needed to understand what conditions in the host environment are
sufficient to reduce s^2U levels. In our study doxycycline treatment only moderately
lowered s^2U tRNA modification levels (**Figure 1**). However, given the broad
downregulation of the 2-thiolation pathway in response to gentamicin (**Extended Data
Figure 1**), we expect that gentamicin may have a stronger impact on modulating s^2U
levels and promoting survival.

Given that s^2U levels are modulated in response to sulfur availability[29], it is likely that
conditions within the host environment, other than antibiotic stress, also result in
lowered s^2U levels within bacterial pathogens. This highlights that the mechanism we
have defined here may be one used broadly by bacteria to adapt to conditions where it
is beneficial to slow growth, such as low nutrient conditions. Fast growing cells need to
have a high number of ribosomes since protein synthesis is limited by the availability of
free ribosomes [42], and ribosome abundance is linearly correlated with growth rate[43].
However, during slow growth, it is beneficial for bacteria to slow ribosome protein
synthesis since ribosome biogenesis and translation are energetically demanding
processes. Therefore, it is interesting to speculate that the impact of s^2U on ribosomal
protein levels we see here could be a previously unknown mechanism by which bacteria
reduce translation of ribosomal proteins in response to conditions where it is beneficial
to slow growth and save energy.

Surprisingly we detected some changes in other tRNA modifications (**Figure 1**), some
of which are modifications also found on ribosomal RNA (rRNA). Particularly under
starvation conditions, ribosomal RNA is more highly degraded, which could explain why
we detected increased levels of modifications commonly found on rRNA in Δ*tusB* [40,41].
However, it is unlikely that rRNA degradation is responsible for lower ribosomal protein
levels in Δ*tusB*. 16S rRNA levels were not different by qPCR in WT versus Δ*tusB* (**see
Supplemental Data 7**). Additionally, given the reduced capability of Δ*tusB* to degrade
GFP (**Extended Data Figure 5**), along with downregulation of the ClpXP protease, it is
unlikely that ribosomal proteins are being readily degraded in Δ*tusB*. Instead, the level
at which ribosomal proteins are downregulated correlates with the enrichment of codons

that rely on s^{2}U for proper decoding, indicating that these codons bias their translation
in the absence of s^{2}U.

It is important to note that we also saw metabolic changes in Δ*tusB* that could be
contributing to the slowed growth and antibiotic tolerance phenotypes (**Figure 3**).
However, if reduced metabolism were the main contributing factor, we would expect to
broadly see antibiotic tolerance[44-46]. Instead, loss of s^{2}U seems to induce tolerance to
only antibiotics that target transcription and translation (**Figure 2**), consistent with our
model that loss of s^{2}U is impacting the translational capacity of the cell. It is unclear
what drives these metabolic changes in Δ*tusB*, but interestingly, others have noted
similar metabolic changes in response to loss of s^{2}U in *Saccharomyces cerevisiae* and
have postulated that 2-thiolation acts as a signal to the cell that nutrients are low [25,29].
Loss of s^{2}U thus results in a compensatory increase in synthesis of sulfur-containing
amino acids[25]. Additionally, carbon metabolism is altered and shifts away from oxidative
phosphorylation and towards carbohydrate storage[25], similar to what we see in our
transcriptional and proteomic analysis of Δ*tusB* (**Figure 3**).

In summary, here we identified a novel mechanism by which changes in s^{2}U tRNA
modification levels reduce translation of ribosomal proteins and induce slowed growth
and antibiotic tolerance. Future work will be needed to identify conditions sufficient to
lower tRNA modification levels, to further bolster the evidence that modulating s^{2}U
levels is a phenotypic response to stress that promotes bacterial survival.
504
Methods and Materials
Bacterial strains
All experiments were done with the *Yersinia pseudotuberculosis* IP2666 strain[15].
Deletion strains are generated by amplifying 1kb each of the 5' and 3' regions flanking
the gene of interest (homology arms). The 5' homology arm includes the start codon
and 6 base pairs of the beginning of the gene and the 3' homology arm includes 6 base
pairs of the end of the gene and the stop codon. These homology arms are fused
together and cloned into the pSR47s suicide vector[47]. The construct is subsequently
conjugated into WT *Yersinia pseudotuberculosis* (for Δ*tusB* Δ*relA*, the *relA* pSR47s
deletion construct was conjugated into the Δ*tusB Yersinia pseudotuberculosis* strain).
Sucrose selects for bacteria that have complete integration of the insert (two
recombination events)[47]. Deletion of the gene was confirmed by PCR. The *tusB* rescue
strain was generated by amplifying the gene and 1kb upstream and downstream of the
gene from WT chromosomal DNA as a single PCR product. This was inserted into
pSR47s and conjugated into the Δ*tusB Yersinia pseudotuberculosis* strain. Rescued
clones were confirmed by PCR. *yrvO-mnmA* complemented and control strains were
generated by transforming the plasmid constructs (pDS145 *yrvO-mnmA*, pDS220
yrvO(C325A)-*mnmA*, and pARA13)[30] into Δ*tusB Yersinia* by electroporation[47].
523
Antibiotic susceptibility assays
To assay viability of exponential phase bacteria, overnight cultures (16h, at 26°C with
rotation) were sub-cultured into 2mL LB and grown for 2h at 26°C before diluting again
in 2mL LB to a starting bacterial concentration of about 10^4 CFUs/mL. Cultures were
treated with antibiotics at the MIC for bacteriostatic antibiotics (1µg/mL doxycycline,

25µg/mL chloramphenicol) or MBC$_{90}$ concentration for bactericidal antibiotics (2µg/mL
gentamicin, 2µg/mL rifampicin, or 0.1µg/mL ciprofloxacin) or left untreated. Cultures
were grown for 4h at 37°C. Samples were plated for CFUs every 2h. For *yrvO-mnmA*
experiments overnight cultures and subcultures were treated with 0.2% arabinose
throughout the entire experiment to induce expression of the construct or as a control.
534
Small RNA extraction and processing to purify tRNA
Overnight cultures (16h, at 26°C with rotation) were diluted into 50mL LB and grown at
37°C for 2h and then approximately 10^9 CFUs were pelleted. Bacterial pellets were
stored at -80°C before further processing. 1mL of Trizol was added to each cell pellet
and samples were rigorously vortexed for 30 seconds, followed by room temperature
incubation for 5 minutes. 200µL of chloroform was added to each sample followed by
shaking by hand to mix and room temperature incubation for 3 minutes. After
centrifugation at 12000xg for 15mins at 4°C, the aqueous phase was isolated and
brought to a concentration of 30% ethanol to precipitate large RNAs, then applied to a
spin column from the Purelink miRNA isolation kit (Thermo). After centrifugation at
12000xg for 1 minute, the flowthrough containing small RNAs was collected, brought to
an ethanol concentration of 70%, and applied to a second spin column. Samples were
washed twice with the kit's wash buffer and spun dry, followed by elution in nuclease-
free water. A nanodrop was used to determine RNA concentration and purity and quality
was assessed by Agilent 2100 Bioanalyzer Nanochip. 1µg RNA was digested at 37°C
for 6 hours in a 50µL reaction containing 5mM Tris-HCl pH 8.0, 2.5mM MgCl2, 0.1 µg
/mL coformycin (adenosine deaminase inhibitor), 0.1mM deferoxamine (antioxidant), 0.1
mM BHT (antioxidant), 0.083U/µL benzonase, 0.1U/µL calf intestinal alkaline
phosphatase, 0.003 U/µL phosphodiesterase I, and 50nM of 15N-dA internal standard.
554
LC-MS/MS of ribonucleoside modifications
Three technical replicate injections of 200ng for each sample were analyzed by
chromatography-coupled triple-quadrupole mass-spectrometry (LC-MS/MS). A Waters
Acuity BEH C18 column (50 x 2.1 mm inner diameter, 1.7 micron particle size) was
used with an Agilent 1290 HPLC system at 25C, 0.3mL/min flow rate, in a gradient of
Buffer A (0.02% formic acid in water) and Buffer B (0.02% formic acid in 70%
acetonitrile) (**see Supplemental Table 1**). The HPLC was coupled to an Agilent 6495
Triple Quad mass spectrometer with an electrospray ionization source in positive mode
with 200C dry gas temperature, 11L/min gas flow, 20psi nebulizer pressure, 300C
sheath gas temperature, 12L/min sheath gas temperature, 3000V capillary voltage, and
0V nozzle voltage. Product ions were detected using multiple reaction monitoring
(MRM) mode, with retention times and optimal collision energies determined by
synthetic standards (see **Supplemental Table 2**). The sum of 260nm UV signals of the
4 canonical nucleotides, as measured by an in-line UV detector, was used as a
normalization factor.
570
qPCR to measure transcriptional regulation of s^{2}U modification pathway in
response to antibiotics
Cultures of bacteria were grown at 37°C for 2h and treated with the MIC of antibiotics
(1µg/mL doxycycline, 25µg/mL chloramphenicol, or 15µg/mL gentamicin) or left

untreated. Approximately 10^7 CFUs were pelleted and then resuspended in Buffer RLT
(QIAGEN) + β-mercaptoethanol. RNA was isolated using the RNeasy kit (QIAGEN).
DNA was removed using the on-column DNase digestion kit (QIAGEN). RNA was
reverse transcribed using the Protoscript II First Strand cDNA synthesis kit (NEB).
Approximately 30ng of cDNA was used as a template for qPCR using SYBR Green
PCR Master Mix according to the manufacturer's protocol (Applied Biosystems).
Relative transcript levels were measured using the DCT and $2^{-\Delta CT}$ method (using 16S
transcript levels as an endogenous control).
583
RNA-seq
Overnight cultures (16h, at 26°C with rotation) were diluted into 4mL fresh LB and
grown at 37°C for 2h. Approximately 10^7 CFUs were pelleted and then resuspended in
Buffer RLT (QIAGEN) + β-mercaptoethanol. RNA was isolated using the RNeasy kit
(QIAGEN). DNA was removed using the on-column DNase digestion kit (QIAGEN).
Purified RNA was shipped to Novogene (Novogene Corporation Inc), for library
preparation, sequencing, and bioinformatic analysis. Sequencing reads were aligned to
IP2666 *Yersinia pseudotuberculosis* genome assembly GCF_003814345.1. DESeq2
was used to determine significant differences in transcript levels between Δ*tusB* and
WT samples. Genes were considered significantly differentially regulated if there was
greater than 2-fold change in transcript levels and based on an adjusted p-value of
0.05 (p_{adj}, adjusted p-value based on transcript size).
596
Protein extraction
Briefly, overnight cultures (16h at 26°C with rotation) were diluted into 50mL fresh LB
and then grown for 2 hours at 37C. Approximately 10^9 CFUs were pelleted and stored
in -80C. Bacterial pellets were subsequently washed in PBS then resuspended in 4X
cell pellet volume of Bacterial Protein Extraction Buffer (BPER, Thermo-Fisher). After
incubation for 15 minutes at room temperature, samples were centrifuged at 15,000xg
for 5 mins at 4C. Supernatants (soluble fraction) and pellets (insoluble fraction) were
saved at -80C. Pellets were subsequently processed to extract the insoluble fraction by
washing with 1:10 diluted BPER and resuspending in 8x cell pellet volume of Inclusion
Body Solubilization Reagent (Thermo-Fisher). Samples were vortexed for 1 minute and
shaken on a platform shaker for 30 minutes at room temperature before being sonicated
in an ice water bath for a total of 5 minutes (30 seconds on pulse, with 10 seconds off in
between pulses). Samples were then centrifuged at 13,000xg for 15min at 4C and the
supernatant was saved at -80°C.
611
LC-MS/MS to detect relative protein expression changes in Δ*tusB* vs. WT
40μg of protein for each sample was processed for mass spectrometry using SP3
cleanup and on-bead trypsin/LysC digest to remove contaminants and digest samples
into peptides. Peptides were subsequently labelled with tandem mass tags (TMT).
Peptides were separated by HPLC (Thermo-Fisher Vanquish NEO UHPLC) using a
self-packed fused silica column over a 90 minute gradient. Eluted peptides were
sprayed into an Exploris 480 mass spectrometer. Survey scans (full ms) were acquired
on an Orbi-trap within 375-1,500 m/z using a data dependent mode. The full scan MS
was followed by MS/MS for the top 15 peptides in each cycle with dynamic exclusion of

45s. Precursor ions were individually isolated with 0.7 Da and fragmented (MS/MS)
using an HCD activation collision energy of 36. Precursor and fragment ions were
analyzed at a resolution of 120,000 and 30,000, respectively. Reporter ions at resolution
of 45,000 (Turbo TMT). Raw mass spectral data files were searched using Proteome
Discoverer (version 3.1, Thermo Scientific) and the GCF_003814345.1 Yersinia
pseudotuberculosis IP2666pBI genome assembly. Search parameters were 10 ppm
mass tolerance for precursor ions. Fixed modification was carbamidomethylation of
cysteine, TMTpro on Nterm, TMTpro on lysine residues. Variable modification was
oxidation on methionine residues (maximum of 3). A false discovery rate (FDR) of 1%
was used to validate high confidence PSMs. Normalized protein abundances were
averaged across three biological replicates. Protein abundance ratios are calculated by
normalizing the average protein abundance of $\Delta tusB$ to the average protein abundance
of WT. Pairwise comparisons were done with One-way ANOVA using a p-value of 0.05
and 2-fold change in protein abundance as cut offs for significantly differentially
regulated proteins.
636
Detection of misincorporation of amino acids in $\Delta tusB$ vs. WT
A preliminary LC/MS-MS run was done on one biological replicate using the soluble and
insoluble fractions from WT and $\Delta tusB$ samples, similar to above (with TMT labeling).
Raw mass spectral data files were searched in Proteome Discoverer v3.1
(ThermoFisher Scientific) using 3 mutations as variable modifications: Glutamate (E)
changed to Aspartate (D), Glutamine (Q) changed to Histidine (H), and Lysine (K)
changed to Asparagine (N). Peptides were prioritized based on relative abundance in
each sample and selected in order to have a good representation of all the codons we
expected there to be changes at (CAA, GAA, GAG, AAA, no mutations were identified
at AAG), and codons we expected to not see increased changes at in WT compared to
$\Delta tusB$ (CAG). A subsequent LC/MS-MS run (without TMT labeling) was done to
discover additional peptides with potential misincorporations before targeting these
peptides by data-independent acquisition (DIA) to determine the relative abundance of
modified and unmodified peptides in each sample.
10µg of protein for each sample was processed using SP3 buffer exchange and on-
bead trypsin/LysC digest. Peptides were desalted on OASIS HLB uElution plates
(Waters), according to manufacturer's instructions. Peptides were separated by a 100-
minute linear gradient. MS1 scans were acquired in the Orbitrap detector of the Exploris
480 mass spectrometer from 300-1250 m/z with the following settings: RF lens setting
of 50%, 120,000 resolution at 200 m/z, normalized AGC of 101%, maximum injection
time set to Auto, and Easy-IC internal lockmass calibration turned on. Precursor ions in
each window were fragmented by HCD at 30% NCE. DIA product ion (MS2) scans were
acquired in the Orbitrap detector using the following settings: 140–1800 m/z, 30,000
resolution at 200 m/z, normalized AGC of 1000%, and a maximum injection time of 54
ms.
Raw data files were searched with Chimerys against a customized *Yersinia*
pseudotuberculosis UniProt FASTA database (proteome accession UP000255087, 1263
entries), with the addition of sequences of the OmpA protein (UniProt accession
P38399) containing E307D, Q309H, and Q273H substitutions. Search criteria were
tryptic cleavage (maximum 2 missed), peptide length 7-30, peptide charge 2-6, 20 ppm

fragment ion mass tolerance, Cys carbamidomethylation as a fixed modification, and
Met oxidation as a variable modification (max 3/peptide). Peptide identifications were
validated by Chimerys at 1% false discovery rate (FDR) based on an auto-concatenated
decoy database search.
671

Ribosome profiling
Ribosome profiling was done as described in Mohamed et al.[33]. Overnight cultures (16h,
at 26°C with rotation) were diluted into 200mL LB and grown at 37°C for 5 hours before
100mL (Approximately 10^10 CFUs) was flash frozen by spraying into liquid nitrogen.
Frozen droplets of bacterial culture were stored at -80°C before 50g of frozen droplets
were lysed by cryo-mill. Lysate was cleared by centrifugation at 9,000xg at 4°C for
10mins and then ribosomes were pelleted over a sucrose cushion. Approximately 12.5
AU of sample was digested with MNase to remove RNA not protected by ribosomes
and then 70S monosomes were isolated by fractionation over a 10%-50% sucrose
gradient. RNA (or ribosome protected fragments) was isolated by phenol chloroform
extraction and then samples were run on a 15% TBE Urea gel to selectively isolate
fragments within the size range of 15-45nt, which has been previously shown to be the
size range of fragments protected by bacterial ribosomes[34]. cDNA libraries were
prepared, sequenced and reads were mapped to the IP2666 *Yersinia*
pseudotuberculosis GCF_003814345.1 genome assembly. All reads that mapped to
ribosomal RNA, tRNA, and non-coding RNA were excluded. Ribosome occupancy was
measured at the single codon level at the A site of the ribosome. Pause scores were
calculated by taking the average ribosome density at a codon in a given gene and
normalizing it to the average ribosome density across that gene for all genes in the
genome.
692

Codon bias analysis
Codon usage was calculated for each gene in the *Yersinia pseudotuberculosis* IP2666
genome by measuring the frequency of each codon used in each gene, otherwise
known as gene codon frequency. To calculate codon Z-scores, the gene-specific codon
frequency was normalized to the average global usage (all ORFs in genome) of that
codon and the standard deviation of that codon. Codon Z-scores for all up and
downregulated proteins (2-fold change in protein abundance and p-value <0.05), were
put into a data matrix and analyzed by principal component analysis. Principal
component regression analysis was done using the log2fold change in protein
abundance as the dependent variable. To analyze codon usage among genes that were
regulated differently at the protein level versus the transcript level, we analyzed genes
that were, 2-fold upregulated at the protein level and 1-fold or lower downregulated at
the transcript level, and 2-fold downregulated at the protein level and 1-fold or higher
upregulated at the transcript level. All graphing and analysis was performed in
GraphPad Prism.
708

Codon-optimized GFP reporter
The *gfp* DNA sequence, with all CAA codons switched for CAG, was synthesized by
Genewiz. The CAG optimized *gfp* was then cloned into pMMB207267 downstream of
the P$_{tac}$ (IPTG inducible) promoter and then transformed into WT and Δ*tusB Yersinia* by

electroporation. Overnight cultures (16h, at 26°C with rotation) were sub-cultured into
4mL LB +1mM IPTG or untreated and grown for 8 hours at 37°C. Every 2h a 200ul
sample was taken to measure optical density (OD600) and fluorescence (480/520) in a
black walled 96 well plate with plate reader. GFP fluorescence is normalized to OD600
to account for bacterial density. At 4h, 1mL of sample was taken to isolate RNA to
quantify *gfp* transcript levels by qPCR as described above.

Acknowledgments

We thank the Davis lab for their support and feedback on this project. We could not
have done this work without the help of Bob Cole, Josh Smith, and Tatiana Boronina at
the Mass spectrometry core at Johns Hopkins School of Medicine. We could also not
have done the ribosome profiling without the help of Allen Buskirk, Annie Campbell, and
Rachel Green at Johns Hopkins School of Medicine. We also thank Patricia Dos Santos
for her advice and for generously sharing plasmids with us. We would also like to thank
Bob Ernst for his help coming up with the idea for the codon-optimized reporter and
Satoshi Kimura for his advice on quantifying tRNA modification levels. This work was
supported by NIH (NIAID) grants AI154116 and AI175307 to K.M.D. K.L.C. was also
supported by training grant AI007417-26 through NIAID.

Author contributions
Writing – original draft: K.L.C; **Writing- review and editing:** K.M.D;
Conceptualization: K.L.C, K.M.D, P.C.D, T.J.B,; **Investigation:** K.L.C, A.M;
Methodology: T.J.B, P.C.D; **Formal Analysis:** K.L.C, T.J.B, A.M; **Funding Acquisition:** K.M.D

Declaration of interests
The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies
The authors declare that no AI or AI-assisted technologies were used in the writing process.

Extended Data Figures

Extended Data Figure 1. *tusB* is downregulated in response to doxycycline while multiple other genes in the 2-thiolation pathway are downregulated in response to gentamicin. Log phase bacterial cultures were grown at 37°C and treated with either 1µg/mL doxycycline, 15µg/mL gentamicin, or left untreated. After 2h RNA was isolated and relative transcript abundance of *tusA, tusB, tusC, tusD, tusE,* and *mnmA* was quantified by qPCR using 16S transcript levels as an endogenous control. Three untreated biological replicates were done in parallel with doxycycline experiments and three untreated biological replicates were done in parallel with gentamicin experiments. All six biological replicates for untreated samples and the three biological replicates for the treated samples are shown (dots). Values are normalized to the average transcript expression of the three untreated controls that were done in parallel for each experiment. Lines indicate mean relative transcript levels. Kruskal-Wallis test with Dunn's multiple comparisons test, * p< 0.05, * p<0.01.

Extended Data Figure 2. Expression of *yrvO-mnmA* restores growth and antibiotic
susceptibility of Δ*tusB*. Log phase bacterial cultures were grown at 37°C with arabinose
induction (0.2% arabinose) and sampled every 2h to enumerate CFUs (colony forming units). At
the 0h timepoint, cultures were treated with either 1µg/mL doxycycline, 25µg/mL
chloramphenicol, 2µg/mL gentamicin, 2µg/mL rifampicin, 0.1µg/mL ciprofloxacin, or left
untreated. Fold change in CFUs are shown by normalizing each timepoint to the 0h timepoint.
Two-way ANOVA with Tukey's multiple comparisons. Asterisks denote significant p-values as
follows: **** p <0.0001 and * p < 0.050. **A.** Δ*tusB* + pDS145 (pBAD plasmid containing yrvO-
mnmA under the regulation of P$_{ara}$ promoter), three biological replicates same as shown in
Figure 2. **B.** Δ*tusB* + pDS220 (pBAD plasmid yrvO(C325A)-mnmA under the regulation of the
P$_{ara}$ promoter), three biological replicates. **C.** Δ*tusB* + pARA13 (pBAD empty vector control), two
biological replicates.
772
773

774
775
Extended Data Figure 3. Deletion of *relA* does not restore growth or doxycycline
susceptibility of Δ*tusB*. Log phase bacterial cultures were grown at 37°C and sampled every
2h to enumerate CFUs (colony forming units). At the 0h timepoint, cultures were treated with
either 1µg/mL doxycycline or left untreated. Fold change in CFUs are shown by normalizing
each timepoint to the 0h timepoint. Two-way ANOVA with Tukey's multiple comparisons.
Asterisks denote significant p-values as follows: **** p <0.0001. WT and Δ*tusB* doxycycline
replicates are the same as shown in Figure 2.
783

A

Gene	Modification	Codon sequence	Annotated Sequence	q-Value (by Search Engine): CHIMERYS
ompA	WT		[R].GEGQANPVTGNTCDNVKPR.[A]	3.18471E-05
	Gln309 to His	CAA	[R].GEG**H**ANPVTGNTCDNVKPR.[A]	0.001650377
ompA	WT		[R].IGQPAPNLALSQR.[R]	3.18471E-05
	Gln273 to His	CAA	[R].IG**H**PAPNLALSQR.[R]	3.18471E-05
elongation factor Tu	WT		[K].NMITGAAQMDGAILVVAATDGPMPQTR.[E]	3.18471E-05
	Gln98 to His	CAG	[K].NMITGAA**H**MDGAILVVAATDGPMPQTR.[E]	3.18471E-05

B

784
785 **Extended Data Figure 4. Loss of s^{2}U may increase frequency of amino acid**
786 **misincorporation at CAA codons. A.** Table shows representative peptides and the
787 modification (Glutamine Q -> Histidine H) detected in red. **B.** The abundance of each peptide
788 was quantified by LC-MS/MS. The abundance ratio of modified to unmodified peptides in $\Delta tusB$
789 is normalized to the modified/unmodified abundance ratio in WT.
790
791

792
793
Extended Data Figure 5. GFP protein decay is slower in $\Delta tusB$. Overnight cultures were
treated with 1mM IPTG to induce expression of *gfp*. After 16-18h of growth at 26°C, samples
were resuspended in PBS and treated with 50 μg/mL of kanamycin (MIC) to stop translation.
Fluorescence (480/520) was measured every 5 minutes for 120 minutes in a plate reader.
Percent signal of the starting GFP signal (0h) is shown overtime. Half-life is calculated using the

one phase decay equation using 25% GFP signal as a constraint for the plateau (limit of
detection).
801

802
Extended Data Figure 6. Predictor pause scores are not higher among downregulated
proteins or ribosomal proteins for random codons where ribosomes do not pause at. Five
codons that the ribosome does not pause more frequently at in Δ*tusB* were selected to calculate
the cumulative pause predictor score as a control. Cumulative pause predictor scores were
calculated by taking the sum of Ala (GCA), Arg (GAG), Asn (AAC), Cys (TGC), Gly (GGA),
codon Z-scores weighted by their ribosome pause score. A. Cumulative pause predictor scores
of upregulated proteins and downregulated proteins. Mann Whitney test, * $p < 0.05$. B.
Cumulative pause predictor scores of ribosomal proteins and all other downregulated proteins
*** $p = 0.001$
812
813

814
815 **Extended Data Figure 7. Sequences were ribosomes pause frequently at are conserved**
816 **among bacterial species. A.** Representative image of ribosome density across a sequence in
817 the *rpmG* (ribosomal subunit L33) gene. **B.** Amino acid sequences of *rpmG* were pulled from
818 UniProt for multiple different bacterial species. Highlighted are the residues ribosomes are
819 predicted to pause more frequently at.
820
821

Name	Gene	RNA fold change	RNA log2fold change	p_{adj} (RNA)	Protein abundance ratio	Protein log2fold change	p-value (protein)
30S ribosomal protein S1	rpsA	1.36604399	0.45000423	0.00048574	0.348	-1.5228408	0.04431992
30S ribosomal protein S10	rpsJ	NA	NA	NA	0.547	-0.8703873	0.00234407
30S ribosomal protein S11	rpsK	NA	NA	NA	0.418	-1.2584252	0.00172815
30S ribosomal protein S12	rpsL	NA	NA	NA	0.32	-1.6438562	0.02426175
30S ribosomal protein S13	rpsM	NA	NA	NA	0.255	-1.9714308	0.0000995855
30S ribosomal protein S14	rpsN	NA	NA	NA	0.235	-2.0892673	0.00487384
30S ribosomal protein S15	rpsO	0.83619814	-0.2581282	0.10124687	0.206	-2.2792838	0.00757242

30S ribosomal protein S16	rpsP	NA	NA	NA	0.269	-1.8943219	0.01653084
30S ribosomal protein S17	rpsQ	1.37130023	0.45556044	0.02096267	0.275	-1.8624965	0.00587481
30S ribosomal protein S18	rpsR	NA	NA	NA	0.382	-1.3883555	0.11602157
30S ribosomal protein S19	rpsS	NA	NA	NA	0.263	-1.9268653	0.00020635
30S ribosomal protein S2	rpsB	0.73455791	-0.4450458	0.00429957	0.395	-1.3400754	0.20138705
30S ribosomal protein S20	rpsT	NA	NA	NA	0.271	-1.8836352	0.00042677
30S ribosomal protein S21	rpsU	0.68565073	-0.5444782	0.00013317	0.196	-2.3510744	0.00026588
30S ribosomal protein S3	rpsC	NA	NA	NA	0.371	-1.4305089	0.11174822
30S ribosomal protein S4	rpsD	NA	NA	NA	0.463	-1.1109159	0.00152108
30S ribosomal protein S5	rpsE	NA	NA	NA	0.387	-1.3695945	0.06083988
30S ribosomal protein S6	rpsF	NA	NA	NA	0.761	-0.3940316	0.59048579
30S ribosomal protein S7	rpsG	NA	NA	NA	0.338	-1.5649048	0.00030091
30S ribosomal protein S8	rpsH	NA	NA	NA	0.466	-1.1015981	0.00433927
30S ribosomal protein S9	rpsI	NA	NA	NA	0.332	-1.5907449	0.11311814
50S ribosomal protein L1	rplA	NA	NA	NA	0.449	-1.1552126	0.00419147
50S ribosomal protein L10	rplJ	0.7324748	-0.4491423	0.02923479	0.59	-0.7612131	0.07999833
50S ribosomal protein L11	rplK	NA	NA	NA	0.975	-0.0365259	0.97643422
50S ribosomal protein L13	rplM	0.74307233	-0.428427	0.00129496	0.298	-1.7466158	0.00246695
50S	rplN	NA	NA	NA	0.465	-1.1046974	0.00269274

ribosomal protein L14							
50S ribosomal protein L15	rplO	NA	NA	NA	0.388	-1.3658714	0.00252536
50S ribosomal protein L16	rplP	NA	NA	NA	0.485	-1.0439433	0.0004594
50S ribosomal protein L17	rplQ	NA	NA	NA	0.266	-1.9105018	0.00038774
50S ribosomal protein L18	rplR	NA	NA	NA	0.356	-1.4900509	0.0424635
50S ribosomal protein L19	rplS	NA	NA	NA	0.331	-1.5950969	0.00154372
50S ribosomal protein L2	rplB	NA	NA	NA	0.398	-1.3291597	0.00019869
50S ribosomal protein L20	rplT	NA	NA	NA	0.343	-1.5437195	0.00074125
50S ribosomal protein L21	rplU	0.64519681	-0.6322143	0.00013194	0.385	-1.3770696	0.0066521
50S ribosomal protein L22	rplV	NA	NA	NA	0.347	-1.5269924	0.00107999
50S ribosomal protein L23	rplW	NA	NA	NA	0.289	-1.7908586	0.00217311
50S ribosomal protein L24	rplX	NA	NA	NA	0.201	-2.3147326	0.00026907
50S ribosomal protein L25	rplY	0.57933516	-0.7875324	0.00119562	0.66	-0.5994621	0.12471253
50S ribosomal protein L27	rpmA	0.63606651	-0.6528387	0.00336469	0.356	-1.4900509	0.0000180606
50S ribosomal protein L28	rpmB	NA	NA	NA	0.281	-1.831358	0.07549257
50S ribosomal protein L29	rpmC	1.27133672	0.34643557	0.10006984	0.283	-1.821126	0.24651617
50S ribosomal protein L3	rplC	NA	NA	NA	0.25	-2	0.00010761
50S ribosomal protein L30	rpmD	NA	NA	NA	0.275	-1.8624965	0.00819852
50S ribosomal	rpmE	NA	NA	NA	0.518	-0.948976	0.06601131

protein L31							
50S ribosomal protein L32	rpmF	NA	NA	NA	0.688	-0.5395195	0.47978783
50S ribosomal protein L33	rpmG	NA	NA	NA	0.162	-2.6259343	0.00143651
50S ribosomal protein L34	rpmH	0.82309168	-0.2808809	0.03025432	0.546	-0.8730271	0.44592677
50S ribosomal protein L35	rpmI	NA	NA	NA	0.189	-2.4035419	0.00363371
50S ribosomal protein L36	rpmJ	NA	NA	NA	0.294	-1.7661119	0.00088671
50S ribosomal protein L4	rplD	NA	NA	NA	0.311	-1.6850135	0.00139106
50S ribosomal protein L5	rplE	NA	NA	NA	0.474	-1.077041	0.03034906
50S ribosomal protein L6	rplF	NA	NA	NA	0.623	-0.6826959	0.00160103
50S ribosomal protein L7/L12	rplL	0.834444	-0.2610968	0.08072009	0.562	-0.831358	0.50418816
50S ribosomal protein L9	rplI	NA	NA	NA	0.454	-1.1392358	0.07865002

822
823
Extended Data Table 1. Ribosomal protein genes are downregulated at the protein level
in Δ*tusB* (vs WT), but few are differentially regulated at the transcript level. All 54
ribosomal protein genes and their transcript abundance in Δ*tusB* compared to WT (as measured
by RNA-seq) and their protein abundance in Δ*tusB* compared to WT (as measured by mass
spectrometry). NA denotes when there was no significant difference (by p-value) detected by
RNA-seq.
830
831
832 **Supplemental Information**
833
Supplemental Tables 1 and 2 Buffer gradient and compound table for LC-MS/MS
analysis (related to Figure 1)
Supplemental Data 1. tRNA modification analysis (related to Figure 1)
Supplemental Data 2. Antibiotic susceptibility assays (related to Figure 1 and Extended
Figures 2 and 3)
Supplemental Data 3. KEGG Analysis (related to Figure 3)
Supplemental Data 4. Ribosome pause scores (related to Figure 4)

Supplemental Data 5. CAG optimized GFP fluorescence and qPCR analysis (related to Figure 5)

Supplemental Data 6. RNA-seq, mass spectrometry, and codon usage analysis (related to Figures 3 and 6).

Supplemental Data 7. qPCR analysis for expression of s^2U pathway genes with doxycycline and gentamicin treatment (related to Extended Data Figure 1)

Supplemental Data 8. Detected peptides with amino acid misincorporations (related to Extended Data Figure 4)

Supplemental Data 9. GFP decay in WT and Δ*tusB* (related to Extended Data Figure 5)

References

(1) Balaban, N. Q.; Helaine, S.; Lewis, K.; Ackermann, M.; Aldridge, B.; Andersson, D. I.; Brynildsen, M. P.; Bumann, D.; Camilli, A.; Collins, J. J.; et al. Definitions and guidelines for research on antibiotic persistence. *Nature Reviews Microbiology* **2019**, *17* (7), 441-448. DOI: 10.1038/s41579-019-0196-3.

(2) Brauner, A.; Fridman, O.; Gefen, O.; Balaban, N. Distinguishing between resistance, tolerance and persistence to antibiotic treatment. **2016**, *14* (5), 320-330. DOI: 10.1038/nrmicro.2016.34.

(3) Ronneau S; Hill P; S, H. Antibiotic persistence and tolerance: not just one and the same. *Curr Opin Microbiol* **2021**, *64*, 76-81.

(4) Fauvart M; De Groote V; J, M. Role of persister cells in chronic infections: clinical relevance and perspectives on anti-persister therapies. *J Med Microbiol* **2011**, *60*, 699-709.

(5) Levin-Reisman, I.; Ronin, I.; Gefen, O.; Braniss, I.; Shoresh, N.; Balaban, N. Antibiotic tolerance facilitates the evolution of resistance. *Science* **2017**, *355* (6327), 826-839. DOI: 10.1126/science.aaj2191.

(6) Windels, E.; Michiels, J.; Fauvart, M.; Wenseleers, T.; Van den Berg, B.; Michiels, J. Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival and mutation rates. *ISME J* **2019**, *13* (5), 1239-1251. DOI: 10.1038/s41396-019-0344-9.

(7) Santi, I.; Manfredi, P.; Maffei, E.; Egil, A.; Jenal, U. Evolution of Antibiotic Tolerance Shapes Resistance Development in Chronic *Pseudomonas aeruginosa* Infections. *mBio* **2021**, *12* (1). DOI: 10.1128/mBio.03482-20.

(8) Bakkeren E; Huisman JS; Fattinger SA; Hausmann A; Furter M; Egli A; Slack E; Sellin ME; Bonhoeffer S; Regoes RR; et al. *Salmonella* persisters promote the spread of antibiotic resistance plasmids in the gut. . *Nature* **2019**, *573* (7773). DOI: 10.1038/s41586-019-1521-8.

(9) Isberg RR; Voorhis DL; S., F. Identification of invasin: a protein that allows enteric bacteria to penetrate cultured mammalian cells. . *Cell* **1987**, *50* (5). DOI: 10.1016/0092-8674(87)90335-7.

(10) Davis, K. M.; Mohammadi, S.; Isberg, R. R. Community behavior and spatial
regulation within a bacterial microcolony in deep tissue sites serves to protect against
host attack. . *Cell host & microbe* **2015**, *17* (1), 21-31. DOI: 10.1016/j.chom.2014.11.
(11) Barnes, P. D.; Bergman, M. A.; Mecsas, J.; Isberg, R. R. *Yersinia*
pseudotuberculosis disseminates directly from a replicating bacterial pool in the
intestine. . *J Exp Med* **2006**, *203* (6), 1591-1601. DOI: 10.1084/jem.20060905.
(12) Fasciano, A. C.; Dasanayake, G. S.; Estes, M. K.; Zachos, N. C.; Breault, D. T.;
Isberg, R. R.; Tan, S.; Mecsas, J. *Yersinia pseudotuberculosis* YopE prevents uptake by
M cells and instigates M cell extrusion in human ileal enteroid-derived monolayers. *Gut*
microbes **2021**, *13* (1). DOI: 10.1080/19490976.2021.1988390.
(13) Bliska J; Brodsky I; Mecsas, J. Role of the *Yersinia pseudotuberculosis* Virulence
Plasmid in Pathogen-Phagocyte Interactions in Mesenteric Lymph Nodes. *EcoSal Plus*
2021, (2). DOI: 10.1128/ecosalplus.ESP-0014-2021.
(14) Raneses, J. R.; Ellison, A. L.; Liu, B.; Davis, K. M. Subpopulations of Stressed
Yersinia pseudotuberculosis Preferentially Survive Doxycycline Treatment within Host
Tissues. *mBio* **2020**, *11* (4). DOI: 10.1128/mBio.00901-20.
(15) Alvarez-Manzo, H.; Davidson, R.; Van Cauwelaert de Wyels, J.; Cotten, K.;
Nguyen, B.; Xiao, M.; Zhu, Z.; Anthony, j.; van Opijnen, T.; Davis, K. *Yersinia*
pseudotuberculosis doxycycline tolerance strategies include modulating expression of
genes involved in cell permeability and tRNA modifications. *PLoS Pathogens* **2022**, *18*
(5). DOI: 10.1371/journal.ppat.1010556.
(16) Yoshiho, I.; Naoki, S.; Jun-ichi, K.; Akiko, N.; Tsutomu, S. Mechanistic Insights into
Sulfur Relay by Multiple Sulfur Mediators Involved in Thiouridine Biosynthesis at tRNA
Wobble Positions. *Molecular Cell* **2006**, *21* (1), 97-108. DOI:
10.1016/j.molcel.2005.11.001.
(17) Numata, T.; Fukai, S.; Ikeuchi, Y.; Suzuki, T.; Nureki, O. Structural basis for sulfur
relay to RNA mediated by heterohexameric TusBCD complex. *Structure* **2006**, *14* (2),
357-366. DOI: 10.1016/j.str.2005.11.009.
(18) Li, G.-W. How do bacteria tune translation efficiency? *Current Opinion in*
Microbiology **2015**, *24*, 66-71. DOI: 10.1016/j.mib.2015.01.
(19) Plotkin, J. B.; Kudla, G. Synonymous but not the same: the causes and
consequences of codon bias. *Nat Rev Genet* **2011**, *12* (1), 32-42. DOI:
10.1038/nrg2899.
(20) Klumpp, S.; Dong, J.; Hwa, T. On Ribosome Load, Codon Bias and Protein
Abundance. *PLoS One* **2012**, *7* (11). DOI: 10.1371/journal.pone.0048542.
(21) Endres, L.; Dedon, P.; Begley, T. Codon-biased translation can be regulated by
wobble-base tRNA modification systems during cellular stress responses. *RNA Biology*
2015, *12* (6). DOI: 10.1080/15476286.2015.1031947.
(22) de Crécy-Lagard, V.; Jaroch, M. Functions of Bacterial tRNA Modifications: From
Ubiquity to Diversity. *Trends in microbiology* **2021**, *29* (1), 41-53. DOI:
10.1016/j.tim.2020.06.010.
(23) Y.H, C.; McBee M.; Babu I.R.; Hia F.; Lin W.; Zhao W.; Cao J.; Dziergowska A.;
Malkiewicz A.; Begley T.J.; et al. tRNA-mediated codon-biased translation in
mycobacterial hypoxic persistence. *Nature communications* **2016**, *7*. DOI:
10.1038/ncomms13302.

(24) Gupta, R.; Walvekar, A.; Liang, S.; Rashida, Z.; Shah, P.; Laxman, S. A tRNA
modification balances carbon and nitrogen metabolism by regulating phosphate
homeostasis. *eLife* **2019**, *8*. DOI: 10.7554/eLife.44795.
(25) Nedialkova, D.; Leidel, S. Optimization of Codon Translation Rates via tRNA
Modifications Maintains Proteome Integrity. *Cell* **2015**, *161* (7), 1606-1618. DOI:
10.1016/j.cell.2015.05.022.
(26) Small-Saunders, J.; Sinha, A.; Bloxham, T.; Hagenah, L.; Sun, G.; Preiser, P.;
Dedon, P.; Fidock, D. tRNA modification reprogramming contributes to artemisinin
resistance in *Plasmodium falciparum*. *Nature Microbiology* **2024**, *9*, 1483-1498. DOI:
10.1038/s41564-024-01664-3.
(27) Aubee, J.; Olu, M.; Thompson, K. TrmL and TusA Are Necessary for *rpoS* and MiaA
Is Required for *hfq*Expression in *Escherichia coli*. *Biomolecules* **2017**, *7* (2). DOI:
10.3390/biom7020039.
(28) Yildiz, T.; Leimkuhler, S. TusA Is a Versatile Protein That Links Translation
Efficiency to Cell Division in *Escherichia coli*. *Journal of Bacteriology* **2021**, *203* (7). DOI:
10.1128/JB.00659-20.
(29) Edwards, A.; Black, K.; Dos Santos, P. Sulfur Availability Impacts Accumulation of
the 2-Thiouridine tRNA Modification in *Bacillus subtilis*. *Journal of Bacteriology* **2022**,
204 (5). DOI: 10.1128/jb.00009-22.
(30) Black, K.; Dos Santos, P. Abbreviated Pathway for Biosynthesis of 2-Thiouridine in
Bacillus subtilis. *Journal of Bacteriology* **2015**, *197* (11), 1952-1962. DOI:
10.1128/JB.02625-14.
(31) Rodriguez-Hernandez, A.; Spears, J.; Gatson, K.; Limbach, P.; Gamper, H.; Hou,
Y.-m.; Kaiser, R.; Agris, P.; Perona, J. Structural and Mechanistic Basis for Enhanced
Translational Efficiency by 2-Thiouridine at the tRNA Anticodon Wobble Position.
Journal of Molecular Biology **2013**, *425* (20), 3888-3906. DOI:
10.1016/j.jmb.2013.05.018.
(32) Lemke, J. J.; Sanchez-Vazquez, P.; Burgos, H. L.; Hedberg, G.; Ross, W.; Gourse,
R. L. Direct regulation of *Escherichia coli* ribosomal protein promoters by the
transcription factors ppGpp and DksA. *PNAS* **2011**, *108* (44), 5712–5717. DOI:
10.1073/pnas.1019383108
(33) Mohammad F; AR., B. Protocol for Ribosome Profiling in Bacteria. *Bio Protocols*
2019, *20* (9). DOI: 10.21769/BioProtoc.3468.
(34) Mohammad, F.; Green, R.; Buskirk, A. A systematically-revised ribosome profiling
method for bacteria reveals pauses at single-codon resolution. *eLife* **2019**, *6*. DOI:
10.7554/eLife.42591.
(35) Abe, T.; Ikemura, T.; Sugahara, J.; Kanai, A.; Ohara, Y.; Uehara, H.; Kinouchi, M.;
Kanaya, S.; Yamada, Y.; Muto, A.; et al. tRNADB-CE 2011: tRNA gene database curated
manually by experts. *Nucleic Acids Research* **2011**, *39* (1), D210–D213. DOI:
10.1093/nar/gkq1007.
(36) Zinshteyn, B.; Gilbert, W. Loss of a conserved tRNA anticodon modification
perturbs cellular signaling. *PLoS Genet* **2013**, *9* (8). DOI:
10.1371/journal.pgen.1003675.
(37) Gottesman S, R. E., Zhou Y, Sauer RT. The ClpXP and ClpAP proteases degrade
proteins with carboxy-terminal peptide tails added by the SsrA-tagging system. *Genes
Dev* **1998**, *12*, 1338-1347.

(38) Karzai AW; Roche ED; RT, S. The SsrA-SmpB system for protein tagging, directed
degradation and ribosome rescue. *Nature Structural Biology* **2000**, *7*, 449–455.
(39) Doyle, F.; Leonardi, A.; Endres, L.; Tenenbaum, S.; Dedon, P.; Begley, T. Gene- and
genome-based analysis of significant codon patterns in yeast, rat and mice genomes
with the CUT Codon UTilization tool. *Methods* **2016**, *107*, 98-109. DOI:
10.1016/j.ymeth.2016.05.010.
(40) de Crécy-Lagard V, B. Z.; Y, Y.; G, B.; J, B.; JM, B.; PC, D.; P, H.; KR, H.; D., K. Are
bacterial processes dependent on global ribosome pausing affected by tRNA
modification defects? *J Mol Biol* **2025**, *9* (169107). DOI: 10.1016/j.jmb.2025.169107.
(40) Basturea GN; Zundel MA; MP, D. Degradation of ribosomal RNA during starvation:
comparison to quality control during steady-state growth and a role for RNase PH. *RNA*
2011, *17* (2), 338-345. DOI: 10.1261/rna.2448911.
(41) Zundel, M.; Basturea, G.; Deutscher, M. Initiation of ribosome degradation during
starvation in *Escherichia coli*. *RNA* **2009**, *15* (5). DOI: 10.1261/rna.1381309.
(42) J, V.; MA, S.; MD, R.; S, P. Synthesis of proteins in *Escherichia coli* is limited by the
concentration of free ribosomes. Expression from reporter genes does not always
reflect functional mRNA levels. *J Mol Biol* **1993**, *231* (3), 678-688. DOI:
10.1006/jmbi.1993.1319.
(43) RL, G.; T, G.; MS, B.; JA, A.; W., R. rRNA transcription and growth rate-dependent
regulation of ribosome synthesis in *Escherichia coli*. . *Annu Rev Microbiol* **1996**, *50*,
Annu Rev Microbiol. DOI: 10.1146/annurev.micro.50.1.645.
(44) Beam J; Wagner N; Shook J; Bahnson E; Fowler Jr V; Rowe S; B, C. Macrophage-
Produced Peroxynitrite Induces Antibiotic Tolerance and Supersedes Intrinsic
Mechanisms of Persister Formation. *Infection and Immunity* **2021**, *89*.
(45) Rowe S; Wagner N; Li L; Beam J; Wilkinson A; Radlinski L; Zhang Q; Miao E; B, C.
Reactive oxygen species induce antibiotic tolerance during systemic *Staphylococcus*
aureus infection. *Nature Microbiology* **2020**, *5*, 282-290.
(46) Peyrusson F; Varet H; Nguyen T; Legendre R; Sismeiro O; Coppée J; Wolz C;
Tenson T; F., V. B. Intracellular *Staphylococcus aureus* persisters upon antibiotic
exposure. *Nature Communications* **2020**, *11*.
(47) Davidson RK; KM, D. *Yersinia pseudotuberculosis*: Cultivation, Storage, and
Methods for Introducing DNA. *Curr Protoc Microbiol.* **2020**, *59* (1). DOI:
10.1002/cpmc.122. PMID: 33079471.
1007
1008
1009