Multifaceted mRNA analysis using programmed RNA cleavage by

Mucilaginibacter paludis **Argonaute**

Lizhi Liu, Eric A. Hunt, Eric J. Wolf, Ivan R. Corrêa Jr., Erbay Yigit, Sebastian Grünberg

Argonaute (Ago) endonucleases are ubiquitous across all domains of life and, like Cas nucleases, can be programmed by guide oligonucleotides to target and cleave specific nucleic acid sequences. However, unlike Cas enzymes, Agos are not restricted by sequence requirements such as protospacer adjacent motifs (PAMs) enhancing their utility as molecular tools. This study demonstrates diverse applications of the DNA-guided RNA-endonuclease *Mucilaginibacter paludis* Argonaute (Mpa Ago) for *in vitro* mRNA processing, e.g., 3' end polishing, and for mRNA analysis in combination with liquid chromatography tandem mass spectrometry (LC-MS/MS). Applications of the latter include the comprehensive assessment of 5' capping efficiency, 3' poly(A) tail length and composition, one-pot multiplexed cleavage for RNA sequencing, modification detection, and fingerprinting.

Main

Like CRISPR-Cas nucleases, Agos are conserved endonucleases that utilize nucleic acid guides to recognize, bind, and cleave their substrates. However, unlike Cas nucleases, Ago proteins are present across all domains of life[1-3]. In eukaryotes, Agos are loaded with small viral RNAs, endogenous miRNAs, or piRNAs, functioning in RNA interference (RNAi) pathways as components of the RNA-induced silencing complex (RISC)[4-8]. They play crucial roles in antiviral defense, posttranscriptional regulation of gene expression, and suppression of transposon activity[9-11].

Prokaryotic Agos (pAgos) are classified into long and short based on their domain architecture[2, 3]. Long pAgos consist of six domains (N-terminal, L1, PAZ, L2, MID, and PIWI) and share structural similarities with eukaryotic Agos (eAgos). However, some long pAgos are catalytically inactive due to disruptions in their catalytic tetrad,

DEDX (X = D/H/N/K), which coordinate divalent metal cations, often Mg^{2+}, that are

crucial for activity[1, 2, 12-14]. Short pAgos contain only the MID and PIWI domains and lack

the ability to cut nucleic acids due to mutations in the catalytic tetrad[15]. Short pAgos and

catalytically inactive long pAgos often associate with other effector proteins which

function in bacterial defense systems[16]. In contrast, active long pAgos have been

reported to be directly involved in cellular defense against foreign genetic elements as

well as in genome replication and recombination[12, 17-21].

Active pAgos load short guides to form a nucleoprotein complex that

subsequently binds and cleaves complementary target sequences between guide

nucleotides 10 and 11, resulting in cleavage products with 3'-OH and 5'-phosphate

ends[22]. Unlike eAgos, which exclusively bind RNA guides to cleave RNA targets, most

known active long pAgos utilize short DNA guides to target DNA sequences[17, 18, 23-30].

Few pAgos have been reported to use RNA guides to cleave DNA[14, 31-34], and still fewer

using RNA guides to target RNA[33, 35, 36]. Recently, several pAgos have been reported to

use DNA guides to preferentially cleave RNA over DNA[28, 37, 38].

The ability to direct endonuclease activity to a specific location with short nucleic

acid guides makes pAgos highly versatile and valuable molecular tools with many

applications in biotechnology. When compared to Cas nucleases, the lack of sequence

restrictions such as PAMs and the predominant use of short DNA guides versus RNA

guides make pAgos a versatile and economical alternative, especially for *in vitro*

applications. DNA-targeting pAgos are already used in the diagnosis of pathogens and

disease related rare mutations, DNA assembly, super-resolution microscopy, and the

detection of single and double stranded DNA sequences[39]. However, applications that

utilize pAgos to specifically target RNA have not yet been reported.

In this work, we describe RNA-specific applications of MpaAgo, a DNA-guided

pAgo that preferentially targets RNA substrates[37]. We demonstrate that MpaAgo can be

used to efficiently cleave a target RNA at 24 sites simultaneously in a multiplex one-pot

reaction, allowing for subsequent RNA sequencing using intact mass spectrometry with

up to >90% sequence coverage. Custom guide mixtures generating characteristic RNA

fragments enable RNA fingerprinting – a fast and unambiguous way to detect and

identify RNAs by liquid chromatography. We also show that MpaAgo can be used in 5'

cap analysis of mRNAs, allowing for the precise determination and relative

quantification of the cap incorporation in a workflow that can be adapted to virtually any

5' untranslated region (UTR) sequence and can be performed in under one hour. Lastly,

we demonstrate that MpaAgo can be guided to and specifically cleave a sequence

embedded in the poly(A) tail of an *in vitro* transcribed mRNA to generate mRNA

populations with homogenized 3' ends, removing non-templated additions introduced by

RNA polymerases. This also allows for the accurate analysis of the poly(A) tail length

and identification of unwanted potential extensions.

Results

MpaAgo efficiently and specifically cleaves mRNA

To assess the potential of pAgos to manipulate therapeutically relevant RNA

substrates, we sought to determine the specificity and cleavage efficiency of mRNA

substrates by MpaAgo, a DNA-guided endoribonuclease from *Mucilaginibacter paludis*.

MpaAgo has previously been described as a programmable endoribonuclease with

strong guide-mediated substrate specificity that is capable of cleaving RNA substrates

of up to 352 nt in length[37]. We tested site-specific RNA cleavage by MpaAgo on two

different *in vitro* transcribed (IVT) mRNAs: pSG90 IVT RNA, which represents a

shortened version of the replication-competent Sindbis virus genome[40] (1678 nt), and

Fluc (*Firefly luciferase*) IVT RNA (1766 nt). We designed 16 nt long 5'-phosphorylated

guides targeting unique sequences in the pSG90 and Fluc IVT RNAs, spaced 100 or 72

nt apart throughout the complete mRNA sequences. When cleaving the pSG90 RNA

with MpaAgo using single guides in separate reactions, we observed cleavage

efficiencies between 71% and 92% compared to a control reaction containing only

MpaAgo without guide (Fig. 1A). Compared to pSG90 RNA, MpaAgo displayed a higher

cleavage efficiency of Fluc RNA, resulting in near complete cleavage of the substrate

mRNA for most guides (Fig. 1B).

To challenge MpaAgo further, we tested cleavage of a uridine-depleted version

of the human erythropoietin (Epo) IVT RNA with a GC content (excluding the poly(A)

tail) of 59% (pSG95; Fig. S1). The ability to cleave GC-rich substrates is a desirable

property, as uridine depletion and/or modification has been shown to reduce

immunogenicity and is commonly applied to RNA therapeutics design[41]. All but one

guide enabled cleavage with efficiencies close to those observed in the other two tested

mRNAs (between 71% to 89%); the outlier, guide 95.2, only cleaved ~ 50% of the

substrate, possibly due to local secondary structures that were not fully resolved under

the conditions tested.

To assess the possibility of off-target cleavage by MpaAgo, we selected guides

targeting sequences in Fluc RNA and used them to target pSG90 RNA. We did not

detect cleavage, whether in the presence of guides (Fig. S2), further corroborating the

highly specific nature of cleavage. Previous work reported that guide length may affect

cleavage efficiency on short RNA substrates[37]. We repeated cleavage of Fluc IVT RNA

with MpaAgo using 18 nt DNA guides and found similar cleavage efficiencies compared

to that using 16 nt guides, with results varying between 61% and complete cleavage

(Fig. S3).

To expand on the initial characterization of MpaAgo by Li et al.[37], we tested the

efficiency of MpaAgo to cleave long RNAs (> 1000 nt) at various temperatures and

incubation times (Fig. S4A). We observed very little cleavage of the 1678 nt pSG90 IVT

RNA at 4°C even after a 16-hour incubation (Fig. S4B). Increasing the temperature to

24°C and 37°C significantly increased cleavage efficiency, with much of the substrate

being cut after 30 min incubation at 37°C. Cleavage by MpaAgo was highly effective at

50°C, with most of the substrate completely cleaved after 10 min. Importantly, both

MpaAgo and a guide were required for specific cleavage as neither MpaAgo nor the

guide alone induced cleavage of the substrate RNA (Fig. S4B). MpaAgo-independent

degradation of the RNA substrate was observed at 37 or 50°C for 60 min or longer

incubation times (Figs. S4A, right panel and S4B), likely due to Mg^{2+} catalyzed

hydrolysis of phosphodiester bonds[42].

We next tested if the DNA guides could be efficiently removed using DNA

exonucleases, which would allow DNA-free downstream processes. *E. coli* exonuclease

I, which is a 3'-5' exonuclease with strong preference for single stranded DNA, readily

degraded most of the DNA guide from the pAgo reaction (Fig. S5, lane 2). We

hypothesize that the small amount of remaining guide may be bound to and thus

protected by the MpaAgo protein.

Multiplexed cleavage of mRNA

After confirming that pAgos specifically and efficiently cleave RNAs longer than

1000 nt, we proceeded to investigate whether MpaAgo could generate the

corresponding cleavage fragments in a single reaction when provided with a pool of

guides targeting multiple locations along an RNA (Fig. 2A). Consistent with the single

cleavage reactions shown in Fig. 1A, incubating MpaAgo with two guides (90.1 and

90.2) resulted in specific cleavage of the full-length pSG90 RNA at two individual sites

(Fig. 2B). The length of the resulting ~100 nt cleavage fragment (Fig. 2B, lane 4) is

consistent with the distance between the target sequences of the two guides. To

completely cleave the full-length Fluc RNA into unique 72 nt fragments (plus a 29 nt 5'

fragment and an 81 nt 3' fragment), we combined 24 16 nt DNA guides in a single

reaction. Using a ratio of substrate to enzyme to total guide pool of 0.1:1:1, complete

cleavage of Fluc IVT RNA was achieved (Fig. 2C, experiment 2). However, several

fragments of sizes >72 nt were still visible, suggesting that not all guides enabled

complete cleavage of their respective target sequences. The addition of 0.5% bovine

serum albumin (BSA) to the reaction enhanced cleavage efficiency, collapsing the

banding pattern to only three bands besides the unique 5' and 3' end fragments: the

desired 72 nt fragments, one prominent band running at 144 nt, and one faint fragment

with a size of ~ 220 nt (Fig. 2C, experiment 4). The 144 nt and 220 nt bands likely

represent fragments with one or two missing cleavages, indicating that cleavage was

still incomplete. We then tested additional supplements to further increase cleavage

efficiency. In the absence of BSA, the addition of an 1x enhancer mix containing 0.5 M

betaine, 20 mM potassium glutamate, 20% glycerol, and 5% PEG 3350 to the reactions

resulted in a similar fragment distribution as did the addition of 0.5% BSA (Fig. 2C,

experiment 5). Combining the enhancer mix with 0.5% BSA resulted in the complete

cleavage of Fluc mRNA into the programmed RNA fragment sizes (Fig. 2C, experiment

3). Cleaved fragments from the reactions containing the guide pool (experiments 2-5)

were submitted to LC-MS/MS analysis, indicating sequence coverages between 79 and

88% (Fig. 2D and Fig S6). Interestingly, even the reaction without additives that showed

only partial cleavage resulted in 79.2% Fluc mRNA sequence coverage (Fig. 2D and

Fig. S6, experiment 2), suggesting that complete digestion of the substrate mRNA may

not be a prerequisite for high mapping coverage.

Recent data suggest that MpaAgo can also use longer DNA guides to cleave

RNA[37]. We thus decided to use 18 nt guides to test if longer guides could increase the

specificity of MpaAgo targeting and cleavage. While the 16 nt guides were selected to

generate uniform 72 nt RNA fragments, the 18 nt guides were designed to generate

laddering RNA fragments ranging from 69 to 81 nt and thus simplify oligonucleotide

analysis by LC-MS/MS. Under conditions similar to those used with the 16 nt guides,

Fluc and pSG90 IVT RNAs were incubated with MpaAgo and their respective 18 nt

guide pools (Fig. S7A). Consistently, the addition of the enhancer mix significantly

increased RNA site-specific cleavage, collapsing most of the RNA fragment sizes to

around 60-80 nt (Fig. S7A, compare experiments 2 and 3, and experiments 5 and 6).

Sequence coverage for pSG90 and Fluc mRNAs ranged from 78 to 95%, based on LC-

MS/MS analysis (Figs. S7B and S7C).

To assess whether an RNA modification commonly used in therapeutic RNAs

affect MpaAgo cleavage accuracy and efficiency, we incubated it with a Fluc IVT RNA in

which all uridines were replaced with N1-methylpseudouridine (m$^1\Psi$) and the same 18

nt guide pool as above. Gel electrophoresis revealed partially cleaved higher molecular

weight fragments even in the presence of the enhancer mix (Fig. S8A), suggesting that

m$^1\Psi$ may affect the cleavage efficiency. Even though cleavage of the fully m$^1\Psi$-

modified Fluc was not complete, the ensemble of identified cleavage products was

sufficient to generate RNA mapping coverages of around 77% (without enhancer mix)

and 86% (with enhancer mix), as determined by LC-MS/MS analysis (Figs. S8B and

S8C), which were only marginally lower than those of unmodified Fluc (78 and 95%,

respectively).

Overall, these data suggest that MpaAgo can be combined with multiple guides

in a single reaction to cleave long RNA substrates into fragments of defined sizes,

effectively reducing the complexity of the fragment pool (compared to non-specific RNA

chemical cleavage or commonly used RNases with short recognition motifs) and

enabling high sequence mapping coverage using oligonucleotide LC-MS/MS analysis.

We next investigated whether pooled guide DNAs could be employed to

simultaneously target multiple RNA species in a one-pot reaction. The resulting

cleavage products then serve as species-specific molecular fingerprints, which can be

detected by LC-MS/MS to include or exclude RNA species. In reactions containing Fluc

or pSG90 IVT RNA—individually or in combination—we introduced Mpa Ago along with

a guide set designed to produce three distinct 72 nt fragments from Fluc and a single

100 nt fragment from pSG90 (Fig. S9A and S9B). Following cleavage, the reactions

were analyzed by LC-MS/MS. In reactions containing only one RNA species, the

corresponding target fragments from either Fluc (Fig. S9C) or pSG90 (Fig. S9D) were

successfully enriched and identified over unassigned Ago-independent background

fragments. Notably, in the reaction containing both RNA species, specific fragments

derived from both Fluc and pSG90 were identified (Fig. S9E), indicating that Mpa Ago in

combination with multiplexed guide sets enables precise RNA fingerprinting and thus

identification of specific species in mixed RNA populations.

Generation of custom RNA ladders using pAgos

The ability of pAgos to cleave a long RNA substrate into fragments of

predetermined lengths defined by sequence-specific guides opens up the possibility of

generating custom RNA ladders containing fragments of desired lengths. To this end,

we synthesized a 1576 nt IVT RNA featuring six unique sequences of lengths ranging

from 56 to 514 nt that were separated by a specific 20 nt sequence (5'-

…CCUACCGUUAAUGACAGGUC… -3', Fig. S10A). A single guide targeting this 20 nt

region enabled simultaneous MpaAgo-mediated cleavage at sites between the adjacent

sequences, releasing the six sequences from the parental RNA, thus generating a set of

fragments with pre-defined lengths. Like in the reactions with multiplexed guides, the

addition of the enhancer mix increased the cleavage efficiency and resulted in near

complete cleavage of the substrate into a clean and precise RNA ladder, as shown on a

denaturing 6% TBE-urea gel (Fig. S10B).

RNA 5' cap analysis using MpaAgo

The *N7*-methyl guanosine cap structure at the 5' end of eukaryotic mRNAs is

important for nuclear export, stability, and translation of mRNA *in situ*. Determining the

efficiency of mRNA 5' capping is an essential quality control step in modern mRNA

vaccine/therapeutics manufacturing[43-45]. We sought to investigate whether pAgos could

be favorably deployed for mRNA 5' cap analysis. Capillary electrophoresis (CE) was

used to assess cleavage efficiency and determine if pAgos could be programmed to

cleave near the 5' end of an RNA. A 5'-FAM labeled synthetic 33 nt oligoribonucleotide

with a sequence representing the 5' UTR sequence of Fluc mRNA was incubated with

MpaAgo, and with one of four guides designed to induce cleavage 20 nt downstream of

the 5' end with either a complementary (g1A) or a non-complementary (g1C, g1G, g1T)

5'-most nucleotide (Fig. S11A). We found that MpaAgo efficiently and precisely cleaved

at the intended site, regardless of the identity of the 5' nucleotide, resulting in the

expected 20 nt cleavage fragment (Fig. S11B).

Next, we extended this approach to full-length mRNAs. To facilitate capping

efficiency testing in a straightforward and cost-effective gel-based assay, we designed

guides that specifically targeted a site either 29 nt (Fluc mRNA) or 30 nt (pSG90 mRNA)

downstream of their 5' end (Fig. 3A). In both cases, the capped and un-capped

cleavage products showed sufficient difference in mobility on a 15% denaturing

polyacrylamide gel to observe successful capping (Figs. 3B and 3C).

To gain more detailed insight into the nature of the cap structure and estimate

the capping efficiency, we turned to LC-MS/MS for the analysis of MpaAgo-mediated

cleavage products. We targeted the 5' Fluc-UTR using three distinct guides designed to

generate 29 nt (guide Fluc_A1), 26 nt (guide Fluc_A1.2), and 32 nt (guide D891) 5'

fragments, respectively. We observed precise Fluc 5' cleavage, producing fragments of

the expected lengths 29-mer (90%), 26-mer (87%) and 32-mer (73%) (Fig. S12A).

Some low levels of MpaAgo- and guide-independent 5 and 12 nt fragments were also

detected (6-20%), which likely represent abortive transcripts (Fig. S12B). The cleavage

of the pSG90 5' UTR, too, was highly precise, predominantly yielding the intended 30 nt

(D831 guide, >92%) or 21 nt (D892 guide, >94%) 5' fragments (Fig. S12C). Similarly,

MpaAgo- and guide-independent 5 and 12 nt fragments were the only other notable

fragments detected. As expected, uncapped Fluc mRNA predominantly exhibited 5'

triphosphate termini (94%) and fewer diphosphate (5%) ends. In contrast, capped Fluc

mRNA was composed exclusively of the Cap-0 (Me^7Gppp) structure (Fig. 3D). Similarly,

uncapped pSG90 mRNA consisted of 94% triphosphate and 4% diphosphate 5' ends,

whereas capped pSG90 mRNA comprised a mixture of 85% Cap-0, 7% diphosphate,

and 3% triphosphate species (Fig. 3E).

These data show that MpaAgo can be used to successfully and quickly access

the capping status and efficiency – either in a simple, cost-effective gel-based assay or

by precise LC-MS/MS analysis.

Homogenization of RNA 3' ends and poly(A) tail analysis

Undesired 3' extensions by RNA polymerases are a common byproduct of *in*

vitro mRNA synthesis using T7 and other RNA polymerases and were first reported in

the 1980s[46, 47]. Importantly, these 3' extensions have been reported to negatively impact

the efficacy of therapeutic mRNAs[48-50].

We envisioned that MpaAgo could be employed to selectively target and cut a

specific sequence incorporated into the poly(A) tail of an mRNA to remove unwanted

extensions. Such 3' end trimming would generate a mRNA population with homogenous

3' ends (Fig. 4A). We utilized a slightly modified version of the poly(A) tail sequence

utilized in the BNT162b2 vaccine[51], consisting of a stretch of 30 adenosines, followed by

a non-poly(A) sequence (GCATATGACTAAAAAACAT), and 60 terminal adenosines to

a 610 nt IVT RNA sequence (pSG95). Guides D758-D761 were designed to induce

MpaAgo cleavage in the non-poly(A) sequence (Fig. S13). After cleavage, the reaction

products were resolved on a denaturing polyacrylamide gel. We found that guides

D758, D760, and D761 enabled the complete cleavage of the full-length substrate at the

respective target site, while a small fraction of the substrate remained uncleaved in the

reaction with guide D759 (Fig. 4B). These data show that MpaAgo successfully

homogenized the 3' ends of IVT products, efficiently cleaving a target embedded in the

poly(A) tail.

Most current high-resolution methods to access poly(A) tail length and

composition include the release of the poly(A) tail by RNase T1, which cleaves 3' of

guanosines[52]. However, when we attempted to use RNase T1 to measure poly(A) tail

length of *in vitro* transcribed pSG120 mRNA, we found that RNase T1 also cleaved

adenosines in a concentration-dependent matter (Fig. S14), necessitating optimization

of RNase T1 concentration and digestion conditions for reliable poly(A) tail

measurements. Here we show that for a more detailed analysis of poly(A) extensions

and/or measurement of the poly(A) tail length using LC-MS/MS, MpaAgo can either be

programmed to cleave directly upstream of the poly(A) tail (Fig. 4C) or a target

sequence embedded in the tail as described above. The poly(A) tail in these

experiments was slightly shortened to accommodate for the resolution of the LC-

MS/MS, and to assure full capture of non-templated additions by T7 RNA polymerase

without exceeding the range of the LC-MS/MS. MpaAgo was complemented with guides

targeting a sequence upstream of the poly(A) tail (D834) or a sequence in the poly(A)

tail (D758), resulting in 91 or 56 nt 3' cleavage fragments, respectively (Fig. 4D).

Cleavage reactions were performed as described above and the cleavage products

analyzed by intact LC-MS/MS. Reactions with either guide allowed for the sequence

determination of the 3' cleavage fragments. As expected, in addition to the encoded

poly(A) sequence, we observed multiple non-templated addition events by T7 RNA

polymerase. In fact, we found that the majority of fragments had one or more additions

to the encoded sequence (Fig. 4E). A significant fraction of fragments from both

cleavage reactions were found to contain a single additional cytosine, an additional

cytosine with one to three adenines, or one to three adenines without cytosine (Tables

S1 and S2). These findings agree with what has been reported for non-templated T7

RNA polymerase additions[49].

These data show that MpaAgo in combination with a guide targeting the poly(A)

tail can be used to precisely cleave the 3' end of mRNAs, thus creating mRNA

molecules with homogenous ends. The 3' cleavage fragments can be further analyzed

by LC-MS/MS for poly(A) tail length measurement and assessment of non-templated

additions by RNA polymerases.

Discussion

Although the properties of DNA-targeting pAgos have long been acknowledged

and utilized in biotechnological applications, RNA-targeting pAgos have largely been

overlooked, likely due to their relative scarcity and limited characterization. In this study,

we demonstrate that MpaAgo can serve as a powerful molecular tool, significantly

enhancing or simplifying existing analytical workflows, such as RNA sequencing,

fingerprinting, 5' cap, and poly(A) tail analysis. Mpa Ago also enables the generation of

mRNA populations with homogeneous mRNA 3' ends.

We show that pAgos can be programmed with multiplexed guides in a single

reaction to specifically and efficiently cleave at multiple sequences in a target mRNA.

This suggests that pAgos may be used to address currently unresolved challenges in

the analysis, quality control, and sequencing of long RNA molecules and therapeutics,

such as self-amplifying RNAs (saRNAs), which exceed 10 kilobases in length. Current

methods for mRNA analysis, such as digestion with sequence-specific

endoribonucleases like RNase T1 or RNase 4, produce short RNA fragments (typically

10-20 nucleotides in length) that are subsequently analyzed by LC-MS/MS[53-56].

However, fragmenting mRNAs longer than 5 kilobases into such short fragments results

in a complex mixture that is often challenging to deconvolute for full sequence

reconstruction and/or modification detection. A possible solution to this problem is to

increase the cleavage fragment size to a length that is short enough for mass

spectrometry detection analysis, but long enough to reduce the complexity of the

oligoribonucleotide pool for RNA sequencing by LC-MS/MS.

In our initial experiments aimed at targeting unique sites in Fluc IVT RNA using

MpaAgo and multiplexed guides, the guides were designed to produce uniform

cleavage fragments with a length of 72 nt. We found that addition of 0.5% BSA slightly

increased the sequence coverage of the Fluc mRNA sequence, possibly by reducing

non-specific binding of the loaded pAgo complexes to the RNA and other surfaces.

Addition of an enhancer mix significantly increased the cleavage efficiency, and thus the

controlled fragmentation of the RNA target, increasing the overall mapping coverage.

We believe that the crowding agents in the enhancer mix facilitate the targeting of

loaded pAgos to their RNA targets by increasing their proximity and thus their ability to

interact in solution. Interestingly, adding the enhancer mix in combination with 0.5%

BSA led to a reduction of coverage, albeit the gel analysis suggested more efficient

cleavage. We hypothesize that the crowding agents increased non-specific binding of

BSA to the RNA, potentially blocking access for MpaAgo to its target sites. Thus, BSA

may be omitted for MpaAgo cleavage experiments that contain crowding agents (e.g.,

as part of the enhancer mix). Another factor potentially responsible for incomplete

sequence coverage is the uniform length of the cleavage fragments, as fragments with

the same length may not be identifiable/distinguishable due to identical or very similar

masses.

Off-target cleavage of MpaAgo due to mismatched guide binding or slippage may

also negatively affect fragment analysis. We thus set to design guides that a) were

comprised of 18 nt instead of 16 nt to increase specificity, and b) induced cleavage into

fragments of various lengths, sorted in size-bins ranging from 69 to 81 nt in length, to

facilitate data deconvolution. While the band pattern on the gel suggested less efficient

cleavage of the Fluc mRNA with the 18 nt guides (compare Fig. 2C, experiment 1 and

Fig. S7A, experiment 2), the sequence coverage in reactions without the enhancer mix

was comparable (~ 79%). Reactions containing the enhancer mix reproducibly achieved

slightly higher coverage (~95%) using 18 nt guides than that of 16 nt guides (~ 88%),

suggesting that increasing guide specificity and cleavage of the substrate RNA into

binned lengths may be beneficial for RNA sequence analysis. Upon closer inspection of

the recovered fragments, we noticed that we were unable to detect fragments spanning

Fluc mRNA positions 1056 to 1128 (16 nt guides) and 1064 to 1145 (18 nt guides) in

any of our experiments. RNA structure modelling using MXfold2[57] and RNAfold[58] of the

respective sequence and flanking regions did not indicate any obvious structural

components that may inhibit pAgo binding (data not shown). The presence of the

flanking fragments indicate that successful cleavage must have occurred. Sequencing

of the IVT template ruled out the possibility of base mutations that could alter the mass

of the expected fragments, making them unidentifiable by LC-MS/MS. We hypothesize

that the missing fragments may be too similar in mass to other cleavage products that

cannot be isolated in the data analysis. We are aware that the RNA sequences tested in

this study are shorter than those of saRNAs, and that the maximal number of combined

guides tested was 24. However, our results suggest that this application could

successfully be used with clinically relevant RNA substrates, especially since full

replacement of uridine with $m^1\Psi$, a replacement often found in therapeutic mRNAs to

minimize the host immune response, only had a moderate effect on the sequence

coverage.

pAgos in combination with a smaller pool of multiplexed guides that cleave RNA

substrates to generate a unique, characteristic, and identifiable fragment pattern can be

used to fingerprint RNAs. This may be useful to confirm the presence of certain RNAs in

an RNA mixture (e.g., to analyze the RNA molecule distribution in RNA pools packaged

in liquid nanoparticles) or quick detection of a specific target RNA in a biological sample

for diagnostic workflows. A similar approach was recently developed using the HD3

deoxyribozyme to fragment RNA[59]. Due to its single turnover character, this method

requires multiple rounds of cleavage and re-annealing of the HD3 deoxyribozyme pool,

which requires up to 24 h total digestion time for complex substrates. This is much

slower than the 15 min of cleavage time in the MpaAgo reaction. In addition, the target

site choice is limited to a particular set of trinucleotide motifs at which HD3 cleaves, a

limitation that does not apply to MpaAgo cleavage.

In eukaryotic mRNA, the *N*7-methyl guanosine cap structure is a 5' end

modification that is important for nuclear export, RNA stability, and translation. Thus,

effective capping of mRNA is an essential part of mRNA-based vaccine and therapeutic

synthesis. Current methods of cap analysis combine LC-MS/MS analysis of a defined 5'

end fragment of a mRNA that was generated by selective cleavage using either *trans*

acting deoxyribozymes[60-62], and ribozymes[63], or a DNA probe-directed approach using

either the sequence-nonspecific Ribonuclease H (RNase H)[64], or sequence specific

RNases (e.g., RNase 4, MC1, and RNase T1[65, 66]) to cleave the 5' UTR at a specific

position and release the cap-containing 5' fragment.

Our data show that guided MpaAgo can efficiently release the cap-containing 5'

termini of mRNAs, enabling the analysis of mRNA capping efficiency and permitting the

determination of the cap structure comparably to other approaches, such as RNase H

or sequence-specific RNases. Contrary to non-guided endoribonucleases, pAgos

exhibit lower turnover[23, 30], thereby requiring comparatively high enzyme concentrations

for effective activity. However, there are significant advantages of using MpaAgo for cap

analysis. First, it eliminates the need for a blocking antisense oligo or the

presence/incorporation of an endoribonuclease recognition sequence for precise

cleavage. Successful 5' fragment generation by sequence-specific endoribonucleases

(e.g., RNase T1, RNase 4) requires blocking of all the respective target sites directly

downstream of the 5' end by hybridization of a DNA oligo. In addition, the release of the

protected fragment is dependent on the presence of a recognition site directly

downstream of the duplex. Second, sequence-specific RNases will fragment the entire

mRNA downstream of the hybridized probe, in some workflows necessitating the

enrichment of the 5' cleavage fragments. Moreover, the additional step of hybridizing of

a blocking oligo is not needed with MpaAgo. The use of MpaAgo for 5' end analysis

may also allow for the detection of endogenous 5' ends with poorly defined transcription

start sites or post-transcriptional modifications[67], in which case the use of a blocking

oligo may not be sufficient to protect the heterogeneous 5' ends. Additionally, the

transient alignment of the short DNA guide with the RNA target at 50°C may mitigate

the influence of RNA structure in the 5' UTR sequence, reducing bias and increasing

binding/cleavage efficiency in MpaAgo workflows.

A potential caveat of cap analysis with sequence-specific RNases is that it

generates RNA/DNA duplexes with a 3'-phosphate and/or 2',3'-cyclic-phosphate

overhang. In contrast, the release of single-stranded MpaAgo cleavage products

facilitates subsequent 3' end modifications of the fragments, such as fluorescent

labeling through fill-in extension using DNA polymerases as in RNase H-based cap

analysis methods[64], or enzymatic ligation of the fragments to other nucleic acid

sequences.

In addition to requiring a significantly shorter reaction time compared to other

workflows (15 min vs. 60 min), MpaAgo also facilitates 5' cap analysis by producing

cleavage products that do not need to be enriched for LC-MS/MS analysis. In fact, the

generation of only two cleavage products (a short 5' fragment and a long 3' fragment)

allows for an easy and cheap capping efficiency assessment using PAGE, making this

analysis accessible to laboratories without LC-MS/MS capabilities. Recently, a group

reported the use of a nucleolytic toxin from a type III toxin/antitoxin system that

recognizes a specific 5 nt target sequence to cleave and subsequently analyze 5'

mRNA ends for cap detection[68]. Theoretically, mRNA cleavage with Toxin III

endoribonucleases also generates only two fragments that enable gel-based capping

assessment, and they do not require a DNA guide or oligo for sequence specific

cleavage. However, this approach necessitates sequence optimization of the mRNA to

remove other potential Toxin III cleavage sites, and – more importantly - a change of the

5' UTR sequence to incorporate the target sequence, which may be limiting for certain

mRNA constructs. In addition, type III toxins possess a relative specificity rather than an

absolute specificity, meaning that they are prone to off-target cleavage at similar

sequence motifs[69]. This can result in the generation of additional fragments, which may

interfere with cap analysis.

One of the shortcomings of using RNase H for cap analysis is its heterogeneous

cleavage that results in additional fragments at positions directly upstream or

downstream of the desired cleavage site. Optimization of the DNA oligo sequence is

often required to reduce off-target cleavage[64]. We show that there is only negligible off-

target cleavage by MpaAgo as tested with four different guides on two different mRNAs

by CE and LC-MS/MS.

We also show that site-directed cleavage by MpaAgo is a superior tool when

analyzing RNA 3' ends. The 3' poly(A) tail directly affects mRNA stability and translation

efficiency, making it an essential component of native and therapeutic mRNAs. While

several low-resolution methods exist that can identify the presence of a poly(A) tail,

there are only limited options for detailed analysis of the poly(A) tail length and

composition. In addition to next generation sequencing, other high-resolution methods

include digestion of the RNA substrate with RNase T1, which cleaves after every

guanine, followed by capillary gel electrophoresis or LC-MS/MS[70, 71]. Site-specific

removal of the poly(A) tail by MpaAgo for subsequent analysis addresses several

shortcomings of digestion with RNase T1. Modern mRNA therapeutics and vaccines

incorporate G-containing, non-A islands into the poly(A) tails to increase RNA

polymerase accuracy (e.g., Pfizer BNT162b2). Guanines in these sequences will be

targeted by RNase T1 and generate a diverse pool of poly(A) fragments which

complicates analysis. In addition, we show here that RNase T1 can non-specifically

degrade poly(A) regions in a concentration dependent manner. MpaAgo on the other

hand can be directed to precisely cleave either the complete poly(A) tail or fragments

thereof when targeting the non-A islands. This generates defined cleavage products,

which abolishes the need for subsequent purification or enrichment. Furthermore, it

simplifies the subsequent analysis by enabling noise-free, single-nucleotide resolution

analysis of the poly(A) tail via LC-MS/MS.

In addition, MpaAgo uniquely enables 3' end polishing of IVT RNA products to

generate RNA populations with homogenous 3' ends. Non-templated addition of

nucleotides is a known feature of RNA polymerases *in vitro* that can lead to unwanted

consequences (e.g., providing a source for double-stranded RNA (dsRNA) formation).

Targeted trimming and thus homogenizing of 3' RNA ends with MpaAgo may prevent

and/or significantly reduce the formation of immunogenic dsRNA in RNA manufacturing,

addressing a main concern in RNA therapeutics development. Since the MpaAgo

enzyme, the short DNA guide, and the cleaved RNA extension can easily be removed

from the homogenous RNA pool by proteinase K/DNase I treatment, and/or a sizing

column, this workflow could be incorporated into large scale RNA synthesis workflows.

Taken together, we show that programmed cleavage of mRNA by MpaAgo

enables faster, more precise, and straightforward workflows for multiple important RNA

analysis workflows, like 5' capping analysis, RNA sequence/modification mapping, and

poly(A) tail analysis – all essential for therapeutic RNA quality assessment. In addition,

we show that MpaAgo can be used to generate RNA populations with homogeneous 3'

ends, which may help to minimize the presence of immunogenic dsRNA and thus

address one of the most critical hurdles in therapeutic RNA production.

Methods

In vitro RNA cleavage assay

200 ng of Fluc RNA (~1.8 kb, ~0.37 pmol), a shortened version of the pTsin

saRNA (pSG90; ~1.7 kb, ~0.37 pmol), or Epo RNA with a modified sequence (pSG95;

~0.7 kb, ~0.86 pmol), all synthesized using *in vitro* transcription (IVT) (New England

Biolabs, #E2040) according to the manufacturer's instructions, were cleaved with

MpaAgo in a 10 μL reaction at 50°C for either 15 minutes (single and double cleavages,

1:5 substrate: Ago molar ratio) or 30 minutes (multiplexing, 1:10 substrate: Ago ratio) in

1x ThermoPol® reaction buffer (New England Biolabs #B9004S). The Ago:guide molar

ratios were 1:5 for single and double cleavages, 1:1 for multiplexing experiments, and

1:10 for cleaving polycistronic RNA. DNA guides were pooled using the Echo acoustic

liquid handler (Beckman Coulter) and phosphorylated using T4 polynucleotide kinase

(New England Biolabs #M0201S) according to the manufacturer's protocols. For

multiplexing experiments and when indicated, a 1x enhancer mix containing 0.5 M

betaine, 20 mM potassium glutamate, 20% glycerol, and 5% PEG3350 was added to

the reactions. BSA was added to 0.5% (v/v) when indicated. Removal of DNA guides

was performed by adding 20 U of exonuclease I (New England Biolabs, #M0293),

followed by incubation at 37°C for 15 minutes. Cleavage experiments were stopped by

the addition of 0.08 U of proteinase K (New England Biolabs #P8107S) at room

temperature for 5-10 minutes. After adding RNA loading dye (New England Biolabs,

#B0363), cleaved RNAs were resolved by gel electrophoresis on 6% TBE-urea gels

(Thermo Fisher Scientific #EC6265), stained by 0.5x SYBR Gold (Thermo Fisher

Scientific #S11494), and analyzed using the Typhoon biomolecule imager (Cytiva) or

Odyssey M imager (Licor Bio). For mass spectrometry analysis, reactions were scaled

up 10-fold in ThermoPol® reaction buffer (i.e., 2 μg of RNA) at 50°C for 30 minutes.

Cleavage products were then column-purified (New England Biolabs #T2030S) and

eluted in 30 μL of nuclease-free water. Purified RNA was subsequently filtered with

Ultrafree-MC Centrifugal Filter (Millipore Sigma # UFC30GV0S) for 5 min at 10k x g

prior to LC-MS/MS analysis. For RNA fingerprinting, cleavage assays were performed

as described above, but with a limited set of guides targeting Fluc IVT RNA and pSG90

IVT RNA, generating three 72 nt or one 100 nt RNA species-specific cleavage

fragments, respectively. 3.5 pmol of Fluc IVT RNA, 3.5 pmol of pSG90 IVT RNA, or a

mix of 1.75 pmol Fluc IVT RNA and 1.75 pmol pSG90 IVT RNA were incubated with

17.5 pmol of the guide pool in the presence or absence of 17.5 pmol MpaAgo in 1x

ThermoPol® reaction buffer and the enhancer mix for 30 minutes at 50°C. Reactions

were stopped by the addition of 0.8 units proteinase K and incubation at 24°C for 10

minutes. Following a column cleanup, samples were either ran on a denaturing 6%

TBE-urea gel or analyzed by LC-MS/MS.

Capping analysis

For 5' mRNA cap analysis experiments, 50 μg (~ 83 pmol) or 25 μg (~ 44 pmol)

of Fluc or pSG90 *in vitro* transcribed mRNA, respectively, were initially denatured for 5

min at 65°C and immediately put on ice. Both mRNAs were capped using FCE (New

England Biolabs #M2081S) according to the manufacturer's manual. The capping

reactions were subsequently column purified (New England Biolabs #T2040S). 200 μl

reactions containing 3.5 pmol of capped and uncapped Fluc or pSG90 mRNA were then

incubated for 15 min at 50°C in 1x ThermoPol® reaction buffer, 30% glycerol, 160 units

of Murine RNase Inhibitor (New England Biolabs #M0314S), 67.4 pmol MpaAgo, and

66.4 pmol guide (A1: /5Phos/ATCTCCTTCTTAAAGT, A1.2: /5Phos/

TCCTTCTTAAAGTTAAAC, D891: /5Phos/ TATATCTCCTTCTTAA targeting Fluc

mRNA, or D831: /5Phos/GTCGGCTGTTTGATTC, D892: /5Phos/

TTGATTCAATAGTGTG targeting pSG90 mRNA). Reactions were incubated at 50°C.

After 15 min, the reactions were put on ice, 120 units of Exonuclease I (New England

Biolabs #M0293S) was added, and the reactions incubated at 37°C for 15 min to

remove unbound DNA guides. The reactions were stopped by the addition of EDTA to

20 mM and incubation at room temperature for 5 min. Finally, reactions were column

purified and submitted for LC-MS/MS analysis.

3' end homogenization and poly(A) tail analysis

For the analysis of the poly(A) tail composition and length and to homogenize the

3' ends of a mRNA population, MpaAgo reactions were scaled up to a 400 μl reaction

volume. 67 pmol of *in vitro* transcribed pSG95 (723 nt) or pSG120 (698 nt) mRNAs

were incubated with 162 pmol MpaAgo, 800 pmol D758

(/5Phos/TATGTTTTTTAGTCATA) or D834 (/5Phos/TTTTTTTATTAGGAGCAG) guides

in 1x ThermoPol® reaction buffer with 640 units of Murine RNase Inhibitor for 30 min at

50°C. Reactions were stopped by the addition of 1.6 units proteinase K and 10 min

incubation at room temperature. Reactions were column purified, followed by

enrichment of the poly(A) containing cleavage fragments. The volume of the purified

cleavage fragments was adjusted to 50 μl, then 1 volume (50 μl) of high salt buffer (20

mM Tris/Cl pH 7.5, 0.5 M LiCl$_2$, 10 mM EDTA) was added. 250 μl of oligo d(T)$_{25}$

magnetic beads (New England Biolabs #S1419S) were washed once with 100 μl low

salt buffer (20 mM Tris/Cl pH 7.5, 50 mM LiCl$_2$, 10 mM EDTA), and twice with high salt

buffer. The RNA was added to the beads and incubated at room temperature on a

rotator for 20 min. Bead-bound RNA was washed once with low salt buffer, twice with

high salt buffer, and one final time with low salt buffer. To elute the enriched RNA, the

beads were incubated with 30 μl of H$_2$O for 2 min at 50°C in a heat shaker (400 rpm).

The eluted RNA was recovered, incubated for 5 min at 65°C, and spun through

Ultrafree-MC Centrifugal Filter (Millipore Sigma # UFC30GV0S) for 5 min at 10k x g.

The enriched and filtered samples were then subjected to LC-MS/MS analysis.

References

1. Makarova, K.S., Wolf, Y.I., van der Oost, J. & Koonin, E.V. Prokaryotic homologs
of Argonaute proteins are predicted to function as key components of a novel
system of defense against mobile genetic elements. *Biol Direct* **4**, 29 (2009).
2. Ryazansky, S., Kulbachinskiy, A. & Aravin, A.A. The Expanded Universe of
Prokaryotic Argonaute Proteins. *mBio* **9** (2018).
3. Swarts, D.C. et al. The evolutionary journey of Argonaute proteins. *Nat Struct Mol*
Biol **21**, 743-753 (2014).
4. Nakanishi, K. Anatomy of RISC: how do small RNAs and chaperones activate
Argonaute proteins? *Wiley Interdiscip Rev RNA* **7**, 637-660 (2016).
5. Bobadilla Ugarte, P., Barendse, P. & Swarts, D.C. Argonaute proteins confer
immunity in all domains of life. *Curr Opin Microbiol* **74**, 102313 (2023).
6. Joshua-Tor, L. & Hannon, G.J. Ancestral roles of small RNAs: an Ago-centric
perspective. *Cold Spring Harb Perspect Biol* **3**, a003772 (2011).
7. Olina, A.V., Kulbachinskiy, A.V., Aravin, A.A. & Esyunina, D.M. Argonaute
Proteins and Mechanisms of RNA Interference in Eukaryotes and Prokaryotes.
Biochemistry (Mosc) **83**, 483-497 (2018).
8. Ozata, D.M., Gainetdinov, I., Zoch, A., O'Carroll, D. & Zamore, P.D. PIWI-
interacting RNAs: small RNAs with big functions. *Nat Rev Genet* **20**, 89-108
(2019).
9. Zlotorynski, E. Cell quiescence sends argonaute-2 to transposons. *Nat Rev Mol*
Cell Biol **25**, 84 (2024).
10. Underwood, C.J. & Martienssen, R.A. Argonautes team up to silence
transposable elements in Arabidopsis. *EMBO J* **34**, 579-580 (2015).
11. Baulcombe, D. RNA silencing in plants. *Nature* **431**, 356-363 (2004).
12. Lisitskaya, L., Aravin, A.A. & Kulbachinskiy, A. DNA interference and beyond:
structure and functions of prokaryotic Argonaute proteins. *Nat Commun* **9**, 5165
(2018).
13. Nakanishi, K., Weinberg, D.E., Bartel, D.P. & Patel, D.J. Structure of yeast
Argonaute with guide RNA. *Nature* **486**, 368-374 (2012).
14. Kaya, E. et al. A bacterial Argonaute with noncanonical guide RNA specificity.
Proc Natl Acad Sci U S A **113**, 4057-4062 (2016).
15. Koopal, B., Mutte, S.K. & Swarts, D.C. A long look at short prokaryotic
Argonautes. *Trends Cell Biol* **33**, 605-618 (2023).
16. Zaremba, M. et al. Short prokaryotic Argonautes provide defence against
incoming mobile genetic elements through NAD(+) depletion. *Nat Microbiol* **7**,
1857-1869 (2022).
17. Swarts, D.C. et al. DNA-guided DNA interference by a prokaryotic Argonaute.
Nature **507**, 258-261 (2014).
18. Swarts, D.C. et al. Argonaute of the archaeon Pyrococcus furiosus is a DNA-
guided nuclease that targets cognate DNA. *Nucleic Acids Res* **43**, 5120-5129
(2015).
19. Kuzmenko, A. et al. DNA targeting and interference by a bacterial Argonaute
nuclease. *Nature* **587**, 632-637 (2020).
20. Jolly, S.M. et al. Thermus thermophilus Argonaute Functions in the Completion of
DNA Replication. *Cell* **182**, 1545-1559 e1518 (2020).

21. Fu, L. et al. The prokaryotic Argonaute proteins enhance homology sequence-directed recombination in bacteria. *Nucleic Acids Res* **47**, 3568-3579 (2019).
22. Wu, J., Yang, J., Cho, W.C. & Zheng, Y. Argonaute proteins: Structural features, functions and emerging roles. *J Adv Res* **24**, 317-324 (2020).
23. Kuzmenko, A., Yudin, D., Ryazansky, S., Kulbachinskiy, A. & Aravin, A.A. Programmable DNA cleavage by Ago nucleases from mesophilic bacteria Clostridium butyricum and Limnothrix rosea. *Nucleic Acids Res* **47**, 5822-5836 (2019).
24. Olina, A. et al. Genome-wide DNA sampling by Ago nuclease from the cyanobacterium Synechococcus elongatus. *RNA Biol* **17**, 677-688 (2020).
25. Sheng, G. et al. Structure-based cleavage mechanism of Thermus thermophilus Argonaute DNA guide strand-mediated DNA target cleavage. *Proc Natl Acad Sci U S A* **111**, 652-657 (2014).
26. Zander, A. et al. Guide-independent DNA cleavage by archaeal Argonaute from Methanocaldococcus jannaschii. *Nat Microbiol* **2**, 17034 (2017).
27. Kropocheva, E., Kuzmenko, A., Aravin, A.A., Esyunina, D. & Kulbachinskiy, A. A programmable pAgo nuclease with universal guide and target specificity from the mesophilic bacterium Kurthia massiliensis. *Nucleic Acids Res* **49**, 4054-4065 (2021).
28. Liu, Y. et al. A programmable omnipotent Argonaute nuclease from mesophilic bacteria Kurthia massiliensis. *Nucleic Acids Res* **49**, 1597-1608 (2021).
29. Cao, Y. et al. Argonaute proteins from human gastrointestinal bacteria catalyze DNA-guided cleavage of single- and double-stranded DNA at 37 degrees C. *Cell Discov* **5**, 38 (2019).
30. Hegge, J.W. et al. DNA-guided DNA cleavage at moderate temperatures by Clostridium butyricum Argonaute. *Nucleic Acids Res* **47**, 5809-5821 (2019).
31. Agapov, A. et al. Prokaryotic Argonaute nuclease cooperates with co-encoded RNase to acquire guide RNAs and target invader DNA. *Nucleic Acids Res* **52**, 5895-5911 (2024).
32. Liu, Y. et al. Accommodation of Helical Imperfections in Rhodobacter sphaeroides Argonaute Ternary Complexes with Guide RNA and Target DNA. *Cell Rep* **24**, 453-462 (2018).
33. Doxzen, K.W. & Doudna, J.A. DNA recognition by an RNA-guided bacterial Argonaute. *PLoS One* **12**, e0177097 (2017).
34. Miyoshi, T., Ito, K., Murakami, R. & Uchiumi, T. Structural basis for the recognition of guide RNA and target DNA heteroduplex by Argonaute. *Nat Commun* **7**, 11846 (2016).
35. Bastiaanssen, C. et al. RNA-guided RNA silencing by an Asgard archaeal Argonaute. *Nat Commun* **15**, 5499 (2024).
36. Lapinaite, A., Doudna, J.A. & Cate, J.H.D. Programmable RNA recognition using a CRISPR-associated Argonaute. *Proc Natl Acad Sci U S A* **115**, 3368-3373 (2018).
37. Li, W. et al. A programmable pAgo nuclease with RNA target preference from the psychrotolerant bacterium Mucilaginibacter paludis. *Nucleic Acids Res* **50**, 5226-5238 (2022).

38.   Lisitskaya, L. et al. Programmable RNA targeting by bacterial Argonaute
656   nucleases with unconventional guide binding and cleavage specificity. *Nat*
657   *Commun* **13**, 4624 (2022).
658   39.   Graver, B.A., Chakravarty, N. & Solomon, K.V. Prokaryotic Argonautes for in vivo
659   biotechnology and molecular diagnostics. *Trends Biotechnol* **42**, 61-73 (2024).
660   40.   Strauss, E.G., Rice, C.M. & Strauss, J.H. Complete nucleotide sequence of the
661   genomic RNA of Sindbis virus. *Virology* **133**, 92-110 (1984).
662   41.   Vaidyanathan, S. et al. Uridine Depletion and Chemical Modification Increase
663   Cas9 mRNA Activity and Reduce Immunogenicity without HPLC Purification. *Mol*
664   *Ther Nucleic Acids* **12**, 530-542 (2018).
665   42.   AbouHaidar, M.G. & Ivanov, I.G. Non-enzymatic RNA hydrolysis promoted by the
666   combined catalytic activity of buffers and magnesium ions. *Z Naturforsch C J*
667   *Biosci* **54**, 542-548 (1999).
668   43.   Ramaswamy, S. et al. Systemic delivery of factor IX messenger RNA for protein
669   replacement therapy. *Proc Natl Acad Sci U S A* **114**, E1941-E1950 (2017).
670   44.   Pardi, N., Hogan, M.J., Porter, F.W. & Weissman, D. mRNA vaccines - a new era
671   in vaccinology. *Nat Rev Drug Discov* **17**, 261-279 (2018).
672   45.   VanBlargan, L.A. et al. An mRNA Vaccine Protects Mice against Multiple Tick-
673   Transmitted Flavivirus Infections. *Cell Rep* **25**, 3382-3392 e3383 (2018).
674   46.   Melton, D.A. et al. Efficient in vitro synthesis of biologically active RNA and RNA
675   hybridization probes from plasmids containing a bacteriophage SP6 promoter.
676   *Nucleic Acids Res* **12**, 7035-7056 (1984).
677   47.   Milligan, J.F., Groebe, D.R., Witherell, G.W. & Uhlenbeck, O.C.
678   Oligoribonucleotide synthesis using T7 RNA polymerase and synthetic DNA
679   templates. *Nucleic Acids Res* **15**, 8783-8798 (1987).
680   48.   Triana-Alonso, F.J., Dabrowski, M., Wadzack, J. & Nierhaus, K.H. Self-coded 3'-
681   extension of run-off transcripts produces aberrant products during in vitro
682   transcription with T7 RNA polymerase. *J Biol Chem* **270**, 6298-6307 (1995).
683   49.   Gholamalipour, Y., Karunanayake Mudiyanselage, A. & Martin, C.T. 3' end
684   additions by T7 RNA polymerase are RNA self-templated, distributive and diverse
685   in character-RNA-Seq analyses. *Nucleic Acids Res* **46**, 9253-9263 (2018).
686   50.   Hur, S. Double-Stranded RNA Sensors and Modulators in Innate Immunity. *Annu*
687   *Rev Immunol* **37**, 349-375 (2019).
688   51.   Vogel, A.B. et al. BNT162b vaccines protect rhesus macaques from SARS-CoV-
689   2. *Nature* **592**, 283-289 (2021).
690   52.   Brouze, A., Krawczyk, P.S., Dziembowski, A. & Mroczek, S. Measuring the tail:
691   Methods for poly(A) tail profiling. *Wiley Interdiscip Rev RNA* **14**, e1737 (2023).
692   53.   Jiang, T. et al. Oligonucleotide Sequence Mapping of Large Therapeutic mRNAs
693   via Parallel Ribonuclease Digestions and LC-MS/MS. *Anal Chem* **91**, 8500-8506
694   (2019).
695   54.   Vanhinsbergh, C.J. et al. Characterization and Sequence Mapping of Large RNA
696   and mRNA Therapeutics Using Mass Spectrometry. *Anal Chem* **94**, 7339-7349
697   (2022).
698   55.   Wolf, E.J. et al. Human RNase 4 improves mRNA sequence characterization by
699   LC-MS/MS. *Nucleic Acids Res* **50**, e106 (2022).

56. Gau, B.C. et al. Oligonucleotide mapping via mass spectrometry to enable comprehensive primary structure characterization of an mRNA vaccine against SARS-CoV-2. *Sci Rep* **13**, 9038 (2023).

57. Sato, K., Akiyama, M. & Sakakibara, Y. RNA secondary structure prediction using deep learning with thermodynamic integration. *Nat Commun* **12**, 941 (2021).

58. Lorenz, R. et al. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011).

59. Mutchek, S. et al. Strand-Cleaving Deoxyribozymes Enable the Mid-Down Sequencing of mRNA by Mass Spectrometry with No Front-End Separations. *Anal Chem* **97**, 2972-2980 (2025).

60. Bird, J.G. et al. Highly efficient 5' capping of mitochondrial RNA with NAD(+) and NADH by yeast and human mitochondrial RNA polymerase. *Elife* **7** (2018).

61. Cairns, M.J., King, A. & Sun, L.Q. Optimisation of the 10-23 DNAzyme-substrate pairing interactions enhanced RNA cleavage activity at purine-cytosine target sites. *Nucleic Acids Res* **31**, 2883-2889 (2003).

62. Vlatkovic, I. et al. Ribozyme Assays to Quantify the Capping Efficiency of In Vitro-Transcribed mRNA. *Pharmaceutics* **14** (2022).

63. Inagaki, M. et al. Cap analogs with a hydrophobic photocleavable tag enable facile purification of fully capped mRNA with various cap structures. *Nat Commun* **14**, 2657 (2023).

64. Chan, S.H. et al. RNase H-based analysis of synthetic mRNA 5' cap incorporation. *RNA* **28**, 1144-1155 (2022).

65. Wolf, E.J., Dai, N., Chan, S.H. & Correa, I.R., Jr. Selective Characterization of mRNA 5' End-Capping by DNA Probe-Directed Enrichment with Site-Specific Endoribonucleases. *ACS Pharmacol Transl Sci* **6**, 1692-1702 (2023).

66. Nwokeoji, A.O., Chou, T. & Nwokeoji, E.A. Low Resource Integrated Platform for Production and Analysis of Capped mRNA. *ACS Synth Biol* **12**, 329-339 (2023).

67. Ni, T. et al. A paired-end sequencing strategy to map the complex landscape of transcription initiation. *Nat Methods* **7**, 521-527 (2010).

68. Rothweiler, U. et al. Using nucleolytic toxins as restriction enzymes enables new RNA applications. *Nucleic Acids Res* **52**, e90 (2024).

69. Guegler, C.K. & Laub, M.T. Shutoff of host transcription triggers a toxin-antitoxin system to cleave phage RNA and abort infection. *Mol Cell* **81**, 2361-2373 e2369 (2021).

70. Di Grandi, D. et al. A single-nucleotide resolution capillary gel electrophoresis workflow for poly(A) tail characterization in the development of mRNA therapeutics and vaccines. *J Pharm Biomed Anal* **236**, 115692 (2023).

71. Beverly, M., Hagen, C. & Slack, O. Poly A tail length analysis of in vitro transcribed mRNA by LC-MS. *Anal Bioanal Chem* **410**, 1667-1677 (2018).

Acknowledgements

The authors want to thank Katherine Marks for purifying the pAgo variants tested in this study. We also want to thank Thomas C. Evans Jr. and Eric Cantor for their advice and strong support of this research.

Ethics declarations

At the time of this work, the authors were employed by New England Biolabs,

Inc., which funded this work and is the manufacturer of reagents, some of which were

used in this manuscript. Even though the authors were employed and funded by New

England Biolabs, Inc., this did not detract from the objectivity of data generation or its

interpretation.

Figure legends

Figure 1. MpaAgo efficiently and specifically cleaves mRNAs.

(A) pSG90 IVT RNA (~1.7 kb) was subject to MpaAgo cleavage with single

complementary DNA guides in ThermoPol® reaction buffer at 50°C for 15 minutes. The

substrate:Ago molar ratio was 1:5, and the Ago:guide molar ratio was 1:5. 16 nt guides

were 5' phosphorylated and were designed for MpaAgo to cleave 100 nt apart from

each other. Cleavage products were analyzed on a denaturing 6% TBE-urea gel stained

with SYBR Gold. Guide identities and cleavage efficiencies (%) are indicated above the

corresponding lanes. **(B)** Fluc IVT RNA (~1.8 kb) was subject to MpaAgo cleavage with

single complementary DNA guides. Cleavage reactions were performed under the same

conditions as in (A). 16 nt guides for Fluc RNA were also 5' phosphorylated but were

designed for MpaAgo to cleave 72 nt apart from each other. Cleavage products were

analyzed on a denaturing 6% TBE-urea gel stained with SYBR Gold. Guide identities

and cleavage efficiencies (%) were indicated above the corresponding lanes.

Figure 2. Guide multiplexing with MpaAgo produces RNA fragments suitable for

LC-MS/MS analysis.

(A) Schematic showing guide multiplexing with pAgo. Guides targeting different regions

of an RNA are pooled and used in a single reaction to cleave an RNA into smaller

fragments compatible with LC-MS/MS analysis. **(B)** pSG90 IVT RNA (~1.7 kb) was

subject to MpaAgo cleavage with single guides (lanes 2 and 3) or two 5' phosphorylated

guides in a single reaction (lane 4), at 50°C for 30 minutes. The substrate:Ago molar

ratio was 1:5, and the Ago:guide molar ratio was 1:5. Cleavage products were analyzed

on a denaturing 6% TBE-urea gel stained with SYBR Gold. The resulting short cleavage

fragments are indicated by asterisks. The target sites for the two guides were 100 nt

apart, producing a new 100 nt fragment (black asterisk) when used together. This

fragment migrated faster than the 100 nt marker likely due to different buffer

compositions or unresolved secondary structure elements. **(C)** Fluc IVT RNA (~1.8 kb)

was subjected to MpaAgo cleavage with a pool of 24 5' phosphorylated guides, with or

without 0.5% BSA or enhancer mix at 50°C for 30 minutes as indicated above the gel.

The substrate:Ago molar ratio was 1:10, and the Ago:guide molar ratio was 1:1.

Cleavage products were analyzed on a denaturing 6% TBE-urea gel stained with SYBR

Gold. The distance between target sites of adjacent guides was 72 nt, such that

complete cleavage would produce fragments of a uniform 72 nt length in addition to

unique 5' and 3' fragments of 29 nt and 81 nt, respectively. Bands ≥ 72 nt represent

incomplete cleavage fragments. **(D)** Cleavage products from (C) were analyzed by LC-

MS/MS, and the sequence coverage of Fluc RNA (%) are shown in a bar graph.

Experimental conditions: 1. no guides; 2. a pool of 24 guides, no additives; 3. 24 guides

together with 0.5% BSA and the enhancer mix; 4. 24 guides with 0.5% BSA; and 5. 24

guides with the enhancer mix.

Figure 3. 5' cap analysis with MpaAgo

(A) Schematic of MpaAgo facilitated 5' cap analysis, in which a 5' phosphorylated guide

targets MpaAgo cleavage to a site close to the 5' end of a mRNA. Upon substrate

release, the resulting fragments are analyzed via LC-MS/MS. **(B)** 5' cleavage fragments

(29 nt) of reactions containing Fluc mRNA that was either capped in vitro (+) or

uncapped (-) were resolved on a 15% TBE-urea gel. Capped fragments migrate slower

in the gel compared to uncapped fragments. The 16 nt guide DNA is indicated. **(C)** 5'

cleavage fragments (30 nt) of reactions containing pSG90 mRNA that was either

capped in vitro (+) or uncapped (-) were resolved on a 15% TBE-urea gel. **(D)** and **(E)**

Heatmap showing the relative intensity of detected 5' fragments starting with a

triphosphate (3p), diphosphate (2p), guanosine triphosphate (Gppp), cap 0

(1Me7Gppp), N⁷,2'-O-dimethylguanosine triphosphate cap (2Me7Gppp) for Fluc mRNA

(D), and pSG90 mRNA (E).

Figure 4. mRNA 3' end processing by MpaAgo

(A) Schematic representation of mRNA 3' end trimming. A non-poly(A) sequence island

in the poly(A) tail of a mRNA is targeted and cleaved by MpaAgo, resulting in a mRNA

population with homogenized 3' ends, while the 3' fragments can be further analyzed for

their composition. **(B)** A 723 nt IVT RNA (pSG95) containing a non-poly(A) island was

cleaved at four adjacent sites, using four guides (D758-D761 as indicated above the

gel, Fig. S13), and the cleavage products were resolved on a denaturing 6% TBE-urea

gel. **(C)** Schema of a reaction where MpaAgo is used to remove the complete poly(A)

tail for analysis of the full poly(A) tail by LC-MS/MS. **(D)** Sequences of the 3' region of

the pSG120 IVT RNA and two guides (D758, D834). The respective cleavage sites and

the lengths of the resulting 3' fragments are indicated. **(E)** LC-MS/MS analysis of the 3'

fragments post cleavage using the D758 (left panel) and D834 (right panel) guides. The

mass in kDa of the fragments is depicted on the x axis versus the relative intensity on

the y axis. The lengths in nt are indicated above each bar.

Figure 1

A

B

Figure 2

A

pAgo

pool of guides
targeting distinct sites

3' ———— 5'

5' ——— RNA ········ 3'

5' ········ 3'

*RNA fragments with MS compatible lengths

LC-MS/MS

B

pSG90 saRNA

cleavage
fragments

guides (16 nt)

C

no guides
24 guides
24 guides + BSA
enhancer mix
24 guides + BSA
enhancer mix

Fluc mRNA

144 nt

81 nt (3' fragment)
72 nt

29 nt (5' fragment)

guides (16 nt)

Experiment   1   2   3   4   5

D

% sequence coverage

Experiment

Figure 3

A

B Fluc mRNA

cap   +   −

29 nt +/− cap

16 nt guide

C pSG90 mRNA

cap   +   −

30 nt +/− cap

16 nt guide

D Fluc mRNA

Relative Intensity

0.00   0.25   0.50   0.75

1Me7Gppp

Gppp

2p

3p

cap   +   −

E pSG90 mRNA

Relative Intensity

0.00   0.25   0.50   0.75

2Me7Gppp

1Me7Gppp

Gppp

2p

3p

cap   +   −

Figure 4

A
pAgo cleavage in poly(A) tail sequence

mRNA ————————·····———— AAAAAAA━━━━AAxxxxxxx

polyA tail containing Ago target sequence
and unwanted extensions

mRNA ————————·····———— AAAAAAA━━

RNA population with
homogenous 3' end/length

AAxxxxxxx

- removal of cleaved 3' extensions
- analysis of poly(A) tail
composition/length

C
pAgo cleavage upstream of poly(A) tail sequence

mRNA ————————·····———— AAAAAAA━━━━AAxxxxxxx

polyA tail containing Ago target sequence
and unwanted extensions

mRNA ————————·····————

AAAAAAA━━━━AAxxxxxxx

- analysis of poly(A) tail
composition/length

B

guide, -, D758, D759, D760, D761, -

1000, 800, 600, 400, 300, 200, 100

pSG95 mRNA

cleavage products

guides

D

D834 — 91 nt →   D758 — 56 nt →

polyA tail

...GCUCCUAAUAAAAAAAAAAAAAAAAAAAAAAAAGCAUAUGACUAAAAAACAUAAAAAAAAAAAAA...

GACGAGGATTATTTTTTT-P (D834)   **ATACTGATTTTTTGTAT-P** (D758)

E

RNA fragments after pAgo/D758 cleavage

(plot: Relative Intensity vs Mass (Da), peaks labeled 55, 56, 56, 57, 57, 58, 58, 59, 59, 60)

RNA fragments after pAgo/D834 cleavage

(plot: Relative Intensity vs Mass (Da), peaks labeled 91, 92, 92, 92, 93, 93, 93, 94, 94)