Neural trajectories improve motor precision

WeiHsien Lee

Neurosurgery Department, University of Kansas Medical Center, Kansas City, Kansas, USA;

Department of Rehabilitation Medicine, University of Kansas Medical Center, Kansas City, Kansas, USA

Xavier Scherschligt

Neurosurgery Department, University of Kansas Medical Center, Kansas City, Kansas, USA;

Cell Biology and Physiology Department, University of Kansas Medical Center, Kansas City, Kansas, USA

Matthew Nishimoto

Neurosurgery Department, University of Kansas Medical Center, Kansas City, Kansas, USA;

Department of Rehabilitation Medicine, University of Kansas Medical Center, Kansas City, Kansas, USA

Adam G. Rouse†

Neurosurgery Department, University of Kansas Medical Center, Kansas City, Kansas, USA;

Cell Biology and Physiology Department, University of Kansas Medical Center, Kansas City, Kansas, USA

Bioengineering Program, University of Kansas, Lawrence, Kansas City, USA;

Electrical Engineering and Computer Science Department, University of Kansas, Lawrence, Kansas City, USA;

Abstract

Populations of neurons in motor cortex signal voluntary movement. Most classic neural encoding models and current brain-computer interface decoders assume individual neurons sum together along a neural dimension to represent movement features such as velocity or force. However, large population neural analyses continue to identify trajectories of neural activity evolving with time that traverse multiple dimensions. Explanations for these neural trajectories typically focus on how cortical circuits processes learn, organize, and implement movements. However, descriptions of how these neural trajectories might improve performance, and specifically motor precision, are lacking. In this study, we proposed and tested a computational model that highlights the role of neural trajectories, through the selective co-activation and selective timing of firing rates across the neural populations, for improving motor precision. Our model uses experimental results from a center-out reaching task as inspiration to create several physiologically realistic models for the neural encoding of movement. Using a recurrent neural network to simulate how a downstream population of neurons might receive such information, like the spinal cord and motor units, we show that movements are more accurate when neural information specific to the phase and/or amplitude of movement are incorporated across time instead of an instantaneous, velocity-only tuning model. Our finding suggests that precise motor control arises from spatiotemporal recruitment of neural populations that create distinct neural trajectories. We anticipate our results will significantly impact not only how neural encoding of movement in motor cortex is described but also future understating for how brain networks communicate information for planning and executing movements. Our model also provides potential inspiration for how to incorporate selective activation across a neural population to improve future brain-computer interfaces.

Main

Introduction

The primary motor cortex (M1) contains neurons that modulate their firing rate to help orchestrate motor output through descending recruitment of muscles. Computational models that fit M1 neural activity to various behavioral outputs such as limb kinematics, muscle force, speed, and target location show reasonable performance in a subset of movements [1,2]. Almost always, a linear summation of neural activity along certain neural dimensions is described as mapping to movement features. We believe this simplified readout assumption might be a key limitation to understanding how the brain represents movement, especially differences between large, fast movements and small, precise movements.

Point-to-point arm movements are executed with bell-shaped velocity profiles, characterized by smooth accelerations and decelerations [3–5]. In addition to representing reach direction (Georgopoulos et al, 1986), Moran and colleagues [7,8] showed that individual neurons in M1 modulate their firing rates with movement speed, producing neural activations that closely mirrored the bell-shaped profile characteristic of reaching movements. This relationship has

shaped the foundation of many neural decoding models, including those used in brain-computer interfaces (BCIs), where firing rates are typically modeled as encoding movement velocity — composed of direction and speed —as the primary variable to reconstruct motor intent [9–11]. By extension, these linear models assume all the same neurons increase their firing proportionally when greater speed is required to reach further targets.

However, these linear models typically do not generalize well to diverse sets of movements, and actual neural activity almost always has much richer spatial-temporal dynamics than could be explained by a simple readout of movement features from this neural activity [12,13]. Several explanations have previously been given for why individual M1 neurons have diverse activations over time compared to the bell-shaped kinematic features. Spatiotemporal patterns of neural firing rates have been proposed to be due to the complex muscle activations needed for movement, involving agonists, antagonists, and fixators [1,14]. Additionally, the motor cortex shows significant activity unrelated to movement, preparing for action in the 'null space' before muscle activity begins [15,16]. Coordinated cortical neural signals across many neurons is also potentially crucial for motor control, learning, and adaptation, making motor signals robust to neuron loss. Despite the growing understanding of variance in neural encoding, all these models still usually simplify downstream communication of these internal processes to a single neural dimension, using linear weighting and summation of firing rates for a degree-of-freedom of motor output [14].

The assumption that neural encoding can be reduced to a proportional output signal overlooks important aspects of neural recruitment. As the intended reach distance increases, the peak speed of movement increases [17] and greater force is required, which in turn demands the recruitment of additional muscle fibers. In the spinal cord, Henneman's size principle [18] describes the orderly and serial activation of motor neurons based on cell size rather than proportional activation of all motor neurons. Smaller neurons drive slow-twitch fibers for lower-force movements, while larger neurons only engage fast-twitch fibers for higher-force demands. Similar patterns of selective recruitment, albeit less orderly, have been observed centrally: for instance, some pyramidal tract neurons in M1 fire more robustly during slow, precise movements than during fast, ballistic ones [19]. More recently, cortical to motor unit recruitment has been shown to be functionally tuned depending on task conditions[20–22]. In addition to different recruitment patterns, motor cortex neurons also range in their timing with peaks in firing rate from 200-50 ms before peak hand velocity occurs [7,23]. These combined findings suggest that increases in force or speed may not produce a uniform and proportional scaling of activity across the same cortical population, but rather selective co-activation and/or selective timing when subpopulations are engaged.

Building on this perspective, we hypothesize that to precisely execute different amplitude movements, the motor cortex may not simply scale firing rates within a fixed neural ensemble but that selective recruitment might be important. We propose that selectively engaging subpopulations both in their co-activation and timing improves the estimation of reach distance in the presence of neural noise. To test this hypothesis, we analyzed behavioral and neural data from a non-human primate performing reaches to two different distances in the same direction.

These initial findings reveal that reach distance modulates not just the magnitude but varies the
composition and timing of the active neural ensemble. Motivated by this observation, we
developed a computational framework to investigate why the motor cortex may adopt this
strategy. We provide evidence that selective co-activation of distinct neural subpopulations with
different timings based on reach distance increases the separability between movement types in
the neural population code. We demonstrate that both *which* neuronal subpopulations are
engaged and *when* they are engaged may be critical for the most accurate motor representation.
This expanded representational space may allow brain-computer interfaces to decode movement
amplitude and movement dynamics with a more precise prediction of motor intent.

Results

Distinct neural dimensions that encode reaching behavior

We examined the neural subspaces that vary the most with given reach features of: i) velocity, ii)
condition-invariant, and iii) reach distance, on reaches to two targets. Fig. 1A illustrates the task
for the two targets analyzed, where the monkey performed a reach to different distances (near
and far) in the same direction.   Linear regression identified the two neural dimensions that best
correlated with reach velocity (A-C). Demixed Principal Component Analysis (dPCA) identified
neural dimensions with both the most variance that were condition-invariant (D-F) and that best
separated the two reach distances across time (G-I). By examining the three neural trajectories in
their separate subspaces (C, F, I), distinct neural features are observed. The velocity
representation shows neural firing rates correlated to the velocity of the reach as expected.
However, we also observe two other distinct neural features that are less directly correlated with
movement. The condition-invariant activity (F) shows that subsets of neurons are selectively
active at different times, progressing through the two different neural dimensions. This activity is
invariant between the near and far reach. The amplitude x time dimensions show neural
dimensions that separate the two reach distances across time (I). Despite these being reaches in
the same direction, this demonstrates that neurons are selectively co-activated in unique patterns
for reaches to the two different distances. Given the design of our dPC analysis, it is not
surprising that it identified two separate subspaces with neural activity that was either i)
condition-invariant or ii) distinct between the two reach distances. Notably, the amount of
variance in these neural subspaces was larger than the velocity subspace identified by regression.
Thus, these firing rate changes are not only present but larger in size to those dedicated to
representing velocity. Together, these findings suggest that reaches are encoded across multiple
neural dimensions—some dimensions representing amplitude and others capturing temporal
evolution—in addition to more classically described velocity tuned activity. These unique
dimensions differ from a classical encoding model where a neural dimension representing force
or velocity would carry direction and speed information together. Instead of most neural activity
occurring in the top row of Fig. 1, additional condition-invariant and amplitude specific neural
activity is present.

Improved Decoding Performance with Higher-Dimensional Neuron Encoding Models

Using the condition-invariant and amplitude specific neural activity as inspiration, we designed a neural simulation to test whether this type of neural activity can improve the accuracy of reaches. Synthetic firing rates were generated to encode reaches all in the same direction but for different distances. Four different neural encoding models of speed were used to generate the synthetic firing rates: (1) a proportional activation model; (2) a selective timing model with neurons activated differentially in the timing of their peak firing relative to peak movement speed; (3) a selective co-activation model created with preferentially activated neurons within distinct neural subpopulations; and (4) a combined model with both selective co-activation and timing. Fig. 2 shows the firing rates and neural trajectories under same neural subspace for the proposed neural encoding models.

Fig. 2, model 1, shows the proportional activation model where all neurons activate proportionally to the distance of movement required, whether a near target (red) or far target (blue). This velocity tuning model utilizes a single neural dimension to represent movement along a single movement degree-of-freedom as motor cortex is often classically modeled. The magnitude along the single neural dimension indicates the speed along the single degree-of-freedom. Notably, this model creates neural trajectories that lay directly on top of each other for the different amplitude movements.

For our other encoding models, we added neural dynamics that occur in multiple dimensions. Model ii utilizes selective timing of neurons (Fig. 2, model 2). The condition-invariant, cyclic neural activity we observed in Figure 1F inspired this model. By varying the lead and lag of various neurons, the current movement state, such as acceleration, peak velocity, or deceleration phases along those trajectories can be identified. While inspired by dPCA analysis of condition-invariant activity, our modeling adds this condition-invariant timing to neurons that are also varying their firing rate amplitude with reach velocity. This introduces selective timing of neural activity into neural activity that also encodes the velocity of the intended reach. Model 3 utilized a selective co-activation encoding scheme for amplitude. The separate neural dimensions for the Amplitude x Time activity identified in Fig. 1I inspired this model. In the 2D neural space, the small amplitude movement occurs in one neural dimension while the largest movement occurs in an orthogonal, second dimension. Finally, we investigated model 4 incorporating both selective co-activation and timing which we hypothesize most closely mimics the various neural features naturally observed. Here, the neural trajectories are a summation of the 2) co-activation and 3) timing models with their mixture encoding the movement information.

We trained a similarly structured MLP (Multilayer Perceptron) – LSTM (Long-short term Memory) Recurrent Neural Network (RNN) mixture model for each encoding model to provide movement decoding (Fig. 3). We used the same decoding architecture to directly compare performance across the four encoding models. The same white noise levels were added to each synthetic neural encoding models.

Comparable Performance Improvements with Selective Co-activation and Selective Timing Models

The results of our simulation, summarized in Fig. 4, show that both the selective co-activation
and selective timing neural encoding models lead to improved decoding of movement velocity
compared to the proportional model (blue line). In particular, the selective co-activation model
(orange line), where distinct subpopulations of neurons are preferentially activated with varying
peak firing rates depending on whether the target is near or far, significantly enhances decoding
performance across all tested noise levels. A similar improvement was also observed when using
a temporal progression within the neural dynamics to signal the phase of movement. Fig. 4 also
showed better performance of the selective timing model (green line) compared to the
proportional activation model (blue line). In this scenario, the decoder presumably uses the
current neural trajectory location to identify the phase of movement-acceleration or deceleration-
and better integrates the decoded velocity across time to more accurately predict movement,
highlighting how the precise timing of neural activity can lead to increased motor precision.

We next examined whether selective timing model outperforms selective co-activation model or
vice versa. Here we compared the decoding performance between selective timing (green line)
and selective co-activation model (orange line) in Fig. 4. The results revealed no significant
performance differences (Cliff's delta < 0.75) between the selective timing model and the
selective co-activation models when there was moderate noise or above ($\sigma > 0.075$), suggesting
similar performance improvements are possible with either encoding scheme.

206 **Encoding both spatial and temporal information provides the best neural representation of**
movement

Finally, we examined the encoding scheme when selective timing and co-activation is combined.
This combined model has neural trajectories with a mixture of amplitude and timing neural
features to embed movement information. Similar to the amplitude encoding scheme, the angle
of the trajectories indicates the speed of movement. Additionally, the cyclic pattern within each
trajectory allows the movement phase to be identified. Together, both selective timing and co-
activation enable a spatiotemporal encoding. As shown in Fig. 4, the selective timing + co-
activation model provides decoding performance better than the other three proposed models
(Cliffs delta > 0.75) except at very low noise levels ($\sigma < 0.075$).

217 **Similar tradeoff between selective co-activation and timing**

To better quantify the effect on encoding quality of selective timing, co-activation, or both, we
analyzed a parameter sweep across varying differences in timings and co-activations as shown in
Fig. 5. The parameter sweep was performed under a fixed noise level ($\sigma = 0.3$) within the
encoding model. The decoding performance (R^2) is presented as a heatmap across the
parameters. The darker blue indicates better decoding performance under that given parameter.
The result shows an improvement in decoding performance as both timing differences (in ms)
and the degrees between the pair of small and large neural co-activation dimensions increase.
Here we define the dimensionality by altering the angle between two representational neural
dimensions within a 2D neural manifold [24]. The dimensional degree that represents encoding of

amplitude between 0 to 90 would then indicate how much the two representational neural modes overlap with each other with a ranging dimensionality between 1D (fully overlapped and can explain neural variance in one single dimension) and 2D (orthogonal to each other, utilizing 2 neural dimension). We observe the effects of selective co-activation and timing are synergistic. In general, only a small amount of difference in co-activation or timing is needed to realize much of the performance improvement when the other feature difference is large.

Discussion

Our results underscore the critical role of selective recruitment of neurons in encoding movement information. The proposed selective co-activation and timing models demonstrate significantly improved decoding performance compared to a proportional activation model for different reach speeds. By including the selective information of neuron co-activation, the resulting neural trajectories become more distinct, allowing for better separation of movement types. This idea has some similarities to the idea of motor cortex reducing "tangling" of neural trajectories, so different movements are unambiguously represented in the dynamic neural system [25]. Here, the advantage of these separated neural features is evident where movements of varying speeds need to be differentiated. Our decoder could predict the upcoming peak speed at an early stage by analyzing the initial neural trajectory leaving the origin (idle state) before the neural activity peaks in its depth of modulation (the furthest neural state from the origin) (Fig. 2). Subsequently, the higher-dimensional models provide bigger neural differences between conditions, facilitating a more robust representation of neural activity resilient to noise.

The idea of neural trajectory encoding is similar in several ways to the idea of movement fragments presented by Hatsopoulos and colleagues (Hatsopoulos et al., 2007; Saleh et al., 2012). Like us, their work identified the importance of time-varying tuning of individual neurons to upcoming movement. It also recognized the importance of holistic submovements rather than thinking of cortex representing upcoming movement at a single instance of time. Their model, however, focused more on how a single neuron might have a preferred, often curved, movement trajectory with different directions at various future timepoints. Our work focuses on how multiple neural dimensions might better represent upcoming movements—even straight-line reaches— more accurately.

Our results highlight the utility of multiple neural dimensions for improving decoding performance. Our selective co-activation model performs comparably to the selective timing model, suggesting that either temporal variation or specific neural co-activations can enhance decoding. The combined encoding model, which combines both spatial and temporal features, emerges as the most comprehensive representation of movement. By integrating selective timing and co-activation information into higher dimensional neural spaces, this model adds a critical layer of precision, allowing for the differentiation of movement states such as acceleration, peak velocity, and deceleration. Again, this makes it easier to predict both speed and movement states even at movement initiation. The neural trajectories enable a more detailed representation of the temporal progression of movement besides the instantaneous speed. These findings align with our hypothesis that natural brain encoding leverages both spatial and temporal neural features to optimize movement representation.

Our neural trajectory results also provide a tantalizing result when considering not only motor output but interarea brain communication. Key concepts of the cerebellum are internal models and an efference copy leading to a teaching signal [28–30]. While this teaching signal is often conceived separate from a motor output signal, a predictable neural trajectory has the beneficial property that the early phase of the trajectory signals upcoming movement. This combined representation of current and future movement could provide information to cerebellum circuits about how an executed movement compares to the original motor plan. Mixed selectivity in cortical neurons may facilitate communication across brain regions by embedding contextual and motor variables into high-dimensional representations, enabling downstream structures to flexibly decode and compare motor plans [31]. Motor action selection by the basal ganglia [32,33] and other brain structures that gate motor output might also be aided by neural trajectories. The initiation of several simultaneous neural trajectories could represent possible motor actions to structures like the basal ganglia that can then be either reinforced for execution or suppressed to select a single desired motor action.

Limitations and Future Directions

While our findings provide compelling evidence for the advantages of high-dimensional encoding models, several limitations should be acknowledged. Artificial neural networks demonstrate remarkable ability to infer hidden structures in neural activity and predict movement states with high accuracy. However, they often lack in identifying biophysically realistic encoding mechanisms employed by the brain. For instance, are spatial and temporal encoding strategies interchangeable or synergistic? Our results suggest that the selective co-activation and temporal encoding schemes may be equally effective and potentially integrated within the neural system, pointing to a gap in understanding for when selectively activated neurons and temporal information might be utilized to optimize movement control. The combined encoding model suggests that these strategies are synergistic in our specific model. However, further studies and modeling are needed to understand the relative contribution of these neural features under different conditions and when one might be preferred to the other.

Finally, while the encoding models tested here were robust under controlled noise levels, their performance in more complex, naturalistic scenarios with variable noise requires further investigation. For instance, as a simplified illustrative model, our rigid neural trajectory encoding model limits online correction. The entire movement and amplitude are completely planned as the trajectories diverge at the very start of movement. While this has advantages for both movement accuracy and interarea communication, it likely limits the ability to adapt to external perturbations. More studies—both experimental and computational—of neural trajectories during perturbations are needed to understand how trajectory encoding is utilized while still providing online correction. To address this, future research should focus on refining neural network models to not only predict movement states but also developing explainable decoding architectures that can be directly compared to experimentally validated neural processes. Advancements in model interpretability and techniques for disentangling spatial and temporal contributions could provide deeper insights. Moreover, further exploration of higher-dimensional selective co-activation and timing encoding schemes could reveal whether and how the brain leverages both strategies synergistically.

Methods

Behavior task

We utilized a Center-Out Amplitude (COT Amplitude) task to characterize neural activity correlated to movement behavior with different directions and reaching distances. A male, six year old 10 kg rhesus macaque participated in this study. All animal care and research procedures adhered to the Guide for the Care and Use of Laboratory Animals. Approval was granted by the Institutional Animal Care and Use Committee at the University of Kansas Medical Center.

The monkey was trained to perform the COT Amplitude task using an exoskeleton robotic arm to control a cursor on a 47'' LED display provided by NHP KinArm (BKIN Technologies, Kingston, ON). Research subjects were body-restrained in the KinArm chair, as well as forearm and elbow restrained on the exoskeleton arm. The exoskeleton arm could move freely at the elbow and shoulder on a 2-Dimensional plane parallel to the ground in this task. The cursor on the screen represented the center of hand position in the exoskeleton arm which can freely move within a 1000-by-1000 pixel workspace. A circular cursor indicated a single point in the workspace. Dexterit-E data acquisition and experimental control was customized within a Simulink software environment to record the cursor position and trial event information and was down sampled to 100 Hz for analysis with the neural data.

We recorded 13 sessions of neural recordings in a time span of 1 month. The monkey successfully completed a total of 8867 trials across the recording sessions. During the COT Amplitude task, the monkey was instructed to perform a center-out reach to 1 of the 8 peripheral targets in four reaching directions equally spaces by 90° around the origin after obtaining and holding the cursor at the center target for 300-500 ms with a radius of 1.25 cm. Each of the reach directions includes two targets with different reach distances, a closer target and a further target. In this study, we are only using neural and behavioral data from one of the four directions to show the different encoding schemes between the two reaching distances to simplify and align with the proposed hypothesis and simulation models.

Linear regression

We implemented linear regression to examine how neural firing rates evolved in relation to cursor movement X and Y velocity during one direction of reaching in COT Amplitude task. To begin, we extracted trial-averaged firing rates from 1,228 neurons during near and far reaches and aligned them with the corresponding conditions. We then performed principal component analysis (PCA) on this matrix to reduce dimensionality, capturing dominant features of population activity across time and movement contexts.

To link neural dynamics with behavior, we next fit linear regression models that predicted each kinematic feature—X and Y velocity—based on the top 15 neural principal components. This allowed us to assess how well low-dimensional neural trajectories captured behavioral variation. By projecting each behavioral variable back onto the neural state space, we visualized how

351 motor parameters were encoded over time within the population activity. This approach revealed
352 behaviorally tuned dimensions in the neural state space that reflected continuous encoding of
353 movement features.

dPCA analysis

We applied demixed principal component analysis (dPCA) to identify and emphasize the key
features of complex population activity. This method decomposes the population activity into a
limited number of demixed components that capture the most variance in the data related to the
specified task features such as time and reach distance.

We first applied principal component analysis (PCA) to reduce the dimensionality of the firing
rate data to 15 dimensions. PCA was performed on the trial-averaged firing rates within a time
window spanning from 300 ms before to 300 ms after movement onset. The neural activity
matrix was reshaped and projected onto principal components (PCs) using the computed
eigenvectors, resulting in a low-dimensional representation of neural dynamics. The PCA scores
were then reorganized into a structured format suitable for dPCA, preserving the original data
dimensions while enabling task-specific decomposition. The neural data were arranged into a
tensor of size $N \times T \times A$, where N represents the number of neural PCs, T time points and A
amplitudes. This transformation allowed dPCA to systematically separate neural dimensions that
encode task parameters, distinguishing between condition-dependent neural dynamics that vary
with time.

Once dPCA was applied, the projected neural trajectories were then computed by multiplying the
PCA-transformed neural activity with the dPCA demixing weights. We analyzed neural
population activity in these distinct component spaces derived from dPCA, focusing on three key
representations: (1) condition-invariant time-varying neural dynamics, reflecting the shared
temporal structure of neural activity across conditions; (2) condition-dependent representations,
where neural activity was modulated by reach amplitude, highlighting task-specific encoding;
and (3) time × amplitude interactions, which revealed how movement speed influenced neural
dynamics over time.

Neural simulation

In the study, we simulated motor cortical activity to generate movements. The simulated task
mimicked the kinematics of a single direction reaching task from movement onset to target
acquisition. A bell-shaped speed profile like those observed in natural point-to-point reaches was
used [34] without considering other factors such as position or acceleration.

We simulated 1000 reaching trials, each with a randomly assigned velocity peak within a range
of 0.25 to 1 arbitrary unit. Each trial had a duration of 200 ms with sampling of 5 ms. Despite
various magnitudes of the movements, the speed profiles across all trials shared the same
symmetric gaussian shape with a standard deviation of 33.3 ms. Example movement speed
profiles are shown in Figure 1 with the i) red curve representing a small magnitude reaching

movement with lower speed and ii) blue curve for a large magnitude reaching movement with
high speed. In our simplified task, all movements were the same duration with the speed
amplitude directly related to the distance reached. These same 1000 simulated reaching trials
were used for all the various simulated neural models.

Speed encoding models

We designed four neural encoding models of speed to describe the neural activity based on
selective activation of neurons: (1) a **proportional activation** model, in which all neurons scale
their firing rates linearly with movement velocity; (2) a **selective timing** model, in which all
neurons encode speed proportionally, but differ in the timing of their peak firing relative to peak
movement speed; (3) a **selective co-activation** model, where distinct neural subpopulations are
preferentially activated depending on movement amplitude (near vs. far), with all neurons
peaking at the same time relative to movement; and (4) a selective timing + co-activation model
that includes both selective co-activation and timing for each neuron across the subpopulation.
The toy neuron models were tuned solely to velocity for simplicity. The same Gaussian white
noise was added to the firing rates of all neural models. In the proportional activation model, the
firing rate was generated by multiplying the actual speed vector and the speed tuning weights of
each neuron with following equation:

$$f(t - \tau) = b_0 + b_v vel(t) + \sigma$$

Where f is the neural firing rate; b_0 is the baseline firing rate; b_v is the speed encoding weight
vector across the neurons; vel is the actual movement speed across each trial.

In the selective co-activation model, we used two orthogonal neural dimensions of tuning
weights to encode speed. These two vectors combine together to generate the neural activity as
follows:

$$f = b_0 + b_{v,1} * \sqrt{1 - \alpha} * scaled_{factor}\, vel(t) + b_{v,2} * \sqrt{\alpha} * scaled_{factor}\, vel(t)$$

Now, $b_{v,1}$ and $b_{v,2}$ define the two weight vectors across the neurons creating two neural
dimensions of encoding. The α value is a scaling term that is 0 for the smallest speed movement
possible and 1 for the largest speed movements.

The selective timing model introduces a change in time lags for the different neurons tuning to
speed. We manipulated the shift of the time across the neurons to create a population code with
the same velocity shape as other models to represent the same velocity profile. The neural
activity would then be described as follow

$$vel_{drift} = normpdf(\mu_t, \sigma_t)$$

$$f = peakvel \cdot (vel_{drift} \cdot f_{shape})$$

Where $velshape_{drift}$ define the time lag of different neurons, f_{shape} applies the bell-shape
speed profile encoded in the neural activities.

Finally, we tested a combination model with a linear sum of both selective co-activation
and timing model with various weights defined for both the amplitude and phase encoding as
described above.

Neural Trajectory

Principal component analysis (PCA, pca function in MATLAB; MathWorks, 2021) was used to
visualize the neural dynamics and identify the dominant patterns within the lower dimensional
latent space. We projected the spike firing rates into a single PCA subspace across time to get
neural trajectories and imply neural dynamics for given movements within each encoding model.

The neural trajectory demonstration of two movement types (small and large) are color-coded the
same as above in speed profiles (Fig. 1). These trajectories allow us to visualize the distinct
neural patterns between various movements. The trajectories indicate the transition of neural
states in the first two principal component state space over the trial time (200ms).

Our hypothesis revealed distinct neural encoding models for movement. A proportional
activation model showed neural activity linearly related to speed along a single dimension (Fig.
2, model 1). In the selective co-activation model, neural trajectories occupied different regions of
neural space for varying movement types, indicating higher dimensional encoding (Fig. 2, model
2). A selective timing model introduced cyclic patterns in neural space, distinguishing movement
phases, such as acceleration and deceleration (Fig. 2, model 3). Combining selective co-
activation and timing models enhanced movement information, allowing inference of movement
type and phase within the neural trajectory (Fig. 2, model 4).

RNN decoder design

We employed an RNN mixture model which integrates a MLP and a LSTM network to train on
the hypothesized four encoding models of neural activities.

The MLP serves as an initial processing stage, reorganizing the neural data and providing a more
structured interpretation for the subsequent LSTM network. It consists of two fully connected
layers with ReLU activation functions, designed to capture the non-linear relationship in the raw
neural data. The last layer readout in MLP, preserving the dimensions of neural data, is then
passed down to the subsequent LSTM network as input. The non-linear regression process
within the hidden layers of MLP can be described as follow.

Let $x \in \mathbb{R}^n$ represent the input neural data. The MLP consists of L layers, each defined by a set
of weights $W^{(l)}$ and biases $b^{(l)}$ for layer l , where $l \in \{1,2, \dots, L\}$.

The non-linear transformation at each layer can be described as:

$$h^{(l)} = \phi\big(W^{(l)}h^{(l-1)} + b^{(l)}\big)$$

where:

• $h^{(0)} = x,$
• $h^{(l)}$ is the output of the l -th layer,

• $\phi(\cdot)$ is non-linear activation function ReLU (Rectified Linear Unit), defined as $\phi(z) =$
$\max(0, z)$.

The final output of the MLP, $h^{(L)}$, provides the encoded representation of the neural data, which
is then passed to the LSTM network.

The LSTM network comprises two hidden layers with tanh activation functions and sigmoid
gates to manage the flow of information across time steps and finally infer the speed over time
steps at the output.

We optimized the decoding model's hyperparameters by conducting a comprehensive sweep over
weights and biases. This process involved systematically varying parameters including learning
rate, number of units per layer, regularization coefficients, and initialization methods.
Performance metrics, including accuracy and loss, were used to evaluate each configuration. The
optimal hyperparameter set was selected based on its ability to enhance the capacity to capture
temporal dynamics in the neural data, resulting in improved predictive performance.

Quantification of Decoding performance

We used the correlation of determination (R^2) to evaluate how well velocity can be predicted
using the same deep learning architecture across different encoding models. A single-trial R^2 is
calculated to assess the performance of the MLP-LSTM model on individual trials. This metric
provides a detailed measure of the model's predictive accuracy, showing how effectively it
captures the relationship between neural activity and movement speed within each trial. By
examining single-trial R^2, we gain insight into the model's ability to decode speed from neural
activity and account for variability observed in single-trial movement patterns.

To evaluate overall performance, a trial-averaged R^2 is calculated by averaging the R^2 values
across all trials for each prediction made by the MLP-LSTM model. This trial-averaged R^2
provides a comprehensive measure of the model's generalization capability, reducing the
influence of noise and trial-to-trial variability. By aggregating the R^2 values, we emphasize the
predictive power of each encoding model under varying conditions, highlighting their ability to
capture the relationship between neural activity and movement dynamics effectively.

Statistics

We performed multiple iterations using random neural tunings in the encoding model and
randomized initial conditions in the similarly structured MLP-LSTM decoding architecture to
test for repeatability and robustness of the observed model differences. The white neural noise in
the encoding model was also randomly generated but the same noise instance was used across
the four encoding neural models to minimize the effect of randomized noise on model
differences. Models were trained through the similarly structured MLP-LSTM decoding
architecture to get a confidential interval of the performance difference between those encoding
models and identify outliers during training.

We used Wilcoxon signed-rank test as a non-parametric statistical tool to conduct pairwise
comparison between decoding performance of the four encoding models. As is common with

computational models with many iterations, the Wilcoxon test showed statistically significant
mean differences in decoding performance ($p \ll 0.001$) between all pairs of models at each noise
level. We then chose to calculate the Cliff's delta values for each comparison groups to test for
meaningful effect size differences in decoding performance. We chose a Cliff's delta greater than
0.75 as a meaningful difference in effect size. This corresponds to one model outperforming the
other model 75% of the time. We also present the 25 to 75 percentiles for R^2 values across trials,
allowing us to quantify the variability and reliability of the model predictions.

Figures

Figure 1. Neural subspace for reaches to two target distances.

We applied a linear regression model for velocity and demixed Principal Component Analysis
(dPCA) to neural population activity recorded during reaches to targets of two different distances
to isolate and visualize behaviorally relative neural dynamics in three subspaces. The X/Y
velocity regression model identifies the neural spaces where velocity information is best
predicted. A) Neural dimension showing trial-averaged firing rates over time best predicting X-
velocity. Red lines show the trial average movement trajectories towards the near targets. Blue
lines show the trial average movement trajectories towards the far targets. B) Neural dimension
showing trial-averaged firing rates over time best predicting Y-velocity. C) Neural trajectories
best predicting velocity in the 2D neural space, first and second dimension indicate velocity
prediction in X, Y reach dimensions respectively. The Condition-Invariant activity identified
with dPCA. D) First neural dimension in the dPCA CI space, showing condition invariant
average firing rate over time. E) Second neural dimension in the dPCA CI space, showing
condition invariant average firing rate over time. F) Neural trajectories in the condition-invariant
subspace combining D and E. The Amplitude x Time dimensions identified with dPCA. G) First
neural dimension in the dPCA amplitude component space, showing amplitude dependent
average firing rate over time for near and far reaches. H) First neural dimension in the dPCA
amplitude × time interaction space, showing average firing rate over time for near and far
reaches. I. Neural trajectories in the 2D dPCA space composed of G and H. Each subplots shows

neural states from 400 ms before movement onset (box) to 250 ms after movement onset
(triangle). All color-coded by targets similarly shown in fig 1.A.

540
Figure 2. Firing rates and neural subspace for simulated encoding models.

Firing rates of simulated neurons for 4 encoding models. The firing rates are shown for near
(left) and far (right) distance reaches with velocity scaled to maintain the same reach duration.
First, in proportional activation, all neurons scale their firing rate in a linear relationship with
velocity as assumed by a linear neural encoding model. Next, subpopulations of neurons are
selectively co-activated with different peak firing rates depending on whether the target is near or
far. All neurons have the same timing relative to movement. Third, the selective timing of peak
firing rates relative to peak speed is varied. The peak firing rates are proportional to speed for all
neurons in this selective timing model. Finally, a combined model includes both selective co-
activation and timing for each neuron across the population. Neurons in each row are identified
by the same color for near and far reaches.

Figure 3. RNN mixture model architecture diagram.

This diagram shows the architecture of the Multi-Layer Perceptron (MLP) and a Long short-term
memory (LSTM) mixture model we used as the speed information decoder. The MLP consists of
one fully connected hidden layer to capture non-linear relationships with hidden cells (green
circles) and pass the output as the input (yellow circles) of LSTM model. Where LSTM consists of
2 hidden recurrent layers (recurrent cells color-coded in blue) and one output cell (orange
circle).

Decoding performances over noise level

Figure 4. R2 performance plot

The correlation coefficient (R^2) indicates the decoding performance of the propose encoding models, proportional model (red), selective co-activation model (orange), selective timing model (green), selective timing + co-activation model (red), over different noise levels. The average R^2 of the model's performance is color-coded in bold color lines and the 25-75 percentile range of performances over bootstrapping is in shaded area with corresponding color keys. The proportional model was significantly worse at all noise levels (Cliffs' delta > 0.75). Selective timing + co-activation model was better than all other models (Cliffs' delta > 0.75) except for being equivalent to selective co-activation at low noise levels (σ= 0-0.075). selective timing and selective co-activation model were also equivalent (Cliffs' delta < 0.75) at low noise levels (σ= 0-0.05).

Figure 5. Selective timing/ co-activation model parameter sweep analysis on a fixed noise level.

A heatmap of the parameter sweep performed across timing and co-activation levels,
quantifying the improvement in encoding both dimensional and phase information at various
level. The proportion of neuron co-activation that's encoded in the sweep is determined by
degrees. 0 degree indicates no encoding of movement amplitude, and 90 degree indicates two
orthogonal dimensions of the neural space to encode movement amplitude. Different times lags
indicate the variance in time delays across neurons relative to movement during encoding.
Darker blue color indicates a better performance in the model. The parameter sweep is
performed on encoding models at a single noise level ($\sigma = 0.3$).

Data and code availability.

The data that support the findings of this study are available from the corresponding authors upon request. The code used for the analyses will be publicly available through *github*.

Funding:

HHS | NIH |NINDS R00NS101127 (AGR)

Frank & Evangeline Thompson Opportunity Fund (AGR)

Steve Palermo Foundation (AGR)

Acknowledgements:

Author contributions
Contributions:

	WL	XS	MN	AR
Designed research	v	v	v	v
Performed research	v			
Contributed unpublished reagents/analytic tools				
Analyzed data	v	v	v	v
Wrote the first draft of the paper	v			v
Edited the paper	v	v	v	v
Wrote the paper	v	v	v	v

Reference

1.   Todorov, E. Direct cortical control of muscle activation in voluntary arm movements: a model. *Nature* **3**, 391–398 (2000).

2.   Tanaka, H. & Sejnowski, T. J. Computing reaching dynamics in motor cortex with Cartesian spatial coordinates. *J Neurophysiol* **109**, 1182–1201 (2013).

3.   Georgopoulos, A. P., Kalaska, J. F. & Massey, J. T. *Spatial Trajectories and Reaction Times of Aimed Movements: Effects of Practice, Uncertainty, and Change in Target Location*. *JOURNALOF NEUROPHYSIOL~GY* vol. 46 (1981).

4.   Morasso, P. *Ex Mental Bran Research Spatial Control of Arm Movements**. *Exp Brain Res* vol. 42 (1981).

5.   Soechting, J. F. *Effect of Target Size on Spatial and Temporal Characteristics of a Pointing Movement in Man*. *Exp Brain Res* vol. 54 (1984).

6.   A P Georgopoulos, A. B. S. R. E. K. Neuronal population coding of movement direction. **1986**, (1986).

7.   Moran, D. W. & Schwartz, A. B. *Motor Cortical Representation of Speed and Direction During Reaching*. (1999).

8.   Wang, W., Chan, S. S., Heldman, D. A. & Moran, D. W. Motor cortical representation of position and velocity during reaching. *J Neurophysiol* **97**, 4258–4270 (2007).

9.   Carmena, J. M. *et al.* Learning to control a brain-machine interface for reaching and grasping by primates. *PLoS Biol* **1**, 193–208 (2003).

10.   Simeral, J. D., Kim, S. P., Black, M. J., Donoghue, J. P. & Hochberg, L. R. Neural control of cursor trajectory and click by a human with tetraplegia 1000 days after implant of an intracortical microelectrode array. in *Journal of Neural Engineering* vol. 8 (2011).

11.   Willsey, M. S. *et al.* Real-time brain-machine interface in non-human primates achieves high-velocity prosthetic finger movements using a shallow feedforward neural network decoder. *Nat Commun* **13**, (2022).

12.   Churchland, M. M. *et al.* Neural population dynamics during reaching. *Nature* **487**, 51–56 (2012).

13.   Michaels, J. A., Dann, B. & Scherberger, H. Neural Population Dynamics during Reaching Are Better Explained by a Dynamical System than Representational Tuning. *PLoS Comput Biol* **12**, (2016).

14.   Sussillo, D., Churchland, M. M., Kaufman, M. T. & Shenoy, K. V. A neural network that finds a naturalistic solution for the production of muscle activity. *Nat Neurosci* **18**, 1025–1033 (2015).

15.   Kaufman, M. T., Churchland, M. M., Ryu, S. I. & Shenoy, K. V. Cortical activity in the null space: Permitting preparation without movement. *Nat Neurosci* **17**, 440–448 (2014).

16.　Elsayed, G. F., Lara, A. H., Kaufman, M. T., Churchland, M. M. & Cunningham, J. P. Reorganization
between preparatory and movement population responses in motor cortex. *Nat Commun* **7**, 1–15
(2016).

17.　Fitts, P. M. *THE INFORMATION CAPACITY OF THE HUMAN MOTOR SYSTEM IN CONTROLLING THE*
AMPLITUDE OF MOVEMENT 1. *Journal of Experimental Psychology* vol. 47 (1954).

18.　Henneman, E. *Relation between Size of Neurons and Their Susceptibility to Discharge*. *New Series*
vol. 27 (1957).

19.　Fromm' And, C. & Evarts, E. V. *RELATION OF SIZE AND ACTIVITY OF MOTOR CORTEX PYRAMIDAL*
TRACT NEURONS DURING SKILLED MOVEMENTS IN THE MONKEY. *The Journal of Neurosci ence*
vol. 1 (1981).

20.　Davidson, A. G., Chan, V., O'Dell, R. & Schieber, M. H. Rapid changes in throughput from single
motor cortex neurons to muscle activity. *Science* **318**, 1934–7 (2007).

21.　Griffin, D. M., Hoffman, D. S. & Strick, P. L. Corticomotoneuronal cells are 'functionally tuned'.
Science (1979) **350**, 667–670 (2015).

22.　Marshall, N. J. *et al.* Flexible neural control of motor units. *Nat Neurosci* **25**, 1492–1504 (2022).

23.　Cheney, P. D., Fetz, E. E., Cheney, P. D. & Fetz, E. E. *Functional Classes of Primate*
Corticomotoneuronal Cells and Their Relation to Active Force. *JOURNALOFNEUROPHYSIOLOGY*
vol. 44 (1980).

24.　Gallego, J. A., Perich, M. G., Miller, L. E. & Solla, S. A. Neural Manifolds for the Control of
Movement. *Neuron* vol. 94 978–984 Preprint at https://doi.org/10.1016/j.neuron.2017.05.025
(2017).

25.　Russo, A. A. *et al.* Motor Cortex Embeds Muscle-like Commands in an Untangled Population
Response. *Neuron* **97**, 953-966.e8 (2018).

26.　Hatsopoulos, N. G., Xu, Q. & Amit, Y. Encoding of movement fragments in the motor cortex.
Journal of Neuroscience **27**, 5105–5114 (2007).

27.　Saleh, M., Takahashi, K. & Hatsopoulos, N. G. Encoding of coordinated reach and grasp
trajectories in primary motor cortex. *The Journal of neuroscience* **32**, 1220–32 (2012).

28.　Marr, D. A theory of cerebellar cortex. *J Physiol* **202**, 437–470 (1969).

29.　Sanger, T. D., Yamashita, O. & Kawato, M. Expansion coding and computation in the cerebellum:
50 years after the Marr–Albus codon theory. *Journal of Physiology* **598**, 913–928 (2020).

30.　Albus, J. S. A theory of cerebellar function. *Math Biosci* **10**, 25–61 (1971).

31.　Rigotti, M. *et al.* The importance of mixed selectivity in complex cognitive tasks. *Nature* **497**, 585–
590 (2013).

32.   Stephenson-Jones, M., Samuelsson, E., Ericsson, J., Robertson, B. & Grillner, S. Evolutionary
conservation of the basal ganglia as a common vertebrate mechanism for action selection.
Current Biology **21**, 1081–1091 (2011).

33.   Mink, J. W. The Basal Ganglia and Involuntary Movements. *Arch Neurol* **60**, 1365 (2003).

34.   Kang, T., He, J. & Tillery, S. I. H. Determining natural arm configuration along a reaching trajectory.
Experimental brain research. Experimentelle Hirnforschung. Expérimentation cérébrale **167**, 352–
361 (2005).

683