1

Identifying and prioritising conflicts between human and wildlife interests in Great Britain

8   Kate L. Palphramand[1]*, Dave Cowan[1], Daniel A. Warren[1] & Graham C. Smith[1]

12   [1] Animal and Plant Health Agency, York, YO41 1LZ, UK

14   *Corresponding author

15   E-mail: kate.palphramand@apha.gov.uk (KP)

Abstract

We reviewed the scientific literature to identify human-wildlife conflicts associated with British terrestrial mammals to summarise their magnitude and, where possible, costs. Conflicts were classified as either economic, health, environmental or social, further divided into 32 subcategories. We applied a Generic Impact Scoring System (GISS), whereby individual conflicts were scored on a scale of 1-5 (minimal to highest known impact); these were used to determine the potential impact (sum of all conflicts) and actual impact (sum of all conflicts corrected for geographical distribution) of individual mammals so they could be ranked on their level of impact. Forty-eight mammals were involved in a total of 200 separate conflicts. The highest ranked conflicts (scoring 5) were all measurable on an economic scale (involving rabbit, badger, brown rat and grey squirrel). We estimated a conservative annual economic cost of human-wildlife conflicts in excess of £0.5 billion, which could exceed £1 billion if health and environmental conflicts could be monetarised. The most common conflicts were health related (reservoir of disease and zoonotic disease), although generally, health conflicts were scored low (mostly 1 or 2). Overall, there was no statistical difference in impact score depending on status (native or non-native); however, when considered by subcategory, non-native species scored higher in relation to damage to woodland, agricultural interests and property, and reservoir of zoonotic disease, whilst native species caused more conflict with construction and development, and predation of livestock. Non-native species had a higher potential impact score compared to native species, but not actual impact. However, actual impact was predicted to increase, relative to population size, at a higher rate for non-native species compared to native, highlighting the need for surveillance and monitoring. Where mammals represent a cost to some, they may provide a benefit to others, e.g. fox predation of rabbits, and

40   as such may require a holistic approach to management. We concluded that a GISS approach

41   is a useful tool for identifying conflict species and could form part of a package of measures to

42   ensure a balance between the need to conserve and enhance biodiversity, whilst also

43   considering the interests of those directly affected by human-wildlife conflict, is achieved.

Introduction

Co-habitation between humans and wildlife occurs on more than half of the earth's surface [1]

and consequently human-wildlife interactions form an integral part of many ecosystems

worldwide. Interactions between people and wildlife are often varied and complex and whilst

the presence of wildlife may have beneficial impacts on human interests, there is a continued

focus on conflict narratives that dominates how human-wildlife interactions are viewed [2].

Human-wildlife conflict is commonly described as conflict occurring between people and

wildlife [3], presence and/or actions of humans or wildlife that have an adverse effect on the

other [4] or threats posed by wildlife to economic activities, human life, recreation [5] or

ecosystem services. Factors such as the intensification of agriculture and increasing human

population have had major impacts on ecosystems worldwide [6]. Agricultural growth is

expected to accelerate, resulting in an increase in farmland and subsequent decrease in the

available habitat for wildlife [7]. Wildlife damage to crops is often cited as a major global conflict

[3, 8, 9], which will continue to worsen as habitat loss precipitates an increase in human-

wildlife encounters. Population growth and habitat encroachment may also increase the risk of

disease transmission, which is of particular concern where wildlife pose zoonotic risks. For

example, the first human cases of severe acute respiratory syndrome coronavirus (SARS-CoV)

in Guangdong province, China, in 2002, were linked to wildlife found at a live animal market [10,

11]. Subsequent serological surveys from residents living close to bat caves in Yunnan, China,

which is geographically disparate to Guangdong, were found to be positive for SARS-related

coronavirus (SARSr-CoV), indicating that spillover events may have occurred in zoonotic CoVs

[12].

Conflict issues can arise with several types of problematic species, including native pests,

invasive native species and invasive non-native species [13]. These often induce debates on

how or whether humans and wildlife can co-exist [7], and may involve complex decisions

where the conservation status of a species conflicts with its impacts on humans [3]. Addressing

and categorising the adverse effects of human-wildlife conflicts is recognised as a challenging

issue for conservation and development efforts [14]. There is a general consensus that the

negative effects created by some wildlife species, particularly invasive non-native species,

have to be reduced [15]. Where conflicts arise between wildlife and human interests, there is

often the need to intervene to reduce or mitigate the impacts and therefore identifying what

conflicts are occurring and what their impacts are is essential for informing decision-making

and resource allocation [16]. In 2022, the UK agreed to a global policy framework at the

Conference of Parties (COP15) to the Convention on Biological Diversity that created a package

of international measures to address the loss of global biodiversity and to restore natural

ecosystems. One of the targets identified for reducing threats to biodiversity was managing

human-wildlife conflicts, which in turn would feed into other elements of the framework, such

as enhancing sustainability in agriculture, fisheries and forestry [17]. However, identifying which

species are causing conflict and subsequently which should be targeted first can be challenging

and controversial [15].

Invasive non-native species are a major threat to biodiversity worldwide, severely

impacting areas of high endemism [18], whilst also posing a serious threat to public health and

economic activities. As of 2019, 3,224 plants and animals had been introduced to Great Britain,

costing the UK economy almost £4 billion annually [16]; 10-15% of these are considered

invasive with an estimated 188 invasive non-native animals recorded in 2020 [19]. Tackling

invasive non-native species is a key part of achieving the UK government's aims under its 25-

year Environment plan [20] as well as the UK's obligation as a signatory of the Convention on

Biological Diversity (https://www.cbd.int/convention/text/). However, even managing conflict

caused by invasive non-native species can be controversial, resulting in debates on

achievability, efficiency, social fairness and ethical implications [21]. For example, the removal

of hedgehogs from South Uist, where they were introduced to control garden pests but

consequently contributed significantly to the decline of wading shorebirds [22] resulted in

widespread criticism and opposition, given that hedgehogs are a popular icon of British wildlife

and are thought to be declining nationally [21].

Much of the available literature on wildlife conflicts in the UK, both in terms of economic

and biodiversity impacts, focuses on invasive non-native species [e.g. 16, 23]. However, there

is evidence to suggest that the most invasive non-native species are not necessarily the most

damaging [24, 25], therefore there is the need to consider also the impacts of native species,

some of which can cause considerable human-wildlife conflict. For example, badgers are a

known reservoir host of *Mycobacterium bovis*, the causative agent of bovine tuberculosis,

which poses a risk to livestock health and is a serious economic problem for the UK cattle

industry [26]. Issues such as this highlight the challenge of conflict resolution for iconic British

species, which can be both controversial and associated with complex legislative

considerations. It is necessary, however, to consider all problematic species to address the

UK's commitments to improving biodiversity as part of government plans or global policy

frameworks.

Whilst policy frameworks are essential for providing goals and targets to reverse

biodiversity loss, they do not necessarily incorporate a process by which these targets, such as

managing human-wildlife conflict, can be identified in the first place, nor which should be

prioritised. Assessing conflicts between species is further complicated by the fact that impacts

are context dependent [27], which requires that the measure of impact is standardised to

quantify and compare differences between species, taxonomic groups and localities [28].

When addressing the biodiversity impacts of invasive non-native species, Nentwig et al. [15]

suggested prioritising actions against the alien species with a high impact potential of becoming

established or spreading, using a method to quantify impacts in a way that could be compared

among alien species within a taxonomic group. They devised a generic impact scoring system

(GISS) to compare relevant impact categories that had been reported in the scientific literature,

applying a scoring system to quantify impacts within the categories and rank species according

to their impact scores. The GISS has since been modified and applied to a variety of

classifications including alien fish [29], terrestrial arthropods [30], aquatic invertebrates [31],

plants [32, 33] and alien birds [34].

The main aim of this paper was to identify and categorise human-wildlife conflicts

associated with British mammals in order to summarise their magnitude and, where possible,

their costs. The research augmented a review of human-wildlife conflicts produced over a

decade ago (APHA, unpublished data) in response to a now withdrawn Defra policy framework

on Wildlife Management in England [35]. Only terrestrial mammals (including bats) were

assessed as they involve some of the most controversial and sensitive issues as well as

complex legislative considerations. To determine whether the available evidence reflected the

level of risk to human interests, or the welfare or conservation status of an affected species, a

modified GISS was applied to all conflicts so that mammals could be ranked according to their

level of impact. Identifying species with the highest impact scores has the potential to assist

policy development by enabling prioritisation of the most damaging species. In addition, the

compilation of a comprehensive audit of British wildlife conflicts highlights where further areas

of research might be required and identifies the early emergence of potential issues.

Materials and methods

Literature searches and screening

Searches were undertaken using Scopus (an abstract and citation database of peer-reviewed

scientific literature) for literature published up until and including 2024. Additional searches

were conducted using Google Scholar limited to the first 200 returns [36] to provide additional

evidence where required. Separate searches were carried out for each of the 66 terrestrial

mammals incorporating a combination of unique species terms and common keywords within

the title, abstract and keywords of the scientific literature. The following search terms were

applied to each species: [{common name} AND {Latin name}] AND [{England} OR {Scotland}

OR {Wales} OR {Great Britain} OR] AND [{conflict} OR {pest} OR {damage} OR {impact} OR

{management} OR {disease} OR {zoono*} OR {animal health} OR {health} OR {conservation} OR

{predat*}]. The minimum requirements for a literature resource to be included in the analysis

were the presence of an accessible abstract and that the conflict occurred in Great Britain. Only

articles written in English were included.

All articles returned during an individual search were subjected to a number of screening

processes. A rapid first screening of the title assessed documents for any likely relevance to the

species concerned and its potential conflicts. All potentially relevant articles or ambiguous

titles were screened further by examining the abstracts in more detail or the full article, if

necessary, resulting in a definitive list of conflicts for each species. In many cases, single

articles were applicable to multiple conflicts and/or more than one species and therefore may

have been added retrospectively to a species reference catalogue.

Categorising and scoring conflicts

Conflicts were categorised depending on whether they were an economic, health,

environmental or social impact. Within these four main categories, conflicts were further

assigned to 32 subcategories (economic=15, health=6, environmental=7, social=4).

Subcategories were not weighted, therefore all subcategories were considered as of equal

importance. Some species were involved in multiple conflicts within multiple categories, which

were all captured as distinct entities. A comprehensive report was compiled for all species

identified as being involved in human-wildlife conflict. An impact scale scoring system modified

from Nentwig et al. [15] was applied to estimate the magnitude of each conflict at the national

level for Great Britain, rated independently by three reviewers (KP, DC, GS). The level of impact

was quantified on a scale of 1 (minimal impact known) to 5 (the highest possible impact: Table

1), with a value of confidence (low=1, medium=2 or high=3: Table 2) also assigned to each

conflict (taken from Hawkins et al. [37]). Mammals for which no data were available, no impacts

were known or impacts were not detectable were excluded from the analysis. The median of

the three independent assignments of impact scale and confidence rating are reported for each

conflict.

Table 1. Criteria used to describe the scale of impacts in each of the four categories of conflict.

Scale of impact	A. Economic impact	B. Health impact	C. Environmental impact	D. Social impact
1	<£10K / yr	Local mild, short-term, reversible effects to individuals	Local, short-term, population loss, no significant ecosystem effect	Minimal social disruption
2	£10K – 100K / yr	Mild short-term, reversible effects to identifiable groups, localised	Some ecosystem impact, reversible changes, localised	Significant concern expressed at local level
3	£100K – 1M / yr	Minor irreversible effects and/or larger numbers covered by reversible effects, localised	Measurable long-term damage to populations and ecosystem, but little spread, no extinction	Temporary changes to normal activities at local level
4	£1M – 10M / yr	Significant irreversible effects locally or reversible effects over a large area	Long-term irreversible ecosystem change, spreading beyond local area	Some permanent change of activity locally, concern over a wider area
5	>£10M / yr	Widespread, severe, long-term, irreversible health effects	Widespread, long-term population loss or extinction, affecting several species with serious ecosystem effects	Long-term social change, significant loss of employment, migration from area

179   **Table 2.** Criteria used for the confidence ratings of the impact scale assignments.

Confidence level	Example evidence
High (3) (approx. 90% chance of impact scale assignment being correct)	There is direct relevant observational evidence to support the assessment; **and** Impacts are recorded at the typical spatial scale over which the conflict can be characterised; **and** There are reliable / good quality data sources on the impacts of the conflict; **and** The interpretation of the data / information is straightforward; **and** Data / information is not controversial or contradictory
Medium (2) (approx. 65-75% chance of impact scale assignment being correct)	There is some direct observational evidence to support the assessment but some information is inferred; **and/or** Impacts are recorded at a spatial scale, which may not be relevant to the scale over which conflict can be characterized, but extrapolation or downscaling of the data to relevant scales is considered reliable, or to embrace little uncertainty; **and/or** The interpretation of the data is to some extent ambiguous or contradictory
Low (1) (approx. 35% chance of impact scale assignment being correct)	There is no direct observational evidence to support the assessment, e.g. only inferred data have been used as supporting evidence; **and/or** Impacts are recorded at a spatial scale, which is unlikely to be relevant to the scale over which conflict can be characterized; extrapolation or downscaling of the data to relevant scales is considered unreliable potentially leading to significant uncertainties; **and/or** Evidence is poor and difficult to interpret, e.g. because it is strongly ambiguous; **and/or** The information sources are considered to be of low quality or contain information that is unreliable

Various species characteristics were also included in the analyses to determine whether

they were associated with a conflict impact (modified from Nentwig et al. [15]). Impact scores

were considered as the maximum impact for a particular species in Great Britain and did not

account for their current distribution. Overall impacts were compared between species by

summing the median scores of all conflicts and within each category ("potential impact"). To

consider the distribution of a species, potential impact values were multiplied by the proportion

of hectads (10 kilometre square resolution) each species occupies in Great Britain (derived

from Crawley et al. [38]). The potential impact scores, corrected for the area occupied can be

considered as the "actual impact" a species causes in Great Britain. Estimated population sizes

were derived mostly from Mathews et al. [39], typical body weights were taken mostly from

Harris & Yalden [40]. Where information was missing, relevant online sources were used (see

S2 Table). The overall biomass for each species was calculated by multiplying population

estimates and body mass. Population estimates and/or distribution data could not be reliably

established for three species (feral cat, feral ferret, greater white-toothed shrew), therefore they

were excluded from some of the statistical analyses.

Statistical analysis

All statistical analyses were performed using the R statistical programme version 4.2.2 [41] and

the RStudio Integrated Development Environment (IDE: version 2022.12.0, Posit Software,

PBC). Given the nature of the response variables, initial analyses (to assess whether

impact/confidence scores assigned by the three reviewers differed overall and in relation to the

conflict species assessed) were performed using an ordinal logistic regression model fitted

using the *ordinal::clm* function version 2023.12-4. [42]. Species were aggregated based on

status (native/non-native/feral) and taxonomic order, with an interaction term between each

aggregate type and reviewer ID included. Additional analyses were performed to assess

whether impact scores assigned by the reviewers differed across impact categories (i.e.

economic, environmental, health and social), across subcategories, between native/non-native

species and between taxonomic orders, including interaction terms. (N.B. feral species were

removed from further analyses due to their low number of conflicts). Data were expanded to

account for the disparity in the number of subcategories described for each species from the

literature, so that for each species, a total of 32 subcategories were present, with an impact and

confidence score of zero (0), representing no available data, assigned to absent subcategories.

Differences in potential and actual impact scores were analysed using generalised linear

models (GLM), using the *glmmTMB* package version 1.1.10 [43]. Diagnostic testing of these

analyses was performed using the *DHARMa* package version 0.4.7 [44]. Prior to analysis, data

were modified to account for species whereby an actual impact score could not be reliably

established, with a score of zero (0) assigned to indicate insufficient evidence of impact. Feral

ferret was subsequently excluded from the analyses, given a lack of data regarding population

size and geographic distribution. Analyses assessed whether estimates of total impact (actual

or potential) differed in relation to native/non-native status and taxonomic order, when

modelled independently. Several covariate predictors were also specified independently in

each model, including population size, body mass (kg) and biomass (kg). Models consisted of

GLMs, fitted with a gaussian error distribution and "identity" link function, selected given that

model residuals conformed to key assumptions (i.e. linearity, normality, homogeneity of

variance), when total impact estimates were logarithmically transformed [$\log(n+1)$]. For each

model, a two-way interaction term was specified between species aggregation type (either

native/non-native status or taxonomic order) and each covariate predictor (population size,

body mass, biomass). Due to issues with model performance, population size, body mass and

biomass were also log-transformed [log(n+1)].

Additional analyses examined how estimates of impact (actual or potential) might differ

across the main impact categories (economic, health, environment, social); these were

performed using a GLM fitted with a Tweedie distribution, which is appropriate when analysing

a continuous response variable consisting of a combination of zeroes and positive values.

These models compared impact across categories in relation to native/non-native status and

order, as well as population size, body mass and biomass. Interaction terms involving impact

category, species aggregation type and each covariate predictor were included with respect to

native/non-native species only. Analyses of impact, in relation to impact categories and order

and covariate predictors, were limited to main order effects only, due to persistent issues with

model performance (i.e. failure of models to converge), when considering any possible

interaction between both predictor variables. This is likely due to the complexity of the models

(impact category=4 levels; taxonomic order=6 levels; total number of factorial

combinations=24) when considering interaction terms and the complete separation of certain

impact categories across certain taxonomic orders, caused by the presence of absolute zeroes

(e.g. social impact and Rodentia).

Results

General literature searches

A total of 13,148 articles were returned for potential human-wildlife conflicts involving the 66

terrestrial mammals in the UK. Further screening identified 405 useful articles for 48 mammals

belonging to seven taxonomic groups (Rodentia: n=15; Carnivora: n=11; Artiodactyla: n=8;

Chiroptera: n=6; Eulipotyphla: n=4; Lagomorpha: n=3; Perissodactyla: n=1). The first available

record was from 1947 and the latest was from 2024. The majority of articles (n=367, 91%) were

published within the last three decades (2014-2024: n=176, 44%; 2004-2013: n=110, 27%;

1994-2003: n=81, 20%).

A detailed description of all of the conflicts is in S1 File. A summary of characteristics

(native/non-native/feral status, population size, body mass, biomass) and conflicts for each

species is in S2 Table. The periods during which conflicts were reported in the scientific

literature (i.e. identified during the original review only ("pre-2014"), during the original and

current review ("pre-2014 and 2014 to present") and during the current review only ("2014 to

present") are also shown. The median impact scores and confidence ratings are shown after

each conflict in parentheses.

Overview of conflicts

Conflicts and status

Twenty-nine of the 48 mammals (61%) are native to the British Isles, 17 (35%) are non-native

(which includes 10 that have been introduced in recent times and seven that are considered

naturalised) and two (4%) are feral (species that are considered wild but are either managed or

may have human dependence). Of the 48 mammals, 40 (83%) caused at least one economic

conflict, 36 (75%) caused at least one health conflict, 23 (48%) caused at least one

environmental conflict and eight (17%) caused at least one social conflict (Fig 1). When

compared by status, 86% (25/29) of native mammals caused economic conflict, 66% (19/29)

caused health conflict and 35% (10/29) caused environmental conflict. In comparison, 82%

(14/17) of non-native mammals caused economic conflict, 88% (15/17) caused health conflict

and 71% (12/17) caused environmental conflict. Social conflicts accounted for approximately

17% for both native (5/29) and non-native (3/17) species. The proportion of feral mammals

causing conflict was either 50% (economic and environment) or 100% (health), although this

was attributable to only two species (feral cat, Dartmoor/Exmoor pony).

Fig 1. Percentage of mammals within each status causing conflict within the four main
categories. Data labels denote the number of mammals within each status.

Frequency of conflicts

In total, 200 separate conflicts were identified, with 103 attributable to native mammals (52%),

92 to non-native mammals (46%) and five to feral mammals (3%). Ninety-one of the 200

conflicts (45%) were economic impacts (Appendix 1a), 71 (35%) were health impacts

(Appendix 1b), 29 (15%) were environmental impacts (Appendix 1c) and nine (5%) were social

impacts (Appendix 1d). A similar proportion of conflicts were recorded for both native and non-

native species within all four categories (Fig 2). For native mammals, economic impacts

accounted for 47% (48/103), health impacts accounted for 36% (37/103), environmental

impacts accounted for 11% (12/103) and social impacts accounted for 6% (6/103). For non-

native mammals, economic impacts accounted for 45% (42/92), health impacts accounted for

36% (32/92), environmental impacts accounted for 16% (15/92) and social impacts accounted

for 3% (3/92). Feral mammals were involved in considerably less conflicts, mostly attributable

to health and environmental impacts.

Fig 2. The percentage of conflicts within each category depending on status. Data labels denote
the number of conflicts within each category. Native mammals: n=103; non-native mammals:
n=92; feral mammals: n=5.

    The total number of mammals causing conflict within a particular subcategory ranged from

one to 30 (Appendix 1a-1d). Two of the four most common conflicts were health related

(reservoir of zoonotic disease, n=30; reservoir of disease, n=22) followed by two economic

conflicts (damage to agricultural interests, n=19; damage to forestry, n=15), with a similar

number of native and non-native mammals causing conflict in these subcategories. However,

when these data were analysed separately according to status, non-native mammals

accounted proportionately more than native mammals in all four subcategories (Fig 3).

Fig 3. The percentage of native and non-native mammals causing conflict within the four most common conflict subcategories. Data labels denote the number of mammals. Native: n=29; non-native: n=17.

Impact scores and ranking of conflicts

Reproducibility of the scoring

Of the 200 conflicts, the three reviewers agreed on the impact score in 92 cases (46%); there was a maximum deviation of one score in 89 cases (44%), a maximum deviation of two scores in 18 cases (9%) and a maximum deviation of three scores in 1 case (1%). Overall, there was no statistically significant difference between the independently assigned impact scores (χ^2(2)=3.057, p=0.217). However, confidence scores were found to differ statistically between reviewers (χ^2(2)=144.54, p<0.001); all three agreed in 33 of the 200 conflicts (17%), two agreed in 131 cases (65%), with a deviation of one or two assigned by the third reviewer, and all three assigned a different confidence score in 36 cases (18%). This suggests that the reviewers were largely consistent in their impact scoring of British mammals but not necessarily in the confidence of the assignments.

Conflict scoring and ranking

Using the median impact scores across the three reviewers, seven of the 200 conflicts (4%)

were assessed as impact score 5, 20 (10%) as impact score 4, 30 (15%) as impact score 3, 88

(44%) as impact score 2 and 55 (27%) as impact score 1 (Table 4a-4d). Six of the seven

conflicts assessed as having the highest impact score of 5 (all with confidence rating of high)

were economic and one was a health conflict. These were (in order of highest approximate

economic cost): (1) rabbit damage to agricultural interests (£120 million), (2) badger

involvement in the transmission of bTB to cattle (>£100 million), (3) brown rat damage to

agricultural interests (£54 million), (4) rabbit damage to forestry (£43 million) (5) badger damage

to agricultural interests (£39 million) (6) grey squirrel damage to forestry (£28 million, not

including carbon capture estimates) and (7) brown rat damage to property (£14 million, based

on local authority costs only). In all four categories, the majority of conflicts were scored as

impact 2 (economic: 35% (32/91); health: 46% (33/71); environmental: 55% (16/29); social:

78% (7/9)). Despite two health conflicts being the most common amongst all mammals,

generally, health conflicts were scored low, with over 90% (64/71) scoring either a 1 or 2.

 Table 4a. Total number of mammals causing conflict within the 15 economic subcategories and their median impact score. Status categories:
 Na=native; NN=non-native; Fe=feral.

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Damage to forestry (15)	Grey squirrel, rabbit	Roe deer, red deer, fallow deer	Japanese sika, brown hare, edible dormouse, field vole	Muntjac, feral goat, mountain hare, beaver, bank vole	Chinese water deer	Na - 6 NN - 9
Damage to agricultural interests (19)	Badger, rabbit, brown rat	Roe deer, fallow deer, mole, house mouse	Red deer	Japanese sika, Chinese water deer, muntjac, wild boar, brown hare, mountain hare, grey squirrel, beaver, wood mouse	Edible dormouse, field vole	Na - 9 NN - 10
Damage to vehicles (RTAs) (9)	-	Roe deer, fallow deer, badger	Red deer, muntjac	Japanese sika, Chinese water deer, wild boar, Exmoor/Dartmoor pony	-	Na - 4 NN - 4 Fe - 1
Competition with livestock (3)	-	-	-	-	Red deer, Japanese sika, feral goat	Na - 1 NN - 2
Predation of game species (11)	-	Red fox	-	Stoat, badger, American mink, hedgehog, brown rat	Wild cat, pine marten, feral ferret, weasel, polecat	Na - 8 NN - 3
Predation/impact on commercial fisheries (3)	-	-	-	Otter, American mink	Beaver	Na - 2 NN - 1
Predation of poultry (4)	-	-	-	American mink	Pine marten, feral ferret, polecat	Na - 2 NN - 2
Damage to property (6)	Brown rat	Badger, grey squirrel, house mouse	Edible dormouse	Natterer's bat	-	Na - 2 NN - 4
Damage to amenities and domestic grassland (4)	-	Rabbit	American mink, mole	Badger	-	Na - 2 NN=2
Conflicts with construction, infrastructure and development (9)	-	Badger	Natterer's, serotine, common pipistrelle, soprano pipistrelle, brown long-eared bat	Beaver, hazel dormouse, water vole	-	Na - 9
Predation of livestock (1)	-	Red fox	-	-	-	Na - 1
Conflicts with transport (2)	-	-	Rabbit, brown rat	-	-	NN - 2
Sustaining predators harmful to game (2)	-	-	-	-	Field vole, Orkney vole	Na - 1 NN - 1
Damage to property as a result of poaching (2)				Brown hare	Mountain hare	Na - 1 NN - 1
Conflicts with utilities (1)	-	Brown rat	-	-	-	NN - 1
Total per impact score	**6**	**18**	**17**	**32**	**18**	**91**

Table 4b. Total number of mammals causing conflict within the six health subcategories and their median impact score. Status categories: Na=native; NN=non-native; F=feral.

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Health risks from RTAs (9)	-	-	Roe deer, wild boar	Red deer, Japanese sika, fallow deer, muntjac, badger	Chinese water deer, Exmoor/Dartmoor pony	**Na - 4** **NN - 4** **Fe - 1**
Reservoir of disease (22)	-	-	Rabbit, grey squirrel	Roe deer, red deer, fallow deer, muntjac, wild boar, feral cat, badger, red fox, field vole, wood mouse, house mouse	Japanese sika, Chinese water deer, feral goat, stoat, Daubenton's, hedgehog, brown hare, red squirrel, beaver	**Na - 12** **NN - 9** **Fe - 1**
bTB transmission (8)	Badger	-	-	Fallow deer, muntjac, wild boar	Roe deer, red deer, Japanese sika, red fox	**Na - 5** **NN - 3**
Reservoir of zoonotic disease (30)	-	Brown rat	House mouse	Roe deer, red deer, fallow deer, muntjac, American mink, red fox, Daubenton's, serotine, hedgehog, rabbit, bank vole, field vole, wood mouse	Japanese sika, Chinese water deer, feral goat, polecat, common shrew, greater white-toothed shrew, brown hare, mountain hare, red squirrel, grey squirrel, beaver, edible dormouse, yellow-necked mouse, Skomer vole, black rat	**Na - 15** **NN - 15**
Threat to livestock where ground is undermined (1)	-	-	-	Mole	-	**Na - 1**
Threats of branches falling (1)	-	-	-	-	Grey squirrel	**NN - 1**
Total per impact score	**1**	**1**	**5**	**33**	**31**	**71**

346 **Table 4c.** Total number of mammals causing conflict within the seven environmental subcategories and their median impact score. Status categories:
347 Na=native; NN=non-native; F=feral.

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Damage to conservation habitats (9)	-	-	Roe dee, red deer	Japanese sika, fallow deer, muntjac, feral goat	Chinese water deer, wild boar, bank vole	**Na - 4** **NN - 5**
Threat to biodiversity (6)	-	Grey squirrel	-	Roe deer, red deer, Japanese sika, fallow deer, greater white-toothed shrew	-	**Na - 2** **NN - 4**
Threat to genetic integrity (3)	-	-	Feral cat	Japanese sika, feral ferret	-	**NN - 2** **Fe - 1**
Competition with or predation of wildlife (including protected or conservation species) (7)	-	-	Badger, American mink, red fox	Feral cat, otter, pine marten, stoat	-	**Na - 5** **NN - 1** **Fe - 1**
Predation of important ground-nesting birds and island ecosystem damage (2)	-	-	Hedgehog, brown rat	-	-	**Na - 1** **NN - 1**
Sustaining predators harmful to important conservation species (1)				Rabbit	-	**NN - 1**
Damage to island ecosystems (1)	-	-	-	-	Black rat	**NN - 1**
Total per impact score	**0**	**1**	**8**	**16**	**4**	**29**

349 **Table 4d.** Total number of mammals causing conflict within the four social subcategories and their median impact score. Status categories: Na=native;
350 NN=non-native; F=feral.

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Public nuisance (4)	-	-	-	Muntjac, feral goat, rabbit	Wild boar	**Na - 1** **NN - 3**
Domestic nuisance (2)	-	-	-	Pine marten	Red fox	**Na - 2**
Predation of domestic pets (1)	-	-	-	Red fox	-	**Na - 1**
Domestic nuisance & hygiene (2)	-	-	-	Natterer's, serotine	-	**Na - 2**
Total per impact score	**0**	**0**	**0**	**7**	**2**	**9**

Analysis of impact scores in relation to native/non-native status and conflict category,

found a statistically significant effect of category, when considered as a main order effect

($\chi^2(3)$=62.90, p<0.001) but not status ($\chi^2(1)$=0.421, p=0.516). Reviewers assigned a greater

score in relation to economic impacts, followed by environmental, health and then social

impacts (Fig 4). When conflict category was considered in relation to status, there was no

statistically significant interaction term between the two ($\chi^2(3)$=4.839, p=0.184).

Fig 4. The proportion of conflicts depending on impact score (1-5) within the four categories.
Data labels denote to total number of conflicts within each impact score. Economic: n=91,
health: n=71, health: n=29, social: n=9.

Analysis of impact scores, in relation to native/non-native status and conflict

subcategories, showed both as having a statistically significant effect ($\chi^2(1)$=33.63, p<0.001;

$\chi^2(31)$=640.18, p<0.001, respectively), with a significant two-way interaction term also

identified ($\chi^2(31)$=155.56, p<0.001). Across all economic impacts, non-native species received

a statistically significantly greater impact score than natives in relation to damage to woodland,

damage to agricultural interests, damage to property, conflict with transport and conflict with

utilities (all p<0.05). Conversely, native species received a significantly higher score due to

conflicts with construction and development and the predation of livestock (both $p<0.001$).

Across health impacts, the model predicted a statistically greater impact score for non-native

species as reservoirs of zoonotic diseases only ($p=0.008$). With regards to the environment,

non-native species received a statistically significant greater impact score for damage to

conservation habitats and their threat to biodiversity and genetic integrity (all $p<0.05$). In terms

of social impacts, non-native species received a greater impact than natives with respect to

public nuisance only ($p<0.01$).

Potential impact

The number of separate conflicts per species ranged from one to eleven (Table 5), with sika deer

involved in the most conflicts. Species scoring highest in potential total impact (i.e. the sum of

all median conflict scores for a particular species) were badger, rabbit, roe deer, fallow deer and

red deer. Generally, these species also scored highly within the separate categories (except for

social conflicts), with badgers scoring highest in both potential economic and health impacts.

The highest potential environmental impact was produced by sika deer, followed by roe deer,

red deer and feral cat. The highest potential social impact was observed in the red fox, followed

by rabbit, wild boar, pine marten, Natterer's and serotine bat.

Table 5. Distribution (i.e. the proportion of Great Britain each mammal is found, based on presence in 10x10km squares), total number of conflicts
and impact scores (potential and actual) in total and within each of the four categories (economic, health, environmental, social) for all conflict
mammals. Mammals are listed in order of highest total actual impact. – denotes missing data.

Species	Distribution	No. conflicts	Potential impact					Actual impact				
			Total potential impact	Economic	Health	Environment	Social	Total actual impact	Economic	Health	Environment	Social
Badger	0.746	10	33	21	9	3	0	24.624	15.670	6.716	2.239	0
Rabbit	0.833	8	26	17	5	2	2	21.649	14.155	4.163	1.665	1.665
Roe deer	0.726	9	25	12	8	5	0	18.151	8.713	5.808	3.630	0
Red fox	0.800	8	19	8	5	3	3	15.207	6.403	4.002	2.401	2.401
Grey squirrel	0.583	7	20	11	5	4	0	11.653	6.409	2.913	2.331	0
Brown rat	0.529	7	21	14	4	3	0	11.105	7.403	2.115	1.586	0
Red deer	0.412	10	23	11	7	5	0	9.480	4.534	2.885	2.061	0
Fallow deer	0.347	9	24	12	8	4	0	8.317	4.158	2.772	1.386	0
American mink	0.533	6	14	9	2	3	0	7.457	4.794	1.065	1.598	0
Mole	0.807	3	9	7	2	0	0	7.266	5.651	1.615	0	0
Hedgehog	0.824	4	8	2	3	3	0	6.592	1.648	2.472	2.472	0
Brown hare	0.680	5	9	7	2	0	0	6.119	4.759	1.360	0	0
Muntjac	0.334	9	17	7	8	2	0	5.673	2.336	2.669	0.667	0
Field vole	0.564	5	9	5	4	0	0	5.075	2.819	2.256	0	0
Wood mouse	0.560	3	6	2	4	0	0	3.360	1.120	2.240	0	0
Otter	0.782	2	4	2	0	2	0	3.126	1.563	0	1.563	0
House mouse	0.235	4	13	8	5	0	0	3.060	1.883	1.177	0	0
Stoat	0.597	3	5	2	1	2	0	2.983	1.193	0.597	1.193	0
Natterer's	0.411	3	7	5	0	0	2	2.880	2.057	0	0	0.823
Sika deer	0.134	11	19	8	5	6	0	2.540	1.069	0.668	0.802	0
Bank vole	0.446	3	5	2	2	1	0	2.229	0.892	0.892	0.446	0
Common pipistrelle	0.674	1	3	3	0	0	0	2.021	2.021	0	0	0
Soprano pipistrelle	0.619	1	3	3	0	0	0	1.858	1.858	0	0	0
Brown long-eared bat	0.600	1	3	3	0	0	0	1.799	1.799	0	0	0
Serotine	0.246	3	7	3	2	0	2	1.721	0.738	0.492	0	0.492
Daubenton's	0.491	2	3	0	3	0	0	1.472	0	1.472	0	0
Pine marten	0.171	4	6	2	0	2	2	1.025	0.342	0	0.342	0.342

Table 5 cont.

Species	Distribution	No. conflicts	Total potential impact	Economic	Health	Environment	Social	Total actual impact	Economic	Health	Environment	Social
				Potential impact				**Actual impact**				
Mountain hare	0.162	4	6	5	1	0	0	0.973	0.811	0.162	0	0
Water vole	0.425	1	2	2	0	0	0	0.850	0.850	0	0	0
Polecat	0.269	3	3	2	1	0	0	0.807	0.538	0.269	0	0
Weasel	0.530	1	1	1	0	0	0	0.530	0.530	0	0	0
Common shrew	0.502	1	1	0	1	0	0	0.502	0	0.502	0	0
Red squirrel	0.247	2	2	0	2	0	0	0.494	0	0.494	0	0
Feral goat	0.023	6	13	4	7	1	1	0.209	0.070	0.047	0.047	0.047
Hazel dormouse	0.199	1	2	2	0	0	0	0.397	0.397	0	0	0
Chinese water deer	0.042	7	9	5	3	1	0	0.375	0.208	0.125	0.042	0
Wild boar	0.031	7	9	3	2	2	2	0.402	0.124	0.216	0.03	0.031
Beaver	0.018	6	9	7	2	0	0	0.166	0.129	0.037	0	0
Yellow-necked mouse	0.129	1	1	0	1	0	0	0.129	0	0.129	0	0
Edible dormouse	0.008	4	8	7	1	0	0	0.064	0.056	0.008	0	0
Wildcat	0.039	1	1	1	0	0	0	0.039	0.039	0	0	0
Exmoor/Dartmoor pony	0.005	2	3	2	1	0	0	0.016	0.011	0.005	0	0
Black rat	0.006	2	2	0	1	1	0	0.011	0	0.006	0.006	0
Skomer vole	0.004	1	1	0	1	0	0	0.004	0	0.004	0	0
Orkney vole	0.003	1	1	1	0	0	0	0.003	0.003	0	0	0
Feral cat	-	3	7	0	2	5	0	-	-	-	-	-
Feral ferret	-	3	4	2	0	2	0	-	-	-	-	-
Greater white-toothed shrew	-	2	3	0	1	2	0	-	-	-	-	-

Analyses identified a statistically significant difference between native/non-native species

(χ^2(1)=4.235, p=0.040), with the latter predicted as having a higher potential impact overall

(β=0.513, z ratio=2.058, p=0.046). When specified as main order effects, total potential impact

was found to increase as a function of body mass (β=0.335, t ratio=3.598, p<0.001) and

biomass (β=0.184, t ratio=4.706, p<0.001) but not population size (β=0.040, t ratio=1.012,

p=0.311). However, no statistically significant interaction term was identified (p>0.05),

suggesting that differences in the rate of change for total potential impact were comparable

between native and non-native species, relative to changes in population size, body mass and

biomass.

There was a statistically significant difference between taxonomic orders (χ^2(5)=18.468,

p=0.002), with Artiodactyla having a higher potential impact compared to Chiroptera (β=1.129,

t ratio=3.661, p=0.009) and Rodentia (β=1.231, t ratio=3.236, p=0.028), whilst all other

taxonomic comparisons were non-significant (Fig 5). Inclusion of covariate predictors as main

order effects, revealed a statistically significant effect of population size (χ^2(1)=7.259, p=0.007)

and biomass (χ^2(1)=39.467, p<0.001), with total potential impact predicted to increase as a

function of each (population size: β=0.096, t ratio=2.499, p<0.001; biomass: β=0.199, t

ratio=5.641, p<0.001).

Figure 5. Average potential impacts within the four categories across orders (excluding Perissodactyla).

Actual impact

Species scoring the highest actual impact were badger, rabbit, roe deer and red fox (Table 5).

Generally, these species also scored highly within the separate categories, with badgers scoring

highest in both actual economic and health impacts. Roe deer scored the highest actual

environmental impact, followed by hedgehog, red fox, grey squirrel, badger and red deer. The

highest potential social impact was observed in the red fox, followed by rabbit and Natterer's

bat.

Analysis of actual impact estimates showed no statistically significant difference between

native/non-native species (χ^2(1)=0.056, p=0.813). When considered as main order effects,

total actual impact was predicted to increase significantly in relation to population size

(β=0.094 t ratio=1.980, p=0.048) and biomass (β=0.215, t ratio=4.691, p<0.001). When

considered in relation to status, there was no statistically significant interaction for biomass

(χ^2(1)=0.298, p=0.585) suggesting that the rate of increase was comparable across native and

non-native species. Body mass was found to have no statistically significant effect on total

actual impact when considered as a main order effect ($\chi^2(1)$=1.741, p=0.187) or through

interaction with status ($\chi^2(1)$=0.088, p=0.767). A statistically significant interaction term was

identified between status and population size, with total actual impact predicted to increase,

relative to population size, at a significantly higher rate for non-native species when compared

to natives (β=0.179, t ratio=2.120, p=0.034).

There was no statistical difference, in relation to actual impact scores, between the

taxonomic orders ($\chi^2(5)$=5.622, p=0.345: Fig 6).

Figure 6. Average actual impacts within the four categories across orders (excluding
Perissodactyla).

Conflict change over time

For many mammals, a conflict occurring before 2014 was still being reported in the scientific

literature within the last decade (S2 Table). Twenty-seven of the 46 (59%) conflicts reported

only before 2014 were economic, followed by 10 (22%) health, 5 environment (11%) and 4

social (8%). In contrast, 25 of the 45 (55%) conflicts reported from 2014 to 2024 were health

(55%), mostly attributable to reservoir of disease or reservoir of zoonotic disease, followed by

9 environment (20%), 8 economic (18%) and 3 social (7%).

Discussion

Negative interactions between people and wildlife are more often than not viewed as human–

wildlife conflict, with socio-economic factors cited amongst the issues influencing people's

perception of wildlife [45]. In this paper, we estimated a conservative total annual economic

cost of human-wildlife conflicts in Great Britain (as reported in S1 File) in excess of £0.5 billion,

which would likely exceed £1 billion if health and environmental conflicts could be

monetarised. All but one of the seven conflicts assessed as having the highest impact score of

5 involved economic conflicts. The one health conflict scored as impact 5 (the involvement of

badgers in the transmission of bTB), was also measurable on an economic scale, suggesting

that conflicts associated with a monetary value are inherently easier to quantify compared with

those that are not. To account for potential bias associated with individual conflicts, we applied

a prioritisation approach (generic impact scoring system (GISS)) to evaluate the overall impact

of conflict mammals. In some cases, mammals causing the greatest economic damage, such

as rabbit and badger, were ranked highest overall for their actual impact. However, this approach

also identified species that did not necessarily cause the highest individual impacts; for

example, roe deer and red fox were ranked third and fourth overall, respectively, for actual

impact despite neither receiving the highest impact score of 5 for any of their identified conflicts.

The emphasis of legislation associated with wildlife in the UK has moved from protecting

human economic interests (e.g. Prevention of Damage by Pests Act 1949 and Pests Act 1954),

to promoting animal welfare (e.g. Wildlife and Countryside Act 1981 and Animal Welfare Act

2006) and protecting biodiversity (e.g. Habitats Directive 1992, Environment Act 2021). There

is likely to be an increasing trend for conflicts arising between promoting biodiversity and human

economic interests, as human activities continue to expand into existing wildlife habitats. Four

of the five mammals identified as causing the greatest actual economic and health impact

(badger, rabbit, roe deer), environmental impact (roe deer, hedgehog, red fox), and social impact

(red fox, rabbit) are native, which leads to increasing conflict where specific legislation applies

to a protected species. For example, badgers are protected under the Protection of Badgers Act

1992 but are the focus of ongoing controversy regarding management options for controlling the

spread of *Mycobacterium bovis* to cattle in the UK, which is estimated to cost the taxpayer over

£100 million annually [46]. Likewise, some bat species are widely distributed throughout Great

Britain and often associated with conflicts with buildings or construction; as bats and their

roosts are legally protected under the Wildlife and Countryside Act 1981 and the Conservation

of Habitats and Species Regulations 2010, this can lead to potentially complex and costly

resolution.

Although native mammals were ranked amongst the highest scoring impacts in all four

categories, overall, there was no statistical difference in the impact scores between native and

non-native species. However, there was a disproportionate amount of conflict in specific areas,

with non-native species perceived as causing significantly more damage to woodlands,

conservation habitats and agricultural interests, and a greater reservoir of zoonotic disease

compared to native species. Rabbits caused the highest economic conflict compared to all

other mammals with their impact on agriculture, and ranked second overall for total actual

impact. As such, this may highlight rabbits as a priority species for human-wildlife resolution.

However, rabbit populations have declined significantly in the UK in recent years, with severe

local population reductions reported [47], which have led to concerns regarding the impact of

such population reductions on other species, such as carnivores, that rely on them as a food

source [48]. In contrast, grey squirrel abundance and distribution has expanded rapidly since

they were introduced to Great Britain in the late 19th century [49]. They are considered a pest

species in the UK and as such are afforded no protection under the Wildlife and Countryside

Act 1981. As part of its 25-year Environment plan [20], the government has committed to

managing grey squirrels to protect woodlands and red squirrel populations in the UK. While the

general consensus is that grey squirrels must be managed to protect biodiversity within the UK,

removing healthy wildlife can be controversial and may face opposition from pressure groups,

which can be key to the success or failure of an eradication project [50].

Although the majority of reported conflicts were economic, the two most common

conflicts were health related (reservoir of zoonotic disease and reservoir of disease). More than

60% of human pathogens are zoonotic in origin [51] with urbanisation and anthropogenic

factors, amongst others, thought to have greatly influenced the emergence and distribution of

zoonoses [52]. Generally, though, the impact score of health conflicts were deemed low

(mostly 1 or 2), therefore we assumed that most health impacts were likely local, mild, short-

to mid-term and with reversible effects (see Table 1). This highlights the importance of

surveillance, which is crucial to preventing and controlling the spread of disease, in particular

zoonotic disease [52]. This is also pertinent to exotic diseases, which are not reported here as

only current diseases causing conflict in Britian were assessed. Whilst there is extensive

research on the pathology and epidemiology of disease in a wide range of British mammals, as

reported in S1 File, it is evident that for some mammals, there is an inference of disease

presence and/or zoonotic potential or a general lack of information (e.g. some hantaviruses [53]

and Ljungan virus [54] in rodents; *Babesia* spp. in deer [55] and badgers [56]), which suggests

further research may be warranted. Understanding disease dynamics in British wildlife is

intrinsically linked to biodiversity conservation, as anthropogenic activities such as urbanisation

and habitat destruction, are likely to lead to increased interaction between humans and wildlife

[57].

In many cases, mammals that caused the most conflict were amongst the most widely

distributed throughout the British Isles, e.g. badger, rabbit, roe deer and red fox. Our analyses

identified that total actual impact tended to increase as a function of biomass and population

size, therefore a GISS approach may be particularly beneficial for monitoring recently

introduced species, whether native or non-native, which have the potential to expand beyond

their current range. Beavers were reintroduced to Great Britain around 20 years ago and are

considered keystone species due to their ability to alter ecosystem structure and function

through their engineering activities [58, 59]. Whilst beavers are generally considered as having

a positive impact on biodiversity, we identified six potential conflicts of interest with humans;

whilst they ranked 38[th] overall based on total actual impact, likely owing to their current limited

distribution, they ranked joint 12[th] overall based on potential impact. Whilst the reintroduction

of beavers across Great Britain is currently carefully managed, highlighting and monitoring

potential issues at this stage is essential for ensuring the status of this species remains positive

rather than possible future perception as a pest species.

The overall assessments of the conflicts reported here were made at the national level for

Great Britain. Local impacts may be of considerable importance for those affected, for instance

those injured in road traffic accidents, and thus a national score of 1 or 2 should not be taken to

indicate that the conflict is not important or worth mitigating at a local level. Furthermore, it is

important to recognise that the impacts are not necessarily uniformly distributed geographically

or equitably shared. For example, Artiodactyla were predicted to receive significantly higher

potential impact scores compared to other taxonomic orders. The total potential impact for five

of the six deer species were amongst the highest of all the mammals, although the actual

impact for some deer species may be lower depending on their distribution. For example, sika

deer were involved in the most conflicts of all the mammals but were ranked 20[th] overall for

actual impact, suggesting its impacts are localised given its limited distribution.

Human-wildlife conflict has long been associated with negative impacts on human

interests, although it is evident that interactions between people and wildlife are more varied

and complex [2]. More recently, the emphasis has shifted from conflict between humans and

wildlife and its mitigation, to a coexistence approach [14], or at least one that constrains

conflict or adapts shared spaces to mitigate conflict. Whilst there is a general consensus that

some species must be managed or even eradicated (e.g. invasive non-native species: Nentwig

et al. [15]), for others, particularly native species, a more holistic approach may be required. In

some cases, mammals that represent a conflict and cost to some, may also provide a benefit

to others. In Great Britain, the red fox has been widely persecuted due to its perceived status as

a pest species [60]. Their impact on agriculture has been estimated to cost £20 million annually,

most of which is attributable to direct and indirect effects of livestock predation [61]. However,

fox predation on rabbits, which have a much more significant impact on agriculture, is estimated

to create an indirect economic benefit to farmers of at least £11 million annually and may even

be approaching an economically neutral impact on agriculture [61].

The findings reported here must be viewed in light of several limitations. Although an

extensive literature search was conducted to identify conflicts associated with British wildlife,

it is likely that some conflicts were not identified or cost estimates were not available, and

therefore not reported. Whilst the conflict descriptions given in S1 File are not exhaustive and

based only on the searches returned, they likely represent reasonable overviews and are

sufficiently robust to provide an overall audit of the current national importance of conflicts

between human and wildlife interests in Great Britain. However, individual assignments of scale

may need to be revised as further information emerges. Conflicts that were identified before

2014 only are more than likely still occurring or may not be reported as frequently in the

scientific literature. Likewise, conflicts reported after 2014 only may have been occurring prior

to this and again, either missed in the original literature searches or less frequently reported.

The majority of "new" conflicts we identified were disease related, which may reflect a shift of

focus leading to increased reporting in this area, possibly in light of several international public

health emergencies of zoonotic origin within the last ten years, such as Middle East respiratory

syndrome (MERS), Ebola and Covid-19 pandemics [62].

Another limitation of our findings is the lack of data for some species. Feral species were

excluded from some of the analyses due to the small number of conflicts in Great Britain,

although their impacts can be significant. Data regarding feral cats are limited due to the

difficulty in differentiating between feral and domestic cats [63], the uncertainties of feral cat

population estimates (based mostly on urban populations) and the lack of known distribution

within Great Britain. Cats are one of the most ubiquitous mammals globally (see McDonald &

Skillings [64]) and one of the most destructive, with free-ranging cats suggested as having a

major impact on the mortality of several bird species in the UK [65]. They are also the biggest

threat to the survival of Scottish wildcats due to introgression [66]. Given their domesticated

origins, feral cats have no native range and their ubiquitous adverse impacts on native wildlife

around the world has led to them being labelled amongst the world's worst invasive species

[67]. From a biodiversity conservation perspective, a wealth of evidence suggests that feral cats

should be controlled and ideally removed from places where their impact is detrimental, often

as an urgent priority [68].

When applying the GISS in this paper, none of the conflicts were weighted, i.e. they were all

considered of equal importance. This follows Nentwig et al. [15], who stated that in the absence

of objective scientific criteria to consider one impact category more important than the other,

selective weights were not applied. They also stated that deciding relative impact weights is a

social process rather than a scientific one. It is possible to apply individual weights to conflicts,

for example, by considering stakeholder values (e.g. Kumschick et al. [69]), which will likely

differ according to the relevance of a particular conflict to a particular group. The scoring system

applied here is thus considered robust to the uncertainty of impact knowledge but could be

adjusted with weight scores to account for specific social values, such as stakeholder interests,

in the future [15].

Conclusions

The prioritisation approach we applied here is a useful tool for identifying individual conflicts for

individual species but also provides a broader assessment of where human-wildlife impacts

may be affecting socio-economic and biodiversity interests. It is clear that it should not be relied

upon in isolation to identify priority species for mitigation purposes, and all other available

evidence or opinion from conservation groups, stakeholder groups, public perception and other

interested parties, should be considered. Ideally, any decision making, whether relevant to

government policies or conservation organisations, should consider ensuring a balance

between the need to conserve and enhance biodiversity, the interests of the general public and

the interests of those directly affected [35].

Acknowledgements

We are grateful to Eloise Cunningham for her assistance with the literature searches.

Author contributions

Conceptualisation and methodology: DC, KLP. Analysed the data: KP, DW, DC, GC. Writing-original draft: KP. Writing-review & editing: KP, DC, DW, GC.

References

1.   Locke H, Ellis EC, Venter O, Schuster R, Ma K, Shen X, et al. Three global conditions for biodiversity conservation and sustainable use: an implementation framework. National Science Review. 2019;6:1080-82.

2.   Hill CM. Conflict is integral to human-wildlife coexistence. Frontiers in Conservation Science. 2021;2:734314.

3.   Woodroffe R, Thirgood S, Rabinowitz A. People and wildlife: conflict or coexistence? Cambridge, UK: Cambridge University Press; 2005.

4.   Conover MR. Resolving human-wildlife conflicts: the science of wildlife damage management: CRC Press, Boca Raton, FL; 2002.

5.   Treves A, Karanth KU. Human-carnivore conflict and perspectives on carnivore management worldwide. Conservation Biology. 2003;17:1491–99.

6.   Tilman D. Global environmental impacts of agricultural expansion: the need for sustainable and efficient practices. The Proceedings of the National Academy of Sciences. 1999;96:5995–6000.

7.   Nyhus PJ. Human-wildlife conflict and co-existence. Annual Review of Environment and

Resources. 2016;41:143-71.

8.   Pérez E, Pacheco LF. Damage by large mammals to subsistence crops within a

protected area in a montane forest of Bolivia. Crop Protection. 2006;25:933-39.

9.   Fang L, Hong Y, Zhou Z, Chen W. The frequency and severity of crop damage by wildlife

in rural Beijing, China. Forest Policy and Economics. 2021;124:102379.

10.   Drexler JF, Corman VM, Drosten C. Ecology, evolution and classification of bat

coronaviruses in the aftermath of SARS. Antiviral Research. 2014;101:45-56.

11.   Song Z, Xu Y, Bao L, Zhang L, Yu P, Qu Y, et al. From SARS to MERS, thrusting

coronaviruses into the spotlight. Viruses. 2019;11.

12.   Hao YJ, Wang YL, Wang MY, Zhou L, Shi JY, Cao JM, et al. The origins of COVID-19

pandemic: A brief overview. Transboundary and Emerging Diseases. 2022;69(6):3181-197.

13.   Cassini MH. Human-wildlife conflicts: does origin matter? . Animals. 2022;12:1-10.

14.   Gross EM, Jayasinghe N, Dahal S, Tenzin S, Klenzendorf S, Vannelli K, et al. C2C—

conflict to coexistence: a global approach to manage human–wildlife conflict for coexistence.

Conservation Science and Practice. 2024;7:e13292.

15.   Nentwig W, Künel E, Bacher S. A generic impact-scoring system applied to alien

mammals in Europe. Conservation Biology. 2010;24:302–11.

16.   Eschen R, Kadzamira M, Stutz S, Ogunmodede A, Djeddour D, Shaw R, et al. An updated

assessment of the direct costs of invasive non native species to the United Kingdom. Biological

Invasions. 2023;25:3265-76.

17.   The Wildlife Trust. Local to global: the Global Biodiversity Framework and what the UK

needs to do to implement it. 2024; https://www.wildlifetrusts.org/sites/default/files/2024-

09/UK%20and%20Global%20Biodiversity%20Framework%20-

%20The%20Wildlife%20Trusts%202024.pdf.

18.   Dueñas MA, Hemming DJ, Roberts A, Diaz-Soltero H. The threat of invasive species to

IUCN-listed critically endangered species: a systematic review. Global Ecology and

Conservation. 2021;26:e01476.

19.   NNSS.   GB   non-native   species   report   card   2020.   2020;

https://www.nonnativespecies.org/assets/NNSIP-Scorecard-2020.pdf.

20.   HMGovernment. A green future: our 25 year plan to improve the environment. 2018;

https://assets.publishing.service.gov.uk/media/65fd713d65ca2f00117da89e/CD1.H_HM_Go

vernment_A_Green_Future_Our_25_Year_Plan_to_Improve_the_Environment.pdf; 2018.

21.   Crowley SL, Hinchliffe S, McDonald RA. Conflict in invasive species management.

Frontiers in Ecology and the Environment. 2017;15:133-41.

22.   Jackson DB, Green RE. The importance of the introduced hedgehog (*Erinaceus*

europaeus) as a predator of the eggs of waders (Charadrii) on machair in South Uist, Scotland.

Biological Conservation. 2000;93:333–48.

23.   Manchester SJ, Bullock JM. The impacts of non-native species on UK biodiversity and

the effectiveness of control. Journal of Applied Ecology. 2000;37:845-64.

24.   Ricciardi A, Cohen J. The invasiveness of an introduced species does not predict its

impact. Biological Invasions. 2007;9:309–15.

25.   Ricciardi A, Hoopes MF, Marchetti MP, Lockwood JL. Progress toward understanding the

ecological impacts of nonnative species. Ecological Monographs,. 2013;83:263–82.

26.   Swift BMC, Barron ES, Christley R, Corbetta D, Grau-Roma L, Jewell C, et al.

Tuberculosis in badgers where the bovine tuberculosis epidemic is expanding in cattle in

England. Scientific Reports. 2021;11:20995.

27.   Kumschick S, Gaertner M, Vilà M, Essl F, Jeschke JM, Pyšek P, et al. Ecological impacts

of alien species: quantification, scope, caveats, and recommendations. Bioscience.

2015;65:55–63.

28.   Nentwig W, Bacher S, Pyšek P. The generic impact scoring system (GISS): a

standardized tool to quantify the impacts of alien species. Environmental Monitoring and

Assessment. 2016;188:315.

29.   Van der Veer G, Nentwig W. Environmental and economic impact assessment of alien

and invasive fish species in Europe using the generic impact scoring system. Ecology

Freshwater Fish. 2014(4):646-56.

30.   Vaes-Petignat S, Nentwig W. Environmental and economic impact of alien terrestrial

arthropods in Europe. NeoBiota. 2014;22:23–42.

31.   Laverty C, Nentwig W, Dick JTA, Lucy FE. Alien aquatics in Europe: assessing the relative

environmental and socio-economic impacts of invasive aquatic macroinvertebrates and other

taxa. Management of Biological Invasions. 2015;6:341–50.

32.   Novoa A, Kumschick S, Richardson DM, Rouget M, Wilson JRU. Native range size and

growth form in Cactaceae predicts invasiveness and impact. NeoBiota. 2016;30:75-90.

33.   Sohrabi S, Pergl J, Pyšek P, Foxcroft LC, Gherekhloo J. Quantifying the potential impact

of alien plants of Iran using the Generic Impact Scoring System (GISS) and Environmental

Impact Classification for Alien Taxa (EICAT). Biological Invasions. 2021;23:2435-449.

34.   Evans T, Blackburn TM, Jeschke JM, Probert AF, Bacher S. Application of the Socio-

Economic Impact Classification for Alien Taxa (SEICAT) to a global assessment of alien bird

impacts. NeoBiota. 2020;62:123-42.

35.   Defra. Wildlife management in England: a policy making framework for resolving human-

wildlife   conflicts   (withdrawn).   2010;

https://www.thenbs.com/publicationindex/documents/details?Pub=DEFRA&DocId=295502.

36.   Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of Google Scholar in evidence

reviews and its applicability to grey literature searching. PLOS One. 2015;7:e013823.

37.   Hawkins CL, Bacher S, Essl F, Hulme PE, Jeschke JM, Kühn I, et al. Framework and

guidelines for implementing the proposed IUCN Environmental Impact Classification for Alien

Taxa (EICAT). Diversity and Distributions. 2015;21:1360-63.

38.   Crawley D, Coomber F, Kubasiewicz L, Harrower C, Evans P, Waggitt J, et al. Atlas of the

mammals of Great Britain and Northern Ireland. The Mammal Society: Pelagic Publishing, UK;

2020. 192 p.

39.   Mathews F, Kubasiewicz LM, Gurnell J, Harrower CA, McDonald RA, Shore RF. A review

of the population and conservation status of British mammals: technical summary. A report by

the Mammal Society under contract to Natural England, Natural Resources Wales and Scottish

Natural Heritage. Peterborough: Natural England; 2018.

40.   Harris S, Yalden DW. Mammals of the British Isles: handbook 4th edition. Southampton:

The Mammal Society; 2008.

41.   R Core Team. R: A language and environment for statistics computing. In: Computing

RFfS, editor. Vienna, Austria2022.

42.   Christensen R. *ordinal* - Regression Models for Ordinal Data. R package version

2023.12-4.1. 2023; https://cran.r-project.org/web/packages/ordinal/ordinal.pdf.

43.   Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al.

glmmTMB balances speed and flexibility among packages for zero-inflated generalised linear

mixed modelling. The R Journal. 2017;9:378-400.

44.   Hartig F. *DHARMa*: Residual diagnostics for hierarchical (multi-level/mixed) regression

models.   R   package   version   0.4.7.   2024;   https://cran.r-

project.org/web/packages/DHARMa/vignettes/DHARMa.html.

45.   Bhatia S, Redpath SM, Suryawanshi K, Mishra C. Beyond conflict: exploring the

spectrum of human–wildlife interactions and their underlying mechanisms. Oryx.

2020;54(5):1-8.

46.   Defra. Government to end badger cull with new TB eradication strategy. 2024;

https://www.gov.uk/government/news/government-to-end-badger-cull-with-new-tb-

eradication-

strategy#:~:text=The%20new%20strategy%20will%20mark,the%20end%20of%20this%20p

arliament.

47.   Petrovan SO, Ward AI, P. W. Detectability counts when assessing populations for

biodiversity targets. PLOS One. 2011;6:8.

48.   Sainsbury KA, Shore RF, Schofield H, Croose E, Campbell RD, McDonald RA. Recent

history, current status, conservation and management of native mammalian carnivore species

in Great Britain. Mammal Review. 2019;49(2):171-88.

49.   Gurnell J, Hare EJ. Rodents: Order Rodentia. In: Harris S, Yalden DW, editors. Mammals

of the British Isles. Southampton: The Mammal Society; 2008. p. 55-199.

50.   Bremner A, Park K. Public attitudes to the management of invasive non-native species

in Scotland. Biological Conservation. 2007;139:306-14.

51.   Taylor LH, Latham SM, Woolhouse ME. Risk factors for human disease emergence.

Philosophical Transactions of the Royal Society of London Series B, Biological Sciences.

2001;356:983–89.

52. Rahman T, Sobur A, Islam S, Ievy S, Hossain J, El Zowalaty ME, et al. Zoonotic diseases:

etiology, impact, and control. Microorganisms. 2020;8:1405.

53. Vapalahti O, Mustonen J, Lundkvist Å, Henttonen H, Plyusnin A, Vaheri A. Hantavirus

Infections in Europe. Lancet Infectious Diseases. 2003;3:653–61.

54. Salisbury AM, Begon M, Dove W, Niklasson B, Stewart JP. Ljungan virus is endemic in

rodents in the UK. Archives of Virology. 2014;159:547-51.

55. Gray A, Capewell P, Zadoks R, Taggart MA, French AS, Katzer F, et al. Wild deer in the

United Kingdom are a potential reservoir for the livestock parasite *Babesia divergens*. Current

Research in Parasitology & Vector-Borne Diseases. 2021;1:100019.

56. Guardone L, Ebani VV, Verin R, Nardoni S, Consolazione A, Bennett M, et al. Molecular

detection of arthropod-borne pathogens in Eurasian badgers (*Meles meles*) from the United

Kingdom. Animals. 2020;10:446.

57. Bedenham G, Kirk A, Luhano U, Shields A. The importance of biodiversity risks: link to

zoonotic diseases. British Actuarial Journal. 2022;27(e10):1-9.

58. Puttock A, Graham HA, Cunliffe AM, Elliott M, Brazier RE. Eurasian beaver activity

increases water storage, attenuates flow and mitigates diffuse pollution from intensively-

managed grasslands. Science of the Total Environment. 2017;576:430-43.

59. Auster RE, Puttock A, Brazier R. Unravelling perceptions of Eurasian beaver

reintroduction in Great Britain. Area. 2019;52:364-75.

60. Macdonald DW, Johnson PJ. Foxes in the landscape: hunting, control, and economics.

In: Macdonald DW, Feber RE, editors. Wildlife Conservation on Farmland: Volume 2, Conflict in

the Countryside. Oxford: Oxford University Press; 2015. p. 20-46.

61. Baker P, Harris S, White PCL. After the hunt: the future for foxes in Britain. London:

International Fund for Animal Welfare. 2006; https://mknhs.org.uk/wp-

content/uploads/2015/08/IFAW-after-the-hunt.pdf; 2006.

62. Sun ZS, Wan EY, Agbana YL, Zhao HQ, Yin JX, Jiang TG, et al. Global One Health index for

zoonoses: A performance assessment in 160 countries and territories. iScience.

2024;27:109297.

63. Hand A. Estimating feral cat densities using distance sampling in an urban environment.

Ecological Evolution. 2019;9:2699–705.

64. McDonald JL, Skillings E. Human influences shape the first spatially explicit national

estimate of urban unowned cat abundance. Scientific Reports. 2021;11:20216.

65. Baker PJ, Molony SE, Stone E, Cuthill IC, Harris S. Cats about town: is predation by free-

ranging pet cats Felis catus likely to affect urban bird populations? . Ibis. 2008;150:86-99.

66. Fredrikson A. Of wildcats and wild cats: troubling species-based conservation in the

Anthropocene. Environment and Planning D: Society and Space. 2016;34(4):1-25.

67. Lowe S, Browne M, Boudjelas S. 100 of the world's worst invasive alien species: a

selection from the Global Invasive Species database. Invasive Species Specialist Group (ISSG),

International Union for Conservation of Nature. 2000;

https://portals.iucn.org/library/sites/library/files/documents/2000-126.pdf.

68. Trouwborst A, McCormack PC, Camacho EM. Domestic cats and their impacts on

biodiversity: A blind spot in the application of nature conservation law. People and Nature.

2019;2:235-50.

69. Kumschick S, Bacher S, Dawson W, Heikkila J, Sendek A, Pluess T, et al. A conceptual

framework for prioritization of invasive alien species for management according to their impact.

NeoBiota. 2012;15:69-100.

Supporting information

S1 File. Detailed description of all of the conflicts identified in the scientific literature for 48 mammals of the British Isles.

S2 Table. List of species causing conflict, including order, status, population size, body mass, biomass and type of conflicts recorded over time (pre-2014 only; both pre-2014 and 2014 to present; 2014 to present only). The median impact score (1=minimal known impact to 5=highest known impact) and confidence rating (1=low; 2=medium; 3=high) are shown after each conflict [x,x]. The highest impact scores are shown in bold.