1

3   Identifying and prioritising conflicts between human and wildlife interests in Great

4   Britain

8   Kate L. Palphramand[1]*, Dave Cowan[1], Daniel A. Warren[1] & Graham C. Smith[1]

12   [1] Animal and Plant Health Agency, York, YO41 1LZ, UK

14   *Corresponding author

15   E-mail: kate.palphramand@apha.gov.uk (KP)

Abstract

We reviewed the scientific literature to identify human-wildlife conflicts associated with British terrestrial mammals to summarise their magnitude and, where possible, costs. Conflicts were classified as either economic, health, environmental or social, further divided into 32 subcategories. We applied a Generic Impact Scoring System (GISS), whereby individual conflicts were scored on a scale of 1-5 (minimal to highest known impact); these were used to determine the potential impact (sum of all conflicts) and actual impact (sum of all conflicts corrected for geographical distribution) of individual mammals so they could be ranked on their level of impact. Forty-eight mammals were involved in a total of 200 separate conflicts. The highest ranked conflicts (scoring 5) were all measurable on an economic scale (involving rabbit, badger, brown rat and grey squirrel). We estimated a conservative annual economic cost of human-wildlife conflicts in excess of £0.5 billion, which could exceed £1 billion if health and environmental conflicts could be monetarised. The most common conflicts were health related (reservoir of disease and zoonotic disease), although generally, health conflicts were scored low (mostly 1 or 2). Overall, there was no statistical difference in impact score depending on status (native or non-native); however, when considered by subcategory, non-native species scored higher in relation to damage to woodland, agricultural interests and property, and reservoir of zoonotic disease, whilst native species caused more conflict with construction and development, and predation of livestock. Non-native species had a higher potential impact score compared to native species, but not actual impact. However, actual impact was predicted to increase, relative to population size, at a higher rate for non-native species compared to native, highlighting the need for surveillance and monitoring. Where mammals represent a

cost to some, they may provide a benefit to others, e.g. fox predation of rabbits, and

as such may require a holistic approach to management. We concluded that a GISS

approach is a useful tool for identifying conflict species and could form part of a

package of measures to ensure a balance between the need to conserve and enhance

biodiversity, whilst also considering the interests of those directly affected by human-

wildlife conflict, is achieved.

47 Introduction

Co-habitation between humans and wildlife occurs on more than half of the earth's

surface [1] and consequently human-wildlife interactions form an integral part of many

ecosystems worldwide. Interactions between people and wildlife are often varied and

complex and whilst the presence of wildlife may have beneficial impacts on human

interests, there is a continued focus on conflict narratives that dominates how human-

wildlife interactions are viewed [2].

Human-wildlife conflict is commonly described as conflict occurring between

people and wildlife [3], presence and/or actions of humans or wildlife that have an

adverse effect on the other [4] or threats posed by wildlife to economic activities,

human life, recreation [5] or ecosystem services. Factors such as the intensification of

agriculture and increasing human population have had major impacts on ecosystems

worldwide [6]. Agricultural growth is expected to accelerate, resulting in an increase in

farmland and subsequent decrease in the available habitat for wildlife [7]. Wildlife

damage to crops is often cited as a major global conflict [3, 8, 9], which will continue

to worsen as habitat loss precipitates an increase in human-wildlife encounters.

Population growth and habitat encroachment may also increase the risk of disease

transmission, which is of particular concern where wildlife pose zoonotic risks. For

example, the first human cases of severe acute respiratory syndrome coronavirus

(SARS-CoV) in Guangdong province, China, in 2002, were linked to wildlife found at

a live animal market [10, 11]. Subsequent serological surveys from residents living

close to bat caves in Yunnan, China, which is geographically disparate to Guangdong,

were found to be positive for SARS-related coronavirus (SARSr-CoV), indicating that

spillover events may have occurred in zoonotic CoVs [12].

Conflict issues can arise with several types of problematic species, including

native pests, invasive native species and invasive non-native species [13]. These often

induce debates on how or whether humans and wildlife can co-exist [7], and may

involve complex decisions where the conservation status of a species conflicts with its

impacts on humans [3]. Addressing and categorising the adverse effects of human-

wildlife conflicts is recognised as a challenging issue for conservation and

development efforts [14]. There is a general consensus that the negative effects

created by some wildlife species, particularly invasive non-native species, have to be

reduced [15]. Where conflicts arise between wildlife and human interests, there is often

the need to intervene to reduce or mitigate the impacts and therefore identifying what

conflicts are occurring and what their impacts are is essential for informing decision-

making and resource allocation [16]. In 2022, the UK agreed to a global policy

framework at the Conference of Parties (COP15) to the Convention on Biological

Diversity that created a package of international measures to address the loss of

global biodiversity and to restore natural ecosystems. One of the targets identified for

reducing threats to biodiversity was managing human-wildlife conflicts, which in turn

would feed into other elements of the framework, such as enhancing sustainability in

agriculture, fisheries and forestry [17]. However, identifying which species are causing

conflict and subsequently which should be targeted first can be challenging and
controversial [15].

Invasive non-native species are a major threat to biodiversity worldwide, severely
impacting areas of high endemism [18], whilst also posing a serious threat to public
health and economic activities. As of 2019, 3,224 plants and animals had been
introduced to Great Britain, costing the UK economy almost £4 billion annually [16];
10-15% of these are considered invasive with an estimated 188 invasive non-native
animals recorded in 2020 [19]. Tackling invasive non-native species is a key part of
achieving the UK government's aims under its 25-year Environment plan [20] as well
as the UK's obligation as a signatory of the Convention on Biological Diversity
(https://www.cbd.int/convention/text/). However, even managing conflict caused by
invasive non-native species can be controversial, resulting in debates on achievability,
efficiency, social fairness and ethical implications [21]. For example, the removal of
hedgehogs from South Uist, where they were introduced to control garden pests but
consequently contributed significantly to the decline of wading shorebirds [22] resulted
in widespread criticism and opposition, given that hedgehogs are a popular icon of
British wildlife and are thought to be declining nationally [21].

Much of the available literature on wildlife conflicts in the UK, both in terms of
economic and biodiversity impacts, focuses on invasive non-native species [e.g. 16,
23]. However, there is evidence to suggest that the most invasive non-native species
are not necessarily the most damaging [24, 25], therefore there is the need to consider
also the impacts of native species, some of which can cause considerable human-
wildlife conflict. For example, badgers are a known reservoir host of *Mycobacterium*
bovis, the causative agent of bovine tuberculosis, which poses a risk to livestock health
and is a serious economic problem for the UK cattle industry [26]. Issues such as this

highlight the challenge of conflict resolution for iconic British species, which can be

both controversial and associated with complex legislative considerations. It is

necessary, however, to consider all problematic species to address the UK's

commitments to improving biodiversity as part of government plans or global policy

frameworks.

Whilst policy frameworks are essential for providing goals and targets to reverse

biodiversity loss, they do not necessarily incorporate a process by which these targets,

such as managing human-wildlife conflict, can be identified in the first place, nor which

should be prioritised. Assessing conflicts between species is further complicated by

the fact that impacts are context dependent [27], which requires that the measure of

impact is standardised to quantify and compare differences between species,

taxonomic groups and localities [28]. When addressing the biodiversity impacts of

invasive non-native species, Nentwig et al. [15] suggested prioritising actions against

the alien species with a high impact potential of becoming established or spreading,

using a method to quantify impacts in a way that could be compared among alien

species within a taxonomic group. They devised a generic impact scoring system

(GISS) to compare relevant impact categories that had been reported in the scientific

literature, applying a scoring system to quantify impacts within the categories and rank

species according to their impact scores. The GISS has since been modified and

applied to a variety of classifications including alien fish [29], terrestrial arthropods [30],

aquatic invertebrates [31], plants [32, 33] and alien birds [34].

The main aim of this paper was to identify and categorise human-wildlife conflicts

associated with British mammals in order to summarise their magnitude and, where

possible, their costs. The research augmented a review of human-wildlife conflicts

produced over a decade ago (APHA, unpublished data) in response to a now

withdrawn Defra policy framework on Wildlife Management in England [35]. Only

terrestrial mammals (including bats) were assessed as they involve some of the most

controversial and sensitive issues as well as complex legislative considerations. To

determine whether the available evidence reflected the level of risk to human interests,

or the welfare or conservation status of an affected species, a modified GISS was

applied to all conflicts so that mammals could be ranked according to their level of

impact. Identifying species with the highest impact scores has the potential to assist

policy development by enabling prioritisation of the most damaging species. In

addition, the compilation of a comprehensive audit of British wildlife conflicts highlights

where further areas of research might be required and identifies the early emergence

of potential issues.

Materials and methods

Literature searches and screening

Searches were undertaken using Scopus (an abstract and citation database of peer-

reviewed scientific literature) for literature published up until and including 2024.

Additional searches were conducted using Google Scholar limited to the first 200

returns [36] to provide additional evidence where required. Separate searches were

carried out for each of the 66 terrestrial mammals incorporating a combination of

unique species terms and common keywords within the title, abstract and keywords

of the scientific literature. The following search terms were applied to each species:

[{common name} AND {Latin name}] AND [{England} OR {Scotland} OR {Wales} OR

{Great Britain} OR] AND [{conflict} OR {pest} OR {damage} OR {impact} OR

{management} OR {disease} OR {zoono*} OR {animal health} OR {health} OR

{conservation} OR {predat*}]. The minimum requirements for a literature resource to

be included in the analysis were the presence of an accessible abstract and that the

conflict occurred in Great Britain. Only articles written in English were included.

All articles returned during an individual search were subjected to a number of

screening processes. A rapid first screening of the title assessed documents for any

likely relevance to the species concerned and its potential conflicts. All potentially

relevant articles or ambiguous titles were screened further by examining the abstracts

in more detail or the full article, if necessary, resulting in a definitive list of conflicts for

each species. In many cases, single articles were applicable to multiple conflicts

and/or more than one species and therefore may have been added retrospectively to

a species reference catalogue.

173   Categorising and scoring conflicts

Conflicts were categorised depending on whether they were an economic, health,

environmental or social impact. Within these four main categories, conflicts were

further assigned to 32 subcategories (economic=15, health=6, environmental=7,

social=4). Subcategories were not weighted, therefore all subcategories were

considered as of equal importance. Some species were involved in multiple conflicts

within multiple categories, which were all captured as distinct entities. A

comprehensive report was compiled for all species identified as being involved in

human-wildlife conflict. An impact scale scoring system modified from Nentwig et al.

[15] was applied to estimate the magnitude of each conflict at the national level for

Great Britain, rated independently by three reviewers (KP, DC, GS). The level of

impact was quantified on a scale of 1 (minimal impact known) to 5 (the highest possible

impact: Table 1), with a value of confidence (low=1, medium=2 or high=3: Table 2)

also assigned to each conflict (taken from Hawkins et al. [37]). Mammals for which no

data were available, no impacts were known or impacts were not detectable were

excluded from the analysis. The median of the three independent assignments of

impact scale and confidence rating are reported for each conflict.

Table 1. Criteria used to describe the scale of impacts in each of the four
categories of conflict.

Scale of impact	A. Economic impact	B. Health impact	C. Environmental impact	D. Social impact
1	<£10K / yr	Local mild, short-term, reversible effects to individuals	Local, short-term, population loss, no significant ecosystem effect	Minimal social disruption
2	£10K – 100K / yr	Mild short-term, reversible effects to identifiable groups, localised	Some ecosystem impact, reversible changes, localised	Significant concern expressed at local level
3	£100K – 1M / yr	Minor irreversible effects and/or larger numbers covered by reversible effects, localised	Measurable long-term damage to populations and ecosystem, but little spread, no extinction	Temporary changes to normal activities at local level
4	£1M – 10M / yr	Significant irreversible effects locally or reversible effects over a large area	Long-term irreversible ecosystem change, spreading beyond local area	Some permanent change of activity locally, concern over a wider area
5	>£10M / yr	Widespread, severe, long-term, irreversible health effects	Widespread, long-term population loss or extinction, affecting several species with serious ecosystem effects	Long-term social change, significant loss of employment, migration from area

195   **Table 2. Criteria used for the confidence ratings of the impact scale**
196   **assignments.**

Confidence level	Example evidence
High (3) (approx. 90% chance of impact scale assignment being correct)	There is direct relevant observational evidence to support the assessment; **and** Impacts are recorded at the typical spatial scale over which the conflict can be characterised; **and** There are reliable / good quality data sources on the impacts of the conflict; **and** The interpretation of the data / information is straightforward; **and** Data / information is not controversial or contradictory
Medium (2) (approx. 65-75% chance of impact scale assignment being correct)	There is some direct observational evidence to support the assessment but some information is inferred; **and/or** Impacts are recorded at a spatial scale, which may not be relevant to the scale over which conflict can be characterized, but extrapolation or downscaling of the data to relevant scales is considered reliable, or to embrace little uncertainty; **and/or** The interpretation of the data is to some extent ambiguous or contradictory
Low (1) (approx. 35% chance of impact scale assignment being correct)	There is no direct observational evidence to support the assessment, e.g. only inferred data have been used as supporting evidence; **and/or** Impacts are recorded at a spatial scale, which is unlikely to be relevant to the scale over which conflict can be characterized; extrapolation or downscaling of the data to relevant scales is considered unreliable potentially leading to significant uncertainties; **and/or** Evidence is poor and difficult to interpret, e.g. because it is strongly ambiguous; **and/or** The information sources are considered to be of low quality or contain information that is unreliable

Various species characteristics were also included in the analyses to determine

whether they were associated with a conflict impact (modified from Nentwig et al. [15]).

Impact scores were considered as the maximum impact for a particular species in

Great Britain and did not account for their current distribution. Overall impacts were

compared between species by summing the median scores of all conflicts and within

each category ("potential impact"). To consider the distribution of a species, potential

impact values were multiplied by the proportion of hectads (10 kilometre square

resolution) each species occupies in Great Britain (derived from Crawley et al. [38]).

The potential impact scores, corrected for the area occupied can be considered as the

"actual impact" a species causes in Great Britain. Estimated population sizes were

derived mostly from Mathews et al. [39], typical body weights were taken mostly from

Harris & Yalden [40]. Where information was missing, relevant online sources were

used (see S2 Table). The overall biomass for each species was calculated by

multiplying population estimates and body mass. Population estimates and/or

distribution data could not be reliably established for three species (feral cat, feral

ferret, greater white-toothed shrew), therefore they were excluded from some of the

statistical analyses.

215   Statistical analysis

All statistical analyses were performed using the R statistical programme version 4.2.2

[41] and the RStudio Integrated Development Environment (IDE: version 2022.12.0,

Posit Software, PBC). Given the nature of the response variables, initial analyses (to

assess whether impact/confidence scores assigned by the three reviewers differed

overall and in relation to the conflict species assessed) were performed using an

ordinal logistic regression model fitted using the *ordinal::clm* function version 2023.12-

4. [42]. Species were aggregated based on status (native/non-native/feral) and

taxonomic order, with an interaction term between each aggregate type and reviewer

ID included. Additional analyses were performed to assess whether impact scores

assigned by the reviewers differed across impact categories (i.e. economic,

environmental, health and social), across subcategories, between native/non-native

species and between taxonomic orders, including interaction terms. (N.B. feral species

were removed from further analyses due to their low number of conflicts). Data were

expanded to account for the disparity in the number of subcategories described for

each species from the literature, so that for each species, a total of 32 subcategories

were present, with an impact and confidence score of zero (0), representing no

available data, assigned to absent subcategories.

Differences in potential and actual impact scores were analysed using generalised

linear models (GLM), using the *glmmTMB* package version 1.1.10 [43]. Diagnostic

testing of these analyses was performed using the *DHARMa* package version 0.4.7

[44]. Prior to analysis, data were modified to account for species whereby an actual

impact score could not be reliably established, with a score of zero (0) assigned to

indicate insufficient evidence of impact. Feral ferret was subsequently excluded from

the analyses, given a lack of data regarding population size and geographic

distribution. Analyses assessed whether estimates of total impact (actual or potential)

differed in relation to native/non-native status and taxonomic order, when modelled

independently. Several covariate predictors were also specified independently in each

model, including population size, body mass (kg) and biomass (kg). Models consisted

of GLMs, fitted with a gaussian error distribution and "identity" link function, selected

given that model residuals conformed to key assumptions (i.e. linearity, normality,

homogeneity of variance), when total impact estimates were logarithmically

transformed [log(n+1)]. For each model, a two-way interaction term was specified

between species aggregation type (either native/non-native status or taxonomic order)

and each covariate predictor (population size, body mass, biomass). Due to issues

with model performance, population size, body mass and biomass were also log-

transformed [log(n+1)].

Additional analyses examined how estimates of impact (actual or potential) might

differ across the main impact categories (economic, health, environment, social);

these were performed using a GLM fitted with a Tweedie distribution, which is

appropriate when analysing a continuous response variable consisting of a

combination of zeroes and positive values. These models compared impact across

categories in relation to native/non-native status and order, as well as population size,

body mass and biomass. Interaction terms involving impact category, species

aggregation type and each covariate predictor were included with respect to

native/non-native species only. Analyses of impact, in relation to impact categories

and order and covariate predictors, were limited to main order effects only, due to

persistent issues with model performance (i.e. failure of models to converge), when

considering any possible interaction between both predictor variables. This is likely

due to the complexity of the models (impact category=4 levels; taxonomic order=6

levels; total number of factorial combinations=24) when considering interaction terms

and the complete separation of certain impact categories across certain taxonomic

orders, caused by the presence of absolute zeroes (e.g. social impact and Rodentia).

Results

General literature searches

A total of 13,148 articles were returned for potential human-wildlife conflicts involving

the 66 terrestrial mammals in the UK. Further screening identified 405 useful articles

for 48 mammals belonging to seven taxonomic groups (Rodentia: n=15; Carnivora:

n=11; Artiodactyla: n=8; Chiroptera: n=6; Eulipotyphla: n=4; Lagomorpha: n=3;

Perissodactyla: n=1). The first available record was from 1947 and the latest was from

2024. The majority of articles (n=367, 91%) were published within the last three

decades (2014-2024: n=176, 44%; 2004-2013: n=110, 27%; 1994-2003: n=81, 20%).

A detailed description of all of the conflicts is in S1 File. A summary of

characteristics (native/non-native/feral status, population size, body mass, biomass)

and conflicts for each species is in S2 Table. The periods during which conflicts were

reported in the scientific literature (i.e. identified during the original review only ("pre-

2014"), during the original and current review ("pre-2014 and 2014 to present") and

during the current review only ("2014 to present") are also shown. The median impact

scores and confidence ratings are shown after each conflict in parentheses.

Overview of conflicts

Conflicts and status

Twenty-nine of the 48 mammals (61%) are native to the British Isles, 17 (35%) are

non-native (which includes 10 that have been introduced in recent times and seven

that are considered naturalised) and two (4%) are feral (species that are considered

wild but are either managed or may have human dependence). Of the 48 mammals,

40 (83%) caused at least one economic conflict, 36 (75%) caused at least one health

conflict, 23 (48%) caused at least one environmental conflict and eight (17%) caused

at least one social conflict (Fig 1). When compared by status, 86% (25/29) of native

mammals caused economic conflict, 66% (19/29) caused health conflict and 35%

(10/29) caused environmental conflict. In comparison, 82% (14/17) of non-native

mammals caused economic conflict, 88% (15/17) caused health conflict and 71%

(12/17) caused environmental conflict. Social conflicts accounted for approximately

17% for both native (5/29) and non-native (3/17) species. The proportion of feral

mammals causing conflict was either 50% (economic and environment) or 100%

(health), although this was attributable to only two species (feral cat, Dartmoor/Exmoor

pony).

Fig 1. Percentage of mammals within each status causing conflict within the

four main categories. Data labels denote the number of mammals within each

status.

Frequency of conflicts

In total, 200 separate conflicts were identified, with 103 attributable to native mammals

(52%), 92 to non-native mammals (46%) and five to feral mammals (3%). Ninety-one

of the 200 conflicts (45%) were economic impacts (Appendix 1a), 71 (35%) were

health impacts (Appendix 1b), 29 (15%) were environmental impacts (Appendix 1c)

and nine (5%) were social impacts (Appendix 1d). A similar proportion of conflicts were

recorded for both native and non-native species within all four categories (Fig 2). For

native mammals, economic impacts accounted for 47% (48/103), health impacts

accounted for 36% (37/103), environmental impacts accounted for 11% (12/103) and

social impacts accounted for 6% (6/103). For non-native mammals, economic impacts

accounted for 45% (42/92), health impacts accounted for 36% (32/92), environmental

impacts accounted for 16% (15/92) and social impacts accounted for 3% (3/92). Feral

mammals were involved in considerably less conflicts, mostly attributable to health

and environmental impacts.

Fig 2. The percentage of conflicts within each category depending on status. Data labels denote the number of conflicts within each category. Native mammals: n=103; non-native mammals: n=92; feral mammals: n=5.

The total number of mammals causing conflict within a particular subcategory ranged from one to 30 (Appendix 1a-1d). Two of the four most common conflicts were health related (reservoir of zoonotic disease, n=30; reservoir of disease, n=22) followed by two economic conflicts (damage to agricultural interests, n=19; damage to forestry, n=15), with a similar number of native and non-native mammals causing conflict in these subcategories. However, when these data were analysed separately according to status, non-native mammals accounted proportionately more than native mammals in all four subcategories (Fig 3).

Fig 3. The percentage of native and non-native mammals causing conflict within the four most common conflict subcategories. Data labels denote the number of mammals. Native: n=29; non-native: n=17.

Impact scores and ranking of conflicts

Reproducibility of the scoring

Of the 200 conflicts, the three reviewers agreed on the impact score in 92 cases (46%); there was a maximum deviation of one score in 89 cases (44%), a maximum deviation of two scores in 18 cases (9%) and a maximum deviation of three scores in 1 case (1%). Overall, there was no statistically significant difference between the independently assigned impact scores (χ^2(2)=3.057, p=0.217). However, confidence scores were found to differ statistically between reviewers (χ^2(2)=144.54, p<0.001); all three agreed in 33 of the 200 conflicts (17%), two agreed in 131 cases (65%), with a

341 deviation of one or two assigned by the third reviewer, and all three assigned a
342 different confidence score in 36 cases (18%). This suggests that the reviewers were
343 largely consistent in their impact scoring of British mammals but not necessarily in the
344 confidence of the assignments.

Conflict scoring and ranking

Using the median impact scores across the three reviewers, seven of the 200 conflicts
(4%) were assessed as impact score 5, 20 (10%) as impact score 4, 30 (15%) as
impact score 3, 88 (44%) as impact score 2 and 55 (27%) as impact score 1 (Table
4a-4d). Six of the seven conflicts assessed as having the highest impact score of 5
(all with confidence rating of high) were economic and one was a health conflict. These
were (in order of highest approximate economic cost): (1) rabbit damage to agricultural
interests (£120 million), (2) badger involvement in the transmission of bTB to cattle
(>£100 million), (3) brown rat damage to agricultural interests (£54 million), (4) rabbit
damage to forestry (£43 million) (5) badger damage to agricultural interests (£39
million) (6) grey squirrel damage to forestry (£28 million, not including carbon capture
estimates) and (7) brown rat damage to property (£14 million, based on local authority
costs only). In all four categories, the majority of conflicts were scored as impact 2
(economic: 35% (32/91); health: 46% (33/71); environmental: 55% (16/29); social:
78% (7/9)). Despite two health conflicts being the most common amongst all
mammals, generally, health conflicts were scored low, with over 90% (64/71) scoring
either a 1 or 2.

Table 4a. Total number of mammals causing conflict within the 15 economic subcategories and their median impact score.
Status categories: Na=native; NN=non-native; Fe=feral.

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Damage to forestry (15)	Grey squirrel, rabbit	Roe deer, red deer, fallow deer	Japanese sika, brown hare, edible dormouse, field vole	Muntjac, feral goat, mountain hare, beaver, bank vole	Chinese water deer	**Na - 6** **NN - 9**
Damage to agricultural interests (19)	Badger, rabbit, brown rat	Roe deer, fallow deer, mole, house mouse	Red deer	Japanese sika, Chinese water deer, muntjac, wild boar, brown hare, mountain hare, grey squirrel, beaver, wood mouse	Edible dormouse, field vole	**Na - 9** **NN - 10**
Damage to vehicles (RTAs) (9)	-	Roe deer, fallow deer, badger	Red deer, muntjac	Japanese sika, Chinese water deer, wild boar, Exmoor/Dartmoor pony	-	**Na - 4** **NN - 4** **Fe - 1**
Competition with livestock (3)	-	-	-	-	Red deer, Japanese sika, feral goat	**Na - 1** **NN - 2**
Predation of game species (11)	-	Red fox	-	Stoat, badger, American mink, hedgehog, brown rat	Wild cat, pine marten, feral ferret, weasel, polecat	**Na - 8** **NN - 3**
Predation/impact on commercial fisheries (3)	-	-	-	Otter, American mink	Beaver	**Na - 2** **NN - 1**
Predation of poultry (4)	-	-	-	American mink	Pine marten, feral ferret, polecat	**Na - 2** **NN - 2**
Damage to property (6)	Brown rat	Badger, grey squirrel, house mouse	Edible dormouse	Natterer's bat	-	**Na - 2** **NN - 4**
Damage to amenities and domestic grassland (4)	-	Rabbit	American mink, mole	Badger	-	**Na - 2** **NN=2**
Conflicts with construction, infrastructure and development (9)	-	Badger	Natterer's, serotine, common pipistrelle, soprano pipistrelle, brown long-eared bat	Beaver, hazel dormouse, water vole	-	**Na - 9**
Predation of livestock (1)	-	Red fox	-	-	-	**Na - 1**
Conflicts with transport (2)	-	-	Rabbit, brown rat	-	-	**NN - 2**
Sustaining predators harmful to game (2)	-	-	-	-	Field vole, Orkney vole	**Na - 1** **NN - 1**

				Brown hare	Mountain hare	**Na - 1**
Damage to property as a result of poaching (2)				Brown hare	Mountain hare	**Na - 1** **NN - 1**
Conflicts with utilities (1)	-	Brown rat	-	-	-	**NN - 1**
Total per impact score	**6**	**18**	**17**	**32**	**18**	**91**

366 **Table 4b. Total number of mammals causing conflict within the six health subcategories and their median impact score. Status**
367 **categories: Na=native; NN=non-native; F=feral.**

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Health risks from RTAs (9)	-	-	Roe deer, wild boar	Red deer, Japanese sika, fallow deer, muntjac, badger	Chinese water deer, Exmoor/Dartmoor pony	**Na - 4** **NN - 4** **Fe - 1**
Reservoir of disease (22)	-	-	Rabbit, grey squirrel	Roe deer, red deer, fallow deer, muntjac, wild boar, feral cat, badger, red fox, field vole, wood mouse, house mouse	Japanese sika, Chinese water deer, feral goat, stoat, Daubenton's, hedgehog, brown hare, red squirrel, beaver	**Na - 12** **NN - 9** **Fe - 1**
bTB transmission (8)	Badger	-	-	Fallow deer, muntjac, wild boar	Roe deer, red deer, Japanese sika, red fox	**Na - 5** **NN - 3**
Reservoir of zoonotic disease (30)	-	Brown rat	House mouse	Roe deer, red deer, fallow deer, muntjac, American mink, red fox, Daubenton's, serotine, hedgehog, rabbit, bank vole, field vole, wood mouse	Japanese sika, Chinese water deer, feral goat, polecat, common shrew, greater white-toothed shrew, brown hare, mountain hare, red squirrel, grey squirrel, beaver, edible dormouse, yellow-necked mouse, Skomer vole, black rat	**Na - 15** **NN - 15**
Threat to livestock where ground is undermined (1)	-	-	-	Mole	-	**Na - 1**
Threats of branches falling (1)	-	-	-	-	Grey squirrel	**NN - 1**
Total per impact score	**1**	**1**	**5**	**33**	**31**	**71**

370 **Table 4c. Total number of mammals causing conflict within the seven environmental subcategories and their median impact**
371 **score. Status categories: Na=native; NN=non-native; F=feral.**

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Damage to conservation habitats (9)	-	-	Roe dee, red deer	Japanese sika, fallow deer, muntjac, feral goat	Chinese water deer, wild boar, bank vole	**Na - 4** **NN - 5**
Threat to biodiversity (6)	-	Grey squirrel	-	Roe deer, red deer, Japanese sika, fallow deer, greater white-toothed shrew	-	**Na - 2** **NN - 4**
Threat to genetic integrity (3)	-	-	Feral cat	Japanese sika, feral ferret	-	**NN - 2** **Fe - 1**
Competition with or predation of wildlife (including protected or conservation species) (7)	-	-	Badger, American mink, red fox	Feral cat, otter, pine marten, stoat	-	**Na - 5** **NN - 1** **Fe - 1**
Predation of important ground-nesting birds and island ecosystem damage (2)	-	-	Hedgehog, brown rat	-	-	**Na - 1** **NN - 1**
Sustaining predators harmful to important conservation species (1)				Rabbit	-	**NN - 1**
Damage to island ecosystems (1)	-	-	-	-	Black rat	**NN - 1**
Total per impact score	**0**	**1**	**8**	**16**	**4**	**29**

373 **Table 4d. Total number of mammals causing conflict within the four social subcategories and their median impact score. Status**
374 **categories: Na=native; NN=non-native; F=feral.**

Conflict (total mammals)	Mammal impact scores within each subcategory					Total per status
	Impact 5	Impact 4	Impact 3	Impact 2	Impact 1	
Public nuisance (4)	-	-	-	Muntjac, feral goat, rabbit	Wild boar	**Na - 1** **NN - 3**
Domestic nuisance (2)	-	-	-	Pine marten	Red fox	**Na - 2**
Predation of domestic pets (1)	-	-	-	Red fox	-	**Na - 1**
Domestic nuisance & hygiene (2)	-	-	-	Natterer's, serotine	-	**Na - 2**
Total per impact score	**0**	**0**	**0**	**7**	**2**	**9**

Analysis of impact scores in relation to native/non-native status and conflict

category, found a statistically significant effect of category, when considered as a main

order effect ($\chi^2(3)$=62.90, p<0.001) but not status ($\chi^2(1)$=0.421, p=0.516). Reviewers

assigned a greater score in relation to economic impacts, followed by environmental,

health and then social impacts (Fig 4). When conflict category was considered in

relation to status, there was no statistically significant interaction term between the two

($\chi^2(3)$=4.839, p=0.184)

Fig 4. The proportion of conflicts depending on impact score (1-5) within the

four categories. Data labels denote to total number of conflicts within each

impact score. Economic: n=91, health: n=71, health: n=29, social: n=9.

Analysis of impact scores, in relation to native/non-native status and conflict

subcategories, showed both as having a statistically significant effect ($\chi^2(1)$=33.63,

p<0.001; $\chi^2(31)$=640.18, p<0.001, respectively), with a significant two-way interaction

term also identified ($\chi^2(31)$=155.56, p<0.001). Across all economic impacts, non-

native species received a statistically significantly greater impact score than natives in

relation to damage to woodland, damage to agricultural interests, damage to property,

conflict with transport and conflict with utilities (all p<0.05). Conversely, native species

received a significantly higher score due to conflicts with construction and

development and the predation of livestock (both p<0.001). Across health impacts, the

model predicted a statistically greater impact score for non-native species as

reservoirs of zoonotic diseases only (p=0.008). With regards to the environment, non-

native species received a statistically significant greater impact score for damage to

conservation habitats and their threat to biodiversity and genetic integrity (all p<0.05).

In terms of social impacts, non-native species received a greater impact than natives

with respect to public nuisance only ($p<0.01$).

Potential impact

The number of separate conflicts per species ranged from one to eleven (Table 5),

with sika deer involved in the most conflicts. Species scoring highest in potential total

impact (i.e. the sum of all median conflict scores for a particular species) were badger,

rabbit, roe deer, fallow deer and red deer. Generally, these species also scored highly

within the separate categories (except for social conflicts), with badgers scoring

highest in both potential economic and health impacts. The highest potential

environmental impact was produced by sika deer, followed by roe deer, red deer and

feral cat. The highest potential social impact was observed in the red fox, followed by

rabbit, wild boar, pine marten, Natterer's and serotine bat.

Table 5. Distribution (i.e. the proportion of Great Britain each mammal is found, based on presence in 10x10km squares),
total number of conflicts and impact scores (potential and actual) in total and within each of the four categories (economic,
health, environmental, social) for all conflict mammals. Mammals are listed in order of highest total actual impact. – denotes
missing data.

Species	Distribution	No. conflicts	Potential impact					Actual impact				
			Total potential impact	Economic	Health	Environment	Social	Total actual impact	Economic	Health	Environment	Social
Badger	0.746	10	33	21	9	3	0	24.624	15.670	6.716	2.239	0
Rabbit	0.833	8	26	17	5	2	2	21.649	14.155	4.163	1.665	1.665
Roe deer	0.726	9	25	12	8	5	0	18.151	8.713	5.808	3.630	0
Red fox	0.800	8	19	8	5	3	3	15.207	6.403	4.002	2.401	2.401
Grey squirrel	0.583	7	20	11	5	4	0	11.653	6.409	2.913	2.331	0
Brown rat	0.529	7	21	14	4	3	0	11.105	7.403	2.115	1.586	0
Red deer	0.412	10	23	11	7	5	0	9.480	4.534	2.885	2.061	0
Fallow deer	0.347	9	24	12	8	4	0	8.317	4.158	2.772	1.386	0
American mink	0.533	6	14	9	2	3	0	7.457	4.794	1.065	1.598	0
Mole	0.807	3	9	7	2	0	0	7.266	5.651	1.615	0	0
Hedgehog	0.824	4	8	2	3	3	0	6.592	1.648	2.472	2.472	0
Brown hare	0.680	5	9	7	2	0	0	6.119	4.759	1.360	0	0
Muntjac	0.334	9	17	7	8	2	0	5.673	2.336	2.669	0.667	0
Field vole	0.564	5	9	5	4	0	0	5.075	2.819	2.256	0	0
Wood mouse	0.560	3	6	2	4	0	0	3.360	1.120	2.240	0	0
Otter	0.782	2	4	2	0	2	0	3.126	1.563	0	1.563	0
House mouse	0.235	4	13	8	5	0	0	3.060	1.883	1.177	0	0
Stoat	0.597	3	5	2	1	2	0	2.983	1.193	0.597	1.193	0
Natterer's	0.411	3	7	5	0	0	2	2.880	2.057	0	0	0.823
Sika deer	0.134	11	19	8	5	6	0	2.540	1.069	0.668	0.802	0
Bank vole	0.446	3	5	2	2	1	0	2.229	0.892	0.892	0.446	0
Common pipistrelle	0.674	1	3	3	0	0	0	2.021	2.021	0	0	0
Soprano pipistrelle	0.619	1	3	3	0	0	0	1.858	1.858	0	0	0
Brown long-eared bat	0.600	1	3	3	0	0	0	1.799	1.799	0	0	0
Serotine	0.246	3	7	3	2	0	2	1.721	0.738	0.492	0	0.492
Daubenton's	0.491	2	3	0	3	0	0	1.472	0	1.472	0	0
Pine marten	0.171	4	6	2	0	2	2	1.025	0.342	0	0.342	0.342

417 **Table 5 cont.**

Species	Distribution	No. conflicts	Total potential impact	Economic	Health	Environment	Social	Total actual impact	Economic	Health	Environment	Social
				Potential impact				Actual impact				
Mountain hare	0.162	4	6	5	1	0	0	0.973	0.811	0.162	0	0
Water vole	0.425	1	2	2	0	0	0	0.850	0.850	0	0	0
Polecat	0.269	3	3	2	1	0	0	0.807	0.538	0.269	0	0
Weasel	0.530	1	1	1	0	0	0	0.530	0.530	0	0	0
Common shrew	0.502	1	1	0	1	0	0	0.502	0	0.502	0	0
Red squirrel	0.247	2	2	0	2	0	0	0.494	0	0.494	0	0
Feral goat	0.023	6	13	4	7	1	1	0.209	0.070	0.047	0.047	0.047
Hazel dormouse	0.199	1	2	2	0	0	0	0.397	0.397	0	0	0
Chinese water deer	0.042	7	9	5	3	1	0	0.375	0.208	0.125	0.042	0
Wild boar	0.031	7	9	3	2	2	2	0.402	0.124	0.216	0.03	0.031
Beaver	0.018	6	9	7	2	0	0	0.166	0.129	0.037	0	0
Yellow-necked mouse	0.129	1	1	0	1	0	0	0.129	0	0.129	0	0
Edible dormouse	0.008	4	8	7	1	0	0	0.064	0.056	0.008	0	0
Wildcat	0.039	1	1	1	0	0	0	0.039	0.039	0	0	0
Exmoor/Dartmoor pony	0.005	2	3	2	1	0	0	0.016	0.011	0.005	0	0
Black rat	0.006	2	2	0	1	1	0	0.011	0	0.006	0.006	0
Skomer vole	0.004	1	1	0	1	0	0	0.004	0	0.004	0	0
Orkney vole	0.003	1	1	1	0	0	0	0.003	0.003	0	0	0
Feral cat	-	3	7	0	2	5	0	-	-	-	-	-
Feral ferret	-	3	4	2	0	2	0	-	-	-	-	-
Greater white-toothed shrew	-	2	3	0	1	2	0	-	-	-	-	-

Analyses identified a statistically significant difference between native/non-native

species (χ^2(1)=4.235, p=0.040), with the latter predicted as having a higher potential

impact overall (β=0.513, z ratio=2.058, p=0.046). When specified as main order

effects, total potential impact was found to increase as a function of body mass

(β=0.335, t ratio=3.598, $p<0.001$) and biomass (β=0.184, t ratio=4.706, $p<0.001$) but

not population size (β=0.040, t ratio=1.012, p=0.311). However, no statistically

significant interaction term was identified ($p>0.05$), suggesting that differences in the

rate of change for total potential impact were comparable between native and non-

native species, relative to changes in population size, body mass and biomass.

There was a statistically significant difference between taxonomic orders

(χ^2(5)=18.468, p=0.002), with Artiodactyla having a higher potential impact compared

to Chiroptera (β=1.129, t ratio=3.661, p=0.009) and Rodentia (β=1.231, t ratio=3.236,

p=0.028), whilst all other taxonomic comparisons were non-significant (Fig 5).

Inclusion of covariate predictors as main order effects, revealed a statistically

significant effect of population size (χ^2(1)=7.259, p=0.007) and biomass (χ^2(1)=39.467,

$p<0.001$), with total potential impact predicted to increase as a function of each

(population size: β=0.096, t ratio=2.499, $p<0.001$; biomass: β=0.199, t ratio=5.641,

$p<0.001$).

Figure 5. Average potential impacts within the four categories across orders

(excluding Perissodactyla).

Actual impact

Species scoring the highest actual impact were badger, rabbit, roe deer and red fox

(Table 5). Generally, these species also scored highly within the separate categories,

with badgers scoring highest in both actual economic and health impacts. Roe deer

scored the highest actual environmental impact, followed by hedgehog, red fox, grey

squirrel, badger and red deer. The highest potential social impact was observed in the

red fox, followed by rabbit and Natterer's bat.

Analysis of actual impact estimates showed no statistically significant difference

between native/non-native species ($\chi^2(1)$=0.056, p=0.813). When considered as main

order effects, total actual impact was predicted to increase significantly in relation to

population size (β=0.094 t ratio=1.980, p=0.048) and biomass (β=0.215, t ratio=4.691,

p<0.001). When considered in relation to status, there was no statistically significant

interaction for biomass ($\chi^2(1)$=0.298, p=0.585) suggesting that the rate of increase

was comparable across native and non-native species. Body mass was found to have

no statistically significant effect on total actual impact when considered as a main order

effect ($\chi^2(1)$=1.741, p=0.187) or through interaction with status ($\chi^2(1)$=0.088, p=0.767).

A statistically significant interaction term was identified between status and population

size, with total actual impact predicted to increase, relative to population size, at a

significantly higher rate for non-native species when compared to natives (β=0.179, t

ratio=2.120, p=0.034).

There was no statistical difference, in relation to actual impact scores, between

the taxonomic orders ($\chi^2(5)$=5.622, p=0.345: Fig 6).

Figure 6. Average actual impacts within the four categories across orders

(excluding Perissodactyla).

Conflict change over time

For many mammals, a conflict occurring before 2014 was still being reported in the

scientific literature within the last decade (S2 Table). Twenty-seven of the 46 (59%)

conflicts reported only before 2014 were economic, followed by 10 (22%) health, 5 environment (11%) and 4 social (8%). In contrast, 25 of the 45 (55%) conflicts reported from 2014 to 2024 were health (55%), mostly attributable to reservoir of disease or reservoir of zoonotic disease, followed by 9 environment (20%), 8 economic (18%) and 3 social (7%).

Discussion

Negative interactions between people and wildlife are more often than not viewed as human–wildlife conflict, with socio-economic factors cited amongst the issues influencing people's perception of wildlife [45]. In this paper, we estimated a conservative total annual economic cost of human-wildlife conflicts in Great Britain (as reported in S1 File) in excess of £0.5 billion, which would likely exceed £1 billion if health and environmental conflicts could be monetarised. All but one of the seven conflicts assessed as having the highest impact score of 5 involved economic conflicts. The one health conflict scored as impact 5 (the involvement of badgers in the transmission of bTB), was also measurable on an economic scale, suggesting that conflicts associated with a monetary value are inherently easier to quantify compared with those that are not. To account for potential bias associated with individual conflicts, we applied a prioritisation approach (generic impact scoring system (GISS)) to evaluate the overall impact of conflict mammals. In some cases, mammals causing the greatest economic damage, such as rabbit and badger, were ranked highest overall for their actual impact. However, this approach also identified species that did not necessarily cause the highest individual impacts; for example, roe deer and red fox were ranked third and fourth overall, respectively, for actual impact despite neither receiving the highest impact score of 5 for any of their identified conflicts.

The emphasis of legislation associated with wildlife in the UK has moved from

protecting human economic interests (e.g. Prevention of Damage by Pests Act 1949

and Pests Act 1954), to promoting animal welfare (e.g. Wildlife and Countryside Act

1981 and Animal Welfare Act 2006) and protecting biodiversity (e.g. Habitats Directive

1992, Environment Act 2021). There is likely to be an increasing trend for conflicts

arising between promoting biodiversity and human economic interests, as human

activities continue to expand into existing wildlife habitats. Four of the five mammals

identified as causing the greatest actual economic and health impact (badger, rabbit,

roe deer), environmental impact (roe deer, hedgehog, red fox), and social impact (red

fox, rabbit) are native, which leads to increasing conflict where specific legislation

applies to a protected species. For example, badgers are protected under the

Protection of Badgers Act 1992 but are the focus of ongoing controversy regarding

management options for controlling the spread of *Mycobacterium bovis* to cattle in the

UK, which is estimated to cost the taxpayer over £100 million annually [46]. Likewise,

some bat species are widely distributed throughout Great Britain and often associated

with conflicts with buildings or construction; as bats and their roosts are legally

protected under the Wildlife and Countryside Act 1981 and the Conservation of

Habitats and Species Regulations 2010, this can lead to potentially complex and costly

resolution.

Although native mammals were ranked amongst the highest scoring impacts in all

four categories, overall, there was no statistical difference in the impact scores

between native and non-native species. However, there was a disproportionate

amount of conflict in specific areas, with non-native species perceived as causing

significantly more damage to woodlands, conservation habitats and agricultural

interests, and a greater reservoir of zoonotic disease compared to native species.

Rabbits caused the highest economic conflict compared to all other mammals with

their impact on agriculture, and ranked second overall for total actual impact. As such,

this may highlight rabbits as a priority species for human-wildlife resolution. However,

rabbit populations have declined significantly in the UK in recent years, with severe

local population reductions reported [47], which have led to concerns regarding the

impact of such population reductions on other species, such as carnivores, that rely

on them as a food source [48]. In contrast, grey squirrel abundance and distribution

has expanded rapidly since they were introduced to Great Britain in the late 19[th]

century [49]. They are considered a pest species in the UK and as such are afforded

no protection under the Wildlife and Countryside Act 1981. As part of its 25-year

Environment plan [20], the government has committed to managing grey squirrels to

protect woodlands and red squirrel populations in the UK. While the general

consensus is that grey squirrels must be managed to protect biodiversity within the

UK, removing healthy wildlife can be controversial and may face opposition from

pressure groups, which can be key to the success or failure of an eradication project

[50].

Although the majority of reported conflicts were economic, the two most common

conflicts were health related (reservoir of zoonotic disease and reservoir of disease).

More than 60% of human pathogens are zoonotic in origin [51] with urbanisation and

anthropogenic factors, amongst others, thought to have greatly influenced the

emergence and distribution of zoonoses [52]. Generally, though, the impact score of

health conflicts were deemed low (mostly 1 or 2), therefore we assumed that most

health impacts were likely local, mild, short- to mid-term and with reversible effects

(see Table 1). This highlights the importance of surveillance, which is crucial to

preventing and controlling the spread of disease, in particular zoonotic disease [52].

This is also pertinent to exotic diseases, which are not reported here as only current

diseases causing conflict in Britian were assessed. Whilst there is extensive research

on the pathology and epidemiology of disease in a wide range of British mammals, as

reported in S1 File, it is evident that for some mammals, there is an inference of

disease presence and/or zoonotic potential or a general lack of information (e.g. some

hantaviruses [53] and Ljungan virus [54] in rodents; *Babesia* spp. in deer [55] and

badgers [56]), which suggests further research may be warranted. Understanding

disease dynamics in British wildlife is intrinsically linked to biodiversity conservation,

as anthropogenic activities such as urbanisation and habitat destruction, are likely to

lead to increased interaction between humans and wildlife [57].

In many cases, mammals that caused the most conflict were amongst the most

widely distributed throughout the British Isles, e.g. badger, rabbit, roe deer and red

fox. Our analyses identified that total actual impact tended to increase as a function of

biomass and population size, therefore a GISS approach may be particularly beneficial

for monitoring recently introduced species, whether native or non-native, which have

the potential to expand beyond their current range. Beavers were reintroduced to

Great Britain around 20 years ago and are considered keystone species due to their

ability to alter ecosystem structure and function through their engineering activities [58,

59]. Whilst beavers are generally considered as having a positive impact on

biodiversity, we identified six potential conflicts of interest with humans; whilst they

ranked 38th overall based on total actual impact, likely owing to their current limited

distribution, they ranked joint 12th overall based on potential impact. Whilst the

reintroduction of beavers across Great Britain is currently carefully managed,

highlighting and monitoring potential issues at this stage is essential for ensuring the

status of this species remains positive rather than possible future perception as a pest
species.

The overall assessments of the conflicts reported here were made at the national
level for Great Britain. Local impacts may be of considerable importance for those
affected, for instance those injured in road traffic accidents, and thus a national score
of 1 or 2 should not be taken to indicate that the conflict is not important or worth
mitigating at a local level. Furthermore, it is important to recognise that the impacts
are not necessarily uniformly distributed geographically or equitably shared. For
example, Artiodactyla were predicted to receive significantly higher potential impact
scores compared to other taxonomic orders. The total potential impact for five of the
six deer species were amongst the highest of all the mammals, although the actual
impact for some deer species may be lower depending on their distribution. For
example, sika deer were involved in the most conflicts of all the mammals but were
ranked 20th overall for actual impact, suggesting its impacts are localised given its
limited distribution.

Human-wildlife conflict has long been associated with negative impacts on human
interests, although it is evident that interactions between people and wildlife are more
varied and complex [2]. More recently, the emphasis has shifted from conflict between
humans and wildlife and its mitigation, to a coexistence approach [14], or at least one
that constrains conflict or adapts shared spaces to mitigate conflict. Whilst there is a
general consensus that some species must be managed or even eradicated (e.g.
invasive non-native species: Nentwig et al. [15]), for others, particularly native species,
a more holistic approach may be required. In some cases, mammals that represent a
conflict and cost to some, may also provide a benefit to others. In Great Britain, the
red fox has been widely persecuted due to its perceived status as a pest species [60].

Their impact on agriculture has been estimated to cost £20 million annually, most of

which is attributable to direct and indirect effects of livestock predation [61]. However,

fox predation on rabbits, which have a much more significant impact on agriculture, is

estimated to create an indirect economic benefit to farmers of at least £11 million

annually and may even be approaching an economically neutral impact on agriculture

[61].

The findings reported here must be viewed in light of several limitations. Although

an extensive literature search was conducted to identify conflicts associated with

British wildlife, it is likely that some conflicts were not identified or cost estimates were

not available, and therefore not reported. Whilst the conflict descriptions given in S1

File are not exhaustive and based only on the searches returned, they likely represent

reasonable overviews and are sufficiently robust to provide an overall audit of the

current national importance of conflicts between human and wildlife interests in Great

Britain. However, individual assignments of scale may need to be revised as further

information emerges. Conflicts that were identified before 2014 only are more than

likely still occurring or may not be reported as frequently in the scientific literature.

Likewise, conflicts reported after 2014 only may have been occurring prior to this and

again, either missed in the original literature searches or less frequently reported. The

majority of "new" conflicts we identified were disease related, which may reflect a shift

of focus leading to increased reporting in this area, possibly in light of several

international public health emergencies of zoonotic origin within the last ten years,

such as Middle East respiratory syndrome (MERS), Ebola and Covid-19 pandemics

[62].

Another limitation of our findings is the lack of data for some species. Feral species

were excluded from some of the analyses due to the small number of conflicts in Great

Britain, although their impacts can be significant. Data regarding feral cats are limited

due to the difficulty in differentiating between feral and domestic cats [63], the

uncertainties of feral cat population estimates (based mostly on urban populations)

and the lack of known distribution within Great Britain. Cats are one of the most

ubiquitous mammals globally (see McDonald & Skillings [64]) and one of the most

destructive, with free-ranging cats suggested as having a major impact on the mortality

of several bird species in the UK [65]. They are also the biggest threat to the survival

of Scottish wildcats due to introgression [66]. Given their domesticated origins, feral

cats have no native range and their ubiquitous adverse impacts on native wildlife

around the world has led to them being labelled amongst the world's worst invasive

species [67]. From a biodiversity conservation perspective, a wealth of evidence

suggests that feral cats should be controlled and ideally removed from places where

their impact is detrimental, often as an urgent priority [68].

When applying the GISS in this paper, none of the conflicts were weighted, i.e.

they were all considered of equal importance. This follows Nentwig et al. [15], who

stated that in the absence of objective scientific criteria to consider one impact

category more important than the other, selective weights were not applied. They also

stated that deciding relative impact weights is a social process rather than a scientific

one. It is possible to apply individual weights to conflicts, for example, by considering

stakeholder values (e.g. Kumschick et al. [69]), which will likely differ according to the

relevance of a particular conflict to a particular group. The scoring system applied here

is thus considered robust to the uncertainty of impact knowledge but could be adjusted

with weight scores to account for specific social values, such as stakeholder interests,

in the future [15].

Conclusions

The prioritisation approach we applied here is a useful tool for identifying individual conflicts for individual species but also provides a broader assessment of where human-wildlife impacts may be affecting socio-economic and biodiversity interests. It is clear that it should not be relied upon in isolation to identify priority species for mitigation purposes, and all other available evidence or opinion from conservation groups, stakeholder groups, public perception and other interested parties, should be considered. Ideally, any decision making, whether relevant to government policies or conservation organisations, should consider ensuring a balance between the need to conserve and enhance biodiversity, the interests of the general public and the interests of those directly affected [35].

Acknowledgements

We are grateful to Eloise Cunningham for her assistance with the literature searches.

Author contributions

Conceptualisation and methodology: DC, KLP. Analysed the data: KP, DW, DC, GC. Writing-original draft: KP. Writing-review & editing: KP, DC, DW, GC.

References

1.   Locke H, Ellis EC, Venter O, Schuster R, Ma K, Shen X, et al. Three global conditions for biodiversity conservation and sustainable use: an implementation framework. National Science Review. 2019;6:1080-82.

2.   Hill CM. Conflict is integral to human-wildlife coexistence. Frontiers in

Conservation Science. 2021;2:734314.

3.   Woodroffe R, Thirgood S, Rabinowitz A. People and wildlife: conflict or

coexistence? Cambridge, UK: Cambridge University Press; 2005.

4.   Conover MR. Resolving human-wildlife conflicts: the science of wildlife damage

management: CRC Press, Boca Raton, FL; 2002.

5.   Treves A, Karanth KU. Human-carnivore conflict and perspectives on carnivore

management worldwide. Conservation Biology. 2003;17:1491–99.

6.   Tilman D. Global environmental impacts of agricultural expansion: the need for

sustainable and efficient practices. The Proceedings of the National Academy of

Sciences. 1999;96:5995–6000.

7.   Nyhus PJ. Human-wildlife conflict and co-existence. Annual Review of

Environment and Resources. 2016;41:143-71.

8.   Pérez E, Pacheco LF. Damage by large mammals to subsistence crops within

a protected area in a montane forest of Bolivia. Crop Protection. 2006;25:933-39.

9.   Fang L, Hong Y, Zhou Z, Chen W. The frequency and severity of crop damage

by wildlife in rural Beijing, China. Forest Policy and Economics. 2021;124:102379.

10.   Drexler JF, Corman VM, Drosten C. Ecology, evolution and classification of bat

coronaviruses in the aftermath of SARS. Antiviral Research. 2014;101:45-56.

11.   Song Z, Xu Y, Bao L, Zhang L, Yu P, Qu Y, et al. From SARS to MERS,

thrusting coronaviruses into the spotlight. Viruses. 2019;11.

12.   Hao YJ, Wang YL, Wang MY, Zhou L, Shi JY, Cao JM, et al. The origins of

COVID-19 pandemic: A brief overview. Transboundary and Emerging Diseases.

2022;69(6):3181-197.

13.   Cassini MH. Human-wildlife conflicts: does origin matter? . Animals. 2022;12:1-

10.

14.   Gross EM, Jayasinghe N, Dahal S, Tenzin S, Klenzendorf S, Vannelli K, et al.

C2C—conflict to coexistence: a global approach to manage human–wildlife conflict for

coexistence. Conservation Science and Practice. 2024;7:e13292.

15.   Nentwig W, Künel E, Bacher S. A generic impact-scoring system applied to

alien mammals in Europe. Conservation Biology. 2010;24:302–11.

16.   Eschen R, Kadzamira M, Stutz S, Ogunmodede A, Djeddour D, Shaw R, et al.

An updated assessment of the direct costs of invasive non native species to the United

Kingdom. Biological Invasions. 2023;25:3265-76.

17.   The Wildlife Trust. Local to global: the Global Biodiversity Framework and what

the   UK   needs   to   do   to   implement   it.   2024;

https://www.wildlifetrusts.org/sites/default/files/2024-

09/UK%20and%20Global%20Biodiversity%20Framework%20-

%20The%20Wildlife%20Trusts%202024.pdf.

18.   Dueñas MA, Hemming DJ, Roberts A, Diaz-Soltero H. The threat of invasive

species to IUCN-listed critically endangered species: a systematic review. Global

Ecology and Conservation. 2021;26:e01476.

19.   NNSS.   GB   non-native   species   report   card   2020.   2020;

https://www.nonnativespecies.org/assets/NNSIP-Scorecard-2020.pdf.

20.   HMGovernment. A green future: our 25 year plan to improve the environment.

2018;

https://assets.publishing.service.gov.uk/media/65fd713d65ca2f00117da89e/CD1.H_

HM_Government_A_Green_Future_Our_25_Year_Plan_to_Improve_the_Environme

nt.pdf; 2018.

21. Crowley SL, Hinchliffe S, McDonald RA. Conflict in invasive species

management. Frontiers in Ecology and the Environment. 2017;15:133-41.

22. Jackson DB, Green RE. The importance of the introduced hedgehog

(*Erinaceus europaeus*) as a predator of the eggs of waders (Charadrii) on machair in

South Uist, Scotland. Biological Conservation. 2000;93:333–48.

23. Manchester SJ, Bullock JM. The impacts of non-native species on UK

biodiversity and the effectiveness of control. Journal of Applied Ecology. 2000;37:845-

64.

24. Ricciardi A, Cohen J. The invasiveness of an introduced species does not

predict its impact. Biological Invasions. 2007;9:309–15.

25. Ricciardi A, Hoopes MF, Marchetti MP, Lockwood JL. Progress toward

understanding the ecological impacts of nonnative species. Ecological Monographs,.

2013;83:263–82.

26. Swift BMC, Barron ES, Christley R, Corbetta D, Grau-Roma L, Jewell C, et al.

Tuberculosis in badgers where the bovine tuberculosis epidemic is expanding in cattle

in England. Scientific Reports. 2021;11:20995.

27. Kumschick S, Gaertner M, Vilà M, Essl F, Jeschke JM, Pyšek P, et al.

Ecological impacts of alien species: quantification, scope, caveats, and

recommendations. Bioscience. 2015;65:55–63.

28. Nentwig W, Bacher S, Pyšek P. The generic impact scoring system (GISS): a

standardized tool to quantify the impacts of alien species. Environmental Monitoring

and Assessment. 2016;188:315.

29. Van der Veer G, Nentwig W. Environmental and economic impact assessment

of alien and invasive fish species in Europe using the generic impact scoring system.

Ecology Freshwater Fish. 2014(4):646-56.

30.   Vaes-Petignat S, Nentwig W. Environmental and economic impact of alien

terrestrial arthropods in Europe. NeoBiota. 2014;22:23–42.

31.   Laverty C, Nentwig W, Dick JTA, Lucy FE. Alien aquatics in Europe: assessing

the relative environmental and socio-economic impacts of invasive aquatic

macroinvertebrates and other taxa. Management of Biological Invasions. 2015;6:341–

50.

32.   Novoa A, Kumschick S, Richardson DM, Rouget M, Wilson JRU. Native range

size and growth form in Cactaceae predicts invasiveness and impact. NeoBiota.

2016;30:75-90.

33.   Sohrabi S, Pergl J, Pyšek P, Foxcroft LC, Gherekhloo J. Quantifying the

potential impact of alien plants of Iran using the Generic Impact Scoring System

(GISS) and Environmental Impact Classification for Alien Taxa (EICAT). Biological

Invasions. 2021;23:2435-449.

34.   Evans T, Blackburn TM, Jeschke JM, Probert AF, Bacher S. Application of the

Socio-Economic Impact Classification for Alien Taxa (SEICAT) to a global assessment

of alien bird impacts. NeoBiota. 2020;62:123-42.

35.   Defra. Wildlife management in England: a policy making framework for

resolving human-wildlife conflicts (withdrawn). 2010;

https://www.thenbs.com/publicationindex/documents/details?Pub=DEFRA&DocId=2

95502.

36.   Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of Google Scholar in

evidence reviews and its applicability to grey literature searching. PLOS One.

2015;7:e013823.

37. Hawkins CL, Bacher S, Essl F, Hulme PE, Jeschke JM, Kühn I, et al.

Framework and guidelines for implementing the proposed IUCN Environmental Impact

Classification for Alien Taxa (EICAT). Diversity and Distributions. 2015;21:1360-63.

38. Crawley D, Coomber F, Kubasiewicz L, Harrower C, Evans P, Waggitt J, et al.

Atlas of the mammals of Great Britain and Northern Ireland. The Mammal Society:

Pelagic Publishing, UK; 2020. 192 p.

39. Mathews F, Kubasiewicz LM, Gurnell J, Harrower CA, McDonald RA, Shore

RF. A review of the population and conservation status of British mammals: technical

summary. A report by the Mammal Society under contract to Natural England, Natural

Resources Wales and Scottish Natural Heritage. Peterborough: Natural England;

2018.

40. Harris S, Yalden DW. Mammals of the British Isles: handbook 4th edition.

Southampton: The Mammal Society; 2008.

41. R Core Team. R: A language and environment for statistics computing. In:

Computing RFfS, editor. Vienna, Austria2022.

42. Christensen R. *ordinal* - Regression Models for Ordinal Data. R package

version 2023.12-4.1. 2023; https://cran.r-

project.org/web/packages/ordinal/ordinal.pdf.

43. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen

A, et al. *glmmTMB* balances speed and flexibility among packages for zero-inflated

generalised linear mixed modelling. The R Journal. 2017;9:378-400.

44. Hartig F. *DHARMa*: Residual diagnostics for hierarchical (multi-level/mixed)

regression models. R package version 0.4.7. 2024; https://cran.r-

project.org/web/packages/DHARMa/vignettes/DHARMa.html.

45.   Bhatia S, Redpath SM, Suryawanshi K, Mishra C. Beyond conflict: exploring

the spectrum of human–wildlife interactions and their underlying mechanisms. Oryx.

2020;54(5):1-8.

46.   Defra. Government to end badger cull with new TB eradication strategy. 2024;

https://www.gov.uk/government/news/government-to-end-badger-cull-with-new-tb-

eradication-

strategy#:~:text=The%20new%20strategy%20will%20mark,the%20end%20of%20thi

s%20parliament.

47.   Petrovan SO, Ward AI, P. W. Detectability counts when assessing populations

for biodiversity targets. PLOS One. 2011;6:8.

48.   Sainsbury KA, Shore RF, Schofield H, Croose E, Campbell RD, McDonald RA.

Recent history, current status, conservation and management of native mammalian

carnivore species in Great Britain. Mammal Review. 2019;49(2):171-88.

49.   Gurnell J, Hare EJ. Rodents: Order Rodentia. In: Harris S, Yalden DW, editors.

Mammals of the British Isles. Southampton: The Mammal Society; 2008. p. 55-199.

50.   Bremner A, Park K. Public attitudes to the management of invasive non-native

species in Scotland. Biological Conservation. 2007;139:306-14.

51.   Taylor LH, Latham SM, Woolhouse ME. Risk factors for human disease

emergence. Philosophical Transactions of the Royal Society of London Series B,

Biological Sciences. 2001;356:983–89.

52.   Rahman T, Sobur A, Islam S, Ievy S, Hossain J, El Zowalaty ME, et al. Zoonotic

diseases: etiology, impact, and control. Microorganisms. 2020;8:1405.

53.   Vapalahti O, Mustonen J, Lundkvist Å, Henttonen H, Plyusnin A, Vaheri A.

Hantavirus Infections in Europe. Lancet Infectious Diseases. 2003;3:653–61.

54.   Salisbury AM, Begon M, Dove W, Niklasson B, Stewart JP. Ljungan virus is

endemic in rodents in the UK. Archives of Virology. 2014;159:547-51.

55.   Gray A, Capewell P, Zadoks R, Taggart MA, French AS, Katzer F, et al. Wild

deer in the United Kingdom are a potential reservoir for the livestock parasite *Babesia*

divergens.   Current   Research   in   Parasitology   &   Vector-Borne   Diseases.

2021;1:100019.

56.   Guardone L, Ebani VV, Verin R, Nardoni S, Consolazione A, Bennett M, et al.

Molecular detection of arthropod-borne pathogens in Eurasian badgers (*Meles meles*)

from the United Kingdom. Animals. 2020;10:446.

57.   Bedenham G, Kirk A, Luhano U, Shields A. The importance of biodiversity risks:

link to zoonotic diseases. British Actuarial Journal. 2022;27(e10):1-9.

58.   Puttock A, Graham HA, Cunliffe AM, Elliott M, Brazier RE. Eurasian beaver

activity increases water storage, attenuates flow and mitigates diffuse pollution from

intensively-managed grasslands. Science of the Total Environment. 2017;576:430-43.

59.   Auster RE, Puttock A, Brazier R. Unravelling perceptions of Eurasian beaver

reintroduction in Great Britain. Area. 2019;52:364-75.

60.   Macdonald DW, Johnson PJ. Foxes in the landscape: hunting, control, and

economics. In: Macdonald DW, Feber RE, editors. Wildlife Conservation on Farmland:

Volume 2, Conflict in the Countryside. Oxford: Oxford University Press; 2015. p. 20-

46.

61.   Baker P, Harris S, White PCL. After the hunt: the future for foxes in Britain.

London: International Fund for Animal Welfare. 2006; https://mknhs.org.uk/wp-

content/uploads/2015/08/IFAW-after-the-hunt.pdf; 2006.

62. Sun ZS, Wan EY, Agbana YL, Zhao HQ, Yin JX, Jiang TG, et al. Global One

Health index for zoonoses: A performance assessment in 160 countries and territories.

iScience. 2024;27:109297.

63. Hand A. Estimating feral cat densities using distance sampling in an urban

environment. Ecological Evolution. 2019;9:2699–705.

64. McDonald JL, Skillings E. Human influences shape the first spatially explicit

national estimate of urban unowned cat abundance. Scientific Reports.

2021;11:20216.

65. Baker PJ, Molony SE, Stone E, Cuthill IC, Harris S. Cats about town: is

predation by free-ranging pet cats Felis catus likely to affect urban bird populations? .

Ibis. 2008;150:86-99.

66. Fredrikson A. Of wildcats and wild cats: troubling species-based conservation

in the Anthropocene. Environment and Planning D: Society and Space. 2016;34(4):1-

25.

67. Lowe S, Browne M, Boudjelas S. 100 of the world's worst invasive alien

species: a selection from the Global Invasive Species database. Invasive Species

Specialist Group (ISSG), International Union for Conservation of Nature. 2000;

https://portals.iucn.org/library/sites/library/files/documents/2000-126.pdf.

68. Trouwborst A, McCormack PC, Camacho EM. Domestic cats and their impacts

on biodiversity: A blind spot in the application of nature conservation law. People and

Nature. 2019;2:235-50.

69. Kumschick S, Bacher S, Dawson W, Heikkila J, Sendek A, Pluess T, et al. A

conceptual framework for prioritization of invasive alien species for management

according to their impact. NeoBiota. 2012;15:69-100.

Supporting information

S1 File. Detailed description of all of the conflicts identified in the scientific

literature for 48 mammals of the British Isles.

S2 Table. List of species causing conflict, including order, status, population

size, body mass, biomass and type of conflicts recorded over time (pre-2014

only; both pre-2014 and 2014 to present; 2014 to present only). The median

impact score (1=minimal known impact to 5=highest known impact) and

confidence rating (1=low; 2=medium; 3=high) are shown after each conflict [x,x].

The highest impact scores are shown in bold.

Figure

Figure

Figure

Figure

Figure

Figure