DcaP-Family Porins are Required for Carboxylic Acid Utilization and Infection in *Acinetobacter*

*baumannii*

Hannah R. Noel[1], Nikita R. Kesav[1], Jonathan D. Winkelman[2], Lauren D. Palmer[1]*

[1] Department of Microbiology and Immunology, University of Illinois Chicago, Chicago, IL, USA

[2] Trestle LLC, Milwaukee, WI, USA

*Corresponding author

Lauren D. Palmer

835 S Wolcott Ave

MSB E703

Chicago, IL 60612

ldpalmer@uic.edu

**ABSTRACT**

*Acinetobacter baumannii* is a pathogen of concern and a leading cause of multidrug-

resistant healthcare-associated infections. The *A. baumannii* outer membrane is a barrier

to antimicrobials and host defenses but must also allow essential nutrients to permeate. Here,

we investigate the functional importance of the putative DcaP-family outer membrane porins,

which include a proposed vaccine target in *A. baumannii*. All *A. baumannii* genomes surveyed

encode multiple DcaP family porins, which we classify in four classes based on protein

sequence phylogeny (DcaP1-4). DcaP proteins encoded by species in other genera could

not be mapped to these DcaP classes and phylogenetic analysis suggests DcaP1-4

proteins diversified within *Acinetobacter*. Phenotypic array assays and additional experiments

show that the DcaP proteins were necessary for growth on multiple di- and tri-carboxylic acids

as sole carbon sources, including citric acid and tricarballylic acid. Specifically, DcaP3 was

required; however, DcaP1, DcaP2, and DcaP4 were all able to complement the total *dcaP*

knockout to varying degrees, suggesting partial functional overlap. Finally, a mutant lacking all

DcaP proteins was attenuated in the liver and spleen in a mouse model of bloodstream infection

and complemented by expression of DcaP3. However, a Δ*dcaP3* mutant had no defect,

demonstrating functional redundancy among DcaP proteins during infection. These findings

provide insight on how *A. baumannii* acquires nutrients through the outer membrane barrier and

show DcaP proteins are important during infection in specific host niches, validating their

potential as a therapeutic target.

**INTRODUCTION**

*Acinetobacter baumannii* is a pathogen of critical concern and a major cause of

healthcare-associated infections. Isolates of *A. baumannii* are often multidrug resistant (MDR)

or extensively drug resistant (XDR) and exhibit resistance to first and last line antibiotics, such

as meropenem and colistin, respectively (1). The Centers for Disease Control and the World

Health Organization have therefore identified *A. baumannii* as an urgent public health threat,

calling for new avenues of treatment (2–4). The bacterial cell envelope is the primary barrier

from the environment and protects bacteria from environmental stress including antibiotics and

host defenses; however, the cell envelope must also allow acquisition of essential nutrients and

cofactors (5,6). The outer membrane (OM) of *A. baumannii* is thought to be less permeable

compared to other Gram-negative bacteria such as *E. coli*, contributing to intrinsic antibiotic

resistance (7). How small molecule nutrients required for growth transit the impermeable OM in

*A. baumannii* is incompletely understood.

The OM of *A. baumannii* is an asymmetric lipid bilayer where the inner leaflet is

composed of glycerophospholipids and the outer leaflet of lipooligosaccharides (LOS) (5,8,9).

The OM bilayer is occupied by proteins that serve a variety of functions, including nutrient influx.

Nutrients typically pass through the OM via active transport by TonB-dependent transporters or

passive diffusion through porins, which are transmembrane channels that allow passage of

small molecules (6,10). The primary porin in *A. baumannii* is OmpA, which shows slow

molecular transport in liposome swelling assays, contributing to the low permeability barrier of

the OM (11–13). However, the primary function of OmpA is thought to be structural, providing a

physical tether between the OM and the peptidoglycan cell wall in *A. baumannii* and other

species (14,15). *A. baumannii* encodes multiple other porins including CarO, an OM porin that

selectively allows influx of ornithine and other basic amino acids (16), and the Occ family porins,

responsible for the influx of compounds like benzoate, hydroxycinnamate, and other aromatic

molecules (17–19). The protein DcaP is a predicted OM porin widely distributed across

*Acinetobacter* species named for its localization within a genetic locus for utilization of

dicarboxylic acids (*dca*) (20). Notably, a DcaP family protein is among the most abundant OM

proteins in *A. baumannii* during rat and mouse infections (21). Due to this fact, multiple groups

have studied a DcaP-like protein, which we define below as DcaP3, as a promising vaccine

candidate for protection against *A. baumannii* infection (21–25). However, the biological role of

DcaP family proteins remains unclear and their role in dicarboxylic acid utilization has not been

experimentally interrogated.

First identified in Parke *et al*. as part of the dicarboxylic acid catabolic operon of

*Acinetobacter baylyi* ADP1, *dcaP* (putative dicarboxylic acid porin) and DcaP-like proteins were

thought to be specific to the Moraxellaceae family (20). The structure for one DcaP-like protein

has been solved, presenting as a homo-trimeric 16-stranded beta-barrel protein with an

extended periplasmic N-terminus that forms a coiled coil (21). Due to the genetic localization

and induction of expression with the dicarboxylic acid adipate, *A. baylyi* DcaP porins were

proposed to be nonspecific porins that facilitate the uptake of dicarboxylic acids (20); however, a

role for DcaP porins in carboxylic acid catabolism has not been reported. Molecular dynamics

and electrophysiology studies determined a DcaP-like protein to have an affinity for anionic

compounds as substrates, such as the clinically relevant β-lactamase inhibitor sulbactam (21).

In other studies, DcaP-like proteins have been implicated in biofilm formation and mucin

catabolism (26,27). Here, we describe four classes of DcaP-family porins in *Acinetobacter* and

investigate their role in carbon source utilization in *A. baumannii* ATCC 17978 and AB5075

strains.

**RESULTS**

***Acinetobacter spp. encode multiple DcaP-family porins***

The first DcaP was described in *A. baylyi* as part of a dicarboxylic acid catabolic operon

(20), and DcaP-like proteins have been identified in various *A. baumannii* clinical isolates and

type-strains (21,23). Proteins designated DcaP-like in *A. baumannii* ATCC 17978 are

homologous to *A. baylyi* DcaP, sharing ~30-35% amino acid identity by Clustal Omega multiple

sequence alignment (28). However, given the abundance of DcaP-family porins within individual

*Acinetobacter* genomes, we hypothesized that these proteins could be further classified based

on protein sequence. Using DcaP and DcaP-like protein sequences from a set of deduplicated

*A. baumannii* genomes, four distinct classes of DcaP-family porins were identified that cluster

independently, including the original *A. baylyi* DcaP which we display as DcaP1 (Fig. 1A, S1A).

We next determined the distribution of DcaP-family porins across the *Acinetobacter* genus.

Using a set of 250 deduplicated *Acinetobacter* genomes, 235 of which were *A. baumannii*,

DcaP-family porins were shown to be widely conserved across the genus (Fig. S1B).

Additionally, *Acinetobacter* species within the *A. calcoaceticus/baumannii* (*acb*) complex/clade

of pathogenic *Acinetobacter* appeared to be enriched for having multiple DcaP-family porins

compared to non-pathogenic *Acinetobacter* species (Fig. S1B). Among the *A. baumannii*

genomes, over 130 of the strains encode four *dcaP* genes (Fig. 1B). In fact, no *A. baumannii*

genome analyzed encoded fewer than two *dcaP* genes (Fig. 1B). The type-strain *A. baumannii*

ATCC 17978 used for most studies here encodes *dcaP1*, *dcaP2*, and *dcaP3* encoded in three

distinct loci (Fig. S2A). To determine if the *dcaP* genes were equally distributed across the

species, the number of each DcaP subfamily were tallied for each genome and organized based

on how many total DcaP-family porins were present. The four classes of DcaP-family porins

were equally likely to be encoded in a given *A. baumannii* genome (Fig. 1B). For example, in

most cases where a genome encodes four DcaP-family porins, it was most likely to have one of

each variant. Alternatively, in a genome encoding five DcaP-family porins, DcaP2 was most

likely to be duplicated. Finally, *A. baumannii* AB5075 encodes *dcaP1-3* and also encodes *dcaP4*

in another distinct locus (Fig. S2A). These data suggest that DcaP-family porins are encoded

across the *Acinetobacter* genus, and enriched in pathogenic *Acinetobacter,* including *A.*

*baumannii*.

**_DcaP proteins from other bacterial species are distinct from Acinetobacter DcaP proteins_**

To investigate the evolutionary history of DcaP-family porins in _Acinetobacter_ and

determine if DcaP diversified within _Acinetobacter_ or across taxa, orthologues in other bacterial

taxa were identified through KEGG. Bacterial taxa were selected for further analysis if the DcaP-

like protein was encoded in more than half of the genomes for a given genus (29,30). A rooted

phylogenomic tree was generated to demonstrate the evolutionary distance between

_Acinetobacter_ and the bacteria containing a DcaP orthologue (Fig. 2A). Orthologues of DcaP-

family porins were found in distantly related organisms, such as _Shewanella spp._ and

_Xanthomonas spp.,_ albeit generally in fewer numbers (Fig. 2A). These DcaP orthologues could

not be further classified into one of the four DcaP classes defined in _A. baumannii_. This

suggests that the four DcaP classes of proteins defined here are unique to _Acinetobacter spp_.

To better understand the relationship between _Acinetobacter_ DcaP-family porins and proteins

designated DcaP-like from distantly related species, an unrooted tree of protein sequences was

generated. This analysis revealed that DcaP-family porins from AB5075, the included reference

_A. baumannii_ strain which encodes DcaP1-4, were more similar to each other than to non-

_Acinetobacter_ DcaP-family porins (Fig. 2B). This suggests that diversification of the four DcaP

classes defined here occurred within _Acinetobacter_. In contrast to _A. baumannii_, multiple DcaP-

like proteins encoded in other organisms were sequence divergent, such as in

_Pseudoxanthomonas daejeonensis_, _Stenotrophomonas spp._, _Shewanella psychrotolerans_, and

_Marinobacter spp._ (Fig 2B). Comparison of the species phylogeny and DcaP protein phylogeny

suggest there may have been occurrences of horizontal gene transfer in other species outside

_Acinetobacter_. Overall, these data suggest that DcaP-like proteins are encoded by species

across Pseudomonodota and that DcaP diversification has occurred within the _Acinetobacter_

clade.

***DcaP family porins are necessary for growth on select di- and tri-carboxylic acids***

Next, we hypothesized that the DcaP-family porins are involved in small molecule

uptake. One study suggested that DcaP3 may translocate the clinically relevant beta-lactamase

inhibitor sulbactam based on electrophysiology and applied field simulation data (21). Many *A.*

*baumannii* isolates are intrinsically resistant to the beta-lactam ampicillin due to the expression

of beta-lactamases and sulbactam has been shown to resensitize resistant isolates to beta-

lactam antibiotics (31,32). We reasoned that if DcaP-family porins facilitated sulbactam entry, a

mutant lacking all *dcaP* genes would exclude sulbactam from entering the cell and result in

increased resistance to the beta-lactam ampicillin while a wildtype strain would exhibit re-

sensitization to ampicillin. We therefore constructed a triple mutant in ATCC 17978 lacking all

encoded DcaP proteins, Δ*dcaP1*Δ*dcaP2*Δ*dcaP3* (ΔΔΔ*dcaP1-3*). However, the wildtype and the

ΔΔΔ*dcaP1-3* mutant strains exhibited similar susceptibility to ampicillin/sulbactam treatment

(Fig. S2). These data suggest that DcaP-family porins are not required for sulbactam activity.

We next hypothesized that DcaP-family porins may be important for growth on certain

nutrient sources such as dicarboxylic acids as previously suggested (20). The growth of

wildtype and ΔΔΔ*dcaP1-3* strains were compared using the high-throughput Biolog PM system

to screen for carbon source utilization. Multiple compounds were identified on which ΔΔΔ*dcaP1-*

*3* grew more than wildtype and compounds on which wildtype was able to grow but ΔΔΔ*dcaP1-3*

had a defect (Fig. 3A-B). The screen suggested ΔΔΔ*dcaP1-3* had increased growth compared

to wildtype on amino acids such as glutamic acid and glutaminic acid (Fig. 3A-B); however,

validation showed that there was no difference in growth between wildtype and ΔΔΔ*dcaP1-3*

(Fig. S3B). Glutamic acid was thus used as a control for subsequent assays to ensure growth of

strains in minimal media. Butyric acid also appeared to better support growth of ΔΔΔ*dcaP1-3*

and was analyzed in subsequent assays. Many of the compounds on which WT grew better

than ΔΔΔ*dcaP1-3* were carboxylic acids, including simple carboxylic acids, hydroxy carboxylic

acids, and sugar acids (Fig. 3A-B). These data suggest that DcaP-family porins may allow the

influx of carboxylic acid-containing compounds, as previously predicted (20).

***DcaP3 is important for growth on multiple carboxylic acid carbon sources***

To verify the Biolog results and further determine functional importance of the DcaP

classes, growth curves using ΔΔΔ*dcaP1-3*, Δ*dcaP1*Δ*dcaP2*, Δ*dcaP1*Δ*dcaP3*, and

Δ*dcaP2*Δ*dcaP3* and carboxylic acids as sole carbon sources were performed. The strains were

also assayed on glutamic acid and succinic acid as controls for growth in minimal media, and

butyric acid as a potential carbon source in which ΔΔΔ*dcaP1-3* grew to higher yield than WT.

There were no differences in growth in glutamic acid, and potential minor defects for growth of

the ΔΔΔ*dcaP1-3* strain in succinic acid and butyric acid. This showed that the screen did not

identify any conditions where the ΔΔΔ*dcaP1-3* strain grew more than WT. By contrast, mutants

lacking DcaP3 were defective for growth on citric acid, tricarballylic acid, and had a minor defect

on mucic acid compared to the WT (Fig. 4A). We were unable to identify conditions in which

DcaP1 or DcaP2 were required for growth. Similarly, a single Δ*dcaP3* mutation conferred a

defect in growth on citric acid, tricarballylic acid, and mucic acid, whereas single mutants of

Δ*dcaP1* or Δ*dcaP2* showed no defect (Fig. 4B).

A recent study in *A. baylyi* found that tricarballylic acid metabolism is closely linked with

the metabolism of *cis*- and *trans*-aconitic acid (33). The genetic organization of the tricarballylic

acid catabolic genes is conserved in *A. baumannii* compared to *A. baylyi* (Fig. S4A). Since the

data thus far suggest DcaP3 is important for growth on tricarballylic acid, we hypothesized that

DcaP3 may also be important for *cis*- and *trans*-aconitic acid growth. Mutants lacking DcaP3

had a defect for growth on both *cis*- and *trans*-aconitic acid (Fig. 4A) and the Δ*dcaP3* single

mutant had a defect in growth on both *cis*- and *trans*-aconitic acid compared to wildtype (Fig.

4B). To highlight the similarities in the carbon sources in which DcaP3 is important for growth,

the chemical structures for each carbon source are outlined in Fig. 4C. Given the lack of

identifiable phenotype for ΔdcaP1 or ΔdcaP2, transcript reads from previously published

datasets were analyzed to determine if *dcaP1* or *dcaP2* are expressed (34). In *A. baumannii*

17978, *dcaP3* was expressed at a high level in untreated lysogeny broth while *dcaP1* and

*dcaP2* were not expressed; in *A. baumannii* AB5075, *dcaP3* was similarly highly expressed

while *dcaP1*, *dcaP2*, and *dcaP4* were not expressed (Fig. 4D) (34). Similar results were

observed in another *A. baumannii* 17978 dataset (35). Taken together, these data suggest that

DcaP3 is highly expressed and important for growth using multiple di- and tri-carboxylic acids as

sole carbon sources.

***DcaP3 is required for growth on di- and tri- carboxylic acids in a recent clinical isolate***

To test if DcaP3 is important for growth on di- and tri-carboxylic acids in a more recent

clinical isolate, a ΔdcaP3 mutant was generated in *A. baumannii* AB5075, a member of the

widely circulating clonal complex 1 that was isolated in 2008 and also serves as a commonly

used reference strain (36). An *A. baumannii* AB5075 mutant lacking *dcaP3* was defective for

growth on citric acid, tricarballylic acid, mucic acid, and *cis*- and *trans*-aconitic acid compared to

wildtype (Fig. 5). This suggests that the requirement for DcaP3 for growth on select carboxylic

acids as sole carbon sources *in vitro* is conserved among *A. baumannii* isolates.

***DcaP1-4 can complement ΔΔΔdcaP for growth on some carboxylic acids***

Thus far, genetic disruption in *A. baumannii* ATCC 17978 and AB5075 suggested DcaP3 was

the primary DcaP important for utilization of certain di- and tri-carboxylic acids. However,

published expression analysis suggests only DcaP3 is expressed in laboratory conditions (Fig.

4D) (34,35). We hypothesized that individual DcaP proteins may complement the *A. baumannii*

17978 ΔΔΔdcaP1-3 growth defects if expressed from the *dcaP3* promoter. Therefore,

chromosomal mini-Tn*7* (mTn*7*) complementation constructs were constructed expressing *A.*

*baumannii* 17978 *dcaP1*, *dcaP2*, or *dcaP3* or *A. baumannii* AB5075 *dcaP4* from the *A.*

*baumannii* 17978 *dcaP3* promoter. Consistent with previous observations, the ΔΔΔ*dcaP* mutant

was unable to grow using citric acid, tricarballylic acid, or *cis*- and *trans*-aconitic acid and had a

slight defect on succinic acid, butyric acid, and mucic acid (Fig. 6). Growth was restored in

succinic acid and butyric acid by expressing *dcaP4* (Fig. 6)*,* suggesting DcaP4 can contribute to

succinic acid and butyric acid transport. Similarly, growth on citric acid, mucic acid, tricarballylic

acid, and trans- and cis-aconitic acid was restored to the ΔΔΔ*dcaP1-3* mutant by expressing

*dcaP2, dcaP3*, or *dcaP4* (Fig. 6). This suggests that DcaP2, DcaP3, and DcaP4 have

overlapping functions in at least some *A. baumannii* strains and conditions. Expression of *dcaP1*

in the ΔΔΔ*dcaP1-3* strain did not restore growth in succinic acid, butyric acid, citric acid, mucic

acid, or tricarballylic acid but did restore growth in *trans*- and *cis*-aconitic acid, suggesting a

largely non-overlapping functional role for DcaP1.

***DcaP3 is sufficient but not required for virulence in the liver and spleen during***

***bloodstream infection of mice***

Carbon source utilization can be a critical determinant of pathogenesis for *A. baumannii*

and other bacterial species in different host niches and DcaP3 has been proposed as an *A.*

*baumannii* vaccine candidate (21–25,37,38). Therefore, we sought to determine if DcaP-family

porins are important for *A. baumannii* pathogenesis in a murine model of bloodstream infection.

There was no defect of the ΔΔΔ*dcaP1-3* mutant compared to wildtype in lung, kidney, or heart

(Fig. 7A). By contrast, there was a defect for the ΔΔΔ*dcaP1-3* mutant compared to wildtype in

the spleen and liver (Fig. 7A). The defect in the spleen and liver was reversed by

complementing *dcaP3* in the chromosome of the ΔΔΔ*dcaP1-3* mutant, showing DcaP3 is

sufficient for proliferation in the liver and spleen (Fig. 7B). However, a single Δ*dcaP3* mutant had

no defect during infection (Fig. 7C). This suggests that in the host, DcaP1 and/or DcaP2 are

expressed and able to functionally complement the lack of DcaP3 in *A. baumannii* 17978.

Together these data show that DcaP-family porins are important in specific host niches during

infection.

**DISCUSSION**

The OM protects *A. baumannii* from antimicrobials and other host stresses but must

permit influx of nutrients essential for replication. How nutrients pass through the *A. baumannii*

OM is not well understood. Here, we designate classes of a family of predicted porins, DcaP1-4,

that are widely distributed across *Acinetobacter spp.* and show DcaP-family porins are required

for growth on multiple carboxylic acids.

DcaP-family porins have been identified in genetic screens for biofilm formation,

desiccation, and mucin catabolism, largely referred to as DcaP-like proteins without further

distinction (26,27,39–41). We show that *A. baumannii* DcaP-family porins can be grouped into

four classes based on protein sequence (Fig. 1A, S1A). DcaP-encoding genes are broadly

conserved among *Acinetobacter* spp, and particularly in *A. baumannii* and the *acb* complex (Fig.

1B, Fig. S1B). While DcaP-family porins can be found in distantly related organisms (Fig. 2A),

the DcaP1-4 classes we define in *A. baumannii* cluster together (Fig. 2B). These findings

suggest DcaP1-4 diversified within the *Acinetobacter* clade.

Antibiotic resistance in *A. baumannii* is a critical public health problem due to the

emergence of XDR isolates. Various porins in *A. baumannii* have been implicated in the uptake

of toxic molecules including antibiotics (12,42–44). DcaP3 was proposed to allow entry for the

beta-lactamase inhibitor sulbactam (21). Here, we show that a ΔΔΔ*dcaP1-3* mutant displays

similar susceptibility to ampicillin/sulbactam as the wildtype *A. baumannii* 17978, suggesting

that DcaP-family porins are not required for sulbactam function (Fig. S2). However, additional

studies would be required to determine whether sulbactam may enter via DcaP-family porins in

addition to other outer membrane porins.

The transport of essential nutrients across the cell envelope is incompletely understood

in *A. baumannii*. In general, *A. baumannii* utilizes organic acids and amino acids as carbon

sources and does not utilize sugars such as glucose (37,45). Accordingly, multiple OM channels

have been identified that are important for growth on specific carbon sources including ornithine,

arginine, glutamic acid, and benzoic acid (16–19,46). One study in ATCC 19606 found that

*dcaP1* (*A1S_1380*) was upregulated in the presence of mucin (26), suggesting that the host

environment may affect *dcaP* gene regulation. In a Biolog screen, we identified multiple

carboxylic acid carbon sources that supported significantly more growth of the wildtype

compared to the ΔΔΔ*dcaP1-3* mutant (Fig. 3), suggesting DcaP-family porins are important for

their diffusion through the OM.

The data here suggest DcaP3 is responsible for the influx of carboxylic acid carbon

sources citric acid, mucic acid, tricarballylic acid, and *cis*- and *trans*- aconitic acid in *A.*

*baumannii* ATCC 17978 and AB5075 in the *in vitro* conditions tested, as mutants lacking DcaP3

show decreased growth on these substrates (Fig. 4-5). Phylogenomics analysis found three *A.*

*baumannii* strains lacking *dcaP3*: *A. baumannii* E-011922, *A. baumannii* VB82, and *A.*

*baumannii* RBH2 (Fig. S1B) (47–49). Of these, carbon utilization data were reported for *A.*

*baumannii* E-011922, an environmental isolate that retains the ability to grow on citric acid and

mucic acid despite not encoding DcaP3 (48). Similarly, *A. baylyi* can grow on citric acid and

tricarballylic acid and does not encode DcaP3 but rather DcaP1, DcaP2, and DcaP4 (33). This

suggests that in strains that lack DcaP3, other DcaP-family porins are likely expressed at higher

abundance and allow transport of these carboxylic acids.

Citric acid is an intermediate of the tricarboxylic acid (TCA) cycle, a ubiquitous pathway

for energy generation. Studies quantifying citric acid in rat and mouse tissues report it is 0.15-

0.3 μmol/g in the liver and 0.27-0.45 μmol/g in the kidney (50–52), suggesting citric acid is

abundant in some tissues. Despite this, many bacteria cannot utilize citric acid as a carbon

source and how citric acid permeates the OM is unknown (53). Mutants in *P. aeruginosa* OpdH

were defective for growth on *cis*-aconitic acid, but not on citric acid, suggesting they encode

other citric acid transporters (54,55). A ΔΔΔ*dcaP1-3* mutant displayed a strong defect for growth

on citric acid that can be rescued by complementing with *dcaP2*, *dcaP3* or *dcaP4* (Fig. 4A, 4B,

Fig. 5). Therefore, encoding multiple porins for OM citric acid passage may be conserved

across bacterial genera.

Mucic acid, or galactaric acid, is a sugar acid derived from the oxidation of galactose and

is unrelated to mucin. Mucic acid is not typically found in the host; but, some antibiotic

treatments result in increased mucic acid in the human gut (56–58). However, only a subset of

*A. baumannii* strains, including ATCC 17978, encode the D-galactaric acid dehydrogenase

required for growth on mucic acid (59). The digestive tract is a reservoir for *A. baumannii*

colonization that increases the potential for infection and transmission and *A. baumannii* ATCC

17978 can asymptomatically colonize the mouse gut (60–65). Thus, mucic acid may be used as

carbon source by some *A. baumannii* in the inflamed gut. Our data suggest that DcaP2, DcaP3

or DcaP4 are important for growth on mucic acid.

The catabolism of *cis*- and *trans*-aconitic acid has been described in *A. baylyi*, and the

catabolism of tricarballylic acid in *A. baylyi* and *Salmonella* (33,66,67). However, the OM porin

required for growth on these carbon sources in *Acinetobacter* was unknown. Our data suggest

that DcaP2, DcaP3, and DcaP4 porins are important for growth on tricarballylic acid, and that

DcaP1 can also support growth on *cis*- and *trans*-aconitic acid.

Overall, our data suggests that DcaP3 is the major porin for passive transport of specific

carboxylic acid carbon sources in the *in vitro* conditions tested. The requirement for DcaP3 in

growth on select di- and tri-carboxylic acids is conserved in the more recent clinical isolate *A.*

*baumannii* AB5075, as a Δ*dcaP3* mutant had reduced growth in these conditions (Fig. 6).

Published RNA sequencing data show that *dcaP3* is the only *dcaP* gene expressed under

laboratory conditions tested, consistent with the defects for the Δ*dcaP3* single mutants in *A.*

*baumannii* 17978 and AB5075 strain backgrounds (Fig. 4B, Fig. 5) (34,35).

During infection, bacteria must scavenge nutrients from the host to replicate. The *A.*

*baumannii* ATCC 17978 ΔΔΔ*dcaP1-3* mutant had a defect compared to wildtype in the mouse

liver and spleen that was complemented by expression of *dcaP3* (Fig. 7A-B). However, the

Δ*dcaP3* single mutant had no defect (Fig. 7C). Therefore, DcaP3 passive import of carboxylic

acid carbon sources is sufficient but not necessary for pathogenesis in the liver and spleen. This

suggests the host environment may lead to increased expression of other *dcaP* genes;

regulation of *dcaP* expression and the molecular basis for substrate specificity is an area for

future study. Additionally, future study will be required to dissect the relevant carboxylic acid

carbon sources and DcaP substrates important for *A. baumannii* infection. There is little known

about the role for these carboxylic acid carbon sources during bacterial infection. Citric acid has

been shown to serve as a signaling molecule for *Staphylococcus aureus* pathogenesis; citric

acid was increased in the lung at 4 weeks during *Mycobacterium tuberculosis* infection, and

citric acid utilization promoted *Vibrio cholerae* small intestine colonization (68,69). The *Klebsiella*

*pneumoniae* citric acid synthase GltA has been shown to be important due to amino acid

biosynthesis required in the spleen, liver, and gut, but dispensable in the bloodstream (70).

Succinic acid has also been shown to be an important carbon source for *Pseudomonas*

*aeruginosa* during infection (71). Alternatively, DcaP-family porins may play another role in

pathogenesis. For example, porins in *A. baumannii* have demonstrated roles for the adhesion or

invasion of eukaryotic cells (13,72–75). DcaP3 has been proposed as a vaccine candidate due

to its high expression in infection models and immunogenicity (21–25). These data suggest that

vaccine strategies to functionally block DcaP proteins must block multiple DcaP proteins to

affect *A. baumannii* virulence.

In summary, we further classify a family of porins previously termed DcaP-like into four

classes in *Acinetobacter* spp. This classification allows for the differentiation between DcaP-like

proteins and the potential to determine individual functions for DcaP1-4 in *Acinetobacter* spp.

DcaP3 was critically important for growth on citric acid, tricarballylic acid, mucic acid, and *cis*-

and *trans*-aconitic acid as sole carbon sources. In a murine model of bloodstream infection,

DcaP3 was sufficient but not necessary for infection, suggesting functional overlap among DcaP

porins in the host. Expressing *dcaP2*, *dcaP3*, or *dcaP4* complemented growth of the *A.*

*baumannii* 17978 ΔΔΔ*dcaP1-3* mutant on all carbon sources tested and expression of *dcaP1*

complemented growth on *trans-* and *cis*-aconitic acid (Fig. 5). Our data therefore provides a

deeper understanding of how *A. baumannii* acquires nutrients including in the host.

**MATERIALS AND METHODS**

<u>Bacterial strain construction and growth</u>

All bacterial strains and plasmids used in this study are listed in Tables S1 and S2,

respectively. The wildtype strain was *A. baumannii* ATCC 17978VU or AB5075 (36,76,77).

Oligonucleotides are listed in Table S3 and were purchased from Integrated DNA Technologies

(Coralville, IA). Strains were grown in lysogeny broth (LB; Miller) or on LB with 1.5% (wt/vol)

agar unless indicated otherwise. Antibiotics were used at the following concentrations:

carbenicillin (Carb), 75 mg/L; kanamycin (Kan), 40 mg/L; chloramphenicol, 15 mg/L; hygromycin

(Hyg) 300 mg/L. For deletion constructs, 1000 bp of 3' and 5' flanking DNA to the gene of

interest was amplified using Q5 High Fidelity 2x Master Mix [New England Biolabs (NEB),

Ipswich, MA] from *A. baumannii* ATCC 17978VU or AB5075. The vector pFLP2 was digested

with BamHI and KpnI restriction enzymes (NEB, Ipswitch, MA) and gel purified (78). The

kanamycin or hygromycin cassette was amplified from pKD4 or pLDP357, respectively (79). A

hygromycin-resistant excisable cassette was generated by replacing kanamycin resistance in

pKD4 with hygromycin resistance amplified from an *A. baumannii* AB5075 transposon T101

mutant (36,77) with XR185/XR186 oligonucleotides and ligation with NEBuilder® HiFi DNA

Assembly Master Mix (NEB, Ipswitch, MA) to generate pLDP357. Fragments were ligated

together using HiFi ligation mix (NEB, Ipswitch, MA), transformed into chemically competent *E.*

*coli* DH5$\alpha$, and plated to selective media. Transformants were confirmed to contain the insert by

PCR. To generate the knockout strains, a triparental conjugation using helper *E. coli* strain

HB101 containing pRK2013 was performed. Single-crossover merodiploids containing the

integrated pFLP2 vector were chosen based on a Kan$^R$ Carb$^R$ or Hyg$^R$ and sucrose-sensitive

(10% w/v; Suc$^S$) phenotype. Putative merodiploids displaying were verified by PCR.

Merodiploids were then grown on LB agar, resuspended, and plated to LB agar or LB agar with

10% sucrose to select for colonies with the second crossover event. Colonies were then

screened for a Kan$^R$ Carb$^S$ Suc$^R$ or Hyg$^R$ Suc$^S$ phenotype. Mutants were confirmed via PCR.

For chromosomal integration constructs, DNA from ATCC 17978VU or AB5075 was

amplified using Q5 High Fidelity 2x Master Mix (NEB, Ipswich, MA). The mTn*7* pKNOCK vector

was digested with BamHI and KpnI prior to ligation using HiFi ligation mix (NEB, Ipswitch, MA)

and transformation into chemically competent *E. coli* DH5$\alpha$ $\lambda$*pir116*$^+$ (80). To generate

chromosomally integrated mTn7 constructs, a modified four-way parental mating was performed

using helper *E. coli* strain HB101 containing pRK2013 and *E. coli* DH5$\alpha$ containing pTNS2 (81).

Colonies were screened by PCR for integration into the correct chromosomal locus at the *att*

site. All plasmid constructs were verified by whole plasmid sequencing before transforming into

*A. baumannii* (Plasmidsaurus, South San Francisco, CA).

The *A. baumannii* AB5075 wildtype and $\Delta$*dcaP3*::hyg mutant were verified to have the

virulent/opaque (VIR-O) phenotype on 1/2X LB agar plates (82,83).

<u>Phylogenetic tree construction</u>

To identify DcaP orthologs and classify them into hierarchical orthogroups (HOGs),

OrthoFinder v2.5.5 was used (84). Two datasets were analyzed: (i) a 253-strain dataset

including 235 *A. baumannii* strains deduplicated as previously described (85); (ii) a 28-species

dataset that included multiple *Acinetobacter* and a broader range of gammaproteobacteria

rooted by one alphaproteobacterial. The protein accession numbers and genome accession

numbers are listed in Table S4.

Non-Acinetobacter species were selected based on the conservation of DcaP orthologs

across the genus or species indicated by KEGG. Specific organisms were selected to portray a

representative distribution of DcaP orthologs across bacteria, a group of 22 species. The

FastTree v2.1.11-derived (86), manually rooted species tree was used as an input for

OrthoFinder HOG inference (84,87), ensuring accurate classification of gene duplication and

speciation events. The resulting DcaP classifications were visualized and annotated in iTOL

(88). To infer a species phylogeny, a set of 120 universal bacterial marker genes (Bac120) was

identified from each proteome using HMMER3 (hmmsearch, v3.3), following the methodology

described in Parks *et al*. 2018 (89). The majority of species had only one significant hit for each

of the 120 protein HMMs; if a species had more than one hit, the best-scoring hit for that marker

was extracted.

Each of the 120 marker proteins was independently aligned to its respective HMM profile

using hmmalign (HMMER v3.3). Alignments were trimmed to remove poorly aligned regions and

concatenated into a single concatenated supermatrix for each species. To improve alignment

quality, columns were filtered based on the following criteria: (i) highly gapped regions (>80%

gaps) were removed; (ii) completely homogenous columns were excluded to reduce

redundancy.

Maximum likelihood species trees were inferred from the filtered concatenated alignment

using FastTree v2.1.11, with the LG model of amino acid evolution. The resulting unrooted trees

were manually rooted and annotated in iTOL using *Brevundimonas subvibrioides* or *E. coli* as

outgroups for the smaller and larger datasets.

DcaP protein phylogeny

DcaP homologs were extracted from their respective proteomes by selecting sequences

within the DcaP HOG at the root of all *Acinetobacter* species in the 255-species dataset in

Orthofinder. To ensure consistent domain identification, sequences were aligned to the Pfam

Hidden Markov Model (HMM) PF19577.4, corresponding to the DcaP OM protein family.

Alignments were trimmed to remove poorly aligned residues prior to phylogenetic inference. A

maximum likelihood phylogenetic tree was constructed using FastTree v2.1.11 with the LG

model of amino acid evolution. The inferred clusters were identical to OrthoFinder's DcaP

subfamilies, except for a small cluster of DcaP proteins that OrthoFinder grouped with DcaP1,

while FastTree grouped the same cluster with DcaP3.

To identify putative orthologs of DcaP proteins from the 22 species identified above and

*A. baumannii* AB5075, proteomes from genomic assemblies were downloaded from the NCBI

Datasets database. Two complementary approaches were employed to verify orthology. First,

HMMER v3.4 was used to search each proteome with the Pfam profile HMM PF19577,

corresponding to the DcaP protein family. Significant hits were retained as candidate DcaP

homologs. To independently assess orthology and reconstruct gene relationships, OrthoFinder

v2.5.5 was run on all 23 proteomes, which also produced a rooted gene tree for the DcaP

family. The resulting DcaP sequences were realigned to the PF19577 HMM using hmmalign,

and poorly aligned or unaligned amino acids were trimmed. A maximum likelihood phylogeny

was then inferred using RAxML v8.2.12, under the LG amino acid substitution model.

Amino acid sequence alignment and percent identity were determined using the Clustal

Omega program (1.2.4) through UniProt and the DcaP-family porin secondary structure was

based on PDB entry 6EUS (28,90,91).

Carbon source utilization screen using Biolog plates

Plates PM1 and PM2A were obtained from Biolog for carbon source utilization testing. A

110 $\mu$L aliquot of defined M9 minimal media lacking carbon was dispensed into every well and

resuspended. Then, 100 $\mu$L was transferred from the Biolog plates into sterile flat-bottom 96-

well plates. Overnight bacterial cultures were diluted 100-fold in sterile PBS before 1 $\mu$L of

diluted culture was inoculated into every well. Plates were incubated at 37°C with shaking and

growth was monitored by optical density at 600 nm ($OD_{600}$) in an EPOCH2 or SynergyH1 BioTek

plate reader (Winooski, VT). Background absorbance was subtracted before area under the

curve calculations for each isolate and each compound were calculated and the difference

between wildtype and $\Delta\Delta\Delta dcaP1$-$3$ was analyzed. Compounds were included if any isolate of

either wildtype or $\Delta\Delta\Delta dcaP1$-$3$ reached an $OD_{600}$ of 0.15 or higher.

<u>Growth curve analyses</u>

M9 minimal medium without glucose was supplemented 0.1X Vishniac's trace minerals

(92) with the specified carbon source at a concentration to maintain the carbon equivalency of a

4-carbon compound at 16.5 mM. Cultures were inoculated with a single bacterial colony and

grown in 3 mL LB at 37°C with shaking at 180 rpm for 8-16 hours. Growth curves were

performed by diluting LB cultures 100-fold in sterile PBS and inoculating 1 $\mu$L diluted culture into

99 $\mu$L media in a 96-well flat bottom plate. Growth was monitored as described above.

<u>Murine bloodstream infection</u>

Six-week-old female and male ND4 Swiss Webster mice were purchased from

Envigo/Harlan Laboratory. Mice were boarded in a temperature-controlled environment with

14:10 h light:dark cycles and food and water were provided as needed. Mice were acclimated to

the facility for 1 week prior to infection. Mice were anesthetized with isoflurane and inoculated

retro-orbitally with approximately 3 × $10^8$ CFU in a 50 $\mu$L bacterial suspension of a 1:1 mixture of

*A. baumannii* ATCC 17978VU wildtype and ΔΔΔ*dcaP1-3*::Kn mutant, ΔΔΔ*dcaP1-3*::Kn+*dcaP3*,

or Δ*dcaP3*::Kn. For wildtype versus ΔΔΔ*dcaP1-3,* 5 of the mice were infected with wildtype

ATCC 17978VU and 5 were infected with ATCC 17978 mTn*7*(Carb$^R$); for the wildtype versus

Δ*dcaP3*, wildtype was ATCC 17978 mTn*7*(Carb$^R$). All mice infected with the complement strain

were infected with ATCC 17978VU. The inoculum dose was determined by serial dilution and

plating on selective agar media. Mice were euthanized at 24 h post infection by

$CO_2$ asphyxiation, and the organs were excised aseptically. Tissues were homogenized in PBS

using a Bullet Blender (Next Advance, Troy, NY), and all samples were serially diluted and

plated on LB and LB with carbenicillin or kanamycin selective agar plates for bacterial

enumeration. All animal care protocols were approved by the University of Illinois Chicago

Institutional Animal Care and Use Committee (IACUC; protocol number 23–119) in accordance

with the Animal Care Policies of UIC, the Animal Welfare Act, the National Institutes of Health,

and the American Veterinary Medical Association (AVMA). Animals were humanely euthanized

consistent with the AVMA guidelines.

Data reporting, statistical analysis, and figure preparation

Each measurement was taken from a distinct biological sample (e.g., bacterial culture

from a single colony or an individual mouse). Data processing and statistical analyses were

performed using Microsoft Excel 16.98 and GraphPad Prism 10.4.2 or 10.6.1. Statistical tests

used are indicated in each figure legend. Figures were prepared in Adobe Illustrator 25.2.3 and

30.2.1.

Author contributions

H.R.N. – project conceptualization, methodology, data collection, data analysis, and writing.

N.R.K. – data collection, data analysis, and writing. J.D.W – bioinformatics, data analysis, and

writing. L.D.P. – project conceptualization, methodology, data analysis, writing, funding, and

resources.

## ACKNOWLEDGMENTS

499    We thank the members of the Palmer laboratory for critical reading of the manuscript, Xiaomei

500    Ren for construction of pLDP357, and funding from the National Institutes of Health

501    R01AI189516 to L.D.P. and the Michael Reese Research and Education Foundation Pioneer in

502    Research Award.

**REFERENCES**

1. Clark NM, Zhanel GG, Lynch JPI. Emergence of antimicrobial resistance among *Acinetobacter* species: a global threat. Curr Opin Crit Care. 2016 Oct;22(5):491. doi:10.1097/MCC.0000000000000337

2. World Health Organization. No time to wait: securing the future from drug-resistant infections [Report to the Secretary-General of the United Nations]. World Health Organization and Interagency Coordination Group on Antimicrobial Resistance; 2019 Apr. Report No.

3. Weiner LM, Webb AK, Limbago B, Dudeck MA, Patel J, Kallen AJ, et al. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011–2014. Infect Control Hosp Epidemiol. 2016 Nov;37(11):1288–301. doi:10.1017/ice.2016.174

4. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis. 2018 Mar 1;18(3):318–27. doi:10.1016/S1473-3099(17)30753-3

5. Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. Cold Spring Harb Perspect Biol. 2010 May;2(5):a000414. doi:10.1101/cshperspect.a000414 PubMed PMID: 20452953; PubMed Central PMCID: PMC2857177.

6. Nikaido H. Molecular basis of bacterial outer membrane permeability revisited. Microbiol Mol Biol Rev. 2003 Dec;67(4):593–656. doi:10.1128/mmbr.67.4.593-656.2003

7. Geisinger E, Huo W, Hernandez-Bird J, Isberg RR. *Acinetobacter baumannii*: envelope determinants that control drug resistance, virulence, and surface variability. Annu Rev Microbiol. 2019;73(1):481–506. doi:10.1146/annurev-micro-020518-115714 PubMed PMID: 31206345.

8. Kenyon JJ, Hall RM. Variation in the complex carbohydrate biosynthesis loci of *Acinetobacter baumannii* genomes. PLOS ONE. 2013 Apr 16;8(4):e62160. doi:10.1371/journal.pone.0062160

9. Kenyon JJ, Nigro SJ, Hall RM. Variation in the OC locus of *Acinetobacter baumannii* genomes predicts extensive structural diversity in the lipooligosaccharide. PLOS ONE. 2014 Sep 23;9(9):e107833. doi:10.1371/journal.pone.0107833

10. Silale A, Berg B van den. TonB-dependent transport across the bacterial outer membrane. Annu Rev Microbiol. 2023 Sep 15;77(Volume 77, 2023):67–88. doi:10.1146/annurev-micro-032421-111116

11. Sugawara E, Nikaido H. OmpA is the principal nonspecific slow porin of *Acinetobacter baumannii*. J Bacteriol. 2012 Jul 18;194(15):4089–96. doi:10.1128/jb.00435-12

12. Iyer R, Moussa SH, Durand-Réville TF, Tommasi R, Miller A. *Acinetobacter baumannii*
OmpA Is a selective antibiotic permeant porin. ACS Infect Dis. 2018 Mar 9;4(3):373–81.
doi:10.1021/acsinfecdis.7b00168

13. Scribano D, Cheri E, Pompilio A, Di Bonaventura G, Belli M, Cristina M, et al. *Acinetobacter*
*baumannii* OmpA-like porins: functional characterization of bacterial physiology, antibiotic-
resistance, and virulence. Commun Biol. 2024 Aug 6;7(1):1–11. doi:10.1038/s42003-024-
06645-0

14. Samsudin F, Ortiz-Suarez ML, Piggot TJ, Bond PJ, Khalid S. OmpA: a flexible clamp for
bacterial cell wall attachment. Structure. 2016 Dec 6;24(12):2227–35.
doi:10.1016/j.str.2016.10.009

15. Park JS, Lee WC, Yeo KJ, Ryu KS, Kumarasiri M, Hesek D, et al. Mechanism of anchoring
of OmpA protein to the cell wall peptidoglycan of the gram-negative bacterial outer
membrane. FASEB J. 2012 Jan;26(1):219. doi:10.1096/fj.11-188425 PubMed PMID:
21965596.

16. Mussi MA, Relling VM, Limansky AS, Viale AM. CarO, an *Acinetobacter baumannii* outer
membrane protein involved in carbapenem resistance, is essential for l-ornithine uptake.
FEBS Lett. 2007;581(29):5573–8. doi:10.1016/j.febslet.2007.10.063

17. Zahn M, Bhamidimarri SP, Baslé A, Winterhalter M, van den Berg B. Structural insights into
outer membrane permeability of *Acinetobacter baumannii*. Structure. 2016 Feb 2;24(2):221–
31. doi:10.1016/j.str.2015.12.009

18. Clark TJ, Momany C, Neidle EL. The *benPK* operon, proposed to play a role in transport, is
part of a regulon for benzoate catabolism in *Acinetobacter* sp. strain ADP1. Microbiology.
2002;148(4):1213–23. doi:10.1099/00221287-148-4-1213

19. Smith MA, Weaver VB, Young DM, Ornston LN. Genes for chlorogenate and
hydroxycinnamate catabolism (hca) are linked to functionally related genes in the *dca-pca-*
*qui-pob-hca* chromosomal cluster of *Acinetobacter* sp. strain ADP1. Appl Environ Microbiol.
2003 Jan;69(1):524–32. doi:10.1128/AEM.69.1.524-532.2003

20. Parke D, Garcia MA, Ornston LN. Cloning and genetic characterization of *dca* genes
required for β-oxidation of straight-chain dicarboxylic acids in *Acinetobacter* sp. strain ADP1.
Appl Environ Microbiol. 2001 Oct;67(10):4817–27. doi:10.1128/AEM.67.10.4817-4827.2001

21. Bhamidimarri SP, Zahn M, Prajapati JD, Schleberger C, Söderholm S, Hoover J, et al. A
multidisciplinary approach toward identification of antibiotic scaffolds for *Acinetobacter*
*baumannii*. Structure. 2019 Feb 5;27(2):268-280.e6. doi:10.1016/j.str.2018.10.021

22. Raoufi Z, Abdollahi S, Armand R. DcaP porin and its epitope-based subunit promise
effective vaccines against *Acinetobacter baumannii*; *in-silico* and *in-vivo* approaches. Microb
Pathog. 2022 Jan;162:105346. doi:10.1016/j.micpath.2021.105346 PubMed PMID:
34864145.

23. Fereshteh S, Ajdary S, Sepehr A, Bolourchi N, Barzi SM, Haririzadeh Jouriani F, et al.
Immunization with recombinant DcaP-like protein and AbOmpA revealed protections against
sepsis infection of multi-drug resistant *Acinetobacter baumannii* ST2Pas in a C57BL/6

mouse model. Microb Pathog. 2023 Jan;174:105882. doi:10.1016/j.micpath.2022.105882
PubMed PMID: 36403713.

24. Khalid K, Irum S, Ullah SR, Andleeb S. In-silico vaccine design based on a novel vaccine
candidate against infections caused by *Acinetobacter baumannii*. Int J Pept Res Ther. 2021
Dec 2;28(1):16. doi:10.1007/s10989-021-10316-7

25. Fereshteh S, Abdoli S, Shahcheraghi F, Ajdary S, Nazari M, Badmasti F. New putative
vaccine candidates against *Acinetobacter baumannii* using the reverse vaccinology method.
Microb Pathog. 2020 Jun 1;143:104114. doi:10.1016/j.micpath.2020.104114

26. Ohneck EJ, Arivett BA, Fiester SE, Wood CR, Metz ML, Simeone GM, et al. Mucin acts as a
nutrient source and a signal for the differential expression of genes coding for cellular
processes and virulence factors in *Acinetobacter baumannii*. PLoS ONE. 2018 Jan
8;13(1):e0190599. doi:10.1371/journal.pone.0190599 PubMed PMID: 29309434; PubMed
Central PMCID: PMC5757984.

27. Cabral MP, Soares NC, Aranda J, Parreira JR, Rumbo C, Poza M, et al. Proteomic and
functional analyses reveal a unique lifestyle for *Acinetobacter baumannii* biofilms and a key
role for histidine metabolism. J Proteome Res. 2011 Aug 5;10(8):3399–417.
doi:10.1021/pr101299j

28. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011
Oct 11;7:539. doi:10.1038/msb.2011.75 PubMed PMID: 21988835; PubMed Central PMCID:
PMC3261699.

29. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res.
2000 Jan 1;28(1):27–30. doi:10.1093/nar/28.1.27 PubMed PMID: 10592173; PubMed
Central PMCID: PMC102409.

30. Kanehisa M, Furumichi M, Sato Y, Matsuura Y, Ishiguro-Watanabe M. KEGG: biological
systems database as a model of the real world. Nucleic Acids Res. 2025 Jan
6;53(D1):D672–7. doi:10.1093/nar/gkae909 PubMed PMID: 39417505; PubMed Central
PMCID: PMC11701520.

31. Ji J, Du X, Chen Y, Fu Y, Wang H, Yu Y. In vitro activity of sulbactam in combination with
imipenem, meropenem, panipenem or cefoperazone against clinical isolates of
*Acinetobacter baumannii*. Int J Antimicrob Agents. 2013 Apr 1;41(4):400–1.
doi:10.1016/j.ijantimicag.2012.12.014

32. Choi JY, Soo Park Y, Cho CH, Seon Park Y, Shin SY, Song YG, et al. Synergic in-vitro
activity of imipenem and sulbactam against *Acinetobacter baumannii*. Clin Microbiol Infect.
2004 Dec 1;10(12):1098–101. doi:10.1111/j.1469-0691.2004.00987.x

33. Baugh AC, Defalco JB, Duscent-Maitland CV, Tumen-Velasquez MP, Laniohan NS, Figatner
K, et al. Regulation of tricarboxylate transport and metabolism in *Acinetobacter baylyi* ADP1.
Appl Environ Microbiol. 2024 Jan 30;90(2):e02111-23. doi:10.1128/aem.02111-23

34. Rokes AB, Santos-Lopez A, Cooper VS. History shapes regulatory and evolutionary
responses to tigecycline in two reference strains of *Acinetobacter baumannii*. Microbiology.

2025 Jun;171(6):001570. doi:10.1099/mic.0.001570 PubMed PMID: 40489210; PubMed
Central PMCID: PMC12149411.

35. Kröger C, MacKenzie KD, Alshabib EY, Kirzinger MWB, Suchan DM, Chao TC, et al. The
primary transcriptome, small RNAs and regulation of antimicrobial resistance in
*Acinetobacter baumannii* ATCC 17978. Nucleic Acids Res. 2018 Oct 12;46(18):9684–98.
doi:10.1093/nar/gky603 PubMed PMID: 29986115; PubMed Central PMCID: PMC6182133.

36. Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, et al. AB5075, a
highly virulent isolate of *Acinetobacter baumannii*, as a model strain for the evaluation of
pathogenesis and antimicrobial treatments. mBio. 2014 May 27;5(3):e01076-01014.
doi:10.1128/mBio.01076-14 PubMed PMID: 24865555; PubMed Central PMCID:
PMC4045072.

37. Ren X, Palmer LD. *Acinetobacter* metabolism in infection and antimicrobial resistance.
Infect Immun. 2023 May 16;91(6):e00433-22. doi:10.1128/iai.00433-22

38. Barber MF, Fitzgerald JR. Mechanisms of host adaptation by bacterial pathogens. FEMS
Microbiol Rev. 2024 Jul 1;48(4):fuae019. doi:10.1093/femsre/fuae019

39. Rumbo-Feal S, Gómez MJ, Gayoso C, Álvarez-Fraga L, Cabral MP, Aransay AM, et al.
Whole transcriptome analysis of *Acinetobacter baumannii* assessed by RNA-sequencing
reveals different mRNA expression profiles in biofilm compared to planktonic cells. PLoS
ONE. 2013 Aug 30;8(8):e72968. doi:10.1371/journal.pone.0072968 PubMed PMID:
24023660; PubMed Central PMCID: PMC3758355.

40. Shin JH, Lee HW, Kim SM, Kim J. Proteomic analysis of *Acinetobacter baumannii* in biofilm
and planktonic growth mode. J Microbiol. 2009 Dec 1;47(6):728–35. doi:10.1007/s12275-
009-0158-y

41. Gayoso CM, Mateos J, Méndez JA, Fernández-Puente P, Rumbo C, Tomás M, et al.
Molecular mechanisms involved in the response to desiccation stress and persistence in
*Acinetobacter baumannii*. J Proteome Res. 2014 Feb 7;13(2):460–76.
doi:10.1021/pr400603f PubMed PMID: 24299215.

42. Liu J, Chen S, Zhao B, Li G, Ma T. A novel FadL homolog, AltL, mediates transport of long-
chain alkanes and fatty acids in *Acinetobacter venetianus* RAG-1. Appl Environ Microbiol.
2022 Sep 28;88(20):e01294-22. doi:10.1128/aem.01294-22

43. Mussi MA, Limansky AS, Viale AM. Acquisition of resistance to carbapenems in multidrug-
resistant clinical strains of *Acinetobacter baumannii*: natural insertional inactivation of a
gene encoding a member of a novel family of β-barrel outer membrane proteins. Antimicrob
Agents Chemother. 2005 Apr;49(4):1432–40. doi:10.1128/AAC.49.4.1432-1440.2005
PubMed PMID: 15793123; PubMed Central PMCID: PMC1068641.

44. Catel-Ferreira M, Coadou G, Molle V, Mugnier P, Nordmann P, Siroy A, et al. Structure-
function relationships of CarO, the carbapenem resistance-associated outer membrane
protein of *Acinetobacter baumannii*. J Antimicrob Chemother. 2011 Sep;66(9):2053–6.
doi:10.1093/jac/dkr267 PubMed PMID: 21705362.

45. Juni E. Genetics and physiology of *Acinetobacter*. Annu Rev Microbiol. 1978;32:349–71.
doi:10.1146/annurev.mi.32.100178.002025 PubMed PMID: 360969.

46. Zahn M, D'Agostino T, Eren E, Baslé A, Ceccarelli M, van den Berg B. Small-molecule
transport by CarO, an abundant eight-stranded β-Barrel outer membrane protein from
*Acinetobacter baumannii*. J Mol Biol. 2015 Jul 17;427(14):2329–39.
doi:10.1016/j.jmb.2015.03.016

47. Ambrose SJ, Hamidian M, Hall RM. The extensively antibiotic resistant ST111 *Acinetobacter*
*baumannii* isolate RBH2 carries an extensive mobile element complement of plasmids,
transposons and insertion sequences. Plasmid. 2023 Sep;128:102707.
doi:10.1016/j.plasmid.2023.102707

48. Hamidian M, Maharjan RP, Farrugia DN, Delgado NN, Dinh H, Short FL, et al. Genomic and
phenotypic analyses of diverse non-clinical *Acinetobacter baumannii* strains reveals strain-
specific virulence and resistance capacity. Microb Genomics. 2022 Feb;8(2):000765.
doi:10.1099/mgen.0.000765 PubMed PMID: 35166651; PubMed Central PMCID:
PMC8942024.

49. Prity FT, Tobin LA, Maharajan R, Paulsen IT, Cain AK, Hamidian M. The evolutionary tale of
eight novel plasmids in a colistin-resistant environmental *Acinetobacter baumannii* isolate.
Microb Genomics. 2023 May;9(5):mgen001010. doi:10.1099/mgen.0.001010 PubMed
PMID: 37171842; PubMed Central PMCID: PMC10272872.

50. Spencer AF, Lowenstein JM. Citrate content of liver and kidney of rat in various metabolic
states and in fluoroacetate poisoning. Biochem J. 1967 May;103(2):342–8.
doi:10.1042/bj1030342 PubMed PMID: 6032977; PubMed Central PMCID: PMC1270414.

51. Ross KL, Eisenstein RS. Iron deficiency decreases mitochondrial aconitase abundance and
citrate concentration without affecting tricarboxylic acid cycle capacity in rat liver. J Nutr.
2002 Apr;132(4):643–51. doi:10.1093/jn/132.4.643

52. Tanner GA, Tanner JA. Citrate therapy for polycystic kidney disease in rats. Kidney Int. 2000
Nov;58(5):1859–69. doi:10.1111/j.1523-1755.2000.00357.x PubMed PMID: 11044205.

53. Blount ZD, Borland CZ, Lenski RE. Historical contingency and the evolution of a key
innovation in an experimental population of Escherichia coli. Proc Natl Acad Sci U S A. 2008
Jun 10;105(23):7899–906. doi:10.1073/pnas.0803151105 PubMed PMID: 18524956;
PubMed Central PMCID: PMC2430337.

54. Tamber S, Ochs MM, Hancock REW. Role of the novel OprD family of porins in nutrient
uptake in *Pseudomonas aeruginosa*. J Bacteriol. 2006 Jan;188(1):45–54.
doi:10.1128/JB.188.1.45-54.2006 PubMed PMID: 16352820; PubMed Central PMCID:
PMC1317591.

55. Tamber S, Maier E, Benz R, Hancock REW. Characterization of OpdH, a *Pseudomonas*
*aeruginosa* porin involved in the uptake of tricarboxylates. J Bacteriol. 2007
Feb;189(3):929–39. doi:10.1128/JB.01296-06 PubMed PMID: 17114261; PubMed Central
PMCID: PMC1797325.

56. Hwang I, Park YJ, Kim YR, Kim YN, Ka S, Lee HY, et al. Alteration of gut microbiota by
vancomycin and bacitracin improves insulin resistance via glucagon-like peptide 1 in diet-
induced obesity. FASEB J. 2015;29(6):2397–411. doi:10.1096/fj.14-265983

57. Theriot CM, Koenigsknecht MJ, Carlson PE, Hatton GE, Nelson AM, Li B, et al. Antibiotic-
induced shifts in the mouse gut microbiome and metabolome increase susceptibility to
*Clostridium difficile* infection. Nat Commun. 2014;5:3114. doi:10.1038/ncomms4114
PubMed PMID: 24445449; PubMed Central PMCID: PMC3950275.

58. Faber F, Tran L, Byndloss MX, Lopez CA, Velazquez EM, Kerrinnes T, et al. Host-mediated
sugar oxidation promotes post-antibiotic pathogen expansion. Nature. 2016 Jun
30;534(7609):697–9. doi:10.1038/nature18597 PubMed PMID: 27309805; PubMed Central
PMCID: PMC4939260.

59. Farrugia DN, Elbourne LDH, Hassan KA, Eijkelkamp BA, Tetu SG, Brown MH, et al. The
complete genome and phenome of a community-acquired *Acinetobacter baumannii*. PLOS
ONE. 2013 Mar 19;8(3):e58628. doi:10.1371/journal.pone.0058628

60. Timsit JF, Garrait V, Misset B, Goldstein FW, Renaud B, Carlet J. The digestive tract is a
major site for *Acinetobacter baumannii* colonization in intensive care unit patients. J Infect
Dis. 1993 Nov 1;168(5):1336–7. doi:10.1093/infdis/168.5.1336

61. Corbella X, Pujol M, Ayats J, Sendra M, Ardanuy C, Domínguez MA, et al. Relevance of
digestive tract colonization in the epidemiology of nosocomial infections due to multiresistant
*Acinetobacter baumannii*. Clin Infect Dis. 1996 Aug 1;23(2):329–34.
doi:10.1093/clinids/23.2.329

62. Wong SC, Chen JHK, Chau PH, So SYC, AuYeung CHY, Yuen LLH, et al. Gastrointestinal
colonization of carbapenem-resistant *Acinetobacter baumannii*: what is the implication for
infection control? Antibiotics. 2022 Sep 22;11(10):1297. doi:10.3390/antibiotics11101297
PubMed PMID: 36289955; PubMed Central PMCID: PMC9598245.

63. Harris AD, Johnson JK, Pineles L, O'Hara LM, Bonomo RA, Thom KA. Patient-to-patient
transmission of *Acinetobacter baumannii* gastrointestinal colonization in the intensive care
unit. Antimicrob Agents Chemother. 2019 Jul 25;63(8):10.1128/aac.00392-19.
doi:10.1128/aac.00392-19

64. Zheng Y, Xu N, Pang J, Han H, Yang H, Qin W, et al. Colonization with extensively drug-
resistant *Acinetobacter baumannii* and prognosis in critically ill patients: an observational
cohort study. Front Med. 2021;8:667776. doi:10.3389/fmed.2021.667776 PubMed PMID:
33996866; PubMed Central PMCID: PMC8119758.

65. Ren X, Clark RM, Bansah DA, Varner EN, Tiffany CR, Jaswal K, et al. Amino acid
competition shapes *Acinetobacter baumannii* gut carriage. BioRxiv Prepr Serv Biol. 2024
Oct 19;2024.10.19.619093. doi:10.1101/2024.10.19.619093 PubMed PMID: 39502362;
PubMed Central PMCID: PMC11537318.

66. Lewis JA, Stamper LW, Escalante-Semerena JC. Regulation of expression of the
tricarballylate utilization operon (*tcuABC*) of *Salmonella enterica*. Res Microbiol. 2009
Apr;160(3):179–86. doi:10.1016/j.resmic.2009.01.001 PubMed PMID: 19284970; PubMed
Central PMCID: PMC2692759.

67. Lewis JA, Horswill AR, Schwem BE, Escalante-Semerena JC. The tricarballylate utilization
(*tcuRABC*) genes of *Salmonella enterica* serovar Typhimurium LT2. J Bacteriol. 2004
Mar;186(6):1629–37. doi:10.1128/JB.186.6.1629-1637.2004 PubMed PMID: 14996793;
PubMed Central PMCID: PMC355976.

68. Chen F, Zhao Q, Yang Z, Chen R, Pan H, Wang Y, et al. Citrate serves as a signal molecule
to modulate carbon metabolism and iron homeostasis in *Staphylococcus aureus.* PLoS
Pathog. 2024 Jul;20(7):e1012425. doi:10.1371/journal.ppat.1012425 PubMed PMID:
39078849; PubMed Central PMCID: PMC11315280.

69. Liu M, Hao G, Li Z, Zhou Y, Garcia-Sillas R, Li J, et al. CitAB two-component system-
regulated citrate utilization contributes to *Vibrio cholerae* competitiveness with the gut
microbiota. Payne SM, editor. Infect Immun. 2019 Mar;87(3):e00746-18.
doi:10.1128/IAI.00746-18

70. Vornhagen J, Sun Y, Breen P, Forsyth V, Zhao L, Mobley HLT, et al. The *Klebsiella*
*pneumoniae* citrate synthase gene, *gltA*, influences site specific fitness during infection.
PLoS Pathog. 2019 Aug;15(8):e1008010. doi:10.1371/journal.ppat.1008010 PubMed PMID:
31449551; PubMed Central PMCID: PMC6730947.

71. Riquelme SA, Prince A. *Pseudomonas aeruginosa* consumption of airway metabolites
promotes lung infection. Pathogens. 2021 Jul 29;10(8):957.
doi:10.3390/pathogens10080957

72. Gaddy JA, Tomaras AP, Actis LA. The *Acinetobacter baumannii* 19606 OmpA protein plays a
role in biofilm formation on abiotic surfaces and in the interaction of this pathogen with
eukaryotic cells. Infect Immun. 2009 Aug;77(8):3150–60. doi:10.1128/IAI.00096-09 PubMed
PMID: 19470746; PubMed Central PMCID: PMC2715673.

73. Choi CH, Lee JS, Lee YC, Park TI, Lee JC. *Acinetobacter baumannii* invades epithelial cells
and outer membrane protein A mediates interactions with epithelial cells. BMC Microbiol.
2008 Dec 10;8:216. doi:10.1186/1471-2180-8-216 PubMed PMID: 19068136; PubMed
Central PMCID: PMC2615016.

74. Choi CH, Hyun SH, Lee JY, Lee JS, Lee YS, Kim SA, et al. *Acinetobacter baumannii* outer
membrane protein A targets the nucleus and induces cytotoxicity. Cell Microbiol.
2008;10(2):309–19. doi:10.1111/j.1462-5822.2007.01041.x

75. Labrador-Herrera G, Pérez-Pulido AJ, Álvarez-Marín R, Casimiro-Soriguer CS, Cebrero-
Cangueiro T, Morán-Barrio J, et al. Virulence role of the outer membrane protein CarO in
carbapenem-resistant *Acinetobacter baumannii*. Virulence. 2020 Dec;11(1):1727–37.
doi:10.1080/21505594.2020.1855912 PubMed PMID: 33300460; PubMed Central PMCID:
PMC7733888.

76. Wijers CDM, Pham L, Menon S, Boyd KL, Noel HR, Skaar E, et al. Identification of two
variants of *Acinetobacter baumannii* 17978 with distinct genotypes and phenotypes. Infect
Immun. 2021 Aug 30;IAI0045421. doi:10.1128/IAI.00454-21 PubMed PMID: 34460288.

77. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, et al. Resources for
genetic and genomic analysis of emerging pathogen *Acinetobacter baumannii*. J Bacteriol.
2015 Jun 15;197(12):2027–35. Located at: 25845845. doi:10.1128/JB.00131-15

78. Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP. A broad-host-range Flp-FRT
recombination system for site-specific excision of chromosomally-located DNA sequences:
application for isolation of unmarked *Pseudomonas aeruginosa* mutants. Gene. 1998 May
28;212(1):77–86. doi:10.1016/S0378-1119(98)00130-9

79. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli*
K-12 using PCR products. Proc Natl Acad Sci U S A. 2000 Jun 6;97(12):6640–5.
doi:10.1073/pnas.120163297 PubMed PMID: 10829079; PubMed Central PMCID:
PMC18686.

80. Carruthers MD, Nicholson PA, Tracy EN, Munson RS Jr. Acinetobacter baumannii utilizes a
type VI secretion system for bacterial competition. PLoS One. 2013;8(3):e59388. Located
at: 23527179. doi:10.1371/journal.pone.0059388

81. Kumar A, Dalton C, Cortez-Cordova J, Schweizer HP. Mini-Tn7 vectors as genetic tools for
single copy gene cloning in *Acinetobacter baumannii*. J Microbiol Methods. 2010
Sep;82(3):296–300. doi:10.1016/j.mimet.2010.07.002 PubMed PMID: 20638421.

82. Tipton KA, Dimitrova D, Rather PN. Phase-variable control of multiple phenotypes in
*Acinetobacter baumannii* strain AB5075. J Bacteriol. 2015 Aug 1;197(15):2593–9.
doi:10.1128/JB.00188-15 PubMed PMID: 26013481; PubMed Central PMCID:
PMC4518826.

83. Chin CY, Tipton KA, Farokhyfar M, Burd EM, Weiss DS, Rather PN. A high-frequency
phenotypic switch links bacterial virulence and environmental survival in *Acinetobacter
baumannii*. Nat Microbiol. 2018 May;3(5):563–9. doi:10.1038/s41564-018-0151-5 PubMed
PMID: 29693659; PubMed Central PMCID: PMC5921939.

84. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019 Nov 14;20(1):238. doi:10.1186/s13059-019-1832-y

85. Noel HR, Keerthi S, Ren X, Winkelman JD, Troutman JM, Palmer LD. Genetic synergy
between *Acinetobacter baumannii* undecaprenyl phosphate biosynthesis and the Mla
system impacts cell envelope and antimicrobial resistance. mBio. 2024 Feb
16;15(3):e02804-23. doi:10.1128/mbio.02804-23

86. Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for
large alignments. PLOS ONE. 2010 Mar 10;5(3):e9490. doi:10.1371/journal.pone.0009490

87. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. Genome Biol. 2015 Aug 6;16(1):157.
doi:10.1186/s13059-015-0721-2

88. Letunic I, Bork P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display
and annotation. Bioinforma Oxf Engl. 2007 Jan 1;23(1):127–8.
doi:10.1093/bioinformatics/btl529 PubMed PMID: 17050570.

89. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, et al. A
standardized bacterial taxonomy based on genome phylogeny substantially revises the tree
of life. Nat Biotechnol. 2018 Nov;36(10):996–1004. doi:10.1038/nbt.4229

90. Bhamidimarri SP, Zahn M, Prajapati JD, Schleberger C, Söderholm S, Hoover J, et al. A
multidisciplinary approach toward identification of antibiotic scaffolds for *Acinetobacter*
*baumannii*. Structure. 2019 Feb;27(2):268-280.e6. doi:10.1016/j.str.2018.10.021

91. The UniProt Consortium, Bateman A, Martin MJ, Orchard S, Magrane M, Adesina A, et al.
UniProt: the Universal Protein Knowledgebase in 2025. Nucleic Acids Res. 2025 Jan
6;53(D1):D609–17. doi:10.1093/nar/gkae1010

92. Vishniac W, Santer M. The thiobacilli. Bacteriol Rev. 1957 Sep;21(3):195–213.
doi:10.1128/br.21.3.195-213.1957 PubMed PMID: 13471458; PubMed Central PMCID:
PMC180898.

     **FIGURES**

834    **Figure 1.** ***A. baumannii* DcaP proteins cluster in four classes and are widely distributed**
835    **among *Acinetobacter*. (A)** An unrooted tree of DcaP protein sequences from 255
836    *Acinetobacter* genomes (235 *A. baumannii* genomes). **(B)** Manual counts and distribution of
837    individual DcaP proteins binned by the total number of DcaP proteins in an *A. baumannii*
838    genome.
839

**Figure 2.** *A. baumannii* **DcaP-family proteins cluster together compared to non-*Acinetobacter* DcaP proteins. (A)** A rooted tree of DcaP containing species outside of *Acinetobacter* and some *Acinetobacter spp.* depicting class, order, and number of encoded DcaP proteins. **(B)** An unrooted tree of DcaP protein sequences from non-*Acinetobacter* species with AB5075 as the *A. baumannii* reference. Tree scale in amino acid substitutions.

**Figure 3. Wildtype *A. baumannii* grows better than ΔΔΔ*dcaP1-3* mutant on certain di- and**
**tricarboxylic acids as sole carbon sources.** Biolog plates PM1 and PM2a were resuspended
in M9 minimal media without a carbon source and inoculated with either WT or the ΔΔΔ*dcaP1-3*
mutant. (**A**) Difference in area under the curve (ΔAUC) was calculated for WT – ΔΔΔ*dcaP1-3*
mutant. Each point is from one experiment with 1 biological replicate per strain. Significance is
by one sample t-test compared to 0. (**B**) Maximum optical density at 600 nm (OD$_{600}$) is shown.
Significance is by two-way ANOVA with Sidak's multiple comparisons. Experiments were
conducted three times with an n=1 for a total of n=3. Data are mean +/- SD.  *$P$ < 0.05, **$P$ <
0.01, ***P < 0.001, ****P < 0.0001
857

**Figure 4. DcaP3 is important for growth on the carboxylic acid carbon sources citric acid,**
**tricarballylic acid, mucic acid, and *cis*- and *trans*- aconitic acid. (A)** Wildtype, double, and
the triple ΔΔΔ*dcaP1-3* mutants were grown in M9 media with the indicated compound as the
sole carbon source. Growth was monitored by optical density at 600 nm (OD$_{600}$). **(B)** Wildtype,
the ΔΔΔ*dcaP1-3* mutant, and the ΔΔΔ*dcaP1-3* mutant complemented with *dcaP3* were grown on
the indicated compound as the sole carbon source. **(C)** The chemical structures of the relevant
carbon source compounds. **(D)** Expression of the *dcaP* genes in *A. baumannii* 17978 and

AB5075 from (34). Average transcripts per million (Tpm) reads. (A-B) Data are mean +/- SEM
and n=2-3. Experiments were repeated at least twice with similar results.

868
869 **Figure 5. *A. baumannii* AB5075 Δ*dcaP3* is defective for growth on citric acid, tricarballylic**
870 **acid, mucic acid, and *cis*- and *trans*-aconitic acid.** *A. baumannii* AB5075 wildtype (WT) and
871 Δ*dcaP3* mutant strains were grown in M9 media with the indicated compounds as the sole
872 carbon source. Growth was monitored by optical density at 600 nm ($OD_{600}$). Data are mean +/-
873 SEM and n=3. Experiments were repeated twice with similar results.
874

875
**Figure 6. DcaP1, DcaP2, DcaP3, and DcaP4 can complement growth of *A. baumannii***
**ATCC 17978 ΔΔΔ*dcaP1-3* with certain sole carbon sources.** Wildtype, the ΔΔΔ*dcaP1-3*
mutant, and the ΔΔΔ*dcaP1-3* mutant complemented with *dcaP1, dcaP2, dcaP3,* or *A. baumannii*
AB5075 *dcaP4* were grown in M9 media with the indicated compound as the sole carbon
source. Growth was monitored by optical density at 600 nm (OD$_{600}$). Data are mean +/- SEM
and n=3. Experiments were repeated at least twice with similar results.
882

883
**Figure 7. DcaP3 is important for infecting the liver and spleen in a model of bloodstream**
**infection. (A-C)** Wildtype (WT) and the indicated strains were retro-orbitally inoculated into
Swiss Webster mice in a 1:1 ratio. At 24 h post-infection, organs were harvested and bacterial
CFU were enumerated. The dotted lines indicate the limit of detection. Each point represents
one strain enumerated from one mouse; n=9-10. Circle symbols are female mice and square
symbols are male mice. (A-B) Data are combined from two experiments. (C) Data are from one
experiment. Statistics are Wilcoxon non-parametric test. *$P < 0.05$.

Supplementary information for

**DcaP-Family Porins are Required for Carboxylic Acid Utilization and Infection in *Acinetobacter baumannii***

Hannah R. Noel[1], Nikita R. Kesav[1], Jonathan D. Winkelman[2], Lauren D. Palmer[1]

[1] Department of Microbiology and Immunology, University of Illinois Chicago, Chicago, IL, USA
[2] Trestle LLC, Milwaukee, WI, USA

Correspondence: ldpalmer@uic.edu

**This PDF includes:**
Figures S1 to S4
Tables S1 to S3
Supplemental methods

**Figure S1. DcaP proteins are conserved across *Acinetobacter* (A)** *A. baumannii* DcaP proteins clustering with bootstrap values displayed. **(B)** Phylogenetic tree of *Acinetobacter* strains and clades depicting prevalence of individual DcaP proteins. Color coded boxes indicate the presence, absence, or duplication of individual DcaP proteins. Clades within *Acinetobacter* are indicated by the outermost color bar circling the tree. The accession numbers of the data depicted are in Table S4.

**Figure S2.** ***dcaP* genomic neighborhoods and DcaP protein alignments. (A)** The genomic loci of *dcaP1-4* are depicted. **(B)** CLUSTAL Protein alignment of DcaP proteins from ABUW 5075 (1). Secondary structure modeled after PDB entry 6EUS (2).

**Figure S3. Checkerboard assay of ampicillin and sulbactam for wild-type and triple DcaP mutant and growth on glutamic acid and glutaminic acid.**
**(A)** A 96-well plate was filled with stocks of LB with ampicillin and sulbactam and serially diluted to obtain 100 μL of unique concentrations for each compound in each well before inoculation with 1 μL of indicated bacterial strain. Growth was scored as 1, strong turbidity; 0.5, mild turbidity; 0, little to no turbidity. Experiment was repeated 3 times with an n=2 for a total n=6. Means are shown. **(B)** Wildtype and the triple ΔΔΔ*dcaP1-3* mutant were grown in M9 media with the indicated compound as the sole carbon source. Growth was monitored by optical density at 600 nm ($OD_{600}$). Data are mean +/- standard error of the mean (SEM). N=3 and data are representative of at least 2 independent experiments.

**Figure S4. The genetic organization of the *tcu* metabolic locus in *Acinetobacter baylyi* ADP1 and *A. baumannii* ATCC 17978.** Amino acid identity was determined by CLUSTAL Protein alignment using UniProt (1, 3).

**Table S1. Strains**

| Strain | Source | Identifier |
|---|---|---|
| *A. baumannii* ATCC 17978VU (WT) | ATCC | LP486 |
| *A. baumannii* ATCC 17978VU *att*::mTn7(empty); Carb**R** | Ren and Clark *et al.*, 2025 (4) | LP526 |
| *A. baumannii* ATCC 17978VU Δ*dcaP1::kan* (*ACX60_11260*) | This manuscript | LP1191 |
| *A. baumannii* ATCC 17978VU Δ*dcaP2::kan* (*ACX60_10665*) | This manuscript | LP1190 |
| *A. baumannii* ATCC 17978VU Δ*dcaP3::kan* (*ACX60_03745*) | This manuscript | LP1189 |
| *A. baumannii* ATCC 17978 Δ*dcaP1*Δ*dcaP2*::kan | This manuscript | LP1253 |
| *A. baumannii* ATCC 17978VU Δ*dcaP1* Δ*dcaP3* Δ*dcaP2*::kan | This manuscript | LP1308 |
| *A. baumannii* ATCC 17978VU Δ*dcaP3*Δ*dcaP1*::kan | This manuscript | LP1302 |
| *A. baumannii* ATCC 17978VU Δ*dcaP3*Δ*dcaP2*::kan | This manuscript | LP1227 |
| *A. baumannii* ATCC 17978VU Δ*dcaP1* Δ*dcaP3* Δ*dcaP2*::kan *att*::mTn7(*dcaP3p-dcaP3*) | This manuscript | LP1310 |
| *A. baumannii* ATCC 17978VU Δ*dcaP1* Δ*dcaP3* Δ*dcaP2*::kan *att*::mTn7(*dcaP3p-dcaP4(ABUW_2463)*) | This manuscript | LP1314 |
| *A. baumannii* AB5075 | Colin Manoil (University of Washington) (5, 6) | LP345 |
| *A. baumannii* AB5075 Δ*dcaP3*::hyg (*ABUW_RS04045*) | This manuscript | LP1315 |
| *A. baumannii* ATCC 17978VU Δ*dcaP1* Δ*dcaP3* Δ*dcaP2*::kan *att*::mTn7(*dcaP3p-dcaP1*) | This manuscript | LP1368 |
| *A. baumannii* ATCC 17978VU Δ*dcaP1* Δ*dcaP3* Δ*dcaP2*::kan *att*::mTn7(*dcaP3p-dcaP1*) | This manuscript | LP1369 |

**Table S2. Plasmids**

| Use or Phenotype | Source | Name | Identifier |
|---|---|---|---|
| Kan**R**; mobilization helper plasmid | Figurski and Helinski, 1979 (7) | pRK2013 | |
| Carb**R**; plasmid encoding mTn7 transposition pathway | Choi *et al.*, 2005 (8) | pTNS2 | |
| Carb**R**; allelic exchange vector with sucrose sensitivity | Hoang *et al.*, 1998 (9) | pFLP2 | |
| Contains an FRT-flanked kanamycin resistance gene | Datsenko and Wanner, 2000 (10) | pKD4 | |
| Contains a mTn7 with carbenicillin resistance gene | Carruthers *et al.*, 2013 (11) and Alexeyev, 1999 (12) | pKNOCK | |
| Carb**R**; construct to complement *dcaP3* | This manuscript | pKNOCK-*dcaP3* | pLDP343 |
| Carb**R**; express the AB5075 DcaP4 in ATCC 17978 | This manuscript | pKNOCK-*dcaP4* | pLDP359 |
| Carb**R** Kan**R**; deletion construct for DcaP1 | This manuscript | pFLP2-*dcaP1* | pFLP2-11260 |
| Carb**R** Kan**R**; deletion construct for DcaP2 | This manuscript | pFLP2-*dcaP2* | pFLP2-10665 |
| Carb**R** Kan**R**; deletion construct for DcaP3 | This manuscript | pFLP2-*dcaP3* | pFLP2-03745 |
| pKD4 with hygromycin^R replacing kanamycin^R cassette | This manuscript | pKD4-Hyg | pLDP357 |
| Carb**R** Hyg**R**; deletion construct for DcaP3 in AB5075 | This manuscript | pFLP2-5075*dcaP3* | pLDP360 |

| | | | |
|---|---|---|---|
| Carb^R; construct to express *dcaP1* | This manuscript | pKNOCK-*dcaP1* | pLDP401 |
| Carb^R; construct to express *dcaP2* | This manuscript | pKNOCK-*dcaP2* | pLDP402 |

| | | | |
|---|---|---|---|
| Carb[R]; construct to express *dcaP1* | This manuscript | pKNOCK-*dcaP1* | pLDP401 |
| Carb[R]; construct to express *dcaP2* | This manuscript | pKNOCK-*dcaP2* | pLDP402 |

**Table S3. Oligonucleotides**

All oligonucleotides were purchased from IDT (Coralville, IA)

| Sequence | Identifier | Name |
|---|---|---|
| tgaacggcaggtatatgtgatggg | LP52 | pFLP2 fwd |
| ccatgattacgaattcgagc | LP54 | pFLP2 rev |
| gtgtaggctggagctgcttc | LP154 | pKD4-FRTfrag_F |
| catatgaatatcctccttagttcctattc | LP155 | pKD4-FRTfrag_R |
| ttcctttagcagcccttg | LP181 | FRT_Frag_102_R |
| gatctcatgctggagttc | LP183 | FRT_Frag_1384_F |
| cttctctcaaaataaggaacttcg | HN355 | 03745 up up fwd |
| caatccaagtcacttggtaac | HN356 | 03745 dn dn rev |
| aaaaggatcgatcctctagaggatcacatcaccatgcttggg | HN357 | 03745 up fwd |
| acttcgaagcagctccagcctacaccaaagactcctaaaaatatattttatgc | HN358 | 03745 up rev |
| aggaactaaggaggatattcatatgtaacgattttttagctaaaaatagaaaag | HN359 | 03745 dn fwd |
| atgattacgaattcgagctcggtacagttgaaaatatcttgtttagaaagc | HN360 | 03745 dn rev |
| aaaaggatcgatcctctagaggatccgttgtatttaaaattttcatcatg | HN285 | pFLP-11260-up-fwd |
| attctctagaaagtataggaacttctcaaaagctcggtcaaag | HN286 | pFLP-11260-up-rev |
| aggaactaaggaggatattcatatgtcccgtttattttctggtca | HN287 | pFLP-11260-dn-fwd |
| atgattacgaattcgagctcggtacttcaacatgatagcgagttatg | HN288 | pFLP-11260-dn-rev |
| gcaaagaacacatttcacac | HN318 | 11260 up up rev |
| ccagagcaagtatctaatcagac | HN319 | 11260 dn dn fwd |
| ctggtgcagaagatattgaag | HN351 | 10665_dcaP_up up fwd |
| cagacttcttaggtcgtgc | HN352 | 10665 dcaP dn dn rev |
| aaaaggatcgatcctctagaggatccaatgctgaagtagtgatgag | HN353 | pFLP2-10665-up fwd |
| acttcgaagcagctccagcctacactggatataaaagatttccaatatgtgc | HN354 | pFLP2-10665-up rev |
| aggaactaaggaggatattcatatgaatccggttaaaaatattgatttagg | HN363 | pFLP2-10665-dn-fwd |
| atgattacgaattcgagctcggtacggtgtatttctgtagctcgt | HN364 | pFLP2-10665-dn-rev |
| tcatgcatgagctcactagtggatcggttctgtaaaccaaaagt | HN377 | pKNOCK-03745_fwd |
| ggcctgcaaggccttcgcgaggtacttagaaactatatttagccattaaac | HN378 | pKNOCK-03745_rev |
| tcatgcatgagctcactagtggatcgattggatattagcctaaagtct | HN400 | dcaP3 promoter fwd |
| ctgcagctaaaattaatttttcatcaaagactcctaaaaatatattttatgc | HN401 | dcaP3 promoter rev |
| taaaatatattttaggagtctttgatgaaaaaattaattttagctgcag | HN402 | dcaP4 fwd |
| ggcctgcaaggccttcgcgaggtacttagaatttatacatcgatacaaag | HN403 | dcaP4 rev |
| aacttcggaataggaacttcggaaatgtgcgcggaaccccctatttgtttattttt | XR185 | hyg-pKD-F |
| ttcgaaccccagagtcccgctcaggcgccgggggcggtgtccggc | XR186 | hyg-pKD-R |
| aaaaggatcgatcctctagaggatcacatcaccatgcttgg | HN385 | 5075 dcaP3 up fwd |
| acttcgaagcagctccagcctacaccaaagactcctaaaaatatattttatg | HN386 | 5075 dcaP3 up rev |
| aggaactaaggaggatattcatatggaacgattttttagctaaaaatagaaaag | HN387 | 5075 dcaP3 dn fwd |
| atgattacgaattcgagctcggtacaaaagttgaaaatatcttgtttagaaag | HN388 | 5075 dcaP3 dn rev |
| gaacttcgttcttggttaatc | HN389 | 5075 dcaP3 up up fwd |
| ctttacaacccgagttattatc | HN390 | 5075 dcaP3 dn dn rev |

| | | |
|---|---|---|
| tcatgcatgagctcactagtggatcgattggatattagcctaaagtctaa | NK5 | pKNOCK-dcaP3p-dcap2 FWD |
| tatttttcacaggcattccccacatcaaagactcctaaaaatatattttatgc | NK6 | pKNOCK-dcaP3p-dcap2 REV |
| taaaatatattttaggagtctttgatgtgggggaatgcctgtgaaaaata | NK7 | pKNOCK-dcaP3p-dcap2 FWD |
| ggcctgcaaggccttcgcgaggtacttagaaattatattttgctaataaatttatgc gt | NK8 | pKNOCK-dcaP3p-dcap2 REV |
| ccgttgtatttaaaattttcatcatcaaagactcctaaaaatatattttatgc | NK10 | pKNOCK-dcaP3p-dcaP1 REV |
| cataaaatatattttaggagtctttgatgatgaaaatttaaatacaacggt | NK11 | pKNOCK-dcaP3p-dcaP1 FWD |
| ggcctgcaaggccttcgcgaggtacttaaaaatcataactcgctatcatg | NK12 | pKNOCK-dcaP3p-dcaP1 REV |

**Supplementary Methods**

<u>Checkerboard assay</u>

In a 96-well plate, each well was filled with 100 µl LB. Stock solutions of ampicillin (100 mg/mL) and sulbactam (30 mg/mL) were diluted into LB for a final concentration of 1024 µg/mL and 32 µg/mL, respectively. An additional solution of 2048 µg/mL ampicillin was prepared. 100 µl aliquots of 1024 µg/mL ampicillin were placed in row A, wells 1-11, and 2024 µg/mL ampicillin in well 12. 2-fold serial dilutions were performed beginning in row A and ending in row H, discarding the remaining 100 µl. 100 µl aliquots of 32 µg/mL sulbactam were placed in every well in column 12, and 2-fold serial dilutions were performed beginning in column 12 and ending in column 2, discarding the remaining 100 µl. The maximum concentration for each antibiotic was therefore 512 µg/mL ampicillin and 16 µg/mL sulbactam. Then, 1 µl overnight culture of *A. baumannii* ATCC 17978 wild-type or the ΔΔΔ*dcaP* mutant was inoculated into each well before incubation at 37°C with shaking at 180 rpm for 8-16 hours. The following day, plates were scored for growth or no growth phenotypes. Wells with strong optical turbidity were assigned a value of '1' and wells with mild turbidity or mild turbidity with clumped cells were assigned a value of '0.5'. Wells with no visual growth or fully clumped cells were assigned a value of '0.'

**Supplementary References**

1. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol 7:539.

2. Bhamidimarri SP, Zahn M, Prajapati JD, Schleberger C, Söderholm S, Hoover J, West J, Kleinekathöfer U, Bumann D, Winterhalter M, Van Den Berg B. 2019. A multidisciplinary approach toward identification of antibiotic scaffolds for *Acinetobacter baumannii*. Structure 27:268-280.e6.

3. The UniProt Consortium, Bateman A, Martin M-J, Orchard S, Magrane M, Adesina A, Ahmad S, Bowler-Barnett EH, Bye-A-Jee H, Carpentier D, Denny P, Fan J, Garmiri P, Gonzales LJDC, Hussein A, Ignatchenko A, Insana G, Ishtiaq R, Joshi V, Jyothi D, Kandasaamy S, Lock A, Luciani A, Luo J, Lussi Y, Marin JSM, Raposo P, Rice DL, Santos R, Speretta E, Stephenson J, Totoo P, Tyagi N, Urakova N, Vasudev P, Warner K, Wijerathne S, Yu CW-H, Zaru R, Bridge AJ, Aimo L, Argoud-Puy G, Auchincloss AH, Axelsen KB, Bansal P, Baratin D, Batista Neto TM, Blatter M-C, Bolleman JT, Boutet E, Breuza L, Gil BC, Casals-Casas C, Echioukh KC, Coudert E, Cuche B, De Castro E, Estreicher A, Famiglietti ML, Feuermann M, Gasteiger E, Gaudet P, Gehant S, Gerritsen V, Gos A, Gruaz N, Hulo C, Hyka-Nouspikel N, Jungo F, Kerhornou A, Mercier PL, Lieberherr D, Masson P, Morgat A, Paesano S, Pedruzzi I, Pilbout S, Pourcel L, Poux S, Pozzato M, Pruess M, Redaschi N, Rivoire C, Sigrist CJA, Sonesson K, Sundaram S, Sveshnikova A, Wu CH, Arighi CN, Chen C, Chen Y, Huang H, Laiho K, Lehvaslaiho M, McGarvey P, Natale DA, Ross K, Vinayaka CR, Wang Y, Zhang J. 2025. UniProt: the Universal Protein Knowledgebase in 2025. Nucleic Acids Research 53:D609–D617.

4. Ren X, Clark RM, Bansah DA, Varner EN, Tiffany CR, Jaswal K, Geary JH, Todd OA, Winkelman JD, Friedman ES, Jarrett RN, Zemel BS, Wu GD, Zackular JP, DePas WH, Behnsen J, Palmer LD. 2025. Amino acid competition shapes *Acinetobacter baumannii* gut carriage. Cell Host Microbe 33:1396-1411.e9.

5. Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, Gancz HY, Corey BW, Moon JK, Si Y, Owen MT, Hallock JD, Kwak YI, Summers A, Li CZ, Rasko DA, Penwell WF, Honnold CL, Wise MC, Waterman PE, Lesho EP, Stewart RL, Actis LA, Palys TJ, Craft DW, Zurawski DV. 2014. AB5075, a highly virulent isolate of *Acinetobacter baumannii*, as a model strain for the evaluation of pathogenesis and antimicrobial treatments. mBio 5:e01076-14.

6. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse HK, Zurawski DV, Brittnacher MJ, Manoil C. 2015. Resources for genetic and genomic analysis of emerging pathogen *Acinetobacter baumannii*. J Bacteriol 197:2027–35.

7. Figurski DH, Helinski DR. 1979. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. Proc Natl Acad Sci U S A 76:1648–1652.

8. Choi K-H, Gaynor JB, White KG, Lopez C, Bosio CM, Karkhoff-Schweizer RR, Schweizer HP. 2005. A Tn7-based broad-range bacterial cloning and expression system. Nat Methods 2:443–448.

9. Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP. 1998. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. Gene 212:77–86.

10. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640–6645.

11. Carruthers MD, Nicholson PA, Tracy EN, Jr RSM. 2013. *Acinetobacter baumannii* utilizes a type VI secretion system for bacterial competition. PLOS ONE 8:e59388.

12. Alexeyev MF. 1999. The pKNOCK series of broad-host-range mobilizable suicide vectors for gene knockout and targeted DNA insertion into the chromosome of gram-negative bacteria. Biotechniques 26:824–826, 828.