DcaP-Family Porins are Required for Carboxylic Acid Catabolism in *Acinetobacter baumannii*

Hannah R. Noel[1], Jonathan D. Winkelman[2], Lauren D. Palmer[1]*

[1] Department of Microbiology and Immunology, University of Illinois Chicago, Chicago, IL, USA

[2] Trestle LLC, Milwaukee, WI, USA

*Corresponding author

Lauren D. Palmer

835 S Wolcott Ave

MSB E703

Chicago, IL 60612

ldpalmer@uic.edu

ABSTRACT

Acinetobacter baumannii is a pathogen of concern and a leading cause of multidrug-resistant healthcare-associated infections. The *A. baumannii* outer membrane is a barrier to antimicrobials and host defenses but must also allow essential nutrients to permeate. Here, we investigate the functional importance of the putative DcaP-family outer membrane porins, which include a proposed vaccine target in *A. baumannii*. All *A. baumannii* genomes surveyed encode multiple DcaP family porins, which we classify in four classes based on protein sequence phylogeny (DcaP1-4). DcaP proteins encoded by species in other genera could not be mapped to these DcaP classes and phylogenetic analysis suggests DcaP1-4 proteins diversified within *Acinetobacter*. Phenotypic array assays and additional experiments show that DcaP3 was necessary for growth on multiple di- and tri-carboxylic acids as sole carbon sources, including citric acid and tricarballylic acid. Finally, a mutant lacking all DcaP proteins was attenuated in the liver and spleen in a mouse model of bloodstream infection and complemented by expression of DcaP3. These findings provide insight on how *A. baumannii* acquires nutrients through the outer membrane barrier and show DcaP3 is important during infection in specific host niches, validating its potential as a therapeutic target.

INTRODUCTION

Acinetobacter baumannii is a pathogen of critical concern and a major cause of healthcare-associated infections. Isolates of *A. baumannii* are often multidrug resistant (MDR) or extensively drug resistant (XDR) and exhibit resistance to first and last line antibiotics, such as meropenem and colistin, respectively (1). The Centers for Disease Control and the World Health Organization have therefore identified *A. baumannii* as an urgent public health threat, calling for new avenues of treatment (2–4). The bacterial cell envelope is the primary barrier from the environment and protects bacteria from environmental stress including antibiotics and host defenses; however, the cell envelope must also allow acquisition of essential nutrients and cofactors (5,6). The outer membrane (OM) of *A. baumannii* is thought to be less permeable compared to other Gram-negative bacteria such as *E. coli*, contributing to intrinsic antibiotic resistance (7). How small molecule nutrients required for growth transit the impermeable OM in *A. baumannii* is incompletely understood.

The OM of *A. baumannii* is an asymmetric lipid bilayer where the inner leaflet is composed of glycerophospholipids and the outer leaflet of lipooligosaccharides (LOS) (5,8,9). The OM bilayer is occupied by proteins that serve a variety of functions, including nutrient influx. Nutrients typically pass through the OM via active transport by TonB-dependent transporters or passive diffusion through porins which are transmembrane channels that allow passage of small molecules (6,10). The primary porin in *A. baumannii* is OmpA, which shows slow molecular transport in liposome swelling assays, contributing to the low permeability barrier of the OM (11–13). However, the primary function of OmpA is thought to be structural, providing a physical tether between the OM and the peptidoglycan cell wall in *A. baumannii* and other species (14,15). *A. baumannii* encodes multiple other porins including CarO, an OM porin that selectively allows influx of ornithine and other basic amino acids (16), and the Occ family porins, responsible for the influx of compounds like benzoate, hydroxycinnamate, and other aromatic molecules (17–19). The protein DcaP is a predicted OM porin widely distributed across

Acinetobacter species named for its localization within a genetic locus for utilization of

dicarboxylic acids (*dca*) (20). Notably, a DcaP family protein is among the most abundant OM

proteins in *A. baumannii* during rat and mouse infections (21). Due to this fact, multiple groups

have studied a DcaP-like protein as a promising vaccine candidate for protection against *A.*

baumannii infection (21–25). However, the biological role of DcaP family proteins remains

unclear.

First identified in Parke *et al*. as part of the dicarboxylic acid catabolic operon of

Acinetobacter baylyi ADP1, *dcaP* (dicarboxylic acid porin) and DcaP-like proteins were thought

to be specific to the Moraxellaceae family (20). The structure for one DcaP family protein has

been solved, presenting as a homo-trimeric 16-stranded beta-barrel protein with an extended

periplasmic N-terminus that forms a coiled coil (21). Due to the genetic localization and

induction of expression with the dicarboxylic acid adipate, *A. baylyi* DcaP-like proteins were

proposed to be nonspecific porins that facilitate the uptake of dicarboxylic acids (20); however, a

role for DcaP-like proteins in carboxylic acid catabolism has not been reported. Molecular

dynamics and electrophysiology studies determined a DcaP-like protein to have an affinity for

anionic compounds as substrates, such as the clinically relevant β-lactamase inhibitor

sulbactam (21). In other studies, DcaP-like proteins have been implicated in biofilm formation

and mucin catabolism (26,27). Here, we describe four classes of DcaP-like proteins in

Acinetobacter and investigate their role in carbon source utilization.

RESULTS

Acinetobacter spp. encode multiple DcaP-family porins

The first DcaP was described in *A. baylyi* as part of a dicarboxylic acid catabolic operon

(20), and DcaP-like proteins have been identified in various *A. baumannii* clinical isolates and

type-strains (21,23). Proteins designated DcaP-like in *A. baumannii* ATCC 17978 are

homologous to *A. baylyi* DcaP, sharing ~30-35% amino acid identity by Clustal Omega multiple

sequence alignment (28). However, given the abundance of DcaP-like proteins within individual

Acinetobacter genomes, we hypothesized that these proteins could be further classified based

on protein sequence. Using DcaP and DcaP-like sequences from a set of deduplicated *A.*

baumannii genomes, four distinct classes of DcaP-like proteins were identified that cluster

independently, including the original DcaP which we display as DcaP1 (Fig. 1A, S1A). We next

determined the distribution of DcaP-like proteins across the *Acinetobacter* genus. Using a set of

250 deduplicated *Acinetobacter* genomes, most of which were *A. baumannii*, DcaP proteins

were shown to be widely conserved across the genus (Fig. S1B). Additionally, *Acinetobacter*

species within the *A. calcoaceticus/baumannii* (*acb*) complex/clade of pathogenic *Acinetobacter*

appeared to be enriched for having multiple DcaP proteins compared to non-pathogenic

Acinetobacter species (Fig. S1B). Among the *A. baumannii* genomes, over 130 of the strains

encode four *dcaP* genes (Fig. 1B). In fact, no *A. baumannii* genome analyzed encoded fewer

than two *dcaP* genes (Fig. 1B). To determine if the *dcaP* genes were equally distributed across

the species, the number of each DcaP subfamily were tallied for each genome and organized

based on how many total DcaP proteins were present. The four classes of DcaP proteins were

equally likely to be encoded in a given *A. baumannii* genome (Fig. 1B). For example, in most

cases where a genome encodes four DcaP proteins, it was most likely to have one of each

variant. Alternatively, in a genome encoding five DcaP proteins, DcaP2 was most likely to be

duplicated. These data suggest that DcaP proteins are encoded across the *Acinetobacter*

genus, and enriched in pathogenic *Acinetobacter,* including *A. baumannii*.

DcaP proteins from other bacterial species are distinct from Acinetobacter DcaP proteins

To investigate the evolutionary history of DcaP proteins in *Acinetobacter* and determine if

DcaP diversified within *Acinetobacter* or across phyla, orthologues in other bacterial phyla were

identified through KEGG. Bacterial phyla were selected for further analysis if the DcaP-like

protein was encoded in more than half of the genomes for a given genus (29,30). A rooted

phylogenomic tree was generated to demonstrate the evolutionary distance between

Acinetobacter and the bacteria containing a DcaP orthologue (Fig. 2A). Orthologues of DcaP

proteins were found in distantly related organisms, such as *Shewanella spp.* and *Xanthomonas*

spp., albeit generally in fewer numbers (Fig. 2A). These DcaP orthologues could not be further

classified into one of the four DcaP classes defined in *A. baumannii*. This suggests that the four

DcaP classes of proteins defined here are unique to *Acinetobacter spp*. To better understand

the relationship between *Acinetobacter* DcaP proteins and proteins designated DcaP-like from

distantly related species, an unrooted tree of protein sequences was generated. This analysis

revealed that DcaP proteins from AB5075, the included reference *A. baumannii* strain which

encodes DcaP1-4, were more similar to each other than to non-*Acinetobacter* DcaP proteins

(Fig. 2B). This suggests that diversification of the four DcaP classes defined here occurred

within *Acinetobacter*. In contrast to *A. baumannii*, multiple DcaP-like proteins encoded in other

organisms were sequence divergent, such as in *Pseudoxanthomonas daejeonensis*,

Stenotrophomonas spp., *Shewanella psychrotolerans*, and *Marinobacter spp.* (Fig 2B).

Comparison of the species phylogeny and DcaP protein phylogeny suggest there may have

been occurrences of horizontal gene transfer in other species outside *Acinetobacter*. Overall,

these data suggest that DcaP-like proteins are encoded by species across Proteobacteria and

that DcaP diversification has occurred within the *Acinetobacter* clade.

DcaP family porins are necessary for growth on select di- and tri-carboxylic acids

Next, we hypothesized that the DcaP-family porins are involved in small molecule

uptake. One study suggested that DcaP3 may translocate the clinically relevant beta-lactamase

inhibitor sulbactam based on electrophysiology and applied field simulation data (21).

Sulbactam is a therapeutic often used in combination with beta-lactam antibiotics like ampicillin

for bacteria with beta-lactam resistance. Many *A. baumannii* isolates are intrinsically resistant to

ampicillin due to the expression of beta-lactamases and sulbactam has been shown to

resensitize resistant isolates to beta-lactam antibiotics (31,32). We reasoned that if DcaP

proteins facilitated sulbactam entry, a ΔΔΔ*dcaP* mutant would exclude sulbactam from entering

the cell and result increased resistance to the beta-lactam ampicillin while a wild-type strain

would exhibit a re-sensitization to ampicillin. However, the wild-type and the ΔΔΔ*dcaP* mutant

strains exhibited similar sensitivities to ampicillin/sulbactam treatment (Fig. S2). These data

suggest that DcaP proteins are not required for sulbactam activity.

We next hypothesized that DcaP porins may be important for growth on certain nutrient

sources such as dicarboxylic acids as previously suggested (20). The growth of wild-type and

ΔΔΔ*dcaP* strains were compared using the high-throughput Biolog PM system to screen for

carbon source usage. Multiple compounds were identified on which wildtype was able to grow

but ΔΔΔ*dcaP* had a defect, and compounds where ΔΔΔ*dcaP* grew more than wildtype (Fig. 3A-

B). Many of the compounds in which WT grew better than ΔΔΔ*dcaP* were carboxylic acids,

including simple carboxylic acids, hydroxy carboxylic acids, and sugar acids (Fig. 3A-B).

Conversely, compounds in which the mutant outgrew WT were largely amino acids and their

derivatives (Fig. 3A-B). These data suggest that DcaP porins may allow the influx of carboxylic

acid-containing compounds, as previously predicted (20).

DcaP3 is important for growth on multiple carboxylic acid carbon sources

To verify the Biolog results and further determine specificity of the DcaP classes, growth

curves using ΔΔΔ*dcaP,* Δ*dcaP1*Δ*dcaP2*, Δ*dcaP1*Δ*dcaP3*, and Δ*dcaP2*Δ*dcaP3* and the carboxylic

acids as sole carbon sources were performed. Mutants lacking DcaP3 were defective for growth

on citric acid, tricarballylic acid, and mucic acid compared to the wild-type strain (Fig. 4A). We

were unable to identify conditions in which DcaP1 or DcaP2 were required for growth.

Complementation experiments showed that DcaP3 expression was sufficient to restore growth

on these substrates to near wild-type levels (Fig. 4B). Similarly, a single Δ*dcaP3* mutation

conferred a defect in growth on citric acid, tricarballylic acid, and mucic acid, whereas single

mutants of ∆*dcaP1* or ∆*dcaP2* showed no defect (Fig. S3).

A recent study in *A. baylyi* found that tricarballylic acid metabolism is closely linked with

the metabolism of *cis*- and *trans*-aconitic acid (33). The genetic organization of the tricarballylic

acid catabolic genes are conserved in *A. baumannii* compared to *A. baylyi* (Fig. 4C). Since the

data thus far suggest DcaP3 is necessary for growth on tricarballylic acid, we hypothesized that

DcaP3 may also be important for *cis*- and *trans*-aconitic acid growth. Mutants lacking DcaP3

had a defect for growth on both *cis*- and *trans*-aconitic acid (Fig. 4A). Complementing DcaP3 in

the chromosome of the triple mutant ∆∆∆*dcaP* restored growth to near wild-type-like levels,

confirming the role of DcaP3 in growth on these carbon sources (Fig. 4B). Finally, the ∆*dcaP3*

single mutant had a defect in growth on both *cis*- and *trans*-aconitic acid compared to wildtype

(Fig. S3). To highlight the similarities in the carbon sources in which DcaP3 is important for

growth, the chemical structures for each carbon source are outlined in Fig. 4D. Taken together,

these data suggest that DcaP3 is important for growth using multiple di- and tri-carboxylic acids

as sole carbon sources.

DcaP4 complements ∆dcaP3 for growth on carboxylic acids

Thus far, experiments were performed with *A. baumannii* ATCC 17978 which does not

encode DcaP4; thus, the potential role for DcaP4 in growth using carboxylic acids was unclear.

A. baylyi can grow on citric acid and tricarballylic acid and does not encode DcaP3 but rather

DcaP1, DcaP2, and DcaP4 (33). We hypothesized that DcaP4 could be also allow influx of

carboxylic acids. To test this, the DcaP4 from *A. baumannii* AB5075, which shares 77% amino

acid identity to *A. baylyi* DcaP4, was expressed from the 17978 DcaP3 promoter in the

∆∆∆*dcaP A. baumannii* mutant. Consistent with previous observations, the ∆∆∆*dcaP* mutant was

unable to grow using citric acid, tricarballylic acid, mucic acid, or *cis*- and *trans*-aconitic acid

(Fig. 5). Growth was restored to the ΔΔΔ*dcaP* mutant by expressing DcaP4 chromosomally from

the DcaP3 promoter, suggesting that DcaP4 and DcaP3 have overlapping functions in some *A.*

baumannii strains (Fig. 5).

DcaP3 is required for growth on di- and tri- carboxylic acids in a clinical isolate

189   To test if DcaP3 is important for growth on di- and tri-carboxylic acids in a more recent

clinical isolate, a Δ*dcaP3* mutant was generated in the more recent clinical isolate AB5075 and

similar growth curves were performed. Mutants lacking *dcaP3* were defective for growth on citric

acid, tricarballylic acid, mucic acid, and *cis-* and *trans*-aconitic acid compared to wild-type (Fig.

6). This suggests that the requirement for DcaP3 for growth on select carboxylic acids as sole

carbon sources is conserved among *A. baumannii* isolates.

DcaP3 is required for virulence in the liver and spleen during bloodstream infection of mice

197   Carbon source utilization can be a critical determinant of pathogenesis for *A. baumannii*

and other bacterial species in different host niches (34,35). Therefore, we sought to determine if

DcaP function is important for *A. baumannii* pathogenesis. To test this, murine bloodstream

infections were performed, co-inoculating wild-type and the ΔΔΔ*dcaP* mutant or wild-type and

the ΔΔΔ*dcaP* mutant complemented with DcaP3. We hypothesized that there may be a host

niche in which DcaP proteins are important for *A. baumannii* replication, leading to a defect for

ΔΔΔ*dcaP* compared to wildtype. There was no defect of the ΔΔΔ*dcaP* mutant compared to

wildtype in lung, kidney, or heart (Fig. 7A). By contrast, there was a defect for the ΔΔΔ*dcaP*

mutant compared to wildtype in the spleen and liver (Fig. 7A). The defect in the spleen and liver

was reversed by complementing *dcaP3* in the chromosome of the ΔΔΔ*dcaP* mutant, showing

DcaP3 is important for replication in the liver and spleen (Fig. 7B). Together these data suggest

that carboxylic acid permeability by DcaP3 is important in specific host niches during infection.

DISCUSSION

The OM protects *A. baumannii* from antimicrobials and other host stresses but must

permit influx of nutrients essential for replication. How nutrients pass through the *A. baumannii*

OM is not well understood. Here, we designate classes of a family of predicted porins, DcaP1-4,

that are widely distributed across *Acinetobacter spp.* and show DcaP3 or DcaP4 are required for

growth on multiple carboxylic acids.

DcaP-like porins have been identified in genetic screens for biofilm formation,

desiccation, and mucin catabolism, largely referred to as DcaP-like proteins without further

distinction (26,27,36–38). We show that *A. baumannii* DcaP-like proteins can be grouped into

four classes based on protein sequence (Fig. 1A, S1A). The presence of DcaP-encoding genes

is conserved among *Acinetobacter* spp, and particularly in *A. baumannii* and the *acb* complex

(Fig. 1B, Fig. S1B). While DcaP proteins can be found in distantly related organisms (Fig. 2A),

the DcaP1-4 classes we define in *A. baumannii* cluster together (Fig. 2B). These findings

suggest DcaP1-4 diversified within the *Acinetobacter* clade.

Antibiotic resistance in *A. baumannii* is a critical public health problem due to the

emergence of XDR isolates. Various porins in *A. baumannii* have been implicated in the uptake

of toxic molecules including antibiotics (12,39–41). DcaP3 was proposed to allow entry for the

beta-lactamase inhibitor sulbactam (21). Here, we show encoding DcaP-like proteins has no

impact on sensitivity to ampicillin in the presence of sulbactam under the conditions tested (Fig.

S2). These data suggest DcaP-like proteins may not be required for function of sulbactam.

The transport of essential nutrients across the cell envelope is incompletely understood

in *A. baumannii*. In general, *A. baumannii* utilizes organic acids and amino acids as carbon

sources and cannot utilize sugars such as glucose (34). Accordingly, multiple OM channels

have been identified that are important for growth on specific carbon sources including ornithine,

arginine, glutamic acid, and benzoic acid (16–19,42). One study in ATCC 19606 found that

DcaP1 (*A1S_1380*) was upregulated in the presence of mucin, suggesting DcaP1 may play a

role in the influx of mucin components (26). In a Biolog screen, we identified multiple carboxylic

acid carbon sources that supported significantly more growth of the wildtype compared to the

ΔΔΔ*dcaP* mutant (Fig. 3), suggesting DcaP-family proteins are important for their diffusion

through the OM.

The data suggest DcaP3 is responsible for the influx of carboxylic acid carbon sources

citric acid, mucic acid, tricarballylic acid, and *cis*- and *trans*- aconitic acid in *A. baumannii* ATCC

17978, as mutants lacking DcaP3 show decreased growth on these substrates (Fig. 4A-B, S3).

However, the Δ*dcaP3* mutant grew more than the ΔΔΔ*dcaP* mutant in ATCC 17978, suggesting

DcaP1 or DcaP2 may play a minor role in growth with these carbon sources (Fig. S3). Citric

acid is an intermediate of the tricarboxylic acid (TCA) cycle, a ubiquitous pathway for energy

generation. Despite this, many bacteria cannot utilize citric acid as a carbon source and how

citric acid permeates the OM is unknown (43). Mutants in *P. aeruginosa* OpdH were defective

for growth on *cis*-aconitic acid, but not on citric acid, suggesting they encode other citric acid

transporters (44,45). A ΔΔΔ*dcaP* mutant displayed a strong defect for growth on citric acid that

can be rescued by complementing with DcaP3 or DcaP4 (Fig. 4A, 4B, Fig. 5). Therefore,

encoding multiple porins for OM citric acid passage may be conserved across bacterial genera.

Mucic acid, or galactaric acid, is a sugar acid derived from the oxidation of galactose and

is unrelated to mucin. Mucic acid is not typically found in the host; but, some antibiotic

treatments result in increased mucic acid in the human gut (46–48). However, only a subset of

A. baumannii strains, including ATCC 17978, encode the D-galactaric acid dehydrogenase

required for growth on mucic acid (49). The digestive tract is a reservoir for *A. baumannii*

colonization that increases the potential for infection and transmission and *A. baumannii* ATCC

17978 can asymptomatically colonize the mouse gut (50–55). Thus, mucic acid may be used as

carbon source by some *A. baumannii* in the inflamed gut. Our data suggest that DcaP3 or

DcaP4 are important for growth on mucic acid.

The catabolism of *cis*- and *trans*-aconitic acid has been described in *A. baylyi*, and the

catabolism of tricarballylic acid in *A. baylyi* and *Salmonella* (33,56,57). However, the OM porin

required for growth on these carbon sources in *Acinetobacter* is unknown. Our data suggest that

DcaP3 or DcaP4 are important for growth on tricarballylic acid, and *cis*- and *trans*-aconitic acid.

Taken together, our data suggests that DcaP3 and DcaP4 are the major porins for passive

transport of specific carboxylic acid carbon sources.

The requirement for DcaP3 in growth on select di- and tri-carboxylic acids is conserved

in the more recent clinical isolate *A. baumannii* AB5075, as a Δ*dcaP3* mutant has reduced

growth in these conditions (Fig. 6). However, this was surprising as *A. baumannii* AB5075

encodes all four DcaP proteins, including DcaP4, which complemented growth in an *A.*

baumannii ATCC 17978 ΔΔΔ*dcaP* mutant (Fig. 5). We speculate that *dcaP4* may not be

expressed under these conditions.

During infection, bacteria must scavenge nutrients from the host to replicate. The *A.*

baumannii ATCC 17978 ΔΔΔ*dcaP* mutant had a defect compared to wildtype in the mouse liver

and spleen that was complemented by expression of *dcaP3* (Fig. 7A-B). This suggests DcaP3

passive import of carboxylic acid carbon sources is important for pathogenesis in the liver and

spleen. Alternatively, DcaP3 may play another role in pathogenesis. For example, porins in *A.*

baumannii have demonstrated roles for the adhesion or invasion of eukaryotic cells (13,58–61).

DcaP3 is the DcaP-like protein that is a proposed vaccine candidate (21–25). These data

suggest that neutralizing DcaP3 may also promote *A. baumannii* clearance.

In summary, we further classify a family of porins previously termed DcaP-like into four

classes in *Acinetobacter* spp. This classification allows for the differentiation between DcaP-like

proteins and the potential to determine individual functions for DcaP1-4 in *Acinetobacter* spp.

DcaP3 was critically important for growth on citric acid, tricarballylic acid, mucic acid, and *cis*-

and *trans*-aconitic acid as sole carbon sources. A ΔΔΔ*dcaP* mutant was attenuated in a

competitive murine systemic infection and complemented by expression of DcaP3, suggesting a

role for DcaP3 in pathogenesis and validating its potential as a therapeutic target. Our data

therefore provides a deeper understanding of how *A. baumannii* acquires nutrients including in

the host.

MATERIALS AND METHODS

<u>Bacterial strain construction and growth</u>

294   All bacterial strains and plasmids used in this study are listed in Tables S1 and S2,

respectively. The wild-type strain was *A. baumannii* ATCC 17978VU or AB5075 (62).

Oligonucleotides are listed in Table S3 and were purchased from Integrated DNA Technologies

(Coralville, IA). Strains were grown in Luria-Bertani lysogeny broth (LB; Miller) or on LB with

1.5% (wt/vol) agar unless indicated otherwise. Antibiotics were used at the following

concentrations: carbenicillin (Carb), 75 mg/L; kanamycin (Kan), 40 mg/L; chloramphenicol, 15

mg/L; hygromycin (Hyg) 300 mg/L. For deletion constructs, 1000 bp of 3' and 5' flanking DNA to

the gene of interest was amplified using Q5 High Fidelity 2x Master Mix [New England Biolabs

(NEB), Ipswich, MA] from *A. baumannii* ATCC 17978VU or AB5075. The vector pFLP2 was

digested with BamHI and KpnI restriction enzymes (NEB, Ipswitch, MA) and gel purified. The

kanamycin or hygromycin cassette was amplified from pKD4. Fragments were ligated together

using HiFi ligation mix (NEB, Ipswitch, MA), transformed into chemically competent *E. coli*

DH5α, and plated to selective media. Transformants were confirmed to contain the insert by

PCR. To generate the knockout strains, a triparental conjugation using helper *E. coli* strain

HB101 containing pRK2013 was performed. Single-crossover merodiploids containing the

integrated pFLP2 vector were chosen based on a Kan^R Carb^R or Hyg^R and sucrose-sensitive

(10% w/v; Suc^S) phenotype. Putative merodiploids displaying were verified by PCR.

Merodiploids were then grown on LB agar, resuspended, and plated to LB agar or LB agar with

10% sucrose to select for colonies with the second crossover event. Colonies were then

screened for a Kan^R Carb^S Suc^R or Hyg^R Suc^S phenotype. Mutants were confirmed via PCR.

For chromosomal integration constructs, DNA from ATCC 17978VU or AB5075 was

amplified using Q5 High Fidelity 2x Master Mix (NEB, Ipswich, MA). The mTn*7* pKNOCK vector

was digested with BamHI and KpnI prior to ligation using HiFi ligation mix (NEB, Ipswitch, MA)

and transformation into chemically competent *E. coli* DH5α λ*pir116*$^+$. To generate

chromosomally integrated mTn7 constructs, a modified four-way parental mating was performed

using helper *E. coli* strain HB101 containing pRK2013 and *E. coli* DH5α containing pTNS2 (63).

Colonies were screened by PCR for integration into the correct chromosomal locus at the *att*

site. All plasmid constructs were verified by whole plasmid sequencing before transforming into

A. baumannii (Plasmidsaurus, South San Francisco, CA).

<u>Phylogenetic tree construction</u>

To identify DcaP orthologs and classify them into hierarchical orthogroups (HOGs),

OrthoFinder v2.5.5 was used (64). Two datasets were analyzed: (i) a 253-strain dataset

including 235 *A. baumannii* strains deduplicated as previously described (65); (ii) a 28-species

dataset that included multiple *Acinetobacter* and a broader range of gammaproteobacteria

rooted by one alphaproteobacterial. The protein accession numbers and genome accession

numbers are listed in Table S4.

Non-Acinetobacter species were selected based on the conservation of DcaP orthologs

across the genus or species indicated by KEGG. Specific organisms were selected to portray a

representative distribution of DcaP orthologs across bacteria, a group of 22 species. The

FastTree v2.1.11-derived (66), manually rooted species tree was used as an input for

OrthoFinder HOG inference (64,67), ensuring accurate classification of gene duplication and

speciation events. The resulting DcaP classifications were visualized and annotated in iTOL

(68). To infer a species phylogeny, a set of 120 universal bacterial marker genes (Bac120) was

identified from each proteome using HMMER3 (hmmsearch, v3.3), following the methodology

described in Parks *et al.* 2018 (69). The majority of species had only one significant hit for each

of the 120 protein HMMs; if a species had more than one hit, the best-scoring hit for that marker

was extracted.

Each of the 120 marker proteins was independently aligned to its respective HMM profile

using hmmalign (HMMER v3.3). Alignments were trimmed to remove poorly aligned regions and

concatenated into a single concatenated supermatrix for each species. To improve alignment

quality, columns were filtered based on the following criteria: (i) highly gapped regions (>80%

gaps) were removed; (ii) completely homogenous columns were excluded to reduce

redundancy.

Maximum likelihood species trees were inferred from the filtered concatenated alignment

using FastTree v2.1.11, with the LG model of amino acid evolution. The resulting unrooted trees

were manually rooted and annotated in iTOL using *Brevundimonas subvibrioides* or *E.coli* as

outgroups for the smaller and larger datasets.

DcaP protein phylogeny

DcaP homologs were extracted from their respective proteomes by selecting sequences

within the DcaP HOG at the root of all *Acinetobacter* species in the 255-species dataset in

Orthofinder. To ensure consistent domain identification, sequences were aligned to the Pfam

Hidden Markov Model (HMM) PF19577.4, corresponding to the DcaP OM protein family.

Alignments were trimmed to remove poorly aligned residues prior to phylogenetic inference. A

maximum likelihood phylogenetic tree was constructed using FastTree v2.1.11 with the LG

model of amino acid evolution. The inferred clusters were identical to OrthoFinder's DcaP

subfamilies, except for a small cluster of DcaP proteins that OrthoFinder grouped with DcaP1,

while FastTree grouped the same cluster with DcaP3.

To identify putative orthologs of DcaP proteins from the 22 species identified above and

A. baumannii AB5075, proteomes from genomic assemblies were downloaded from the NCBI

Datasets database. Two complementary approaches were employed to verify orthology. First,

HMMER v3.4 was used to search each proteome with the Pfam profile HMM PF19577,

corresponding to the DcaP protein family. Significant hits were retained as candidate DcaP

homologs. To independently assess orthology and reconstruct gene relationships, OrthoFinder

v2.5.5 was run on all 23 proteomes, which also produced a rooted gene tree for the DcaP

family. The resulting DcaP sequences were realigned to the PF19577 HMM using hmmalign,

and poorly aligned or unaligned amino acids were trimmed. A maximum likelihood phylogeny

was then inferred using RAxML v8.2.12, under the LG amino acid substitution model.

Amino acid sequence alignment and percent identity were determined using the Clustal

Omega program (28).

<u>Carbon source utilization screen using Biolog plates</u>

Plates PM1 and PM2A were obtained from Biolog for carbon source utilization testing. A

110 μL aliquot of defined M9 minimal media lacking carbon was dispensed into every well and

resuspended. Then, 100 μL was transferred from the Biolog plates into sterile flat-bottom 96-

well plates. Overnight bacterial cultures were diluted 100-fold in sterile PBS before 1 μL of

diluted culture was inoculated into every well. Plates were incubated at 37°C with shaking and

growth was monitored by optical density at 600 nm (OD_{600}) in an EPOCH2 or SynergyH1 BioTek

plate reader (Winooski, VT). Background absorbance was subtracted before area under the

curve calculations for each isolate and each compound were calculated and the difference

between wildtype and $\Delta\Delta\Delta dcaP$ was analyzed. Compounds were included if any isolate of

either wildtype or $\Delta\Delta\Delta dcaP$ reached an OD_{600} of 0.15 or higher.

Growth curve analyses

M9 minimal medium without glucose was supplemented 0.1X Vishniac's trace minerals (70) with the specified carbon source at a concentration to maintain the carbon equivalency of a 4-carbon compound at 16.5 mM. Cultures were inoculated with a single bacterial colony and grown in 3 mL LB media overnight at 37°C with shaking at 180 rpm for 8-16 hours. Growth curves were performed by diluting overnight cultures 100-fold in sterile PBS and inoculating 1 μL diluted culture into 99 μL media in a 96-well flat bottom plate. Growth was monitored as described above.

Murine bloodstream infection

Six-week-old female ND4 Swiss Webster mice were purchased from Envigo/Harlan Laboratory. Mice were boarded in a temperature-controlled environment with 14:10 h light:dark cycles and food and water were provided as needed. Mice were acclimated to the facility for 1 week prior to infection. Mice were anesthetized with isoflurane and inoculated retro-orbitally with approximately 3×10^8 CFU in a 50 µL bacterial suspension of a 1:1 mixture of *A. baumannii* ATCC 17978VU wildtype and ΔΔΔ*dcaP* mutant or ΔΔΔ*dcaP+dcaP3*. For wildtype versus ΔΔΔ*dcaP,* 5 of the mice were infected with wild-type ATCC 17978VU and 5 were infected with ATCC 17978 mTn7(Carb^R). All mice infected with the complement strain were infected with ATCC 17978VU. The inoculum dose was determined by serial dilution and plating on selective agar media. Mice were euthanized at 24 h post infection by CO_2 asphyxiation, and the organs were excised aseptically. Tissues were homogenized in PBS using a Bullet Blender (Next Advance, Troy, NY), and all samples were serially diluted and plated on LB and LB with carbenicillin or kanamycin selective agar plates for bacterial enumeration. All animal care protocols were approved by the University of Illinois Chicago Institutional Animal Care and Use

Committee (IACUC; protocol number 23–119) in accordance with the Animal Care Policies of

UIC, the Animal Welfare Act, the National Institutes of Health, and the American Veterinary

Medical Association (AVMA). Animals were humanely euthanized consistent with the AVMA

guidelines.

Data reporting, statistical analysis, and figure preparation

Each measurement was taken from a distinct biological sample (e.g., bacterial culture

from a single colony or an individual mouse). Data processing and statistical analyses were

performed using Microsoft Excel 16.98 and GraphPad Prism 10.4.2. Statistical tests used are

indicated in each figure legend. Figures were prepared in Adobe Illustrator 25.2.3.

Author contributions

H.R.N. – project conceptualization, methodology, data collection, data analysis, and writing.

J.D.W – bioinformatics, data analysis, and writing. L.D.P. – project conceptualization,

methodology, data analysis, writing, funding, and resources.

ACKNOWLEDGMENTS

We thank the members of the Palmer laboratory for critical reading of the manuscript and The

Michael Reese Research and Education Foundation Pioneer in Research Award for funding.

REFERENCES

1. Clark NM, Zhanel GG, Lynch JPI. Emergence of antimicrobial resistance among *Acinetobacter* species: a global threat. Curr Opin Crit Care. 2016 Oct;22(5):491.

2. World Health Organization. No time to wait: securing the future from drug-resistant infections. World Health Organization and Interagency Coordination Group on Antimicrobial Resistance; 2019 Apr.

3. Weiner LM, Webb AK, Limbago B, Dudeck MA, Patel J, Kallen AJ, et al. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011–2014. Infect Control Hosp Epidemiol. 2016 Nov;37(11):1288–301.

4. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis. 2018 Mar 1;18(3):318–27.

5. Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. Cold Spring Harb Perspect Biol. 2010 May;2(5):a000414.

6. Nikaido H. Molecular basis of bacterial outer membrane permeability revisited. Microbiol Mol Biol Rev. 2003 Dec;67(4):593–656.

7. Geisinger E, Huo W, Hernandez-Bird J, Isberg RR. *Acinetobacter baumannii*: envelope determinants that control drug resistance, virulence, and surface variability. Annu Rev Microbiol. 2019;73(1):481–506.

8. Kenyon JJ, Hall RM. Variation in the complex carbohydrate biosynthesis loci of *Acinetobacter baumannii* genomes. PLOS ONE. 2013 Apr 16;8(4):e62160.

9. Kenyon JJ, Nigro SJ, Hall RM. Variation in the OC locus of *Acinetobacter baumannii* genomes predicts extensive structural diversity in the lipooligosaccharide. PLOS ONE. 2014 Sep 23;9(9):e107833.

10. Silale A, Berg B van den. TonB-dependent transport across the bacterial outer membrane. Annu Rev Microbiol. 2023 Sep 15;77(Volume 77, 2023):67–88.

11. Sugawara E, Nikaido H. OmpA is the principal nonspecific slow porin of *Acinetobacter baumannii*. J Bacteriol. 2012 Jul 18;194(15):4089–96.

12. Iyer R, Moussa SH, Durand-Réville TF, Tommasi R, Miller A. *Acinetobacter baumannii* OmpA Is a selective antibiotic permeant porin. ACS Infect Dis. 2018 Mar 9;4(3):373–81.

13. Scribano D, Cheri E, Pompilio A, Di Bonaventura G, Belli M, Cristina M, et al. *Acinetobacter baumannii* OmpA-like porins: functional characterization of bacterial physiology, antibiotic-resistance, and virulence. Commun Biol. 2024 Aug 6;7(1):1–11.

14. Samsudin F, Ortiz-Suarez ML, Piggot TJ, Bond PJ, Khalid S. OmpA: a flexible clamp for bacterial cell wall attachment. Structure. 2016 Dec 6;24(12):2227–35.

15. Park JS, Lee WC, Yeo KJ, Ryu KS, Kumarasiri M, Hesek D, et al. Mechanism of anchoring
of OmpA protein to the cell wall peptidoglycan of the gram-negative bacterial outer
membrane. FASEB J. 2012 Jan;26(1):219.

16. Mussi MA, Relling VM, Limansky AS, Viale AM. CarO, an *Acinetobacter baumannii* outer
membrane protein involved in carbapenem resistance, is essential for l-ornithine uptake.
FEBS Lett. 2007;581(29):5573–8.

17. Zahn M, Bhamidimarri SP, Baslé A, Winterhalter M, van den Berg B. Structural insights into
outer membrane permeability of *Acinetobacter baumannii*. Structure. 2016 Feb 2;24(2):221–
31.

18. Clark TJ, Momany C, Neidle EL. The *benPK* operon, proposed to play a role in transport, is
part of a regulon for benzoate catabolism in *Acinetobacter* sp. strain ADP1. Microbiology.
2002;148(4):1213–23.

19. Smith MA, Weaver VB, Young DM, Ornston LN. Genes for chlorogenate and
hydroxycinnamate catabolism (hca) are linked to functionally related genes in the *dca-pca-*
qui-pob-hca chromosomal cluster of *Acinetobacter* sp. strain ADP1. Appl Environ Microbiol.
2003 Jan;69(1):524–32.

20. Parke D, Garcia MA, Ornston LN. Cloning and genetic characterization of *dca* genes
required for β-oxidation of straight-chain dicarboxylic acids in *Acinetobacter* sp. strain ADP1.
Appl Environ Microbiol. 2001 Oct;67(10):4817–27.

21. Bhamidimarri SP, Zahn M, Prajapati JD, Schleberger C, Söderholm S, Hoover J, et al. A
multidisciplinary approach toward identification of antibiotic scaffolds for *Acinetobacter*
baumannii. Structure. 2019 Feb 5;27(2):268-280.e6.

22. Raoufi Z, Abdollahi S, Armand R. DcaP porin and its epitope-based subunit promise
effective vaccines against *Acinetobacter baumannii*; *in-silico* and *in-vivo* approaches. Microb
Pathog. 2022 Jan;162:105346.

23. Fereshteh S, Ajdary S, Sepehr A, Bolourchi N, Barzi SM, Haririzadeh Jouriani F, et al.
Immunization with recombinant DcaP-like protein and AbOmpA revealed protections against
sepsis infection of multi-drug resistant *Acinetobacter baumannii* ST2Pas in a C57BL/6
mouse model. Microb Pathog. 2023 Jan;174:105882.

24. Khalid K, Irum S, Ullah SR, Andleeb S. In-silico vaccine design based on a novel vaccine
candidate against infections caused by *Acinetobacter baumannii*. Int J Pept Res Ther. 2021
Dec 2;28(1):16.

25. Fereshteh S, Abdoli S, Shahcheraghi F, Ajdary S, Nazari M, Badmasti F. New putative
vaccine candidates against *Acinetobacter baumannii* using the reverse vaccinology method.
Microb Pathog. 2020 Jun 1;143:104114.

26. Ohneck EJ, Arivett BA, Fiester SE, Wood CR, Metz ML, Simeone GM, et al. Mucin acts as a
nutrient source and a signal for the differential expression of genes coding for cellular
processes and virulence factors in *Acinetobacter baumannii*. PLoS ONE. 2018 Jan
8;13(1):e0190599.

27. Cabral MP, Soares NC, Aranda J, Parreira JR, Rumbo C, Poza M, et al. Proteomic and
functional analyses reveal a unique lifestyle for *Acinetobacter baumannii* biofilms and a key
role for histidine metabolism. J Proteome Res. 2011 Aug 5;10(8):3399–417.

28. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011
Oct 11;7:539.

29. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res.
2000 Jan 1;28(1):27–30.

30. Kanehisa M, Furumichi M, Sato Y, Matsuura Y, Ishiguro-Watanabe M. KEGG: biological
systems database as a model of the real world. Nucleic Acids Res. 2025 Jan
6;53(D1):D672–7.

31. Ji J, Du X, Chen Y, Fu Y, Wang H, Yu Y. In vitro activity of sulbactam in combination with
imipenem, meropenem, panipenem or cefoperazone against clinical isolates of
Acinetobacter baumannii. Int J Antimicrob Agents. 2013 Apr 1;41(4):400–1.

32. Choi JY, Soo Park Y, Cho CH, Seon Park Y, Shin SY, Song YG, et al. Synergic in-vitro
activity of imipenem and sulbactam against *Acinetobacter baumannii*. Clin Microbiol Infect.
2004 Dec 1;10(12):1098–101.

33. Baugh AC, Defalco JB, Duscent-Maitland CV, Tumen-Velasquez MP, Laniohan NS, Figatner
K, et al. Regulation of tricarboxylate transport and metabolism in *Acinetobacter baylyi* ADP1.
Appl Environ Microbiol. 2024 Jan 30;90(2):e02111-23.

34. Ren X, Palmer LD. *Acinetobacter* metabolism in infection and antimicrobial resistance.
Infect Immun. 2023 May 16;91(6):e00433-22.

35. Barber MF, Fitzgerald JR. Mechanisms of host adaptation by bacterial pathogens. FEMS
Microbiol Rev. 2024 Jul 1;48(4):fuae019.

36. Rumbo-Feal S, Gómez MJ, Gayoso C, Álvarez-Fraga L, Cabral MP, Aransay AM, et al.
Whole transcriptome analysis of *Acinetobacter baumannii* assessed by RNA-sequencing
reveals different mRNA expression profiles in biofilm compared to planktonic cells. PLoS
ONE. 2013 Aug 30;8(8):e72968.

37. Shin JH, Lee HW, Kim SM, Kim J. Proteomic analysis of *Acinetobacter baumannii* in biofilm
and planktonic growth mode. J Microbiol. 2009 Dec 1;47(6):728–35.

38. Gayoso CM, Mateos J, Méndez JA, Fernández-Puente P, Rumbo C, Tomás M, et al.
Molecular mechanisms involved in the response to desiccation stress and persistence in
Acinetobacter baumannii. J Proteome Res. 2014 Feb 7;13(2):460–76.

39. Liu J, Chen S, Zhao B, Li G, Ma T. A novel FadL homolog, AltL, mediates transport of long-
chain alkanes and fatty acids in *Acinetobacter venetianus* RAG-1. Appl Environ Microbiol.
2022 Sep 28;88(20):e01294-22.

40. Mussi MA, Limansky AS, Viale AM. Acquisition of resistance to carbapenems in multidrug-
resistant clinical strains of *Acinetobacter baumannii*: natural insertional inactivation of a

gene encoding a member of a novel family of β-barrel outer membrane proteins. Antimicrob
Agents Chemother. 2005 Apr;49(4):1432–40.

41. Catel-Ferreira M, Coadou G, Molle V, Mugnier P, Nordmann P, Siroy A, et al. Structure-
function relationships of CarO, the carbapenem resistance-associated outer membrane
protein of *Acinetobacter baumannii*. J Antimicrob Chemother. 2011 Sep;66(9):2053–6.

42. Zahn M, D'Agostino T, Eren E, Baslé A, Ceccarelli M, van den Berg B. Small-molecule
transport by CarO, an abundant eight-stranded β-Barrel outer membrane protein from
Acinetobacter baumannii. J Mol Biol. 2015 Jul 17;427(14):2329–39.

43. Blount ZD, Borland CZ, Lenski RE. Historical contingency and the evolution of a key
innovation in an experimental population of Escherichia coli. Proc Natl Acad Sci U S A. 2008
Jun 10;105(23):7899–906.

44. Tamber S, Ochs MM, Hancock REW. Role of the novel OprD family of porins in nutrient
uptake in *Pseudomonas aeruginosa*. J Bacteriol. 2006 Jan;188(1):45–54.

45. Tamber S, Maier E, Benz R, Hancock REW. Characterization of OpdH, a *Pseudomonas*
aeruginosa porin involved in the uptake of tricarboxylates. J Bacteriol. 2007
Feb;189(3):929–39.

46. Hwang I, Park YJ, Kim YR, Kim YN, Ka S, Lee HY, et al. Alteration of gut microbiota by
vancomycin and bacitracin improves insulin resistance via glucagon-like peptide 1 in diet-
induced obesity. FASEB J. 2015;29(6):2397–411.

47. Theriot CM, Koenigsknecht MJ, Carlson PE, Hatton GE, Nelson AM, Li B, et al. Antibiotic-
induced shifts in the mouse gut microbiome and metabolome increase susceptibility to
Clostridium difficile infection. Nat Commun. 2014;5:3114.

48. Faber F, Tran L, Byndloss MX, Lopez CA, Velazquez EM, Kerrinnes T, et al. Host-mediated
sugar oxidation promotes post-antibiotic pathogen expansion. Nature. 2016 Jun
30;534(7609):697–9.

49. Farrugia DN, Elbourne LDH, Hassan KA, Eijkelkamp BA, Tetu SG, Brown MH, et al. The
complete genome and phenome of a community-acquired *Acinetobacter baumannii*. PLOS
ONE. 2013 Mar 19;8(3):e58628.

50. Timsit JF, Garrait V, Misset B, Goldstein FW, Renaud B, Carlet J. The digestive tract is a
major site for *Acinetobacter baumannii* colonization in intensive care unit patients. J Infect
Dis. 1993 Nov 1;168(5):1336–7.

51. Corbella X, Pujol M, Ayats J, Sendra M, Ardanuy C, Domínguez MA, et al. Relevance of
digestive tract colonization in the epidemiology of nosocomial infections due to multiresistant
Acinetobacter baumannii. Clin Infect Dis. 1996 Aug 1;23(2):329–34.

52. Wong SC, Chen JHK, Chau PH, So SYC, AuYeung CHY, Yuen LLH, et al. Gastrointestinal
colonization of carbapenem-resistant *Acinetobacter baumannii*: what is the implication for
infection control? Antibiotics. 2022 Sep 22;11(10):1297.

53. Harris AD, Johnson JK, Pineles L, O'Hara LM, Bonomo RA, Thom KA. Patient-to-patient
transmission of *Acinetobacter baumannii* gastrointestinal colonization in the intensive care
unit. Antimicrob Agents Chemother. 2019 Jul 25;63(8):10.1128/aac.00392-19.

54. Zheng Y, Xu N, Pang J, Han H, Yang H, Qin W, et al. Colonization with extensively drug-
resistant *Acinetobacter baumannii* and prognosis in critically ill patients: an observational
cohort study. Front Med. 2021;8:667776.

55. Ren X, Clark RM, Bansah DA, Varner EN, Tiffany CR, Jaswal K, et al. Amino acid
competition shapes *Acinetobacter baumannii* gut carriage. BioRxiv Prepr Serv Biol. 2024
Oct 19;2024.10.19.619093.

56. Lewis JA, Stamper LW, Escalante-Semerena JC. Regulation of expression of the
tricarballylate utilization operon (*tcuABC*) of *Salmonella enterica*. Res Microbiol. 2009
Apr;160(3):179–86.

57. Lewis JA, Horswill AR, Schwem BE, Escalante-Semerena JC. The tricarballylate utilization
(*tcuRABC*) genes of *Salmonella enterica* serovar Typhimurium LT2. J Bacteriol. 2004
Mar;186(6):1629–37.

58. Gaddy JA, Tomaras AP, Actis LA. The *Acinetobacter baumannii* 19606 OmpA protein plays a
role in biofilm formation on abiotic surfaces and in the interaction of this pathogen with
eukaryotic cells. Infect Immun. 2009 Aug;77(8):3150–60.

59. Choi CH, Lee JS, Lee YC, Park TI, Lee JC. *Acinetobacter baumannii* invades epithelial cells
and outer membrane protein A mediates interactions with epithelial cells. BMC Microbiol.
2008 Dec 10;8:216.

60. Choi CH, Hyun SH, Lee JY, Lee JS, Lee YS, Kim SA, et al. *Acinetobacter baumannii* outer
membrane protein A targets the nucleus and induces cytotoxicity. Cell Microbiol.
2008;10(2):309–19.

61. Labrador-Herrera G, Pérez-Pulido AJ, Álvarez-Marín R, Casimiro-Soriguer CS, Cebrero-
Cangueiro T, Morán-Barrio J, et al. Virulence role of the outer membrane protein CarO in
carbapenem-resistant *Acinetobacter baumannii*. Virulence. 2020 Dec;11(1):1727–37.

62. Wijers CDM, Pham L, Menon S, Boyd KL, Noel HR, Skaar E, et al. Identification of two
variants of *Acinetobacter baumannii* 17978 with distinct genotypes and phenotypes. Infect
Immun. 2021 Aug 30;IAI0045421.

63. Kumar A, Dalton C, Cortez-Cordova J, Schweizer HP. Mini-Tn7 vectors as genetic tools for
single copy gene cloning in *Acinetobacter baumannii*. J Microbiol Methods. 2010
Sep;82(3):296–300.

64. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019 Nov 14;20(1):238.

65. Noel HR, Keerthi S, Ren X, Winkelman JD, Troutman JM, Palmer LD. Genetic synergy
between *Acinetobacter baumannii* undecaprenyl phosphate biosynthesis and the Mla
system impacts cell envelope and antimicrobial resistance. mBio. 2024 Feb
16;15(3):e02804-23.

66. Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for
large alignments. PLOS ONE. 2010 Mar 10;5(3):e9490.

67. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. Genome Biol. 2015 Aug 6;16(1):157.

68. Letunic I, Bork P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display
and annotation. Bioinforma Oxf Engl. 2007 Jan 1;23(1):127–8.

69. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, et al. A
standardized bacterial taxonomy based on genome phylogeny substantially revises the tree
of life. Nat Biotechnol. 2018 Nov;36(10):996–1004.

70. Vishniac W, Santer M. The thiobacilli. Bacteriol Rev. 1957 Sep;21(3):195–213.

FIGURES

636
637   **Figure 1. *A. baumannii* DcaP proteins cluster in four classes and are widely distributed**
638   **among *Acinetobacter*. (A)** An unrooted tree of DcaP protein sequences from 255
639   *Acinetobacter* genomes (235 *A. baumannii* genomes). **(B)** Manual counts and distribution of
640   individual DcaP proteins binned by the total number of DcaP proteins in an *A. baumannii*
641   genome.
642

643
Figure 2. ***A. baumannii* DcaP-family proteins cluster together compared to non-**
***Acinetobacter* DcaP proteins. (A)** A rooted tree of DcaP containing species outside of
Acinetobacter and some *Acinetobacter spp.* depicting class, order, and number of encoded
DcaP proteins. **(B)** An unrooted tree of DcaP protein sequences from non-*Acinetobacter* species
with AB5075 as the *A. baumannii* reference. Tree scale in amino acid substitutions.

Figure 3. Wild-type *A. baumannii* grows better than ∆∆∆*dcaP* mutant on certain di- and tricarboxylic acids as sole carbon sources. Biolog plates PM1 and PM2a were resuspended in M9 minimal media without a carbon source and inoculated with either WT or the ∆∆∆*dcaP* mutant. (**A**) Difference in area under the curve (∆AUC) was calculated for WT – ∆∆∆*dcaP* mutant. Each point is from one experiment with 1 biological replicate per strain. Significance is by one sample t-test compared to 0. (**B**) Maximum optical density at 600 nm (OD_{600}) is shown. Significance is by two-way ANOVA with Sidak's multiple comparisons. Experiments were conducted three times with an n=1 for a total of n=3. Data are mean +/- SD. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$

Figure 4. DcaP3 is important for growth on the carboxylic acid carbon sources citric acid, tricarballylic acid, mucic acid, and *cis*- and *trans*- aconitic acid. (A) Wildtype, double, and the triple ΔΔΔ*dcaP* mutants were grown in M9 media with the indicated compound as the sole carbon source. Optical density at 600 nm (OD$_{600}$) is shown. **(B)** Wildtype, the ΔΔΔ*dcaP* mutant, and the ΔΔΔ*dcaP* mutant complemented with *dcaP3* were grown on the indicated compound as the sole carbon source. **(C)** The genetic organization of the *tcu* metabolic locus and *dcaP3* in *A. baumannii* ATCC 17978. **(D)** The chemical structures of the relevant carbon source compounds. Data are mean +/- SEM and n=3. Experiments were repeated twice with similar results.

670
Figure 5. *A. baumannii* AB5075 DcaP4 complements growth of *A. baumannii* ATCC 17978
Δ*dcaP1-3* on citric acid, tricarballylic acid, mucic acid, and *cis*- and *trans*-aconitic acid.
Wildtype, the ΔΔΔ*dcaP* mutant, and the ΔΔΔ*dcaP* mutant complemented with *dcaP4* were grown
in M9 media with the indicated compound as the sole carbon source. Optical density at 600 nm
(OD_{600}) is shown. Data are mean +/- SEM and n=3. Experiments were repeated twice with
similar results.
677

678
679 **Figure 6. *A. baumannii* AB5075 Δ*dcaP3* is defective for growth on citric acid, tricarballylic**
680 **acid, mucic acid, and *cis*- and *trans*-aconitic acid.** *A. baumannii* AB5075 wild-type and
681 Δ*dcaP3* mutant strains were grown in M9 media with the indicated compound as the sole
682 carbon source. Optical density at 600 nm (OD_{600}) is shown. Data are mean +/- SEM and n=3.
683 Experiments were repeated twice with similar results.
684

685
Figure 7. DcaP3 is important for infecting the liver and spleen in a model of bloodstream infection. (A-B) Wild-type and the indicated strains were retro-orbitally inoculated into Swiss Webster mice in a 1:1 ratio. At 24 h post-infection, organs were harvested and bacterial CFU were enumerated. The dotted lines indicate the limit of detection. Each point represents one strain enumerated from one mouse; n=9-10. Data are combined from two experiments. Statistics are Wilcoxon non-parametric test. *$P < 0.05$.