Full title:

Developmental and Aging Changes in Brain Network Switching Dynamics Revealed by EEG Phase
Synchronization

Short title:

Brain Network Switching Dynamics in Development and Aging

Authors:

Dionysios Perdikis[1,2]*, Rita Sleimen-Malkoun[3]*, Viktor Müller[4], Viktor Jirsa[1]

[1]Aix-Marseille Univ, Inserm, INS, Institut de Neurosciences des Systèmes, Marseille, France

[2]Berlin Institute of Health at Charité, Universitätsmedizin Berlin, Berlin, Germany, Brain Simulation
Section, Department of Neurology with Experimental Neurology, Charité, Universitätsmedizin Berlin,
Corporate member of Freie Universität Berlin and Humboldt Universität zu Berlin, Berlin, Germany

[3]Aix-Marseille Univ, CNRS, ISM, Institut des Sciences du Mouvement, Marseille, France

[4]Max Planck Institute for Human Development, Center for Lifespan Psychology, Berlin, Germany

*These authors contributed equally to this work

Address correspondence to:

Dionysios Perdikis, Berlin Institute of Health, Brain Simulation Section, Department of Neurology with
Experimental Neurology at Charité, Universitätsmedizin Berlin, Robert-Koch-Platz 4, 10115 Berlin,
Germany, dionysios.perdikis@bih-charite.de.

RSM and DP were supported by A*Midex (Aix-Marseille Initiative of Excellence) through project
'Coord-Age' (#ANR-11-IDEX-0001-02).

VJ acknowledges the support of EU's Horizon Europe Programme under the Specific Grant Agreement
No. 101147319 (EBRAINS 2.0 Project), Specific Grant Agreement No. 101137289 (Virtual Brain Twin
Project), the support of Agence Nationale de la Recherche under France 2030, reference No. ANR-24-
RRII-0005 (NAUTILUS). The funders had no role in study design, data collection and analysis, decision
to publish, or preparation of the manuscript.

Abstract

Adaptive behavior depends on the brain's capacity to vary its activity across multiple spatial and temporal scales. Yet, how distinct facets of this variability evolve from childhood to older adulthood remains poorly understood, limiting mechanistic models of neurocognitive aging. Here, we characterize lifespan neural variability using an integrated empirical-computational approach. We analyzed high-density EEG cohort data spanning 111 healthy individuals aged 9–75 years, recorded at rest and during passive and attended auditory oddball stimulation task. We extracted scale-dependent measures of EEG fluctuations amplitude and entropy, together with millisecond-resolved phase-synchrony networks in the 2–20 Hz range. Multi-condition partial least squares decomposition analysis revealed two independent lifespan trajectories. First, slow-frequency power, variance and complexity at longer timescales declined monotonically with age, indicating a progressive dampening of low-frequency fluctuations and large-scale coherence. Second, the temporal organization of phase-synchrony reconfigurations followed an inverted U-trend: young adults exhibited the slowest yet most diverse switching—characterized by low mean but high variance and low kurtosis of jump lengths at 2–6 Hz and the opposite pattern at 8–20 Hz—whereas children and older adults showed faster, more stereotyped dynamics. To mechanistically account for these patterns, we fitted a ten-node phase-oscillator model constrained by the human structural connectome. Only an intermediate, metastable coupling regime reproduced the empirical combination of reduced low-frequency variability and maximally heterogeneous synchrony dynamics observed in young adults, while deviations toward weaker or stronger coupling mimicked the children's and older adults' profiles. Our results demonstrate that development and aging entail changes in the switching dynamics of EEG phase synchronization, by differentially sculpting stationary and transient aspects of neural variability. This establishes time-resolved phase-synchrony metrics as sensitive, mechanistically grounded markers of neurocognitive status across the lifespan.

Author summary [150-200]

Brain activations fluctuate and synchronize according to functionally relevant patterns, both at rest and during task performance. Here we investigated how these fluctuations change from childhood to older adulthood and what do these changes reveal about healthy development and aging. Using electrophysiological brain activity recordings from 111 participants aged 9 to 75 years during rest and a cognitive task, we discovered two key developmental trajectories. First, a linear trend reflecting mainly a gradual weakening of slower brain rhythms with age. Second, an inverted U-shaped trend suggesting that neural flexibility peaks in young adults, who switch between neural synchronization states in a uniquely diverse yet controlled ways, whereas children and older adults showed more rigid and predictable patterns. To explain this, we developed a computational model. The simulated data confirmed that young adults operate in a "sweet spot" of connectivity—balanced enough to support dynamic coordination without becoming unstable or overly rigid—unlike in childhood and older age, where connectivity becomes either too weak or too strong. Our work shows that age fundamentally reshapes how flexibly brain regions communicate. This gives us sensitive markers for tracking brain health across life.

Keywords

Brain dynamics, Development, Aging, EEG, Complexity, Variability, Phase synchronization networks, Functional connectivity, Lifespan, Virtual Brain.

Introduction

Neurodevelopment and aging form a continuum of structural re-wiring and functional reconfiguration that reshapes performance and adaptability from childhood to late adulthood. Yet because childhood, adulthood, and older adulthood are often examined in isolation, an integrated lifespan perspective remains elusive. Rapid advances in structural and functional imaging, high-density electrophysiology and computational modelling now let us probe neural networks with unprecedented spatiotemporal resolution and link their dynamics to behavior, age and functional status [1- 3]. Leveraging these tools, the present study combines resting-state and task-based electroencephalography with generative modelling to trace how core features of brain dynamics evolve across the full spectrum of maturation, adulthood, and aging.

We focus on moment-to-moment variability and synchrony in the EEG because the brain, like any complex multiscale system, exhibits "one-to-many" and "many-to-one" mappings between activity patterns and behavior [4- 6]. The resulting fluctuations are not mere noise; they are stochastic excursions around an underlying deterministic manifold that encode biologically meaningful information [7-9]. As activity unfolds, these excursions trace low-dimensional trajectories—Structured Flows on Manifolds (SFM) [10-12]—whose geometry operates across multiple timescales and underpins distinct cognitive and sensorimotor functions [13]. Development and aging should therefore remodel the shape of these flows in coordination with behavioral change [14]. Independent of age, each individual possesses a finite yet plastic repertoire of neuro-behavioral states, constrained by network architecture and sculpted by experience, that can be flexibly reconfigured to meet environmental demands. Treating variability as a window onto these structured flows provides a unified framework that links bottom-up structural constraints with top-down functional adaptation across the lifespan.

At the structural level, we now understand far more about how the brain wires—and rewires—itself to acquire new skills, maintain function, and remain adaptable across the lifespan [15, 16]. Parallel advances in experimental design, quantitative analysis, and large-scale modelling have turned moment-to-moment neural dynamics into a powerful lens on brain and behavior. Function can be captured as the patterns of activity expressed within and between networks—and across states from rest to task—revealing how circuits integrate, segregate, and reconfigure [17-24]. Whether evoked by a task or emerging spontaneously, these dynamics carry the imprint of each person's neurobehavioral repertoire and their capacity to switch efficiently between functional modes.

Most research tracks development and ageing in separate cohorts [25- 30], with only a handful covering the full lifespan [31- 35]. Theoretical frameworks converge on a noise-driven view in which stochastic perturbations expand the brain's repertoire and push healthy systems toward a metastable "sweet spot" that maximizes integration–segregation balance [36-38]. Empirically, this regime appears as variability that is midway between randomness and rigidity and often predicts higher performance and more adaptable behavior [9, 39-42]. EEG studies exploit its millisecond resolution to capture such dynamics, but scalp signals blur spatial detail; multivariate and multiscale analyses therefore play a key role in disentangling local versus global processing [43-45]. Ultimately, understanding lifespan change requires linking these functional signatures to their structural underpinnings—a relationship that is far from linear given the complex, bidirectional interactions between anatomy and dynamics [46-48].

Although late adulthood is marked by a well-documented decline in white-matter integrity and inter-regional connectivity, lifespan change is not captured by a single linear descent. Instead, structural maturation and senescence follow two archetypal trajectories—one monotonic, the other non-monotonic—whose expression depends on the region or network examined [15, 16, 49]. Normalized gray-matter volume, along with most subcortical nuclei, decreases steadily from childhood onward, whereas the amygdala remains relatively stable, and the hippocampus traces a pronounced inverted U trajectory— maturing later and declining earlier. Douaud and colleagues [16] further identified an inverted-U component that peaks near 40 years and implicates transmodal cortical regions that both

develop and degenerate ahead of the rest of the brain, rendering them especially vulnerable to age-
related pathology. Similar midlife apices have been observed in graph-theoretic metrics of structural
networks [50] and in white-matter tractography [15, 49].

Functional connectivity exhibits the same duality. Linear declines coexist with non-linear, inverted-U
profiles whose timing and magnitude are system-specific [49, 51]. A recent large-scale analysis [52]
located inflection points around mid-life: global mean connectivity peaks near 40 years, while its
variance tops out a decade earlier. Primary sensorimotor circuits reach maturity sooner and begin to
wane earlier than association networks that support complex cognition, which in turn show greater
susceptibility to later-life decline.

Since the two, monotonic and non-monotonic, trends of change are found in different structures and
in interregional functional connectivity of the brain, they may be associated with different control
mechanisms and neural dynamics with implications in different functional domains. Indeed, similar
trends were found in behavioral studies also, although evidence is sparser and more scattered
regarding lifespan changes [53-57]. This is also the case regarding changes in brain activations, that are
more commonly studied separately during early development (infancy-childhood, e.g., [45]),
maturation (infancy/childhood-adulthood, e.g., [58]) and aging (adulthood-old age, e.g. [29]).

Because development and aging are seldom tracked within a single experiment, lifespan inferences
have largely been drawn post-hoc by stitching together separate studies that presume a common
theoretical footing [59]. Only a handful cover childhood through old age in one design [34, 60, 61], and
even these are hard to compare because they deploy heterogeneous metrics—linear or non-linear,
univariate or multivariate—while variously treating EEG as stationary or explicitly time-varying.
Traditional work emphasizes static, linear indices such as power spectra or peak-alpha frequency [62,
63], whereas more recent studies exploit non-linear and time-resolved approaches—fractal
dimension, multiscale entropy, or sliding-window connectivity—to capture the evolving complexity of
brain activity across age [41, 64-66]. This methodological diversity underscores the need for
integrative, cross-sectional analyses that span the full developmental arc and are anchored in
mechanistic models.

Here, we analyze EEG data in four age groups: younger children (*YC*, mean age = 9.9 years), older
children (*OC*, mean age = 12 years), younger adults (*YA*, mean age = 22.7 years), and older adults (*OA*,
mean age = 67.8 years). We capitalize on both resting-state and task-evoked EEG activations to show
that neural variability follows both monotonic decline and inverted-U trajectories, echoing known
structural and connectivity changes. Stationary features are quantified with multiscale entropy and
multivariate statistics, whereas non-stationary features are captured through time-resolved phase-
synchronization dynamics. Finally, we demonstrate that a network of coupled phase oscillators
reproduces these non-stationary signatures, providing a mechanistic framework for understanding
how brain variability is modulated by age—and, by extension, by disease.

Methods

Participants

The studied sample consisted of 24 younger children (*YC*, mean age = 9.9, SD = 0.6, age range = 9.0–
10.8 years, 13 females), 28 older children (*OC*, mean age = 12.0, SD = 0.6, age range = 11.0–12.8 years,
14 females), 31 young adults (*YA*, mean age = 22.7, SD = 1.6, age range = 18.8–25.1 years, 14 females),
and 28 old adults (*OA*, mean age = 67.8, SD = 3.0, age range = 63.9–74.5 years, 14 females).

Ethics Statement

The study has been approved by the ethics committee of Saarland University and has therefore been
performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. All
participants volunteered for this experiment and gave their written informed consent prior to their

inclusion in the study. In the case of children, the parents did give consent for their children to
participate in the study. Details that might disclose the identity of the subjects under study were
omitted.

Table 1 Mean, standard deviation, minimum and maximum numbers of EEG segments per group and
condition included in the analysis

Groups	Conditions	Mean	Standard Deviation	Minimum	Maximum
YC	REC	7.0	1.3	4	8
	UOT	21.2	3.4	14	25
	AOT	22.6	1.9	19	25
OC	REC	7.6	0.8	5	8
	UOT	22.8	2.0	18	25
	AOT	22.9	2.0	18	25
YA	REC	7.8	0.6	5	8
	UOT	24.0	2.0	15	25
	AOT	23.0	3.1	11	25
OA	REC	7.4	1.1	12	25
	UOT	22.3	3.4	12	25
	AOT	21.6	3.1	15	25

Procedure

The EEG measurement began with a 3-minute resting state recording (1.5 minutes with eyes closed,
and 1.5 minutes with eyes open) and was followed by the auditory oddball task. During the task,
participants heard two different tone beeps: a frequent (probability of 0.8) 1000 Hz tone as a standard
stimulus and a rare (probability of 0.2) 800 Hz tone as a deviant stimulus. Stimuli were presented
binaurally at 70 dB SPL with duration of 70 ms (including 10 ms rise and fall time) and inter-stimulus
intervals ranging from 1200 to 1500 ms. There were two different experimental conditions: passive
listening (unattended) and active counting (attended). In the first condition, participants merely
listened to the tone beeps without any response, whereas in the second condition, they had to attend
to stimuli, count the deviant tones and report their number after the end of the session. For this study
we considered three conditions, all with eyes closed: resting state (*REC*, i.e., *Rest* with *Eyes Closed*),
auditory oddball task without counting (*UOT*, i.e., *Unattended Oddball Task*) and auditory oddball with
counting (*AOT*, i.e., *Attended Oddball Task*). The condition of resting state with eyes open was not
included since it differed largely in its frequency content compared to all other conditions, which
interfered with tasks contrasts. Instead, we focused on studying task differences under comparable
conditions along the axis of increasing cognitive and task demands.

EEG recordings and preprocessing

The electroencephalogram (EEG) was recorded with a sampling rate of 500 *Hz* in a frequency band
ranging from 0.5 to 100 *Hz*. The electrodes were placed according to the international 10–10 system.
Data were downsampled to a sampling rate of 250 *Hz*, segmented in artifact free 10 s segments (i.e.,
comprising N_t = 2500 data points each), and mean centered within segments before further analysis.

Table 1 shows the statistics of the resulting number of segments included in the analysis for each
condition and group.

A more detailed description of the participants' cohort, the experimental procedure and the EEG
recording and preprocessing can be found in [33, 34, 41].

We proceed with describing the analysis pipeline leading to the various metrics used in this study. The
pipeline is sketched out in **Fig. 1** for the characteristic example of a data segment of a young adult at
rest with eyes closed.

Univariate variability metrics

The following variability metrics have been applied to all data segments and MATLAB (The Mathworks
Inc.) was used for all computations. We computed Power spectrum (*P*), standard deviation (*σ*),
Multiscale Entropy (*MSE*), and an extended version of the latter (*MSEn*), in an identical way to [41].
MSE is a representative metric for the complexity of a signal across a large range of temporal scales
[67, 68]. The *MSE* algorithm combines the computation of *sample entropy* (*SampEn*) as introduced in
[69] with a coarse graining procedure that acts like downsampling with a Finite Impulse Response (FIR)
low-pass filter, in order to compute this entropy for different timescales. By normalizing with the
standard deviation at each time scale in the case of *MSEn*, we aim at controlling for the confounding
effect of decreasing standard deviation with increasing coarse-graining and, thus, improving the
interpretability of our results. A previous study [41] has shown that this modified metric provides
additional information on EEG variability (see also [70]) for a similar approach). Characteristic examples
of these four metrics for channels Fz, Cz, Pz, and O1 are depicted in **Fig. 1**, *Panel B*.

Phase synchronization metrics

We selected N_{ch} = 19 channels out of the available 58 and applied to all data segments $x(t)$ a
continuous Morlet wavelet transform with cycle number *c*=7:

$$WT(t,f) = \frac{1}{\sqrt{f}} \int x(t) \, A_f e^{-t^2/2\sigma_t^2} e^{i2\pi ft} dt,$$

where $i = \sqrt{-1}$, f is the center frequency that spanned 10 values from 2 to 20 Hz in steps of 2 Hz,
$\sigma_t = c/2\pi f$, and A_f is a normalization constant, chosen such as to retain the signal's amplitude values
for a given time-frequency point. Thus, the wavelet duration was given by $2\sigma_t = c/\pi f$, its wavelength
by $WL = 2\pi\sigma_t$ and its spectral bandwidth by $2\sigma_f$, where $\sigma_f = 1/WL = 1/2\pi\sigma_t = f/c$. Then, we
extracted the phase $\varphi(t,f)$ from the complex $WT(t,f)$ and computed *Circular Correlation Coefficient*
(*CCorr*; see [71]), as a measure of *phase synchronization* (*PS*), among all $N_{con} = N_{ch}*(N_{ch}-1)/2 = 171$ pairs
of channels for each data segment, within a sliding, full-overlapping, window of length 1.004 seconds
or 251 data points. CCR was computed as:

$$CCorr_{i,j}(t,f) = \frac{\sum_{t-125}^{t+125} \sin(\varphi_i(t,f)-\overline{\varphi_i}) \sin(\varphi_j(t,f)-\overline{\varphi_j})}{\sqrt{\sum_{t-125}^{t+125} \sin^2(\varphi_i(t,f)-\overline{\varphi_i}) \sin^2(\varphi_j(t,f)-\overline{\varphi_j})}},$$

where *i, j* denotes here indexes of two different channels, the time variable *t* takes integer values and
corresponds to the center of the sliding window, and $\overline{\varphi_{i,j}}$ denotes the sample mean of the phase within
a sliding window. *CCorr* takes values between 1 (for perfect in phase synchronization) and -1 for perfect
antiphase -180°- synchronization), whether small values around 0 signify orthogonality of the two
signals as a lack of synchronization. Thus, this measure evaluates phase covariance between the two
signals and not phase difference constancy (as for instance phase locking statistics [72]), being a
suitable measure of EEG functional connectivity. **Fig. 1**, *Panel C* depicts characteristic examples of *CCorr*
time series for all combinations of channels Fz, Cz, Pz, and O1 at 10 Hz and *Panel D* for the whole
network of N_{con} connections.

Fig. 1. Data analysis pipeline exemplified. *Panel A:* Time series of four channels (*Fz, Cz, Pz, O1*, out of 58 in total of the international 10–10 system) of a characteristic 10 sec segment of a young adult (*OA*) participant from resting state with eyes closed (*REC*). *Panel B:* Univariate variability metrics for the data of *Panel A* (same channels and colors); *power spectrum* (*P*; top left, x-axis is frequency in Hz), *standard deviation* (*σ*; bottom left, x-axis is time scale in sec), *normalized multiscale entropy* (*MSEn*; bottom left, x-axis is time scale in sec) and *multiscale entropy* (*MSE*; top right, x-axis similar to *MSEn*). *Panel C: Phase synchronization* (*Circular Correlation Coefficient*; *CCorr*) of the data of *Panel A* at 10 Hz, with a sliding, fully overlapping, window of 1.004 sec. Note that there is a rich dynamic leading to a non-stationary (in the wide sense) signal. *Panel D:* Similar to *Panel C* but for 19 channels, which leads to 171 connections (y axis). *Panel E:* Normalized Euclidean norm of the *phase synchronization* networks ($|PS(t,f)|$) at each time point and frequency. Colors correspond to the different frequency bins as explained by the colorbar. The mean ($\mu_{|PS|(f)|}$) and standard deviation ($\sigma_{|PS(f)|}$) for each frequency across time were computed for statistical comparisons. *Panel F: Phase synchronization dynamics* (*PSD*) at 10 Hz computed as the correlation distance among the phase synchronization (*PS*) networks of *Panel D*, at every two distinct time points, leading to a time versus time plot. Note the rich dynamic, according to which there is an alternation of time intervals with small distance (in bluish colors) versus large distance (in reddish colors) changes in the topology of the *PS* networks. The thick dotted black line indicates approximately the time scales (diagonals of the *PSD* matrix) that were chosen for the statistical analysis, in the range 1.004-2 sec. *Panel G:* Considering *PSD* as a random walk in the 171-dimensional space of possible *PS* networks, time sequences of *jump lengths* ($JL(t,\tau)$) for selected time scales τ are plotted, for the indicated time scale range, i.e., the corresponding diagonals of the *PSD* matrix (see the legend for the exact time scales shown in different colors). *Panel H:* Probability distributions of the *JL* sequences ($p_{JL(\tau, 10\ Hz)}$) of *Panel G*, computed with the MATLAB function *ksdensity*() at 25 equidistant points, and normalized with the maximum value, only for demonstration. The sample mean ($\mu_{JL(\tau,f)}$), standard deviation ($\sigma_{JL(\tau,f)}$) and kurtosis ($k_{JL(\tau,f)}$) of such *JL* sequences are computed for further statistical analysis. The x-axis in *Panels A*, *C-G* is the same and corresponds to time. For all phase synchronization graphs (*Panels C-G*), time points correspond to sliding window centers.

Phase synchronization norm. Subsequently, we computed for each time point the normalized Euclidian norm of the phase synchronization across all connections among channels, as:

$$|PS(t,f)| = \frac{1}{N_{con}} \sqrt{\sum_{k}^{N_{con}} CCorr_k^2(t,f)},$$

where the index *k* spans all N_{con} connections (i.e., we treat a synchronization network $PS(t,f)$ at a particular time-frequency point as a vector of N_{con} elements). Then, we computed the mean ($\mu_{|PS(f)|}$) and standard deviation ($\sigma_{|PS(f)|}$) of this quantity across time for each data segment. Thus, we obtained measures of the mean and variability of the norm (magnitude) of the phase synchronization networks across time for each frequency. **Fig. 1**, *Panel E* depicts characteristic *PS* time series for all frequencies.

Phase synchronization dynamics. Next, we evaluated *phase synchronization dynamics* (*PSD*), i.e., the fluctuations of those synchronization networks across time, focusing on the changes only on their topology. For that purpose, we calculated correlation distance among every pair of synchronization networks at time points separated in time by a time scale factor *τ*:

$$PSD(t, t+\tau, f) = 1 - corr\big(PS(t,f), PS(t+\tau, f)\big),$$

where operator *corr*(.) corresponds to the correlation coefficient, which is independent of the magnitude (norm) of $PS(t,f)$. **Fig. 1**, *Panel F* depicts a characteristic example of a PSD matrix at 10 Hz. Treating *PSD* as a random walk in the N_{con}th space of possible synchronization networks (*PS*), we selected time scales in the range $\tau \in [1.004, 2]$ sec, i.e., spanning almost 1 sec of non-overlapping windows, using all possible steps of 0.004 ms. Thus, we defined fluctuations' *jump lengths* $JL(\tau,f)$, as the (correlation) distances *PSD* have moved in *τ* time for all time points *t*, following:

$$JL(\tau, f) = \{PSD(t, t+\tau, f)\}, t \in [0.5\tau + 0.004, 9.5]\ s,$$

i.e., each *JL*(*τ,f*) vector of data points corresponds to the *τ*-th diagonal of the $PSD(t, t+\tau, f)$ matrix. Characteristic *JL*(*τ,f*) vectors for *f* = 10 Hz and a few selected time scales *τ* are plotted across time *t* in **Fig. 1**, *Panel G*, whereas *Panel H* depicts their probability distributions computed with MATLAB

296 function *ksdensity()* at 25 equidistant points, only to demonstrate their long-tailed form. Finally, we
297 computed the mean ($\mu_{JL(\tau,f)}$), the standard deviation ($\sigma_{JL(\tau,f)}$), and the kurtosis $k_{JL(\tau,f)}$ of $JL(\tau,f)$. $\mu_{JL(\tau,f)}$ can
298 be interpreted as the average speed of *PSD*, $\sigma_{JL(\tau,f)}$ evaluates the *PSD* variability, whereas a leptokurtic
299 distribution of $JL(\tau,f)$, as evidenced by a high $k_{JL(\tau,f)}$, characterizes *PSD*, as "switching dynamics", i.e.,
300 dynamics where very short or very long jumps are more likely than in a normal (Gaussian) distribution.

Partial Least Squares (PLS) statistical analysis

Partial Least Squares [73, 74] is a multivariate statistical method that aims at revealing the relationship
between two blocks of datasets, i.e., on the one hand, the metrics computed in this study, and, on the
other hand, some vectors coding for the experimental design. Thus, it is a method suitable for
explorative investigation of spatial and/or time distributed signal changes by combining information
across different signal dimensions. The method is based on a decomposition of the covariance of the
two blocks in a set of new variables that optimally -in a least square sense- relate them, and, therefore,
explain as much of covariance with as few dimensions as possible. In this study, we used mean
centering Task PLS ("mc-Task PLS"), which allows us to explore the contrast of the experimental design
that optimally covaries with our metrics. The method starts by constructing a brain data matrix for
each experimental group. Condition averages are taken across conditions, group matrices are
concatenated, and the grand average is removed (all processing is performed in parallel for each data
element). The resulting matrix undergoes a singular value decomposition. The output of the mc-Task
PLS yields three matrices: i) the *task design latent variables*, i.e., the *saliences of the contrasts*
describing the relations among the conditions and groups of our design for each contrast, ii) the *brain*
latent variables, i.e., *element saliences* that are proportional to the covariance of each data element
with each one of the task contrasts, and iii) *singular values* that are proportional to the covariance
explained by each contrast (the ratio of each squared singular value by the sum of squares of all
singular values of all LVs describes the percentage of covariance each LV accounts for). The number of
resulting singular values, one for each contrast, depends on the degrees of freedom of the design. In
our design we had four groups, namely younger children ('*YC*'), older children ('*OC*'), young adults ('*YA*')
and old adults ('*OA*') participants, and three conditions, i.e., '*REC*', '*UOT*' and '*AOT*' as explained above.
Thus, mc-Task PLS returned, *number of conditions* x *number of groups* -1 = 3*4 – 1 = 11 orthonormal
contrasts. Furthermore, we computed *brain scores* that indicate the strength of the task effect of each
contrast per participant and condition. In other words, the brain score of a particular participant for a
specific contrast and condition is the covariance of the brain data of this participant for that condition
with the corresponding *brain latent variable* vector of the contrast in question. PLS addresses the
problem of multiple comparisons for statistical significance via a permutation test and the problem of
element-wise reliability via a bootstrap resampling test. The permutation test is performed on the
singular values with resampling of the initial data matrices across conditions and groups without
replacement. This permutation test yields a *p*-value for each *task latent variable*, i.e., for each contrast.
We used an alpha value of significance equal to 0.05 throughout this study. For the bootstrap test, the
initial data matrix is resampled with replacement within conditions and groups. For the contrasts we
present here, we plotted the *task latent variables*, together with the mean brain scores and their
intervals of 95% confidence that are derived from the bootstrap test, after mean-centering and
normalizing with the respective singular value. Conditions or groups, the confidence intervals of which
do not overlap, are reliably distinguished by a contrast. For the *brain latent variables*, we calculated
bootstrap ratios by dividing each element with its standard error as computed by the corresponding
bootstrap sample distribution. Bootstrap ratios greater than 2.5758 approximate the 99th two-tailed
percentile for a particular element (see Z-score table).

Simulation

342 We defined a network of $N = 10$ phase oscillators, coupled mutually via a coupling function derived
343 from synaptic couplings [75]. The model system reads

$$344 \qquad \frac{d\theta_i}{dt} = \omega\left(1 - a\cos\theta_i + (1 - \cos\theta_i)\left(\frac{K}{N}\sum_{\substack{j=1,\\j\neq i}}^{N} w_{ij}\cos\theta_j\right) + Q\eta_i(t)\right),$$

where θ_i is the phase state variable of each oscillator i, $\omega = 2\pi f$ with $f = 10$Hz is the central angular
frequency, a is a parameter controlling the stability of the system (stability decreases for increasing a),
K is the global coupling strength parameter, and $w_{ij} = w_{ji} \geq 0$ is the symmetric connectivity weight
between each pair of oscillators in the network, whereas $\eta_i(t)$ is white Gaussian noise with strength
$Q = 0.01$. For the connectivity weights w_{ij} we used a publicly available The Virtual Brain [76] dataset
with values corresponding to numbers of white matter tracts, derived from Diffusion Tensor Imaging
tractography. We selected a sub-circuit of the first $N = 10$ nodes and normalized the connectivity
matrix by dividing them with their 95th percentile to get the resulting weights w_{ij} (see **Fig. S1** in the
Supplementary Information). We performed simulations for different combinations of parameters'
values $a \in \{0.0, 0.1, 0.25, 0.5, 0.75, 0.9, 0.95, 1.0\}$ and $K \in \{0.0, 0.1, 1.0, 2.0, 5.0, 10.0, 100.0\}$. For each
simulation, we integrated the system with the Euler scheme and a time step of dt = 10 ms for 11 sec,
removing the first 1 sec to discard transients. Then we applied to the resulting $\theta_i(t)$ time series a
similar analysis to the EEG data, namely, after downsampling to 250 Hz, we computed the phase
synchronization time series $CCorr_{i,j}(t, 10\,Hz)$ with a sliding, partly-overlapping, window of length 1
sec (250 data points) in steps of 100 ms, and, subsequently, the phase synchronization dynamics
matrices $PSD(t, t + \tau, 10\,Hz)$ with $\tau \in [1, 2]$ sec. PSD matrices and selected time series for
characteristic points of the above two dimensional parameters' space are depicted in **Fig. 10**. Finally,
for the intermediate pair of parameters' values $a = 0.5, K = 2.0$ we clustered the transient phase
synchronization networks based on the PSD matrix as a (correlation) distance metric among time
windows. For this purpose, we used the function *cluster.kmedoids* of the Python3 package *pyclustering*
[77]. The selection of the specific pair of parameters, as well as the number of clusters (three), and the
initial medoids, as inputs to the clustering algorithm, were based on visual inspection of the respective
PSD matrix. Then, we plotted PSD as a 3-dimensional trajectory (**Fig. 10**, *panel C*), for which the
position of each point (corresponding to the phase synchronization network at time window *t*) depicts
the correlation $1 - PSD(t, m_k, 10\,Hz)$ to each one of the clusters medoids' synchronization networks,
where m_k is the medoid of cluster k for $k = \{1,2,3\}$. Note that, given that the self-correlation distance
$PSD(m_k, m_k, 10\,Hz) = 0$, it follows that $1 - PSD(m_k, m_k, 10\,Hz) = 1$.

Results

Univariate variability metrics

The mean curves with standard error intervals of Power spectrum (*P*), standard deviation (σ),
Multiscale Entropy (*MSE*), and an extended version of the latter (*MSEn*) are shown in **Fig. 2**, allowing
for comparisons between age groups under the same task condition. The mc-Task PLS analyses, to
which we turn next, revealed one significant latent variable (LV) for σ and *MSE* and two such variables
for *P* and *MSEn*, according to the permutation test. The first LV for all univariate variability metrics was
significant ($p < 0.001$) and explained most of the covariance (approximately 71.8%, 97.6%, 96.4 and
83.4% for *P*, σ, *MSE* and *MSEn*, respectively). As can be seen in **Fig. 3** through the task LV and
normalized brain scores' graphs, this LV reliably distinguishes children from adults for all four metrics
with negligible condition differences. Statistical trends, although not significant, exist between *YC* and
OC for *MSE* and *YA* and *OA* for the three other metrics (stronger for *MSEn*). The direction of all these
effects is the same both for aging and development, sketching, in general, a monotonic change with
age across lifespan. We observe that adults have generally less power than children (see bootstrap
ratios of the brain LV in **Fig. 4**), an effect that is stronger for fronto-central channels and frequencies
below 10 Hz. For frequencies above 20 Hz, the effect diminishes or even reverses for some fronto-
central channels (e.g., *C6*, *F8*, *F7*), but with statistically unreliable bootstrap ratios. Accordingly, adults
have a lower standard deviation than children for all time scales, the effect being stronger for more

anterior channels and longer time scales. As for the entropic metrics, adults exhibit higher (lower) *MSE* for short (long) time scales than children, the crossing time scale point being around 20 ms, which relates to the notch filter around 50 Hz. This effect is also stronger for fronto-central channels. *MSEn* follows closely the results of *MSE*, but with a few notable differences —beyond the marked aging effect between *YA* and *OA* described above. Specifically, the *MSEn* effect is stronger for central channels, and the crossing point is at longer time scales (around 50 ms for anterior channels and 40 ms for posterior ones). It is also worth noting that normalizing with the magnitude of variability (i.e., σ) weakens the effect of the notch filter in comparison with *MSE*. As for the second LV of the *P* and *MSEn* that was significant ($p < 0.003$ and $p < 0.015$, respectively) and explained approximately 18.9% and 9.7% of the covariance, respectively, it mainly contrasts *OC* and *YA* to *YC* and *OA*, i.e., it sketches an inverted U-shape component with reverse effects for development and aging (see **Fig. 5**). Turning to the corresponding brain LV (**Fig. 6**), *OC* and *YA* exhibit more power and higher *MSEn* for a short range of frequencies (approximately 6-24 Hz) and time scales (20-50 ms), respectively, than *YC* and *OA*. The effect is stronger and wider at more posterior channels for *P*, and at more anterior ones for *MSEn*.

Fig. 2. Group and condition's means and standard error intervals of univariate variability metrics. From top to bottom: *power spectra* (*P*), *standard deviation* (*σ*)), *multiscale entropy* (*MSE*), and *normalized multiscale entropy* (*MSEn*), for channel *Cz*, are shown for all conditions from left to right columns ('*YC*' –magentish colors, '*OC*' –reddish colors, '*YA*' –greenish colors, and '*OA*' –blueish colors) with darker colors for increasing attentional demands ('*REC*', '*UOT*', '*AOT*'). Thick lines and areas of faded colors represent the means and the standard error intervals, respectively. Horizontal axes depict frequency for *P,* and timescale for *σ*, *MSE* and *MSEn*.

Fig. 3. First task latent variable and normalized brain scores of univariate variability metrics: monotonic component. Each panel shows the weights of the first *task latent variable* (asterisks), and the normalized brain scores with 95% confidence intervals (bars and error bars) of the metrics (*P*, top left, *σ* top right, *MSE* bottom left, *MSEn* bottom right). Each bar corresponds to a group and condition combination, with groups being arranged in increasing age from left to right, and conditions arranged in an order of increasing attention demands (i.e., from '*REC*' to '*AOT*'), also from left to right. Color conventions are identical to **Fig. 2**. The name of each metric, together with the corresponding *p*-value (as derived from the permutation test for significance) and the percentage of covariance explained by this latent variable are shown at the title of the respective panel. Non-overlapping confidence intervals signify that conditions and/or groups are separated reliably by the contrast. Thus, we note that for all metrics except for *MSEn*, this latent variable separates mainly adults with children, whereas for *MSEn* young adults are also distinguished being in between children and old adults. In general, these latent variables sketch a monotonic change across lifespan.

Fig. 4. First brain latent variable of univariate variability metrics: monotonic component. The panels show how
much each data *element*, i.e., a metric's data point, covaries with the task contrasts of **Fig. 3** for *P*, *σ*, *MSE* and
MSEn, from left to right, in terms of bootstrap ratios. Absolute values larger than 2.5758 approximate the 99th
two-tailed percentile. The vertical axis for all panels depicts channels arranged from top to bottom, starting from
frontal and left hemisphere channels to occipital and right hemisphere ones. The horizontal axes depict
frequency for *P* and timescale for *σ*, *MSE* and *MSEn*, whereas the titles present information on the statistical
significance and the percentage of covariance explained, similarly to **Fig. 3**. Positive (negative) bootstrap ratios
signify points that exhibit a positive (negative) correlation with the task latent variables of **Fig. 3**, i.e., they
correspond to elements that were higher for adults (children, respectively).

Fig. 5. Second task latent variable and normalized brain scores of univariate variability metrics. Each panel
shows the weights of the second *task latent variable* (asterisks), and the normalized brain scores with 95%
confidence intervals (bars and error bars) of *P* (left) and *MSEn* (right). Color and other conventions are identical
to **Fig. 3**. An inverted U-shape change of reverse effects for development and aging, i.e., contrasting 'YC' and 'OA'
to 'OC' and 'YA', is significant for *P* and *MSEn*.

Phase synchronization metrics

Phase synchronization norm. **Fig. 7** shows the mean values with standard error intervals of the time-
averaged sample *mean* ($\mu_{|PS(f)|}$) and *standard deviation* ($\sigma_{|PS(f)|}$) of the *phase synchronization norm* for
each frequency bin ($|PS(f)|$), plotted across age groups under the same task condition. The respective
mc-Task PLS analyses revealed only one significant LV for each metric ($p < 0.001$) that explained
approximately 95.6% of the covariance for $\mu_{|PS(f)|}$ and 86% for $\sigma_{|PS(f)|}$. The LV for $\mu_{|PS(f)|}$ reliably contrasts
adults to children, with an additional—but statistically unreliable—difference between *YC* and *OC*. In
contrast, the LV for $\sigma_{|PS(f)|}$ contrasts reliably *OA* to *YC*, while OC and YA fall in between, without being
reliably differentiated from either extreme age groups (see **Fig. 8**, top row with task LV and normalized
brain scores). Both contrasts correspond to generally increasing effects across conditions for their
respective metrics, that are statistically reliable across all frequencies for $\mu_{|PS(f)|}$ —particularly for
frequencies up to 12 Hz —and for $\sigma_{|PS(f)|}$ within the 2–12 Hz range, excluding 10 Hz. This frequency-
specific reliability is illustrated by the brain LV bootstrap ratios shown in the bottom row of **Fig. 8**.

P brain LV₂,
18.9154%, p < 0.003

MSEn brain LV₂,
9.6553%, p < 0.015

Frequency (Hz)

Time scale (sec)

Fig. 6. Second brain latent variable of univariate variability metrics: U-shape component. The panels show how much each data *element*, i.e., a metric's data point, covaries with the task contrasts of **Fig. 5** that correspond to the (inverted) U-shape-like change across lifespan for *P* and *MSEn*, in terms of bootstrap ratios. Axes and other conventions are identical to **Fig. 4**. Positive (negative) values signify points with positive (negative) correlation with the task latent variables of **Fig. 5**, i.e., with values that were higher (lower) for '*OC*' and '*YA*' ('*YC*' and '*OA*', respectively).

Phase synchronization dynamics. The situation is quite different, though, for the statistics of *phase synchronization dynamics* (*PSD*), i.e., of the fluctuations in the topology of the *phase synchronization* networks. Figures of means and *t*-scores of the sample *mean* ($\mu_{JL(\tau,f)}$), the *standard deviation* ($\sigma_{JL(\tau,f)}$), and the *kurtosis* ($k_{JL(\tau,f)}$) of the *jump lengths JL(τ,f)* of these fluctuations are shown in **Fig. S2-7** of the Supporting Information. The first LV of the mc-Task PLS analyses was significant ($p < 0.001$) and explained approximately 96.7% of the covariance for $\mu_{JL(\tau,f)}$, 55.7% for $\sigma_{JL(\tau,f)}$, and 83.3% for $k_{JL(\tau,f)}$. This LV reliably contrasts *YA* with children across all three metrics, with *OA'* being somewhere in between; their separation from either *YA* or children is marginal and not statistically reliable (**Fig. 9**, top row). Thus, this component exhibits an inverted U-shaped trajectory across the lifespan—reflecting opposing effects during development and aging. Turning to the brain LV (**Fig. 9**, bottom row), we observe the following age-related patterns: *YA* exhibit lower $\mu_{JL(\tau,f)}$ values across all frequencies (the effect being more pronounced at higher frequencies); *YA* exhibit higher $\sigma_{JL(\tau,f)}$ values across frequencies in the range 2-6 Hz and lower values in 8-20 Hz (reliable only in 14-16 Hz); *YA* exhibit higher $k_{JL(\tau,f)}$ values for frequencies in the range 8-20 Hz and lower values in 2-6 Hz (reliable only in 2-4 Hz). Overall, the *PSD* of *YA* is generally slower, more variable at frequencies below 6 Hz, and more leptokurtic at frequencies above 6 Hz, compared to all other groups—most notably in contrast to children. Finally, we identified that $\sigma_{JL(\tau,f)}$ and $k_{JL(\tau,f)}$ exhibit a second significant LV ($p < 0.003$ for $\sigma_{JL(\tau,f)}$ and $p < 0.006$ for

485 $k_{JL(\tau,f)}$) that explained approximately 33.3% and 6.8% of the covariance, respectively (see **Fig. S8** in the
486 Supporting Information).

Fig. 7. Group and condition's means and standard error intervals of the metrics of the phase synchronization norm. The sample mean $\mu_{|PS(f)|}$ (top row) and standard deviation $\sigma_{|PS(f)|}$ (bottom row) of the *phase synchronization norm* ($|PS(t,f)|$), computed across time for each frequency (x-axes), are shown. Group and condition colors correspond to those in **Fig. 2**. Columns correspond to conditions of increasing attentional demands from left to right ('*REC*', '*UOT*', '*AOT*').

Simulation: Non-stationary synchronization dynamics in a network of coupled phase oscillators

Computational simulations of a network of $N = 10$ coupled phase oscillators, each with a center
frequency of 10 Hz, exhibit slow (approximately on a time scale of 1 sec) intermittent fluctuations in
synchronization (see **Fig. 1**). The synchronization dynamics depends crucially on the degrees of
multistability of the phasing in the network and the stability of the individual attracting states. As the
parameters for stability a and global coupling strength K alter the landscape, the degree of variability
of the emergent synchronization changes systematically. For low a and/or low K, synchronization
dynamics (*PSD*) is highly suppressed as it stays close to the same attractor. For intermediate values,
rapidly changing and fluid, but also persistent *PSD* appears. For high a and/or K, the *PSD* saturates,
exhibiting too fast changes of too small jump length amplitude. In the intermediate regime a scenario
emerges resembling qualitatively the *PSD* patterns as observed empirically in the EEG signals,
displaying rates of synchronization changes on the time scale of seconds. For that scenario, *k*-medoids
clustering of the transient phase synchronization networks using the *PSD* matrix as a (correlation)
distance metric reveals three differentiated clusters of phase synchronization networks, i.e., three
distinct network "states" which persist and reappear during the 10 sec interval of the simulation.
Mapping *PSD* to a 3-dimensional trajectory in a space spanned by correlations of each synchronization
network at time window t to one of the three clusters' medoids, illustrates PSD dynamics as a
stochastic exploration of the space of synchronization network "states".

Fig. 8. First task latent variable, normalized brain scores and brain latent variable of the phase synchronization norm: monotonic component. *Top row:* Each panel shows the weights of the first *task latent variable* (asterisks), and the normalized brain scores with 95% confidence intervals (bars and error bars) *variable* of $\mu_{|PS(f)|}$ (left) and $\sigma_{|PS(f)|}$ (right). Color and other conventions are identical to **Fig. 3**. We observe generally monotonically increasing effects with age that reliably contrast adults to children for $\mu_{|PS(f)|}$, and old adults to young children, with old children and young adults being somewhere in between, for $\sigma_{|PS(f)|}$. *Bottom row:* Each panel shows the bootstrap ratios of the brain latent variable of $\mu_{|PS(f)|}$ (left) and $\sigma_{|PS(f)|}$ (right; note the threshold of the two-tailed 99th percentile with the red dotted line). All bootstrap ratios are positive, i.e., they correspond to elements that correlate positively with contrasts of the top row, i.e., where adults have higher values, for $\mu_{|PS(f)|}$, and where old adults have higher values for $\sigma_{|PS(f)|}$.

Discussion

Changes in brain-signal variability across the lifespan—spanning childhood, adulthood, and older adulthood—illuminate normative development and reveal windows of resilience or vulnerability that may serve as biomarkers of cognitive health or frailty. Yet even the most richly analyzed datasets cannot, on their own, expose the principles that connect neuronal activity to the computations it supports [78]. Deeper insight arises when neuroimaging and advanced analytics are integrated with computational modelling anchored in a mechanistic theory of brain function [12].

Adopting such a lifespan framework, we analyzed resting-state and task-evoked EEG from four age groups (9–75 years) using univariate statistics and time-resolved phase-synchrony metrics. Across both states we found two co-existing age trajectories: a monotonic decline and a pronounced inverted U-shaped trend. Follow-up simulations showed that a network of coupled phase oscillators can reproduce these non-stationary synchrony patterns, offering a mechanistic lens on how neural variability is tuned by development, maturation, and aging. We now detail the implications of these findings.

Fig. 9. First task latent variable, normalized brain scores and brain latent variable of the phase synchronization
dynamics metrics: U-shape component. *Top row:* Each panel shows the weights of the first *task latent* (asterisks),
and the normalized brain scores with 95% confidence intervals (bars and error bars) *variable* of $\mu_{JL(\tau,f)}$ (left), $\sigma_{JL(\tau,f)}$
(center) and $k_{JL(\tau,f)}$ (right). Color and other conventions are identical to **Fig. 3**. We observe for all metrics inverted
U-shape components with age, with reverse effects for development and aging that reliably contrast young
adults to children, with old adults being somewhere in between. *Bottom row:* Each panel shows the bootstrap
ratios of the brain latent variable of $\mu_{JL(\tau,f)}$ (left), $\sigma_{JL(\tau,f)}$ (center) and $k_{JL(\tau,f)}$ (right). Positive (negative) values
correspond to elements that correlate positively (negatively) with contrasts of the top row, i.e., where young
adults (children) have higher (lower) values (respectively).

Fig. 10. Simulations for illustration of the theoretical concepts used to interpret the results. In this figure, we show how a toy model of ten coupled phase oscillators (of central frequency f=10 Hz) can exhibit non-stationary phase synchronization dynamics, which evolves along distinct network "states" on a slow time scale. We present simulations for different values of the parameters of stability a (for values {0.1, 0.5, 1.0}; stability decreases for increasing a), and global coupling scaling K (for values {0.1, 2.0, 10.0}). *Panel A:* From top to bottom, we plot for a=0.5, K=2.0 the simulation time series of (a) the cosine of phase $\vartheta_i(t)$ and (b) the cosine of $\vartheta_i(t) - 2\pi f t$, as well as (c) the phase synchronization ($CCorr_{i,j}(t, 10Hz)$), with a sliding window (of 1 sec, in steps of τ=100 ms). In plots (a-c) t stands for time (points for (a,b), and windows for (c)), and $i, j \in [1, 10]$ are the indices of the phase oscillators for each synchronization index (plotted in distinct colored lines). For the bottom plot (c), warm (cold) colors stand for stronger positive (negative) synchronization (respectively; see also the color bar on the right), whereas rectangular dotted lines depict time intervals, the synchronization networks of which, have been classified to three distinct clusters (see below), each one of a different color (green, dark red, pink; no rectangles are depicted for windows smaller than 200 ms, i.e., two subsequent sliding steps). The middle plot (b), for which the linear central frequency phase increment has been subtracted from the phase time series, serves to reveal the slow modulation of the phases $\vartheta_i(t)$. *Panel B:* We show Phase Synchronization Dynamics (*PSD*) of $CCorr_{i,j}(t, 10Hz)$, for the different values of parameter a (K), changing in the vertical (horizontal) direction (respectively). The color of each pixel of the *PSD(t, t+τ, 10Hz)* images depict the correlation distance between the synchronization networks of two distinct time windows. Therefore, both axes depict time (windows), and each diagonal pixel line away from the central one corresponds to a distinct time scale τ. At the central panel with a=0.5, K=2.0 note the qualitative similarity with the EEG data PSD matrix of **Fig. 1**. The rectangular, colored dotted lines along the central diagonal enclose time windows that correspond to the distinct clusters of synchronization networks. *Panel C:* We demonstrate the synchronization dynamics for a=0.5, K=2.0 as a stochastic exploration of a 3-dimensional space spanned by characteristic networks. We clustered the transient phase synchronization networks with a k-medoids algorithm using the *PSD* matrix as a (correlation) distance metric (see *Methods*). Thus, the positions of the plotted points depict the correlation 1 - *PSD(t, m_k, 10Hz)* of each network at time window t to each one of the three cluster medoids m_k of the synchronization networks. Note that, given that the self-correlation distance *PSD(m_k, m_k, 10Hz)* = 0, we have 1 - *PSD(m_k, m_k, 10Hz)* = 1 for the three cluster medoids.

Brain dynamics as revealed by multiscale EEG fluctuations

Independent of the experimental condition (rest or task), the univariate metrics analyses–Power spectrum (P), standard deviation (σ), Multiscale Entropy (MSE), normalized Multiscale Entropy (MSEn)–brought forward a clear distinction between children and adults, with the suggestion of a monotonic lifespan change when going from the younger to the older age. This included a decrease in the power and in the amplitude of fluctuations for slow frequencies (below 10 Hz) and longer time scales, respectively. These effects were also reflected in reduced EEG fluctuations' complexity with age resulting from significantly lower entropy at longer time scales that mirror slower processes (> 20 ms for MSE; >50 ms for MSEn) in the adults compared to the children's groups. These changes were most pronounced at frontal electrode sites. Interestingly, for two of the studied metrics, *P* and *MSEn*, the dominant monotonic trend co-existed with a non-linear trend (inverted U-shaped) opposing older children and younger adults to younger children and older adults. The inverted U trend concerned an intermediate range of frequencies (6-24 Hz) and time scales (20-50 ms), with more power and higher entropy for the intermediate age groups compared to the youngest children's and the oldest adults' groups. This result was prevailing at the posterior channels for P and the fronto-central channels for MSEn.

The observed decrease in slow-frequency power (below 10 Hz) and the reduced complexity at longer timescales align with previously reported developmental EEG patterns, with delta and theta power declining from childhood to adulthood, along with brain maturation processes [33, 79]. The frontal predominance of these changes is particularly consistent with the late development of the prefrontal cortex, which continues maturing well into the third decade of life. The finding of decreased multiscale entropy at longer timescales (>20-50ms) in adults versus children completes previous findings that focused on shorter timescales [28, 59] and concluded an increase in brain signal variability with development. Here, we show that EEG variability in terms of magnitude (σ) and complex time-

structure is lower in children than in adults. The link could be made with a possible shift in how the brain processes information (less locally and more distributed) with age [45].

A noteworthy feature of our data is that monotonic decline and inverted U-shaped trajectories coexist within the same EEG recordings. Such duality has been predicted by theory and documented with MRI connectivity but rarely verified in lifespan EEG. Here, the inverted U pattern emerges at intermediate frequencies (≈ 6–24 Hz) and timescales of 20–50 ms, hinting that specific network operations reach optimal efficiency in young adulthood before tapering with age. One plausible contributor is the transient surge in posterior beta activity observed during cognitive maturation, whose scalp distribution overlaps the classic posterior-dominant alpha topography [80].

Brain dynamics as revealed by EEG phase synchronization dynamics

The analysis of the EEG synchronization dynamics afforded a richer account of the lifespan changes as reflected by the differences between the studied age groups. We found first that, across time and for each frequency, the mean and the variability of the magnitude of the phase synchronization networks increased with age (cf. [35]). This was the case for all frequencies, but more reliably so for the lower ones (<10 Hz). The mean of the phase synchronization norm contrasted between children and adults, while its standard deviation opposed young children to older adults, with old children and young adults being undistinguishable between the two former groups. As the above-discussed univariate metrics, these two descriptive bivariate metrics of phase synchronization point out a monotonic change in brain dynamics, with a stronger and more variable functional (phase synchronization) connectivity when going from early childhood to aging.

Moving to the analysis of phase synchronization dynamics (PSD), we could extract a different trajectory of how age affects the functioning of the brain by altering the topology of the phase synchronization networks following a non-linear inverted U-like trend opposing development to aging. Independently of the task condition, younger adults were reliably distinguished from all the three other age groups for all the jump length distribution parameters. Namely, they showed a lower mean jump length for all frequencies, suggesting a slower switching dynamic compared to the children and older adults' groups. They were also found to be more variable and less leptokurtic in their switching dynamics than the other groups for low frequency bins (2-6 Hz), and the reverse for the higher frequencies (8-20 Hz). Accordingly, we can conclude that development and aging modulate the PSD in opposite directions modifying thereby the expression of the transient fluctuations of the functional (phase synchronization) networks. This discriminating transient dynamic can be captured best through measures that do not rely on or assume stationarity of the brain's functional connectivity. Our results are highly consistent with [61], reinforcing the inverted U-shaped model of EEG phase synchronization across the lifespan and the value of dynamic, non-stationary analysis. Our additional focus on the detailed transient switching dynamics (PSD) provides a complementary perspective, offering finer granularity on how development and aging shape the flexibility and variability of brain network connectivity. The finding that younger adults show distinct switching dynamics –slower, more variable switching at low frequencies but less variable at higher frequencies– suggests an optimal dynamic regime in young adulthood. This aligns with theoretical frameworks proposing that the mature healthy brain operates in a metastable state balancing integration and segregation [35, 38, 81].

Modeling the synchronization dynamics in a network of coupled phase oscillators

Earlier work has shown that large-scale "cascades" of spontaneous activity—bursts that sweep through distributed networks—are detectable in both EEG and fMRI and reorganize systematically with age [9, 14, 81-84]. These phenomena have been reproduced in whole-brain simulations, confirmed in animal models, and observed with simultaneous multimodal recordings. Cascades tend to cluster during epochs when the underlying dynamics are quasi-stationary, implying that they both signal and stabilize favorable network states.

We capture an analogous signature here, albeit in a deliberately pared-down form: our analyses focus exclusively on phase relationships. Excluding amplitude inevitably misses waxing-and-waning envelope

modulations, but a band-limited, phase-only approach is standard in EEG research and maps cleanly onto networks of Kuramoto oscillators. Within this framework, cascades appear as brief windows of heightened inter-regional coherence—transient "tiles'' in the temporal mosaic of brain activity.

fMRI studies link richly cascaded, or "fluid," dynamics to robust cognitive performance; this fluidity wanes with advancing age and correlates with declining cognition [14]. Our EEG findings echo that pattern, showing that time-resolved phase synchrony provides an accessible proxy for the same health-related dynamical regime.

Conclusion

Our results show that both the amplitude of EEG fluctuations and the tempo of phase synchrony change with age along two distinct trajectories. Aggregate, "static" indices—means and standard deviations—decline roughly linearly from childhood to older adulthood, whereas transient, time-resolved metrics of network switching follow a pronounced inverted U-shaped trajectory, peaking in young adulthood. Detailed jump-length analyses (mean, variance, kurtosis across frequencies) reveal that young adults execute slower, more selective transitions between functional states, consistent with an optimally tuned balance between stability and flexibility. Children appear to be learning this control regime, while older adults show a drift toward faster, less discriminating switches, suggesting a loss of the optimized switching dynamics established at the earlier stage of life.

These electrophysiological patterns mirror the dual monotonic and inverted U trends long reported in structural and functional MRI, implying a common mechanistic underpinning. Within the framework of Structured Flows on Manifolds, our findings link micro-scale neural dynamics to macro-scale network architecture, providing a coherent view across imaging modalities and timescales. A critical next step is to extend this approach to simultaneous behavioral recordings, allowing us to trace how brain and behavior co-evolve across the human lifespan.

References

1. Loosen AM, Kato A, Gu X. Revisiting the role of computational neuroimaging in the era of integrative neuroscience. Neuropsychopharmacology. 2025;50(1):103–113. doi:10.1038/s41386-024-01946-8.

2. Steinmetz NA, Koch C, Harris KD, Carandini M. Challenges and opportunities for large-scale electrophysiology with Neuropixels probes. Curr Opin Neurobiol. 2018 Jun;50:92–100. doi:10.1016/j.conb.2018.01.009. Epub 2018 Feb 13. PMID: 29444488; PMCID: PMC5999351.

3. Sun L, Zhao T, Liang X, Xia M, Li Q, Liao X, Gong G, Wang Q, Pang C, Yu Q, Bi Y, Chen P, Chen R, Chen Y, Chen T, Cheng J, Cheng Y, Cui Z, Dai Z, Deng Y, Ding Y, Dong Q, Duan D, Gao JH, Gong Q, Han Y, Han Z, Huang CC, Huang R, Huo R, Li L, Lin CP, Lin Q, Liu B, Liu C, Liu N, Liu Y, Liu Y, Lu J, Ma L, Men W, Qin S, Qiu J, Qiu S, Si T, Tan S, Tang Y, Tao S, Wang D, Wang F, Wang J, Wang P, Wang X, Wang Y, Wei D, Wu Y, Xie P, Xu X, Xu Y, Xu Z, Yang L, Yuan H, Zeng Z, Zhang H, Zhang X, Zhao G, Zheng Y, Zhong S; Alzheimer's Disease Neuroimaging Initiative; DIDA-MDD Working Group; MCADI; He Y. Human lifespan changes in the brain's functional connectome. Nat Neurosci. 2025 Apr;28(4):891–901. doi:10.1038/s41593-025-01907-4. Epub 2025 Apr 3. PMID:40181189.

4. Changeux J-P, Courrège P, Danchin A. A theory of the epigenesis of neuronal networks by selective stabilization of synapses. Proc Natl Acad Sci U S A. 1973 Oct;70(10):2974–2978. doi: 10.1073/pnas.70.10.2974.

5. Edelman GM, Gally JA. Degeneracy and complexity in biological systems. Proc Natl Acad Sci U S A. 2001;98(24):13763–13768. https://doi.org/10.1073/pnas.231499798.

6. Breakspear M, Jirsa VK. Neuronal dynamics and brain connectivity. In: Jirsa VK, McIntosh AR, editors. *Handbook of Brain Connectivity*. Berlin: Springer; 2007. p. 110–130.

7. Waschke L, Kloosterman NA, Obleser J, Garrett DD. Behavior needs neural variability. Neuron. 2021 Mar 3;109(5):751–766. doi:10.1016/j.neuron.2021.01.023.

8. Zilles K, Amunts K. Individual variability is not noise. Trends Cogn Sci. 2013 Apr;17(4):153–155. doi: 10.1016/j.tics.2013.02.003.

9. Fousek J, Rabuffo G, Gudibanda K, Sheheitli H, Petkoski S, Jirsa V, et al. Symmetry breaking organizes the brain's resting state manifold. *Sci Rep*. 2024;14:31970.

10. Huys R, Perdikis D, Jirsa VK. Functional architectures and structured flows on manifolds: A dynamical framework for motor behavior. Psychol Rev. 2014 Oct;121(3):302–336. doi: 10.1037/a0037014.

11. Pillai AS, Jirsa VK. Symmetry breaking in space-time hierarchies shapes brain dynamics and behavior. Neuron. 2017 Jun 7;94(5):1010–1026. doi: 10.1016/j.neuron.2017.05.013.

12. Jirsa VK, Sheheitli H. Entropy, free energy, symmetry and dynamics in the brain. J Phys Complex. 2022;3(1):015007. doi:10.1088/2632-072X/ac4bec.

13. Brennan C, Proekt A. A quantitative model of conserved macroscopic dynamics predicts future motor commands. eLife. 2019 Jul 11;8:e46814. doi: 10.7554/eLife.46814.

14. Lavanga M, Stumme J, Yalcinkaya BH, Fousek J, Jockwitz C, Sheheitli H, Bittner N, Hashemi M, Petkoski S, Caspers S, Jirsa V. The virtual aging brain: Causal inference supports interhemispheric dedifferentiation in healthy aging. NeuroImage. 2023 Dec 1;283:120403. doi:10.1016/j.neuroimage.2023.120403. Epub 2023 Oct 20. PMID: 37865260.

15. Coupé P, Catheline G, Lanuza E, Manjón JV. Towards a unified analysis of brain maturation and aging across the entire lifespan: A MRI analysis. Hum Brain Mapp. 2019;40(1):215–229. doi: 10.1002/hbm.23743.

16. Douaud G, Groves AR, Tamnes CK, Westlye LT, Duff EP, Engvig A, Walhovd KB, James A, Gass A, Monsch AU, Matthews PM, Fjell AM, Smith SM, Johansen-Berg H. A common brain network links development, aging, and vulnerability to disease. Proc Natl Acad Sci U S A. 2014 Dec 9;111(49):17648–17653. doi: 10.1073/pnas.1410378111.

17. Deco G, Jirsa VK, McIntosh AR. Resting brains never rest: computational insights into potential cognitive architectures. Trends Neurosci. 2013 May;36(5):268–274. doi: 10.1016/j.tins.2013.03.001.

18. Florin E, Baillet S. The brain's resting-state activity is shaped by synchronized cross-frequency coupling of neural oscillations. Neuroimage. 2015 Apr 15;120:85–94. doi: 10.1016/j.neuroimage.2015.01.054.

19. Garrett DD, Samanez-Larkin GR, MacDonald SWS, Lindenberger U, McIntosh AR, Grady CL. Moment-to-moment brain signal variability: a next frontier in human brain mapping? Neurosci Biobehav Rev. 2013 May;37(4):610–624. doi:10.1016/j.neubiorev.2013.02.015.

20. Hansen ECA, Battaglia D, Spiegler A, Deco G, Jirsa VK. Functional connectivity dynamics: modeling the switching behavior of the resting state. NeuroImage. 2015 Feb 1;105:525–535. doi:10.1016/j.neuroimage.2014.11.001.

21. Hutchison RM, Morton JB. Tracking the brain's functional coupling dynamics over development. J Neurosci. 2015 May 6;35(17):6849–6859. doi:10.1523/JNEUROSCI.4638-14.2015.

22. Monti JM, Sanz Perl Y, Tagliazucchi E, Kringelbach ML, Deco G. Fluctuation-dissipation theorem and the discovery of distinctive off-equilibrium signatures of brain states. Phys Rev Research. 2025 Mar 21;7(1):013301. doi:10.1103/PhysRevResearch.7.013301.

23. Schlesinger KJ, Turner BO, Lopez BA, Miller MB, Carlson JM. Age-dependent changes in task-based modular organization of the human brain. NeuroImage. 2017 Feb 1;146:741–762. doi: 10.1016/j.neuroimage.2016.09.001.

24. Shine JM, Poldrack RA. Principles of dynamic network reconfiguration across diverse brain states. NeuroImage. 2018 Oct 15;180(Pt B):396–405. doi: 10.1016/j.neuroimage.2017.08.010.

25. Heisz JJ, Gould M, McIntosh AR. Age-related shift in neural complexity related to task performance and physical activity. J Cogn Neurosci. 2015 Mar;27(3):605–613. doi:10.1162/jocn_a_00725.

26. Koenis MMG, Brouwer RM, van den Heuvel MP, Mandl RCW, van Soelen ILC, Kahn RS, Boomsma DI, Hulshoff Pol HE. Development of the brain's structural network efficiency in early adolescence: A longitudinal DTI twin study. Hum Brain Mapp. 2015 Mar;36(3):1088–1099. doi:10.1002/hbm.22988.

27. Manor B, Costa MD, Hu K, Newton E, Starobinets O, Kang HG, et al. Physiological complexity and system adaptability: evidence from postural control dynamics of older adults. J Appl Physiol. 2010;109(6):1786–91.

28. McIntosh AR, Kovacevic N, Itier RJ. Increased brain signal variability accompanies lower behavioral variability in development. PLoS Comput Biol. 2008 Jul 4;4(7):e1000106. doi:10.1371/journal.pcbi.1000106.

29. McIntosh AR, Vakorin V, Kovacevic N, Wang H, Diaconescu A, Protzner AB. Spatiotemporal dependency of age-related changes in brain signal variability. Cereb Cortex. 2014 Jul;24(7):1806–1817. doi:10.1093/cercor/bht030.

30. Wu K, Taki Y, Sato K, Hashizume H, Sassa Y, Takeuchi H, Thyreau B, He Y, Evans AC, Li X, Kawashima R, Fukuda H. Topological organization of functional brain networks in healthy children: Differences in relation to age, sex, and intelligence. PLoS ONE. 2013 Feb 4;8(2):e55347. doi:10.1371/journal.pone.0055347.

31. Davison EN, Turner BO, Schlesinger KJ, Miller MB, Grafton ST, Bassett DS, Carlson JM. Individual differences in dynamic functional brain connectivity across the human lifespan. PLoS Comput Biol. 2016 Nov 23;12(11):e1005178. doi:10.1371/journal.pcbi.1005178.

32. Müller V, Brehmer Y, von Oertzen T, Li S-C, Lindenberger U. Electrophysiological correlates of selective attention: A lifespan comparison. BMC Neurosci. 2008;9(1):18. https://doi.org/10.1186/1471-2202-9-18.

33. Müller V, Gruber W, Klimesch W, Lindenberger U. Lifespan differences in cortical dynamics of auditory perception. *Dev Sci.* 2009;12(6):839–53. https://doi.org/10.1111/j.1467-7687.2009.00834.x.

34. Müller V, Lindenberger U. Lifespan differences in nonlinear dynamics during rest and auditory oddball performance. Dev Sci. 2012 Jul;15(4):540–556. doi:10.1111/j.1467-7687.2012.01153.x.

35. Müller V, Jirsa V, Perdikis D, Sleimen-Malkoun R, von Oertzen T, Lindenberger U. Lifespan changes in network structure and network topology dynamics during rest and auditory oddball performance. Front Aging Neurosci. 2019 Jun 11;11:138. doi:10.3389/fnagi.2019.00138.

36. Deco G, Jirsa V, McIntosh AR, Sporns O, Kötter R. Key role of coupling, delay, and noise in resting brain fluctuations. Proc Natl Acad Sci U S A. 2009 Jun 2;106(25):10302–10307. doi:10.1073/pnas.0901831106.

37. Garrett DD, Kovacevic N, McIntosh AR, Grady CL. The importance of being variable. J Neurosci. 2011 Mar 23;31(12):4496–4503. doi:10.1523/JNEUROSCI.5641-10.2011.

38. Tognoli E, Kelso JAS. The metastable brain. Neuron. 2014 Jan 8;81(1):35–48. doi:10.1016/j.neuron.2013.12.022.

39. Garrett DD, Kovacevic N, McIntosh AR, Grady CL. The modulation of BOLD variability between cognitive states varies by age and processing speed. Cereb Cortex. 2013 Mar;23(3):684–693. doi:10.1093/cercor/bhs055.

40. Sleimen-Malkoun R, Temprado JJ, Hong SL. Aging induced loss of complexity and dedifferentiation: consequences for coordination dynamics within and between brain, muscular and behavioral levels. Front Aging Neurosci. 2014 Jun 27;6:140. doi:10.3389/fnagi.2014.00140.

41. Sleimen-Malkoun R, Perdikis D, Müller V, Blanc J-L, Huys R, Temprado J-J, Jirsa VK. Brain dynamics of aging: Multiscale variability of EEG signals at rest and during an auditory oddball task. eNeuro. 2015 Jun 3;2(3):ENEURO.0067-14.2015. doi:10.1523/ENEURO.0067-14.2015.

42. Goodman ZT, Nomi JS, Kornfeld S, Bolt T, Saumure RA, Romero C, Bainter SA, Uddin LQ. Brain signal variability and executive functions across the life span. *Network Neuroscience*. 2024;8(1):226–240. doi:10.1162/netn_a_00347.

43. Courtiol J, Perdikis D, Petkoski S, Müller V, Huys R, Sleimen-Malkoun R, Jirsa VK. The multiscale entropy: Guidelines for use and interpretation in brain signal analysis. J Neurosci Methods. 2016 Nov 1;273:175–190. doi:10.1016/j.jneumeth.2016.09.004.

44. Mizuno T, Takahashi T, Cho RY, Kikuchi M, Murata T, Takahashi K, et al. Assessment of EEG dynamical complexity in Alzheimer's disease using multiscale entropy. Clin Neurophysiol. 2010 Sep;121(9):1438–46. doi:10.1016/j.clinph.2010.03.025. PMID: 20400371; PMCID: PMC2914820.

45. Vakorin VA, Lippé S, McIntosh AR. Variability of brain signals processed locally transforms into connectivity in human brain networks. J Neurosci. 2011 Apr 27;31(17):6405–6413. doi:10.1523/JNEUROSCI.3153-10.2011.

46. Fjell AM, Walhovd KB, Reinvang I, Lundervold A, Salat DH, Quinn BT, Fischl B, Dale AM. Selective increase of cortical thickness in high-performing elderly—structural indices of optimal cognitive aging. Neuroimage. 2006;29(3):984–994. doi:10.1016/j.neuroimage.2005.08.007.

47. Jockwitz C, Caspers S, Lux S, Eickhoff SB, Jütten K, Lenzen S, Moebus S, Pundt N, Reid A, Hoffstaedter F, Jöckel KH, Erbel R, Cichon S, Nöthen M, Shah NJ, Zilles K, Amunts K. Influence of age and cognitive performance on resting-state brain networks of older adults in a population-based cohort. Cortex. 2017;89:28–44. doi:10.1016/j.cortex.2017.01.008.

48. Jauny G, Mijalkov M, Canal-Garcia A, Volpe G, Pereira J, Eustache F, Hinault T. Linking structural and functional changes during aging using multilayer brain network analysis. Commun Biol. 2024 Feb 28;7(1):239. doi:10.1038/s42003-024-05927-x.

49. Luo N, Sui J, Abrol A, Lin D, Chen J, Vergara VM, Fu Z, Du Y, Damaraju E, Xu Y, Turner JA, Calhoun VD. Age-related structural and functional variations in 5,967 individuals across the adult lifespan.

Hum Brain Mapp. 2020 May;41(7):1725–1737. doi:10.1002/hbm.24905. Epub 2019 Dec 26. PMID: 31876339; PMCID: PMC7267948.

50. Wu K, Taki Y, Sato K, Kinomura S, Goto R, Okada K, Kawashima R, He Y, Evans AC, Fukuda H. Age-related changes in topological organization of structural brain networks in healthy individuals. Hum Brain Mapp. 2012 Mar;33(3):552–568. doi:10.1002/hbm.21232.

51. Wang L, Su L, Shen H, Hu D. Decoding lifespan changes of the human brain using resting-state functional connectivity MRI. PLoS ONE. 2012 Aug 30;7(8):e44530. doi:10.1371/journal.pone.0044530.

52. Sun L, Zhao T, Liang X, Xia M, Li Q, Liao X, Gong G, Wang Q, Pang C, Yu Q, Bi Y, Chen P, Chen R, Chen Y, Chen T, Cheng J, Cheng Y, Cui Z, Dai Z, Deng Y, Ding Y, Dong Q, Duan D, Gao JH, Gong Q, Han Y, Han Z, Huang CC, Huang R, Huo R, Li L, Lin CP, Lin Q, Liu B, Liu C, Liu N, Liu Y, Liu Y, Lu J, Ma L, Men W, Qin S, Qiu J, Qiu S, Si T, Tan S, Tang Y, Tao S, Wang D, Wang F, Wang J, Wang P, Wang X, Wang Y, Wei D, Wu Y, Xie P, Xu X, Xu Y, Xu Z, Yang L, Yuan H, Zeng Z, Zhang H, Zhang X, Zhao G, Zheng Y, Zhong S, Alzheimer's Disease Neuroimaging Initiative, DIDA-MDD Working Group, MCADI, He Y. Functional connectome through the human life span. bioRxiv. 2023 Sep 13;2023.09.12.557193. doi:10.1101/2023.09.12.557193.

53. Cepeda NJ, Kramer AF, Gonzalez de Sather JCM. Changes in executive control across the life span: examination of task-switching performance. Dev Psychol. 2001 Sep;37(5):715–30. doi:10.1037/0012-1649.37.5.715.

54. Craik FIM, Bialystok E. Cognition through the lifespan: mechanisms of change. Trends Cogn Sci. 2006 Mar;10(3):131–8. doi:10.1016/j.tics.2006.01.007.

55. Hartshorne JK, Germine LT. When does cognitive functioning peak? The asynchronous rise and fall of different cognitive abilities across the life span. Psychol Sci. 2015 Apr;26(4):433–43. doi:10.1177/0956797614567339.

56. Leversen JSR, Haga M, Sigmundsson H. From children to adults: motor performance across the life-span. PLoS One. 2012 Jun 18;7(6):e38830. doi:10.1371/journal.pone.0038830.

57. Schaie KW. Developmental Influences on Adult Intelligence: The Seattle Longitudinal Study. New York: Oxford University Press; 2005.

58. Meyer-Lindenberg A. The evolution of complexity in human brain development: an EEG study. Electroencephalogr Clin Neurophysiol. 1996 Nov;99(5):405–411. doi:10.1016/S0013-4694(96)95699-0.

59. McIntosh AR, Kovacevic N, Lippe S, Garrett D, Grady C, Jirsa V. The development of a noisy brain. Archives Italiennes de Biologie. 2010;148:323–337.

60. Fernández A, Zuluaga P, Abásolo D, Gómez C, Serra A, Méndez MA, Hornero R. Brain oscillatory complexity across the life span. Clin Neurophysiol. 2012 Nov;123(11):2154–2162. doi:10.1016/j.clinph.2012.04.025.

61. Shou G, Yuan H, Cha YH, Li Y, Sclabassi RJ, Sun M. Age-related changes of whole-brain dynamics in spontaneous neuronal coactivations. Sci Rep. 2022;12:12140. doi:10.1038/s41598-022-16125-2.

62. James TM, Burgess AP. Estimating chronological age from the electrical activity of the brain: how EEG-age can be used as a marker of general brain functioning. Psychophysiology. 2025 Mar 16;62(3):e70033. doi:10.1111/psyp.70033.

63. Joffe D, Oakley DS, Arese Lucini F, Palermo FX. Measurements of EEG alpha peak frequencies over the lifespan: validating target ranges on an in-clinic platform. bioRxiv. 2021 Oct 7;2021.10.06.463353. doi:10.1101/2021.10.06.463353.

64. Tou SLJ, Chau T. The fractal dimension of resting state EEG increases over age in children. Cereb Cortex. 2025 Jun;35(6):bhaf138. doi:10.1093/cercor/bhaf138.

65. Vakorin VA, McIntosh AR, Mišić B, Krakovska O, Poulsen C, Martinu K, Paus T. Exploring transient transfer entropy based on a group-wise reconstruction of functional brain networks. PLoS One. 2013;8(3):e57217. doi:10.1371/journal.pone.0057217.

66. Zappasodi F, Marzetti L, Olejarczyk E, Tecchio F, Pizzella V. Age-related changes in electroencephalographic signal complexity. PLoS One. 2015;10(11):e0141995. doi:10.1371/journal.pone.0141995.

67. Costa M, Goldberger AL, Peng CK. Multiscale entropy analysis of complex physiologic time series. Phys Rev Lett. 2002;89(6):068102.

68. Costa M, Goldberger AL, Peng CK. Multiscale entropy analysis of biological signals. Phys Rev E Stat Nonlin Soft Matter Phys. 2005;71(2 Pt 1):021906.

69. Richman JS, Moorman JR. Physiological time-series analysis using approximate entropy and sample entropy. Am J Physiol Heart Circ Physiol. 2000;278(6):H2039–H2049.

70. Valencia JF, Porta A, Vallverdu M, Claria F, Baranowski R, Orłowska-Baranowska E, et al. Refined multiscale entropy: application to 24-h Holter recordings of heart period variability in healthy and aortic stenosis subjects. IEEE Trans Biomed Eng. 2009 Sep;56(9):2202-13. doi: 10.1109/TBME.2009.2021986.

71. Burgess AP. On the interpretation of synchronization in EEG hyperscanning studies: a cautionary note. Front Hum Neurosci. 2013 Dec 24;7:881. doi: 10.3389/fnhum.2013.00881.

72. Lachaux JP, Rodriguez E, Le Van Quyen M, Lutz A, Martinerie J, Varela FJ. Studying single-trials of phase synchronous activity in the brain. Int J Bifurcat Chaos. 2000;10(10):2429–39. doi:10.1142/S0218127400001560

73. McIntosh AR, Bookstein FL, Haxby JV, Grady CL. Spatial pattern analysis of functional brain images using partial least squares. Neuroimage. 1996 Jun;3(3):143-57. doi: 10.1006/nimg.1996.0016.

74. McIntosh AR, Lobaugh NJ. Partial least squares analysis of neuroimaging data: applications and advances. Neuroimage. 2004;23(Suppl 1):S250-63. doi: 10.1016/j.neuroimage.2004.07.020.

75. Roy D, Ghosh A, Jirsa VK. Phase description of spiking neuron networks with global electric and synaptic coupling. Phys Rev E. 2011;83(5):051909. doi:10.1103/physreve.83.051909.

76. Sanz Leon P, Knock SA, Woodman MM, Domide L, Mersmann J, McIntosh AR, Jirsa VK. The Virtual Brain: a simulator of primate brain network dynamics. Front Neuroinform. 2013 Jun 11;7:10. doi: 10.3389/fninf.2013.00010.

77. Novikov AV. PyClustering: Data Mining Library. J Open Source Softw. 2018;3(25):1230. doi:10.21105/joss.01230.

78. Jonas E, Kording KP. Could a Neuroscientist Understand a Microprocessor? PLoS Comput Biol. 2017 Jan;13(1):e1005268. doi:10.1371/journal.pcbi.1005268.

79. Segalowitz SJ, Santesso DL, Jetha MK. Electrophysiological changes during adolescence: a review. Brain Cogn. 2010;72(1):86–100.

80. Getzmann S, Gajewski PD, Schneider D, Drolshagen S, Falkenstein M. Resting-state EEG data before and after cognitive activity across the adult lifespan and a 5-year follow-up. Sci Data. 2024;11:988. doi:10.1038/s41597-024-03797-w.

81. Capouskova K, Zamora-López G, Kringelbach ML, Deco G. Integration and segregation manifolds in the brain ensure cognitive flexibility during tasks and rest. *Hum Brain Mapp*. 2023;44(18):6349–6363. doi:10.1002/hbm.26511. PMCID: PMC10681658.

82. Rabuffo G, Fousek J, Bernard C, Jirsa V. Neuronal cascades shape whole-brain functional dynamics at rest. *eNeuro*. 2021;8(5):ENEURO.0283-21.2021. doi:10.1523/ENEURO.0283-21.2021.

83. Sorrentino P, Troisi Lopez E, Romano A, Granata C, Corsi MC, Sorrentino G, et al. Brain fingerprint is based on the aperiodic, scale-free, neuronal activity. Neuroimage. 2023;277:120260. doi:10.1016/j.neuroimage.2023.120260.

84. Runfola C, Petkoski S, Sheheitli H, Bernard C, McIntosh AR, Jirsa V. A mechanism for the emergence of low-dimensional structures in brain dynamics. *NPJ Syst Biol Appl*. 2025;11(1):32. doi:10.1038/s41540-025-00499-w.

A Young Adult EEG at rest with eyes closed

Fz
Cz
Pz
O1

B Univariability metrics

C Phase synchronization among Fz. Cz. Pz, O1 at 10 Hz

Fz<->Cz
Fz<->Pz
Fz<->O1
Cz<->Pz
Cz<->O1
Pz<->O1

D Phase synchronization (PS(t)) among 19 channels of the 10-20 system -i.e., 171 connections- at 10 Hz

E Phase synchronization norm for each frequency

20 Hz
18 Hz
16 Hz
14 Hz
12 Hz
10 Hz
8 Hz
6 Hz
4 Hz
2 Hz

F Phase synchronization dynamics at 10 Hz: correlation distance among distinct time moments $(1-corr(PS(t_1),PS(t_2))$

G Jump lengths for distinct time scales (JL(t,τ)) at 10 Hz

τ=0.004 sec
τ=0.204 sec
τ=0.404 sec
τ=0.604 sec
τ=0.804 sec
τ=1.004 sec
τ=1.204 sec
τ=1.404 sec
τ=1.604 sec
τ=1.804 sec
τ=2 sec

H JL(τ) probability distributions at 10 Hz