Uncovering *Fusarium* species diversity, distribution and pathogenicity associated

with cotton-corn rotation fields in recently expanded cotton-growing areas of the

Northern High Plains of Texas

Ihsanul Khaliq, and Nolan R. Anderson*

Texas A&M AgriLife Research and Extension Center, Amarillo, TX 79106, U.S.A.

*Corresponding author: Nolan Anderson; E-mail: nolan.anderson@ag.tamu.edu

15 **Keywords:** *Gossypium* species, maize, *Zea mays*, Texas High Plains, *Fusarium*

16 diseases, *Fusarium* diversity, Fusarium wilt, FOV4, semi-arid climate

19 Funding: Cotton Inc. 24-115

20 Texas A&M AgriLife Research Capacity Support

Abstract

The recent expansion of cotton acreage in the Northern High Plains of Texas has

raised concerns about the occurrence of a diverse *Fusarium* community, particularly

Fusarium oxysporum f. sp. *vasinfectum* race 4. We investigated the diversity,

distribution, and pathogenicity of *Fusarium* species in soil samples collected from 26

cotton-corn rotation fields across 10 counties. A total of nine *Fusarium* species were

isolated: *F. incarnatum*, *F. equiseti*, *F. solani*, *F. verticillioides*, *F. proliferatum*, *F.*

clavum, *F. oxysporum*, *F. flocciferum*, and one unidentified *Fusarium* sp. The highest

species diversity (five species) was observed in Sherman County. *Fusarium solani* was

the most frequently isolated species (39%) across all counties, followed by *F. equiseti*

(22%), while *F. flocciferum* was the least frequently isolated species (2%). Greenhouse

pathogenicity assays showed *F. incarnatum* caused significantly higher disease severity

relative to non-inoculated control on aerial parts of cotton ($P=0.0137$), while *Fusarium*

flocciferum, F. proliferatum, and *F. clavum* caused significantly higher disease severity

relative to non-inoculated control on roots ($P=0.01$, $P=0.01$, $P=0.02$, respectively). On

corn, differences in disease severity were not statistically significant ($P>0.05$). *Fusarium*

isolates were recovered from both symptomatic and asymptomatic roots following

pathogenicity trials, suggesting endophytic colonization. This study revealed a complex

Fusarium community capable of cross-infecting cotton and corn, with implications for

regional disease management. The absence of *Fusarium oxysporum* f. sp. *vasinfectum*

race 4 indicates it may not yet have been introduced into the Northern High Plains;

however, continued surveillance is warranted given its detection in New Mexico and

adjacent counties in Texas.

The Northern High Plains are a mixture of rangeland and cropland, comprising

the 22 northernmost counties in Texas (Zivkovic and Hudson 2012). The region faces

severe water scarcity, with an annual mean precipitation of about 558 mm–well below

the US national average of 965 mm per year (BestPlaces 2025; NOAA 2021).

Additionally, the region relies heavily on water drawn from the Ogallala Aquifer to

produce irrigated crops. However, current water use is unsustainable, with annual

extraction exceeding annual recharge (Thayer and McCarl 2018). Approximately 7

million beef cattle are raised annually in beef cattle feedyards in the Texas High Plains

(TCFA 2015). As of March 2025, there were 2.35 million head of cattle and calves on

feed in the Northern High Plains, accounting for 89 percent of the state's total (USDA

National Agricultural Statistics Service 2025). Corn remains the primary feed crop, with

more than 50 percent of the corn grown in Texas coming from the High Plains region

(National IPM Database 2001; Xue et al. 2017). Although corn is the major feed crop,

there has been significant interest in growing cotton as a rotational crop due to water

constraints and the high fertilizer costs associated with corn production (Colaizzi et al.

2009).

Cotton (*Gossypium* species) is the most important fiber crop for the textile

industry and a significant oilseed crop globally. The United States is the world's fourth-

largest cotton producing country (USDA Foreign Agricultural Service 2024), with

cultivation spanning 17 southern states. Texas is the leading producer, accounting for

approximately 40% of the U.S. cotton production in recent years (USDA Economic

Research Service 2022). Although cotton was introduced to the Northern High Plains in

1968, acreage remained below 20,000 ha until 1998, but has expanded substantially

since 2004—including a 126% increase between 2013 and 2019 alone—raising

concerns about diseases that may affect cotton production (Ledbetter 2020; Marek and

Bordovsky 2006).

Key fungal genera implicated in cotton diseases worldwide and in the U.S.

Cotton Belt include *Fusarium*, *Verticillium*, *Rhizoctonia*, and *Pythium*. Among these,

Fusarium, *Rhizoctonia*, and *Pythium* species are primarily associated with the seedling

disease complex (Howell 2002; Watkins 1981), while *Fusarium* species—alone or in

combination with *Alternaria*, *Aspergillus* species, or bacterial pathogens such as

Pantoea agglomerans and *Serratia marcescens*—are frequently linked to seed rot

(Glover et al. 2020; Medrano et al. 2007; Minton and Garber 1983). Diseases caused

by oomycetes and splash dispersed pathogens are not considered a major concern in

the Northern High Plains due to the dry climate and severe water scarcity.

Fusarium wilt, caused by *Fusarium oxysporum* f.sp. *vasinfectum* (FOV), is of

particular concern in the Northern High Plains because cooler temperatures during the

early planting season favor its development (Zhang et al. 2020). Night temperatures in

the Northern Panhandle are among the coolest in cotton growing regions of the U.S.,

with mean annual temperature of about 11°C (Kimmel Jr et al. 2016). FOV was first

detected in the United States in 1892 (Atkinson 1892). Historically, FOV has been

characterized into 8 different races based on variation in pathogenicity to various plant

species, including *Gossypium* species and cultivars. Among these, race 4 (FOV4) is of

particular concern in the U.S. due to its high virulence on both Upland and Pima cotton,

as well as its ability to persist in soil for years as chlamydospores (Hutmacher et al.

2013; Kim et al. 2005). Additionally, unlike other races, FOV4 can cause severe wilting

in the absence of root knot nematode *Meloidogyne incognita.* FOV4 was first detected

in the U.S. in 2001 in California (Kim et al. 2005) and more recently in 2017 in Texas

(Halpern et al. 2018) and 2019 in New Mexico (Zhu et al. 2019). This highly aggressive

race has not yet been reported in the Northern High Plains. However, given the

proximity to New Mexico and adjacent counties in Texas where FOV4 has been

detected (Cianchetta et al. 2015; Diaz et al. 2021), there is great concern about its

potential introduction into the Northern High Plains. Therefore, a comprehensive survey

is urgently needed to investigate the presence or absence of FOV4 in the region.

Several surveys have been conducted across the U.S. Cotton Belt for *Fusarium*

species (Cianchetta et al. 2015; Diaz et al. 2021; Mao and Hu 2019; Zhu et al. 2021).

However, these surveys did not include the Northern High Plains, leaving limited

information on the diversity, distribution and relative severity of *Fusarium* species

affecting cotton in this region. Moreover, these studies were primarily focused on

investigating the presence or absence of FOV4 rather than determining overall

Fusarium species diversity. Several *Fusarium* species are known to cause economically

important diseases in cotton, including wilt, damping-off, boll rot, and root rot (Davis et

al. 2006; Zaki et al. 2021). While *F. oxysporum* and *F. solani* are the most common

species complexes associated with cotton, the relative severity and impact of other

Fusarium species remain poorly understood (Mao and Hu 2019).

Crop rotation has been recognized as an effective practice to reduce nitrogen

fertilizer use and enhance both grain yield and sustainability in agricultural systems

(Shu et al. 2021). However, crop rotation can also inadvertently increase inoculum

levels, thereby elevating the risk of crop diseases (Li et al. 2021). Previous studies have

demonstrated that crop residue serves as a natural reservoir for *Fusarium* species and

may even facilitate genetic exchange (Ward et al. 2008). For instance, in wheat–corn

cropping rotations, continuous wheat cultivation following corn has been shown to

increase the incidence of *Fusarium* stalk rot and ear rot in maize, as well as *Fusarium*

head blight in wheat (Xi et al. 2021; Zhang et al. 2016). In the Northern High Plains,

where cotton and corn are commonly grown in rotation, investigating the pathogenicity

of *Fusarium* species on both crops is crucial. Such studies will provide valuable insights

into the pathogen's persistence, cross-infectivity, and the associated epidemiological

risks.

Greenhouse pathogenicity assays offer several advantages, including reduced

variability resulting from uneven inoculum distribution and external biotic or abiotic

factors. Most pathogenicity assays conducted in greenhouse conditions use root-dip

inoculation method, where two-week-old seedlings' roots are dipped in a spore

suspension for two minutes and then transplanted into healthy soil (Kim et al. 2005).

This method can predispose seedlings to stress during transplantation and does not

reflect host infection in the field (Schoeneweiss 1975). Conversely, pathogenicity

assays that involve directly planting seeds in *Fusarium*-infested soil or adding inoculum

after seedling emergence better reflect infection under field conditions (Diaz et al.

2021).

To address these key gaps, the present study was conducted to: (1) conduct a

comprehensive survey to determine the diversity and distribution of *Fusarium* species

associated with cotton-corn rotation fields in the recently expanded cotton-growing

areas of the Northern High Plains; (2) investigate the presence or absence of FOV4 in

the region; and (3) evaluate the pathogenicity and severity of *Fusarium* species on

cotton and corn under greenhouse conditions.

Materials and Methods

Sampling sites and sampling procedure

With the help of growers and county agents, a total of 26 cotton-corn rotation

fields across 10 different counties in the Northern High Plains of Texas were sampled

between late spring and late fall (May to October) of 2024. Specifically, six fields were

sampled in Sherman County; four fields each in Moore and Deaf Smith counties; three

fields in Hartley County; two fields each in Randall, Hansford, and Carson counties; and

one field each in Potter, Dallam, and Hutchinson counties. Fields selection was based

on growers or county agents' willingness to participate in the study.

In each field, a total of 25 soil cores, each 0.2 m deep, were collected at 5

different locations, with at least 60 m space between sampling points (5 soil cores per

sampling location within a field, 25 soil cores per field). The soil samples from each field

were pooled and placed into cloth bags, kept on ice in an insulated box to protect them

from high temperature and direct sunlight, and carried to the laboratory.

In the laboratory, the samples collected from each field were mixed thoroughly

and air dried for a week to inhibit microbial activity and facilitate grinding and sieving

(Sangalang et al. 1995). The air-dried soil sample for each field was poured into a

Nasco-Asplin soil grinder, pulverized gently to break the clods, and passed through a 2

mm sieve. To prevent cross-contamination between samples, the excess dirt from

grinder's parts were removed with a brush, washed with water, and then cleaned

thoroughly with 70% ethanol and paper towels.

Diversity and distribution of *Fusarium* species in the Northern High Plains

Isolation of single spore isolates from soil

To isolate *Fusarium* species from soil, a total of 20 cc of soil was added to 80 mL of reverse osmosis (RO) water in a small beaker (200 mL) placed on a magnetic stirrer and mixed vigorously to ensure a thorough mixing of the soil. From this soil suspension, a 1 mL aliquot was plated onto each of 5 Petri dishes of Peptone PCNB agar (PPA) medium (1000 mL distilled water, 15 g Peptone, 1 g KH_2PO_4, 0.5 g $MgSO_4 \cdot 7H_2O$, 0.75 g PCNB, 20 g agar, pH adjusted to 5.5–6.5) (Nash and Snyder 1962). The process was repeated with an additional 20 cc of soil using another set of five Petri dishes, allowing us to assess 40 cc of soil for each field using a total of 10 Petri dishes. The Petri dishes were incubated for 3 days on PPA at room temperature (23 ± 2°C) with a 12-hour photoperiod. The colonies developed on PPA were subcultured onto full strength potato dextrose agar (PDA) medium by taking small mycelial plugs, observed under a Leica M205C stereoscope (Leica Camera AG, Wetzlar, Germany) at 160x magnification, from the colony margins.

After incubating for two weeks on PDA, 200 µL of RO water was added to each Petri dish, and conidia were dislodged by gently scrapping the mycelium with a pipette tip. A total of 150 µL of this conidial suspension was then transferred onto 30% water agar and incubated overnight. Single spore isolates were obtained by transferring single germinated conidia, observed under Nikon SM225 microscope (Nikon corp., Tokyo, Japan) at 180x magnification, onto fresh PDA plates.

Genomic DNA extraction, PCR amplification, and sequencing

Isolates were divided into morphotypes based on their gross colony morphology

(e.g., mycelial growth patterns), and their identities were obtained using the nuclear

ribosomal internal transcribed spacer region (ITS rDNA) and the translation elongation

factor 1-α (tef-1α) gene region (Table 1). DNA was extracted using Zymo Quick DNA

Fungal/Bacterial Microprep kit (Zymo Research Corp; Irvine, CA). Amplicons were

obtained using the following primer pairs: ITS1 (TCCGTAGGTGAACCTGCG) × ITS4

(TCCTCCGCTTATTGATATGC) for ITS rDNA (White et al. 1990); and EF-1 (5′-

ATGGGT AAGGARGACAAGAC-3′) × EF-2 (5′-GGARGTACCA GTSATCATG-3′) for tef-

1α gene region (O'Donnell et al. 1998). The PCR reaction mixture (25 μl reaction)

contained 12.5 μl of Phire Green Hot Start II Master Mix (Thermo scientific), 1.25 μl of

each primer (0. 5 μM), and 1.5 μl of genomic DNA. The PCR amplification program

conditions were as follows: initial denaturation at 98°C for 90 s followed by 35 cycles at

98°C for 5 s, annealing at 58°C for 15 s and extension at 72°C for 15 s, followed by final

extension for 4 minutes at 72°C.

The PCR products were detected using 1% agarose gel in 1x TAE buffer. The

PCR products were cleaned using DNA Clean & Concentrator kit (Zymo Research) and

sent to Eurofins Genomics (Eurofins Genomics LLC, Louisville, Kentucky) for

sequencing with both forward and reverse primers. ABI sequence chromatograms were

edited using SnapGene 8.0 (https://www.snapgene.com/) and exported as FASTA files

that were aligned using MUSCLE alignment (Edgar 2004) in Mega11

(https://www.megasoftware.net/). Consensus sequences were obtained using BioEdit

7.7.1 (http://www.mbio.ncsu.edu/BioEdit/bioedit.html), which were then used to conduct

BLASTn queries of NCBI GenBank (https://www.ncbi.nlm.nih. gov/genbank/) to obtain

preliminary identifications.

Given the regional importance of *F. oxysporum*, two isolates identified as *F.*

oxysporum were additionally amplified using RNA polymerase II second largest subunit

(RPB2) (Liu et al. 1999) with the primer pair: 5f2 (GGGGWGAYCAGAAGAAGGC) × 7cr

(CCCATRGCTTGYTTRCCCAT), using the PCR conditions described above, to further

confirm its identity.

The two isolates identified as *F. oxysporum* in the present study were tested to

determine whether they are FOV4 (and to determine whether they are genotype N, T,

MT, and MiT) using specific primer pairs developed by Bell et al. (2019), following the

protocol described by Wagner et al. (2022). This PCR detection method has been

developed based on Tfo1 insertion event at the NAT gene locus and provides a rapid

alternative approach to vegetative compatibility analysis, which is labour-intensive and

time consuming (Wagner et al. 2022). Specific primer pairs included: FovT-F

(GGCCGATATTGTCGGTCGTA) x FovA-R (TCAACAGACCCTGCACTACG) for the

detection of FOV4; FovP-F (GGCCGATATTGTCGGTCGTA) x FovP-R

(CTCCAGTGCAGTGCTTGGTA) for the detection of FOV4 N genotype; FovP-F

(GGCCGATATTGTCGGTCGTA) x FovT-R (ATCTGTCTTTCGTCGGCAAT) for the

detection of FOV4 T or FOV4 MiT genotype; and FovP-F

(GGCCGATATTGTCGGTCGTA) x FovM-R (CCGCCATATCCACTGAACA) for the

detection of FOV4 MT genotype.

Phylogenetic analyses

Nucleotide sequences generated in this study, along with reference sequences

retrieved from the NCBI GenBank database, were aligned using MUSCLE alignment

(Edgar 2004) in Mega11 (https://www.megasoftware.net/). Alignments were then

trimmed avoiding missing data at either end of alignments. The aligned sequences were

analysed with maximum likelihood bootstrapping (MLBS) using IQ-TREE (http://

www.iqtree.org/) (Nguyen et al. 2015) to determine the best-fit model of molecular

evolution. These models were based on the Bayesian information criterion (BIC) scores

(Chernomor et al. 2016) using ModelFinder (Kalyaanamoorthy et al. 2017). Once the

optimal models were determined, clade support was assessed via 50,000 ML-BS

pseudoreplicates of the data using IQ-TREE.

To determine the phylogenetic placement of *Fusarium oxysporum* sequences

generated in this study relative to known FOV races (Cianchetta et al. 2015; Wang et

al. 2010) and closely related sequences, additional reference sequences representing

recognized FOV races and top BLASTn hits from the NCBI GenBank database were

retrieved and phylogenetic analysis was conducted as described above. DNA

sequences generated in the present study were deposited in NCBI (Table 1).

Pathogenicity trials

A single trial was conducted to test the pathogenicity and severity of seven

isolates, each representing a different *Fusarium* species, on corn and cotton. These

included *F. incarnatum, F. equiseti, F. solani, F. verticillioides, F. proliferatum, F.*

clavum, and F. flocciferum. Given its regional importance, *F. oxysporum* was not

included in the inoculated pathogenicity trials due to regional concerns in relation to

247 potentially working with FOV and the potential risk of its spread into these recently

248 expanded cotton-growing areas.

249 **Pathogenicity of *Fusarium* species on cotton**

250 **Biological material and experimental design**

251  The trial was conducted under greenhouse conditions at Texas A&M AgriLife

252 Research, Bushland, Texas at ambient temperatures (25 ± 5°C). A commonly cultivated

253 variety in the Texas High Plains, FiberMax 868AXTP, with fair resistance to Fusarium

254 wilt was used in this study (BASF 2024). Disease-free seeds of this variety were

255 sourced from a previous trial: seeds were de-linted but received no chemical treatment

256 prior to planting. Briefly, 10-cm pots were filled with a modified media (1 bag of Berger's

257 OM1 potting mix, 0.085 m^3 of organic growing mix, 2.2 kg of sand, and 3 kg of soil). A

258 total of 2-3 disease-free seeds were sown in each pot. Upon germination, seedlings

259 were thinned to one per pot. There were six replicate pots for each *Fusarium* species

260 and a non-inoculated control (total 48 plants), arranged in a completely randomized

261 block design. Throughout the trial period, seedlings were fertilized with water-soluble

262 fertilizer 'Miracle-Gro' as needed.

263 **Inoculum preparation**

264  The sorghum grain inoculation method was used to prepare the inoculum (Yadav

265 et al. 2017). Disease-free sorghum grains were sourced from a previous year's trial.

266 Briefly, sorghum grains were soaked overnight, and 50 g of soaked grains were placed

267 into each 250 mL beaker and sealed with aluminium foil. The beakers were autoclaved

268 at 121°C for 30 minutes on two consecutive days. Once cooled, the sorghum grains in

269 each beaker were inoculated with five agar plugs (5 mm), which were cut from a 7-day-

old colony of specific *Fusarium* isolate grown on PDA. The inoculated beakers were

gently shaken to uniformly distribute the inoculum and then placed inside zip-lock plastic

bags to avoid contamination. The inoculated beakers were incubated for 7 days at room

temperature (23 ± 2°C) with a 12-hour photoperiod and shaken every 2–3 days to

ensure even distribution of the inoculum.

Inoculation method

Cotton seedlings were inoculated at 2–3 leaf stage. Seedlings were watered

thoroughly in the evening prior to inoculation the following morning. To inoculate

seedlings, four 5-cm deep holes were made vertically into the potting medium of each

pot using a wooden rod (18 x 0.8 cm), approximately 2 to 4 cm from the crown of each

plant. A total of 1.25 g inoculum was inserted into each hole, excluding the non-

inoculated controls. The holes were then covered with potting medium, and plants were

watered daily following inoculation.

Disease assessment

Cotton seedlings were observed daily for the appearance of disease symptoms.

Disease severity on aerial plant parts was recorded at 7, 14, 21, and 28 days post-

inoculation using a 0–4 rating scale, where: 0 = healthy; 1 = cotyledons wilted only; 2 =

≤50% true leaves wilted; 3= >50% but ≤90% true leaves wilted; and 4 = all leaves wilted

or plant death (Table S1).

All cotton seedlings were harvested after one month of inoculation. Roots were

washed free of soil and disease severity was recorded using 0–5 rating scale, where 0 =

no symptoms; 1 = <10% of roots discolored; 2 = 10–30% of roots discolored; 3= 31–

50% of roots discolored; 4 = 51–80% of roots discolored; and 5 = 81–100% of roots

discolored (Table S1).

At the end of the trial, the roots of all cotton seedlings were plated onto acidified

PDA (1000 mL RO water, 39 g potato dextrose agar, 1 g streptomycin) to confirm

pathogenicity.

Pathogenicity of *Fusarium* species on corn

Biological material, experimental design, inoculum preparation and inoculation

method

A commonly cultivated drought tolerant, untreated corn hybrid, Blue River

Organic 82-14P, was used in the pathogenicity trial.

The experimental design, inoculum preparation and application method, as well

as cultural practices, were the same as those used in the cotton pathogenicity trial. That

is, a single batch of inoculum was used to inoculate both cotton and corn seedlings at

the same time and similar cultural practices were followed to ensure consistency.

Seedlings were inoculated at the three-leaf stage.

Disease assessment

Corn seedlings were harvested after one month of inoculation. Roots were

washed free of soil and crown and root lesions were rated using a 1–8 rating scale

(Parikh et al. 2018), where: 1 = no visible symptoms, 0% infection, and normal

development; and 8 = seedling barely emerged and could not grow, scanty root but no

emergence, or seed is dead (no emergence and no roots); infection coverage of

seedling is >95% (Table S1).

At the end of the trial, the roots of all corn seedlings were plated onto acidified

PDA (1000 mL RO water, 39 g potato dextrose agar, 1 g streptomycin) to confirm

pathogenicity.

Data Analysis

In the case of disease severity recorded on aerial parts of cotton, the area under

the disease progress stairs (AUDPS) was calculated using the 'audps()' function from

'agricolae' R package (Mendiburu 2023), based on severity ratings recorded across all

four disease assessments. The AUDPS was selected over the traditional area under the

disease progress curve (AUDPC) because the latter underestimates the effect of

disease severity recorded during first and last disease assessments (Simko and Piepho

2012).

Linear mixed models effect models (LMMs) were then fitted using the 'lmer()'

function from the 'lmerTest' R package (Kuznetsova et al. 2022) to investigate

differences in disease severity among *Fusarium* species. The AUDPS was included as

the dependent variable, the predictor *Fusarium* species were included as a fixed effect,

and different blocks were included as a random effect to account for variability across

blocks. Estimated marginal means for each *Fusarium* species were calculated using the

'emmeans()' function from 'emmeans' R package (Lenth 2021). The estimated marginal

means were compared using Tukey-adjusted pairwise comparisons, and groupings

were assigned using the 'cld()' function from the 'multcomp' R package (Hothorn 2020).

These groupings indicated significant differences at $\alpha = 0.05$. Models diagnostics was

performed using the 'DHARMa' R package, which uses a simulation-based approach

via the function 'simulateResiduals' to create readily interpretable scaled residuals for

models (Hartig 2020).

In the case of disease severity recorded on roots of cotton and corn, differences

in disease severity were tested using a nonparametric Kruskal-Wallis test using

'ggbetweenstats()' function from the 'ggstatplot' package (Patil 2021). If significant

differences were found (α = 0.05), pairwise differences were performed using a post hoc

Dunn test.

Results

Diversity and distribution of *Fusarium* species in the Northern High Plains

A total of nine *Fusarium* species were identified: these included *F. incarnatum, F.*

equiseti, F. solani, F. verticillioides, F. proliferatum, F. clavum, F. oxysporum, F.

flocciferum, and *Fusarium* sp. (Table 1, Figure 1). The number of isolates recovered for

each species and their counties of origin are given in Table 1. Colony morphology (e.g.,

mycelial growth patterns) on PDA varied among isolates belonging to the same species,

particularly for *F. solani.* No PCR amplifications were achieved for two *F. oxysporum*

isolates identified in this study using FOV4 specific primers. Phylogenetic analysis of

tef-1α gene region placed these two *F. oxysporum* isolates within a clade containing

reference sequences of *F. oxysporum* f. sp. *asparagi* and *F. oxysporum* f. sp. *lactucae*

(Figure S1). *Fusarium solani* was the most frequently isolated species (39%), followed

by *F. equiseti* (22%), *F. proliferatum* and *F. clavum* (7% each), *F. incarnatum*, *F.*

verticillioides, *F. oxysporum*, and *Fusarium* sp. (5% each), and *F. flocciferum* (2%)

(Table 1).

Although no single county harbored all species isolated in this study, five species

were recovered from Sherman County: *F. incarnatum*, *F. equiseti*, *F. solani*, *F.*

verticillioides, and *Fusarium* sp. (Table 1). This was followed by Moore County (*F.*

incarnatum, *F. equiseti*, *F. solani*, and *F. proliferatum*) and Carson County (*F. equiseti*,

F. solani, *F. oxysporum*, and *F. flocciferum*), each with four species isolated (Table 1).

Two species were restricted to a single county. That is, *F. verticillioides* was isolated

only from Sherman County, while *F. flocciferum* was isolated from Carson County only

(Table 1).

Pathogenicity of *Fusarium* species on cotton

All *Fusarium* isolates tested were pathogenic on cotton; however, disease

severity was largely restricted to root discoloration (Figure 2). No disease was recorded

on non-inoculated control plants. *Fusarium* isolates were re-isolated from all inoculated

roots, both symptomatic and asymptomatic, plated on PDA (i.e., 100% re-isolation rate).

For the aerial parts, the highest disease severity (expressed as AUDPS) was

caused by *F. incarnatum* (31.4), followed by *F. solani* (12.5), while the lowest disease

severity was associated with *F. verticillioides* (7.1) (Figure 3A). Only *F. incarnatum*

caused significantly higher disease severity relative to non-inoculated control (Figure

3B, *P*=0.0137, EMMeans pairwise comparisons). Disease severity associated other

Fusarium species did not differ significantly from the non-inoculated control plants or

from each other (Figure 3B, *P*>0.05, EMMeans pairwise comparisons). DHARMa

diagnostics showed that the data met the model assumptions.

For roots, the Kruskal-Wallis test indicated that there was a significant difference

in disease severity among groups (Figure 4A, χ^2 = 17.58, df = 5, *P* = 0.01). The highest

disease severity, expressed as median rank, was caused by *F. proliferatum* and *F.*

clavum (2 each) (Figure 4A)*. Fusarium flocciferum, F. proliferatum,* and *F. clavum*

caused significantly higher disease severity relative to non-inoculated control (Figure

4A, $P=0.01$, $P=0.01$, $P=0.02$, respectively). However, no significant differences in

disease severity were recorded among different *Fusarium* species (Figure 4A, $P>0.05$).

Pathogenicity of *Fusarium* species on corn

All *Fusarium* isolates tested were pathogenic on corn, causing stalk rot, root

necrosis and brown to black discolorations; however, disease severity did not exceed

20–40%. The highest disease severity, expressed as median rank, was caused by *F.*

proliferatum, F. flocciferum, *F. incarnatum* and *F. verticillioides* (2 each) (Figure 4B). No

disease was recorded on non-inoculated control plants. Like cotton, *Fusarium* species

were re-isolated from all inoculated corn plants, regardless of whether they exhibited

symptoms. Differences in disease severity were not statistically significant among

Fusarium species or relative to non-inoculated control plants (Figure 4B, $P>0.05$).

Discussion

To the best of our knowledge, this study is the first to conduct a comprehensive

survey of the recently expanded cotton-growing areas in the Northern High Plains of

Texas and provided valuable insights into the diversity, distribution, and pathogenicity

of *Fusarium* species associated with cotton-corn rotation fields. A total of nine *Fusarium*

species were identified, with *F. solani* being the most frequently isolated and *F.*

flocciferum the least. Sherman County harboured the highest number of *Fusarium*

species. All tested species were pathogenic on cotton and corn during the pathogenicity

assays conducted under greenhouse conditions.

The high frequency of *F. solani* and *F. equiseti* isolation in the present study

aligns with prior findings from Arizona and Alabama, where they were the most

frequently isolated species from cotton roots and rhizosphere soil (Mao and Hu 2019;

Palmateer et al. 2002). Likewise, *F. solani* was reported as the most predominant

species, isolated from 97% of sorghum and corn fields across central and southeastern

United States (Leslie et al. 1990). Likewise, the least frequent isolation of *F. flocciferum*

in the present study is consistent with the findings from previous studies conducted in

France (Nguyen et al. 2024) and Belgium (Scauflaire et al. 2011), where it was among

the least frequently isolated species in corn agroecosystems. To date, there are no

reports of *F. flocciferum* as a widely distributed pathogen in the U.S. Cotton Belt. These

findings suggest that *F. solani* and *F. equiseti* are habitat generalists, being widely

distributed across the U.S. Cotton Belt, whereas *F. flocciferum* is mostly a habitat

specialist with a narrow distribution range.

The observation that colony morphology (e.g., mycelial growth patterns) on PDA

varied among isolates of the same species, particularly *F. solani,* is consist with finding

from a previous study (Mao and Hu 2019) and suggests that morphological

characteristics alone are insufficient for species identification, emphasizing the need for

sequence-based identification.

The variation in *Fusarium* species diversity observed across counties in this

study is likely caused by a combination of factors, including environmental conditions,

anthropogenic disturbance, variation in the number of fields sampled across counties,

soil types, cultural practices (e.g., crop rotation and tillage), and predisposing factors

such as the presence of nematodes or flooding events (Mai and Abawi 1987;

Steinkellner and Langer 2004). Differences in cotton acreage among counties may have

also contributed to variation in species diversity across counties. Specifically, the

highest species diversity in Sherman, Moore, and Carson counties can partly be

attributed to the substantial increase in cotton acreage between 2013 and 2018, during

which production grew by 301% to over 250,000 hectares (Ledbetter 2020; Marek and

Bordovsky 2006).

The significantly higher disease severity observed on cotton inoculated with *F.*

flocciferum, *F. proliferatum*, and *F. clavum*, and *F. solani* (relative to non-inoculated

control plants), suggests that these species may pose a potential threat to cotton and

corn production in the Northern High Plains. Among these, *F. proliferatum* and *F. solani*

have been previously identified as the most virulent species on cotton (Abdel-Latif et al.

2007; Palmateer et al. 2002). *Fusarium proliferatum* has also been reported as a

pathogen of cotton in New Mexico (Zhu et al. 2019). Disease severity on corn did not

differ significantly between inoculated and control plants or among *Fusarium* species;

however, *F. proliferatum* and *F. verticillioides* have been described as the most virulent

species on corn in previous studies (Okello et al. 2019; Wang et al. 2021). The

pathogenicity of *Fusarium* species on both cotton and corn suggests that they can

overwinter on the stubble of either crop and serve as a primary source of inoculum in

the next spring.

The observation of overall low disease severity on both cotton and corn may

partly be attributed to the application of inoculum after seedling emergence for both

cotton and corn, rather than the use of a root-dip inoculation method. Although the root-

dip method can result in increased disease severity, it does not closely mimic natural

infection under field conditions and may introduce stress during transplantation,

potentially predisposing seedlings to infection (Diaz et al. 2021; Schoeneweiss 1975).

The 100% re-isolation rate of *Fusarium* species from both symptomatic and

asymptomatic roots following pathogenicity trials in the present study supports the

previous findings that systemic colonisation can occur in the absence of visible disease

symptoms (Leslie et al. 1990). This implies that the pathogenicity of *Fusarium* species in

the field may need to be interpreted with caution, as *Fusarium* species may be living in

plants as endophytes or secondary invaders (Leslie et al. 1990). In the case of plant

tissues, this further underscores the importance of sampling both symptomatic and

asymptomatic plant tissues for accurately determining the diversity and distribution of

Fusarium species (Leslie et al. 1990). The present study likely provided an accurate

estimate of *Fusarium* species diversity and distribution because isolates were recovered

from soil samples, which minimized the bias associated with symptom-based tissue

sampling and isolation. Furthermore, a previous study did not find any significant

differences in the number of *Fusarium* species isolated from cotton roots and

rhizosphere soil (Mao and Hu 2019), suggesting that soil-based isolation approaches

can reliably capture species diversity.

This study provided baseline *Fusarium* species in the Northern High Plains of

Texas. The detection of both common (e.g., *F. solani* and *F. equiseti*) and less

frequently reported species (e.g., *F. clavum* and *F. flocciferum*) highlights the diversity

of *Fusarium* species in this region. The non-detection of the highly aggressive FOV4,

using FOV4 specific primers, in present study suggests that the pathogen may not yet

have been introduced into the Northern High Plains. Additionally, the phylogenetic

clustering of the two *F. oxysporum* isolates recovered in the present study with

reference sequences of *F. oxysporum* f. sp. *asparagi* and *F. oxysporum* f. sp. *lactucae*

suggest that they unlikely belong to the forma specialis `vasinfectum`. Therefore, there

is a probability that *F. oxysporum* isolates recovered in this study may have not been

pathogenic on cotton. However, given the known presence of FOV4 in nearby regions

such as El Paso, Hudspeth, and New Mexico, continued surveillance remains essential.

Crop rotation is unlikely to be effective since *Fusarium* species are cross-host

pathogenic and the region is primarily a rangeland where mainly susceptible crops such

as cotton, corn and wheat are grown for fodder. For species with broad niches,

containment and treatment (such as soil solarization) would potentially be required over

vast areas and therefore be impractical. Local containment may still be an option for

species with restricted niches. However, the use of resistant or tolerant cultivars should

be prioritized as a primary disease management strategy.

Since *Fusarium* species diversity can vary with the time of the year (Fulcher et al.

2021), future studies should aim to collect samples across multiple seasons.

Additionally, including counties not sampled in the present study will contribute to a

more comprehensive understanding of *Fusarium* species diversity and distribution

across the Northern High Plains of Texas.

Acknowledgements

We would like to acknowledge Dennis Coker, Marcel Fischbacher, JD Ragland, Kristy

Slough, Josie Strnad, and Laura Taylor for assistance in field selection, and Kristen

Tatro for greenhouse technical support. The work was supported by funding from

Cotton Inc. and Texas A&M AgriLife Research.

Author-Recommended Internet Resources

https://github.com/IhsanKhaliq/HighPlainsFusa

Literature Cited

Abdel-Latif, M., Shehata, Z., Galal, A., Hussein, E., and El-Samawaty, A. 2007.

Pathological studies on *Fusarium* species affecting cotton plants. J. Plant Prod.

32:4395-4408.

Atkinson, G. F. 1892. Some diseases of cotton. Ala. Agric. Exp. Stn. 41:1–65.

BASF. 2024. FM 757AXTP variety. https://agriculture.basf.us/crop-

protection//products/seeds/fibermax/fm-757axtp-variety.html (accessed June 1, 2025).

Bell, A. A., Gu, A., Olvey, J., Wagner, T. A., Tashpulatov, J. J., Prom, S., Quintana, J.,

Nichols, R. L., and Liu, J. 2019. Detection and characterization of *Fusarium oxysporum*

f. sp. *vasinfectum* VCG0114 (Race 4) isolates of diverse geographic origins. Plant Dis.

103:1998-2009.

BestPlaces. 2025. Panhandle, Tx, climate.

https://www.bestplaces.net/climate/city/texas/panhandle (accessed April 14, 2025).

Chernomor, O., Von Haeseler, A., and Minh, B. Q. 2016. Terrace aware data structure

for phylogenomic inference from supermatrices. Syst. Biol. 65:997-1008.

Cianchetta, A. N., Allen, T. W., Hutmacher, R. B., Kemerait, R. C., Kirkpatrick, T. L.,

Lawrence, G. W., Lawrence, K. S., Mueller, J. D., Nichols, R. L., and Olsen, M. W. 2015.

Survey of *Fusarium oxysporum* f. sp. *vasinfectum* in the United States. J. Cotton Sci.

19:328-336.

Colaizzi, P. D., Gowda, P. H., Marek, T., and Porter, D. O. 2009. Irrigation in the Texas

High Plains: A brief history and potential reductions in demand. Irrig. Drain. 58:257-274.

Davis, R., Colyer, P., Rothrock, C., and Kochman, J. 2006. Fusarium wilt of cotton:

population diversity and implications for management. Plant Dis. 90:692-703.

Diaz, J., Garcia, J., Lara, C., Hutmacher, R. B., Ulloa, M., Nichols, R. L., and Ellis, M. L.

2021. Characterization of current *Fusarium oxysporum* f. sp. *vasinfectum* isolates from

cotton in the San Joaquin Valley of California and Lower Valley El Paso, Texas. Plant

Dis. 105:1898-1911.

Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 32:1792-1797.

Fulcher, M. R., Oladipo, E. D., Winans, J. B., Quan, M., and Bergstrom, G. C. 2021.

Structure and diversity of *Fusarium* communities inhabiting non-cultivated grass

inflorescences in New York State. Can. J. Plant Pathol. 43:48-55.

Glover, J. P., Medrano, E. G., Isakeit, T., and Brewer, M. J. 2020. Transmission of cotton

seed and boll rotting bacteria by the verde plant bug (*Hemiptera*: *Miridae*). J. Econ.

Entomol. 113:793-799.

Halpern, H., Bell, A., Wagner, T., Liu, J., Nichols, R., Olvey, J., Woodward, J., Sanogo,

S., Jones, C., and Chan, C. 2018. First report of Fusarium wilt of cotton caused by

Fusarium oxysporum f. sp. *vasinfectum* race 4 in Texas, USA. Plant Dis. 102:446-446.

Hartig, F. 2020. Residual diagnostics for hierarchical (multi-level/mixed) regression

models. R package version 0.4.6. https://CRAN.R-project.org/package=DHARMa

(accessed May 15, 2025).

Hothorn, T. 2020. Multcomp: simultaneous inference in general parametric models. R

package version 1.4.25.

http://ftp5.gwdg.de/pub/misc/cran/web/packages/multcomp/multcomp.pdf (accessed

May 20, 2025).

Howell, C. 2002. Cotton seedling preemergence damping-off incited by *Rhizopus*

oryzae and *Pythium* spp. and its biological control with *Trichoderma* spp.

Phytopathology 92:177-180.

Hutmacher, R. B., Ulloa, M., Wright, S. D., Campbell, B. T., Percy, R., Wallace, T.,

Myers, G., Bourland, F., Weaver, D., and Chee, P. 2013. Elite upland cotton germplasm-

pool assessment of *Fusarium* wilt resistance in California. Agron. J. 105:1635-1644.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., and Jermiin, L. S.

2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat.

Methods 14:587-589.

Kim, Y., Hutmacher, R., and Davis, R. 2005. Characterization of California isolates of

Fusarium oxysporum f. sp. *vasinfectum*. Plant Dis. 89:366-372.

Kimmel Jr, T. M., Nielsen-Gammon, J., Rose, B., and Mogil, H. M. 2016. The weather

and climate of Texas: A big state with big extremes. Weatherwise 69:25-33.

Kuznetsova, A., Brockhoff, P., Christensen, R., and Jensen, S. 2022. lmerTest: tests in

linear mixed effects models. R package version 3.1.3. https://cran.r-

project.org/web/packages/lmerTest/lmerTest.pdf (accessed May 14, 2025).

Ledbetter, K. 2020. Northern High Plains cotton acres more than double in last 5 years.

https://www.cottongrower.com/cotton-production/planting/northern-high-plains-cotton-

acres-more-than-double-in-last-5-years/ (accessed April 25, 2025).

Lenth, R. 2021. EMMEANS: estimated marginal means, aka Least-Squares Means. R

package version 1.8.5. https://github.com/rvlenth/emmeans (accessed April 1, 2025).

Leslie, J. F., Pearson, C., Nelson, P. E., and Toussoun, T. 1990. *Fusarium* spp. from

corn, sorghum, and soybean fields in the central and eastern United States.

Phytopathology 80:343-350.

Li, B., Liang, J., Zhu, Y., Wang, Y., and Jiao, Z. 2021. Epidemiological analysis and

management strategies of *Fusarium* head blight of wheat. Curr. Biotechnol. 11:647-652.

Liu, Y. J., Whelen, S., and Hall, B. D. 1999. Phylogenetic relationships among

ascomycetes: evidence from an RNA polymerse II subunit. Mol Biol Evol. 16:1799-1808.

Mai, W., and Abawi, G. 1987. Interactions among root-knot nematodes and Fusarium

wilt fungi on host plants. Annu. Rev. Phytopathol. 25:317-338.

Mao, Y., and Hu, J. 2019. Survey of *Fusarium oxysporum* f. sp. *vasinfectum* race 4 in

Arizona cotton. Pages 178-184 in: 2019 Proceedings of the Beltwide Cotton

Conferences, The National Cotton Council of America, Memphis, TN.

Marek, T., and Bordovsky, D. 2006. Performance of ten cotton varieties in the Northern

Texas High Plains. Tex. J. Agric. Nat. Resour. 19:48-61.

Medrano, E., Esquivel, J., and Bell, A. 2007. Transmission of cotton seed and boll

rotting bacteria by the southern green stink bug (*Nezara viridula* L.). J. Appl. Microbiol.

103:436-444.

Mendiburu, F. 2023. agricolae: statistical procedures for agricultural research. R

pacakge version 1.3-7. https://cran.r-project.org/web/packages/agricolae/index.html

(accessed April 12, 2025).

Minton, E. B., and Garber, R. H. 1983. Controlling the seedling complex of cotton. Plant

Dis. 67:115-117.

Nash, S. M., and Snyder, W. C. 1962. Quantitative estimations by plate counts of

propagules of the bean root rot *Fusarium* in field soils. Phytopathology 52:567-572.

National IPM Database. 2001. Crop profile for corn in Texas. National IPM Database.

https://ipmdata.ipmcenters.org/documents/cropprofiles/TXcorn.pdf (accessed May 20,

2025). .

Nguyen, L. T., Schmidt, H. A., Von Haeseler, A., and Minh, B. Q. 2015. IQ-TREE: a fast

and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol

Biol Evol. 32:268-274.

Nguyen, T. B. H., Henri-Sanvoisin, A., Coton, M., Le Floch, G., and Picot, A. 2024. Shifts

in *Fusarium* communities and Mycotoxins in maize Residues, soils, and wheat Grains

throughout the wheat cycle: implications for *Fusarium* Head Blight epidemiology.

Microorganisms 12:1783.

NOAA. 2021. National Oceanic and Atmospheric Administration. National Centers for

Environmental Information. U.S. climate normals quick access 1991-2020.

https://www.ncei.noaa.gov/access/us-climate-normals/ (accessed June 15, 2025).

O'Donnell, K., Kistler, H. C., Cigelnik, E., and Ploetz, R. C. 1998. Multiple evolutionary

origins of the fungus causing Panama disease of banana: concordant evidence from

nuclear and mitochondrial gene genealogies. Proc. Natl. Acad. Sci. U. S. A. 95:2044-

2049.

Okello, P. N., Petrović, K., Kontz, B., and Mathew, F. M. 2019. Eight species of *Fusarium*

cause root rot of corn (Zea mays) in South Dakota. Plant Health Prog. 20:38-43.

Palmateer, A., McLean, K., and Morgan-Jones, G. 2002. Relative recovery and

pathogenecity of *Fusarium* species on cotton. in: 2002 Proceedings of the Beltwide

Cotton Conferences, Atlanta, Georgia.

Parikh, L., Kodati, S., Eskelson, M., and Adesemoye, A. 2018. Identification and

pathogenicity of *Fusarium* spp. in row crops in Nebraska. Crop Prot. 108:120-127.

Patil, I. 2021. Visualizations with statistical details: The'ggstatsplot'approach. Journal of

Open Source Software 6:3167.

Sangalang, A., Burgess, L., Backhouse, D., Duff, J., and Wurst, M. 1995.

Mycogeography of *Fusarium* species in soils from tropical, arid and mediterranean

regions of Australia. Mycological Research 99:523-528.

Scauflaire, J., Mahieu, O., Louvieaux, J., Foucart, G., Renard, F., and Munaut, F. 2011.

Biodiversity of *Fusarium* species in ears and stalks of maize plants in Belgium. Eur. J.

Plant Pathol. 131:59-66.

Schoeneweiss, D. F. 1975. Predisposition, stress, and plant disease. Annu. Rev.

Phytopathol. 13:193-211.

Shu, X., Wang, Y., Wang, Y., Ma, Y., Men, M., Zheng, Y., Xue, C., Peng, Z., and Noulas,

C. 2021. Response of soil N_2O emission and nitrogen utilization to organic matter in the

wheat and maize rotation system. Sci. Rep. 11:4396.

Simko, I., and Piepho, H.-P. 2012. The area under the disease progress stairs:

calculation, advantage, and application. Phytopathology 102:381-389.

Steinkellner, S., and Langer, I. 2004. Impact of tillage on the incidence of *Fusarium* spp.

in soil. Plant Soil. 267:13-22.

TCFA. 2015. Cattle feeders resource guide. http://www.tcfa.org/ (accessed May 3,

2025).

Thayer, A. W., and McCarl, B. A. 2018. Water Depletion, climate change, and the Texas

High Plains: a model on the future of irrigation dependent agriculture. AgEcon Search.

https://ageconsearch.umn.edu/record/266615/files/SAEASubmission_ThayerMcCarl.pdf

(accessed June 2, 2025).

USDA Economic Research Service. 2022. Cotton and wool - cotton sector at a glance

https://www.ers.usda.gov/topics/crops/cotton-and-wool/cotton-sector-at-a-glance

(accessed May 20, 2025).

USDA Foreign Agricultural Service. 2024. Production - cotton. U.S. Department of

Agriculture, Foreign Agricultural Service.

https://www.fas.usda.gov/data/production/commodity/2631000 (accessed June 20,

2025).

USDA National Agricultural Statistics Service. 2025. Cattle on feed.

https://www.nass.usda.gov (accessed May 20, 2025).

Wagner, T. A., Bell, A. A., Castles, Z. A., Ali, A., Flores, O., and Liu, J. 2022. Detection,

genotyping and virulence characterization of *Fusarium* wilt race 4 (VCG0114) causing

cotton wilt in three Texas fields. J. Plant Pathol. 170:492-503.

Wang, B., Brubaker, C. L., Summerell, B. A., Thrall, P. H., and Burdon, J. J. 2010. Local

origin of two vegetative compatibility groups of *Fusarium oxysporum* f. sp. *vasinfectum*

in Australia. Evolutionary Applications 3:505-524.

Wang, W., Wang, B., Sun, X., Qi, X., Zhao, C., Chang, X., Khaskheli, M. I., and Gong,

G. 2021. Symptoms and pathogens diversity of corn *Fusarium* sheath rot in Sichuan

province, China. Sci. Rep. 11:2835.

Ward, T. J., Clear, R. M., Rooney, A. P., O'Donnell, K., Gaba, D., Patrick, S., Starkey, D.

E., Gilbert, J., Geiser, D. M., and Nowicki, T. W. 2008. An adaptive evolutionary shift in

Fusarium head blight pathogen populations is driving the rapid spread of more toxigenic

Fusarium graminearum in North America. Fungal Genet. Biol. 45:473-484.

Watkins, G. 1981. Compendium of Cotton Disease. American Phytopathological

Society. St. Paul, MN, USA.

White, T. J., Bruns, T., Lee, S., and Taylor, J. 1990. Amplification and direct sequencing

of fungal ribosomal RNA genes for phylogenetics. Pages 315-322 in: PCR protocols: a

guide to methods and applications. M. Innis, D. Gelfand, J. Sninsky, T. White and eds.

Academic Press, San Diego, California.

Xi, K., Shan, L., Yang, Y., Zhang, G., Zhang, J., and Guo, W. 2021. Species diversity

and chemotypes of *Fusarium* species associated with maize stalk rot in Yunnan

province of southwest China. Front. Microbiol. 12:652062.

Xue, Q., Marek, T. H., Xu, W., and Bell, J. 2017. Irrigated corn production and

management in the Texas High Plains. J. Contemp. Water Res. Educ. 162:31-41.

Yadav, S. L., Ahir, R., Yadav, A., and Meena, S. 2017. Inoculation techniques of

Fusarium Oxysporum on onion. Chem. Rev. Lett. 6:1059-1062.

Zaki, S. A., Ouf, S. A., Aly, A. A., and Abd-Elsalam, K. A. 2021. Fungi involved in

damping-off of cotton seedlings and their differential pathogenicity on two cotton

cultivars. Egypt. J. Bot. 61:911-921.

Zhang, H., Brankovics, B., Luo, W., Xu, J., Xu, J., Guo, C., Guo, J., Jin, S., Chen, W.,

and Feng, J. 2016. Crops are a main driver for species diversity and the toxigenic

potential of *Fusarium* isolates in maize ears in China. World Mycotoxin J. 9:701-716.

Zhang, J., Abdelraheem, A., Zhu, Y., Wheeler, T. A., Dever, J. K., Elkins-Arce, H.,

Nichols, R., and Wedegaertner, T. 2020. Pedigree selection under field conditions within

Acala 1517-08 and its glandless derivatives for development of cotton resistant to

Fusarium wilt caused by *Fusarium oxysporum* f. sp. *vasinfectum* race 4. Euphytica

216:1-21.

Zhu, Y., Abdelraheem, A., Sanogo, S., Wedegaertner, T., Nichols, R., and Zhang, J.

2019. First report of *Fusarium oxysporum* f. sp. *vasinfectum* race 4 causing Fusarium

wilt of cotton in New Mexico, USA. Plant Dis. 104:2679-2679.

Zhu, Y., Abdelraheem, A., Lujan, P., Idowu, J., Sullivan, P., Nichols, R., Wedegaertner,

T., and Zhang, J. 2021. Detection and characterization of *Fusarium* wilt (*Fusarium*

oxysporum f. sp. *vasinfectum*) race 4 causing *Fusarium* wilt of cotton seedlings in New

Mexico. Plant Dis. 105:3353-3367.

Zivkovic, S., and Hudson, d. 2012. Carbon sequestration and carbon management

policy effects on production agriculture in the Texas High Plains. Environ. Manag.

Sustain. Dev. 3:44-60.

Table 1 Isolate identity, collection date, county of origin, host crop history, and GenBank accession numbers for *Fusarium* species isolated in this study.

Isolate	Organism	County	Previous crop	Current crop	Sampling date	GenBank Accession number		
						tef-1α	ITS	RPB2
F1	*F. incarnatum*	Sharman	Corn	Cotton	2024/08/22	PV699158		
F10	*F. incarnatum*	Moore	Corn	Cotton	2024/08/21		PV670408	
F2	*F. equiseti*	Carson	Corn	Cotton	2024/10/24		PV670407	
F14	*F. equiseti*	Carson	Corn	Cotton	2024/10/24	PV699144		
F15	*F. equiseti*	Moore	Corn	Cotton	2024/08/21	PV699145		
F18	*F. equiseti*	Hartley	Corn	Cotton	2024/09/26	PV699148		
F20	*F. equiseti*	Sharman	Corn	Cotton	2024/08/22	PV699152		
F25	*F. equiseti*	Hansford	Cotton	Wheat	2024/05/01	PV699157		
F38	*F. equiseti*	Hutchinson	Fallow	Cotton	2024/05/21	PV717850		
F40	*F. equiseti*	Sharman	Corn	Cotton	2024/08/22	PV717843		
F41	*F. equiseti*	Sharman	Corn	Cotton	2024/08/22	PV717844		
F3	*F. solani*	Deaf Smith	Fallow	Cotton	2024/10/02	PV699137		
F4	*F. solani*	Deaf Smith	Corn	Cotton	2024/10/02	PV699138		
F7	*F. solani*	Deaf Smith	Corn	Cotton	2024/10/02	PV699140		
F8	*F. solani*	Moore	Corn	Cotton	2024/08/21	PV699141		
F9	*F. solani*	Moore	Corn	Cotton	2024/08/21	PV699142		
F19	*F. solani*	Carson	Corn	Cotton	2024/10/24	PV699149		
F21	*F. solani*	Deaf Smith	Corn	Cotton	2024/10/02	PV699153		
F22	*F. solani*	Carson	Corn	Cotton	2024/10/24	PV699154		
F23	*F. solani*	Deaf Smith	Wheat	Cotton	2024/10/02	PV699155		
F30	*F. solani*	Moore	Corn	Cotton	2024/08/21	PV717836		
F31	*F. solani*	Hutchinson	Fallow	Cotton	2024/05/21	PV717837		
F32	*F. solani*	Sharman	Corn	Cotton	2024/08/22	PV717838		
F33	*F. solani*	Dallam	Corn	Cotton	2024/09/26	PV717839		

Isolate	Organism	County	Previous crop	Current crop	Sampling date	tef-1α	ITS	RPB2
F35	*F. solani*	Randall	Corn	Cotton	2024/08/22	PV717840		
F36	*F. solani*	Hartley	Corn	Cotton	2024/09/26	PV717841		
F37	*F. solani*	Carson	Corn	Cotton	2024/10/24	PV717842		
F5	*F. verticillioides*	Sharman	Corn	Cotton	2024/08/22	PV699139		
F42	*F. verticillioides*	Sharman	Corn	Cotton	2024/08/22	PV717845		
F12	*F. proliferatum*	Randall	Corn/Sorghum	Cotton	2024/08/22		PV670409	
F16	*F. proliferatum*	Randall	Corn/Sorghum	Cotton	2024/08/22	PV699146		
F29	*F. proliferatum*	Moore	Corn	Cotton	2024/05/01	PV717848		
F13	*F. clavum*	Hansford	Cotton	Wheat	2024/05/01	PV699143		
F26	*F. clavum*	Randall	Corn/Sorghum	Cotton	2024/05/01	PV717846		
F27	*F. clavum*	Hartley	Corn	Cotton	2024/09/26	PV717835		
F17	*F. oxysporum*	Randall	Corn/Sorghum	Cotton	2024/08/22	PV699147	PV731270	PV699150
F34	*F. oxysporum*	Carson	Corn	Cotton	2024/10/24	PV717849	PV731268	PV699151
F39	*F. flocciferum*	Carson	Corn	Cotton	2024/10/24	PV717851		
F24	*Fusarium sp.*	Sharman	Corn	Cotton	2024/08/22	PV699156		
F28	*Fusarium sp.*	Hansford	Wheat	Cotton	2024/05/01	PV717847		

697 0.04

698 **Figure** 1 Phylogenetic comparison of *Fusarium* species isolated from cotton-corn

699 rotation fields in the Texas High Plains (underlined) in relation to closely related

700 reference species based on the translation elongation factor 1-α (tef-1α) gene region.

0 **1** **2** **3** **4** **5**

704 **Figure 2** Damage on roots of cotton seedlings inoculated with *Fusarium* species. Whole

705 root systems were rated visually using a 0–5 rating scale. Left to right: 0 - no symptoms;

706 1 - <10% of roots discolored; 2 - 10–30% of roots discolored; 3 - 31–50% of roots

707 discolored; 4 - 51–80% of roots discolored; and 5 - 81–100% of roots discolored.

Figure 3. Comparison of the area under the disease progress stairs (AUDPS) based on

severity ratings recorded on aerial parts of cotton seedlings across all four disease

assessments: (A) AUDPS for each *Fusarium* species; (B) a linear mixed effect model fit

showing estimated marginal means using the emmeans R package. Different letters (a,

b) indicate significant differences among treatments at $P{\leq}0.05$.

Figure 4 Comparison of disease severity of different *Fusarium* species on roots of (A)

cotton and (B) corn. Differences in disease severity were compared using a non-

parametric Kruskal–Wallis test.

Table S1 Disease rating scales used for assessing disease severity of *Fusarium*

species in cotton and corn. For cotton, disease severity was recorded on both aerial

parts and roots, whereas for corn, it was recorded on roots only.

Crop	Plant part	Rating	Description	Range (%)
Cotton	**Aerial parts**	0	Healthy	-
		1	Cotyledons wilted only	-
		2	≤50% true leaves wilted	-
		3	>50% but ≤90% true leaves wilted	-
		4	All leaves wilted or plant death	-
	Roots	0	no symptoms	-
		1	< 10% of roots discolored	1–9
		2	10–30% of roots discolored	10–30
		3	31–50% of roots discolored	31–50
		4	51–80% of roots discolored	51–80
		5	81–100% of roots discolored	81–100
Corn	**Roots**	1	No visible symptoms, 0% infection, and normal development	-
		2	Small speck of lesions near the crown and/or superficial root infection. Infection is <10%, and development is normal	1–9
		3	Superficial lesions visible near the crown or stem. Infection is 10–20%, and development is normal	10–20
		4	Deep lesion has advanced from crown and lower roots to the middle stem. Infection is 20–40%, and development is normal	21–40
		5	Deep lesion visible through approximately half the length of the stem and root. Infection is 40–60%, and plant development is affected	40–60

Crop	Plant part	Rating	Description	Range (%)
		6	Deep lesions on most parts of the stem and root, reduced root growth, significantly reduced plant development, and stunting are observed. Infection is 60–85%	60–85
		7	Severe deep lesions covering almost all parts of the stem and root, significantly reduced root growth and plant development, and extensive stunting are observed. Infection is 85–95%	85–95
		8	Seedling barely emerged and could not grow, scanty root but no emergence, or seed is dead (no emergence and no roots). Infection coverage of seedling is >95%	95–100

0.003

Fig. S1 *Phylogenetic* relationships of *Fusarium oxysporum* isolates collected in this

study (highlighted in blue) compared with reference isolates of *F. oxysporum* f.

sp. *vasinfectum* races 1, 2, 3, 4, 6, and 8 (FOV1 to FOV8) and other closely related

species, based on sequences of the translation elongation factor 1-α (tef1-α) gene

region.