Preclinical Study of Cannflavins A and B Action Against Glioblastoma Cells

Jennifer Holborn [a], Hannah Robeson [a,1], Ellis Chartley [a,1], Tiana Gluscevic [a,b,1], Adina Borenstein [a], Colby Perrin [a], Begüm Alural [a], Himain Perera [a], Tariq A. Akhtar [a], Nina Jones [a], Jasmin Lalonde [a,*]

[a] Department of Molecular and Cellular Biology, University of Guelph, 50 Stone Road E, Guelph, ON N1G 2W1, Canada

[b] Canurta Therapeutics, 2425 Matheson Blvd. E, Suite 807, Mississauga, ON, L4W 5K4, Canada

[1] These authors contributed equally to this study.

* Corresponding author: **Jasmin Lalonde**

Email address: jlalon07@uoguelph.ca

Phone: (519) 824-4120 x. 54706

ORCID: 0000-0002-6797-1894 (J. Lalonde)

Keywords: Cannflavins, Cerebral organoids, Flavonoids, Glioblastoma multiform, Invasion, Migration

Abbreviations: AKT, Protein kinase B; BDNF, Brain-Derived Neurotrophic Factor; BSA,

Bovine Serum Albumin; DMEM, Dulbecco's Modified Eagle's Medium; DMSO,

Dimethylsulfoxide; FBS, Fetal Bovine Serum; GBM, Glioblastoma multiform; GFP,

Green Fluorescent Protein; iPSC, induced Pluripotent Stem Cell; MAPK, Mitogen-

Activated Protein Kinase; mTOR, mammalian target of rapamycin; NF-κB, Nuclear factor-

κB; PBS, Phosphate Buffered Saline; RTK, Receptor Tyrosine Kinase; TMZ,

Temozolomide; TrkB, Tropomyosin receptor kinase B.

ABSTRACT

Background: Flavonoids represent a large group of naturally occurring polyphenolic

compounds, many of which have been found to produce valuable biological outcomes,

including action against cancer cells. Glioblastoma multiforme (GBM) is an aggressive

type of brain tumor that is associated with a poor prognosis and has limited treatment

options. Previous findings suggest that cannflavins can produce promising effects for

pancreatic and bladder cancers, but the efficacy of these phytochemicals in attacking brain

tumour cells remains unknown. Our study evaluates the potential of cannflavin A and

cannflavin B against GBM cells using a range of *in vitro* approaches.

Results: We conducted experiments using A-172 and U-87 GBM lines to assess the impact

of cannflavins A and B on cell viability, cycle, migration, and invasion capacity. Our

results revealed a consistent dose-dependent decrease in cell viability in both lines after the

addition of cannflavin B to the culture media. Interestingly, we found that chrysoeriol (the

non-prenylated synthesis precursor of cannflavins in the *Cannabis sativa* plant) only has a

limited adverse effect on survival using the same approach, while techtochrysin (an O-

methylated flavone) has none. Using time-lapse live-cell imaging and a scratch assay, we

also show that cannflavins can inhibit tumor cell migration at concentrations below those

that produce significant cell death. Finally, we found that cannflavin B exhibits anti-

migratory and anti-invasive properties in transwell and tumorsphere assays, underscoring

its multifaceted therapeutic potential.

Conclusion: These findings suggest that cannflavin B holds promise as a therapeutic agent

in the treatment of GBM.

HIGHLIGHTS (3-5 BULLET POINTS)

• Cannflavin B but not cannflavin A limits the viability of A-172 and U-87 GBM cells

in a concentration- and time-dependent manner.

• Using time-lapse live-cell imaging and a scratch assay, we provide evidence that

cannflavins A and B can inhibit GBM cell migration at concentrations below those

that result in a significant reduction in cell viability.

• Cannflavin B exhibits robust anti-migratory and anti-invasive properties in transwell

and tumorsphere assays, underscoring its multifaceted therapeutic potential.

• Although previous work had reported the capacity of cannflavins A and B to interfere

with TrkB and downstream the MAPK/AKT signaling pathways in mouse primary

neurons, the effect of those molecules is limited in GBMs, suggesting that other targets

are engaged to produce the cellular effects.

64 **INTRODUCTION**

65 Flavonoids are naturally occurring polyphenolic molecules that accumulate in dietary items

66 such as fruits, vegetables, and teas. This broad group of phytochemicals has received

67 considerable research attention for over 25 years due to their beneficial anti-inflammatory

68 and anti-oxidative properties (Panche *et al.*, 2016). Many flavonoids have also attracted

69 interest in cancer research due to their potential therapeutic applications. For instance,

70 kaempferol has been found to induce G2/M cell cycle arrest as well as reduce migration

71 and invasion in breast, liver, ovarian, lung, and brain cancers (Gao et al., 2018; Jo et al.,

72 2015; Li et al., 2017; Zhu et al., 2019). Apigenin and luteolin, for their part, have shown

73 promise as anti-cancer agents for breast, prostate, lung, and brain subtypes by suppressing

74 the mitogen-activated protein kinase (MAPK) and Protein kinase B (AKT) signaling

75 pathways that control cell proliferation and survival (Cheng et al., 2013; Erdogan et al.,

76 2016; Huang et al., 2019; Tseng et al., 2017). Finally, quercetin, one of the most consumed

77 dietary flavonoids, induces G2/M phase arrest and significantly increases tumor suppressor

78 protein p53 expression in MDA-MB-453 breast cancer cells (Choi et al., 2008). It also

79 limits angiogenesis and metastasis of other cancer cell types through changes in

80 MAPK/AKT/mTOR signaling pathways (Balakrishnan et al., 2016; Gao et al., 2012; Kim

81 et al., 2019; Lee et al., 2015; Pratheeshkumar et al., 2012). Taken together, these examples

82 support searching for additional flavonoids that also produce robust anti-cancer effects.

83  Cannflavins are prenylated and highly lipophilic metabolites first purified 40 years ago

84 from *Cannabis sativa (C. sativa)* plant extracts (Barrett et al., 1985). As illustrated in

85 **Supplementary Figure 1**, the cannflavin family includes five known members, which are

86 cannflavin A, cannflavin B, cannflavin C, isocannflavin B (also known as FBL-03G), and

8-prenylcannflavin B (also known as dorsmanin D). Several of these molecules have been

shown to produce promising effects applicable to various clinical conditions. For instance,

studies revealed the anti-inflammatory properties of cannflavins A and B through their

ability to inhibit pro-inflammatory mediators and the production of prostaglandin E_2 at a

rate thirty times more potent than aspirin (Barrett et al., 1985; Werz et al., 2014). Consistent

with their anti-inflammatory activity, Eggers and colleagues (2019) have reported

neuroprotective action of cannflavin A in neuronal PC12 cells against amyloid β-induced

cell death by reducing $A\beta_1$-42 aggregation and fibril formation.

Beyond studies concerning inflammation, other research suggests that specific

cannflavin molecules exhibit desirable anti-cancer effects that may involve other pathways.

For instance, isocannflavin B was found to improve survival in preclinical mouse models

of metastatic pancreatic cancer by increasing apoptosis of tumor cells (Moreau et al., 2019),

and cannflavin A has been reported to synergize with agents commonly used to treat

bladder cancer at specific concentrations (Tomko et al., 2022). Despite those promising

results, whether cannflavins can effectively attack brain cancer cells has not been directly

studied.

Glioblastoma multiforme (GBM) is the most aggressive and common primary brain

tumor in adults, characterized by rapid growth and diffuse infiltration into surrounding

healthy brain tissues (Tilak et al., 2021a). The prognosis for GBM patients is extremely

poor, with a median survival of approximately 15 months despite aggressive treatment

(Johnson and O'Neill, 2012; Ostrom et al., 2020). The standard of care for GBM is

maximum safe surgical resection followed by a combination of radiation therapy and

temozolomide (TMZ) chemotherapy, a DNA alkylating agent that induces cell death

through DNA lesions (Stupp et al., 2005; Stupp et al., 2009; Fisher and Adamson, 2021).

However, the inability to fully resect the tumor with surgery, combined with the persistence

of TMZ-resistant cancer cells capable of recapitulating tumors, typically leads to disease

relapse within just a few months (Weller et al., 2013; Birzu et al., 2020). Therefore, there

is a pressing need to identify new agents that could improve patient prognosis.

In this study, we conducted experiments using the A-172 and U-87 GBM cell lines to

assess the effects of cannflavins A and B on cell viability, migration, and invasion. Our

work revealed a decrease in GBM cell viability with increasing concentrations of

cannflavin B. Additionally, analysis of cell migration using a time-lapse live-cell scratch

assay revealed the potential of cannflavins to impede tumor dissemination at concentrations

below those inducing cell death. Specifically, cannflavin B exhibited robust anti-migratory

and anti-invasive properties, as evidenced by transwell and tumorsphere invasion assays,

highlighting its multifaceted therapeutic effects. Finally, as our previous work found

interference of TrkB signaling by cannflavin A and cannflavin B in mouse primary neurons

(Holborn et al., 2023), we also evaluated whether the action of cannflavins on GBM cells

could be tied to reduced activity of this receptor tyrosine kinase (RTK). Together, our

findings underscore the promising anti-cancer properties of cannflavin B against

immortalized GBM cells, suggesting its potential role as a novel therapeutic agent.

MATERIALS AND METHODS

Cell culture and reagents

Malignant male glioma cell lines A-172 and U-87 MG (U-87) were obtained from the

American Type Culture Collection (ATCC, USA, Cat No. CRL-1620 and HTB-14,

respectively). Cells were cultured in DMEM (Gibco, USA, Cat No. 11965-092)

supplemented with 10% FBS (Gibco, Cat No. 12483-020) and 0.5%

penicillin/streptomycin (Gibco, Cat No. 155140-122), at 37°C and 5% CO_2. For treatment

with cannflavins and other tested agents, the molecules were dissolved in DMSO and

applied directly to the culture media.

The BIONi010-A cell line (European Bank for induced Pluripotent Stem Cells,

ECACC Cat No. 66540022) used to prepare cerebral organoids was purchased from Sigma-

Aldrich (St-Louis, MO, USA) and cultured in ExCellerate iPSC Expansion Medium (R&D

Systems, Cat No. CCM036) according to the manufacturer's instructions.

Antibodies and pharmacological compounds

The antibodies recognizing phosphorylated p44/p42 MAPK (Cat No. 4370), AKT (Cat No.

4691), phosphorylated AKT Thr 308 (AKT T308, Cat No. 2965), and phosphorylated AKT

Ser 473 (AKT S473, Cat No. 4060) were acquired from Cell Signaling Technology

(Beverly, MA, USA). The antibody detecting phosphorylated TrkB Tyr705 (TrkB Y705,

Cat No. ab229908) was from Abcam (Waltham, MA, USA), the antibody recognizing p42

MAPK (Erk2, sc-1647) was from Santa Cruz Biotechnology (Santa Cruz, CA), the

GAPDH (Cat No. AB2302) antibody was from Sigma-Aldrich, and the Tubulin β 3 (TUJ1,

Cat No. 801202) antibody was from BioLegend (San Diego, CA, USA). Finally, the GFP

(Cat No. 10262) and cross-absorbed secondary antibodies (horseradish peroxidase- and

fluorophore-conjugated) were from Invitrogen.

Cannflavin A (Cat No. C-291001) and cannflavin B (Cat No. C-291003) were

purchased from TLC Pharmaceutical Standards (Newmarket, ON, Canada). ANA-12 (Cat

No. SML0209) was acquired from Sigma-Aldrich, and chrysoeriol (Cat No. 491-71-4)

from Cayman Chemical (Ann Arbor, MI, USA).

Techtochrysin was biosynthesized as follows: The *O. basilicum* O-methyl transferase

ObFOMT1 (Accession number JQ653275) open reading frame was synthesized by Twist

Bioscience (San Francisco, CA, USA) in the *pET28a* vector system between the *NdeL* and

XhoI sites. This system inserted an N-terminal hexahistidine tag into the coding sequence.

The construct was then introduced into *E. coli* BL21-CodonPlus (DE3)-RIPL. Bacterial

cells harboring the expression vector were cultured in LB media at 37°C to an OD_{600} of

0.6. Isopropyl-β-D-thiogalactoside was added to a final concentration of 0.2 mM, and

cultures were further incubated at 16°C for an additional 16 hours to allow for protein

expression. Thereafter, *E. coli* cells were collected by centrifugation, resuspended in Buffer

A (20 mM Tris-HCl, 500 mM KCl, pH 8.0) and lysed using an Emulsiflex. Cell debris and

remaining intact cells were removed from the solution by centrifugation (20,000 x *g*, 20

minutes at 4°C) and the supernatant was applied to a 1 mL HisTrap FF column (Cytiva,

Marlborough, MA) equilibrated in Buffer B (20 mM Tris-HCl, 500 mM KCl, 20 mM

imidazole, pH 8.0). Proteins bound to the Ni^{2+} affinity matrix were washed with fifteen

column volumes of Buffer B, eluted with one column volume of Buffer C (20 mM Tris-

HCl, 500 mM KCl, 200 mM imidazole, pH 8.0), and then immediately desalted on PD-10

columns (Cytiva) equilibrated with Buffer D (50 mM Tris-HCl, 5 mM $MgCl_2$, pH 8.0, 10%

glycerol). The concentration of recombinant ObFOMT1 was determined by Bradford

Assay using BSA as a standard, and 1 mg was used per enzymatic assay to produce

techtochrysin. Precisely, purified recombinant ObFOMT1 was mixed in 10 mL of assay

solution (1 mM chrysin, 10% w/v hydroxypropyl-β-cyclodextrin, 2 mM S-

adenosylmethionine, and 5 mM $MgCl_2$ in 50 mM Tris-HCl, pH 8.0) and incubated with

shaking for 16 hours at 15°C. Reagents for the assay solution were obtained from Xi'an B-

Thriving I/E Co., Ltd (Xi'an, Shaanxi, China). The enzymatic reaction was terminated by

adding 1 mL of 10% formic acid. Prenylated bibenzyl products were extracted with three

volumes of ethyl acetate three times, evaporated to dryness with gaseous N, and

resuspended in 6 mL of methanol. The sample was chromatographed on an Infinity Lab

Poroshell 120 EC-C18 2.7 μM (ODS2 9.4 x 150 mm) column and resolved by an Agilent

1260 infinity HPLC using a linear gradient with solvent A (45% methanol, 0.1% formic

acid in water) and solvent B (95% methanol, 0.1% formic acid in water) at a flow rate of

2.5 mL/min heated at 30°C. The gradient started at 60% of solvent B and increased to 70%

in 7 minutes, then increased to 100% of solvent B in 1 minute and remained isocratic for 2

minutes. Techtochrysin was detected by absorption at 340 nm and was collected from the

HPLC at 9.3 minutes. Finally, the eluate was dried under gaseous nitrogen, lyophilized,

and techtochrysin quantified relative to a standard curve generated with chrysin. Product

identity was confirmed based on mass spectral fragmentation patterns.

Cell viability assay

Cell viability was assessed using a resazurin reduction assay, as described previously

(Holborn et al., 2023). Briefly, A-172 and U-87 cells were seeded in 96 multi-well plates

at a density of 1.0×10^4 per well and allowed to attach overnight. Cells were then treated

with DMSO and varying concentrations of ANA-12 (1 to 50 µM) or cannflavins (1 to 20

µM) diluted in Opti-MEM reduced serum media (Gibco, Cat No. 31985070) and incubated

at 37°C for the indicated time. Following treatment, resazurin sodium salt (Sigma-Aldrich,

Cat No. R7017) was dissolved in Hank's balanced salt solution (Wisent Inc., Saint-Jean-

Baptiste, QC, Canada, Cat No. 311-513-CL), diluted 1:10 in each well, and cells were

further incubated at 37°C for three hours. Fluorescence was measured on a microplate

reader at 570/590 nm, and a background control was subtracted from each value. All

treatments were completed in triplicate.

The following procedure was used for quantification: First, fluorescence values were

exported from the microplate reader, and the average of three background control wells

with Opti-MEM reduced serum media only was subtracted from each treatment value.

Second, the subtracted values were multiplied by 100 and divided by the average of the

DMSO controls to obtain a percentage relative to the vehicle control. Finally, the average

of three replicates per treatment was used as the final value for one biological replicate. A

minimum of three biological replicates were taken.

Lactate dehydrogenase cytotoxicity assay

Lactate dehydrogenase (LDH) release was measured using the CyQUANT LDH

Cytotoxicity Assay Kit according to the manufacturer's protocol (Invitrogen, Cat No.

C20300). Briefly, A-172 and U-87 cells were seeded in 96-well multiplates and treated

with the tested molecules, using a similar approach to the resazurin reduction assay. Before

the treatment endpoint, 10× Lysis Buffer was applied to control wells for 10 minutes to get

the Maximum LDH Release value. Cell culture supernatant from each treatment was

transferred to a new 96-well multiplate, an equal volume of reaction mixture was added,

mixed, and then the plate was incubated at room temperature for 30 minutes, protected

from light. Absorbance was measured on a microplate reader at 490 nm and 680 nm

(background). All treatments were completed in triplicate.

The following procedure was used for quantification: First, LDH activity was

calculated by subtracting the raw absorbance value measured at 680 nm from the 490 nm

value for each replicate and treatment. The average of three background control wells with

Opti-MEM reduced serum media was subtracted from each treatment value. Finally, the

LDH activity of each technical replicate of the compound-treated wells was divided by the

average of the maximum LDH activity and multiplied by 100 to obtain the percent

cytotoxicity. The average of the triplicates per treatment was taken and used as the final

value for one biological replicate; three biological replicates were completed in total.

Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) assay

The Roche *in Situ* Cell Death Detection Kit, Fluorescein (Sigma-Aldrich, Cat No.

11684795910), was used according to the manufacturer's protocol. Briefly, A-172 cells

were fixed using 4% paraformaldehyde for 15 minutes after treatment, then incubated for

2 minutes in permeabilization solution (0.1% Triton X-100, 0.1% sodium citrate in PBS).

This was followed by a 1-hour incubation in the dark at 37°C with humidity in TUNEL

reaction solution. After rinsing with PBS three times for 10 minutes, cells were incubated

in 4',6-diamidino-2-phenylindole (DAPI; Invitrogen, Cat No. 62247) staining solution for

10 minutes and finally washed in PBS twice for 10 minutes before being mounted onto

slides with ProLong Diamond Antifade (Invitrogen, Cat No. P36930).

Flow cytometry

A-172 cells and U-87 cells were seeded in 6-well plates at a density of 2.5 x 10^5 per well

and allowed to attach overnight. The next day, cells were treated with DMSO control,

cannflavin A, or cannflavin B (1 and 5 µM) for 24 hours. Following that time, the cell

monolayer was dissociated with trypsin, pelleted, resuspended in FACS buffer (PBS

supplemented with 2% FBS and 1 mM EDTA), and frozen fixed at -20°C in 70% ethanol.

Immediately before running samples, fixed cells were thawed and washed twice in FACS

buffer, 50 µL of 100 µg/mL RNAse A (Thermo Fisher Scientific, Cat No. EN0531) in

FACS buffer was added for 30 minutes, then 200 µL of 0.25 µg/mL propidium iodide

(Molecular Probes, Cat No. P-1304) in FACS buffer was added for 10 minutes in the dark.

The DNA contents were analyzed using a Northern Lights cytometer (Cytek Biosciences,

Bethesda, MD, USA).

Data quantification was performed using FlowJo v10.10 software. First, debris was

excluded using SSC-A vs. FSC-A gating strategy, and then singlets were selected using

FSC-H vs. FSC-A. Specifically, the univariate cell cycle platform Watson Pragmatic

algorithm distributed cell cycle peaks to the histograms presented, and data points were

plotted as a percentage of the cell count found in each cell cycle phase (G1, S, G2/M).

Transwell invasion assay

The bottoms of Corning Transwell inserts (New York, USA, Cat No. CLS3442) were

coated with 20 µg/mL fibronectin (Sigma-Aldrich, Cat No. F1141-1MG) at 4°C for 1 hour.

After three washes with PBS, the tops were coated with 0.2 mg/mL Corning Matrigel

matrix (Cat No. 354234) overnight at 37°C. In parallel with the preparation of the

transwells, A-172 cells were seeded at a density of 1.5 x 10^5 cells per well in 6-well plates

and grown to 85-90% confluency for 24 hours. They were then treated overnight with 10

µM cannflavin B or DMSO in DMEM containing 1% FBS media. The next day, 30,000

live cells from each condition were collected and seeded in the upper chamber of each well

with DMEM containing 1% FBS plus cannflavin B (10 µM), and in the bottom chamber

with DMEM containing 10% FBS. Cells were allowed to invade the Matrigel matrix for

24 hours at 37°C and 5% CO_2. Afterward, the transwell membranes containing the invaded

cells were removed. The membranes were stained with a 1% crystal violet solution in 20%

methanol for 20 minutes, washed three times with PBS, and dried for several hours at room

temperature. Representative photographic images of the membranes were taken and then

cut out. Finally, crystal violet was eluted from each membrane with 10% acetic acid for 20

minutes, and the absorbance (at 562 nm) of the eluates was measured to quantify invasion.

Three biological replicates with each treatment run in duplicate were analyzed.

Scratch migration assay

A-172 cells were seeded in a 6-well plate at a density of 2.0 x 10^5 per well and grown to

complete confluency in DMEM supplemented with 10% FBS for 48 hours. Using a P1000

pipette tip, 4 crossing linear wounds were created through the monolayer of cells within

each well. Next, the unattached cells were washed away with PBS twice before being added

back to 1% FBS DMEM supplemented with DMSO only or the treatment molecules.

Baseline images of the wound area were acquired immediately after adding the culture

media using a phase-contrast inverted Nikon Eclipse Ti2-E microscope (Nikon

Instruments, Melville, NY, USA) equipped with a motorized stage and STXF stage top

incubator (Tokai Hit USA Inc., PA, USA) to maintain temperature (37°C), humidity, and

CO_2 (5%, 150 mL/min) constant. Two separate areas per scratch were imaged every 15

minutes over a 48-hour period. NIS-Elements Advanced Research software was used to

run binary area fraction analysis, and the number of migrated cells was counted using

ImageJ v1.54 software. Three biological replicates were performed for each treatment

condition.

Generation of tumorspheres (tumor spheroids)

The tumorsphere protocol was adapted from previous work (Vinci et al., 2015; Tilak et al.,

2021b). Briefly, U-87 cells were passaged at least three times in complete media and then

20,000 cells per well were seeded into a clear round-bottom ultra-low attachment

microplate (Corning, Cat No. 7007). Tumorspheres were allowed to form for 5 days at

37°C and 5% CO_2, with half the media changed on Day 3.

Tumorsphere invasion assay

Individual tumorspheres prepared as mentioned above were embedded in Corning Matrigel

matrix (Cat No. 354234) containing the specified treatment. Four tumorspheres with a

circular shape and no protrusions were chosen per treatment, and half the media was

removed from the well. An equal volume of Matrigel supplemented with treatment at 2x

concentration was added to each well and incubated at 37°C and 5% CO_2 for one hour to

polymerize. Treatments at 1x of the desired treatment concentration were prepared in

complete media and added to each well immediately after incubation. The plate was then

imaged on a Nikon Eclipse Ti2-E inverted microscope equipped with a motorized stage,

image stitching capability, and a 10× objective at 0, 24, 48, and 72 hours. One image per

tumorsphere per treatment (4 tumorspheres and 4 treatments for 16 images at each time

point) was analyzed, and three biological replicates were performed.

The image analysis on original tiff files was performed with ImageJ as follows: For

the area measurement, the elliptical tool was used to encircle each tumorsphere, ensuring

all cells grown out from the original tumorsphere were included in the area measurement.

Three area measurements were taken for each tumorsphere at each time point and averaged.

For the radius measurement, the line tool was used to measure from the center point of the

tumorsphere out to the furthest protrusion from the tumorsphere. Three radius

measurements were taken for each tumorsphere at each time point and averaged. The

average area or radius of each treatment was divided by the DMSO control average at its

corresponding time point to obtain the values used for statistical analysis.

Generation of cerebral organoids

BIONi010-A induced pluripotent stem cells (iPSCs) were cultured on Matrigel-coated

plates (Corning, Cat No. 354277) in ExCellerate iPSC Expansion Medium (R&D Systems,

Cat No. CCM036). Cerebral organoids were generated using the STEMdiff Cerebral

Organoid Kit (StemCell Technologies, Cat No. 08570) as described in the manufacturer's

protocol and Tilak and colleagues (2021b). Briefly, on Day 0, iPSCs were lifted with

Gentle Cell Dissociation Reagent, resuspended in embryoid body (EB) Seeding Medium,

and seeded in an ultra-low attachment, round bottom 96-well plate (Corning, Cat No. 7007)

at a density of 9,000 cells/well. EB Formation Medium was added on Days 2 and 4, and

on Day 5, EBs were transferred to ultra-low attachment 24-well plates (Corning, Cat No.

3473) containing Induction Medium for 48 hours. On Day 7, EBs were encased in droplets

of Matrigel on organoid embedding sheets (StemCell Technologies, Cat No. 08579), which

were allowed to polymerize at 37°C before being washed into ultra-low attachment 6-well

plates (Corning, Cat No. 3471) using 3 mL of Expansion Medium. On Day 10, the

Expansion Medium was replaced with Maturation Medium, and the plate was transferred

to an orbital shaker inside a 37°C, 5% CO_2 incubator. The media was changed three times

per week until Day 60, the beginning of the organoid co-culture assay.

Organoid co-culture assay

The organoid co-culture assay protocol was modified from Tilak and colleagues (2021b).

Tumorspheres were generated similar to above, except with 12,000 GFP-expressing U-87

cells seeded per well. Sixty (60)-day-old organoids were removed from the 6-well plate

and placed individually in 1.5 mL Eppendorf tubes, each containing 4 tumorspheres and

500 μL of Maturation Medium. Tubes were incubated at 37°C and 5% CO_2 inside 10 cm

dishes with the caps open to allow airflow during attachment, while maintaining a sterile

environment. After 48 hours, 50% of the media was replaced, and after 72 hours, organoids

with tumorsphere attached were gently removed from the tubes using a wide-bore pipette

and put back in 6-well plates on the orbital shaker for an additional 24 hours (for the 10-

day experiment) or 8 days (for the 3-day experiment), at which point cannflavin B (5 μM)

or DMSO was added to the well in a complete media change. Media containing cannflavin

B or DMSO was replaced every other day for the duration of the experiment. Fourteen days

after initiating the co-culture, all organoids were removed from the wells, washed three

times with PBS, and fixed in 4% paraformaldehyde overnight at 4°C. Organoids were

washed again with PBS and incubated in a 30% sucrose solution at 4°C for 4 days before

being embedded in a 7.5% gelatin, 10% sucrose solution and snap-frozen in dry ice

submerged in 95% ethanol. Twenty (20) μm sections were obtained using a cryostat (Leica,

CM1860), placed on Vectabond-coated slides (Vector Laboratories, Cat No. SP-1800-7),

and stored at -80°C until immunostaining.

For immunostaining, organoid sections were washed 3 times with PBS and blocked in

10% normal goat serum in PBS at room temperature for 30 minutes. Sections were

incubated overnight at 4°C in blocking solution containing primary antibodies (TUJ1 and

GFP, 1:1,000). Slides were washed again 3 times with PBS and incubated in Alexa Fluor-

488- or Alexa Fluor-594-conjugated secondary antibodies (Invitrogen, Cat Nos. A11039

and A11032) in 5% normal goat serum in PBS for 2 hours at room temperature. Slides

were washed 3 additional times in PBS, counterstained with DAPI, and cover-slipped with

ProLong Glass Antifade mounting reagent (Invitrogen, Cat No. P36984).

Captures were taken using a Nikon Eclipse Ti2-E microscope at 20× magnification.

Exposure for each channel was kept consistent between conditions. To analyze GFP-

positive U-87 cell infiltration in the corresponding cerebral organoid, the following steps

were followed: 1) the perimeter of the cerebral organoid was traced based on the TUJ1

signal, 2) a threshold of 100 was applied to the GFP channel to binarize pixels, and 3) the

377 number of GFP pixels was normalized to the total number of pixels in the TUJ1 channel to

378 provide the proportional count. For estimating tumorsphere area, a section from

379 approximately the middle of each tumorsphere to be analyzed was immunostained, and the

380 perimeter of the GFP signal was traced using the polygon selection tool in ImageJ software

381 by an individual blinded to the experimental conditions.

383 *Western blotting*

384 For western blot analysis, cells were seeded in 6-well plates and grown for 2 to 3 days

385 before treatment. After treatment, cells were collected with radioimmunoprecipitation

386 assay (RIPA) buffer (50 mM Tris [pH 8.0], 0.5% Igepal-CA-630, 0.1% SDS, 0.5%

387 deoxycholic acid, 1 mM EDTA, 300 mM NaCl), supplemented with Phosphatase Inhibitor

388 Cocktail 3 (Sigma-Aldrich, Cat No. P0044) and EDTA-free Protease Inhibitor Cocktail

389 (Sigma-Aldrich, Cat No. 11836170001), and homogenized using QIAshredder columns

390 (QIAGEN, Cat No. 79656). Concentration of all lysates was measured using Pierce BCA

391 Assay Kit (Thermo Fisher Scientific, Cat No. 23225), one volume of 2× Laemmli buffer

392 (100 mM tris-HCl [pH 6.8], 4% SDS, 0.15% bromophenol blue, 20% glycerol, 200 mM β-

393 mercaptoethanol) was added, samples were adjusted to an equal concentration, and finally

394 boiled for 5 minutes. A total of 15 µg of protein was run on a 10% SDS-PAGE gel and

395 transferred to a nitrocellulose membrane. Membranes were incubated in TBST (Tris-

396 buffered saline and 0.1% Tween 20) supplemented with 5% non-fat powdered milk for 30

397 min at room temperature, and then probed overnight with the intended primary antibody at

398 the following dilution: P-TrkB Y705 (1:1,000), P-AKT S473 (1:1,000), P-AKT T308

399 (1:1,000), pan-AKT (1:2,000), P-p44/p42 MAPK (1:1,000), p42 MAPK (1:1,000), and

GAPDH (1:20,000). The next day, membranes were washed with TBST, incubated with

the appropriate secondary antibody, and visualized using enhanced chemiluminescence

(ECL), according to the manufacturer's guidelines (Pierce, Thermo Fisher Scientific). Each

experiment was repeated three times.

The western blots were quantified as follows: First, equal amount of protein sample

was analyzed by SDS-PAGE for each biological replicate. Second, the exposure time of

the film to ECL was the same for each biological replicate. Third, the exposed films were

scanned using a HP LaserJet Pro M377dw scanner in grayscale at a resolution of 300 dpi.

Finally, protein expression was quantified using ImageJ software to determine the mean

grey value of the protein bands, with the background value subtracted. The pixel density

for all data was inverted by subtracting the mean grey value from 255. The final values of

each protein band were taken as a ratio over the corresponding loading control to provide

a normalized value for statistical analysis.

Statistical analysis

All statistical calculations were completed with Prism 10 (GraphPad Software, Boston,

MA, USA). No statistical methods were employed to predetermine the sample size of any

of the presented experiments. Student *t*-test or analysis of variance (ANOVA) plus Tukey's

post hoc test for multiple comparisons was performed where indicated. A value of $p \leq .05$

was considered statistically significant. No test for outliers was conducted, and no data

point was excluded. Unless mentioned otherwise, all results represent the mean ± SEM

from at least three independent experiments ($n = 3$). Sample size was estimated based on

previous studies of a similar nature (Lalonde et al., 2021; Holborn et al.,

2025).

RESULTS

Cannflavin B decreases GBM cell viability

The flavonol kaempferol and flavone apigenin have been shown to significantly reduce the

viability of different cancer cell types in the low micromolar concentration range after 24-

48 hours of exposure (e.g., Gao *et al.*, 2018; Tseng *et al.*, 2017). Except for the work of

Moreau and colleagues (2019) using mouse models of metastatic pancreatic cancer and the

study by Tomko and colleagues (2022), which reported a promising effect of cannflavin A

against bladder carcinoma cells, we know little about the impact of cannflavins on other

cancer types. This prompted us to thoroughly test the effect of cannflavin A and cannflavin

B on the viability of the A-172 and U-87 GBM cell lines. Importantly, we used the small-

molecule non-competitive TrkB antagonist ANA-12 (Cazorla et al., 2011) as a comparison

compound in this analysis because it has been reported to reduce A-172 cell viability in a

dose- and time-dependent manner (Pinheiro et al., 2017). Hence, this represents a relevant

reference measure that also connects with our recent publication demonstrating the

interference of cannflavins A and B on TrkB signaling in mouse primary cortical neurons

(Holborn et al., 2023).

As presented in **Figure 1**, applying cannflavin A and cannflavin B in the 1-20 µM

final concentration range to the culture media of either A-172 (**Fig. 1A-C**) and U-87 (**Fig.**

1D-F) cells produced similar mixed results. Specifically, no significant effect on viability

was detected using a resazurin reduction assay for cannflavin A at any concentration or

time-point, although a trend for reduced viability can be observed in A-172 cells with the

highest concentration (20 µM) at the 24-hour and 48-hour time-points. On the other hand,

cannflavin B produced a robust reduction in viability for both A-172 and U-87 cells, with

the most significant effect observed at higher concentrations (20 µM) and longer treatment

times (48 hours). Interestingly, ANA-12 in the 20-50 µM range also negatively affected

survival in both cell lines, with the notable difference that higher compound concentrations

were required to achieve an impact comparable to that of cannflavin B. This can be best

appreciated at 48 hours in A-172 cells, where 5 µM cannflavin B reduced viability below

50%, while ANA-12 at 10 µM had no statistically significant effect (estimated average cell

viability of 90%). The estimated IC50 for each of the compounds and time-points in this

experiment are presented in the **Supplementary Information**. Together, these results

establish that cannflavin B is more potent at affecting GBM cell survival than cannflavin

A and ANA-12.

To gain a broader view of the potency of cannflavin B towards U-87 cells, we next

performed a similar experiment with two structurally related flavones, namely chrysoeriol,

the non-prenylated synthesis precursor of cannflavins in the *C. sativa* plant (**Fig. S2A**, Rea

et al., 2019), and techtochrysin (**Fig. S2B**), an O-methylated flavone previously discovered

to have effects against SW480 and HCT116 colon cancer cells through inhibition of

Nuclear factor-κB (NF-κB) activity (Park et al., 2015). Interestingly, we found that the

direct application of chrysoeriol to the cell culture medium significantly reduced U-87 cell

viability, but only at a final concentration of 20 µM and for treatment times of 24-48 hours

(**Fig. S2C**). This result suggests that the prenyl group found at the 3' position of the flavone

B-ring that distinguishes cannflavin B from chrysoeriol contributes to the potency of the

latter molecule in this assay. The effect of techtochrysin, however, was the opposite of that

of chrysoeriol. Intriguingly, application of techtochrysin in the range of 5-40 μM for 24-

48 hours significantly enhanced the viability of U-87 cells. In summary, these experiments

reinforce the notion that cannflavin B exerts anti-survival effects on GBM cells at lower

concentrations and more rapidly than closely related flavones.

Cannflavins A and B do not induce necrosis in GBM cells

To assess the extent to which cannflavin B's observed impact on GBM cell viability could

be attributed to necrosis, we performed cell membrane integrity tests with the lactate

dehydrogenase (LDH) release assay. A-172 and U-87 cells treated with 10-20 μM

cannflavin A or cannflavin B for 24 hours presented minimal LDH release (**Fig. S3A, B**).

Specifically, application of cannflavin A resulted in 3.57% LDH release at 10 μM and

2.54% at 20 μM in A-172 cells, and 0.96% at 10 μM and 1.03% at 20 μM in U-87 cells.

For cannflavin B, LDH release was 4.43% at 10 μM and 5.38% at 20 μM in A-172 cells,

and 0.81% at 10 μM and 0.88% at 20 μM in U-87 cells. Those values are comparable to

those obtained with ANA-12 at 10 μM (2.01% for A-172 cells and 1.61% for U-87). In

summary, these results suggest that the adverse effect of cannflavin B on cell viability, as

assessed using the resazurin reduction method, is not due to necrotic cell death at the

specified concentrations and time points.

Cannflavins A and B do not cause apoptosis in the low micromolar range

Our following goals were to assess the effect of cannflavins A and B in the low micromolar

(1 to 10 μM) range on GBM's cell cycle, migration, and capacity to invade. Before

proceeding with those tests, though, we wanted to confirm that neither cannflavin A nor

cannflavin B causes widespread programmed cell death resulting in DNA fragmentation

after at least 24 hours of exposure. Using the highly sensitive TUNEL assay, we observed

that A-172 GBM cells treated with cannflavin A or cannflavin B at a final concentration of

10 µM for 24 hours did not produce a noticeable fluorescein signal indicative of apoptosis

(**Fig. S3C**). This negative result assures that any effect associated with cannflavins at a

concentration of 10 µM or lower, which will be identified next, is distinct from

programmed cell death.

Effect of cannflavins on A-172 and U-87 cell cycle

Various flavonoids have been demonstrated to affect the cell cycle of different cancer cell

types (e.g., Lin et al., 2015; Srivastava et al., 2016; Zhang et al., 2017). This led us to ask

whether cannflavins A and B, at concentrations that do not produce a rapid and significant

anti-viability effect on A-172 and U-87 cells, could alter their cell cycle. Flow cytometry

analysis showed that neither cannflavin A nor cannflavin B affected cell cycle phase after

24 hours of treatment in the concentration range of 1-5 µM (**Fig. 2**). Interestingly, the A-

172 and U-87 cell lines exhibited distinctive distributions of the cell cycle that remained

unchanged with treatment (G1 proportion for the DMSO control was 61.7% and 41.6%,

respectively). In both GBM cell lines, compared to the DMSO control, there was no

statistically significant change in the cell cycle phase upon cannflavin A or cannflavin B

treatment. These results demonstrate that cannflavins in the low micromolar range (1 to 10

µM) do not alter GBM cell cycle.

Cannflavins limit GBM cell migration and invasion

Invasion and metastasis are primarily responsible for mortality associated with solid tumors

(Li et al., 2025). This is the case for GBM, where rapid invasion and damaging infiltration

of the cancer cells into surrounding healthy brain tissue account for most of the poor

outcomes (Barbaro et al., 2022). Therefore, systematically assessing the action of

cannflavins on GBM cell migration and invasion is highly relevant.

To start, we completed an analysis of cell migration using a time-lapse scratch assay

in which cannflavin A (10 μM final concentration), cannflavin B (5 μM final

concentration), ANA-12 (10 μM final concentration), or DMSO alone was applied directly

to the culture media of A-172 cells. Cell migration into the scratch wound area was

monitored over 48 hours (**Fig. 3A** and **Supplementary Video**). Our data revealed striking

differences between wells treated with cannflavins and exposed to DMSO. First, analysis

of the binary area fraction over all time-points suggested a delayed wound closure in

inhibitor-treated conditions and more linear cell movement over time for the DMSO and

cannflavin A conditions compared to the cannflavin B and ANA-12 (**Fig. 3B**). Examining

next just the 24-hour and 48-hour time-points, we found that the normalized wound area

fraction was significantly lower for the three tested molecules compared to DMSO (**Fig.**

3C). Further, considering the data as a percentage of wound closure, significantly less area

was covered by cells after 24 hours and 48 hours exposure to cannflavin B, but only at 48

hours for cannflavin A (**Fig. 3D**). ANA-12 results in that analysis were not different

compared to DMSO. Finally, considering the number of individual cells detectable within

the wound area, significantly fewer were present at the end of the experiment for cannflavin

B (**Fig. 3E**).

Next, we performed a transwell assay to gauge the potential of cannflavin B to limit

A-172's capacity to invade when the cells are exposed to the molecule. A-172 cells were

pre-treated overnight (~16 hours) with 10 µM cannflavin B or DMSO and then plated on

Matrigel-coated inserts with the addition of fresh cannflavin B (10 µM). As presented in

Figure 4, the results from the transwell invasion assay revealed that exposure to cannflavin

B significantly reduced the number of cells that invaded through the Matrigel matrix,

resulting in a statistically significant reduction of more than 50% compared to the vehicle

control. This result supports that cannflavin B at a concentration of 10 µM effectively

inhibits GBM cell migration and invasion, providing initial evidence of its potential against

tumor progression.

Lastly, since U-87 cells have a strong tendency to aggregate and form tumor-like

spheroid formations (tumorspheres) in suspension culture, we leverage this attribute to

provide a different evaluation of the action of cannflavins A and B on the invasion ability

of these GBM cell subtypes. Specifically, we cultured U-87 as 3-dimensional (3D)

tumorspheres and then embedded those within a Matrigel matrix to measure how they

would progress in the outward direction (**Fig. 5A**). Interestingly, adding cannflavin A to

the culture media at a final concentration of 10 µM resulted in a significantly smaller radius

and area of tumorsphere outgrowth at the three examined times (**Fig. 5B, C**). For cannflavin

B, tumorspheres in wells treated at a final concentration of 5 µM presented a significantly

reduced area at 72 hours after the start of the experiment (**Fig. 5B**) and a smaller radius in

the 48-72 hours times (**Fig. 5C**). ANA-12, for its part, was minimally active in this assay

producing only a slight significant effect on radius at 72 hours (**Fig. 5C**). Taken together,

these three experiments provide strong evidence that cannflavins, particularly cannflavin

B, can produce rapid and sustained anti-migration and anti-invasion action against GBM

cells.

Cannflavin B limits the growth and infiltration of U-87 GBM tumorspheres attached to

cerebral organoids

Conventional pre-clinical studies for GBM consist of monolayer tumor cell cultures and

animal approaches, including xenografts and genetically engineered mouse models

(Gómez-Oliva et al., 2021; Haddad et al., 2021). However, these strategies encounter

significant obstacles in recapitulating the complexity of GBM, especially the variety of

tumor cellular states and the cancerous mass microenvironment. The use of brain organoids

derived from human pluripotent stem cells helps overcome some of those challenges and

provides exciting new ways to study GBM tumor formation and invasion in real time

(Linkous et al., 2019; Wang et al., 2023)—an approach that our group has used with

success in recent years (Tilak et al., 2021b; Lau et al., 2025). To test if cannflavin B could

produce therapeutically relevant effects in a more complex *in vitro* system, we conducted

a co-culture experiment where iPSC-derived cerebral organoids matured for 60 days were

mixed with U-87 tumorspheres expressing GFP (**Fig. 6A**). Our pool of co-culture

specimens was then divided in two groups: one where the U-87 tumorspheres attached and

grew unchallenged for 11 days followed by the application of cannflavin B at a final

concentration of 5 µM in the culture media for 3 days (Paradigm 1), and another (Paradigm

2) where the cancer cells attached and grew for 4 days and then 5 µM cannflavin B was

added for 10 days (**Fig. 6B**). Co-cultures treated with DMSO only were included for both

tests as control, and media change with fresh cannflavin B (5 µM) or DMSO was done

every other day for the duration of the experiment. As seen in fluorescence microscopy

captures of immunostained sections (**Figure 6C-F**), U-87 tumorspheres attached to

cerebral organoids in both paradigms. In the two DMSO control conditions (**Fig. 6C, E**), a

distinct intermingling between GBM cells expressing GFP and neurons, as revealed by

TUJ1 immunostaining, can be observed at the edge of the cerebral organoid where the

tumorsphere is attached. Interestingly, less GBM cell penetration into the organoids was

observed in the cannflavin B-treated wells in both tests (**Fig. 6D, F**). To quantify this

phenotype, we determined the proportion of pixels corresponding to the GFP signal (U-87

GBM cells) in their respective cerebral organoid section (**Fig. 6G, H**). Although this

analysis did not reveal statistical significance due to low GBM cell infiltration in some

DMSO control replicates, cannflavin B-treated samples consistently showed very little

GFP signal. Considering this fact, we consider this result highly consistent with other

findings demonstrating the efficacy of cannflavin B against GBM cells. Finally, although

the tumorsphere total area for corresponding sections was not statistically different between

the two conditions in the 3-day (Paradigm 1) experiment (**Fig. 6I**), we found that the U-87

tumorspheres were significantly smaller when cannflavin B was added for 10 days

(Paradigm 2, **Fig. 6J**). Together, these observations collected with a more physiologically

relevant model provide compelling additional data to support an effect of cannflavin B

against the ability of aggressive GBM cells to invade human brain-like tissue.

Cannflavins A and B impact on TrkB signaling in GBM cells

As mentioned previously, a study by Pinheiro and colleagues (2017) reported that the TrkB

inhibitor ANA-12 reduces the viability of A-172 cells, suggesting that the activity of this

RTK may be essential for the growth and survival of this GBM cell line. Since our previous

work using mouse primary cortical neurons also found that cannflavins A and B can

interfere with TrkB signaling (Holborn et al., 2023), we considered it necessary to gain

more information about the effect of cannflavins and ANA-12 on TrkB signaling in A-172

(**Fig. 7**) and U-87 (**Fig. 8**) cells to interpret our functional data. As expected, western blot

analysis revealed that application of cannflavin A at a final concentration of 20 µM (**Fig.**

7A, B) and cannflavin B at 10-20 µM (**Fig. 7F, G**) to the culture media of A-172 cells for

6 hours significantly reduced the phosphorylation of TrkB at tyrosine 705 (Y705), which

is in the catalytic domain and stimulates kinase activity of the receptor when

phosphorylated. Consistent with this result, ANA-12 at a final concentration of 10 µM also

significantly decreased phospho-TrkB Y705 signal (**Fig. 7B, G**). Interestingly, however,

our western blot data also revealed two unexpected and critical results. Firstly, key

downstream TrkB signaling effectors (i.e., p44/p42 MAPK and AKT) were unaffected by

cannflavins and ANA-12 in A-172 cells (**Fig. 7C-E, H-J**). Secondly, cannflavins and

ANA-12 did not impair TrkB phosphorylation and downstream signaling effectors in U-

87 cells, even at the highest tested concentration (**Fig. 8**). Together, these western blot data

force the critical conclusion that any positive results about GBM survival, migration, and

invasion presented above are most likely distinct from TrkB signaling impairment, whether

they were the result of cannflavins or ANA-12 treatment.

DISCUSSION

In this study, we employed various *in vitro* assays to characterize the phenotypic outcomes

of cannflavins A and B on the viability, cell cycle, migration, and invasion capacities of A-

172 and U-87 GBM cells. Of particular interest, our results show a consistent dose-

dependent decrease in cell viability in both lines when cannflavin B is added to the culture

media. No effect on cell cycle by either cannflavin molecule was detected, but we found

that both molecules can reliably limit tumor cell migration below the 10 µM concentration

and that cannflavin B has robust anti-invasive properties based on results from transwell

and tumorsphere assays. Together, these observations underscore the multifaceted

therapeutic potential of cannflavin B against this aggressive type of brain cancer.

Focusing first on our overall cell survival results, it is interesting to speculate about

why cannflavin B shows a higher effect against A-172 and U-87 cells than the other tested

flavonoids (i.e., cannflavin A, chrysoeriol, and techtochrysin). One key factor contributing

to this effect is certainly prenylation, which is considered to increase the lipophilicity of

flavonoids, thereby enhancing their affinity for biological membranes and improving their

interactions with target proteins (Xu et al., 2012; Yang et al., 2015). The reduced potency

of chrysoeriol, which is structurally similar to cannflavins A and B but lacks prenylation

attached to the B-ring (**Fig. S2A, C**), is consistent with this interpretation. That said, the

length and position of the prenyl side-chain are undoubtedly also critical, as we measured

greater bioactivity with cannflavin B, which has a single prenyl group, compared to

cannflavin A, which has two (**Fig. S1**). Finally, another structural-activity feature that may

explain other differences among flavonoids in our study is the B-ring hydroxylation

patterns. In fact, extensive evidence suggests that hydroxylation at specific positions can

markedly enhance bioactivity (Liu et al., 2022; Li et al., 2023). This detail most likely

contributed to the result we obtained in the techtochrysin test (Fig. S2B, D).

In addition, we conducted a western blot analysis to gain insights about the possible

molecular signaling changes induced by cannflavins that could produce the reported

cellular effects. However, the tested cannflavins and ANA-12 consistently reduced TrkB

phosphorylation at Y705 in A-172 cells, but this effect was not observed in U-87 cells. The

divergent TrkB phosphorylation responses observed in A-172 and U-87 cells likely reflect

differences in baseline endogenous TrkB expression (**Fig. S4**) and downstream pathway

coupling. U-87 cells are known to express lower levels of full-length TrkB than A-172

cells, which may explain their limited responsiveness. Most importantly, cannflavins and

ANA-12 had minimal effects on MAPK and AKT, which are well-known contributing

factors to cancer hallmarks, including growth, apoptosis, and invasion. These unexpected

results depart from our previous work on cannflavins A and B in mouse primary cortical

neurons (Holborn et al., 2023) and suggest that different molecular events, influenced by

cannflavins, are responsible for the migration and invasion disruptions observed in GBM

cells in this study. Therefore, future work is needed to elucidate the target engagement of

cannflavins in GBM cells. One unbiased strategy for this task could be to utilize the

Proteome Integral Solubility Alteration (PISA) assay, a high-throughput and mass-

spectrometry-based approach that leverages binding-dependent changes in protein

solubility to identify drug-target interactions (Gaetani et al., 2019; Gaetani and Zubarev,

2023).

On a separate note, an intriguing finding in our study concerns the increased viability

of U-87 cells exposed to techtochrysin (**Fig. S2B, D**)—a flavone molecule that had been

reported to suppress the growth of SW480 and HCT116 human colon cancer cells by

inhibiting NF-κB resulting in the expression of death receptors like DR3, DR4, and Fas

(Park et al., 2015). Although we cannot explain at this point why U-87 cells exposed to this

specific flavone produced an opposite effect to what has been reported with human colon

cancer cells, this bring to our attention the possibility that some of the anti-cancer

phenotypes associated with cannflavins A and B that we found may be linked to disruption

of the transcription factor NF-κB. Indeed, extensive research has implicated NF-κB in

various aspects of cancer formation, growth, treatment response, immune escape, and other

related processes (Taniguchi and Karin, 2018; Zhang et al., 2021; Khan et al., 2024). In

GBM and other gliomas, it has been reported for more than two decades that the aberrant

constitutive activity of this transcription factor contributes to disease initiation and

progression, including the invasion phenotype (Nagaai et al., 2002; Raychaudhuri et al.,

2004; Wang et al., 2004). Considering this, one tantalizing scenario is the disruption by

cannflavins of the constitutive signaling pathway that phosphorylates and promotes

degradation of IκBα by the ubiquitin-proteasome system, ultimately releasing NF-κB

dimers that translocate to the nucleus and modulate target gene expression that contributes

to disease progression (Cahill et al., 2016). While the present study did not directly evaluate

NF-κB activity, these findings raise the possibility that cannflavin B may influence NF-

κB–associated invasion programs. Future work incorporating NF-κB reporter assays and

transcriptomic profiling will be essential to define the signaling changes underlying the

anti-invasive effects observed here. If this is the case, it would be revealing to test the effect

of cannflavin B administration in xenograft or induced mouse models of GBM as used in

previous work targeting aberrant NF-κB activity (Friedmann-Morvinski et al., 2016).

Ultimately, this would provide a direct test of this hypothesis and additional pre-clinical

evidence to promote cannflavin B as a potentially valuable therapeutic agent against GBM.

Lastly, we present evidence that GFP-expressing U-87 cells migrating from a

tumorsphere can be challenged by cannflavin B at low μM levels as they infiltrate human

iPSC-derived cerebral organoids. Although we could not obtain precise quantitative

measures of GBM cell infiltration into the cerebral organoid structure using thin frozen

sections, we found that the tumorsphere size was significantly smaller after 10 days of

exposure to cannflavin B than after exposure to DMSO (vehicle control). This result calls

for a deeper investigation into the therapeutic value of cannflavin B against GBM or the

brain metastasis of other cancer types using live animal approaches, as mentioned above.

Corroborating the effectiveness of cannflavin B at limiting cancer progression with future

animal models would advance our research towards the design and proposal of a clinical

trial with human patients.

In conclusion, the distinctive ability of cannflavins, particularly cannflavin B, to

inhibit GBM cell migration and invasion, combined with their restrained cytotoxicity,

positions them as promising candidates for developing novel therapeutic strategies against

GBM (**Fig. S5**). Future studies should focus on validating these effects in animal models

to understand the pharmacokinetics, bioavailability, and therapeutic index of cannflavins.

Furthermore, exploring the synergistic impact of cannflavins with existing

chemotherapeutic agents or targeted therapies could provide valuable insights into their

potential as adjuvant treatments, particularly in overcoming drug resistance—a significant

challenge in the management of GBM. Combining cannflavins with agents like TMZ

(Stupp et al., 2005; Stupp et al., 2009; Fisher et al., 2013) or the anti-angiogenic agent

bevacizumab (Fu et al., 2023) could enhance their therapeutic efficacy by simultaneously

targeting multiple aspects of tumor growth and invasion.

DATA AVAILABILITY

All data collected for the study are available upon inquiry to the corresponding author.

CONSENT FOR PUBLICATION

Consent to publish the article has been obtained from all authors.

FUNDING

This work was supported by Mitacs (Mitacs Elevate Program, IT29490) and the Canadian

Foundation for Innovation (CFI, 037755). J.H. was also supported by a Queen Elizabeth II

Graduate Scholarship in Science and Technology.

CREdiT AUTHORSHIP CONTRIBUTION STATEMENT

Jennifer Holborn: Conceptualization, Data curation, Formal analysis, Funding

acquisition, Investigation, Writing – original draft, **Hannah Robeson:** Data curation,

Methodology, Writing – review & editing. **Ellis Chartley:** Formal analysis, Methodology,

Writing – review & editing. **Tiana Gluscevic**: Data curation, Methodology. **Adina**

Borenstein: Methodology. **Colby Perrin:** Methodology. **Begüm Alural:** Methodology.

Himain Perera: Visualization, Writing – review & editing. **Tariq A. Akhtar:** Resources.

Nina Jones: Resources, Writing – review & editing. **Jasmin Lalonde:** Conceptualization,

Data curation, Funding acquisition, Supervision, Writing – original draft.

DECLARATION OF COMPETING INTEREST

The other authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

ACKNOWLEDGMENTS

We wish to thank our lab colleagues for their help with this project.

HTTPS://WWW.EDITORIALMANAGER.COM/PHYPLU/.EMENTARY MATERIALS

Fig. S1 Cannflavin family members.

Fig. S2 LDH assay and TUNEL assay.

Fig. S3 Chrysoeriol and techtochrysin survival assay.

Fig. S4 Level of phospho-TrkB Y705 between A-172 and U-87 untreated cells.

Fig. S5 Graphical abstract.

Supplementary Table. Estimated IC50 for anti-survival effects of cannflavins and

ANA-12.

Supplementary Video. Scratch migration assay video.

REFERENCES

Balakrishnan, S., Bhat, F. A., Raja Singh, P., Mukherjee, S., Elumalai, P., Das, S., Patra,

C. R., Arunakaran, J., 2016. Gold nanoparticle–conjugated quercetin inhibits

epithelial–mesenchymal transition, angiogenesis and invasiveness via

EGFR/VEGFR-2-mediated pathway in breast cancer. Cell Prolif. 49, 678–697.

https://doi.org/10.1111/cpr.12296.

Barbaro, M., Blinderman, C.D., Iwamoto, F.M., Kreisl, T.N., Welch, M.R., Odia, Y.,

Donovan L.E., Joanta-Gomez, A.E., Evans, K.A., Lassman, A.B., 2022. Causes of

death and end-of-life care in patients with intracranial high-grade gliomas.

Neurology. 98, e260-e266. https://doi:10.1212/WNL.0000000000013057.

Barrett, M.L., Gordon, D., Evans, F.J., 1985. Isolation from Cannabis sativa L. of

cannflavin—a novel inhibitor of prostaglandin production. Biochem Pharmacol. 34,

2019–2024. http://doi:10.1016/0006-2952(85)90325-9.

Birzu C, French P, Caccese M, Cerretti G, Idbaih A, Zagonel V, Lombardi G., 2020.

Recurrent glioblastoma: From molecular landscape to new treatment perspectives.

Cancers (Basel). 13, 47. https://doi.org/10.3390/cancers13010047.

Cahill, K.E., Morshed, R.A., Yamini, B., 2016. Nuclear factor-κB in glioblastoma:

insights into regulators and targeted therapy. Neuro Oncol. 2016 18, 329–

339. https://doi.org/10.1093/neuonc/nov265.

Cazorla, M., Prémont, J., Mann, A., Girard, N., Kellendonk, C., Rognan, D., 2011.

Identification of a low-molecular weight TrkB antagonist with anxiolytic and

antidepressant activity in mice. J. Clin. Invest. 121, 1846–1857.

https://doi:10.1172/JCI43992.

Cheng, W.Y., Chiao, M.T., Liang, Y.J., Yang, Y.C., Shen, C.C., Yang, C.Y., 2013.

Luteolin inhibits migration of human glioblastoma U-87 MG and T98G cells through

downregulation of Cdc42 expression and PI3K/AKT activity. Mol. Biol. Rep. 40,

5315–5326. https://doi.org/10.1007/s11033-013-2632-1.

Choi, E.J., Bae, S.M., Ahn, W.S., 2008. Antiproliferative effects of quercetin through cell

cycle arrest and apoptosis in human breast cancer MDA-MB-453 cells. Arch. Pharm.

Res. 31, 1281–1285. https://doi.org/10.1007/s12272-001-2107-0.

Friedmann-Morvinski, D., Narasimamurthy, R., Xia, Y., Myskiw, C., Soda, Y., Verma,

I.M., 2016. Targeting NF-κB in glioblastoma: A therapeutic approach. Sci. Adv. 2,

e1501292. https://doi.org/10.1126/sciadv.1501292.

Eggers, C., Fujitani, M., Kato, R., Smid, S., 2019. Novel cannabis flavonoid, cannflavin

A displays both a hormetic and neuroprotective profile against amyloid β-mediated

neurotoxicity in PC12 cells: Comparison with geranylated flavonoids, mimulone and

diplacone. Biochem. Pharmacol., 169. https://doi.org/10.1016/j.bcp.2019.08.011.

Erdogan, S., Doganlar, O., Doganlar, Z.B., Serttas, R., Turkekul, K., Dibirdik, I., Bilir,

A., 2016. The flavonoid apigenin reduces prostate cancer CD44+ stem cell survival

and migration through PI3K/Akt/NF-κB signaling. Life Sci. 162, 77–86.

https://doi.org/10.1016/j.lfs.2016.08.019.

Fisher J.P., Adamson D.C., 2021. Current FDA-approved therapies for high-grade

malignant gliomas. Biomedicines. 9, 324.

https://doi.org/10.3390/biomedicines9030324.

Fu., M., Zhou, Z., Huang, X., Chen, Z., Zhang, L., Zhang, J., Hua. W., Mao, Y., 2023.

Use of Bevacizumab in recurrent glioblastoma: a scoping review and evidence map.

BMC Cancer. 23, 544. https://doi.org/10.1186/s12885-023-11043-6.

Gaetani, M., Sabatier, P., Saei, A.A., Beusch, C.M., Yang, Z., Lundström, S.L., Zubarev,

R.A., 2019. Proteome integral solubility alteration: A high-throughput proteomics

assay for target deconvolution. J. Proteome Res. 18, 4027–4037.

https://doi.org/10.1021/acs.jproteome.9b00500.

Gaetani, M., Zubarev, R.A., 2023. Proteome Integral Solubility Alteration (PISA) for

high-throughput ligand target deconvolution with increased statistical significance

and reduced sample amount. Methods Mol. Biol. 2554, 91–106.

https://doi.org/10.1007/978-1-0716-2624-5_7.

Gao, X., Wang, B., Wei, X., Men, K., Zheng, F., Zhou, Y., Zheng, Y., Gou, M., Huang,

M., Guo, G., Huang, N., Qian, Z., Wei, Y., 2012. Anticancer effect and mechanism

of polymer micelle-encapsulated quercetin on ovarian cancer. Nanoscale, 4, 7021–

7030. https://doi.org/10.1039/c2nr32181e.

Gao, Y., Yin, J., Rankin, G.O., Chen, Y.C., 2018. Kaempferol induces G2/M cell cycle

arrest via checkpoint kinase 2 and promotes apoptosis via death receptors in human

ovarian carcinoma A2780/CP70 cells. Molecules, 23, 1095.

https://doi.org/10.3390/molecules23051095.

Gómez-Oliva, R., Domínguez-García, S., Carrascal, L., Abalos-Martínez, J., Pardillo-

Díaz, R., Verástegui, C., Castro, C., Nunez-Abades, P., Geribaldi-Dolbán, N., 2021.

Evolution of experimental models in the study of glioblastoma: Toward finding

efficient treatments. Front. Oncol., 10, 614295.

https://doi:10.3389/fonc.2020.614295.

Haddad, A.F., Young, J.S., Amara, D., Berger, M.S., Raleigh, D.R., Aghi, M., Butowski,

N.A., 2021. Mouse models of glioblastoma for the evaluation of novel therapeutic

strategies. Neurooncol. Adv., **b**, vdab100. https://doi:10.1093/noajnl/vdab100.

Holborn J, Walczyk-Mooradally A, Perrin C, Alural B, Aitchison C, Borenstein A, Jones

N, Khokhar J.Y., Akhtar T.A., Lalonde J., 2023. Interference of neuronal TrkB

signaling by the cannabis-derived flavonoids cannflavins A and B. Phytomed. Plus.

3, 100410. https://doi.org/10.1016/j.phyplu.2023.100410.

Huang, L., Jin, K., Lan, H., 2019. Luteolin inhibits cell cycle progression and induces

apoptosis of breast cancer cells through downregulation of human telomerase reverse

transcriptase. Oncol. Lett. 17, 3842–3850. https://doi.org/10.3892/ol.2019.10052.

Jo, E., Park, S.J., Choi, Y.S., Jeon, W.K., Kim, B.C., 2015. Kaempferol suppresses

transforming growth factor-β1-induced epithelial-to-mesenchymal transition and

migration of A549 lung cancer cells by inhibiting Akt1-mediated phosphorylation of

Smad3 at threonine-179. Neoplasia. 17, 525–537.

https://doi.org/10.1016/j.neo.2015.06.004.

Johnson, D.R., O'Neill, B.P., 2012. Glioblastoma survival in the United States before and

during the temozolomide era. J. Neuro-Oncol. 107, 359–364.

https://doi.org/10.1007/s11060-011-0749-4.

Khan, A., Zhang, Y., Ma, N., Shi, J., Hou, Y., 2024 NF-κB role on tumor proliferation,

migration, invasion and immune escape. Cancer Gene Ther. 31, 1599–1610.

https://doi.org/10.1038/s41417-024-00811-6.

Kim, S. H., Yoo, E.S., Woo, J.S., Han, S.H., Lee, J.H., Jung, S.H., Kim, H.J., Jung, J.Y.,

2019. Antitumor and apoptotic effects of quercetin on human melanoma cells

involving JNK/P38 MAPK signaling activation. Eur. J. Pharmacol. 860:172568.

https://doi.org/10.1016/j.ejphar.2019.172568.

Lalonde, J., Reis, S.A., Sivakumaran, S., Holland, C.S., Wesseling, H., Sauld, J., Alural,

B., Zhao, W.N., Steen, J., Haggarty, S.J., 2017. Chemogenomic analysis reveals key

role for lysine acetylation in regulating Arc protein stability. Nat. Commun. 8:1659.

https://doi.org/10.1038/s41467-017-01750-7.

Lau, H.R., Alural, B., Martin, C.E., New, L.A., Smith, H.S., Tilak, M., Banerjee, S.L.,

Robeson, H.N., Bisson, N., Gingras, A.C., Lalonde J, Jones, N., 2025. ShcD adaptor

protein drives invasion of triple-negative breast cancer cells by aberrant activation of

EGFR signaling. Mol. Oncol. Mar. 28, https://doi.org/10.1002/1878-0261.70022.

Lee, W.J., Hsiao, M., Chang, J.L., Yang, S.F., Tseng, T.H., Cheng, C.W., Chow, J.M.,

Lin, K.H., Lin, Y.W., Liu, C.C., Lee, L.M., Chien, M.H., 2015. Quercetin induces

mitochondrial-derived apoptosis via reactive oxygen species-mediated ERK

activation in HL-60 leukemia cells and xenograft. Arch. Toxicol. 89, 1103–1117.

https://doi.org/10.1007/s00204-014-1300-0.

Li, Y., Liu, F., Cai, Q., Deng, L., Ouyang, Q., Zhang, X.H.F., Zheng, J., 2025. Invasion

and metastasis in cancer: molecular insights and therapeutic targets. Signal

Transduct. Target. Ther. 10, 57. https://doi.org/10.1038/s41392-025-02148-4.

Li, S., Yan, T., Deng, R., Jiang, X., Xiong, H., Wang, Y., Yu, Q., Wang, X., Chen, C.,

Zhu, Y., 2017. Low dose of kaempferol suppresses the migration and invasion of

triple-negative breast cancer cells by downregulating the activities of RhoA and

Rac1. Onco Targets Ther. 10, 4809–4819. https://doi.org/10.2147/OTT.S140886.

Li, X., Xu, R., Ni, T., 2023. Effects of B-ring structures on binding behavior of flavonols

with proteins: Experimental and molecular docking approaches. J Mol Struct. 1287,

135614. https://doi.org/10.1016/j.molstruc.2023.135614.

Lin, C.H., Chang, C.Y., Lee, K.R., Lin, H.J., Chen, T.H., Wan, L., 2015. Flavones inhibit

breast cancer proliferation through the Akt/FOXO3a signaling pathway. BMC

Cancer. 15, 958. https://doi.org/10.1186/s12885-015-1965-7.

Linkous, A., Balamatsias, D., Snuderl, M., Edwards, L., Miyaguchi, K., Milner, T.,

Reich, B., Cohen-Gould, L., Storaska, A., Nakayama, Y., Schenkein, E., Singhania,

R., Cirigliano, S., Magdeldin, T., Lin, Y., Nanjangud, G., Chadalavada, K., Pisapia,

D., Liston, C., Fine, H.A., 2019. Modeling Patient-Derived Glioblastoma with

Cerebral Organoids. Cell Rep., 26, 3203–3211.

Liu, L., Qian, J., Li, J., Xing, M., Grierson, D., Sun, C., Xu, C., Li, X., Chen, K., 2022.

Hydroxylation decoration patterns of flavonoids in horticultural crops: chemistry,

bioactivity, and biosynthesis. Hortic Res. 9, uhab068.

https://doi.org/10.1093/hr/uhab068.

Moreau, M., Ibeh, U., Decosmo, K., Bih, N., Yasmin-Karim, S., Toyang, N., Lowe, H.,

Ngwa, W., 2019. Flavonoid derivative of cannabis demonstrates therapeutic potential

in preclinical models of metastatic pancreatic cancer. Front. Oncol. 9, 660.

https://doi.org/10.3389/fonc.2019.00660.

Nagai, S., Washiyama, K., Kurimoto, M., Takaku, A., Endo, S., Kumanishi, T., 2002.

Aberrant nuclear factor-kB activity and its participation in the growth of human

malignant astrocytoma. J. Neurosurg. 96, 909–917.

https://doi.org/10.3171/jns.2002.96.5.0909.

Ostrom, Q.T., Patil, N., Cioffi, G., Waite, K., Kruchko, C., Barnholtz-Sloan J.S., 2020.

CBTRUS Statistical Report: Primary brain and other central nervous System tumors

diagnosed in the United States in 2013–2017. Neuro-Oncol., 22, iv1–iv96.

https://doi.org/10.1093/neuonc/noaa200.

Panche, A.N., Diwan, A.D., Chandra, S.R., 2016. Flavonoids: An overview. J. Nutr.

Sci. 5, e47. https://doi:10.1017/jns.2016.41.

Park, M.H., Hong, J.E., Park, E.S., Yoon, H.S., Seo, D.W., Hyun, B.K., 2015. Anticancer

effect of tectochrysin in colon cancer cell via suppression of NF-kappaB activity and

enhancement of death receptor expression. Mol. Cancer. 14:124

https://doi:10.1186/s12943-015-0377-2.

Pinheiro, K. V., Alves, C., Buendia, M., Gil, M. S., Thomaz, A., Schwartsmann, G., De

Farias, C. B., & Roesler, R., 2017. Targeting tyrosine receptor kinase B in gliomas.

Neuro. Oncol. 19, 138–139. https://doi.org/10.1093/neuonc/now199.

Pratheeshkumar, P., Budhraja, A., Son, Y.O., Wang, X., Zhang, Z., Ding, S., Wang, L.,

Hitron, A., Lee, J.C., Xu, M., Chen, G., Luo, J., Shi, X., 2012. Quercetin inhibits

angiogenesis mediated human prostate tumor growth by targeting VEGFR-2

regulated AKT/mTOR/P70S6K signaling pathways. PLoS ONE, 7, e47516.

https://doi.org/10.1371/journal.pone.0047516.

Raychaudhuri, B., Han, Y., Lu, T., Vogelbaum, M.A., 2007. Aberrant constitutive

activation of nuclear factor kappaB in glioblastoma multiforme drives invasive

phenotype. J. Neurooncol. 85, 39–47. https://doi.org/10.1007/s11060-007-9390-7.

Rea, K.A., Casaretto, J.A., Al-Abdul-Wahid, M.S., Sukumaran, A., Geddes-McAlister, J.,

Rothstein, S.J., & Akhtar, T.A., 2019. Biosynthesis of cannflavins A and B from

Cannabis sativa L. *Phytochemistry*, *164*, 162–171.

https://doi.org/10.1016/j.phytochem.2019.05.009.

Srivastava, S., Somasagara, R.R., Hegde, M., Nishana, M., Tadi, S.K., Srivastava, M.,

Choudhary, B., Raghavan, S.C., 2016. Quercetin, a natural flavonoid interacts with

DNA, arrests cell cycle and causes tumor regression by activating mitochondrial

pathway of apoptosis. Sci. Rep. 6, 24049. https://doi.org/10.1038/srep24049.

Stupp, R., Mason, W.P., Bent, M.J. van den, Weller M., Fisher, B., Taphoorn, M.J.B.,

Belanger, K., Brandes, A.A., Marosi, C., Bogdahn, U., Curschmann, J., Janzer, R.C.,

Ludwin, S.K., Gorlia, T., Allgeier, A., Lacombe, D., Cairncross, J.G., Eisenhauer, E.,

Mirimanoff, R.O., European Organisation for Research and Treatment of Cancer

Brain Tumor and Radiotherapy Groups; National Cancer Institute of Canada Clinical

Trials Group., 2005. Radiotherapy plus concomitant and adjuvant temozolomide for

glioblastoma. N. Engl. J. Med. 2352, 987–996.

https://doi.org/10.1056/nejmoa043330.

Stupp, R., Hegi, M.E., Mason, W.P., Bent, M.J. van den, Taphoorn, M.J., Janzer, R.C.,

Ludwin, S.K., Allgeier, A, Fisher, B., Belanger, K., Hau, P., Brandes, A.A.,

Gijtenbeek, J., Marosi, C., Vecht, C.J., Mokhtari, K., Wesseling, P., Villa, S.,

Eisenhauer, E., Gorlia, T., Weller, M., Lacombe, D., Cairncross, J.G., Mirimanoff,

R.O; European Organisation for Research and Treatment of Cancer Brain Tumour

and Radiation Oncology Groups; National Cancer Institute of Canada Clinical Trials

Group, 2009. Effects of radiotherapy with concomitant and adjuvant temozolomide

versus radiotherapy alone on survival in glioblastoma in a randomised phase III

study: 5-year analysis of the EORTC-NCIC trial. Lancet Oncol. 10, 459–466.

https://doi.org/10.1016/s1470-2045(09)70025-7.

Taniguchi, K., Karin, M., 2018. NF-κB, inflammation, immunity and cancer: coming of

age. Nat. Rev. Immunol. 18, 309–324. https://doi.org/10.1038/nri.2017.142.

Tilak, M., Alural, B., Holborn, J., Lalonde, J., Jones, N., 2021a. Receptor tyrosine kinase

signaling and targeting in glioblastoma. Int. J. Mol. Sci. 22, 1831.

https://doi:10.3390/ijms22041831.

Tilak, M., Alural, B., Wismer, S.E., Brasher, M.I., New, L.A., Sheridan, S.D., Perlis,

R.H., Coppolino, M.G., Lalonde, J., Jones, N., 2021b. Adaptor protein ShcD/SHC4

interacts with tie2 receptor to synergistically promote glioma cell invasion. Mol.

Cancer Res. 19, 757–770. https://doi.org/10.1158/1541-7786.MCR-20-0188.

Tomko, A.M., Whynot, E.G., Dupré, D.J., 2022. Anti-cancer properties of cannflavin A

and potential synergistic effects with gemcitabine, cisplatin, and cannabinoids in

bladder cancer. J. Cannabis Res. 4, 41. https://doi.org/10.1186/s42238-022-00151-y.

Tseng, T.H., Chien, M.H., Lin, W.L., Wen, Y.C., Chow, J.M., Chen, C.K., Kuo, T.C.,

Lee, W.J., 2017. Inhibition of MDA-MB-231 breast cancer cell proliferation and

tumor growth by apigenin through induction of G2/M arrest and histone H3

acetylation-mediated p21WAF1/CIP1 expression. Environ. Toxicol. 32, 434–444.

https://doi.org/10.1002/tox.22247.

Vinci, M., Box, C., Eccles, S.A., 2015. Three-dimensional (3D) tumor spheroid invasion

assay. J. Vis. Exp. 99, e52686. https://doi.org/10.3791/52686.

Wang, X., Sun, Y., Zhang, D.Y., Ming, G., Song, H., 2023. Glioblastoma modeling with

3D organoids: progress and challenges. Oxf. Open Neurosci. 2, kvad008.

https://doi:10.1093/oons/kvad008.

Wang, H., Wang, H., Zhang, W., Huang, H.J., Liao, W.S., Fuller, G.N. 2004. Analysis of

the activation status of Akt, NFkappaB, and Stat3 in human diffuse gliomas. Lab

Invest. 84, 941–951. https://doi.org/10.1038/labinvest.3700123.

Weller M, Cloughesy T, Perry J.R., Wick W., 2013. Standards of care for treatment of

recurrent glioblastoma—are we there yet? Neuro Oncol. 15, 4–27.

https://doi.org/10.1093/neuonc/nos273.

Werz, O., Seegers, J., Schaible, A.M., Weinigel, C., Barz, D., Koeberle, A., Allegrone,

G., Pollastro, F., Zampieri, L., Grassi, G., Appendino, G., 2014. Cannflavins from

hemp sprouts, a novel cannabinoid-free hemp food product, target microsomal

prostaglandin E_2 synthase-1 and 5-lipoxygenase. PharmaNutrition. 2, 53–60.

https://doi.org/10.1016/j.phanu.2014.05.001.

Xu, M. J., Wu, B., Ding, T., Chu, J. H., Li, C. Y., Zhang, J., Wu, T., Wu, J., Liu, S. J.,

Liu, S. L., Ju, W. Z., Li, P., 2012. Simultaneous characterization of prenylated

flavonoids and isoflavonoids in Psoralea corylifolia L. by liquid chromatography

with diode-array detection and quadupole time-of-flight mass spectrometry. Rapid

Commun. Mass Spectrom. 26, 2343–2358. https://doi.org/10.1002/rcm.6361.

Yang, X., Jiang, Y., He, J., Sun, J., Chen, F., Zhang, M., Yang, B., 2015. Prenylated

flavonoids, promising nutraceuticals with impressive biological activities. Trends

Food Sci. Technol. 44, 93e104. https://doi.org/10.1016/j.tifs.2015.03.007.

Zhang, Q., Cheng, G., Qiu, H., Wang, Y., Wang, J., Xu, H., Zhang, T., Liu, L., Tao, Y.,

Ren, Z., 2017. Expression of prostate stem cell antigen is downregulated during

flavonoid-induced cytotoxicity in prostate cancer cells. Exp. Ther. Med. 14, 1795–

1801. https://doi.org/10.3892/etm.2017.4638.

Zhang, T., Ma, C., Zhang, Z., Zhang, H., Hu, H., 2021 NF-κB signaling in inflammation

and cancer. MedComm. 2, 618–653. https://doi.org/10.1002/mco2.104.

Zhu, L., Xue, L., 2019. Kaempferol suppresses proliferation and induces cell cycle arrest,

apoptosis, and DNA damage in breast cancer cells. Oncol. Res. 27, 629–634.

https://doi.org/10.3727/096504018X15228018559434.

FIGURE 1

Figure 1. Cannflavin B reduces GBM cell viability. A172 **(A-C)** and U-87 **(D-F)** cells

were treated with increased concentrations of cannflavin A (1-20 µM), cannflavin B (1-20

µM), or ANA-12 (1-50 µM) and cell viability was measured at 24-, 48-, and 72-hour time-

points using the resazurin reduction assay. DMSO was used as a vehicle control.

Application of cannflavin B and ANA-12, but not cannflavin A, reduced the viability of

both A-172 and U-87 GBM cells in a dose- and time-dependent manner. A one-way

ANOVA was used to analyze the data from each cell line, treatment, and time point

independently with DMSO as the reference control. Tukey's honestly significant

difference (HSD) post hoc test: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Graphs represent mean ± SEM ($n = 3$-4).

FIGURE 2

1003

Figure 2. Cannflavins A and B did not affect cell cycle distribution in GBM cell lines.

A-172 **(A-B)** and U-87 **(C-D)** DNA content was determined by gating on the G1, S, and

G2/M phases using flow cytometry and analyzed using the Watson Pragmatic algorithm to

distribute cell cycle phases. A172 **(B)** and U-87 **(D)** cell cycle data are expressed as mean

$\pm$ SEM (n = 3).

FIGURE 3

1010

Figure 3. Cannflavins A and B suppress A-172 cell migration. **(A)** Representative phase-

contrast images of scratch assay showing A-172 GBM cancer cells within a gap created

with a pipette tip at 0 hour and 48 hours. Dashed yellow lines indicate the initial wound

edges. **(B)** Quantification of the binary wound area fraction over time, showing delayed

wound closure in inhibitor-treated conditions. **(C)** Normalized wound area fraction at 24

hours and 48 hours. A two-way ANOVA reveals a significant effect of time ($F_{1,218} = 133.3$,

$p < 0.0001$) as well as treatment ($F_{3,218} = 18.37$, $p < 0.0001$). There is no significant

interaction between time and treatment ($F_{3,218} = 2.094$, $p = 0.1019$). Post hoc comparisons

between DMSO control and each treatment means were conducted using Dunnett's

multiple comparisons. All treatments at all time points produced a significant reduction in

cell migration. **(D)** Percentage of wound closure at 24 hours and 48 hours. A two-way

ANOVA reveals a significant effect of time ($F_{1,218} = 104.2$, $p < 0.0001$) as well as treatment

($F_{3,218} = 16.45$, $p < 0.0001$). There is no significant interaction between time and treatment

($F_{3,218} = 1.885$, $p = 0.133$). Post hoc comparisons between DMSO control and each

treatment means were conducted using Dunnett's multiple comparisons. Cannflavin B

significantly limited closure at both time points, while cannflavin A was significant only

at 48 hours. ANA-12 did not cause a significant effect. **(E)** Total cell counts within the

imaging field at 24 hours and 48 hours. A two-way ANOVA reveals a significant effect of

time ($F_{1,154} = 92.25$, $p < 0.0001$) as well as treatment ($F_{3,154} = 4.46$, $p = 0.0049$). There is

no significant interaction between time and treatment ($F_{3,154} = 0.2556$, $p = 0.8573$). Post

hoc comparisons between DMSO control and each treatment means were conducted using

Dunnett's multiple comparisons. Significantly less cells were found in the wound area with

cannflavin B at 48 hours. No other comparisons were significantly different. All data

represented as mean ± SEM (3 biological replicates with 6-8 images per well; * $p < 0.05$,

** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).

FIGURE 4

1037

Figure 4. Cannflavin B inhibits A-172 glioblastoma cell invasion in a transwell assay.

(A) Representative images of stained migrated A-172 cells following treatment with

DMSO or cannflavin B (10 µM). **(B)** Quantification of migration using crystal violet

elution measured by absorbance at 562 nm. **(C)** Percent invasion between DMSO control

and cannflavin B, showing a significant reduction in cannflavin B-treated cells. Data

represent mean ± SEM, with statistical significance determined by an unpaired two-tailed

t-test (** $p < 0.01$).

FIGURE 5

1046

Figure 5. Cannflavins A and B inhibit the outgrowth of U-87 tumorspheres. **(A)**

Representative phase-contrast images of U-87 tumorspheres at 0 hour and 72 hours post-

treatment with DMSO (vehicle), ANA-12 (10 µM), cannflavin A (10 µM), or cannflavin

B (5 µM) in a tumorsphere invasion assay. Scale bar: 500 µm. The white outline indicates

the borders of tumorsphere outgrowth. **(B)** Quantification of relative tumorsphere area

expansion over time, normalized to the vehicle control. A two-way ANOVA reveals a

significant effect of time ($F_{3,32}$ = 5.949, $p < 0.0024$) as well as treatment ($F_{3,32}$ = 10.13, p

< 0.0001). There is no significant interaction between time and treatment ($F_{9,32}$ = 1.765, p

= 0.1145). Post hoc comparisons between DMSO control and each treatment means were

conducted using Dunnett's multiple comparisons. Ratio was significantly lower at all three

time points for cannflavin A, while cannflavin B was significant only at 72 hours. ANA-

12 did not cause a significant effect. **(C)** Relative tumorsphere radius expansion over time,

showing significant inhibition of invasion upon cannflavin treatment. A two-way ANOVA

reveals a significant effect of time ($F_{3,32}$ = 46.18, $p < 0.0001$) as well as treatment ($F_{3,32}$ =

40.07, $p < 0.0001$). Notably, there is a significant interaction between time and treatment

($F_{9,32}$ = 8.749, $p < 0.0001$), reflecting an earlier effect with cannflavin A, followed by

cannflavin B, and finally ANA-12. Post hoc comparisons between DMSO control and each

treatment means were conducted using Dunnett's multiple comparisons. Data represent

mean ± SEM (n = 3; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$; n.s., not

significant).

FIGURE 6

1068

Figure 6. Cannflavin B reduces the size of attached tumorspheres to a cerebral organoid.

(A) Schematic representation of cerebral organoid and U-87 tumorspheres co-culture

model. Cerebral organoids matured for 60 days before introducing tumorspheres prepared

with U-87 cells expressing GFP. **(B)** Two experimental timelines were tested: 11 days of

co-culture followed by 3 days of applying cannflavin B to the culture media (final

concentration of 5 µM), and 4 days of co-culture followed by 10 days of applying

cannflavin B (final concentration of 5 µM). **(C-F)** Representative capture of organoid with

attached tumorsphere(s) for all conditions. **(G-H)** Proportion of GFP-positive pixels

corresponding to U-87 cells within the cerebral organoid section area for the 3-day **(G)** and

10-day **(H)** conditions. **(I-J)** Estimation of tumorsphere size in the 3-day **(I)** and 10-day **(J)**

conditions. The average tumorsphere area (GFP-positive cells, green) was significantly

smaller after 10 days of cannflavin B treatment than in the corresponding DMSO control.

Data represent mean ± SEM, with statistical significance determined by an unpaired two-

tailed *t*-test (*** $p < 0.001$).

FIGURE 7

1084

Figure 7. Cannflavin A and cannflavin B affect TrkB Y705 activation site but not

downstream signaling in A-172 cells. Western blot analysis from A-172 cells treated for 6

hours with cannflavin A **(A-E)** and cannflavin B **(F-J)** at various concentrations.

Corresponding densitometry quantifications reveal a significant reduction in P-TrkB Y705

signal, but not in P-MAPK and P-AKT. GAPDH (glyceraldehyde-3-phosphate

dehydrogenase) was used as a loading control. Graphs show the mean $\pm$ SEM ($n = 4$) of

the target/GAPDH ratio for cells treated as in **(A)** and **(F)**. Post hoc comparisons between

DMSO control and each treatment means were conducted using Dunnett's multiple

comparisons (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

Figure 8

1095

Figure 8. Cannflavin A and cannflavin B slightly increase TrkB Y705 activation site but

not change downstream signaling in U-87 cells. Western blot analysis from U-87 cells

treated for 6 hours with cannflavin A **(A-E)** and cannflavin B **(F-J)** at various

concentrations. Corresponding densitometry quantifications reveal a significant reduction

in P-TrkB Y705 signal, but not in P-MAPK and P-AKT. GAPDH (glyceraldehyde-3-

phosphate dehydrogenase) was used as a loading control. Graphs show the mean ± SEM

($n = 3$) of the target/GAPDH ratio for cells treated as in **(A)** and **(F)**. Post hoc comparisons

between DMSO control and each treatment means were conducted using Dunnett's

multiple comparisons (* $p < 0.05$).

Supplementary Information for

Preclinical Study of Cannflavins A and B Action Against Glioblastoma Cells

Jennifer Holborn [a], Hannah Robeson [a,1], Ellis Chartley [a,1], Tiana Gluscevic [a,b,1], Adina Borenstein [a], Colby Perrin [a], Begüm Alural [a], Himain Perera [a], Tariq A. Akhtar [a], Nina Jones [a], Jasmin Lalonde [a,*]

* Correspondence to: jlalon07@uoguelph.ca

This PDF file includes:

SUPPLEMENTARY FIGURE 1

Supplementary Figure 1. Members of the cannflavin family. Position of each ring is labeled for cannflavin A.

SUPPLEMENTARY FIGURE 2

Supplementary Figure 2. Effect of chrysoeriol and techtochrysin on U-87 GBM cell survival. **(A)** Chemical structure of chrysoeriol and **(B)** techtochrysin. As proposed by Real and colleagues (2019), *C. sativa* prenyltransferase 3 (CsPT3) catalyzes the biosynthesis of cannflavins from chrysoeriol. **(C-D)** U-87 cells were treated with increased concentrations of chrysoeriol (1-20 µM) or techtochrysin (5-80 µM), and cell viability was measured at the indicated time-points using the resazurin reduction assay. DMSO was used as a vehicle control. Application of chrysoeriol, but not techtochrysin, reduced the viability of U-87 GBM cells in a dose- and time-dependent manner. A one-way ANOVA was used to analyze the data from each cell line, treatment, and time independently with DMSO as the reference control. Tukey's honestly significant

difference (HSD) post hoc test: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Graphs represent

mean ± SEM ($n = 3$).

SUPPLEMENTARY FIGURE 3

A   A-172 LDH Release Assay (24 h)

	% LDH release	SEM
Lysis buffer Positive control	100	–
Vehicle (DMSO) Negative control	1.77	0.65
Cannflavin A 10 µM 20 µM	3.57 2.54	0.84 0.66
Cannflavin B 10 µM 20 µM	4.43 5.38	1.43 1.44
ANA-12 10 µM 50 µM	2.01 11.35	0.48 3.63

B   U-87 LDH Release Assay (24 h)

	% LDH release	SEM
Lysis buffer Positive control	100	–
Vehicle (DMSO) Negative control	1.57	0.29
Cannflavin A 10 µM 20 µM	0.96 1.03	0.09 0.08
Cannflavin B 10 µM 20 µM	0.81 0.88	0.05 0.07
ANA-12 10 µM 50 µM	1.61 10.71	0.26 0.64

C   TUNEL (fluorescein)   DAPI   Light

Supplementary Figure 3. Effects of cannflavins A and B on LDH release and DNA fragmentation. **(A)** A-172 and **(B)** U-87 GBM cells were treated with cannflavin A or cannflavin B at 10 and 20 µM final concentrations, or ANA-12 (10 and 50 µM final concentration), and percent LDH release into cell culture media was quantified using the Invitrogen CyQUANT LDH Cytotoxicity Assay Kit. **(C)** Representative immunofluorescent images of A-172 GBM cells after 24 hours of treatment with cannflavins (10 µM final concentration) probed using TUNEL (fluorescein, green signal represent the occurrence of double-stranded DNA breaks). Nuclei were stained with DAPI (middle panels). Hydrogen peroxide (H_2O_2) was used as a positive control (applied directly to cell culture media for 10 minutes at a final concentration of 100 µM), and DMSO as the vehicle control for cannflavins. Arrowheads in the positive control condition show fragmented nuclei indicative of apoptosis. Scale bar = 50 µm.

Supplementary Figure 4. Level of phospho-TrkB Y705 between A-172 and U-87 untreated cells. **(A)** Representative western blots for phospho-TrkB Y705 and downstream signaling effectors. **(B)** Densitometry quantification shows a significantly higher abundance of phospho-TrkB Y705 in A-172 cells than in U-87 cells. * $p < 0.05$, two-tailed unpaired t-test. Graph represents mean ± SEM ($n = 4$).

Supplementary Figure 5. Graphical abstract.

SUPPLEMENTARY TABLE

A-172		U-87 MG	
Condition	IC$_{50}$ (μM)	Condition	IC$_{50}$ (μM)
Cannflavin A (6 h)	>200	Cannflavin A (6 h)	>200
Cannflavin A (24 h)	173.4	Cannflavin A (24 h)	>200
Cannflavin A (48 h)	88.5	Cannflavin A (48 h)	187.9
Cannflavin B (6 h)	76.1	Cannflavin B (6 h)	63.9
Cannflavin B (24 h)	10.2	Cannflavin B (24 h)	20.2
Cannflavin B (48 h)	4.2	Cannflavin B (48 h)	25.1
ANA-12 (6 h)	168.8	ANA-12 (6 h)	156.2
ANA-12 (24 h)	39	ANA-12 (24 h)	48.0
ANA-12 (48 h)	27.9	ANA-12 (48 h)	49.2

Supplementary Table. The estimated IC$_{50}$ for the survival assay is shown in Figure 1.

SUPPLEMENTARY VIDEO

Supplementary Video. Scratch migration assay video. The green lines represent the initial boundaries ($t = 0$) of the scratched wound.