Distinguishing Protein and Gene Delivery Enables Characterization and Bioengineering of

Extracellular Vesicle-Adeno-Associated Virus Vectors

Jonathan D. Boucher[1,2,3], Devin M. Stranford[1,3,4], Hailey I. Edelstein[1,3,4], Danielle Tullman-Ercek[1-3,5], Neha

P. Kamat[3,5,6], Joshua N. Leonard[1-5,7*]

[1] Department of Chemical and Biological Engineering, Northwestern University, Evanston, Illinois 60208,

United States

[2] Interdisciplinary Biological Sciences Program, Northwestern University, Evanston, Illinois 60208, United

States

[3] Center for Synthetic Biology, Northwestern University, Evanston, Illinois 60208, United States

[4] Syenex Inc, Evanston, Illinois 60201, United States

[5] Chemistry of Life Processes Institute, Northwestern University, Evanston, Illinois 60208, United States

[6] Department of Biomedical Engineering, Northwestern University, Evanston, IL, 60208 USA

[7] Member, Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Evanston, Illinois

60208, United States

Contact Information

*Corresponding author: Joshua N. Leonard, j-leonard@northwestern.edu

ABSTRACT

Adeno-associated virus (AAV) gene therapies have achieved clinical success, with multiple products reaching regulatory approval. Encapsulation of AAV vectors within engineered extracellular vesicles (EVs) is an emerging strategy which could help overcome challenges including pre-existing anti-capsid immunity and the need for controlling targeting and tropism. To guide the development of EV-AAV technologies, we developed an assay for quantifying and controlling for the contribution of pseudotransduction to evaluations of EV-AAV-mediated gene delivery. We developed an AAV vector that switches its transgene output from one reporter to another when acted upon by Cre recombinase expressed in a recipient cell. Using this platform, we investigated EV-AAV transduction as a function of various engineered EV surface modifications. For actively endocytic cells (HEK293FTs), modifications that enhance EV uptake and membrane fusion influence protein delivery but not gene delivery. Conversely, in less endocytic Jurkat T cells, modifications enhancing EV uptake improved both protein and gene delivery. These conclusions held across multiple AAV serotypes. Our results resolve apparent conflicts in prior reports and suggest that effects of enhancing uptake and membrane fusion of EV-AAV vectors are recipient cell type specific. The methods developed here unambiguously dissect EV-AAV transduction mechanisms and can guide future bioengineering of EV-AAV vectors.

INTRODUCTION

Adeno-associated virus (AAV) gene therapies are poised to revolutionize the treatment of genetic diseases. Over the last decade, at least seven AAV-based gene therapies have been approved by the US Food and Drug Administration, with hundreds of clinical trials ongoing.[1] Current efforts focus on challenges such as evading preexisting humoral immunity[2], and reducing off-target tissue transduction[3], which both necessitate higher vector doses that are associated with adverse effects[4,5]. One potential strategy for addressing these needs is encapsulation of AAV within engineered extracellular vesicles (EVs).

EVs are nanoscale, membrane-bound particles secreted by all cells and include exosomes, microvesicles, and particles such as apoptotic bodies.[6] Exosomes are formed through the invagination of endosomal compartments, resulting in multivesicular bodies that fuse with the plasma membrane to release exosomes into the extracellular space. These vesicles protect their cargo—cytoplasmic RNA, proteins, and

other biomolecules—from degradation by nucleases and proteases.[7] Microvesicles bud from the cell

surface and encapsulate a distinct set of biomolecules.[6] When recipient cells take up EVs, some cargo

molecules can be released into the cytoplasm to impact cell state.[8,9] Because EVs mediate intercellular

communication, exhibit lower toxicity and immunogenicity than synthetic delivery vehicles, and have longer

circulation times *in vivo*, they are attractive candidates for therapeutic cargo delivery.[10-12]

An interesting observation is that when AAV particles are assembled in producer cells, secreted

EVs can incorporate AAV particles capable of conferring gene expression in recipient cells, and such EV-

AAV vehicles have distinct properties that are useful for gene delivery.[13] EV-AAVs can mediate gene

delivery to the mouse brain, even when using serotypes not generally able to cross the blood-brain barrier.[14]

EV-AAV1 and EV-AAV9 outperformed their free AAV (standard, non-enveloped AAV particle) counterparts

in transduction of cochlear hair cells in explant cultures.[15] EV-AAV1 encoding *Lhfpl5,* a protein required for

mechanotransduction in outer and inner hair cells and for which loss leads to profound deafness, improved

hearing and movement abnormalities in *Lhfpl5$^{-/-}$* mice.[15] Upon intravitreal injection, EV-AAV2 outperformed

free AAV2 in retinal transduction via delivery of GFP, increasing transduction to cell types that are

transduced by free AAV2 as well as increasing the extent of transduction across different regions of the

retina.[16] EV-AAV8 encoding hF.IX enhanced correction of hemophilia B phenotypes compared to standard

AAV8 vectors in C57BL/6 *F9−/−* mice, demonstrating enhanced liver targeting after intravenous injection.[17]

Both standard AAV8 and EV-AAV8 can transduce a variety of immune cell populations, including CD4^{+} T

cells, CD8^{+} T cells, B cells, macrophages, and dendritic cells after systemic delivery in mice.[18] EV-AAV8

can mediate the expression of human interleukin-2 receptor alpha chain (IL-2Rα) in multiple lymphocyte

populations, suggesting that EV-AAVs could be utilized for *in vivo* cell engineering.[18] In a lung cancer

xenograft mouse model, local injection of EV-AAV6 was much more potent in transducing tumor cells

compared to free AAV6.[19] Upon intratracheal injection in C57BL/6 mice, EV-AAV6 enhanced transduction

of the lung compared to either free AAV6 or a mixture of free AAV6 and EVs, demonstrating that the

enhanced lung transduction by EV-AAV6 was attributable to the direct association of EVs and AAV6.[20] EV-

AAV9 delivering *SERCA2a* significantly improved cardiac function in a mouse model of myocardial

infarction compared to free AAV9.[21] In addition to improving transduction, EV-AAVs resist neutralization by

AAV specific antibodies, suggesting that encapsulation can help address needs for AAV vector

delivery.[14,17,19-25] Moreover, since EVs have been engineered to target specific cell types[26-29], it is possible

that such methods could target EV-AAVs as well.

81   Although EV-AAV vectors are promising, several key mechanistic questions are incompletely

understood. First, the functional consequences of surface-functionalizing EV-AAV vectors on recipient cell

binding and particle uptake are unclear. Second, reports conflict as to how increasing EV-endosomal fusion

modulates transduction by EV-AAV. An early study suggested that functionalizing EV-AAVs by incorporating

the low pH-activated fusogen vesicular stomatitis virus glycoprotein (VSV-G) on the EV surface can diminish

transduction of HEK293T and U87 glioma cells by AAV2, compared to EV-AAV2 lacking VSV-G.[30] This is

surprising as increasing uptake (via VSV-G binding to its receptor, low density lipoprotein receptor, LDL-

R[31]) and fusion-mediated entry of the AAV capsid into the cytosol may be expected to increase transduction.

While a possible explanation is that AAV2 requires processing in the endosome—capsid cleavage by the

endosomal cysteine proteases cathepsins B and L[32], autoproteolytic processing of the capsid at low

pH[33], and exposure of the VP1/2 N-terminal region exposing a phospholipase A2 domain that helps the

virion escape the endosome and traffic to the nucleus[34,35]—that mechanism is untested in EV-AAV delivery.

VSV-G display on EV-AAV2 showed no effect on gene delivery to several other glioma cell lines compared

to non-targeted EV-AAV2[30], suggesting the role of VSV-G may vary by recipient cell type. A related line of

evidence is that display of rabies virus glycoprotein (RVG) on the surface of EV-AAV9 increased

transduction of SH-SY5Y cells and gene delivery to the brain of mice compared to non-targeted EV-AAV9.[22]

RVG binds to the nicotinic acetylcholine receptor (nAChR), neural cell adhesion molecule (NCAM), p75

neurotrophin receptor, integrin β1, and transferrin receptor 1 (TfR1), promoting clathrin-mediated

endocytosis of the particle.[36] Much like VSV-G, upon encountering the low pH of the endosome, RVG

undergoes conformational changes that promote endosomal membrane fusion and escape.[37,38] Given the

similarity in mechanisms used by VSV-G and RVG, the differences in transduction when displayed on EV-

AAVs is surprising and may indicate recipient cell type and/or AAV serotype-specific effects. Thus, although

EV-AAV vectors are promising, successful bioengineering of such vectors would benefit from mechanistic

understanding of delivery mechanisms.

105   A key technical challenge when evaluating engineered EV-AAV-mediated delivery is distinguishing

delivery of protein originating from transgene expression in producer cells (i.e., "pseudotransduction"), from

gene delivery to and resulting transgene expression in recipient cells. This challenge is analogous to the

pseudotransduction observed with lentiviral and retroviral vectors, which are also enveloped and can

incorporate cytosolic proteins from producer cells.[39-42] In certain applications, pseudotransduction may be

useful, as demonstrated in dendritic cell–targeted lentiviral vectors for vaccination, in which incorporated

protein antigens are employed to train induced immune reactions.[43] However, in applications where

quantification of gene delivery or transduction is desired, pseudotransduction can inflate the rates of

transduction reported. Informed in part by prior observations that protein delivery can complicate evaluation

of EV-mediated mRNA delivery[44,45], we hypothesized that controlling for pseudotransduction may be

particularly crucial for measuring EV-AAV transduction because achieving comparable rates of transduction

typically requires higher numbers of particles for EV-AAV compared to lentiviral vectors. Thus,

distinguishing transduction from pseudotransduction (protein delivery) is critical for the engineering and

evaluation of EV-AAV vectors.

In this study, we developed and validated a quantitative assay for disambiguating

pseudotransduction from transduction by EV-AAVs and applied this method to investigate several biological

hypotheses that inform bioengineering. Our assay comprises a modified stoplight system, in which Cre

recombinase is expressed in recipient cells where it can modify an incoming engineered AAV vector to shift

its transgene expression from one fluorescent protein to another. Using this approach, pseudotransduction

(protein delivery) and transduction (gene delivery) can be measured separately. With this system, we

compared various EV surface functionalization strategies that drive uptake and/or membrane fusion with

recipient cells and determined how these engineering choices impact protein and gene delivery. Such

analysis revealed new insights about EV-AAV transduction that would otherwise be obscured by

pseudotransduction. Our assay enables generation and testing of new mechanistic hypotheses for studying

EV-AAV biology and bioengineering these vectors.

METHODS

General DNA assembly

Plasmids were constructed using standard molecular biology techniques. Plasmid assembly was performed

via restriction enzyme cloning. Plasmids were transformed into TOP10 competent Escherichia coli (Thermo

Fisher) and grown at 37 °C. AAV vector (transfer) plasmids were transformed into NEB Stable competent

Escherichia coli (New England Biolabs C3040) and grown at 37 °C. Plasmid cloning was performed

primarily using standard restriction enzyme cloning with restriction enzymes (NEB), T4 DNA Ligase (NEB

M0202L), and Antarctic Phosphatase (NEB M0289L). Plasmids with pcDNA-based backbones were

transformed into chemically competent TOP10 *Escherichia coli* (Thermo Fisher C404010), and cells were

grown at 37°C. Plasmids with AAV transgene vector backbones were transformed into chemically

competent NEB Stable *Escherichia coli* (NEB C3040H), and cells were grown at 30°C.

Source vectors for DNA assembly

The AAV transfer vector (pAAV-GFP SN), and helper plasmid (pHelper) were gifts from David Schaffer

(University of California, Berkeley). The LumiScarlet gene was PCR amplified from pcDNA3-LumiScarlet,

which was a gift from Huiwang Ai (Addgene plasmid # 126623 ; http://n2t.net/addgene:126623 ;

RRID:Addgene_126623)[46], and cloned into the pAAVGFP SN plasmid using AgeI and NotI. pOTTC1032 -

pAAV EF1a Nuc-flox(mCherry)-EGFP (pJB037) was a gift from Brandon Harvey (Addgene plasmid #

112677 ; http://n2t.net/addgene:112677 ; RRID:Addgene_112677).[47] Gene strings for the other Cre-

responsive vectors (pJB038 - No NLS and eGFP, pJB040 - NLS and mNeonGreen, and pJB041 No NLS

and mNeonGreen) were codon optimized and synthesized by ThermoFisher (GeneArt), then cloned into

the pOCTTC1032 plasmid using KpnI and EcoRI restriction enzymes. RepCap plasmids for AAV2

(pRep2Cap2) and AAV6 (pRep2Cap6) were gifts from the Vector Core at the University of Pennsylvania

(Penn Vector Core (RRID: SCR_022432)). pcDNA is plasmid pPD005 (Addgene plasmid # 138749 ;

http://n2t.net/addgene:138749 ; RRID:Addgene_138749).[48] psPAX2 and pMD2.G plasmids were gifted by

William Miller from Northwestern University. pHIE822 was created by introducing point mutations (K47Q,

R354A) in the VSV-G-encoding gene of pMD2.G using site-directed mutagenesis by PCR. pCMV-VSV-

G(P127D)-Myc (pJB042) was a gift from Wesley Sundquist (Addgene plasmid # 80055 ;

http://n2t.net/addgene:80055 ; RRID:Addgene_80055).[49] pHIE963 was created using standard restriction

enzyme cloning to insert a Cre recombinase gene into a third-generation lentiviral transfer vector provided

by Twist Bioscience (pTwist Lenti SFFV PuroR). The Cre recombinase gene was amplified by PCR from

pBS185 CMV-Cre, which was a gift from Brian Sauer (Addgene plasmid # 11916

; http://n2t.net/addgene:11916 ; RRID:Addgene_11916).[50] The Ultra-HEK plasmid is a commercial product

made by Syenex. The plasmids used to produce Ultra-HEK lentiviral vectors are Syenex plasmids: SNX-

mNG-TV3, SNX-Gagpol, SNX-Rev, and SNX-VSV-G. A list of plasmids utilized in this study can be found

in **Supplementary Data 1**. Plasmid sequences can be found in **Supplementary Data 2**.

Plasmid preparation

TOP10 or NEB Stable *E. coli* were grown overnight, shaking at 37°C or 30°C, respectively, in 50-100 mL of

LB media with the appropriate selective antibiotic. The next day, DNA was prepped using a ZymoPURE II

Plasmid Midiprep Kit (Zymo D4201) following the manufacturer's instructions. DNA purity and

concentrations for relevant experiments were measured with a NanoDrop 2000 (Thermo Fisher Scientific).

Cell culture

FreeStyle 293-F cells (Gibco R79007) cells were cultured in FreeStyle 293 Expression Medium (Gibco

12338018) supplemented with 1% penicillin–streptomycin (Gibco 15140-122). HEK293FT cells (Thermo

Fisher R70007) were cultured in Dulbecco's modified Eagle medium (DMEM, Gibco 31600-091)

supplemented with 10% fetal bovine serum (FBS) (Gibco 16140-071), 1% penicillin–streptomycin (Gibco

15140-122) and 4 mM additional L-glutamine (Gibco 25030-081). HEK293T Lenti-X cells (Takara 632180)

were cultured in HEK293FT media formulation supplemented with 1 mm sodium pyruvate from Gibco

(11360070). Jurkat T cells (ATCC TIB-152) were cultured in Roswell Park Memorial Institute medium (RPMI

1640, Gibco 31800-105) supplemented with 10% FBS, 1% penicillin–streptomycin, and 4 mM L-glutamine.

Sublines generated from these cell lines were cultured in the same way. Cells were subcultured at a 1:5 or

1:10 ratio every 2–3 days, using trypsin–EDTA (Gibco 25300-054) to remove adherent cells from the plate.

Freestyle 293-F cells were maintained at 37 °C with 8% CO_2 rotating at 135 rpm on a shaking platform

(Infors HT Multitron – I8000A) with orbital diameter 25 mm. HEK293FT, HEK293T Lenti-X, and Jurkat cells

were maintained at 37 °C with 5% CO_2. Freestyle 293F, HEK293FT, HEK293T Lenti-X and Jurkat cells

tested negative for mycoplasma with the MycoAlert Mycoplasma Detection Kit (Lonza LT07-318). HEK293T

Lenti-X and Jurkat cells tested negative for mycoplasma with the MycoAlert Mycoplasma Detection Kit

(Lonza LT07-318).

Cell line generation

To generate lentivirus, HEK293T Lenti-X cells were plated in 10 cm dishes at a density of 5×10^6 cells per

dish (6.25×10^5 cells per ml). About 24 h later for Lenti-X, cells were transfected with 10 µg of lentiviral

transfer vector plasmid, 8 µg of psPAX2 second generation lentiviral packaging plasmid, and 3 µg pMD2.G

VSV-G-encoding plasmid via calcium phosphate transfection. Briefly, DNA was diluted with sterile H_2O and

added to $CaCl_2$ (2 M) to achieve a final concentration of 0.3 M $CaCl_2$. DNA-containing samples were then

added dropwise to an equal-volume of 2X HEPES-buffered saline (280 mM NaCl, 0.5 M HEPES, 1.5 mM

Na_2HPO_4) and pipetted four times to mix. After 3–4 min, the solution was vigorously pipetted eight times

and 2 mL of transfection reagent per 10 cm dish was added dropwise to cells. The plates were gently swirled

and incubated overnight at 37°C with 5% CO_2. The medium was replaced the morning after transfection

and cells were incubated for an additional 28–30 h. At 28 h post-medium change, lentivirus was collected

from the conditioned medium. The medium was centrifuged at 500 x g for 2 min to clear cells, and the

supernatant was filtered through a 0.45 µm PES pore filter (VWR). Lentivirus was concentrated from the

filtered supernatant by ultracentrifugation in Ultra Clear tubes (Beckman Coulter 344059) at 100,420 x g at

4 °C in a Beckman Coulter Optima L-80 XP ultracentrifuge using an SW41Ti rotor. The supernatant was

aspirated, leaving virus in ~100 µl final volume (~100X concentration), and concentrated lentivirus was left

on ice for at least 30 min before resuspension, then used to transduce ~1×10^5 cells, plated either at the

time of transduction or on the day before. When appropriate, drug selection began 2 d post transduction,

using antibiotic concentrations of 1 µg ml^{-1} puromycin (Invitrogen ant-pr) on HEK293FT cells or 0.2 µg ml^{-1}

puromycin on Jurkat cells. Cells were kept in antibiotics for at least 2 weeks with subculturing every 1–2 d.

Transfection

For transfection of Freestyle 293-F cells, a transfection protocol previously used to make AAV from

suspension HEK cells was utilized.[51] Briefly, 3 d before transfection cells were plated in fresh FreeStyle 293

Expression Medium at a concentration of 1×10^6 cells per mL. On the day of transfection, Freestyle 293-F

cells were spun at 150g x 5 min and resuspended in fresh FreeStyle 293 Expression Medium. The cells

were counted and plated at a density of 1×10^6 cells per mL. Cells were transfected with 1.5 µg of DNA per

million cells by PEI transfection. Plasmid DNA was added to a volume of room temperature (~20°C)

FreeStyle 293 Expression Medium, corresponding to 5% of the total volume in the flask to be transfected.

The tube was mixed to evenly distribute plasmid DNA. 3 µg of PEI (Polysciences, Inc – 24765) per million

cells was added to the tube and vortexed. The tube was then incubated for 10 min at room temperature to

allow for precomplex formation and then added directly to the flask.

EV-AAV production, isolation, and concentration

Freestyle 293-F producer cell lines were transfected as described in Transfection. For EV-AAVs, the AAV

vector plasmid (transfer), packaging plasmid (rep/cap), and adenoviral helper plasmid (pHelper) were

transfected in equal DNA mass ratios up to the maximum DNA amount for the mixture. In transfections used

to make EV-AAV vectors including two types of EV surface protein, the mass ratio of transfected plasmids

was 1:1:1:1.59:1.41 for the AAV vector plasmid (transfer), packaging plasmid (rep/cap), adenoviral helper

plasmid (pHelper), pMD2.G VSV-G plasmid, and Ultra-HEK plasmid respectively. For control vectors in

which one such EV surface protein was omitted, pcDNA was included instead (at the equivalent DNA mass)

to ensure the same total mass of DNA was transfected into each flask. In experiments comparing various

single EV surface functionalizations, an equivalent mass of plasmid encoding each surface protein was

used, with the remaining mass made up by the AAV vector plasmid (transfer), packaging plasmid (rep/cap),

adenoviral helper plasmid (pHelper) at equivalent mass ratios. For exact DNA doses, see **Supplementary**

Data 3. 3 d after transfection, EV-AAVs were collected from the conditioned medium by differential

centrifugation. Briefly, conditioned medium was cleared of debris by centrifugation at 300 x g for 10 min to

remove cells followed by centrifugation at 2,000 x g for 20 min to remove dead cells and apoptotic bodies

in a Sorvall Legend X1R Centrifuge (ThermoFisher Scientific 75004261) with a TX-400 4 x 400 mL Swinging

Bucket Rotor. All centrifugation steps were performed at 4 °C. The supernatant underwent tangential flow

filtration using a KrossFlo KR2i TFF System with a MICROKROS 20CM 500KD filter (Repligen C02-S500-

05-S). Supernatant was flowed through the system at 24.99 mL/min at a constant transmembrane pressure

of 6.0 psi. The supernatant was concentrated to approximately 5 mL by TFF. In between samples, the TFF

was cleaned with at least 50 mL of 0.2 M NaOH that was sterile filtered by a 0.22 μm bottle top filter (Corning

431118). The system was then washed with at least 100 mL of PBS filtered by a 0.1 μm bottle top filter

(Corning 431475). The TFF system was kept in 0.05% sodium azide (EMD SX0299-1) in H_2O to prevent

contamination. After TFF, the EV-AAVs were further concentrated to approximately 0.5 mL (~800X

concentration) in Amicon Ultra-4 100K Centrifugal Filters (Millipore UFC810024) that were pre-washed with

0.1 μm bottle top filtered PBS. 0.5 mL of concentrated EV-AAVs were treated with Benzonase (25 U/mL,

EMD Millipore 70664) supplemented with $MgCl_2$ at a concentration of 2 mM for 1 h at 37 °C. Samples were

then purified to remove free AAV and other non-EV proteins using one of the methods below.

EV iodixanol purification

Iodixanol gradient purification of EVs was modified from a previous report.[21] Briefly, 1x PBS-MK buffer was

prepared by dissolving 26.3 mg of MgCl2 and 14.91 mg of KCl in 100 mL of 1x PBS. The solution was

sterilized by passing it through a 0.22 μm filter (Corning 431118). 1 M NaCl/PBS-MK buffer was prepared

by dissolving 5.84 g of NaCl, 26.3 mg of MgCl2 and 14.91 mg of KCl in 1X PBS in a final volume of 100

mL. The solution was sterilized by passing it through a 0.22 μm filter (Corning 431118). Three iodixanol

(Sigma Alrich D1556) gradients were prepared from these solutions: 25% iodixanol (5 mL of 60% iodixanol,

7 mL of 1x PBS-MK buffer), 40% iodixanol (6.7 mL of 60% iodixanol and 3.3 mL of 1x PBS-MK buffer) and

60% iodixanol (10 mL of 60% iodixanol). These solutions were layered into UltraClear tubes (Beckman

Coulter 344059) in the following order: 1 mL 60% iodixanol, 3 mL 40% iodixanol, and 5 mL 25% iodixanol.

EVs were layered on top of the 25% iodixanol layer. The tubes were spun in a Beckman Coulter Optima L-

80 XP ultracentrifuge with an SW41 Ti rotor at 190,000 x g for 3 h at 18°C. After the spin, EVs were visible

as a white band in the 25% layer and were removed with a syringe with 18G needle. To remove the iodixanol

from the EVs, EVs were placed in a new UltraClear tube and diluted in PBS. The tubes were spun at

120,000 x g for 90 min at 4 °C to pellet the EVs. After the spin, the supernatant was aspirated, and EV

pellets were left in ~100–200 μl of PBS. EVs were incubated on ice for at least 30 min after supernatant

removal before resuspension in a total of 500 μL of PBS.

EV size exclusion chromatography purification

Size exclusion chromatography was performed using qEVoriginal (35 nm) Gen2 columns (Izon ICO-35)

attached to an Automatic Fraction Collector V1 (Izon). We followed the manufacturer's instructions by

equilibrating the SEC column in PBS sterilized with 0.1 µM bottle top filters (Corning 431475). 0.5 mL of

EVs were placed on the column and EV fractions 1-4 were collected. Fractions 1-4 were combined and

concentrated to approximately 250 µL in Amicon Ultra-4 100K Centrifugal Filters (Millipore UFC810024)

that were pre-washed with 0.1 µm bottle top filtered PBS.

EV concentration determination by Nanoparticle Tracking Analysis (NTA)

EV concentration was determined by NanoSight analysis. Samples were diluted in PBS sterilized with 0.1

µM bottle top filters (Corning 431475) to concentrations on the order of 10^8 particles per ml for analysis.

NanoSight analysis was performed on an NS300 (Malvern), software version 3.4. Three 30 s videos were

acquired per sample using a 642 nm laser on a camera level of 14, an infusion rate of 30 and a detection

threshold of 7. Default settings were used for the blur, minimum track length and minimum expected particle

size. EV concentrations were defined as the mean of the concentrations calculated from each video. Size

distributions were generated by the software.

AAV crude cell lysate production

AAV was harvested from the Freestyle 293-F cell lysates of cells transfected to make non-targeted EV-

AAVs (no EV surface modifications). Briefly, 72 h post-transfection the EV-AAV containing supernatant was

removed and the cells were resuspended and washed with PBS sterilized with 0.1 µM bottle top filters

(Corning 431475). The cells were spun at 1000 x g for 7 min in a Sorvall Legend X1R Centrifuge

(ThermoFisher Scientific 75004261) with a TX-400 4 x 400 mL Swinging Bucket Rotor. The supernatant

was removed, and the cells were mixed with AAV lysis buffer corresponding to 2% of the initial culture

volume. AAV lysis buffer was made by combining 2 mL of 5 M NaCl, 0.158 g Tris-HCl and H_2O up to 100

mL. The solution was pH-adjusted to 8.5 and sterile filtered by a 0.22 µm bottle top filter (Corning 431118).

The cell pellet resuspended in AAV lysis buffer was then frozen in a bath of dry ice and ethanol, followed

by thawing in a 37°C bead bath. After each thawing the solution was vortexed vigorously for 30 s. The cell

pellet underwent three rounds of freeze/thaws. Benzonase (50 U/mL, EMD Millipore 70664) was added to

the mixture and incubated for 30 min at 37°C. The solution was transferred to 1.5 mL microcentrifuge tubes

and spun at 7000 rpm for 15 min in a microcentrifuge (Eppendorf Centrifuge 5424). The supernatant,

clarified AAV crude lysate, was transferred to a clean microcentrifuge tube and stored at 4°C until further

processing.

AAV purification by iodixanol gradient centrifugation

1x PBS-MK buffer was prepared by dissolving 26.3 mg of MgCl2 and14.91 mg of KCl in 100 mL of 1x PBS.

The solution was sterilized by passing it through a 0.22 µm filter (Corning 431118). 1 M NaCl/PBS-MK buffer

was prepared by dissolving 5.84 g of NaCl, 26.3 mg of MgCl2 and 14.91 mg of KCl in 1X PBS in a final

volume of 100 mL. The solution was sterilized by passing it through a 0.22 µm filter (Corning 431118). Four

iodixanol (Sigma Alrich D1556) gradients were prepared from these solutions: 15% iodixanol (4.5 mL of

60% iodixanol and 13.5 mL of 1 M NaCl/PBS-MK buffer), 25% iodixanol (5 mL of 60% iodixanol, 7 mL of 1x

PBS-MK buffer and 30 µL of phenol red), 40% iodixanol (6.7 mL of 60% iodixanol and 3.3 mL of 1x PBS-

MK buffer) and 60% iodixanol (10 mL of 60% iodixanol and 45 µL of phenol red). These solutions were

layered into UltraClear ultracentrifuge tubes (Beckman Coulter 344059) in the following order: 1.35 mL 60%

iodixanol, 1.35 mL 40% iodixanol, 1.6 mL 25% iodixanol, 2.7 mL 15% iodixanol, and up to 4 mL of AAV

Crude Lysate. The tubes were spun in a Beckman Coulter Optima L-80 XP ultracentrifuge with an SW41 Ti

rotor at 190,000 x g for 3 h at 18°C. AAV was removed by inserting a syringe with 18G needle at the border

between the 60% and the 40% iodixanol fractions, with the needle facing upward to the lower portion of the

40% iodixanol fraction where the viral particles accumulate. The entire 40% iodixanol layer (transparent)

was removed without disturbing the layer of protein contaminants that accumulate at the interphase

between the 40% and 25% iodixanol layers. To remove the iodixanol this layer was mixed 1:1 by volume

with PBS and added to an Amicon Ultra-15 100K spin filter. The spin filter was spun at 3000 x g until the

volume remaining in the column was less than 1 mL. PBS was added to the column to the full volume and

spun at 3000 x g until 1 mL remained in the column. This was PBS washing was repeated 4 more times to

remove iodixanol and concentrate the AAV to its final volume.

AAV purification by magnetic beads

Free AAV was purified using Dynabeads Capture Select AAVX Magnetic Beads (Thermofisher Scientific

2853522001) by the manufacturer's protocol. Briefly, 40 µL of bead slurry was added to a low protein binding

microcentrifuge tube with 460 µL of PBS. The tubes were placed on a magnetic stand and once the beads

were collected on the side of the tube the supernatant was removed. The beads were then mixed with 500

µL of PBS, the tubes were placed on a magnetic stand, and the supernatant was removed. 1 mL of AAV

crude lysate was added to the beads and the tubes were put on a rotating platform to continuously mix the

beads and lysate for 30 min. After 30 min the beads were collected with the magnetic stand and the

supernatant was removed. 500 µL of PBS was mixed with the beads, the beads were collected on the

magnetic stand, and the supernatant was removed. This PBS wash was repeated again. 20–50 µL of elution

buffer was added to the beads and mixed well. The elution buffer (50 mM citric acid, pH 2.5–3.0) and beads

were incubated at room temperature for 10 min with occasional pipetting. The beads were collected on the

magnetic stand and the supernatant containing purified AAV was collected. The eluted AAV mixture was

neutralized by adding 1 µL of Neutralization Buffer (1 M Tris-HCl, pH 8.7) for each 10 µL of eluate.

Physical titer determination by qPCR

First, 5 µL of the EV-AAV, purified AAV, or clarified crude lysate samples were mixed with 37 µL nuclease-

free H_2O, 5 µL of 10x DNaseI buffer (ThermoFisher B43) and 1 µL of DNaseI (ThermoFisher EN0521) to

eliminate plasmid DNA, incubating at 37°C for 1 h followed by a 15 min inactivation step at 75°C. Samples

were placed on ice. Then, viral genomic DNA (vgDNA) was extracted using the High Pure Viral Nucleic Acid

Kit (Roche 11858874001), following manufacturer instructions. qPCR analysis was performed by amplifying

a sequence within the LumiScarlet reporter using primers CAGGGAGGTTTGACCAGTTT (forward), and

GGATATCGATCTTCAGCCCATT (reverse) or the mCherry reporter using the primers

AGGACGGCGAGTTCATCTA (forward), and CCCATGGTCTTCTTCTGCATTA (reverse). Primers were

designed using IDT's PrimerQuest Tool. A plasmid containing the LumiScarlet transgene (pJB035) or

mCherry transgene (pJB040) were used as a standard; serial dilutions were prepared using EASY dilution

(Takara Bio 9160), plating each dilution in duplicate. The vgDNA samples were diluted in nuclease free

H_2O, plating 4 serial dilutions per sample (in duplicate). For each qPCR reaction (each in a 96-well), 5 µL

of vgDNA sample were mixed with 10 µL Universal SYBR Master Mix 2x (ThermoFisher K0221), 0.15 µL

of 100 µM forward primer, 0.15 µL of 100 µM reverse primer, and 4.7 µL of nuclease free H_2O. qPCR plates

were sealed and spun at 500 x g for 2 min before placing in Bio-Rad C1000 Thermal Cycler with CFX96

Real-Time System. Samples run to detect LumiScarlet were heated to 98°C for 3 min, followed by 40 cycles

of: 98°C for 15 s, 62°C for 30 s. Then, a final step of 0.5°C increments every 5 s from 65°C to 95°C was

performed. Samples run to detect mCherry were heated to 98°C for 3 min, followed by 40 cycles of: 98°C

for 15 s, 55°C for 30 s. Then, a final step of 0.5°C increments every 5 s from 65°C to 95°C was performed.

Transduction assays

For most experiments, EV-AAV or free AAV vectors were diluted in 0.1 µm bottle top filtered PBS so that

100 µL of the dilution contained 10^9 vector genomes. 100 µL of the diluted vector was added to each well

of a 24-well plate. Recipient cells were diluted to 10^4 cells per 900 µL of media and added to the wells with

vector, for a final dosage of 10^5 vector genomes per cell. Negative control wells received 100uL of 0.1 µm

bottle top filtered PBS. After the addition of all vectors and recipient cells, the plate was vigorously mixed

and placed in an incubator. 5 d post transduction, cells were harvested for flow cytometry using 0.05%

Trypsin-EDTA with phenol red for HEK293FT recipient cells for 5 min followed by quenching with medium.

The resulting cell solution was added to at least 2 volumes of FACS buffer (PBS pH 7.4, 2–5 mM EDTA,

0.1% BSA) supplemented with DAPI (to identify live cells) at a final concentration of 3 µM. Jurkat cells were

pipetted from the plate directly into FACS tubes containing FACS buffer supplemented with DAPI. A single

FACs tube of PBS treated cells was added to a tube containing FACS buffer without DAPI to identify live

cells (DAPI negative). Cells were centrifuged (150 × g for 5 min), supernatant was decanted, fresh FACS

buffer, with or without DAPI, was added, and cells were resuspended by briefly vortexing.

For experiments involving lentiviral transduction of HEK293T and HEK293FT by lentiviral vectors including

Ultra-HEK, similar methods were used. Cell lines were harvested, counted, and plated in 96-well, TC

treated, PBS-moated, flat-bottom plates at a density of 10^5 cells/mL in a volume of 100 µL per well (10^4

cells/well). Crude LV supernatant was diluted 5-fold seven times in complete DMEM to generate a seven-

point dilution curve starting from the first dilution. Cells were transduced by adding 100 µL/well of LV (each

dilution) and were incubated for 3 d at 37°C. Before analysis of transduction by flow cytometry, cells were

harvested by aspirating the media, trypsinizing cells for 5 min at room temperature, quenching with

complete DMEM, transferring lifted cells to a non-treated, round-bottom 96-well plate, spinning cells at 150

x g for 5 min, and resuspending in 30 µL of FACS buffer (PBS pH 7.4 with 0.05% BSA and 2 mM EDTA) +

DRAQ7 viability dye (Thermo Fisher D15106, diluted 1000x). DRAQ7 was used to identify live cells (DRAQ7

negative) for analysis.

Microscopy imaging

Microscopy was conducted on a Keyence BZ-X800E fluorescence microscope and a Nikon CFI S Plan

Fluor ELWD 20XC objective (MRH08230). LumiScarlet fluorescence was imaged with a Keyence BZ-X

Filter TexasRed (Keyence, OP-87765). Images were taken using the Keyence BZ Series Viewer Software

(Version 01.03.00.01) in high resolution mode. Scale bars were added to images with the BZ-X800 Analyzer

Software (Version 1.1.2.4).

Lentiviral vector production for Ultra-HEK analysis

Lentivirus (LV) samples were produced in adherent Lenti-X cells seeded at a density of 3.2×10^5 cells/mL

in a volume of 0.5 mL DMEM in 24-well TC-treated plates 24 h prior to transfection. Cells were transfected

with 0.4 µg total DNA comprising 50% transfer vector, 29% gag/pol, 16% rev, and 5% VSV-G for the VSV-

G only composition and 46% transfer vector, 29% gag/pol, 16% rev, 5% VSV-G, and 4% Ultra-HEK for the

VSV-G and Ultra-HEK composition. The transfer vector used encodes constitutive expression of

mNeonGreen. Plasmids used for lentivirus production are Syenex's commercial plasmids: SNX-mNG-TV3,

SNX-Gagpol, SNX-Rev, SNX-VSV-G, and Ultra-HEK. Ultra-HEK is a plasmid designed to increase lentiviral

transduction of HEK293-derived cells. Plasmids and PEI MAX transfection reagent (Polysciences 24765)

were diluted in Opti-MEM and mixed in equal volumes at a 4:1 mass ratio of PEI to DNA and complexed

for 24 min prior to transfection. Medium was changed 12–16 h later. At least 28 h post media change, LV

was harvested from the conditioned medium and centrifuged at 1000 x g for 10 min at 4°C, then aliquoted

and stored at -80°C until use.

Analytical flow cytometry and analysis

Most flow cytometry was performed on a BD LSR Fortessa Special Order Research Product using the

552 nm laser for mCherry or LumiScarlet (610/20 filter) and the 488 nm laser for eGFP or mNeonGreen

(530/30 filter). At least 10,000 live cells were collected per sample for analysis. Data were analyzed using

FlowJo v10 (FlowJo, LLC). Briefly, cells were identified using a forward scatter area (FSC-A) versus side

scatter area (SSC-A) plot and gated for single cells using an FSC-A versus forward scatter height (FSC-H)

plot (**Supplementary Information (Figures S1,S2)**). Live single cells were identified using a DAPI-A versus

SSC-A plot. Gene delivery was determined by the percentage of live single cells that were brighter than

approximately the 0.1% of the brightest cells in the untransduced (PBS treated) cell samples (**Figure S1**).

Protein delivery was quantified by the mean of PE-Texas Red fluorescence of live single cells. This mean

fluorescence intensity was converted to MEPTRs by using UltraRainbow Calibration Particles (Spherotech).

Briefly, to determine the conversion from the arbitrary unit (au) of fluorescence recorded by the flow

cytometer to a standardized fluorescence unit (Molecules of Equivalent PE-TexasRed (MEPTR)),

UltraRainbow Calibration Particles were run as part of each flow cytometry experiment, except the AAV-

LumiScarlet experiment (**Figure 1**). This reagent contains nine subpopulations of beads, each of a specific

size and with a known intensity of various fluorophores, which are provided with calibration data for each

lot. The total bead population was identified by SSC-A vs. FSC-A gating, and subpopulations were identified

through two fluorescent channels (**Figure S3**). A calibration curve was generated for the experimentally

determined au vs. manufacturer supplied MEPTR, and a linear regression was performed with the

constraint that 0 au equals 0 MEPTR. The slope from the regression was used as the conversion factor.

For experiments involving lentiviral transduction of HEK293T and HEK293FT by Ultra-HEK, similar methods

were used. Flow cytometry was performed on a Novocyte Penteon using the 488 nm laser for mNeonGreen

(525/45 filter) and the 637 nm laser for DRAQ7 (725/40 filter). Approximately 3,000 live cells were collected

per sample for analysis. Data were analyzed using FlowJo v10 (FlowJo, LLC). Briefly, cells were identified

using an FSC-A vs SSC-A plot and gated for singlets using an FSC-A vs FSC-H plot. Live cells were

identified using a DRAQ7 negative gate drawn on a non-DRAQ7-stained sample, and transduced cells

were identified by drawing an mNeonGreen positive gate on a non-transduced sample.

Transmission Electron Microscopy (TEM)

For transmission electron microscopy (TEM), samples were diluted in PBS sterilized with 0.1 µm bottle top

filters (Corning 431475) to concentrations on the order of 10^8 particles per µL as determined by Nanoparticle

Tracking Analysis. Grids were plasma treated, and 3 µL of diluted sample were placed on the carbon side

of the grid for 30 s. The grids were then briefly washed on 3 droplets of distilled water and placed on a

droplet of stain solution (2% ammonium molybdate) for a few seconds. The staining solution was blotted

off and the grid was placed on a second droplet of staining solution for 30 s. Finally, the excess staining

solution was removed, and the grid was air-dried. Samples were observed with a JEOL 3200FS TEM at an

acceleration voltage of 300 kV. A K2 camera was used for recording images.

Western blots

For western blots comparing protein in cell lysates and protein in vesicles, a fixed number of vesicles and

a fixed mass of cell lysate were loaded into each well: 4.5×10^8 EVs/lane and 2 µg protein/lane,

respectively. An established western blot protocol[48] was followed with the following modifications. In most

cases, the following reducing Laemmli composition was used to boil samples (60 mM Tris-HCl pH 6.8, 10%

glycerol, 2% sodium dodecylsulphate, 100 mM dithiothreitol (DTT) and 0.01% bromophenol blue); in some

cases, a nonreducing Laemmli composition (without DTT) was used, as previously reported[52]. After transfer,

membranes were blocked whilst rocking for 1 h at room temperature in 5% milk in Tris-buffered saline with

Tween (TBST) (pH: 7.6, 50 mm Tris, 150 mm NaCl, HCl to pH 7.6, 0.1% Tween 20). Primary antibody was

added in 5% milk in TBST, rocking, for 1 h at room temperature and then washed three times with TBST

for 5 min each. Secondary antibody in 5% milk in TBST was added at room temperature for 1 h or overnight

at 4°C. Membranes were then washed three times with TBST for 5 min each. The membrane was incubated

with Clarity Western ECL substrate (Bio-Rad, #1705061) and imaged on an Azure c280 running Azure

cSeries Acquisition software v1.9.5.0606. Specific antibodies, antibody dilution, heating temperature,

heating time and Laemmli composition for each antibody was used as previously reported[52].

Statistical analysis

Statistical tests employed are described in relevant figure legends. Unless otherwise stated, three

independent technical replicates were analyzed per condition, and the mean fluorescence intensity of at

least 10,000 live single cells were analyzed per sample. Each figure represents a single experiment that

was repeated (biological replicate) at least twice. Unless otherwise indicated, error bars represent the

standard error of the mean. GraphPad Prism 10 was used to analyze the data. Multicomparison statistical

analysis was performed using a one-way ANOVA test, followed by Tukey's multiple comparison (i.e.,

honestly significant difference, HSD) test to evaluate specific comparisons. Significance threshold:

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

RESULTS

Distinguishing protein delivery from gene delivery by EV-AAV is challenging

We first sought to quantify the magnitude and dynamics of pseudotransduction for EV-AAV delivery

to HEK293FTs. It is commonly expected that the fluorescence conferred to recipient cells by EV-mediated

delivery of producer cell-derived protein (i.e., pseudotransduction) will decrease over time due to protein

degradation and/or cell division, while fluorescence originating from true transduction (gene delivery and

resulting transgene expression in recipient cells) will persist and increase to steady state. However, the

dynamics of such contributions are poorly explored and are likely to be system specific. To characterize

these phenomena in our model system, we utilized an AAV vector that constitutively expresses a bright

fluorescent protein, LumiScarlet, and we generated both EV-AAV and free AAV preparations (**Figure S4**).

As a control for protein delivery, we included conditions lacking Rep and Cap genes ("No Rep/Cap"). We

also investigated several EV purification methods, reasoning that some fluorescent protein made by

producer cells may be excreted into the media along with free AAVs and EVs, and different EV purification

methods may remove these free proteins to different extents. Both purification methods yielded EVs with

expected protein markers (**Figures S5A,B**), appropriate size distributions (**Figure S5C**), and morphology

by TEM (**Figure S5D**). Purified vectors were dose-normalized (by genome count for EV-AAV and AAV with

No Rep/Cap EVs matched to EV-AAV by particle count) and delivered to HEK293FT cells (which actively

endocytose EVs). Fluorescence was monitored over 7 days by flow cytometry (**Figure 1**) and microscopy

(**Figure S6**). In general, fluorescence attributable to EV-AAV transduction was not readily distinguished

from fluorescence attributable to protein delivery (e.g., No Rep/Cap control). Free AAV2 conferred

distinguishable transgene expression, which was distinguished by quantifying the magnitude of

fluorescence per cell (**Figure 1A, right**). Representative histograms of LumiScarlet fluorescence from each

sample also demonstrate that fluorescence attributable to EV-AAV transduction was not readily

distinguished from fluorescence attributable to protein delivery (**Figure 1B**). This was visually confirmed by

microscopy (**Figure S6A-C**), in which free AAV2 conferred the expected whole cell fluorescence, while EV

conditions exhibit puncta that are readily apparent at 3 days post-treatment and remain detectable at 7

days post-treatment. Our analyses indicate that delivered protein is detectable in recipient cells for up to a

week, such that the time scale for delivered protein depletion and degradation is not readably separated

from the timescale of AAV episomal vector dilution (which may be particularly pronounced in these rapidly

dividing recipient cells). Overall, protein delivery by EV-AAVs may lead to significant pseudotransduction

and is challenging to distinguish from gene delivery regardless of EV purification method.

A Cre recombinase-based dual reporter system enables discrimination between protein and gene

delivery

To address the challenge of pseudotransduction, we developed a novel Cre recombinase (Cre)-

based system to enable quantitative and unambiguous evaluation of gene and protein delivery by EV-AAVs.

We utilized a Cre-responsive AAV genome encoding a constitutive mCherry fluorescent protein that, upon

recombination, is excised to instead drive expression of a green fluorescent protein.[47] This dual-

fluorescence system enables visualization of producer cell-derived protein delivery (mCherry) and gene

delivery (green) in Cre-expressing recipient cells (**Figure 2A**). Given a preceding Cre-responsive construct

developed for a distinct purpose[47], we generated several vectors to identify design choices that are useful

for this application (**Figure 2B**). We explored changing the gene delivery reporter from eGFP to

mNeonGreen, hypothesizing that a brighter fluorophore would help identify gene delivery. We also

hypothesized that removing the nuclear localization sequences (NLS) from the gene delivery reporter

protein may increase accumulation in the recipient cell, again facilitating detection. To evaluate these

designs, we made Cre-expressing recipient HEK293FT cells by lentiviral transduction.

Using this system, we evaluated protein and gene delivery by each reporter vector. Each variant

was packaged as either AAV2 crude lysate—which should include a mix of free AAV2 and cell-derived

fluorescent protein—or a no Rep/Cap control which should include only cell-derived fluorescent protein;

each was delivered to Cre-expressing recipient cells. Removal of the 3 x NLS generally increased mCherry

protein delivery, as expected, since the NLS sequesters some protein in the nucleus of the producer cells

(**Figure 2C**). Greater mCherry fluorescence was observed in Cre-negative than Cre-positive recipient cells,

suggesting that in the absence of Cre, more mCherry gene expression occurs in recipient cells. Most

importantly, green fluorescent protein was detected only with AAV2 preps (i.e., transduction-competent

vectors) and in Cre-expressing recipient cells (**Figure 2D**), validating this reporter system for

unambiguously indicating transduction and de novo transgene expression in recipient cells. Overall, the

mNeonGreen vectors conferred higher percentages of green positive live cells than did eGFP vectors, with

the mNeonGreen vector without an NLS performing the best. On balance, going forward we selected the

vector with mNeonGreen and the 3 x NLS, since it conferred more green-positivity in Cre-expressing

recipient cells compared to eGFP vectors (i.e., greater signal), and it conferred less mCherry fluorescence

compared to vectors without an NLS (i.e., lower background). We carried this validated reporter system

forward to probe EV-AAV bioengineering questions.

EV-AAV2 surface functionalization substantially increases protein delivery but not gene delivery to

HEK293FTs

We next investigated how EV functionalization strategies that have been reported to increase

delivery of lentiviruses or EVs impact protein and gene delivery by EV-AAV. VSV-G is often used in

pseudotyping lentiviruses[53,54] and EVs[55,56], and it increases the protein delivered by EVs.[56] VSV-G binds to

the LDL receptor (LDL-R) on the surface of recipient cells[31], increasing particle uptake as a result. Within

the low pH of the endosome, VSV-G undergoes conformational changes that enable it to promote fusion

between the viral (or EV) membrane and the endosomal membrane to release the vector's luminal contents

into the cytosol of the recipient cell.[57] Interestingly, a prior report indicated that pseudotyping EV-AAV2 with

VSV-G decreased the transduction of HEK293 cells compared to non-targeted EV-AAV2.[30] However, this

analysis employed a luminescence-based plate reader assay that may have been confounded by

pseudotransduction artifacts of protein delivery by the EV-AAVs. To separate the impacts of enhancing

uptake and mediating fusion, we also utilized Ultra-HEK, a commercial product (Syenex) that increases

lentiviral transduction of HEK293-based cells when combined with a fusogen such as VSV-G. Ultra-HEK is

described as interacting with proteins on the surface of recipient HEK293 cells to cause increased uptake

of the particle (but not fusion). In our hands, packing lentiviral vectors with Ultra-HEK led to increased gene

delivery (i.e., functional titer) for delivery to both HEK293T cells and HEK293FT cells (**Figure 3A–C**).

Using our reporter system and this menu of surface functionalization strategies (**Figure 3D**), we

revisited the question of how driving EV-AAV uptake (and/or fusion) impacts transduction efficiency. All

functionalization strategies increased protein delivery, likely due to increased particle uptake by recipient

cells (**Figure 3E**). Ultra-HEK combined with VSV-G functionalization increased protein delivery to the

greatest extent compared to non-targeted EV-AAV2. Since doses of EV-AAVs were normalized by genome

copy, it is possible that equivalent genome doses comprise distinct numbers of EVs. To exclude the

possibility that this could explain the protein delivery trend, we plotted protein delivery by EV dose (in particle

count), and we found that this does not well explain our observations (**Figure S7**). Thus, differences in

protein delivery are best explained by the biological effects of our functionalization strategies. Interestingly,

surface functionalization did not increase gene delivery to the same degree as it did protein delivery (**Figure**

3F). Only the dual functionalization strategy modestly increased the amount of gene delivery to HEKs over

non-targeted vectors, and there was no evidence that fusion mediated by VSV-G enhanced or blocked EV-

AAV gene delivery. This led us to investigate whether these trends extend to other serotypes and recipient

cells.

EV-AAV6 surface functionalization increases protein and gene delivery to Jurkat T cells

We next explored EV-AAV delivery to a cell type with low spontaneous EV uptake. We chose to

use human Jurkat T cells, which are somewhat recalcitrant to EV-based delivery due (at least in part) to

low endocytosis.[28] We transduced Jurkats with the Cre recombinase gene to make a recipient line

compatible with our dual reporter system. We used AAV6 as it infects both primary T cells[59] and Jurkats[60].

We focused on VSV-G functionalization which mediates lentiviral transduction via LDL-R binding and

uptake (**Figure 4A**).[61] Similar to HEK293FT delivery (**Figure 3**), VSV-G functionalization increased protein

delivery to Jurkats compared to non-targeted EV-AAV6 (**Figure 4B**). Surprisingly, and unlike in HEK293FT

recipient cells, VSV-G functionalization also increased gene delivery to Jurkats (**Figure 4C**). This led us to

investigate what mechanism conferred by VSV-G (uptake vs. fusion) contributed to increased gene delivery.

We employed two VSV-G mutants to dissect these mechanisms (**Figure 4D**). P127D retains the ability to

bind to LDL-R but is deficient in membrane fusion.[49,62,63] Conversely, mutations K47Q-R354A block LDL-R

binding but still enable VSV-G to confer membrane fusion.[64-66] Wild-type (WT) and P127D VSV-G vectors

delivered more protein than did K47Q-R354A or non-targeted vectors (**Figure 4E**). Interestingly, the K47Q-

R354A VSV-G conferred less gene delivery than did the WT or P127D VSV-G vectors, indicating that

binding to LDL-R on recipient cells is the dominant mechanism by which VSV-G increases gene delivery to

Jurkat T cells (**Figure 4F**). Notably, P127D VSV-G, which is fusion incompetent, conferred more gene

delivery than did VSV-G WT vectors. Altogether, these results may indicate that for EV-AAV6 delivery to

Jurkats, increased LDL-R binding increases particle uptake and transduction, while membrane fusion limits

transduction.

Recipient cell type but not capsid serotype drives the impact of functionalization on EV-AAV delivery

To reconcile the distinct patterns observed in our HEK293FT and Jurkat delivery experiments, we

investigated whether differences are driven primarily by recipient cell type or AAV serotype. We chose to

investigate transducing HEK293FT cells with AAV6 since AAV6 is much less efficient at infecting

HEK293FTs compared to AAV2.[60] This difference may be due to differences in receptor density on the

surface of HEKs. AAV2 primarily uses heparan sulfate proteoglycan to mediate binding and uptake by

recipient cells.[67] In contrast, AAV6 primarily uses sialic acid as its receptor[67], which may be less abundant

or accessible on HEK293 cells compared to heparan sulfate proteoglycan. Therefore, we hypothesized that

increasing uptake of EV-AAV6 (e.g., via VSV-G functionalization) would increase transduction compared to

both non-targeted EV-AAV6 or free AAV6 (**Figure 5A**). VSV-G functionalization increased protein delivery

compared to non-targeted EV-AAV6, as expected (**Figure 5B**). However, VSV-G mediated uptake (and

fusion) did not increase gene delivery by EV-AAV6 (**Figure 5C**), as was seen for EV-AAV2 (**Figure 3F**).

Thus, increasing uptake and membrane fusion in a cell type that is permissible to EV delivery, such as

HEK293FTs, did not increase gene delivery regardless of AAV capsid choice. With the set of conditions

evaluated, EV surface functionalization strategies conferred cell type-dependent but not capsid-dependent

effects on EV-AAV gene delivery.

DISCUSSION

Engineered EVs are promising delivery vehicles for AAV vectors, enabling tissue-specific targeting and

neutralizing antibody evasion. However, analyzing and bioengineering such vectors is made challenging by

the difficulty associated with distinguishing protein delivery (pseudotransduction) from gene delivery. To

address this need, in this study we developed and validated a new assay that unambiguously distinguishes

gene delivery from protein delivery, while enabling quantification of both phenomena. This assay helped to

generate new insights that inform both interpretation of past studies and identify opportunities for future

investigation.

Our observations provide new insights that help to interpret apparent contradictions in the EV-AAV

literature. Some previous studies report that EV-AAVs are more potent than AAV vectors for *in vivo* delivery

(on a per genome basis)[14-16,22], but other work focused on delivery to neuronal and primary astrocyte cell

cultures seemingly contradicts this finding[68]. It is possible that improved *in vivo* efficacy of EV-AAV particles

is primarily attributable not to the enhanced interaction between the AAV and the target cells, but rather to

shielding of AAVs from neutralizing antibodies and/or biodistribution effects such as increased circulation

time or improved delivery through tissue barriers. Our observations add another layer of possible

explanation, given our finding that target cell type and capsid choice may impact the efficacy of EV-AAV

compared to free AAV. Free AAV2 was significantly more potent for gene delivery to HEK293FTs compared

to non-targeted EV-AAV2 (**Figure 3**). However, non-targeted EV-AAV6 outperformed free AAV6 in gene

delivery to HEK293FTs (**Figure 5**). This suggests that capsid choice may impact the relative gene delivery

efficacy of EV-AAV and free AAV. In contrast, EV-AAV6 outperformed free AAV6 in transduction of Jurkat T

cells (**Figure 4**), suggesting that the target cell may also impact efficacy comparisons between EV-AAV and

free AAV. Overall, our findings indicate that the impact of EV-encapsulation of AAV confers case-specific

effects that should be evaluated for each application, and which may drive application-specific

bioengineering.

This study focused on identifying high level determinants of functional delivery, yet it also identified

opportunities for further subsequent mechanistic investigations. Increasing the uptake of EV-AAV to

HEK293FT recipient cells, both with or without promoting membrane fusion, increased protein delivery, but

this did not substantially increase gene delivery for either AAV2 or AAV6. However, VSV-G functionalization

of EV-AAV6 increased both protein and gene delivery to Jurkat T cells, and this effect was driven primarily

by increased uptake rather than fusion; indeed, blocking VSV-G-mediated fusion enhanced gene delivery.

The basis for these differences is not known. A possible explanation is that AAV requires processing in the

endosome—including capsid cleavage by endosomal cathepsins[32] and acidification-mediated

autoproteolytic processing[33] to expose a phospholipase domain promoting virion escape and gene

delivery[34,35]. This requirement is well-characterized for free AAV2 and may extend to other serotypes. It

would be informative to probe how each step in capsid processing is impacted by encapsulation in EVs. In

addition, directly tracking internalization of both receptors and particles could help to test the mechanisms

offered as speculative explanations here. It will be interesting to extend these studies to the many AAV

capsid variants that exist, including natural serotypes as well as novel forms developed through directed

evolution and protein engineering[69], and to a variety of recipient cell types including those which endocytose

EVs more readily than do HEK293FTs[58]. This work could also enable new AAV bioengineering campaigns

specific to enhancing EV-AAV delivery, such as exploring mutations that destabilize the capsid structure to

enhance transduction after EV fusion and delivery of the AAV core to the cytosol, bypassing endosomal

processing.

An exciting prospect is extending this assay for *in vivo* EV-AAV delivery analyses. The stoplight vector

inspiring our approach was originally employed to detect Cre expression *in vivo*[47]. Given that transgenic

animals can be engineered to express Cre in a tissue-restricted or general fashion[70], our assay could be

employed to unambiguously identify EV-AAV gene delivery in vivo. Such a capability could be particularly

useful for directly comparing AAV and EV-AAV-mediated delivery, and for enabling comparative evaluation

of bioengineering approaches, *in vivo* multiplexed screening, and other efforts for which pseudotransduction

is misleading or reduces signal-to-noise to preclude feasibility. We expect that such work can help advance

the understanding and use of EV-AAV vectors for diverse gene delivery applications.

AUTHOR CONTRIBUTIONS

J.D.B., D.M.S., N.P.K., D.T.E., and J.N.L. conceived the initial project. J.D.B. performed the experiments. D.M.S. and H.I.E. performed the Ultra-HEK lentiviral vector experiment. J.D.B. and J.N.L. planned and analyzed experiments. J.D.B and J.N.L. wrote the manuscript. J.N.L. supervised the work.

ACKNOWLEDGEMENTS

J.D.B. was supported by the National Institutes of Health Training Grant (T32GM008449) through Northwestern University's Biotechnology Training Program. J.D.B was supported by the National Science Foundation Graduate Research Fellowship under grant number (DGE-1842165). This project was partially supported by an award from the McCormick Catalyst Fund (J.N.L.) and by a Cornew Innovation Award from the Chemistry of Life Processes Institute at Northwestern University (J.N.L.). Flow Cytometry was performed in the Northwestern University Flow Cytometry Core Facility supported by Cancer Center Support Grant (NCI CA060553) and in the Single Cell Genomics Facility at Northwestern University (RRID:SCR_026652), graciously supported by the Department of Neurobiology, the Department of Molecular Biosciences, and the NU Office for Research. This work made use of the BioCryo Core (RRID:SCR_021288) of Northwestern University's NUANCE Center, which has received support from the SHyNE Resource (NSF ECCS-2025633), the IIN, and Northwestern's MRSEC program (NSF DMR-2308691). Biological analysis] was performed in the Analytical bioNanoTechnology Equipment Core Facility of the Center for Regenerative Nanomedicine at Northwestern University. ANTEC receives partial support from the Soft and Hybrid Nanotechnology Experimental (SHyNE) Resource (NSF ECCS-2025633) and Feinberg School of Medicine, Northwestern University. This work was supported by the Northwestern University Sanger Sequencing Facility. Any opinion, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. The authors kindly thank the Leonard lab members for useful discussions throughout the planning, experimental, analysis and writing phases of this project. The authors would like to thank Dmitriy Bobrovnikov and Roxi Mitrut for discussions of Cre-responsive AAV vectors.

DECLARATION OF INTERESTS STATEMENT

REFERENCES

Wang, J.-H., Gessler, D. J., Zhan, W., Gallagher, T. L. & Gao, G. Adeno-associated virus as a delivery vector for gene therapy of human diseases. *Signal Transduction and Targeted Therapy* **9**, 78 (2024). https://doi.org:10.1038/s41392-024-01780-w

Earley, J., Piletska, E., Ronzitti, G. & Piletsky, S. Evading and overcoming AAV neutralization in gene therapy. *Trends in Biotechnology* **41**, 836-845 (2023). https://doi.org:https://doi.org/10.1016/j.tibtech.2022.11.006

Srivastava, A. In vivo tissue-tropism of adeno-associated viral vectors. *Current Opinion in Virology* **21**, 75-80 (2016). https://doi.org:https://doi.org/10.1016/j.coviro.2016.08.003

Duan, D. Lethal immunotoxicity in high-dose systemic AAV therapy. *Molecular Therapy* **31**, 3123-3126 (2023). https://doi.org:https://doi.org/10.1016/j.ymthe.2023.10.015

Hordeaux, J. *et al.* High-dose systemic adeno-associated virus vector administration causes liver and sinusoidal endothelial cell injury. *Molecular Therapy* **32**, 952-968 (2024). https://doi.org:https://doi.org/10.1016/j.ymthe.2024.02.002

Raposo, G. & Stoorvogel, W. Extracellular vesicles: exosomes, microvesicles, and friends. *J Cell Biol* **200**, 373-383 (2013). https://doi.org:10.1083/jcb.201211138

Cocucci, E., Racchetti, G. & Meldolesi, J. Shedding microvesicles: artefacts no more. *Trends Cell Biol* **19**, 43-51 (2009). https://doi.org:10.1016/j.tcb.2008.11.003

Valadi, H. *et al.* Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* **9**, 654-659 (2007). https://doi.org:10.1038/ncb1596

Mizrak, A. *et al.* Genetically engineered microvesicles carrying suicide mRNA/protein inhibit schwannoma tumor growth. *Mol Ther* **21**, 101-108 (2013). https://doi.org:10.1038/mt.2012.161

Andaloussi, E. L., Mager, I., Breakefield, X. O. & Wood, M. J. Extracellular vesicles: biology and emerging therapeutic opportunities. *Nature reviews. Drug discovery* **12**, 347-357 (2013). https://doi.org:10.1038/nrd3978

Gyorgy, B., Hung, M. E., Breakefield, X. O. & Leonard, J. N. Therapeutic applications of extracellular vesicles: clinical promise and open questions. *Annual review of pharmacology and toxicology* **55**, 439-464 (2015). https://doi.org:10.1146/annurev-pharmtox-010814-124630

12 Marcus, M. E. & Leonard, J. N. FedExosomes: Engineering Therapeutic Biological Nanoparticles

that Truly Deliver. *Pharmaceuticals-Base* **6**, 659-680 (2013). https://doi.org:10.3390/ph6050659

13 Maguire, C. A. *et al.* Microvesicle-associated AAV vector as a novel gene delivery system. *Mol Ther*

20, 960-971 (2012). https://doi.org:10.1038/mt.2011.303

14 Hudry, E. *et al.* Exosome-associated AAV vector as a robust and convenient neuroscience tool.

Gene Therapy **23**, 380-392 (2016). https://doi.org:10.1038/gt.2016.11

15 György, B. *et al.* Rescue of Hearing by Gene Delivery to Inner-Ear Hair Cells Using Exosome-

Associated AAV. *Molecular Therapy* **25**, 379-391 (2017).

https://doi.org:https://doi.org/10.1016/j.ymthe.2016.12.010

16 Wassmer, S. J., Carvalho, L. S., György, B., Vandenberghe, L. H. & Maguire, C. A. Exosome-

associated AAV2 vector mediates robust gene delivery into the murine retina upon intravitreal

injection. *Scientific Reports* **7**, 45329 (2017). https://doi.org:10.1038/srep45329

17 Meliani, A. *et al.* Enhanced liver gene transfer and evasion of preexisting humoral immunity with

exosome-enveloped AAV vectors. *Blood Advances* **1**, 2019-2031 (2017).

https://doi.org:10.1182/bloodadvances.2017010181

18 Breuer, C. B. *et al.* In vivo engineering of lymphocytes after systemic exosome-associated AAV

delivery. *Scientific Reports* **10**, 4544 (2020). https://doi.org:10.1038/s41598-020-61518-w

19 Liu, B. *et al.* AAV-Containing Exosomes as a Novel Vector for Improved Gene Delivery to Lung

Cancer Cells. *Frontiers in Cell and Developmental Biology* **Volume 9 - 2021** (2021).

https://doi.org:10.3389/fcell.2021.707607

20 Kwak, G. *et al.* Extracellular vesicles enhance pulmonary transduction of stably associated adeno-

associated virus following intratracheal administration. *Journal of Extracellular Vesicles* **12**, 12324

(2023). https://doi.org:https://doi.org/10.1002/jev2.12324

21 Li, X. *et al.* Extracellular Vesicle–Encapsulated Adeno-Associated Viruses for Therapeutic Gene

Delivery to the Heart. *Circulation* **148**, 405-425 (2023).

https://doi.org:10.1161/CIRCULATIONAHA.122.063759

22 György, B., Fitzpatrick, Z., Crommentuijn, M. H. W., Mu, D. & Maguire, C. A. Naturally enveloped

AAV vectors for shielding neutralizing antibodies and robust gene delivery in vivo. *Biomaterials* **35**,

7598-7609 (2014). https://doi.org:https://doi.org/10.1016/j.biomaterials.2014.05.032

23 Cheng, M. *et al.* Neutralizing Antibody Evasion and Transduction with Purified Extracellular Vesicle-

Enveloped Adeno-Associated Virus Vectors. *Human Gene Therapy* **32**, 1457-1470 (2021).

https://doi.org:10.1089/hum.2021.122

24 Cheng, M. *et al.* Probing aspects of extracellular vesicle associated AAV allows increased vector

yield and insight into its transduction and immune-evasive properties. *Mol Ther Methods Clin Dev*

33, 101407 (2025). https://doi.org:10.1016/j.omtm.2025.101407

25 Espinoza, P. *et al.* Metabolic engineering improves transduction efficiency and downstream vector

isolation by altering the lipid composition of extracellular vesicle-enclosed AAV. *Metabolic*

Engineering **88**, 40-49 (2025). https://doi.org:https://doi.org/10.1016/j.ymben.2024.12.003

26 Alvarez-Erviti, L. *et al.* Delivery of siRNA to the mouse brain by systemic injection of targeted

exosomes. *Nat Biotechnol* **29**, 341-345 (2011). https://doi.org:10.1038/nbt.1807

27 Murphy, D. E. *et al.* Extracellular vesicle-based therapeutics: natural versus engineered targeting

and trafficking. *Experimental & molecular medicine* **51**, 1-12 (2019).

28 Stranford, D. M. *et al.* Genetically encoding multiple functionalities into extracellular vesicles for the

targeted delivery of biologics to T cells. *Nature Biomedical Engineering* **8**, 397-414 (2024).

https://doi.org:10.1038/s41551-023-01142-x

29 Wiklander, O. P. B. *et al.* Antibody-displaying extracellular vesicles for targeted cancer therapy.

Nature Biomedical Engineering **8**, 1453-1468 (2024). https://doi.org:10.1038/s41551-024-01214-6

30 Maguire, C. A. *et al.* Microvesicle-associated AAV Vector as a Novel Gene Delivery System.

Molecular Therapy **20**, 960-971 (2012). https://doi.org:https://doi.org/10.1038/mt.2011.303

31 Finkelshtein, D., Werman, A., Novick, D., Barak, S. & Rubinstein, M. LDL receptor and its family

members serve as the cellular receptors for vesicular stomatitis virus. *Proceedings of the National*

Academy of Sciences **110**, 7306-7311 (2013). https://doi.org:doi:10.1073/pnas.1214441110

32 Akache, B. *et al.* A two-hybrid screen identifies cathepsins B and L as uncoating factors for adeno-

associated virus 2 and 8. *Mol Ther* **15**, 330-339 (2007). https://doi.org:10.1038/sj.mt.6300053

33   Salganik, M. *et al.* Evidence for pH-dependent protease activity in the adeno-associated virus

capsid. *J Virol* **86**, 11877-11885 (2012). https://doi.org:10.1128/jvi.01717-12

34   Girod, A. *et al.* The VP1 capsid protein of adeno-associated virus type 2 is carrying a phospholipase

A2   domain   required   for   virus   infectivity.   *J   Gen   Virol*   **83**,   973-978   (2002).

https://doi.org:10.1099/0022-1317-83-5-973

35   Stahnke, S. *et al.* Intrinsic phospholipase A2 activity of adeno-associated virus is involved in

endosomal   escape   of   incoming   particles.   *Virology*   **409**,   77-83   (2011).

https://doi.org:10.1016/j.virol.2010.09.025

36   Guo, Y. *et al.* Early events in rabies virus infection-Attachment, entry, and intracellular trafficking.

Virus Res **263**, 217-225 (2019). https://doi.org:10.1016/j.virusres.2019.02.006

37   Gaudin, Y. Rabies Virus-Induced Membrane Fusion Pathway. *Journal of Cell Biology* **150**, 601-612

(2000). https://doi.org:10.1083/jcb.150.3.601

38   Albertini, A. A., Baquero, E., Ferlin, A. & Gaudin, Y. Molecular and cellular aspects of rhabdovirus

entry. *Viruses* **4**, 117-139 (2012). https://doi.org:10.3390/v4010117

39   Haas, D. L., Case, S. S., Crooks, G. M. & Kohn, D. B. Critical Factors Influencing Stable

Transduction of Human CD34+ Cells with HIV-1-Derived Lentiviral Vectors. *Molecular Therapy* **2**,

71-80 (2000). https://doi.org:https://doi.org/10.1006/mthe.2000.0094

40   Galla, M., Will, E., Kraunus, J., Chen, L. & Baum, C. Retroviral Pseudotransduction for Targeted

Cell   Manipulation.   *Molecular   Cell*   **16**,   309-315   (2004).

https://doi.org:https://doi.org/10.1016/j.molcel.2004.09.023

41   Nash, K. L. & Lever, A. M. L. Green fluorescent protein: green cells do not always indicate gene

expression. *Gene Therapy* **11**, 882-883 (2004). https://doi.org:10.1038/sj.gt.3302246

42   Geering, B., Schmidt-Mende, J., Federzoni, E., Stoeckle, C. & Simon, H.-U. Protein overexpression

following lentiviral infection of primary mature neutrophils is due to pseudotransduction. *Journal of*

Immunological   Methods   **373**,   209-218   (2011).

https://doi.org:https://doi.org/10.1016/j.jim.2011.08.024

Kim, J. T. *et al.* Dendritic cell–targeted lentiviral vector immunization uses pseudotransduction and DNA-mediated STING and cGAS activation. *Science Immunology* **2**, eaal1329 (2017). https://doi.org:doi:10.1126/sciimmunol.aal1329

Hung, M. E. & Leonard, J. N. A platform for actively loading cargo RNA to elucidate limiting steps in EV-mediated delivery. *J Extracell Vesicles* **5**, 31027 (2016). https://doi.org/10.3402/jev.v5.31027

Somiya, M. & Kuroda, S. i. Verification of extracellular vesicle-mediated functional mRNA delivery via RNA editing. *bioRxiv*, 2022.2001.2025.477620 (2022). https://doi.org:10.1101/2022.01.25.477620

Yeh, H. W. *et al.* ATP-Independent Bioluminescent Reporter Variants To Improve in Vivo Imaging. *ACS Chem Biol* **14**, 959-965 (2019). https://doi.org:10.1021/acschembio.9b00150

Bäck, S. *et al.* Neuron-Specific Genome Modification in the Adult Rat Brain Using CRISPR-Cas9 Transgenic Rats. *Neuron* **102**, 105-119.e108 (2019). https://doi.org:https://doi.org/10.1016/j.neuron.2019.01.035

Donahue, P. S. *et al.* The COMET toolkit for composing customizable genetic programs in mammalian cells. *Nature Communications* **11**, 779 (2020). https://doi.org:10.1038/s41467-019-14147-5

Votteler, J. *et al.* Designed proteins induce the formation of nanocage-containing extracellular vesicles. *Nature* **540**, 292-295 (2016). https://doi.org:10.1038/nature20607

Sauer, B. & Henderson, N. Targeted insertion of exogenous DNA into the eukaryotic genome by the Cre recombinase. *New Biol* **2**, 441-449 (1990).

Blessing, D., Déglon, N. & Schneider, B. L. in *Recombinant Protein Expression in Mammalian Cells: Methods and Protocols*   (ed David L. Hacker) 259-274 (Springer New York, 2018).

Gunnels, T. F., Stranford, D. M., Mitrut, R. E., Kamat, N. P. & Leonard, J. N. Elucidating Design Principles for Engineering Cell-Derived Vesicles to Inhibit SARS-CoV-2 Infection. *Small* **18**, 2200125 (2022). https://doi.org:https://doi.org/10.1002/smll.202200125

Gutierrez-Guerrero, A., Cosset, F.-L. & Verhoeyen, E. Lentiviral Vector Pseudotypes: Precious Tools to Improve Gene Modification of Hematopoietic Cells for Research and Gene Therapy. *Viruses* **12**, 1016 (2020).

54   Dautzenberg, I. J. C., Rabelink, M. J. W. E. & Hoeben, R. C. The stability of envelope-pseudotyped
lentiviral vectors. *Gene Therapy* **28**, 89-104 (2021). https://doi.org:10.1038/s41434-020-00193-y

55   Ma, D. *et al.* Engineered extracellular vesicles enable high-efficient delivery of intracellular
therapeutic proteins. *Protein & Cell* **15**, 724-743 (2024). https://doi.org:10.1093/procel/pwae015

56   Meyer, C. *et al.* Pseudotyping exosomes for enhanced protein delivery in mammalian cells. *Int J*
Nanomedicine **12**, 3153-3170 (2017). https://doi.org:10.2147/ijn.S133430

57   Kim, I. S. *et al.* Mechanism of membrane fusion induced by vesicular stomatitis virus G protein.
Proceedings   of   the   National   Academy   of   Sciences **114**,   E28-E36   (2017).
https://doi.org:doi:10.1073/pnas.1618883114

58   Stranford, D. M., Hung, M. E., Gargus, E. S., Shah, R. N. & Leonard, J. N. A Systematic Evaluation
of Factors Affecting Extracellular Vesicle Uptake by Breast Cancer Cells. *Tissue Eng Part A* **23**,
1274-1282 (2017). https://doi.org:10.1089/ten.TEA.2017.0158

59   Wang, J. *et al.* Highly efficient homology-driven genome editing in human T cells by combining zinc-
finger nuclease mRNA and AAV6 donor delivery. *Nucleic Acids Research* **44**, e30-e30 (2015).
https://doi.org:10.1093/nar/gkv1121

60   Ellis, B. L. *et al.* A survey of ex vivo/in vitro transduction efficiency of mammalian primary cells and
cell lines with Nine natural adeno-associated virus (AAV1-9) and one engineered adeno-associated
virus serotype. *Virology Journal* **10**, 74 (2013). https://doi.org:10.1186/1743-422X-10-74

61   Gutierrez-Guerrero, A., Cosset, F. L. & Verhoeyen, E. Lentiviral Vector Pseudotypes: Precious Tools
to Improve Gene Modification of Hematopoietic Cells for Research and Gene Therapy. *Viruses* **12**
(2020). https://doi.org:10.3390/v12091016

62   Fredericksen, B. L. & Whitt, M. A. Vesicular stomatitis virus glycoprotein mutations that affect
membrane fusion activity and abolish virus infectivity. *Journal of Virology* **69**, 1435-1443 (1995).
https://doi.org:10.1128/jvi.69.3.1435-1443.1995

63   Liang, X. *et al.* Engineering of extracellular vesicles for efficient intracellular delivery of multimodal
therapeutics   including   genome   editors.   *Nature   Communications* **16**,   4028   (2025).
https://doi.org:10.1038/s41467-025-59377-y

64 Nikolic, J. *et al.* Structural basis for the recognition of LDL-receptor family members by VSV

glycoprotein. *Nature Communications* **9**, 1029 (2018). https://doi.org:10.1038/s41467-018-03432-

4

65 Dobson, C. S. *et al.* Antigen identification and high-throughput interaction mapping by

reprogramming viral entry. *Nature Methods* **19**, 449-460 (2022). https://doi.org:10.1038/s41592-

022-01436-z

66 Gao, Y. & Bergman, I. Vesicular Stomatitis Virus (VSV) G Glycoprotein Can Be Modified to Create

a Her2/Neu-Targeted VSV That Eliminates Large Implanted Mammary Tumors. *Journal of Virology*

97, e00372-00323 (2023). https://doi.org:10.1128/jvi.00372-23

67 Issa, S. S., Shaimardanova, A. A., Solovyeva, V. V. & Rizvanov, A. A. Various AAV Serotypes and

Their Applications in Gene Therapy: An Overview. *Cells* **12** (2023).

https://doi.org:10.3390/cells12050785

68 Kovács, O. T. *et al.* Impact of Medium-Sized Extracellular Vesicles on the Transduction Efficiency

of Adeno-Associated Viruses in Neuronal and Primary Astrocyte Cell Cultures. *Int J Mol Sci* **22**

(2021). https://doi.org:10.3390/ijms22084221

69 Nisanov, A. M., de Jesús, J. A. R. & Schaffer, D. V. Advances in AAV capsid engineering: Integrating

rational design, directed evolution and machine learning. *Molecular Therapy* **33**, 1937-1945 (2025).

70 Kim, H., Kim, M., Im, S. K. & Fang, S. Mouse Cre-LoxP system: general principles to determine

tissue-specific roles of target genes. *Lab Anim Res* **34**, 147-159 (2018).

https://doi.org:10.5625/lar.2018.34.4.147

FIGURES

Figure 1

885   Figure 2

888   Figure 3

Figure 4

Figure 5

 FIGURE CAPTIONS

Figure 1: EV-AAV pseudotransduction complicates distinguishing protein delivery from gene

delivery. (A) EV-AAV-mediated protein delivery and/or expression in HEK293FT recipient cells. Vector

compositions and EV-AAV purification methods employed to separate EVs and free AAV are summarized

in cartoons at left. Free AAV (which should confer minimal protein delivery) conferred significantly greater

fluorescence than did EV-AAV controls. EV-AAV vectors were generally similar to the No Rep/Cap (protein

delivery only) control at all time points post-delivery. **(B)** Representative histograms of LumiScarlet

fluorescence from samples reported in (A). No subpopulations are apparent than can rigorously distinguish

protein delivery from gene delivery. Experiments were performed in biological triplicate. Data shown are

from one of two independent experiments. Error bars indicate standard error of the mean. Multicomparison

statistical analysis was performed using a two-way ANOVA test, followed by Tukey's multiple comparison

test to evaluate specific comparisons (****p < 0.0001). ns; not significant; PBS: phosphate-buffered saline;

SEC: size exclusion chromatography.

Figure 2: A Cre recombinase-based dual reporter system distinguishes protein delivery and gene

delivery. (A) Schematic depicting the expected function of the stoplight dual-reporter system in which Cre

expressed in recipient cells acts upon incoming AAV genomes to switch them from expression of a red

reporter to a green reporter. Free AAV mediates gene delivery only (green); EV-AAV mediates both protein

delivery (red) and gene delivery (green) and sometimes both (yellow); No Rep/Cap EVs can only mediate

protein delivery (red). **(B)** Cre-responsive plasmids evaluated in this experiment. **(C,D)** Protein delivery (C)

and gene delivery (D) conferred by various vector compositions. Note: some mCherry gene expression

could occur in recipient cells prior to Cre-mediated recombination (or in the possible absence of

recombination in some cells), such that (C) includes both protein delivery and this ambiguous de novo gene

expression. Samples were normalized to include 1e9 vector genomes (for AAV crude lysate conditions) or

a volume-equivalent of the AAV2 crude lysate condition for No Rep/Cap crude lysate conditions (10^4

recipient cells). Experiments were performed in biological triplicate. Data shown are from one of two

independent experiments. Error bars indicate standard error of the mean. Multicomparison statistical

analysis was performed using a two-way ANOVA test, followed by Tukey's multiple comparison test to

evaluate specific comparisons (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). NLS: nuclear

localization sequence; ns: not significant; PBS: phosphate-buffered saline.

Figure 3: EV-AAV2 functionalization increases protein delivery but not gene delivery to HEK293FTs.

(A-C) Validation of Ultra-HEK (Syenex) for enhancing lentiviral delivery to HEK293T and HEK293FT cells.

Relative fold-improvements in functional titer compared to a standard VSV-G-pseudotyped lentiviral vector

are indicated (**D**) Schematic depicting vectors used in (E,F) (Cre-HEK293FT recipient cells). (**E,F**) Protein

(E) and gene (F) delivery to HEK293FT cells (5 days post-delivery). Samples were normalized to include

1e9 vector genomes per condition (10^4 recipient cells). VSV-G and Ultra-HEK-functionalized EV-AAV

increase protein delivery but only modestly increase gene delivery and only when combined. Experiments

were performed in biological triplicate. Data shown are from one of two independent experiments. Error

bars indicate standard error of the mean. Multicomparison statistical analysis was performed using a one-

way ANOVA test, followed by Tukey's multiple comparison test to evaluate specific comparisons (*$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). PBS: phosphate-buffered saline.

Figure 4: VSV-G functionalization of EV-AAV6 increases gene and protein delivery to Jurkat T cells

by increasing uptake. (A) Schematic depicting vectors used in (B,C) (Cre-Jurkat T cell recipient cells).

(**B,C**) Protein (B) and gene (C) delivery to Jurkat T cells (5 days post-delivery). Samples were normalized

to include 1e9 vector genomes per condition (10^4 recipient cells). VSV-G functionalization significantly

increases protein delivery and gene delivery to Jurkats. **(D)** Schematic depicting vectors used in (E,F) (Cre-

Jurkat T cell recipient cells). (**E,F**) Protein (E) and gene (F) delivery to Jurkat T cells (5 days post-delivery).

Samples were normalized to include 2e9 vector genomes per condition (10^4 recipient cells). VSV-G

functionalization significantly increases protein delivery and gene delivery to Jurkats. Protein delivery and

gene delivery is increased only by enhanced binding of recipient cells (VSV-G WT and VSV-G P127D

vectors). Experiments were performed in biological triplicate. Data shown are from one of two independent

experiments. Error bars indicate standard error of the mean. Multicomparison statistical analysis was

performed using a one-way ANOVA test, followed by Tukey's multiple comparison test to evaluate specific

comparisons (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). PBS: phosphate-buffered saline.

Figure 5: Determinants of EV-AAV delivery to HEK293FT cells apply across AAV serotypes. (A)
Schematic depicting vectors used in (B,C) (Cre-HEK293FT recipient cells). (**B,C**) Protein (B) and gene (C)
delivery to HEK29FT cells (5 days post-delivery). Samples were normalized to include 1e9 vector genomes
per condition (10^4 recipient cells). VSV-G functionalization enhances protein delivery by not gene delivery
by EV-AAV6, concordant with observations made using EV-AAV2. Experiments were performed in
biological triplicate. Data shown are from one of two independent experiments. Error bars indicate standard
error of the mean. Multicomparison statistical analysis was performed using a one-way ANOVA test,
followed by Tukey's multiple comparison test to evaluate specific comparisons (*$p < 0.05$, **$p < 0.01$,
***$p < 0.001$, ****$p < 0.0001$). PBS: phosphate-buffered saline.

963