Aerobic scope is sustained through a heatwave in juvenile Atlantic salmon

Lucy Cotgrove[1], Sergey Morozov[2], Miika Raitakivi[3], Evan Sala[3], Jenni M. Prokkola[1]

[1] Natural Resources Institute Finland (Luke), Paavo Havaksen tie 3, 90570 Oulu, Finland

[2] Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental

Sciences, University of Helsinki, Viikinkaari 9, 00014 Helsinki, Finland

[3] Natural Resources Institute Finland (Luke), Vilppulantie 415, 41360 Laukaa, Finland

Corresponding author: lucy.cotgrove@luke.fi

Lay Summary

In juvenile Atlantic salmon, standard metabolic rate increases but there is no difference in maximum

metabolic rate and aerobic scope with increasing temperatures. Considering the increased metabolic

costs of activities at higher temperatures, our results show that juvenile salmon are vulnerable to

deteriorating performance at temperatures commonly experienced in present-day heatwaves.

Abstract

Aquatic ectotherms are vulnerable to heatwave-induced physiological stress, which arises from increased

energy demands and reduced dissolved oxygen content in warmer waters. Understanding thermal physiology

is critical for predicting how commercially and ecologically important populations could be affected by the

increasing risk of rising temperatures. Heatwave risk assessments often examine extremities of time scales:

immediate impacts or long-term consequences. However, little is known about how consistently increasing

mid-term thermal stress shapes aerobic performance in commercially important species such as Atlantic

salmon (*Salmo salar*), which may face heat stress in rivers, especially at the juvenile life stage. By measuring

how salmon juveniles manage their aerobic capacity at 16, 19 and 22°C using intermittent respirometry, we

test if their thermal performance curve shows a decline at temperatures commonly occurring during

heatwaves. Whole-animal metabolism was measured from control individuals kept at 16 °C before and after

the heatwave, and after 4-5 days exposure at 19 and 22°C during the heatwave. We show that standard

metabolic rate increases with temperature, but maximum metabolic rate and aerobic scope do not change

between these temperatures. These findings suggest that juvenile Atlantic salmon may have limited capacity to

increase aerobic performance during moderate heatwaves, leaving them vulnerable to cumulative effects of

oxygen limitation to vital functions such as growth and stress responses. As climate change intensifies,

incorporating thermal performance curves into conservation strategies can be utilized for predicting population

resilience and informing effective management.

Keywords

Respirometry, thermal performance, ecophysiology, energetics

Introduction

With climate change increasing the severity and frequency of heatwaves (IPCC, 2023), organisms may experience physiological thermal stress and even mortality due to extreme temperature fluctuations. Understanding organisms' thermal physiology is therefore critically important for conservation (Clark et al., 2013; Deutsch et al., 2008). Heatwaves frequently surpass the temperature range that species have been adapted to, often cumulatively and repeatedly, imposing selection on thermal performance. This is especially true for ectotherms, whose internal temperature are regulated by the surrounding environment. Some species may demonstrate adaptive or resilient responses to heatwaves including behavioural adjustments, physiological acclimation, or shifts in geographical ranges (Breedveld et al., 2023; Burraco et al., 2020; Jørgensen et al., 2022). However, these responses may not always be sufficient to counteract the negative consequences of extreme weather fluctuation (Perry et al., 2005; Pörtner & Farrell, 2008). Most physiological processes in ectotherms, including development, reproduction, and metabolism, are temperature-dependent (Angilletta et al., 2010; Hochachka & Somero, 1973; Schulte, 2015; Seebacher, 2009). These processes can be described using thermal performance curves (TPCs), which illustrate how a biological rate, such as oxygen consumption or activity per unit time, changes with temperature (Schulte et al., 2011).

TPCs can be estimated using thermal windows, critical temperatures, and preferred temperature ranges (Jørgensen et al., 2021; Ørsted et al., 2022; Peralta-Maraver & Rezende, 2021). TPCs can be used to compare inter- and intraspecies responses to anticipated temperatures under a future climate and how this might vary across life stages (Lefevre et al., 2021; Seebacher & Little, 2021). In water-breathing aquatic ectotherms, such as fishes, TPCs linked to demand, uptake and delivery of oxygen

are particularly important because of the decreasing solubility but increasing oxygen consumption in

warmer water (Little et al., 2020; Little & Seebacher, 2021; Seebacher & Little, 2021; Verberk et al.,

2022). Consequently, TPCs related to metabolic rates and aerobic performance are pivotal to

understanding the effects of heatwaves on fishes.

Metabolic rate is a fundamental aspect of fish physiology, reflecting the energy expenditure

associated with basic life processes such as respiration, digestion, and movement (Brett, 1964; Fry et

al., 1947). Standard metabolic rate (SMR) of an animal represents the baseline rate of aerobic

metabolism required to sustain life in a post-absorptive, resting state, while the maximum capacity for

aerobic performance is set by the maximum metabolic rate (MMR) (Sandblom et al., 2016). The

difference between SMR and MMR represents the aerobic scope (AS) of an animal and theoretically

determines the capacity of aerobic metabolism to support key life-history attributes such as activity,

growth and reproduction, each of which has a specific oxygen cost (Fry, 1971; Fry et al., 1947).

Therefore, reductions in aerobic scope can impair physical performance (Johansen & Jones, 2011;

Pörtner & Farrell, 2008; Priede, 1985). Given that the SMR of ectotherms rises with temperature and

often no such increase is detected in MMR (Fry et al., 1947; Norin et al., 2014; Sandblom et al., 2016),

increases in temperature beyond optimum can result in a decline in AS, as depicted by the TPC

(Farrell, 2016; Fry, 1971; Lefevre et al., 2021). Increased temperature, and decreasing AS, potentially

causes declining growth, health, or survival of individuals (Alfonso et al., 2021; Jutfelt et al., 2021;

Navarro et al., 2019; Sadoul & Vijayan, 2016). Specifically in salmonids, which include economically

important, cold-water adapted species, reduction of AS has already been linked to reduced growth,

survival, or collapse of aerobic capability during heatwaves (Eliason, Wilson, et al., 2013; Pörtner &

Farrell, 2008; Wade et al., 2019).

The Atlantic salmon (*Salmo salar* L.) is faced with supra-optimal temperatures especially at riverine life stages, including juveniles (parr) and adults during their spawning migration, with reports of present-day river temperatures up to 28 °C (O'Sullivan et al., 2023; Strøm et al., 2019). Juvenile Atlantic salmon rear for one to five years in freshwater before migrating to the sea (Aas et al., 2010), with potential to encounter multiple heatwaves. These can have direct effects on survival, but also cascading effects on population dynamics through reduced growth rates. Faster freshwater growth in salmon is linked to higher survival and faster maturation at sea (Hutchings & Jones, 1998; Simpson, 1992; Thorpe, 2007). In some populations, a negative association between freshwater growth and river temperature has been detected (Alioravainen et al., 2023). For the majority of their life cycle at sea, Atlantic salmon will experience temperatures below 8°C (Jensen et al., 2014; Lacroix, 2013; Reddin, 1985). Atlantic salmon tend to avoid temperatures above 15°C (Fisher & Elson, 1950; Johansson et al., 2009; Lacroix, 2013), and their optimal temperature range for feeding and normal behaviour in the wild has been suggested to be 6–20°C, with peak growth rates occurring at 16–17°C in aquaculture conditions (Dwyer & Piper, 1987; Elliott, 1982). Previous work assessing thermal performance in Atlantic salmon has often focussed on upper limits during acute exposure (Desforges et al., 2023 and papers within), or long-term stable temperature increases (Anttila et al., 2014; Del Rio et al., 2021; Hvas et al., 2017). When aerobic scope has been measured in a heatwave-context in a related species, (Casselman et al., 2012) found AS peaked at 17 °C, and decreased until testing finished at 21 °C in juvenile coho salmon *(Oncorhynchus kisutch)*. In post-smolt Atlantic salmon, AS tended to increase (a non-significant effect) as temperature increased from 13 °C to 23 °C, but swimming ability and feeding rate were greatly decreased and mortality increased, when metabolic rates of salmon were tested in groups without a cumulative exposure (Hvas et al., 2017). Further,

thermal variability in acclimation regime had no effect on salmon AS compared to a stable acclimation

when metabolic rates were tested acutely at the same temperature (Morissette et al., 2021). Still,

information on the cumulative effects of thermal fluctuations over several days or weeks, which

better reflect natural heatwave patterns, are limited (Morash et al., 2021; Nuic et al., 2024). This

information is pivotal for the conservation of Atlantic salmon. The diminishing of salmon populations

has already reduced opportunities for commercial and recreational fishery, cultural traditions, and

increased the risk of local extinctions (ICES, 2024, Dadswell et al., 2022)

Here, we test how mid-term increasing temperatures that are prevalent in rivers during heatwaves

impact metabolic rates and AS in hatchery-reared juvenile Atlantic salmon. We hypothesise that SMR

will increase as temperature rises. We expect MMR to show little variation between temperatures,

and therefore a decrease in AS as temperature increases. These findings will provide insight into the

physiological constraints juvenile salmon may face during heatwaves, using an ecologically relevant

temperature regime.

Methods

Fish Husbandry

The experiment was performed with a permit granted by the Finnish Project Authorisation Board (no.

ESAVI/16748/2023). Families of Atlantic salmon were bred using a hatchery broodstock originating

from River Iijoki, maintained by Natural Resources Institute of Finland (Luke), in Taivalkoski, Finland.

The broodstock was established in 2012 from parents that were originally stocked into the river as

juveniles/smolts, and therefore had completed half of their life cycle in the wild. Individuals from the

broodstock were crossed to make full-sib families of offspring in October 2022. The embryos were

incubated in Taivalkoski hatchery in family-specific flow-through trays at natural temperatures of

incoming water from River Ohtaoja, and dead embryos were removed regularly.

Embryos were disinfected by iodine bath (Buffodine) on 9th March 2023, and transported to a LUKE

hatchery in Laukaa, Finland, where experiments were conducted. Survival between transportation

and start of the experiment was 22%. Excess mortality may have been caused by the closeness to the

embryo's hatching date and transporting to the fish farm (no disease was observed in the offspring).

Embryos from three unrelated families were incubated and reared in separate circular tanks

(diameter 80 cm) and supplied with a continuous flow of filtered water from a mix of local lake

Peurunkajärvi and river Peurunkajoki. Feeding of alevins was started with powdered feed (BioMar

Group) when most of the egg yolk had been consumed at approximately 25 May, 2023. Tanks were

cleaned by scrubbing surfaces and siphoning excess food twice a week until 10 June, 2023, and then

daily until the end of the experiment. Feed rations were calculated from growth predictions assuming

feed conversion efficiency of 0.8 (Elliott & Hurley, 1997), and adjusted such that a small amount of

feed was left uneaten. Fish were fed using 12h belt feeders with the daily ration, except for one day

preceding metabolic measurements, when fish were fasted. During the summer and the experiment,

fish were reared under constant light. Rearing temperature raised gradually with natural water

temperatures over approximately three months until the experiment was started; temperature varied

from 9.2 to 15.7 °C between first feeding and experimental period.

At the start of the experiment 27 July, 2023, families were divided into two groups, control and

heatwave, with approximately same number of individuals in each group within a family (N = 24 to 62

within family /tank). Fish were kept in three tanks each for control and heatwave groups. Densities

within families were kept similar between control and heatwave tanks throughout the experiment by

simultaneously removing fish from control tanks to match densities in of heatwave tanks. Control

tanks were kept at close to 16 °C (mean = 16.1, SD = 0.5) throughout the experimental period by

regulating the incoming water flows from the river and the lake with a natural temperature

difference. Temperatures in one control tank (the same water source used in all three tanks) and the

three heatwave tanks were measured every 30 min with Hobo 64K Pendant temperature loggers

(Hobo, Bourne, MA, USA).

Temperatures of the heatwave tanks were regulated via a heating tank to offset the colder

temperatures of the room and incoming water using a temperature controller (TS1000, H-Tronic

GmbH, Germany) with a temperature probe PT-1000 located in each tank, and an Eheim 600 or 300

pump (Eheim, Deizisau, Germany) depending on the distance from the heating tank. The pumps

circulated water through stainless steel coils in the heating tank, in which temperature fluctuated

between 30-35 °C. Heatwave tank water was aerated constantly to prevent declines in dissolved O_2

related to increasing temperatures, as the flow-through rate of fresh water to the tanks was reduced

(approx. 1 L min^{-1}) during the heatwave to maintain temperatures.

Experimental Heatwave

Fish in the heatwave treatment were exposed to three test temperatures: 16 °C, 19 °C, and 22 °C

(Figure 1). First, 16 °C was maintained as a control temperature for eight days, after which 3-8

individuals from each family were randomly picked for measurements of metabolic rate (similar

sample size across temperatures within a family), and moved into a fasting tank, where they were

unfed until the measurements started the following day. Temperature of the fasting tank was the

same as the heatwave tanks. After the measurement of MMR and SMR at 16 °C, temperature was

increased by one degree per day for three days, and maintained for 4-5 days, after which metabolic

rate measurements were taken from a new set of fish at 19 °C. This was repeated and metabolic rate

measurements were taken at 22°C. Due to a malfunctioning temperature controller at 19 °C (after

moving fish to acclimation for B3 and B4), there was an increase in temperature in one of the

heatwave tanks to 21.5 °C (Figure 1). Metabolic rate experiments were conducted once per individual,

and fish were euthanised after the SMR measurement.

To measure MMR, fish were transferred from the fasting tank into 50 L buckets filled with 5 cm of

water at the testing temperature using plastic cups of water to reduce air exposure. Fish were

encouraged to swim by hand around the circular container (Prokkola et al., 2022; Raby et al., 2020).

Chasing lasted 2 min, and all fish were fatigued by the end. The fish were then rapidly transferred to

respirometry chambers using plastic cups with water. Two people chased one fish each at once, and

each batch of max. 16 individuals was processed within 2 h. Fish were left in respirometry chambers

overnight in order to capture the MMR and SMR for each fish, as per (Chabot, McKenzie, et al., 2016).

Chambers were flushed with fresh water from the tank for 5 min every 15 min, allowing fully aerated

water to enter the chamber. The lowest recorded oxygen level in any chamber was 6.7 mg L $^{-1}$ at 22

°C. After SMR measurements, fish were euthanised using an overdose of MS-222, measured, and

weighed.

Immediately before and after metabolic measurements, background respiration was measured in

empty chambers for three measurement cycles (5 min flush, 15 min measurement) and an average of

these slopes was used to adjust the data for MMR and SMR. For those trials which took place

immediately after the previous, the post-trial background data of the previous trial was used as the

pre-trial background values. To prevent bacterial build-up, the system was bleached between

temperature changes, and all chamber parts scrubbed to limit bacterial growth and rinsed thoroughly.

Additionally, the water in the respirometry tank was continuously circulated through a UV filter.

A total of 120 fish were subjected to intermittent flow respirometry after a 24h fasting period to

provide estimates of metabolic rate (SMR, MMR, AS see table S1) (Killen et al., 2021; Svendsen et al.,

2016). Due to death in respirometer chambers, exclusion due to outliers, incomplete data recording,

and due to computer failure, 72 fish were included in the analysis (Table S2).

Respirometer design

For a summary of the respirometry design and measurement protocol, see Supplementary Table 1,

after (Killen et al., 2021). A 16-chamber intermittent flow respirometer, each chamber containing an

individual fish, was submerged in a temperature-regulated water bath (16, 19 & 22 ± 0.1 °C; 200 L) that

was saturated with air. The oxygen content of the water in the chambers was measured every two

seconds using a 4-channel fibre optic oxygen meter with associated oxygen sensors and software

(FireStingO2; PyroScience GmbH, Aachen, Germany). All 16 O_2 sensors were calibrated for 0% oxygen

saturation simultaneously with O_2-free water, made using sodium sulphate, and simultaneous for 100%

oxygen saturation with air-saturated water using an air stone before the first measurements at each

temperature. Temperature-compensation for O_2 saturation was based on PT100 temperature sensors,

which were placed in the middle of the respirometer tank and connected to each of the Firesting

meters. The chambers were shielded from disturbance and light using an opaque plastic cover during

the measurement of SMR. Flush and recirculation pumps were controlled using PumpResp controllers

(4-channel model, FishResp, Finland, https://github.com/embedded-sergey/PumpResp, (2024)). The

temperature of the respirometer tank was maintained using a temperature controller (as in the

heatwave tanks) and a reservoir connected to a TECO 2000 Chiller/heater. The reservoir received

constant inflow of water (approx. 1 L min^{-1}) from the same source as the rearing tanks to maintain good

water quality. Chambers for the respirometer were made using glass tubing (120 mm length, inner

diameter 38 mm, wall thickness 3.2 mm) and plastic caps that were 3D-printed on PA2200 polyamide.

The 'HeiBer'-caps were designed by Heidrikur Bergsson, University of Copenhagen

(https://zenodo.org/record/4062429#.YMSW7h1RVTY). Volume of chambers and tubing was 131.86 ±

1 mL (the exact volumes per chamber are reported in an online data repository, see Data Availability).

Water inside the chambers was mixed by a plastic disk attached to each cap to distribute the flow. The

disks were 3D printed using the same method as the respirometry caps and attached with stainless

steel screws. Caps were sealed using rubber O-rings and connected to the flush and recirculation pumps

and valves with gas-impermeable Tygon tubing (Tygon S3 E-3603, Satint-Gobain, Paris. France). The

recirculation system was confirmed to be waterproof by filling with water and plugging the flush pump

and probe connections. Water was recirculated using one submersible pump (12V DC, 6W 2 L min^{-1}

pump, Qingdao Xinhui Hardware Machinery Co., Ltd., Qingdao) per channel within the recirculation

loop. Chambers were flushed using a second pump (same as previous), using an inflow of water from

the tank into the chambers, and excess water was flushed through the chamber to a flush pipe which

was placed with the end above the water surface. This allowed fully aerated water to enter the

chamber. Flow rate through both the flush line and recirculation loop could be controlled by the valves

attached within the flow. Flow rate of the respirometers was approximately 0.1 (± 0.008) L min^{-1}

without fish. Information about time and pump phase (either flush or measurement) was recorded by

the PumpResp controllers and stored on a computer. Flow had no apparent effect on movement of fish

within the chambers in SMR or MMR measurements.

Data Analysis

All data and statistical analyses were done in R enviroment (R Core Team ,2022). The FishResp

package (Morozov et al., 2019) was used to filter and calculate metabolic rate estimates. Slopes of

oxygen consumption were adjusted for bacterial oxygen consumption using pre-and post-background

measurements, assuming a linear change. For SMR, oxygen consumption (mg O_2 h^{-1}) for each

measurement phase was derived from the slope of linear regression of dissolved oxygen

concentration over time. The SMR slopes were quality filtered and smoothed before analysis to

exclude non-linear declines of oxygen: first, slopes were filtered based on $R^2 > 0.95$, then slopes that

did not meet the R^2 criteria were smoothed using a running mean of 29 seconds (Chabot et al., 2021),

then R^2 filter was re-applied, and finally, linearity was checked visually from all slopes that met the

criteria of $R^2 > 0.95$. The first and last 60 seconds of measurement periods were excluded to account

for mixing of water and changes in flow due to flushing. The mean of the lowest normal distribution

(MLND) was used to estimate SMR for each individual from the extracted slopes (Chabot, Steffensen,

et al., 2016). Slopes for MMR were extracted using a derivative of a polynomial curve fitted on each

measurement (function *smooth.spline,* d.f. = 10) as in the *spline-MMR* method (Prokkola et al., 2022),

i.e. from the beginning of the MMR measurement, when respiration was the highest. The slopes were

then used to calculate MMR (mg O_2 h^{-1}) using *FishResp*. AS was calculated as the difference between

the absolute MMR and SMR.

A linear mixed effect model was fitted (estimated using restricted maximum likelihood and *nloptwrap*

optimizer) to predict SMR, MMR and a non-linear mixed effect model was fitted to predict AS, all of

which were log transformed. Temperature and log-transformed mass as fixed effects (formula:

Metabolic Measurement ~ Temperature + Mass). The model included batch as a random effect

(formula:~1|Batch) (Zuur et al., 2009). Homoscedasticity and normality of residuals were assessed by

visual inspection of residual plots. 95% Confidence Intervals (CIs) and p-values were computed using a

Wald t-distribution approximation. Post-hoc comparisons of temperatures were performed using

Tukey method of comparing estimates, and P value < 0.05 was considered significant. Statistical

models were fitted using *nlme*, *lme4* and *lmerTest* packages in R, and post-hoc analysis was

performed in *emmeans* package (Bates et al., 2015; Kuznetsova et al., 2017; Lenth, 2017; Pinheiro et

al., 1999). For visual representation, SMR, MMR, and AS data were mass-adjusted to account for the

hypo-allometric scaling of metabolic rates by linear regression of $\log_{10}$-transformed metabolic rates

against $\log_{10}$-transformed body mass (Figure 2). Finally, Pearson's correlations were calculated among

the either absolute or mass-adjusted metabolic variables.

Results

In total, the data of 72 fish was used at three temperatures across four consecutive weeks: 16°C,

19°C, and 22°C and 16 °C again (Table 1, Table S3). The metabolic rates and AS scaled positively with

body mass, with scaling exponents from 0.86 to 1.1 (Figure 2, Table 2, Table S4).

We found temperature to be a significant predictor of SMR (16–19 °C: $t(66) = 4.80$, $P < .001$; 16–22 °C:

$t(66) = 7.43$, $P < .001$, Table 3). There were significant differences in SMR between all temperature

groups, except between 19 °C and 22 °C (Figure 3, Table 4). There was a 57% increase in predicted

SMR for 6 °C increase in temperature (Table 4). However, there was no significant change in MMR or

AS between temperatures (Figure 3, Table 2).

Correlations among metabolic variables

At all temperatures, MMR and AS were highly positively correlated ($r > 0.9$) in both raw and mass

adjusted data. Absolute SMR and MMR were positively correlated at 16 and 22 °C. Absolute SMR and

AS were positively correlated at 16 °C. However, mass adjusted SMR and AS were negatively

correlated at 19 and 22 °C (Table S5, Figure S1).

Discussion

This study demonstrates that rising temperatures lead to an increase in standard metabolic rate

(SMR) in juvenile Atlantic salmon. However, there was no change in maximum metabolic rate (MMR)

or, contrary to our hypothesis, in aerobic scope (AS) across the tested temperatures. While an

increase in SMR with temperature is well established in ectotherms due to the effect of temperature

on biochemical rates (Clark et al., 2013; McKenzie et al., 2021; Raby et al., 2016), there are more

mixed results on the relationship of MMR, AS and increasing temperatures in fishes.

The lack of differences between the MMR within temperatures in this study was not surprising, given

that the individuals were similar in age, with identical handling and rearing environment (Farrell et al.,

2009a; Killen et al., 2016). MMR represents the maximum rate at which aerobic physiological

processes can occur, and at 16 °C the salmon may have already reached their maximum capacity for

oxygen uptake and delivery (Clark et al., 2011; McKenzie & Claireaux, 2010; Norin & Clark, 2016;

Sandblom et al., 2016). However, there may be among-individual variation in thermal reaction norms

of SMR and MMR that was not possible to detect with our experimental design, which could explain

why AS was not decreasing even though on average MMR was stable while SMR was increasing (Norin

et al., 2014; Norin & Metcalfe, 2019; Réveillon et al., 2019). In gobies (*Neogobius melanostomus)*,

MMR increased at the very highest temperatures tested; Christensen et al. (2020) speculate that

increases in MMR could be due to the onset of physiologically demanding processes for coping with

high temperatures such as heat shock protein and enzyme production (Christensen et al., 2020;

Loughland & Seebacher, 2020). Other studies have shown that MMR can increase with increasing

temperatures (*Lates clacarifer*: Norin et al 2014; *Dicentrarchus labrax* Claireux et al. 2006)(Claireaux

et al., 2006; Norin et al., 2014), however these studies are based on acute temperature increases. In

O. nerka and *O. kisutch*, MMR continuously increased in acclimation until 22-25 °C, where it plateaued

(Eliason et al., 2011). This plateau may reflect the species' maximum aerobic capacity, similar to what

may be occurring in the present study.

The TPC of AS in fish has been suggested to be a bell shaped curve (Pörtner & Farrell, 2008). For

example, in pink salmon (*O. gorbuscha*), AS peaked at 21 °C and declined with further warming,

although this temperature increase occurred over a maximum of 2 days (Clark et al., 2011). Other

studies note a decrease in AS, which could elude to the post peak decrease of the AS curve, as these

are tested at higher temperatures or after longer acclimations, for example, in a recent study on

salmon parr, AS decreased with temperature from 15 to 21°C after a three-week acclimation (Nuic et

al., 2024). This could suggest long term exposure to a heatwave leading to aerobic collapse. In

contrast in murray cod (*Maccullochella peelii peelii*) and temperatures that increased from 14 – 29 °C,

where fish were acclimated for at least 18 h to each increase, resulted in a continuous increase in

aerobic scope (Clark et al., 2005).

Other studies report a plateau of AS at high temperatures. For example, there were no differences in

AS between acclimation temperatures in minnows (*Phoxinus phoxinus*) after eight months (Závorka et

al., 2020). Morissette et al. (2021) found that below 23 °C, there was no difference between any

metabolic rates after different temperature acclimations, however in their study all metabolic rates

were measured at the same temperature, which complicates the interpretation of the results. In

gobies acclimated for three weeks, AS was seen to plateau between 15 and 28 °C after initially

increasing (Christensen et al., 2021). Further, in Atlantic salmon post-smolts, aerobic scope was

increased with 4-week acclimations at temperatures from 3–13 °C, and plateaued from 13 °C up to a

temperature of 23 °C. Interestingly, fish condition and swim speed dropped and mortality increased

(Hvas et al., 2017). Upper thermal limit of salmon has been widely reported at 23 °C (Elliott & Elliott,

2010), and notable, often cited work (Jensen et al., 1989) found optimal salmon temperature for

growth to be 16 °C. Our results agree with Hvas et al. (2017) as we found that AS plateaued at 16–

22°C when measured at acclimation temperature. However, even though the expectation of reduced

AS with increasing temperature was not met in this study, it is likely that oxygen supply can limit

performance of salmon as temperature increases. For example, maintaining growth and other aerobic

functions with increasing temperature would require an increasing AS, because of the higher costs of

activities such as digestion and assimilation and swimming (Jahn & Seebacher, 2022; Jutfelt et al.,

2021). Also the stress-induced peak oxygen consumption increases exponentially with temperature

(Madaro et al., 2018). An absence of increase suggests physiological systems supporting oxygen

uptake such as cardiac function, gill function and haemoglobin oxygen affinity are no longer able to

scale with increasing thermal pressure (Anttila et al., 2014; Schulte & Healy, 2022). As AS plateaus but

SMR and the costs of aerobic activities continue to increase, fish might become more constrained in

their physiological state and experience cumulative thermal stress, leading to mortality.

AS is proposed to be of ecological significance because it defines the upper limit for oxygen allocation

by a fish to sustain aerobic activities such as foraging, digestion, tissue deposition, migration,

reproduction, and so forth (Claireaux & Lefrançois, 2007; Farrell et al., 2009b; Fry, 1971; H.-O.

Pörtner, 2010; Schulte, 2015). For example, reduced appetite could be behaviourally driven to avoid

anaerobic metabolism, as suggested by the reduction of feed intake during low AS (review by (Jutfelt,

2020). In support of this, reduced growth and lower condition factor at higher temperatures has

previously been documented in Atlantic salmon, when comparing increases to temperatures up to 19

°C (Hevrøy et al., 2015; Kullgren et al., 2013). By measuring AS as an indicator of thermal

performance, we can predict that at high temperatures, Atlantic salmon in the wild would either have

to seek thermal refuge, or compromise aspects of their performance in order to cope with the high

temperatures, which could result in poor growth or health (Koskela et al., 1997).

It is important to note that both TPCs and upper thermal limits are dependent on acclimation

duration and rate of warming (Currie et al., 2014; Lefevre et al., 2021; McKenzie et al., 2021). Most

research in fishes has been conducted with stable thermal profiles, whereby each individual

experiences only one acclimation temperature (Hvas et al., 2017). In contrast, our study accounts for

a cumulative effect of exposure to higher temperatures, using a gradual heating regime. While some

acclimation is expected in the MMR over short-term temperature increases ((Norin & Clark, 2016);

and papers within), each temperature exposure in this study occurred over 10 days, which may not be

enough time to see notable plasticity in salmon's metabolic ceiling. Sandblom et al. (2017) found no

difference in MMR or AS between acute (24 h) or chronic warming (years) regime in perch (*Perca*

fluviatilis), but that all warm-acclimated fish (5 days acclimation, 23 °C) have a higher MMR than fish

acclimated at 18 °C, their reference temperature. However, an ecologically relevant scenario for

salmon in their environments is a heatwave lasting from a few days to few weeks, thus these time

frames are the most relevant to study from a conservation perspective. By testing the cumulative

effect of a heatwave mirroring the temperatures seen in the wild, we can better predict how salmon

could cope in a realistic exposure to sub-lethal temperatures (Morash et al., 2018).

The general expectation is that longer acclimation with a slower increase of temperature should be

beneficial to survival, giving the fish time to make compensatory physiological modifications such as

displaying plasticity in SMR or MMR, enabling AS to be maintained over a broad range of

temperatures. However, in barramundi (*L. calcarifer*) MMR and AS increased when exposed to acute

10 °C warming, due to a higher increase in MMR than SMR, but after 5 weeks of acclimation, AS was

similar, mainly due to a reduction in MMR (Norin et al., 2014). A very similar pattern was observed in

black sea bass (*Centropristis striata)* and common triplefin (*Forsterygion lapillum*), where short term

exposure caused a raise in AS, before it decreased after some weeks of exposure (Khan et al., 2014;

McArley et al., 2017; Slesinger et al., 2019). Additionally, in salmon post smolts, post-stress peak

cortisol levels significantly increased with higher temperatures (Madaro et al., 2018). This suggests

that despite a stable aerobic capacity to cope with temperature increases, costly physiological

mechanisms may be occurring, and therefore a plateau in AS does not necessarily suggest a broad

optimum temperature range.

More accurate predictions of species response to heatwaves under climate change can be achieved

using bioenergetics modelling rather than assuming physiological performance inferred by historical

distribution data (Baltar et al., 2019; Lefevre et al., 2021). For example, in sockeye salmon, the

decrease of AS at high temperatures was associated with a collapse in cardiac performance, and

applying these data to seasonal river temperatures can be used to predict population collapse

(Eliason, Clark, et al., 2013; Eliason et al., 2011; MacNutt et al., 2006). With detailed TPCs, thermal

safety margins can be defined in terms of functional success—such as swimming, feeding, and

reproducing—rather than mortality alone (Pinsky et al., 2019). Interestingly, (Morgan et al., 2022)

discuss plasticity of laboratory fish to temperatures, and show that wild fish have a different TPC to

multigenerational laboratory fish population. Building on this, we propose that our results could be

used to design more detailed studies on individual-level reaction norms and performance metrics at

the temperature range used in this study, which could ultimately inform both management and

conservation strategies.

Recent advances in conservation physiology demonstrate how individual-level physiological metrics

can be used to guide policy and management strategies, as shown in studies on sockeye salmon in

Canada (Patterson et al., 2016). Integrating TPCs and stock specific information, such as life history

genotypes and developmental stages, can inform stakeholders of specific vulnerabilities in

populations and how they vary in location and season (Ward et al., 2017). Temperature-dependent

fisheries management strategies have already been successfully implemented, for example, in

Canadian rivers, and could be adapted globally to account for population-specific TPCs (Van Leeuwen

et al., 2023). Management strategies, such as adjusting harvest timing and identifying thermal refuges

for protection and restoration efforts, could help mitigate the impact of rising temperatures on

vulnerable salmon populations.

In recent years, extreme heat events have had catastrophic impacts on migration success and survival

across species of salmonids, indicating that a better understanding of their thermal performance is

critical for conservation (Baisez et al., 2011; Hinch et al., 2012; Martins et al., 2011, 2012; Muñoz et

al., 2021). Our study showed that aerobic scope was not increased between 16 and 22°C in juvenile

Atlantic salmon despite a significant increase in SMR. This, combined with the increasing metabolic

costs of activities, such as swimming and digestion, indicate that the stable aerobic scope may be

leaving juvenile salmon vulnerable to deteriorating performance at temperatures commonly

experienced in present-day heatwaves.

Ethical Declarations

The experiment was performed with permit granted by the Finnish Project Authorisation Board no.

ESAVI/16748/2023.

Acknowledgements

We are grateful for Jonna Hänninen, Juha Hänninen, and staff at Luke Taivalkoski and Luke Laukaa for

access to fish and help during the experiment, and for Kari Jauhiainen for technical support, and Prof.

Craig Primmer, Dr. Tutku Aykanat, and Dr. Silva Uusi-Heikkilä for access to equipment. We thank Lilian

Redon for feedback on an early version of the manuscript.

Author Contributions

LC and JP: experimental design, conducting experiment, data analysis and writing the manuscript. JP:

supervision, management, and funding acquisition. SM: equipment and experimental design. MR and

ES: fish care, equipment maintenance and assisting with data collection.

Funding

This study was funded by the Research Council of Finland (348965 and 353760), Finnish Cultural

Foundation (00220693) and by the Natural Resources Institute Finland.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability

Data is hosted on Zenodo data repository (https://doi.org/10.5281/zenodo.15308028) and code for

analysis is hosted on Github (https://github.com/jprokkola/Salmon-aerobic-scope-2023-public.git).

References

Aas, Ø., Klemetsen, A., Einum, S., & Skurdal, J. (2010). Atlantic Salmon Ecology. *Salmon Ecological Research*

and Conservation. https://doi.org/10.1002/9781444327755

Alfonso, S., Gesto, M., & Sadoul, B. (2021). Temperature increase and its effects on fish stress physiology in the

context of global warming. *Journal of Fish Biology*, *98*(6), 1496–1508.

https://doi.org/10.1111/jfb.14599

Alioravainen, N., Orell, P., & Erkinaro, J. (2023). Long-Term Trends in Freshwater and Marine Growth Patterns

in Three Sub-Arctic Atlantic Salmon Populations. *Fishes*, *8*(9), 441.

https://doi.org/10.3390/fishes8090441

Angilletta, M. J., Huey, R. B., & Frazier, M. R. (2010). Thermodynamic Effects on Organismal Performance: Is

Hotter Better? *Physiological and Biochemical Zoology*, *83*(2), 197–206. https://doi.org/10.1086/648567

Anttila, K., Jørgensen, S. M., Casselman, M. T., Timmerhaus, G., Farrell, A. P., & Takle, H. (2014). Association

between swimming performance, cardiorespiratory morphometry, and thermal tolerance in Atlantic

salmon (Salmo salar L.). *Frontiers in Marine Science*, *1*. https://doi.org/10.3389/fmars.2014.00076

Baisez, A., Bach, J.-M., Leon, C., Parouty, T., Terrade, R., Hoffmann, M., & Laffaille, P. (2011). Migration delays

and mortality of adult Atlantic salmon Salmo salar en route to spawning grounds on the River Allier,

France. *Endangered Species Research*, *15*(3), 265–270. https://doi.org/10.3354/esr00384

Baltar, F., Bayer, B., Bednarsek, N., Deppeler, S., Escribano, R., Gonzalez, C. E., Hansman, R. L., Mishra, R. K.,

Moran, M. A., Repeta, D. J., Robinson, C., Sintes, E., Tamburini, C., Valentin, L. E., & Herndl, G. J. (2019).

Towards Integrating Evolution, Metabolism, and Climate Change Studies of Marine Ecosystems. *Trends*

in Ecology & Evolution, *34*(11), 1022–1033. https://doi.org/10.1016/j.tree.2019.07.003

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using **lme4**. *Journal*

of Statistical Software, *67*(1). https://doi.org/10.18637/jss.v067.i01

Breedveld, M. C., Devigili, A., Borgheresi, O., & Gasparini, C. (2023). Reproducing in hot water: Experimental

heatwaves deteriorate multiple reproductive traits in a freshwater ectotherm. *Functional Ecology*,

37(4), 989–1004. https://doi.org/10.1111/1365-2435.14279

Brett, J. R. (1964). The respiratory metabolism and swimming performance of young sockeye salmon. *Journal of*

Fish Biology, *21*(5), 1183–1226. https://doi.org/10.1139/f64-103

Burraco, P., Orizaola, G., Monaghan, P., & Metcalfe, N. B. (2020). Climate change and ageing in ectotherms.

Global Change Biology, *26*(10), 5371–5381. https://doi.org/10.1111/gcb.15305

Casselman, M. T., Anttila, K., & Farrell, A. P. (2012). Using maximum heart rate as a rapid screening tool to

determine optimum temperature for aerobic scope in Pacific salmon *Oncorhynchus* spp. *Journal of Fish*

Biology, *80*(2), 358–377. https://doi.org/10.1111/j.1095-8649.2011.03182.x

Chabot, D., McKenzie, D. J., & Craig, J. F. (2016). Metabolic rate in fishes: Definitions, methods and significance

for conservation physiology. *Journal of Fish Biology*, *88*(1), 1–9. https://doi.org/10.1111/jfb.12873

Chabot, D., Steffensen, J. F., & Farrell, A. P. (2016). The determination of standard metabolic rate in fishes.

Journal of Fish Biology, *88*(1), 81–121. https://doi.org/10.1111/jfb.12845

Chabot, D., Zhang, Y., & Farrell, A. P. (2021). Valid oxygen uptake measurements: Using high r^2 values with good

intentions can bias upward the determination of standard metabolic rate. *Journal of Fish Biology*, *98*(5),

1206–1216. https://doi.org/10.1111/jfb.14650

Christensen, E. A. F., Andersen, L. E. J., Bergsson, H., Steffensen, J. F., & Killen, S. S. (2021). Erratum: Shuttle-box

systems for studying preferred environmental ranges by aquatic animals (Conservation Physiology 9:1

(coab028) DOI: 10.1093/conphys/coab028). *Conservation Physiology*, *9*(1), 1–21.

https://doi.org/10.1093/conphys/coab051

Christensen, E. A. F., Norin, T., Tabak, I., Van Deurs, M., & Behrens, J. W. (2020). Effects of temperature on

physiological performance and behavioral thermoregulation in an invasive fish, the round goby. *Journal*

of Experimental Biology, jeb.237669. https://doi.org/10.1242/jeb.237669

Claireaux, G., Couturier, C., & Groison, A. L. (2006). Effect of temperature on maximum swimming speed and

cost of transport in juvenile European sea bass (Dicentrarchus labrax). *Journal of Experimental Biology*,

209(17), 3420–3428. https://doi.org/10.1242/jeb.02346

Claireaux, G., & Lefrançois, C. (2007). Linking environmental variability and fish performance: Integration

through the concept of scope for activity. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *362*(1487), 2031–2041. https://doi.org/10.1098/rstb.2007.2099

Clark, T. D., Jeffries, K. M., Hinch, S. G., & Farrell, A. P. (2011). Exceptional aerobic scope and cardiovascular

performance of pink salmon (*Oncorhynchus gorbuscha*) may underlie resilience in a warming climate.

Journal of Experimental Biology, *214*(18), 3074–3081. https://doi.org/10.1242/jeb.060517

Clark, T. D., Ryan, T., Ingram, B. A., Woakes, A. J., Butler, P. J., & Frappell, P. B. (2005). Factorial Aerobic Scope Is

Independent of Temperature and Primarily Modulated by Heart Rate in Exercising Murray Cod (

Maccullochella peelii peelii). *Physiological and Biochemical Zoology*, *78*(3), 347–355.

https://doi.org/10.1086/430034

Clark, T. D., Sandblom, E., & Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of climate

change: Respirometry, relevance and recommendations. *Journal of Experimental Biology*, *216*(15),

2771–2782. https://doi.org/10.1242/jeb.084251

Currie, S., Schulte, P. M., & Evans, D. H. (2014). Thermal stress. *The Physiology of Fishes*, *4*, 257–279.

Dadswell, M., Spares, A., Reader, J., McLean, M., McDermott, T., Samways, K., & Lilly, J. (2022). The Decline and

Impending Collapse of the Atlantic Salmon (*Salmo salar*) Population in the North Atlantic Ocean: A

Review of Possible Causes. *Reviews in Fisheries Science & Aquaculture*, *30*(2), 215–258.

https://doi.org/10.1080/23308249.2021.1937044

Del Rio, A. M., Mukai, G. N., Martin, B. T., Johnson, R. C., Fangue, N. A., Israel, J. A., & Todgham, A. E. (2021).

Differential sensitivity to warming and hypoxia during development and long-term effects of

developmental exposure in early life stage Chinook salmon. *Conservation Physiology*, *9*(1), coab054.

https://doi.org/10.1093/conphys/coab054

Desforges, J. E., Birnie-Gauvin, K., Jutfelt, F., Gilmour, K. M., Eliason, E. J., Dressler, T. L., McKenzie, D. J., Bates,

A. E., Lawrence, M. J., Fangue, N., & Cooke, S. J. (2023). The ecological relevance of critical thermal

maxima methodology for fishes. *Journal of Fish Biology*, *102*(5), 1000–1016.

https://doi.org/10.1111/jfb.15368

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P. R. (2008).

Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National*

Academy of Sciences, *105*(18), 6668–6672. https://doi.org/10.1073/pnas.0709472105

Dwyer, W. P., & Piper, R. G. (1987). Atlantic Salmon Growth Efficiency as Affected by Temperature. *The*

Progressive Fish-Culturist, *49*(1), 57–59. https://doi.org/10.1577/1548-

8640(1987)49<57:ASGEAA>2.0.CO;2

Eliason, E. J., Clark, T. D., Hague, M. J., Hanson, L. M., Gallagher, Z. S., Jeffries, K. M., Gale, M. K., Patterson, D.

A., Hinch, S. G., & Farrell, A. P. (2011). Differences in Thermal Tolerance Among Sockeye Salmon

Populations. *Science*, *332*(6025), 109–112. https://doi.org/10.1126/science.1199158

Eliason, E. J., Clark, T. D., Hinch, S. G., & Farrell, A. P. (2013). Cardiorespiratory collapse at high temperature in

swimming adult sockeye salmon. *Conservation Physiology*, *1*(1), cot008–cot008.

https://doi.org/10.1093/conphys/cot008

Eliason, E. J., Wilson, S. M., Farrell, A. P., Cooke, S. J., & Hinch, S. G. (2013). Low cardiac and aerobic scope in a

coastal population of sockeye salmon *Oncorhynchus nerka* with a short upriver migration. *Journal of*

Fish Biology, *82*(6), 2104–2112. https://doi.org/10.1111/jfb.12120

Elliott, J. M. (1982). The effects of temperature and ration size on the growth and energetics of salmonids in

captivity. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, *73*(1), 81–91.

https://doi.org/10.1016/0305-0491(82)90202-4

Elliott, J. M., & Elliott, J. A. (2010). Temperature requirements of Atlantic salmon Salmo salar, brown trout

Salmo trutta and Arctic charr Salvelinus alpinus: Predicting the effects of climate change. *Journal of Fish*

Biology, *77*(8), 1793–1817. https://doi.org/10.1111/j.1095-8649.2010.02762.x

Elliott, J. M., & Hurley, M. A. (1997). A functional model for maximum growth of Atlantic Salmon parr, Salmo

salar, from two populations in northwest England. *Functional Ecology*, *11*(5), 592–603.

https://doi.org/10.1046/j.1365-2435.1997.00130.x

Farrell, A. P. (2016). Pragmatic perspective on aerobic scope: Peaking, plummeting, pejus and apportioning.

Journal of Fish Biology, *88*(1), 322–343. https://doi.org/10.1111/jfb.12789

Farrell, A. P., Eliason, E. J., Sandblom, E., & Clark, T. D. (2009a). Fish cardiorespiratory physiology in an era of

climate changeThe present review is one of a series of occasional review articles that have been invited

by the Editors and will feature the broad range of disciplines and expertise represented in our Editorial

Advisory Board. *Canadian Journal of Zoology*, *87*(10), 835–851. https://doi.org/10.1139/Z09-092

Farrell, A. P., Eliason, E. J., Sandblom, E., & Clark, T. D. (2009b). Fish cardiorespiratory physiology in an era of

climate changeThe present review is one of a series of occasional review articles that have been invited

by the Editors and will feature the broad range of disciplines and expertise represented in our Editorial

Advisory Board. *Canadian Journal of Zoology*, *87*(10), 835–851. https://doi.org/10.1139/Z09-092

Fisher, K. C., & Elson, P. F. (1950). The Selected Temperature of Atlantic Salmon and Speckled Trout and the

Effect of Temperature on the Response to an Electrical Stimulus. *Physiological Zoology*, *23*(1), 27–34.

https://doi.org/10.1086/physzool.23.1.30084896

Fry, F. E. J. (1971). The Effect of Environmental Factors on the Physiology of Fish. In W. S. Hoar & D. J. Randall

(Eds.), *Fish Physiology* (Vol. 6, pp. 1–98). Academic Press. https://doi.org/10.1016/S1546-

5098(08)60146-6

Fry, F. E. J., Black, V. S., & Black, E. C. (1947). Influence of temperature on the asphyxiation of young goldfish

(carassius auratus l.) under various tensions of oxygen and carbon dioxide. *The Biological Bulletin*,

92(3), 217–224. https://doi.org/10.2307/1538308

Hevrøy, E. M., Tipsmark, C. K., Remø, S. C., Hansen, T., Fukuda, M., Torgersen, T., Vikeså, V., Olsvik, P. A.,

Waagbø, R., & Shimizu, M. (2015). Role of the GH-IGF-1 system in Atlantic salmon and rainbow trout

postsmolts at elevated water temperature. *Comparative Biochemistry and Physiology Part A: Molecular*

& Integrative Physiology, *188*, 127–138. https://doi.org/10.1016/j.cbpa.2015.06.030

Hinch, S. G., Cooke, S. J., Farrell, A. P., Miller, K. M., Lapointe, M., & Patterson, D. A. (2012). Dead fish

swimming: A review of research on the early migration and high premature mortality in adult Fraser

River sockeye salmon Oncorhynchus nerka. *Journal of Fish Biology*, *81*(2), 576–599.

https://doi.org/10.1111/j.1095-8649.2012.03360.x

Hochachka, P. W., & Somero, G. N. (1973). *Strategies of biochemical adaptation* (Vol. 358). Saunders.

Hutchings, J. A., & Jones, M. E. (1998). Life history variation and growth rate thresholds for maturity in Atlantic

salmon, Salmo salar. *Canadian Journal of Fisheries and Aquatic Sciences*, *55*(S1), 22–47.

https://doi.org/10.1139/d98-004

Hvas, M., Folkedal, O., Imsland, A., & Oppedal, F. (2017). The effect of thermal acclimation on aerobic scope

and critical swimming speed in Atlantic salmon *Salmo salar*. *Journal of Experimental Biology*,

jeb.154021. https://doi.org/10.1242/jeb.154021

ICES. (2023). *Contribution of Working Groups I, II and III to the Sixth Assessment Report of the*

Intergovernmental Panel on Climate Change (Synthesis Report No. 6; First, Climate Change 2023).

Intergovernmental Panel on Climate Change (IPCC). https://doi.org/10.59327/IPCC/AR6-

9789291691647

ICES. (2024a). *Baltic Salmon and Trout Assessment Working Group (WGBAST)* (ICES Scientific Report No. 1).

ICES. https://doi.org/10.17895/ices.pub.25868665.v1

ICES. (2024b). *Working Group on North Atlantic Salmon (WGNAS)* [Report]. ICES Scientific Reports.

https://doi.org/10.17895/ices.pub.25730247.v1

Jahn, M., & Seebacher, F. (2022). Variations in cost of transport and their ecological consequences: A review.

Journal of Experimental Biology, *225*(15), jeb243646. https://doi.org/10.1242/jeb.243646

Jensen, A. J., Karlsson, S., Fiske, P., Hansen, L. P., Østborg, G. M., & Hindar, K. (2014). Origin and life history of

Atlantic salmon (*Salmo salar*) near their northernmost oceanic limit. *Canadian Journal of Fisheries and*

Aquatic Sciences, *71*(11), 1740–1746. https://doi.org/10.1139/cjfas-2014-0169

Johansen, J. L., & Jones, G. P. (2011). Increasing ocean temperature reduces the metabolic performance and

swimming ability of coral reef damselfishes. *Global Change Biology*, *17*(9), 2971–2979.

https://doi.org/10.1111/j.1365-2486.2011.02436.x

Johansson, D., Ruohonen, K., Juell, J.-E., & Oppedal, F. (2009). Swimming depth and thermal history of

individual Atlantic salmon (Salmo salar L.) in production cages under different ambient temperature

conditions. *Aquaculture*, *290*(3–4), 296–303. https://doi.org/10.1016/j.aquaculture.2009.02.022

Jørgensen, L. B., Malte, H., Ørsted, M., Klahn, N. A., & Overgaard, J. (2021). A unifying model to estimate

thermal tolerance limits in ectotherms across static, dynamic and fluctuating exposures to thermal

stress. *Scientific Reports*, *11*(1), 12840. https://doi.org/10.1038/s41598-021-92004-6

Jørgensen, L. B., Ørsted, M., Malte, H., Wang, T., & Overgaard, J. (2022). Extreme escalation of heat failure rates in ectotherms with global warming. *Nature*, *611*(7934), 93–98. https://doi.org/10.1038/s41586-022-05334-4

Jutfelt, F. (2020). Metabolic adaptation to warm water in fish. *Functional Ecology*, *34*(6), 1138–1141. https://doi.org/10.1111/1365-2435.13558

Jutfelt, F., Norin, T., Åsheim, E. R., Rowsey, L. E., Andreassen, A. H., Morgan, R., Clark, T. D., & Speers-Roesch, B. (2021). 'Aerobic scope protection' reduces ectotherm growth under warming. *Functional Ecology*, *35*(7), 1397–1407. https://doi.org/10.1111/1365-2435.13811

Khan, J. R., Pether, S., Bruce, M., Walker, S. P., & Herbert, N. A. (2014). Optimum temperatures for growth and feed conversion in cultured hapuku (*Polyprion oxygeneios*)—Is there a link to aerobic metabolic scope and final temperature preference? *Aquaculture*, *430*, 107–113. https://doi.org/10.1016/j.aquaculture.2014.03.046

Killen, S. S., Christensen, E. A. F., Cortese, D., Závorka, L., Norin, T., Cotgrove, L., Crespel, A., Munson, A., Nati, J. J. H., Papatheodoulou, M., & McKenzie, D. J. (2021). Guidelines for reporting methods to estimate metabolic rates by aquatic intermittent-flow respirometry. *Journal of Experimental Biology*, *224*(18). https://doi.org/10.1242/jeb.242522

Killen, S. S., Glazier, D. S., Rezende, E. L., Clark, T. D., Atkinson, D., Willener, A. S. T., & Halsey, L. G. (2016). Ecological Influences and Morphological Correlates of Resting and Maximal Metabolic Rates across Teleost Fish Species. *The American Naturalist*, *187*(5), 592–606. https://doi.org/10.1086/685893

Koskela, J., Pirhonen, J., & Jobling, M. (1997). Feed intake, growth rate and body composition of juvenile Baltic salmon exposed to different constant temperatures. *Aquaculture International*, *5*, 351–360.

Koskela, J., Pirhonen, J., & Jobling, M. (2005). Variations in feed intake and growth of Baltic salmon and brown trout exposed to continuous light at constant low temperatures. *Journal of Fish Biology*, *50*(4), 837–845. https://doi.org/10.1111/j.1095-8649.1997.tb01976.x

Kullgren, A., Jutfelt, F., Fontanillas, R., Sundell, K., Samuelsson, L., Wiklander, K., Kling, P., Koppe, W., Larsson,

D. G. J., Björnsson, B. T., & Jönsson, E. (2013). The impact of temperature on the metabolome and

endocrine metabolic signals in Atlantic salmon (Salmo salar). *Comparative Biochemistry and Physiology*

Part A: Molecular & Integrative Physiology, *164*(1), 44–53. https://doi.org/10.1016/j.cbpa.2012.10.005

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). **lmerTest** Package: Tests in Linear Mixed Effects

Models. *Journal of Statistical Software*, *82*(13). https://doi.org/10.18637/jss.v082.i13

Lacroix, G. L. (2013). Population-specific ranges of oceanic migration for adult Atlantic salmon (Salmo salar)

documented using pop-up satellite archival tags. *Canadian Journal of Fisheries and Aquatic Sciences*,

70(7), 1011–1030. https://doi.org/10.1139/cjfas-2013-0038

Lefevre, S., Wang, T., & McKenzie, D. J. (2021). The role of mechanistic physiology in investigating impacts of

global warming on fishes. *Journal of Experimental Biology*, *224*. https://doi.org/10.1242/jeb.238840

Lenth, R. V. (2017). *emmeans: Estimated Marginal Means, aka Least-Squares Means* (p. 1.11.0) [Dataset].

https://doi.org/10.32614/CRAN.package.emmeans

Little, A. G., Loughland, I., & Seebacher, F. (2020). What do warming waters mean for fish physiology and

fisheries? *Journal of Fish Biology*, *97*(2), 328–340. https://doi.org/10.1111/jfb.14402

Little, A. G., & Seebacher, F. (2021). Physiological Performance Curves: When Are They Useful? *Frontiers in*

Physiology, *12*, 805102. https://doi.org/10.3389/fphys.2021.805102

Loughland, I., & Seebacher, F. (2020). Differences in oxidative status explain variation in thermal acclimation

capacity between individual mosquitofish (Gambusia holbrooki). *Functional Ecology*, *34*(7), 1380–1390.

https://doi.org/10.1111/1365-2435.13563

MacNutt, M. J., Hinch, S. G., Lee, C. G., Phibbs, J. R., Lotto, A. G., Healey, M. C., & Farrell, A. P. (2006).

Temperature effects on swimming performance, energetics, and aerobic capacities of mature adult

pink salmon (Oncorhynchus gorbuscha) compared with those of sockeye salmon (Oncorhynchus

nerka). *Canadian Journal of Zoology*, *84*(1), 88–97. https://doi.org/10.1139/z05-181

Madaro, A., Folkedal, O., Maiolo, S., Alvanopoulou, M., & Olsen, R. E. (2018). Effects of acclimation

temperature on cortisol and oxygen consumption in Atlantic salmon (*Salmo salar*) post-smolt exposed

to acute stress. *Aquaculture*, *497*, 331–335. https://doi.org/10.1016/j.aquaculture.2018.07.056

Martins, E. G., Hinch, S. G., Patterson, D. A., Hague, M. J., Cooke, S. J., Miller, K. M., Lapointe, M. F., English, K.

K., & Farrell, A. P. (2011). Effects of river temperature and climate warming on stock-specific survival of

adult migrating Fraser River sockeye salmon (Oncorhynchus nerka). *Global Change Biology*, *17*(1), 99–

114. https://doi.org/10.1111/j.1365-2486.2010.02241.x

Martins, E. G., Hinch, S. G., Patterson, D. A., Hague, M. J., Cooke, S. J., Miller, K. M., Robichaud, D., English, K. K.,

& Farrell, A. P. (2012). High river temperature reduces survival of sockeye salmon (Oncorhynchus

nerka) approaching spawning grounds and exacerbates female mortality. *Canadian Journal of Fisheries*

and Aquatic Sciences, *69*(2), 330–342. https://doi.org/10.1139/f2011-154

McArley, T. J., Hickey, A. J. R., & Herbert, N. A. (2017). Chronic warm exposure impairs growth performance and

reduces thermal safety margins in the common triplefin fish (Forsterygion lapillum). *Journal of*

Experimental Biology, *220*(19), 3527–3535. https://doi.org/10.1242/jeb.162099

McKenzie, D. J., & Claireaux, G. (2010). The Effects of Environmental Factors on the Physiology of Aerobic

Exercise. In *Fish Locomotion*. CRC Press.

McKenzie, D. J., Zhang, Y., Eliason, E. J., Schulte, P. M., Claireaux, G., Blasco, F. R., Nati, J. J. H., & Farrell, A. P.

(2021). Intraspecific variation in tolerance of warming in fishes. *Journal of Fish Biology*, *98*(6), 1536–

1555. https://doi.org/10.1111/jfb.14620

Morash, A. J., Neufeld, C., MacCormack, T. J., & Currie, S. (2018). The importance of incorporating natural

thermal variation when evaluating physiological performance in wild species. *Journal of Experimental*

Biology, *221*(14), jeb164673. https://doi.org/10.1242/jeb.164673

Morash, A. J., Speers-Roesch, B., Andrew, S., & Currie, S. (2021). The physiological ups and downs of thermal

variability in temperate freshwater ecosystems. *Journal of Fish Biology*, *98*(6), 1524–1535.

https://doi.org/10.1111/jfb.14655

Morgan, R., Andreassen, A. H., Åsheim, E. R., Finnøen, M. H., Dresler, G., Brembu, T., Loh, A., Miest, J. J., &

Jutfelt, F. (2022). Reduced physiological plasticity in a fish adapted to stable temperatures. *Proceedings*

of the National Academy of Sciences, *119*(22), e2201919119.

https://doi.org/10.1073/pnas.2201919119

Morissette, J., Swart, S., MacCormack, T. J., Currie, S., & Morash, A. J. (2021). Thermal variation near the

thermal optimum does not affect the growth, metabolism or swimming performance in wild Atlantic

salmon Salmo salar. *Journal of Fish Biology*, *98*(6), 1585–1589. https://doi.org/10.1111/jfb.14348

Morozov, S. (2024). *PumpResp* [Computer software]. https://github.com/embedded-sergey/PumpResp

Morozov, S., McCairns, R. J. S., & Merilä, J. (2019). FishResp: R package and GUI application for analysis of

aquatic respirometry data. *Conservation Physiology*, *7*(1), coz003.

https://doi.org/10.1093/conphys/coz003

Muñoz, N. J., Reid, B., Correa, C., Neff, B. D., & Reynolds, J. D. (2021). Non-native Chinook salmon add nutrient

subsidies and functional novelty to Patagonian streams. *Freshwater Biology*, *66*(3), 495–508.

https://doi.org/10.1111/fwb.13655

Navarro, J. M., Paschke, K., Ortiz, A., Vargas-Chacoff, L., Pardo, L. M., & Valdivia, N. (2019). The Antarctic fish

Harpagifer antarcticus under current temperatures and salinities and future scenarios of climate

change. *Progress in Oceanography*, *174*, 37–43. https://doi.org/10.1016/j.pocean.2018.09.001

Norin, T., & Clark, T. D. (2016). Measurement and relevance of maximum metabolic rate. *Journal of Fish*

Biology, *88*, 122–151. https://doi.org/10.1111/jfb.12796

Norin, T., Malte, H., & Clark, T. D. (2014). Aerobic scope does not predict the performance of a tropical

eurythermal fish at elevated temperatures. *Journal of Experimental Biology*, *217*(2), 244–251.

https://doi.org/10.1242/jeb.089755

Norin, T., & Metcalfe, N. B. (2019). *Ecological and evolutionary consequences of metabolic rate plasticity in*

response to environmental change. 1–9.

Nuic, B., Bowden, A., Franklin, C. E., & Cramp, R. L. (2024). Atlantic salmon Salmo salar do not prioritize

digestion when energetic budgets are constrained by warming and hypoxia. *Journal of Fish Biology*,

104(6), 1718–1731. https://doi.org/10.1111/jfb.15693

Ørsted, M., Jørgensen, L. B., & Overgaard, J. (2022). Finding the right thermal limit: A framework to reconcile

ecological, physiological and methodological aspects of CTmax in ectotherms. *Journal of Experimental*

Biology, *225*(19), jeb244514. https://doi.org/10.1242/jeb.244514

O'Sullivan, A. M., Corey, E. M., Collet, E. N., Helminen, J., Curry, R. A., MacIntyre, C., & Linnansaari, T. (2023).

Timing and frequency of high temperature events bend the onset of behavioural thermoregulation in

Atlantic salmon (Salmo salar). *Conservation Physiology*, *11*(1), coac079.

https://doi.org/10.1093/conphys/coac079

Patterson, D. A., Cooke, S. J., Hinch, S. G., Robinson, K. A., Young, N., Farrell, A. P., & Miller, K. M. (2016). A

perspective on physiological studies supporting the provision of scientific advice for the management

of Fraser River sockeye salmon (*Oncorhynchus nerka*). *Conservation Physiology*, *4*(1), cow026.

https://doi.org/10.1093/conphys/cow026

Peralta-Maraver, I., & Rezende, E. L. (2021). Heat tolerance in ectotherms scales predictably with body size.

Nature Climate Change, *11*(1), 58–63. https://doi.org/10.1038/s41558-020-00938-y

Perry, A. L., Low, P. J., Ellis, J. R., & Reynolds, J. D. (2005). Climate Change and Distribution Shifts in Marine

Fishes. *Science*, *308*(5730), 1912–1915. https://doi.org/10.1126/science.1111322

Pinheiro, J., Bates, D., & R Core Team. (1999). *nlme: Linear and Nonlinear Mixed Effects Models* (p. 3.1-168)

[Dataset]. https://doi.org/10.32614/CRAN.package.nlme

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater vulnerability to

warming of marine versus terrestrial ectotherms. *Nature*, *569*(7754), 108–111.

https://doi.org/10.1038/s41586-019-1132-4

Pörtner, H. O., & Farrell, A. P. (2008). Ecology: Physiology and climate change. *Science*, *322*(5902), 690–692.

https://doi.org/10.1126/science.1163156

Pörtner, H.-O. (2010). Oxygen- and capacity-limitation of thermal tolerance: A matrix for integrating climate-

related stressor effects in marine ecosystems. *Journal of Experimental Biology*, *213*(6), 881–893.

https://doi.org/10.1242/jeb.037523

Priede, I. G. (1985). Metabolic Scope in Fishes. In P. Tytler & P. Calow (Eds.), *Fish Energetics* (pp. 33–64).

Springer Netherlands. https://doi.org/10.1007/978-94-011-7918-8_2

Prokkola, J. M., Åsheim, E. R., Morozov, S., Bangura, P., Erkinaro, J., Ruokolainen, A., Primmer, C. R., & Aykanat,

T. (2022). Genetic coupling of life-history and aerobic performance in Atlantic salmon. *Proceedings of*

the Royal Society B: Biological Sciences, *289*(1967), 20212500. https://doi.org/10.1098/rspb.2021.2500

Raby, G. D., Casselman, M. T., Cooke, S. J., Hinch, S. G., Farrell, A. P., & Clark, T. D. (2016). Aerobic scope

increases throughout an ecologically relevant temperature range in coho salmon. *Journal of*

Experimental Biology, jeb.137166. https://doi.org/10.1242/jeb.137166

Raby, G. D., Doherty, C. L. J., Mokdad, A., Pitcher, T. E., & Fisk, A. T. (2020). Post-exercise respirometry

underestimates maximum metabolic rate in juvenile salmon. *Conservation Physiology*, *8*(1), coaa063.

https://doi.org/10.1093/conphys/coaa063

Reddin, D. G. (1985). Atlantic Salmon (*Salmo salar*) On and East of the Grand Bank. *Journal of Northwest*

Atlantic Fishery Science, *6*, 157–164. https://doi.org/10.2960/J.v6.a16

Réveillon, T., Rota, T., Chauvet, É., Lecerf, A., & Sentis, A. (2019). Repeatable inter-individual variation in the

thermal sensitivity of metabolic rate. *Oikos*, *128*(11), 1633–1640. https://doi.org/10.1111/oik.06392

Sadoul, B., & Vijayan, M. M. (2016). Stress and Growth. In C. B. Schreck, L. Tort, A. P. Farrell, & C. J. Brauner

(Eds.), *Fish Physiology* (Vol. 35, pp. 167–205). Academic Press. https://doi.org/10.1016/B978-0-12-

802728-8.00005-9

Sandblom, E., Clark, T. D., Gräns, A., Ekström, A., Brijs, J., Sundström, L. F., Odelström, A., Adill, A., Aho, T., &

Jutfelt, F. (2016). Physiological constraints to climate warming in fish follow principles of plastic floors

and concrete ceilings. *Nature Communications*, *7*(1), 11447. https://doi.org/10.1038/ncomms11447

Schulte, P. M. (2015). The effects of temperature on aerobic metabolism: Towards a mechanistic

understanding of the responses of ectotherms to a changing environment. *Journal of Experimental*

Biology, *218*(12), 1856–1866. https://doi.org/10.1242/jeb.118851

Schulte, P. M., & Healy, T. M. (2022). Chapter 9—Physiological diversity and its importance for fish

conservation and management in the Anthropocene. In S. J. Cooke, N. A. Fangue, A. P. Farrell, C. J.

Brauner, & E. J. Eliason (Eds.), *Fish Physiology* (Vol. 39, pp. 435–477). Academic Press.

https://doi.org/10.1016/bs.fp.2022.04.009

Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal Performance Curves, Phenotypic Plasticity, and the

Time Scales of Temperature Exposure. *Integrative and Comparative Biology*, *51*(5), 691–702.

https://doi.org/10.1093/icb/icr097

Seebacher, F. (2009). Responses to temperature variation: Integration of thermoregulation and metabolism in

vertebrates. *Journal of Experimental Biology*, *212*(18), 2885–2891. https://doi.org/10.1242/jeb.024430

Seebacher, F., & Little, A. G. (2021). Plasticity of Performance Curves in Ectotherms: Individual Variation

Modulates Population Responses to Environmental Change. *Frontiers in Physiology*, *12*.

https://doi.org/10.3389/fphys.2021.733305

Simpson, A. L. (1992). Differences in body size and lipid reserves between maturing and nonmaturing Atlantic

salmon parr, Salmo salar L. *Canadian Journal of Zoology*, *70*(9), 1737–1742.

https://doi.org/10.1139/z92-241

Slesinger, E., Andres, A., Young, R., Seibel, B., Saba, V., Phelan, B., Rosendale, J., Wieczorek, D., & Saba, G.

(2019). The effect of ocean warming on black sea bass (Centropristis striata) aerobic scope and hypoxia

tolerance. *PLOS ONE*, *14*(6), e0218390. https://doi.org/10.1371/journal.pone.0218390

Strøm, J. F., Rikardsen, A. H., Campana, S. E., Righton, D., Carr, J., Aarestrup, K., Stokesbury, M. J. W., Gargan,

P., Javierre, P. C., & Thorstad, E. B. (2019). Ocean predation and mortality of adult Atlantic salmon.

Scientific Reports, *9*(1), 7890. https://doi.org/10.1038/s41598-019-44041-5

Svendsen, M. B. S., Bushnell, P. G., & Steffensen, J. F. (2016). Design and setup of intermittent-flow

respirometry system for aquatic organisms. *Journal of Fish Biology*, *88*(1), 26–50.

https://doi.org/10.1111/jfb.12797

Thorpe, J. E. (2007). Maturation responses of salmonids to changing developmental opportunities. *Marine

Ecology Progress Series*, *335*, 285–288. https://doi.org/10.3354/meps335285

Van Leeuwen, T. E., Lehnert ,Sarah J., Breau ,Cindy, Fitzsimmons ,Michelle, Kelly ,Nicholas I., Dempson ,J. Brian,

Neville ,Victoria M., Young ,Mark, Keefe ,Donald, Bird ,Tomas J., & and Cote, D. (2023). Considerations

for Water Temperature-Related Fishery Closures in Recreational Atlantic Salmon (Salmo salar) Catch

and Release Fisheries: A Case Study from Eastern Canada. *Reviews in Fisheries Science & Aquaculture*,

31(4), 598–619. https://doi.org/10.1080/23308249.2023.2242959

Verberk, W. C. E. P., Sandker, J. F., Van De Pol, I. L. E., Urbina, M. A., Wilson, R. W., McKenzie, D. J., & Leiva, F. P.

(2022). Body mass and cell size shape the tolerance of fishes to low oxygen in a temperature-

dependent manner. *Global Change Biology*, *28*(19), 5695–5707. https://doi.org/10.1111/gcb.16319

Wade, N. M., Clark, T. D., Maynard, B. T., Atherton, S., Wilkinson, R. J., Smullen, R. P., & Taylor, R. S. (2019).

Effects of an unprecedented summer heatwave on the growth performance, flesh colour and plasma

biochemistry of marine cage-farmed Atlantic salmon (*Salmo salar*). *Journal of Thermal Biology*, *80*, 64–

74. https://doi.org/10.1016/j.jtherbio.2018.12.021

Ward, H. G. M., Post, J. R., Lester, N. P., Askey, P. J., & Godin, T. (2017). Empirical evidence of plasticity in life-

history characteristics across climatic and fish density gradients. *Canadian Journal of Fisheries and*

Aquatic Sciences, *74*(4), 464–474. https://doi.org/10.1139/cjfas-2016-0023

Závorka, L., Koeck, B., Armstrong, T. A., Soğanci, M., Crespel, A., & Killen, S. S. (2020). Reduced exploration

capacity despite brain volume increase in warm acclimated common minnow. *The Journal of*

Experimental Biology, *i*(May), jeb.223453. https://doi.org/10.1242/jeb.223453

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and extensions in*

ecology with R. Springer. https://doi.org/10.1007/978-0-387-87458-6

Figure 1: Outline of the experiment temperature regime. Solid lines and points represent temperatures (mean of 16, 19 and 22 °C) of the heatwave tanks (three tanks with independent temperature regulation) and control tanks (three tanks with an identical temperature, 16 °C) during the experiment. Lines fitted using a 40-point rolling average, where each point is one hour. Dashed vertical lines (B1 – B8) indicate timing of metabolic rate measurements: two batches measured each week. Fish were removed from heatwave tanks into H-Tronic controlled acclimation tank 24 hours before testing. Variation in temperature profiles at each step is related to natural mixing in the incoming lake water.

Figure 2: Scatter plots to show the relationship of log transformed metabolic rates with log transformed mass (A: Standard metabolic rate (SMR), B: Maximum Metabolic rate (MMR), C: Aerobic Scope (AS)). Each point represents individual fish, with different colours representing different tested temperatures (16 °C: Blue, 19 °C: Purple, 22 °C: Pink). Goodness of fit lines indicate linear regression and shading shows 95% confidence intervals for each temperature, with orange colour representing fit of all data.

821 **Figure 3: Boxplots to show the relationship between temperature and mass-adjusted**
822 **metabolic rate measurements: SMR (a), MMR (b) and AS (c).** Different point shapes
823 indicate different weeks of measurement and different colours indicate different
824 temperatures. Asterix and brackets show significant differences (* $p < 0.05$ and ** $p < 0.01$).

Table 1: Descriptive statistics for Mass (g) and Length (mm) for juvenile salmon tested at different temperatures. n = number of individuals, $\bar{x}$ = mean metabolic rate, σ = standard deviation, Min = minimum value, Max = maximum value.

Temperature (°C)	n	Mass (g)				Length (mm)			
		$\bar{x}$	σ	Min	Max	$\bar{x}$	σ	Min	Max
16 (week 1)	22	1.42	0.41	1.01	2.59	57.64	5.38	50	69
19	19	1.35	0.35	0.70	1.88	56.84	4.71	47	64
22	16	1.79	0.42	0.88	2.40	61.00	4.35	54	68
16 (week 4)	15	2.56	0.88	1.40	4.32	68.53	8.22	56	82
Total	72	1.72	0.70	0.70	4.32	60.44	7.16	47	82

Table 2: Model summaries for metabolic rate estimates, including fish body mass and temperature (16, 19 and 22 °C) as independent variables. Batch is included as random effect. Brackets indicate standard error, *asterisks* indicates significance values with *p<0.05; **p<0.01; ***p<0.001).

	log10(SMR)	log10(MMR)	log10(AS)
log10(mass)	0.922***	0.978***	1.044***
	(0.057)	(0.120)	(0.229)
19 °C	0.131***	0.011	-0.038
	(0.027)	(0.046)	(0.086)
22 °C	0.198***	0.040	-0.033
	(0.027)	(0.046)	(0.086)
Intercept	-0.704***	-0.152***	-0.342***
	(0.021)	(0.038)	(0.072)
Random Effects			
Intercept SD (Batch)	0.0215	0.0116	0
Residual SD	0.0638	0.1492	0.2878
Marginal R^2 / Conditional R^2	0.849 / 0.864	0.521 / 0.524	0.261 / NA

839 **Table 3: Post-hoc ANOVA contrasts from the results of the SMR model described in Table**
840 **2, comparing the temperature treatments 16 °C, 19 °C, and 22 °C.** SE shows standard
841 error, df degrees of freedom and the associated *t* ratio and *p* (significance) with
842 *p<0.05; **p<0.01; ***p<0.001.

Comparison	Estimate	SE	df	t	p
16 °C - 19 °C	-0.13	0.03	4.66	-4.71	0.014*
16 °C - 22 °C	-0.20	0.03	4.73	-7.42	0.002***
19 °C - 22 °C	-0.07	0.03	4.94	-2.13	0.178

844 Table 4: Table showing predicted SMR (mg O_2 h^{-1}) and 95% confidence intervals (CI) for a
845 mean mass fish (1.72g) at different temperatures.

Temperature (°C)	Predicted SMR (for a 1.72g fish)	Lower CI	Upper CI
16	0.33	0.24	0.44
19	0.44	0.32	0.58
22	0.52	0.39	0.70

846