Aerobic scope is sustained through a heatwave in juvenile Atlantic salmon

Lucy Cotgrove[1], Sergey Morozov[2], Miika Raitakivi[3], Evan Sala[3], Jenni M. Prokkola[1]

1 Natural Resources Institute Finland (Luke), Paavo Havaksen tie 3, 90570 Oulu, Finland

2 Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental

Sciences, University of Helsinki, Viikinkaari 9, 00014 Helsinki, Finland

3 Natural Resources Institute Finland (Luke), Vilppulantie 415, 41360 Laukaa, Finland

Corresponding author: lucy.cotgrove@luke.fi

Total Word Count

9642, including references

Lay Summary

In juvenile Atlantic salmon, standard metabolic rate increases but there is no difference in maximum

metabolic rate and aerobic scope with increasing temperatures. Considering the increased metabolic costs of

activities at higher temperatures, our results show that juvenile salmon are vulnerable to deteriorating

performance at temperatures commonly experienced in present-day heatwaves.

Abstract

Aquatic ectotherms are vulnerable to heatwave-induced physiological stress, which arises from increased energy demands and reduced dissolved oxygen content in warmer waters. Understanding thermal physiology is critical for predicting how commercially and ecologically important populations could be affected by the increasing risk of rising temperatures. Heatwave risk assessments often examine extremities of time scales: immediate impacts or long-term consequences. However, little is known about how consistently increasing mid-term thermal stress shapes aerobic performance in commercially important species such as Atlantic salmon (*Salmo salar*), which may face heat stress in rivers, especially at the juvenile life stage. By measuring how salmon juveniles manage their aerobic capacity at 16, 19 and 22°C using intermittent respirometry, we test if their thermal performance curve shows a decline at temperatures commonly occurring during heatwaves. Whole-animal metabolism was measured from control individuals kept at 16 °C before and after the heatwave, and after 4-5 days exposure at 19 and 22°C during the heatwave. We show that standard metabolic rate increases with temperature, but maximum metabolic rate and aerobic scope do not change between these temperatures. These findings suggest that juvenile Atlantic salmon may have limited capacity to increase aerobic performance during moderate heatwaves, leaving them vulnerable to cumulative effects of oxygen limitation to vital functions such as growth and stress responses. As climate change intensifies, incorporating thermal performance curves into conservation strategies can be utilized for predicting population resilience and informing effective management.

Keywords

Respirometry, thermal performance, ecophysiology, energetics

Introduction

With climate change increasing the severity and frequency of heatwaves (IPCC, 2023), organisms may experience physiological thermal stress and even mortality due to extreme temperature fluctuations. Understanding organisms' thermal physiology is therefore critically important for conservation (Clark et al., 2013; Deutsch et al., 2008). Heatwaves frequently surpass the temperature range that species have been adapted to, often cumulatively and repeatedly, imposing selection on thermal performance. This is especially true for ectotherms, whose internal temperature are regulated by the surrounding environment. Some species may demonstrate adaptive or resilient responses to heatwaves including behavioural adjustments, physiological acclimation, or shifts in geographical ranges (Breedveld et al., 2023; Burraco et al., 2020; Jørgensen et al., 2022). However, these responses may not always be sufficient to counteract the negative consequences of extreme weather fluctuation (Perry et al., 2005; Pörtner & Farrell, 2008). Most physiological processes in ectotherms, including development, reproduction, and metabolism, are temperature-dependent (Angilletta et al., 2010; Hochachka & Somero, 1973; Schulte, 2015; Seebacher, 2009). These processes can be described using thermal performance curves (TPCs), which illustrate how a biological rate, such as oxygen consumption or activity per unit time, changes with temperature (Schulte et al., 2011).

TPCs can be estimated using thermal windows, critical temperatures, and preferred temperature ranges (Jørgensen et al., 2021; Ørsted et al., 2022; Peralta-Maraver & Rezende, 2021). TPCs can be used to compare inter- and intraspecies responses to anticipated temperatures under a future climate and how this might vary across life stages (Lefevre et al., 2021; Seebacher & Little, 2021). In water-breathing aquatic ectotherms, such as fishes, TPCs linked to demand, uptake and delivery of oxygen are particularly important because of the decreasing solubility but increasing oxygen consumption in warmer water (Little et al., 2020; Little & Seebacher, 2021; Seebacher & Little, 2021; Verberk et al., 2022). Consequently, TPCs related to metabolic rates and aerobic performance are pivotal to understanding the effects of heatwaves on fishes.

Metabolic rate is a fundamental aspect of fish physiology, reflecting the energy expenditure associated with basic life processes such as respiration, digestion, and movement (Brett, 1964; Fry et al., 1947). Standard metabolic rate (SMR) of an animal represents the baseline rate of aerobic metabolism required to sustain life in a post-absorptive, resting state, while the maximum capacity for aerobic performance is set by the maximum metabolic rate (MMR) (Sandblom et al., 2016). The difference between SMR and MMR represents the aerobic scope (AS) of an animal and theoretically determines the capacity of aerobic metabolism to support key life-history attributes such as activity, growth and reproduction, each of which has a specific oxygen cost (Fry, 1971; Fry et al., 1947). Therefore, reductions in aerobic scope can impair physical performance (Johansen & Jones, 2011; Pörtner & Farrell, 2008; Priede, 1985). Given that the SMR of ectotherms rises with temperature and often no such increase is detected in MMR (Fry et al., 1947; Norin et al., 2014; Sandblom et al., 2016), increases in temperature beyond optimum can result in a decline in AS (Farrell, 2016; Fry, 1971; Lefevre et al., 2021). However, maintaining growth and other aerobic functions with increasing temperature requires an increasing AS, because of the higher costs of activities such as digestion and assimilation and swimming (Jahn & Seebacher, 2022; Jutfelt et al., 2021). When AS declines at increased temperatures, this may cause declining growth, health, or survival of individuals (Alfonso et al., 2021; Jutfelt et al., 2021; Navarro et al., 2019; Sadoul & Vijayan, 2016). Alternatively, AS can plateau at higher temperatures suggesting that oxygen supply can limit performance as temperature increases (hvas et al 2017, Závorka et al., 2020, Christensen et al 2012). As AS plateaus but SMR and the costs of aerobic activities continue to increase, fish might become more constrained in their physiological state and experience cumulative thermal stress, leading to mortality. Both declines and plateaus of AS pose significant risks for salmonids, which include economically important, cold-water adapted species. In these fish, limitations of AS has already been linked to reduced growth, survival, or collapse of aerobic capability during heatwaves (Hvas 2017 (Eliason, Wilson, et al., 2013; Pörtner & Farrell, 2008; Wade et al., 2019).

The Atlantic salmon (*Salmo salar* L.) is faced with supra-optimal temperatures especially at riverine life stages, including juveniles (parr) and adults during their spawning migration, with reports of present-day river temperatures up to 28 °C (O'Sullivan et al., 2023; Strøm et al., 2019). Juvenile Atlantic salmon rear for one to five years in freshwater before migrating to the sea (Aas et al., 2010), with potential to encounter multiple heatwaves. These can have direct effects on survival, but also cascading effects on population dynamics through reduced growth rates. Faster freshwater growth in salmon is linked to higher survival and faster maturation at sea (Hutchings & Jones, 1998; Simpson, 1992; Thorpe, 2007). In some populations, a negative association between freshwater growth and river temperature has been detected (Alioravainen et al., 2023). For the majority of their life cycle at sea, Atlantic salmon will experience temperatures below 8°C (Jensen et al., 2014; Lacroix, 2013; Reddin, 1985). Atlantic salmon tend to avoid temperatures above 15°C (Fisher & Elson, 1950; Johansson et al., 2009; Lacroix, 2013), and their optimal temperature range for feeding and normal behaviour in the wild has been suggested to be 6–20°C, with peak growth rates occurring at 16–17°C in aquaculture conditions (Dwyer & Piper, 1987; Elliott, 1982). Previous work assessing thermal performance in Atlantic salmon has often focussed on upper limits during acute exposure of CTmax experiments under less than 24 hr(Desforges et al., 2023 and papers within), or long-term stable temperature increases over several months (Anttila et al., 2014; Del Rio et al., 2021; Hvas et al., 2017). When aerobic scope has been measured in a heatwave-context in a related species, (Casselman et al., 2012) found AS peaked at 17 °C, and decreased until testing finished at 21 °C in juvenile Coho salmon *(Oncorhynchus kisutch*). In post-smolt Atlantic salmon, AS tended to increase (a non-significant effect) as temperature increased from 13 °C to 23 °C, but swimming ability and feeding rate were greatly decreased and mortality increased, when metabolic rates of salmon were tested in groups without a cumulative exposure (Hvas et al., 2017). Further, thermal variability in acclimation regime had no effect on salmon AS compared to a stable acclimation when metabolic rates were tested acutely at the same temperature (Morissette et al., 2021). Still, information on the cumulative effects of thermal fluctuations over several days or weeks, which better reflect natural heatwave patterns, are limited (Morash et al., 2021; Nuic et al., 2024). This information is pivotal for the conservation of Atlantic salmon. The

diminishing of salmon populations has already reduced opportunities for commercial and recreational fishery,

cultural traditions, and increased the risk of local extinctions (ICES, 2024, Dadswell et al., 2022)

Here, we test how mid-term increasing temperatures that are prevalent in rivers during heatwaves impact

metabolic rates and AS in hatchery-reared juvenile Atlantic salmon. We hypothesise that SMR will increase as

temperature rises. We expect MMR to show little variation between temperatures, and therefore a decrease

in AS as temperature increases. These findings will provide insight into the physiological constraints juvenile

salmon may face during heatwaves, using an ecologically relevant temperature regime.

Methods

Fish Husbandry

The experiment was performed with a permit granted by the Finnish Project Authorisation Board (no.

ESAVI/16748/2023). Families of Atlantic salmon were bred using a hatchery broodstock originating from

River Iijoki, maintained by Natural Resources Institute of Finland (Luke), in Taivalkoski, Finland. The broodstock

was established in 2012 from parents that were originally stocked into the river as juveniles/smolts, and

therefore had completed half of their life cycle in the wild. Individuals from the broodstock were crossed to

make full-sib families of offspring in October 2022. The embryos were incubated in Taivalkoski hatchery in

family-specific flow-through trays at natural temperatures of incoming water from River Ohtaoja, and dead

embryos were removed regularly.

Embryos were disinfected by iodine bath (Buffodine) on 9th March 2023, and transported to a LUKE hatchery

in Laukaa, Finland, where experiments were conducted. Survival between transportation and start of the

experiment was 22%. Excess mortality may have been caused by the closeness to the embryo's hatching date

and transporting to the fish farm (no disease was observed in the offspring). Embryos from three unrelated

families were incubated and reared in separate circular tanks (diameter 80 cm) and supplied with a continuous

flow of filtered water from a mix of local lake Peurunkajärvi and river Peurunkajoki. Feeding of alevins was started with powdered feed (BioMar Group) when most of the egg yolk had been consumed at approximately 25 May, 2023. Tanks were cleaned by scrubbing surfaces and siphoning excess food twice a week until 10 June, 2023, and then daily until the end of the experiment. Feed rations were calculated from growth predictions assuming feed conversion efficiency of 0.8 (Elliott & Hurley, 1997), and adjusted such that a small amount of feed was left uneaten. Fish were fed using 12h belt feeders with the daily ration, except for one day preceding metabolic measurements, when fish were fasted. During the summer and the experiment, fish were reared under constant light. Rearing temperature raised gradually with natural water temperatures over approximately three months until the experiment was started; temperature varied from 9.2 to 15.7 °C between first feeding and experimental period.

At the start of the experiment 27 July, 2023, families were divided into two groups, control and heatwave, with approximately same number of individuals in each group within a family (N = 24 to 62 within family /tank). Fish were kept in three tanks each for control and heatwave groups. Densities within families were kept similar between control and heatwave tanks throughout the experiment by simultaneously removing fish from control tanks to match densities in of heatwave tanks. Control tanks were kept at close to 16 °C (mean = 16.1, SD = 0.5) throughout the experimental period by regulating the incoming water flows from the river and the lake with a natural temperature difference. Temperatures in one control tank (the same water source used in all three tanks) and the three heatwave tanks were measured every 30 min with Hobo 64K Pendant temperature loggers (Hobo, Bourne, MA, USA).

Temperatures of the heatwave tanks were regulated via a heating tank to offset the colder temperatures of the room and incoming water using a temperature controller (TS1000, H-Tronic GmbH, Germany) with a temperature probe PT-1000 located in each tank, and an Eheim 600 or 300 pump (Eheim, Deizisau, Germany) depending on the distance from the heating tank. The pumps circulated water through stainless steel coils in the heating tank, in which temperature fluctuated between 30-35 °C. Heatwave tank water was aerated

constantly to prevent declines in dissolved O_2 related to increasing temperatures, as the flow-through rate of

fresh water to the tanks was reduced (approx. 1 L min^{-1}) during the heatwave to maintain temperatures.

Experimental Heatwave

Fish in the heatwave treatment were exposed to three test temperatures: 16 °C, 19 °C, and 22 °C (Figure 1).

First, 16 °C was maintained as a control temperature for eight days, after which 3-8 individuals from each

family were randomly picked for measurements of metabolic rate (similar sample size across temperatures

within a family), and moved into a fasting tank, where they were unfed until the measurements started the

following day. Temperature of the fasting tank was the same as the heatwave tanks. After the measurement

of MMR and SMR at 16 °C, temperature was increased by one degree per day for three days, and maintained

for 4-5 days, after which metabolic rate measurements were taken from a new set of fish at 19 °C. This was

repeated and metabolic rate measurements were taken at 22°C. Due to a malfunctioning temperature

controller at 19 °C (after moving fish to acclimation for B3 and B4), there was an increase in temperature in

one of the heatwave tanks to 21.5 °C (Figure 1). Metabolic rate experiments were conducted once per

individual, and fish were euthanised after the SMR measurement.

To measure MMR, fish were transferred from the fasting tank into 50 L buckets filled with 5 cm of water at the

testing temperature using plastic cups of water to reduce air exposure. Fish were encouraged to swim by hand

around the circular container (Prokkola et al., 2022; Raby et al., 2020). Chasing lasted 2 min, and all fish were

unresponsive to the touch of caudal fins, thus determined fatigued. The fish were then rapidly transferred to

respirometry chambers using plastic cups with water. Two people chased one fish each at once, and each

batch of max. 16 individuals was processed within 2 h. Fish were left in respirometry chambers overnight in

order to capture the MMR and SMR for each fish, as per (Chabot, McKenzie, et al., 2016). Chambers were

flushed with fresh water from the tank for 5 min every 15 min, allowing fully aerated water to enter the

chamber. The lowest recorded oxygen level in any chamber was 6.7 mg L $^{-1}$ at 22 °C. After SMR

measurements, fish were euthanised using an overdose of MS-222, measured, and weighed.

Immediately before and after metabolic measurements, background respiration was measured in empty

chambers for three measurement cycles (5 min flush, 15 min measurement) and an average of these slopes

was used to adjust the data for MMR and SMR. For those trials which took place immediately after the

previous, the post-trial background data of the previous trial was used as the pre-trial background values. To

prevent bacterial build-up, the system was bleached between temperature changes, and all chamber parts

scrubbed to limit bacterial growth and rinsed thoroughly. Additionally, the water in the respirometry tank was

continuously circulated through a UV filter.

A total of 120 fish were subjected to intermittent flow respirometry after a 24h fasting period to provide

estimates of metabolic rate (SMR, MMR, AS see table S1) (Killen et al., 2021; Svendsen et al., 2016). Due to

death in respirometer chambers, exclusion due to outliers, incomplete data recording, and due to computer

failure, 72 fish were included in the analysis (Table S2).

Respirometer design

For a summary of the respirometry design and measurement protocol, see Supplementary Table 1, after (Killen

et al., 2021). A 16-chamber intermittent flow respirometer, each chamber containing an individual fish, was

submerged in a temperature-regulated water bath (16, 19 & 22 ± 0.1 °C; 200 L) that was saturated with air. The

oxygen content of the water in the chambers was measured every two seconds using a 4-channel fibre optic

oxygen meter with associated oxygen sensors and software (FireStingO2; PyroScience GmbH, Aachen,

Germany). All 16 O_2 sensors were calibrated for 0% oxygen saturation simultaneously with O_2-free water, made

using sodium sulphate, and simultaneous for 100% oxygen saturation with air-saturated water using an air stone

before the first measurements at each temperature. Temperature-compensation for O_2 saturation was based

on PT100 temperature sensors, which were placed in the middle of the respirometer tank and connected to each of the Firesting meters. The chambers were shielded from disturbance and light using an opaque plastic cover during the measurement of SMR. Flush and recirculation pumps were controlled using PumpResp controllers (4-channel model, FishResp, Finland, https://github.com/embedded-sergey/PumpResp, (2024)). The temperature of the respirometer tank was maintained using a temperature controller (as in the heatwave tanks) and a reservoir connected to a TECO 2000 Chiller/heater. The reservoir received constant inflow of water (approx. 1 L min^{-1}) from the same source as the rearing tanks to maintain good water quality. Chambers for the respirometer were made using glass tubing (120 mm length, inner diameter 38 mm, wall thickness 3.2 mm) and plastic caps that were 3D-printed on PA2200 polyamide. The 'HeiBer'-caps were designed by Heidrikur Bergsson, University of Copenhagen (https://zenodo.org/record/4062429#.YMSW7h1RVTY). Volume of chambers and tubing was 131.86 ± 1 mL (the exact volumes per chamber are reported in an online data repository, see Data Availability). Water inside the chambers was mixed by a plastic disk attached to each cap to distribute the flow. The disks were 3D printed using the same method as the respirometry caps and attached with stainless steel screws. Caps were sealed using rubber O-rings and connected to the flush and recirculation pumps and valves with gas-impermeable Tygon tubing (Tygon S3 E-3603, Satint-Gobain, Paris. France). The recirculation system was confirmed to be waterproof by filling with water and plugging the flush pump and probe connections. Water was recirculated using one submersible pump (12V DC, 6W 2 L min^{-1} pump, Qingdao Xinhui Hardware Machinery Co., Ltd., Qingdao) per channel within the recirculation loop. Chambers were flushed using a second pump (same as previous), using an inflow of water from the tank into the chambers, and excess water was flushed through the chamber to a flush pipe which was placed with the end above the water surface. This allowed fully aerated water to enter the chamber. Flow rate through both the flush line and recirculation loop could be controlled by the valves attached within the flow. Flow rate of the respirometers was approximately 0.1 (± 0.008) L min^{-1} without fish. Information about time and pump phase (either flush or measurement) was recorded by the PumpResp controllers and stored on a computer. Flow had no apparent effect on movement of fish within the chambers in SMR or MMR measurements.

Data Analysis

All data and statistical analyses were done in R environment (R Core Team ,2022). The FishResp package (Morozov et al., 2019) was used to filter and calculate metabolic rate estimates. Slopes of oxygen consumption were adjusted for bacterial oxygen consumption using pre-and post-background measurements, assuming a linear change. For SMR, oxygen consumption (mg O_2 h^{-1}) for each measurement phase was derived from the slope of linear regression of dissolved oxygen concentration over time. The SMR slopes were quality filtered and smoothed before analysis to exclude non-linear declines of oxygen: first, slopes were filtered based on $R^2 > 0.95$, then slopes that did not meet the R^2 criteria were smoothed using a running mean of 29 seconds (Chabot et al., 2021), then R^2 filter was re-applied, and finally, linearity was checked visually from all slopes that met the criteria of $R^2 > 0.95$. The first and last 60 seconds of measurement periods were excluded to account for mixing of water and changes in flow due to flushing. The mean of the lowest normal distribution (MLND) was used to estimate SMR for each individual from the extracted slopes (Chabot, Steffensen, et al., 2016). Slopes for MMR were extracted using a derivative of a polynomial curve fitted on each measurement (function *smooth.spline,* d.f. = 10) as in the *spline-MMR* method (Prokkola et al., 2022), i.e. from the beginning of the MMR measurement, when respiration was the highest. The slopes were then used to calculate MMR (mg O_2 h^{-1}) using *FishResp*. AS was calculated as the difference between the absolute MMR and SMR.

A linear mixed effect model was fitted (estimated using restricted maximum likelihood and *nloptwrap* optimizer) to predict SMR, MMR and a non-linear mixed effect model was fitted to predict AS, all of which were log transformed. Temperature and log-transformed mass as fixed effects (formula: Metabolic Measurement ~ Temperature + Mass). The model included batch as a random effect (formula:~1|Batch) (Zuur et al., 2009). Homoscedasticity and normality of residuals were assessed by visual inspection of residual plots. 95% Confidence Intervals (CIs) and p-values were computed using a Wald t-distribution approximation. Post-hoc comparisons of temperatures were performed using Tukey method of comparing estimates, and P value <

0.05 was considered significant. Statistical models were fitted using *nlme*, *lme4* and *lmerTest* packages in R,

and post-hoc analysis was performed in *emmeans* package (Bates et al., 2015; Kuznetsova et al., 2017; Lenth,

2017; Pinheiro et al., 1999). For visual representation, SMR, MMR, and AS data were mass-adjusted to account

for the hypo-allometric scaling of metabolic rates by linear regression of $\log_{10}$-transformed metabolic rates

against $\log_{10}$-transformed body mass (Figure 2). Finally, Pearson's correlations were calculated among the

either absolute or mass-adjusted metabolic variables.

Results

In total, the data of 72 fish was used at three temperatures across four consecutive weeks: 16°C, 19°C, and

22°C and 16 °C again (Table 1, Table S3). The metabolic rates and AS scaled positively with body mass, with

scaling exponents from 0.86 to 1.1 (Figure 2, Table 2, Table S4).

We found temperature to be a significant predictor of SMR (16–19 °C: $t(66) = 4.80$, $P < .001$; 16–22 °C: $t(66) =$

7.43, $P < .001$, Table 3). There were significant differences in SMR between all temperature groups, except

between 19 °C and 22 °C (Figure 3, Table 4). There was a 57% increase in predicted SMR for 6 °C increase in

temperature (Table 4). However, there was no significant change in MMR or AS between temperatures (Figure

3, Table 2).

Correlations among metabolic variables

At all temperatures, MMR and AS were highly positively correlated ($r > 0.9$) in both raw and mass adjusted

data. Absolute SMR and MMR were positively correlated at 16 and 22 °C. Absolute SMR and AS were positively

correlated at 16 °C. However, mass adjusted SMR and AS were negatively correlated at 19 and 22 °C (Table S5,

Figure S1).

Discussion

This study demonstrates that rising temperatures lead to an increase in standard metabolic rate (SMR) in juvenile Atlantic salmon. However, there was no change in maximum metabolic rate (MMR) or in aerobic scope (AS) across the tested temperatures. While an increase in SMR with temperature is well established in ectotherms due to the effect of temperature on biochemical rates (Clark et al., 2013; McKenzie et al., 2021; Raby et al., 2016), there are more mixed results on the relationship of MMR, AS and increasing temperatures in fishes.

The TPC of AS in fish has been suggested to be a bell shaped curve (Pörtner & Farrell, 2008). For example, in pink salmon (*O. gorbuscha*), AS peaked at 21 °C and declined with further warming, although this temperature increase occurred over a maximum of 2 days (Clark et al., 2011). Other studies note a decrease in AS, which could elude to the post peak decrease of the AS curve, as these are tested at higher temperatures or after longer acclimations, for example, in a recent study on salmon parr, AS decreased with temperature from 15 to 21°C after a three-week acclimation (Nuic et al., 2024). This could suggest long term exposure to a heatwave leading to aerobic collapse. In contrast in murray cod (*Maccullochella peelii peelii*) and temperatures that increased from 14 – 29 °C, where fish were acclimated for at least 18 h to each increase, resulted in a continuous increase in aerobic scope (Clark et al., 2005).

Other studies report a plateau of AS at high temperatures. For example, there were no differences in AS between acclimation temperatures in minnows (*Phoxinus phoxinus*) after eight months (Závorka et al., 2020). Morissette et al. (2021) found that below 23 °C, there was no difference between any metabolic rates after different temperature acclimations, however in their study all metabolic rates were measured at the same temperature, which complicates the interpretation of the results. In gobies acclimated for three weeks, AS was seen to plateau between 15 and 28 °C after initially increasing (Christensen et al., 2021). Further, in Atlantic salmon post-smolts, aerobic scope was increased with 4-week acclimations at temperatures from 3–13 °C, and plateaued from 13 °C up to a temperature of 23 °C. Interestingly, fish condition and swim speed dropped and mortality increased (Hvas et al., 2017). Upper thermal limit of salmon has been widely reported

at 23 °C (Elliott & Elliott, 2010), and notable, often cited work (Jensen et al., 1989) found optimal salmon

temperature for growth to be 16 °C. Our results agree with Hvas et al. (2017) as we found that AS plateaued at

16–22°C when measured at acclimation temperature, and it is likely that oxygen supply can limit performance

of salmon as temperature increases. An absence of increase suggests physiological systems supporting oxygen

uptake such as cardiac function, gill function and haemoglobin oxygen affinity are no longer able to scale with

increasing thermal pressure (Anttila et al., 2014; Schulte & Healy, 2022). As AS plateaus but SMR and the costs

of aerobic activities continue to increase, fish might become more constrained in their physiological state and

experience cumulative thermal stress, leading to mortality.

AS is proposed to be of ecological significance because it defines the upper limit for oxygen allocation by a fish

to sustain aerobic activities such as foraging, digestion, tissue deposition, migration, reproduction, and so

forth (Claireaux & Lefrançois, 2007; Farrell et al., 2009b; Fry, 1971; H.-O. Pörtner, 2010; Schulte, 2015). For

example, reduced appetite could be behaviourally driven to avoid anaerobic metabolism, as suggested by the

reduction of feed intake during low AS (review by (Jutfelt, 2020). In support of this, reduced growth and lower

condition factor at higher temperatures has previously been documented in Atlantic salmon, when comparing

increases to temperatures up to 19 °C (Hevrøy et al., 2015; Kullgren et al., 2013). By measuring AS as an

indicator of thermal performance, we can predict that at high temperatures, Atlantic salmon in the wild would

either have to seek thermal refuge, or compromise aspects of their performance in order to cope with the

high temperatures, which could result in poor growth or health (Koskela et al., 1997).

It is important to note that both TPCs and upper thermal limits are dependent on acclimation duration and

rate of warming (Currie et al., 2014; Lefevre et al., 2021; McKenzie et al., 2021). Most research in fishes has

been conducted with stable thermal profiles, whereby each individual experiences only one acclimation

temperature (Hvas et al., 2017). In contrast, our study accounts for a cumulative effect of exposure to higher

temperatures, using a gradual heating regime. While some acclimation is expected in the MMR over short-

term temperature increases ((Norin & Clark, 2016); and papers within), each temperature exposure in this

study occurred over 10 days, which may not be enough time to see notable plasticity in salmon's metabolic

ceiling. Sandblom et al. (2017) found no difference in MMR or AS between acute (24 h) or chronic warming

(years) regime in perch (*Perca fluviatilis*), but that all warm-acclimated fish (5 days acclimation, 23 °C) have a

higher MMR than fish acclimated at 18 °C, their reference temperature. However, an ecologically relevant

scenario for salmon in their environments is a heatwave lasting from a few days to few weeks, thus these time

frames are the most relevant to study from a conservation perspective. By testing the cumulative effect of a

heatwave mirroring the temperatures seen in the wild, we can better predict how salmon could cope in a

realistic exposure to sub-lethal temperatures (Morash et al., 2018).

The general expectation is that longer acclimation with a slower increase of temperature should be beneficial

to survival, giving the fish time to make compensatory physiological modifications such as displaying plasticity

in SMR or MMR, enabling AS to be maintained over a broad range of temperatures. However, in barramundi

MMR and AS increased when exposed to acute 10 °C warming, due to a higher increase in MMR than SMR, but

after 5 weeks of acclimation, AS was similar, mainly due to a reduction in MMR (Norin et al., 2014). A very

similar pattern was observed in black sea bass (*Centropristis striata)* and common triplefin (*Forsterygion*

lapillum), where short term exposure caused a raise in AS, before it decreased after some weeks of exposure

(Khan et al., 2014; McArley et al., 2017; Slesinger et al., 2019). Additionally, in salmon post smolts, post-stress

peak cortisol levels significantly increased with higher temperatures (Madaro et al., 2018). This suggests that

despite a stable aerobic capacity to cope with temperature increases, costly physiological mechanisms may be

occurring, and therefore a plateau in AS does not necessarily suggest a broad optimum temperature range.

MMR did not differ across temperatures in this study, which may be due to the fish being of similar age and

reared under uniform hatchery conditions (Farrell et al., 2009a; Killen et al., 2016). MMR reflects the

maximum rate at which aerobic physiological processes can occur, and at 16 °C the salmon may already have

been operating near their upper capacity for oxygen uptake and delivery (Clark et al., 2011; McKenzie &

Claireaux, 2010; Norin & Clark, 2016; Sandblom et al., 2016). Although no treatment-level differences in MMR

were detected, individual variation in thermal reaction norms of SMR and MMR may still exist but could not

be resolved with our experimental design. This limitation may help explain why AS did not decline despite SMR

increasing while MMR remained stable (Norin et al., 2014; Norin & Metcalfe, 2019; Réveillon et al., 2019).

Sample size was reduced from 128 to 72 fish due to lost data, largely caused by malfunction of the water

circulation and temperature control system. This uneven sample distribution across temperature treatments

may have limited our ability to detect patterns, particularly at 19 and 22 °C. In addition, one extreme outlier

was removed from SMR, MMR, and AS analyses based on visual inspection of histograms, and while this was a

single fish, its exclusion should be acknowledged. Mortality also occurred in the respirometers (one, three,

and two fish at 16, 19, and 22 °C, respectively), and while this represented <10% of the initial sample size for

each group, removing these data introduces some potential for survival bias. Importantly, our experiment

tested only a single hatchery stock and one juvenile age class; responses could differ in other populations or

life stages, especially given that juvenile parr are often more sensitive than eggs or spawners. Together, these

limitations highlight the need for caution in extrapolating our findings and suggest that broader testing across

stocks, life stages, and larger sample sizes will be important for future studies.

Recent advances in conservation physiology demonstrate how individual-level physiological metrics can be

used to guide policy and management strategies, as shown in studies on Sockeye salmon in Canada (Patterson

et al., 2016). Integrating TPCs and stock specific information, such as life history genotypes and developmental

stages, can inform stakeholders of specific vulnerabilities in populations and how they vary in location and

season (Ward et al., 2017). With detailed TPCs, thermal safety margins can be defined in terms of functional

success—such as swimming, feeding, and reproducing—rather than mortality alone (Pinsky et al., 2019).

Temperature-dependent fisheries management strategies have already been successfully implemented, for

example, in Canadian rivers, and could be adapted globally to account for population-specific TPCs (Van

Leeuwen et al., 2023). Management strategies, such as adjusting harvest timing and identifying thermal

refuges for protection and restoration efforts, could help mitigate the impact of rising temperatures on

vulnerable salmon populations.

In recent years, extreme heat events have had catastrophic impacts on migration success and survival across

species of salmonids, indicating that a better understanding of their thermal performance is critical for

conservation (Baisez et al., 2011; Hinch et al., 2012; Martins et al., 2011, 2012; Muñoz et al., 2021). Our study

showed that aerobic scope was not increased between 16 and 22°C in juvenile Atlantic salmon despite a

significant increase in SMR. This, combined with the increasing metabolic costs of activities, such as swimming

and digestion, indicate that the stable aerobic scope may be leaving juvenile salmon vulnerable to

deteriorating performance at temperatures commonly experienced in present-day heatwaves.

Ethical Declarations

The experiment was performed with permit granted by the Finnish Project Authorisation Board no.

ESAVI/16748/2023.

Acknowledgements

We are grateful for Jonna Hänninen, Juha Hänninen, and staff at Luke Taivalkoski and Luke Laukaa for access

to fish and help during the experiment, and for Kari Jauhiainen for technical support, and Prof. Craig Primmer,

Dr. Tutku Aykanat, and Dr. Silva Uusi-Heikkilä for access to equipment. We thank Lilian Redon for feedback on

an early version of the manuscript.

Author Contributions

LC and JP: experimental design, conducting experiment, data analysis and writing the manuscript. JP:

supervision, management, and funding acquisition. SM: equipment and experimental design. MR and ES: fish

care, equipment maintenance and assisting with data collection.

Funding

This study was funded by the Research Council of Finland (348965 and 353760), Finnish Cultural Foundation

(00220693) and by the Natural Resources Institute Finland.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability

Data is hosted on Zenodo data repository (https://doi.org/10.5281/zenodo.15308028) and code for analysis is

hosted on Github (https://github.com/jprokkola/Salmon-aerobic-scope-2023-public.git).

References

Aas, Ø., Klemetsen, A., Einum, S., & Skurdal, J. (2010). Atlantic Salmon Ecology. *Salmon Ecological Research and*

Conservation. https://doi.org/10.1002/9781444327755

Alfonso, S., Gesto, M., & Sadoul, B. (2021). Temperature increase and its effects on fish stress physiology in the context

of global warming. *Journal of Fish Biology*, *98*(6), 1496–1508. https://doi.org/10.1111/jfb.14599

415   Alioravainen, N., Orell, P., & Erkinaro, J. (2023). Long-Term Trends in Freshwater and Marine Growth Patterns in Three

416   Sub-Arctic Atlantic Salmon Populations. *Fishes*, *8*(9), 441. https://doi.org/10.3390/fishes8090441

417   Angilletta, M. J., Huey, R. B., & Frazier, M. R. (2010). Thermodynamic Effects on Organismal Performance: Is Hotter

418   Better? *Physiological and Biochemical Zoology*, *83*(2), 197–206. https://doi.org/10.1086/648567

419   Anttila, K., Jørgensen, S. M., Casselman, M. T., Timmerhaus, G., Farrell, A. P., & Takle, H. (2014). Association between

420   swimming performance, cardiorespiratory morphometry, and thermal tolerance in Atlantic salmon (Salmo salar

421   L.). *Frontiers in Marine Science*, *1*. https://doi.org/10.3389/fmars.2014.00076

422   Baisez, A., Bach, J.-M., Leon, C., Parouty, T., Terrade, R., Hoffmann, M., & Laffaille, P. (2011). Migration delays and

423   mortality of adult Atlantic salmon Salmo salar en route to spawning grounds on the River Allier, France.

424   *Endangered Species Research*, *15*(3), 265–270. https://doi.org/10.3354/esr00384

425   Baltar, F., Bayer, B., Bednarsek, N., Deppeler, S., Escribano, R., Gonzalez, C. E., Hansman, R. L., Mishra, R. K., Moran, M.

426   A., Repeta, D. J., Robinson, C., Sintes, E., Tamburini, C., Valentin, L. E., & Herndl, G. J. (2019). Towards Integrating

427   Evolution, Metabolism, and Climate Change Studies of Marine Ecosystems. *Trends in Ecology & Evolution*,

428   *34*(11), 1022–1033. https://doi.org/10.1016/j.tree.2019.07.003

429   Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using **lme4**. *Journal of*

430   *Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

431   Breedveld, M. C., Devigili, A., Borgheresi, O., & Gasparini, C. (2023). Reproducing in hot water: Experimental heatwaves

432   deteriorate multiple reproductive traits in a freshwater ectotherm. *Functional Ecology*, *37*(4), 989–1004.

433   https://doi.org/10.1111/1365-2435.14279

434   Brett, J. R. (1964). The respiratory metabolism and swimming performance of young sockeye salmon. *Journal of Fish*

435   *Biology*, *21*(5), 1183–1226. https://doi.org/10.1139/f64-103

436   Burraco, P., Orizaola, G., Monaghan, P., & Metcalfe, N. B. (2020). Climate change and ageing in ectotherms. *Global*

437   *Change Biology*, *26*(10), 5371–5381. https://doi.org/10.1111/gcb.15305

438   Casselman, M. T., Anttila, K., & Farrell, A. P. (2012). Using maximum heart rate as a rapid screening tool to determine

439   optimum temperature for aerobic scope in Pacific salmon *Oncorhynchus* spp. *Journal of Fish Biology*, *80*(2), 358–

440   377. https://doi.org/10.1111/j.1095-8649.2011.03182.x

Chabot, D., McKenzie, D. J., & Craig, J. F. (2016). Metabolic rate in fishes: Definitions, methods and significance for

conservation physiology. *Journal of Fish Biology*, *88*(1), 1–9. https://doi.org/10.1111/jfb.12873

Chabot, D., Steffensen, J. F., & Farrell, A. P. (2016). The determination of standard metabolic rate in fishes. *Journal of*

Fish Biology, *88*(1), 81–121. https://doi.org/10.1111/jfb.12845

Chabot, D., Zhang, Y., & Farrell, A. P. (2021). Valid oxygen uptake measurements: Using high r^2 values with good

intentions can bias upward the determination of standard metabolic rate. *Journal of Fish Biology*, *98*(5), 1206–

1216. https://doi.org/10.1111/jfb.14650

Christensen, E. A. F., Andersen, L. E. J., Bergsson, H., Steffensen, J. F., & Killen, S. S. (2021). Erratum: Shuttle-box systems

for studying preferred environmental ranges by aquatic animals (Conservation Physiology 9:1 (coab028) DOI:

10.1093/conphys/coab028). *Conservation Physiology*, *9*(1), 1–21. https://doi.org/10.1093/conphys/coab051

Christensen, E. A. F., Norin, T., Tabak, I., Van Deurs, M., & Behrens, J. W. (2020). Effects of temperature on physiological

performance and behavioral thermoregulation in an invasive fish, the round goby. *Journal of Experimental*

Biology, jeb.237669. https://doi.org/10.1242/jeb.237669

Claireaux, G., Couturier, C., & Groison, A. L. (2006). Effect of temperature on maximum swimming speed and cost of

transport in juvenile European sea bass (Dicentrarchus labrax). *Journal of Experimental Biology*, *209*(17), 3420–

3428. https://doi.org/10.1242/jeb.02346

Claireaux, G., & Lefrançois, C. (2007). Linking environmental variability and fish performance: Integration through the

concept of scope for activity. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1487),

2031–2041. https://doi.org/10.1098/rstb.2007.2099

Clark, T. D., Jeffries, K. M., Hinch, S. G., & Farrell, A. P. (2011). Exceptional aerobic scope and cardiovascular performance

of pink salmon (*Oncorhynchus gorbuscha*) may underlie resilience in a warming climate. *Journal of*

Experimental Biology, *214*(18), 3074–3081. https://doi.org/10.1242/jeb.060517

Clark, T. D., Ryan, T., Ingram, B. A., Woakes, A. J., Butler, P. J., & Frappell, P. B. (2005). Factorial Aerobic Scope Is

Independent of Temperature and Primarily Modulated by Heart Rate in Exercising Murray Cod (*Maccullochella*

peelii peelii). *Physiological and Biochemical Zoology*, *78*(3), 347–355. https://doi.org/10.1086/430034

Clark, T. D., Sandblom, E., & Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of climate change:

Respirometry, relevance and recommendations. *Journal of Experimental Biology*, *216*(15), 2771–2782.

https://doi.org/10.1242/jeb.084251

Currie, S., Schulte, P. M., & Evans, D. H. (2014). Thermal stress. *The Physiology of Fishes*, *4*, 257–279.

Dadswell, M., Spares, A., Reader, J., McLean, M., McDermott, T., Samways, K., & Lilly, J. (2022). The Decline and

Impending Collapse of the Atlantic Salmon (*Salmo salar*) Population in the North Atlantic Ocean: A Review of

Possible Causes. *Reviews in Fisheries Science & Aquaculture*, *30*(2), 215–258.

https://doi.org/10.1080/23308249.2021.1937044

Del Rio, A. M., Mukai, G. N., Martin, B. T., Johnson, R. C., Fangue, N. A., Israel, J. A., & Todgham, A. E. (2021). Differential

sensitivity to warming and hypoxia during development and long-term effects of developmental exposure in

early life stage Chinook salmon. *Conservation Physiology*, *9*(1), coab054.

https://doi.org/10.1093/conphys/coab054

Desforges, J. E., Birnie-Gauvin, K., Jutfelt, F., Gilmour, K. M., Eliason, E. J., Dressler, T. L., McKenzie, D. J., Bates, A. E.,

Lawrence, M. J., Fangue, N., & Cooke, S. J. (2023). The ecological relevance of critical thermal maxima

methodology for fishes. *Journal of Fish Biology*, *102*(5), 1000–1016. https://doi.org/10.1111/jfb.15368

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P. R. (2008). Impacts

of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*,

105(18), 6668–6672. https://doi.org/10.1073/pnas.0709472105

Dwyer, W. P., & Piper, R. G. (1987). Atlantic Salmon Growth Efficiency as Affected by Temperature. *The Progressive Fish-

Culturist*, *49*(1), 57–59. https://doi.org/10.1577/1548-8640(1987)49<57:ASGEAA>2.0.CO;2

Eliason, E. J., Clark, T. D., Hague, M. J., Hanson, L. M., Gallagher, Z. S., Jeffries, K. M., Gale, M. K., Patterson, D. A., Hinch,

S. G., & Farrell, A. P. (2011). Differences in Thermal Tolerance Among Sockeye Salmon Populations. *Science*,

332(6025), 109–112. https://doi.org/10.1126/science.1199158

Eliason, E. J., Clark, T. D., Hinch, S. G., & Farrell, A. P. (2013). Cardiorespiratory collapse at high temperature in swimming

adult sockeye salmon. *Conservation Physiology*, *1*(1), cot008–cot008. https://doi.org/10.1093/conphys/cot008

Eliason, E. J., Wilson, S. M., Farrell, A. P., Cooke, S. J., & Hinch, S. G. (2013). Low cardiac and aerobic scope in a coastal

population of sockeye salmon *Oncorhynchus nerka* with a short upriver migration. *Journal of Fish Biology*, *82*(6),

2104–2112. https://doi.org/10.1111/jfb.12120

Elliott, J. M. (1982). The effects of temperature and ration size on the growth and energetics of salmonids in captivity.

Comparative Biochemistry and Physiology Part B: Comparative Biochemistry, *73*(1), 81–91.

https://doi.org/10.1016/0305-0491(82)90202-4

Elliott, J. M., & Elliott, J. A. (2010). Temperature requirements of Atlantic salmon Salmo salar, brown trout Salmo trutta

and Arctic charr Salvelinus alpinus: Predicting the effects of climate change. *Journal of Fish Biology*, *77*(8), 1793–

1817. https://doi.org/10.1111/j.1095-8649.2010.02762.x

Elliott, J. M., & Hurley, M. A. (1997). A functional model for maximum growth of Atlantic Salmon parr, Salmo salar, from

two populations in northwest England. *Functional Ecology*, *11*(5), 592–603. https://doi.org/10.1046/j.1365-

2435.1997.00130.x

Farrell, A. P. (2016). Pragmatic perspective on aerobic scope: Peaking, plummeting, pejus and apportioning. *Journal of*

Fish Biology, *88*(1), 322–343. https://doi.org/10.1111/jfb.12789

Farrell, A. P., Eliason, E. J., Sandblom, E., & Clark, T. D. (2009a). Fish cardiorespiratory physiology in an era of climate

changeThe present review is one of a series of occasional review articles that have been invited by the Editors

and will feature the broad range of disciplines and expertise represented in our Editorial Advisory Board.

Canadian Journal of Zoology, *87*(10), 835–851. https://doi.org/10.1139/Z09-092

Farrell, A. P., Eliason, E. J., Sandblom, E., & Clark, T. D. (2009b). Fish cardiorespiratory physiology in an era of climate

changeThe present review is one of a series of occasional review articles that have been invited by the Editors

and will feature the broad range of disciplines and expertise represented in our Editorial Advisory Board.

Canadian Journal of Zoology, *87*(10), 835–851. https://doi.org/10.1139/Z09-092

Fisher, K. C., & Elson, P. F. (1950). The Selected Temperature of Atlantic Salmon and Speckled Trout and the Effect of

Temperature on the Response to an Electrical Stimulus. *Physiological Zoology*, *23*(1), 27–34.

https://doi.org/10.1086/physzool.23.1.30084896

Fry, F. E. J. (1971). The Effect of Environmental Factors on the Physiology of Fish. In W. S. Hoar & D. J. Randall (Eds.), *Fish*

Physiology (Vol. 6, pp. 1–98). Academic Press. https://doi.org/10.1016/S1546-5098(08)60146-6

Fry, F. E. J., Black, V. S., & Black, E. C. (1947). Influence of temperature on the asphyxiation of young goldfish (carassius

auratus l.) under various tensions of oxygen and carbon dioxide. *The Biological Bulletin*, *92*(3), 217–224.

https://doi.org/10.2307/1538308

Hevrøy, E. M., Tipsmark, C. K., Remø, S. C., Hansen, T., Fukuda, M., Torgersen, T., Vikeså, V., Olsvik, P. A., Waagbø, R., &

Shimizu, M. (2015). Role of the GH-IGF-1 system in Atlantic salmon and rainbow trout postsmolts at elevated

water temperature. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *188*,

127–138. https://doi.org/10.1016/j.cbpa.2015.06.030

Hinch, S. G., Cooke, S. J., Farrell, A. P., Miller, K. M., Lapointe, M., & Patterson, D. A. (2012). Dead fish swimming: A

review of research on the early migration and high premature mortality in adult Fraser River sockeye salmon

Oncorhynchus nerka. *Journal of Fish Biology*, *81*(2), 576–599. https://doi.org/10.1111/j.1095-8649.2012.03360.x

Hochachka, P. W., & Somero, G. N. (1973). *Strategies of biochemical adaptation* (Vol. 358). Saunders.

Hutchings, J. A., & Jones, M. E. (1998). Life history variation and growth rate thresholds for maturity in Atlantic salmon,

Salmo salar. *Canadian Journal of Fisheries and Aquatic Sciences*, *55*(S1), 22–47. https://doi.org/10.1139/d98-004

Hvas, M., Folkedal, O., Imsland, A., & Oppedal, F. (2017). The effect of thermal acclimation on aerobic scope and critical

swimming speed in Atlantic salmon *Salmo salar*. *Journal of Experimental Biology*, jeb.154021.

https://doi.org/10.1242/jeb.154021

ICES. (2023). *Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel*

on Climate Change (Synthesis Report No. 6; First, Climate Change 2023). Intergovernmental Panel on Climate

Change (IPCC). https://doi.org/10.59327/IPCC/AR6-9789291691647

ICES. (2024a). *Baltic Salmon and Trout Assessment Working Group (WGBAST)* (ICES Scientific Report No. 1). ICES.

https://doi.org/10.17895/ices.pub.25868665.v1

ICES. (2024b). *Working Group on North Atlantic Salmon (WGNAS)* [Report]. ICES Scientific Reports.

https://doi.org/10.17895/ices.pub.25730247.v1

Jahn, M., & Seebacher, F. (2022). Variations in cost of transport and their ecological consequences: A review. *Journal of*

Experimental Biology, *225*(15), jeb243646. https://doi.org/10.1242/jeb.243646

Jensen, A. J., Karlsson, S., Fiske, P., Hansen, L. P., Østborg, G. M., & Hindar, K. (2014). Origin and life history of Atlantic

salmon (*Salmo salar*) near their northernmost oceanic limit. *Canadian Journal of Fisheries and Aquatic Sciences*,

71(11), 1740–1746. https://doi.org/10.1139/cjfas-2014-0169

Johansen, J. L., & Jones, G. P. (2011). Increasing ocean temperature reduces the metabolic performance and swimming

ability of coral reef damselfishes. *Global Change Biology*, *17*(9), 2971–2979. https://doi.org/10.1111/j.1365-

2486.2011.02436.x

Johansson, D., Ruohonen, K., Juell, J.-E., & Oppedal, F. (2009). Swimming depth and thermal history of individual Atlantic

salmon (Salmo salar L.) in production cages under different ambient temperature conditions. *Aquaculture*,

290(3–4), 296–303. https://doi.org/10.1016/j.aquaculture.2009.02.022

Jørgensen, L. B., Malte, H., Ørsted, M., Klahn, N. A., & Overgaard, J. (2021). A unifying model to estimate thermal

tolerance limits in ectotherms across static, dynamic and fluctuating exposures to thermal stress. *Scientific*

Reports, *11*(1), 12840. https://doi.org/10.1038/s41598-021-92004-6

Jørgensen, L. B., Ørsted, M., Malte, H., Wang, T., & Overgaard, J. (2022). Extreme escalation of heat failure rates in

ectotherms with global warming. *Nature*, *611*(7934), 93–98. https://doi.org/10.1038/s41586-022-05334-4

Jutfelt, F. (2020). Metabolic adaptation to warm water in fish. *Functional Ecology*, *34*(6), 1138–1141.

https://doi.org/10.1111/1365-2435.13558

Jutfelt, F., Norin, T., Åsheim, E. R., Rowsey, L. E., Andreassen, A. H., Morgan, R., Clark, T. D., & Speers-Roesch, B. (2021).

'Aerobic scope protection' reduces ectotherm growth under warming. *Functional Ecology*, *35*(7), 1397–1407.

https://doi.org/10.1111/1365-2435.13811

Khan, J. R., Pether, S., Bruce, M., Walker, S. P., & Herbert, N. A. (2014). Optimum temperatures for growth and feed

conversion in cultured hapuku (*Polyprion oxygeneios*)—Is there a link to aerobic metabolic scope and final

temperature preference? *Aquaculture*, *430*, 107–113. https://doi.org/10.1016/j.aquaculture.2014.03.046

Killen, S. S., Christensen, E. A. F., Cortese, D., Závorka, L., Norin, T., Cotgrove, L., Crespel, A., Munson, A., Nati, J. J. H.,

Papatheodoulou, M., & McKenzie, D. J. (2021). Guidelines for reporting methods to estimate metabolic rates by

aquatic intermittent-flow respirometry. *Journal of Experimental Biology*, *224*(18).

https://doi.org/10.1242/jeb.242522

Killen, S. S., Glazier, D. S., Rezende, E. L., Clark, T. D., Atkinson, D., Willener, A. S. T., & Halsey, L. G. (2016). Ecological

Influences and Morphological Correlates of Resting and Maximal Metabolic Rates across Teleost Fish Species.

The American Naturalist, *187*(5), 592–606. https://doi.org/10.1086/685893

Koskela, J., Pirhonen, J., & Jobling, M. (1997). Feed intake, growth rate and body composition of juvenile Baltic salmon

exposed to different constant temperatures. *Aquaculture International*, *5*, 351–360.

Koskela, J., Pirhonen, J., & Jobling, M. (2005). Variations in feed intake and growth of Baltic salmon and brown trout

exposed to continuous light at constant low temperatures. *Journal of Fish Biology*, *50*(4), 837–845.

https://doi.org/10.1111/j.1095-8649.1997.tb01976.x

Kullgren, A., Jutfelt, F., Fontanillas, R., Sundell, K., Samuelsson, L., Wiklander, K., Kling, P., Koppe, W., Larsson, D. G. J.,

Björnsson, B. T., & Jönsson, E. (2013). The impact of temperature on the metabolome and endocrine metabolic

signals in Atlantic salmon (Salmo salar). *Comparative Biochemistry and Physiology Part A: Molecular &*

Integrative Physiology, *164*(1), 44–53. https://doi.org/10.1016/j.cbpa.2012.10.005

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). **lmerTest** Package: Tests in Linear Mixed Effects Models.

Journal of Statistical Software, *82*(13). https://doi.org/10.18637/jss.v082.i13

Lacroix, G. L. (2013). Population-specific ranges of oceanic migration for adult Atlantic salmon (Salmo salar) documented

using pop-up satellite archival tags. *Canadian Journal of Fisheries and Aquatic Sciences*, *70*(7), 1011–1030.

https://doi.org/10.1139/cjfas-2013-0038

Lefevre, S., Wang, T., & McKenzie, D. J. (2021). The role of mechanistic physiology in investigating impacts of global

warming on fishes. *Journal of Experimental Biology*, *224*. https://doi.org/10.1242/jeb.238840

Lenth, R. V. (2017). *emmeans: Estimated Marginal Means, aka Least-Squares Means* (p. 1.11.0) [Dataset].

https://doi.org/10.32614/CRAN.package.emmeans

Little, A. G., Loughland, I., & Seebacher, F. (2020). What do warming waters mean for fish physiology and fisheries?

Journal of Fish Biology, *97*(2), 328–340. https://doi.org/10.1111/jfb.14402

Little, A. G., & Seebacher, F. (2021). Physiological Performance Curves: When Are They Useful? *Frontiers in Physiology*,

12, 805102. https://doi.org/10.3389/fphys.2021.805102

Loughland, I., & Seebacher, F. (2020). Differences in oxidative status explain variation in thermal acclimation capacity

between individual mosquitofish (Gambusia holbrooki). *Functional Ecology*, *34*(7), 1380–1390.

https://doi.org/10.1111/1365-2435.13563

MacNutt, M. J., Hinch, S. G., Lee, C. G., Phibbs, J. R., Lotto, A. G., Healey, M. C., & Farrell, A. P. (2006). Temperature

effects on swimming performance, energetics, and aerobic capacities of mature adult pink salmon

(Oncorhynchus gorbuscha) compared with those of sockeye salmon (Oncorhynchus nerka). *Canadian Journal of*

Zoology, *84*(1), 88–97. https://doi.org/10.1139/z05-181

Madaro, A., Folkedal, O., Maiolo, S., Alvanopoulou, M., & Olsen, R. E. (2018). Effects of acclimation temperature on

cortisol and oxygen consumption in Atlantic salmon (*Salmo salar*) post-smolt exposed to acute stress.

Aquaculture, *497*, 331–335. https://doi.org/10.1016/j.aquaculture.2018.07.056

Martins, E. G., Hinch, S. G., Patterson, D. A., Hague, M. J., Cooke, S. J., Miller, K. M., Lapointe, M. F., English, K. K., &

Farrell, A. P. (2011). Effects of river temperature and climate warming on stock-specific survival of adult

migrating Fraser River sockeye salmon (Oncorhynchus nerka). *Global Change Biology*, *17*(1), 99–114.

https://doi.org/10.1111/j.1365-2486.2010.02241.x

Martins, E. G., Hinch, S. G., Patterson, D. A., Hague, M. J., Cooke, S. J., Miller, K. M., Robichaud, D., English, K. K., &

Farrell, A. P. (2012). High river temperature reduces survival of sockeye salmon (Oncorhynchus nerka)

approaching spawning grounds and exacerbates female mortality. *Canadian Journal of Fisheries and Aquatic*

Sciences, *69*(2), 330–342. https://doi.org/10.1139/f2011-154

McArley, T. J., Hickey, A. J. R., & Herbert, N. A. (2017). Chronic warm exposure impairs growth performance and reduces

thermal safety margins in the common triplefin fish (Forsterygion lapillum). *Journal of Experimental Biology*,

220(19), 3527–3535. https://doi.org/10.1242/jeb.162099

McKenzie, D. J., & Claireaux, G. (2010). The Effects of Environmental Factors on the Physiology of Aerobic Exercise. In

Fish Locomotion. CRC Press.

McKenzie, D. J., Zhang, Y., Eliason, E. J., Schulte, P. M., Claireaux, G., Blasco, F. R., Nati, J. J. H., & Farrell, A. P. (2021).

Intraspecific variation in tolerance of warming in fishes. *Journal of Fish Biology*, *98*(6), 1536–1555.

https://doi.org/10.1111/jfb.14620

Morash, A. J., Neufeld, C., MacCormack, T. J., & Currie, S. (2018). The importance of incorporating natural thermal

variation when evaluating physiological performance in wild species. *Journal of Experimental Biology*, *221*(14),

jeb164673. https://doi.org/10.1242/jeb.164673

Morash, A. J., Speers-Roesch, B., Andrew, S., & Currie, S. (2021). The physiological ups and downs of thermal variability

in temperate freshwater ecosystems. *Journal of Fish Biology*, *98*(6), 1524–1535.

https://doi.org/10.1111/jfb.14655

Morgan, R., Andreassen, A. H., Åsheim, E. R., Finnøen, M. H., Dresler, G., Brembu, T., Loh, A., Miest, J. J., & Jutfelt, F.

(2022). Reduced physiological plasticity in a fish adapted to stable temperatures. *Proceedings of the National*

Academy of Sciences, *119*(22), e2201919119. https://doi.org/10.1073/pnas.2201919119

Morissette, J., Swart, S., MacCormack, T. J., Currie, S., & Morash, A. J. (2021). Thermal variation near the thermal

optimum does not affect the growth, metabolism or swimming performance in wild Atlantic salmon Salmo salar.

Journal of Fish Biology, *98*(6), 1585–1589. https://doi.org/10.1111/jfb.14348

Morozov, S. (2024). *PumpResp* [Computer software]. https://github.com/embedded-sergey/PumpResp

Morozov, S., McCairns, R. J. S., & Merilä, J. (2019). FishResp: R package and GUI application for analysis of aquatic

respirometry data. *Conservation Physiology*, *7*(1), coz003. https://doi.org/10.1093/conphys/coz003

Muñoz, N. J., Reid, B., Correa, C., Neff, B. D., & Reynolds, J. D. (2021). Non-native Chinook salmon add nutrient subsidies

and functional novelty to Patagonian streams. *Freshwater Biology*, *66*(3), 495–508.

https://doi.org/10.1111/fwb.13655

Navarro, J. M., Paschke, K., Ortiz, A., Vargas-Chacoff, L., Pardo, L. M., & Valdivia, N. (2019). The Antarctic fish Harpagifer

antarcticus under current temperatures and salinities and future scenarios of climate change. *Progress in*

Oceanography, *174*, 37–43. https://doi.org/10.1016/j.pocean.2018.09.001

Norin, T., & Clark, T. D. (2016). Measurement and relevance of maximum metabolic rate. *Journal of Fish Biology*, *88*,

122–151. https://doi.org/10.1111/jfb.12796

Norin, T., Malte, H., & Clark, T. D. (2014). Aerobic scope does not predict the performance of a tropical eurythermal fish

at elevated temperatures. *Journal of Experimental Biology*, *217*(2), 244–251.

https://doi.org/10.1242/jeb.089755

Norin, T., & Metcalfe, N. B. (2019). *Ecological and evolutionary consequences of metabolic rate plasticity in response to*

environmental change. 1–9.

Nuic, B., Bowden, A., Franklin, C. E., & Cramp, R. L. (2024). Atlantic salmon Salmo salar do not prioritize digestion when

energetic budgets are constrained by warming and hypoxia. *Journal of Fish Biology*, *104*(6), 1718–1731.

https://doi.org/10.1111/jfb.15693

Ørsted, M., Jørgensen, L. B., & Overgaard, J. (2022). Finding the right thermal limit: A framework to reconcile ecological,

physiological and methodological aspects of CTmax in ectotherms. *Journal of Experimental Biology*, *225*(19),

jeb244514. https://doi.org/10.1242/jeb.244514

O'Sullivan, A. M., Corey, E. M., Collet, E. N., Helminen, J., Curry, R. A., MacIntyre, C., & Linnansaari, T. (2023). Timing and

frequency of high temperature events bend the onset of behavioural thermoregulation in Atlantic salmon

(Salmo salar). *Conservation Physiology*, *11*(1), coac079. https://doi.org/10.1093/conphys/coac079

Patterson, D. A., Cooke, S. J., Hinch, S. G., Robinson, K. A., Young, N., Farrell, A. P., & Miller, K. M. (2016). A perspective

on physiological studies supporting the provision of scientific advice for the management of Fraser River sockeye

salmon (*Oncorhynchus nerka*). *Conservation Physiology*, *4*(1), cow026.

https://doi.org/10.1093/conphys/cow026

Peralta-Maraver, I., & Rezende, E. L. (2021). Heat tolerance in ectotherms scales predictably with body size. *Nature*

Climate Change, *11*(1), 58–63. https://doi.org/10.1038/s41558-020-00938-y

Perry, A. L., Low, P. J., Ellis, J. R., & Reynolds, J. D. (2005). Climate Change and Distribution Shifts in Marine Fishes.

Science, *308*(5730), 1912–1915. https://doi.org/10.1126/science.1111322

Pinheiro, J., Bates, D., & R Core Team. (1999). *nlme: Linear and Nonlinear Mixed Effects Models* (p. 3.1-168) [Dataset].

https://doi.org/10.32614/CRAN.package.nlme

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater vulnerability to warming of

marine versus terrestrial ectotherms. *Nature*, *569*(7754), 108–111. https://doi.org/10.1038/s41586-019-1132-4

Pörtner, H. O., & Farrell, A. P. (2008). Ecology: Physiology and climate change. *Science*, *322*(5902), 690–692.

https://doi.org/10.1126/science.1163156

Pörtner, H.-O. (2010). Oxygen- and capacity-limitation of thermal tolerance: A matrix for integrating climate-related stressor effects in marine ecosystems. *Journal of Experimental Biology*, *213*(6), 881–893. https://doi.org/10.1242/jeb.037523

Priede, I. G. (1985). Metabolic Scope in Fishes. In P. Tytler & P. Calow (Eds.), *Fish Energetics* (pp. 33–64). Springer Netherlands. https://doi.org/10.1007/978-94-011-7918-8_2

Prokkola, J. M., Åsheim, E. R., Morozov, S., Bangura, P., Erkinaro, J., Ruokolainen, A., Primmer, C. R., & Aykanat, T. (2022). Genetic coupling of life-history and aerobic performance in Atlantic salmon. *Proceedings of the Royal Society B: Biological Sciences*, *289*(1967), 20212500. https://doi.org/10.1098/rspb.2021.2500

Raby, G. D., Casselman, M. T., Cooke, S. J., Hinch, S. G., Farrell, A. P., & Clark, T. D. (2016). Aerobic scope increases throughout an ecologically relevant temperature range in coho salmon. *Journal of Experimental Biology*, jeb.137166. https://doi.org/10.1242/jeb.137166

Raby, G. D., Doherty, C. L. J., Mokdad, A., Pitcher, T. E., & Fisk, A. T. (2020). Post-exercise respirometry underestimates maximum metabolic rate in juvenile salmon. *Conservation Physiology*, *8*(1), coaa063. https://doi.org/10.1093/conphys/coaa063

Reddin, D. G. (1985). Atlantic Salmon (*Salmo salar*) On and East of the Grand Bank. *Journal of Northwest Atlantic Fishery Science*, *6*, 157–164. https://doi.org/10.2960/J.v6.a16

Réveillon, T., Rota, T., Chauvet, É., Lecerf, A., & Sentis, A. (2019). Repeatable inter-individual variation in the thermal sensitivity of metabolic rate. *Oikos*, *128*(11), 1633–1640. https://doi.org/10.1111/oik.06392

Sadoul, B., & Vijayan, M. M. (2016). Stress and Growth. In C. B. Schreck, L. Tort, A. P. Farrell, & C. J. Brauner (Eds.), *Fish Physiology* (Vol. 35, pp. 167–205). Academic Press. https://doi.org/10.1016/B978-0-12-802728-8.00005-9

Sandblom, E., Clark, T. D., Gräns, A., Ekström, A., Brijs, J., Sundström, L. F., Odelström, A., Adill, A., Aho, T., & Jutfelt, F. (2016). Physiological constraints to climate warming in fish follow principles of plastic floors and concrete ceilings. *Nature Communications*, *7*(1), 11447. https://doi.org/10.1038/ncomms11447

Schulte, P. M. (2015). The effects of temperature on aerobic metabolism: Towards a mechanistic understanding of the responses of ectotherms to a changing environment. *Journal of Experimental Biology*, *218*(12), 1856–1866. https://doi.org/10.1242/jeb.118851

Schulte, P. M., & Healy, T. M. (2022). Chapter 9—Physiological diversity and its importance for fish conservation and

management in the Anthropocene. In S. J. Cooke, N. A. Fangue, A. P. Farrell, C. J. Brauner, & E. J. Eliason (Eds.),

Fish Physiology (Vol. 39, pp. 435–477). Academic Press. https://doi.org/10.1016/bs.fp.2022.04.009

Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal Performance Curves, Phenotypic Plasticity, and the Time

Scales of Temperature Exposure. *Integrative and Comparative Biology*, *51*(5), 691–702.

https://doi.org/10.1093/icb/icr097

Seebacher, F. (2009). Responses to temperature variation: Integration of thermoregulation and metabolism in

vertebrates. *Journal of Experimental Biology*, *212*(18), 2885–2891. https://doi.org/10.1242/jeb.024430

Seebacher, F., & Little, A. G. (2021). Plasticity of Performance Curves in Ectotherms: Individual Variation Modulates

Population Responses to Environmental Change. *Frontiers in Physiology*, *12*.

https://doi.org/10.3389/fphys.2021.733305

Simpson, A. L. (1992). Differences in body size and lipid reserves between maturing and nonmaturing Atlantic salmon

parr, Salmo salar L. *Canadian Journal of Zoology*, *70*(9), 1737–1742. https://doi.org/10.1139/z92-241

Slesinger, E., Andres, A., Young, R., Seibel, B., Saba, V., Phelan, B., Rosendale, J., Wieczorek, D., & Saba, G. (2019). The

effect of ocean warming on black sea bass (Centropristis striata) aerobic scope and hypoxia tolerance. *PLOS

ONE*, *14*(6), e0218390. https://doi.org/10.1371/journal.pone.0218390

Strøm, J. F., Rikardsen, A. H., Campana, S. E., Righton, D., Carr, J., Aarestrup, K., Stokesbury, M. J. W., Gargan, P., Javierre,

P. C., & Thorstad, E. B. (2019). Ocean predation and mortality of adult Atlantic salmon. *Scientific Reports*, *9*(1),

7890. https://doi.org/10.1038/s41598-019-44041-5

Svendsen, M. B. S., Bushnell, P. G., & Steffensen, J. F. (2016). Design and setup of intermittent-flow respirometry system

for aquatic organisms. *Journal of Fish Biology*, *88*(1), 26–50. https://doi.org/10.1111/jfb.12797

Thorpe, J. E. (2007). Maturation responses of salmonids to changing developmental opportunities. *Marine Ecology

Progress Series*, *335*, 285–288. https://doi.org/10.3354/meps335285

Van Leeuwen, T. E., Lehnert ,Sarah J., Breau ,Cindy, Fitzsimmons ,Michelle, Kelly ,Nicholas I., Dempson ,J. Brian, Neville

,Victoria M., Young ,Mark, Keefe ,Donald, Bird ,Tomas J., & and Cote, D. (2023). Considerations for Water

Temperature-Related Fishery Closures in Recreational Atlantic Salmon (Salmo salar) Catch and Release Fisheries:

A Case Study from Eastern Canada. *Reviews in Fisheries Science & Aquaculture*, *31*(4), 598–619.

https://doi.org/10.1080/23308249.2023.2242959

Verberk, W. C. E. P., Sandker, J. F., Van De Pol, I. L. E., Urbina, M. A., Wilson, R. W., McKenzie, D. J., & Leiva, F. P. (2022).

Body mass and cell size shape the tolerance of fishes to low oxygen in a temperature-dependent manner. *Global*

Change Biology, *28*(19), 5695–5707. https://doi.org/10.1111/gcb.16319

Wade, N. M., Clark, T. D., Maynard, B. T., Atherton, S., Wilkinson, R. J., Smullen, R. P., & Taylor, R. S. (2019). Effects of an

unprecedented summer heatwave on the growth performance, flesh colour and plasma biochemistry of marine

cage-farmed Atlantic salmon (*Salmo salar*). *Journal of Thermal Biology*, *80*, 64–74.

https://doi.org/10.1016/j.jtherbio.2018.12.021

Ward, H. G. M., Post, J. R., Lester, N. P., Askey, P. J., & Godin, T. (2017). Empirical evidence of plasticity in life-history

characteristics across climatic and fish density gradients. *Canadian Journal of Fisheries and Aquatic Sciences*,

74(4), 464–474. https://doi.org/10.1139/cjfas-2016-0023

Závorka, L., Koeck, B., Armstrong, T. A., Soğanci, M., Crespel, A., & Killen, S. S. (2020). Reduced exploration capacity

despite brain volume increase in warm acclimated common minnow. *The Journal of Experimental Biology*,

i(May), jeb.223453. https://doi.org/10.1242/jeb.223453

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and extensions in ecology*

with R. Springer. https://doi.org/10.1007/978-0-387-87458-6

Table 1: Descriptive statistics for Mass (g) and Length (mm) for juvenile salmon tested at different temperatures. n = number of individuals, x̄ = mean metabolic rate, σ = standard deviation, Min = minimum value, Max = maximum value.

Temperature (°C)	n	Mass (g)				Length (mm)			
		x̄	σ	Min	Max	x̄	σ	Min	Max
16 (week 1)	22	1.42	0.41	1.01	2.59	57.64	5.38	50	69
19	19	1.35	0.35	0.70	1.88	56.84	4.71	47	64
22	16	1.79	0.42	0.88	2.40	61.00	4.35	54	68
16 (week 4)	15	2.56	0.88	1.40	4.32	68.53	8.22	56	82
Total	72	1.72	0.70	0.70	4.32	60.44	7.16	47	82

Table 2: Model summaries for metabolic rate estimates, including fish body mass and temperature (16, 19 and 22 °C) as independent variables. Batch is included as random effect. Brackets indicate standard error, *asterisks* indicates significance values with [*]p<0.05; [**]p<0.01; [***]p<0.001).

	log10(SMR)			log10(MMR)			log10(AS)		
	Est.	*CI*	*p*	*Est.*	*CI*	*p*	*Est.*	*CI*	*p*
log10(mass)	0.922	0.057	<0.001	0.978	0.120	<0.001	1.044	0.229	<0.001
19 °C	0.131	0.027	0.004	0.011	0.046	0.826	-0.038	0.086	0.674
22 °C	0.198	0.027	<0.001	0.040	0.046	0.424	-0.033	0.086	0.717
Intercept	-0.704	0.021	<0.001	-0.152	0.038	0.004	-0.342	0.072	<0.001
Random Effects									
Intercept SD (Batch)	0.0215			0.0116			0.0116		
Residual SD	0.0638			0.1492			0.1492		
Marginal R^2	0.849			0.521			0.521		
Conditional R^2	0.864			0.524			0.524		

Table 3: Post-hoc ANOVA contrasts from the results of the SMR model described in Table 2, comparing the temperature treatments 16 °C, 19 °C, and 22 °C. SE shows standard error, df degrees of freedom and the associated *t* ratio and *p* (significance) with *p<0.05; **p<0.01; ***p<0.001.

Comparison	Estimate	SE	df	t	p
16 °C - 19 °C	-0.13	0.03	4.66	-4.71	**0.014**
16 °C - 22 °C	-0.20	0.03	4.73	-7.42	**0.002**
19 °C - 22 °C	-0.07	0.03	4.94	-2.13	0.178

Table 4: Table showing predicted SMR (mg O_2 h^{-1}) and 95% confidence intervals (CI) for a mean mass fish (1.72g) at different temperatures.

Temperature (°C)	Predicted SMR (for a 1.72g fish)	Lower CI	Upper CI
16	0.33	0.24	0.44
19	0.44	0.32	0.58
22	0.52	0.39	0.70