Telomere-to-telomere reference genome of the common five-lined skink, *Plestiodon*

***fasciatus* (Squamata: Scincidae)**

Jon J. Hoffman[1,2]*

Frank T. Burbrink[1]

R. Alexander Pyron[1,3]

Christopher J. Raxworthy[1]

[1]Department of Herpetology, American Museum of Natural History, New York, NY 10024, USA;

[2]Richard Gilder Graduate School, American Museum of Natural History, New York, NY 10024,

USA

[3]Department of Biological Sciences, The George Washington University, Washington, DC

20052, USA

*Corresponding author: jhoffman1@amnh.org

Keywords: Hi-C; Whole Genome Sequencing; Plestiodon fasciatus; Scincidae; Scincinae

Abstract

Although the publication of high-quality reference genomes is steadily increasing for

vertebrate groups, many clades remain chronically neglected. Skinks (order: Squamata; family:

Scincidae) are one of the most diverse lizard families (1,785 species), yet there are currently just

four published chromosome-level skink genomes. Here, we present the first telomere-to-

telomere, chromosome-level reference genome for one of the most abundant lizards in the

eastern United States, the common five-lined skink (*Plestiodon fasciatus*). Through the

sequencing of RNA, long-read DNA, and Hi-C chromatin interactions, we produced an

annotated reference genome (N50 = 227MB, L50 = 3) consisting of 6 macrochromosome pairs

and 7 microchromosome pairs with 98% of BUSCO genes (lineage: sauropsida; 7480 BUSCO

markers) represented, providing one of the most complete skink genomes to date: *rPleFas1.1*.

Functional annotation predicts 32,520 protein-coding genes (16,100 unique, named genes) with

an average gene length of 9,372bp. Repeat annotations estimate that transposable elements

comprise 46.7% of the genome, for which we show the amount and content is remarkably

conserved across Scincidae.

Introduction

With the advent of whole-genome sequencing, numerous genomic resources are now

available for many clades across the vertebrate tree of life, with birds and mammals comprising

most of these genomes. Despite the abundance of ecological and evolutionary research on

squamates, the availability of high-quality genomes has, until recently, lagged far behind other

amniote groups until recently. There has been a recent drastic push to increase the production of

long-read genomes for many squamate groups (Gable et al., 2023). As a result, the number of

reference-level genomes of NCBI has risen from 115 in 2023 to 292 in Spring of 2025, a ~150%

increase over the last year and a half. Still, there is an underrepresentation of diverse families

such as skinks, geckos, chameleons, and amphisbaenids (Pinto et al., 2023). Generating high-

quality reference genomes across squamates is paramount, as broad sampling is an important

aspect of understanding and preserving a genetic record of biodiversity, hypothesizing species

relationships, and conservation of endangered taxa (Brandies et al., 2019; Worley et al., 2017)

For example, there is a large variation of chromosomal architecture within squamates,

with karyotypes ranging from 2n=16 in the gecko *Gonatodes taniae* to 2n = 62 in the microteiid

Notobrachia ablephara and the chameleon *Rieppeleon brevicaudatus* (Mezzasalma et al., 2024;

Pellegrino et al., 1999; Rovatsos et al., 2017; Schmid et al., 1994). In skinks, the number of

chromosomes ranges from 2n=22 to 2n=32, and the genomic architecture is relatively conserved

(Deweese and Wright, 1970; Giovannotti et al., 2009). Most *Plestiodon* have 13 chromosome

pairs, where 2n=26 (Xu and Zhu, 2024) with 6 macrochromosome pairs and 7 microchromosome

pairs; however, *Plestiodon anthracinus* has 12 chromosome pairs (Hardy et al., 2017).

There is also great diversity of sex-determination systems in squamates (Alam et al.,

2018; Ezaz et al., 2009; Janzen and Phillips, 2006), ranging from temperature-dependent sex

determination to genotypic sex determination with XY/XX (male heterogamy) and ZZ/ZW

(female heterogamy). Sex-determining systems are extremely labile in squamates (Ezaz et al.,

2009; Mezzasalma et al., 2021). For example, it is estimated that there have been 17–25

transitions in sex-determination in geckos, with some genera having multiple systems (Gamble et

al., 2015).

Unlike geckos but similar to iguanas, skinks have a conserved XY sex determining

system (male heterogamy) with difficult-to-distinguish homomorphic sex chromosomes

(Kostmann et al., 2021; Xu et al., 2024). There is little apparent variation in the sex-determining

system of skinks, but this may be an artifact of limited genomic resources (Dodge et al., 2023).

Due to its estimated 80-150Ma conservation in skinks, it has been hypothesized that the XY sex-

determination system may have evolved independently from that of other squamates, such as the

Podarcis mularis (Lacertidae) and *Anolis carolinensis* (Dactyloidae), which are commonly used

in comparative studies (Kostmann et al., 2021).

76   Despite harboring nearly 4% of amniote diversity, there are currently only four

chromosome-level skink genomes, three of which are Australian and all of which are in the

subfamily Lygosominae. The subfamilies Acontinae and Scincinae are not currently represented.

To increase both phylogenetic and biogeographic diversity of genomic resources for skinks, we

present a telomere-to-telomere, chromosome-level annotated reference genome for *Plestiodon*

fasciatus, the first distributed in the Nearctic and in the subfamily Scincinae. The common five-

lined skink, *Plestiodon fasciatus* (Linnaeus 1758), is one of the most abundant lizards of the

eastern U.S and southeastern Canada (Powell et al., 2016). These generalist skinks are small

(total length: 12.5–22.2 cm, maximum snout–vent length: 8.6 cm), found in moist wooded areas,

and reside in cover materials, such as rock crags, logs, and tree bark, but will emerge to

thermoregulate or search for invertebrate prey (Brazeau et al., 2015; Fitch and Fitch, 1954;

Powell et al., 2016). This genome is the fifth chromosome-level in the family Scincidae, which

currently includes *Bassiana duperreyi*, *Spondylurus nitidus, Carinascincus ocellatus,* and *Tiliqua*

scincoides, each of which shares a common ancestor with *Plestiodon* ~115Ma (Title et al., 2024).

Materials & Methods

Sample Acquisition

We collected an individual *Plestiodon fasciatus* in Allegan County, Michigan, and

euthanized with MS-222 prior to tissue collection following Michigan DNR permits and

approved IACUC protocols at the American Museum of Natural History (AMNH). We sampled

liver, lung, heart, skeletal muscle, kidney, and skin from the individual and stored them in NAP

buffer to preserve the RNA. Due to a suboptimal PacBio Revio whole-genome sequencing effort

of NAP preserved liver tissue from this individual, an individual female *Plestiodon fasciatus* was

collected in McCracken County, Kentucky, with Kentucky Department of Fish and Wildlife

permits to provide a blood sample (stored in EDTA) for genomic sequencing. Despite the 650

km between these localities, there are no major biogeographical barriers (Soltis et al., 2006)

Furthermore, there is little mitochondrial phylogeographic structure within the population

(Howes and Lougheed, 2008) and therefore likely little genomic variation, though this has not

been tested. Both individuals are vouchered specimens in the Herpetology Collections at the

American Museum of Natural History (AMNH) as AMNH-179334 (Allegan County, MI) and

AMNH-179327 (McCracken County, KY).

RNA extraction and sequencing

For RNA sequencing, we sent six tissues (liver, lung, heart, skeletal muscle, kidney, and

skin) to Azenta/Genewiz for extraction, library preparation with poly(A) selection to target

eukaryotic strand-specific mRNA, and sequencing on an Illumina NovaSeq 2x150bp, generating

~100M paired-end reads. All RNA sequences passed the initial quality check run with FastQC

(Brown et al., 2017). Sequencing adapters in the resulting sequences were filtered and trimmed

with trimmomatic using the default settings (Bolger et al., 2014).

 Genomic DNA extraction and sequencing

Genomic DNA was extracted from blood stored in EDTA with the Qiagen MagAttract

High Molecular Weight DNA kit following their 'Manual Purification of High-Molecular Weight

Genomic DNA from Whole Blood' protocol from the MagAttract HMW DNA Handbook in the

AMNH ICG. The extracted sample was then sent to Azenta/Genewiz, where it was sequenced

with PacBio Revio HiFi sequencing on 1 SMRT cell, which typically results in ~15 million reads

and ~100GB of data, depending on the quality of the input sample. We expected a coverage of

~66x based on an estimated 1.5GB genome size. The sequencing generated ~6.3 million read

pairs and ~79GB of data, with an average read length of ~12,000bp, resulting in ~53x coverage.

We generated Hi-C data with Phase Genomics using the Proximo® Hi-C animal genome

scaffolding platform from a collected blood sample. Proximity-ligated fragments were sequenced

on an Illumina NovaSeq to produce paired-end reads. The sequencing generated 300 million

read-pairs, of which 56% were high quality, yielding 1,721,863 read-pairs per contig that were

'usable' and mapped to different >5kb contigs.

Draft Genome assembly

A draft genome assembly was made from the HiFi long-read sequencing using Hifiasm, a

haplotype-resolved de-novo assembly tool for PacBio HiFi reads (Cheng et al., 2021). Hifiasm

was run without the Hi-C sequences as its inclusion led to a drastic increase of contigs, from 34

without Hi-C to ~6,000 with Hi-C, potentially due to the low-divergence repeat content of the

genome. Due to the high content of low-divergence repeats, we soft-masked the draft genome

with the *Earlgrey* pipeline (Baril et al., 2024) prior to incorporating the Hi-C data.

Hi-C incorporation

To incorporate the Hi-C sequencing with the draft genome assembly, we initially aligned

the Hi-C sequences to the HiFi draft genome with a Burrow-Wheeler Alignment (BWA) (Li,

2013). Then, we processed the resulting alignments with *SAMtools* (Danecek et al., 2021) to

remove duplicate sequences. We then scaffolded the assembly using the *de novo YAHS* assembly

pipeline (Zhou et al., 2023). From there, we used Juicer Tools to generate a Hi-C contact map

(Durand et al., 2016). Finally, we visualized the scaffolding of the chromosome-level assembly

with Juicebox Assembly Tools (Robinson et al., 2018).

TE and gene annotation

We annotated the final assembly by first modelling and quantifying transposable elements

(TEs) by soft-masking the genome with *EarlGrey* v4.1.1, a fully automated TE annotation

pipeline (Baril et al., 2024) with the default settings, which includes a 100bp minimum length

and 10 iterations of the BLAST, extract, and extend process. Next, we functionally annotated the

predicted gene regions using the train, predict, update, fix, and annotate steps of the *funannotate*

pipeline (Palmer and Stajich, 2020). The 'train' step aligns RNA-seq data, assembles it with

Trinity (Grabherr et al., 2011), and runs *PASA*, which models gene structures based on alignments

of expressed transcripts (Haas et al., 2003). The 'predict' step uses *PASA* gene models to train

Augustus, a *de novo* gene finder (Stanke et al., 2008), prior to running *EvidenceModeler* (Haas et

al., 2008). We included RNA-seq data from the liver, lung, heart, skeletal muscle, kidney, and

skin as evidence for *EvidenceModeler*. We used the Tetrapoda BUSCO database with

Taeniopygia guttata as the seed species for the 'predict' step and kept the default options for each

step. The 'update' step fixes gene models that disagree with RNA-seq data, which are corrected

in the 'fix' step. Prior to running the 'annotate' step, we ran *InterProScan* (Jones et al., 2014) to

run predicted genes against the InterPro database for gene families and downloaded *Eggnog-*

mapper (Cantalapiedra et al., 2021) locally to be run during the 'annotate' step. The 'annotate'

step incorporates the generated data into an annotated genome. We used the default functional

annotation databases in the annotation step.

Synteny analysis

To review genomic synteny of the *Plestiodon fasciatus* genome and other chromosome-

level squamate assemblies, we created a custom pipeline called *Synk*

(https://github.com/jomhoff/Synk) that uses the output files from *compleasm* (Huang and Li,

2023) and isolates the BUSCO genes to create comparative text files, and uses *RIdeogram* (Hao

et al., 2020) to plot the syntenic chromosomal regions from BUSCO genes between species in

one script. Here, we show synteny between *P. fasciatus*, *Bassiana duperreyi*, *Spondylurus*

nitidus, Carinascincus ocellatus, and *Tiliqua scincoides.*

Results and Discussion

Assembly of the P. fasciatus genome

After completing *Hifiasm* with the long-read PacBio sequences, the draft assembly of the

genome was close to complete, with an L50 of 3 and a total of 34 contigs, 9 of which represented

near telomere-to-telomere chromosomes (**Table 1**). With the Hi-C data incorporated, we

successfully generated a chromosome-level assembly of *Plestiodon fasciatus* with 18 unplaced

scaffolds that we refer to as *rPleFas1.1* after Vertebrate Genome Project naming rules (Rhie et al.

2021) (**Table 1; Figure 1**). One of the 18 unplaced scaffolds includes the complete

mitochondrial genome. The other 17 unplaced scaffolds range in size from 8,262bp to

1,241,752bp and consist of unplaced TEs and mRNA sequences. For the complete automsomal

genome, the presence of telomeres was estimated with the characteristic rise in GC content at the

ends of chromosomes (**Figure 1**) and confirmed with *tidk*, a toolkit for identifying telomeres

(Brown et al., 2025). There is also a large spike in GC content in the middle of the second

chromosome, which corresponds to a repeat-dense region. This is the first skink genome to

include all chromosomes with completed telomeres, as well as the highest BUSCO scores

(*Plestiodon fasciatus:* 98.17%; *Bassiana duperreyi*: 97.21%; *Spondylurus nitidus:* 98.05%;

Carinascincus ocellatus: 97.54%; *Tiliqua scincoides*: 94.97%)

	Metric	Draft Genome	Hi-C Informed Genome
Basic Stats	Total assembly size (bp)	1,537,706,373	1,537,706,373
	Number of contigs	34	34
	Number of scaffolds	34	31
	GC content (%)	45.6	45.6
Contiguity	Contigs N50 (bp)	166,278,287	166,278,287
	Contigs L50 (bp)	3	3
	Largest Contig (bp)	244,253,184	244,253,184
	Scaffolds N50 (bp)	166,278,287	227,050,938
	Scaffolds L50	3	2
	Largest Scaffold (bp)	244,253,184	304,269,845
Completeness	BUSCO Complete (%)	98.17	98.17
	Missing BUSCOs (%)	1.24	1.24
	Sequencing coverage (X)	53	116 or 53??
	Telomere to Telomere Chromosomes	6	13

Table 1: Assembly statistics of the genome before and after Hi-C incorporation

Annotation of protein-coding genes and transposable elements in the P. fasciatus genome

The resulting annotating from the *funannotate* pipeline consists of an estimated 32,520

genes. Filtering for named genes and removing isoforms resulted in 16,100 unique genes with

common names (**Table 2**). Like other squamates, the genome of *P. fasciatus* has a high

proportion of repetitive regions which account for 46.7% of the genome (Pasquesi et al., 2018).

23.3% of genome is unclassified repeats, while DNA repeats and Long Interspersed Nuclear

Elements (LINEs) respectively make up 7.0% and 9.3% of the genome (**Figure 2**). The Kimura

2-parameter distance between the repeat sequences produces an approximately bimodal

distribution, with a peak indicating a large degree of moderately diverged repeats and a peak

indicating recently diverged repeats (**Figure 2**). The more diverged peak is mainly driven by

unclassified repeats. Interestingly, this a pattern that is seen in all five skinks tested. Furthermore,

the more recently diverged peak is driven by a large proportion of both DNA and LINE repeats

and is also conserved across the species analyzed (**Figure 2**).

Although the impact of TEs on the diversification and adaptation of squamates has yet to

be studied in detail, it has been hypothesized that variation in TEs can impact phenotypic

adaptation through TE domestication, exaptation, host-gene regulation, formation of retrogenes,

and genomic plasticity (reviewed in Schrader and Schmitz, 2019), and that TEs contribute

significantly to genomic variation (Catlin and Josephs, 2022). Since TEs are highly mobile

across the genome, are abundant, labile, and have been major players in the evolution of

eukaryotic genomes (Bowen and Jordan, 2002; Oliver and Greene, 2009), we expect variation of

TEs among distantly diverged clades. Despite originating around ~115mya, the skinks in this

study display remarkable conservation of TE composition, especially when considering the

variation seen in younger clades, such as extant mammals (Platt et al., 2018) and plethodontid

salamanders (Sun et al., 2012).

Category	Metric	Value
Gene Content	Number of protein-coding genes	32,520
	Unique genes with common names	16,100
Gene Structure	Average gene length (bp)	9,372.34
	Total number of exons	252,300
	Average exon length (bp)	255.27
	Multiple-exon transcripts	25,572
	Single-exon transcripts	7,372
Functional Annotation	Genes with GO term annotations	21,578
	Genes with InterProScan hits	24,316
	Genes with eggNOG annotations	26,007
	Genes with Pfam domains	19,449
	CAZyme-annotated genes	319
	MEROPS-annotated genes	1,143

Table 2: Annotation Statistics of the Genome after the *Funannotate* pipeline

Synteny of scincid genomes

Macrochromosomes are conserved across the five skink species analyzed, aside from a large inversion on chromosome 1 between *Tiliqua scincoides* and all other species (**Figure 3**). Compared to the other scincids, *P. fasciatus* has fewer microchromosomes, where chromosomes 14, 15, and 16 in *S. nitidus* and *T. scincoides* are syntenic with chromosome 8 in *P. fasciatus.*

There is also a rearrangement creating syntenic blocks relating chromosomes 6 and 7 in *S.*

nitidus and *T. scincoides* to chromosome 5 in *P. fasciatus*.

Due to the cryptic, homogametic XY sex-determining system in skinks, little is known

about the position of sex-determining regions on the chromosomes; however, hypothetical sex

chromosomes have been identified in *Bassiana duperryi* (Dissanayake et al., 2020; Hanrahan et

al., 2025). Here, we show that a block of chromosome 5 is syntenic with the X chromosome in *B.*

duperreyi, indicating a potential location for sex-determining regions in *P. fasciatus* (**Figure 3**).

Despite some preliminary species-specific work, more research is required to further classify

sex-determination in skinks, ideally with population-level sampling of populations with

numerous representatives from both sexes across Scincidae.

Conclusion

We present a high-quality, telomere-to-telomere, annotated chromosome-level assembly

of the North American common five-lined skink *Plestiodon fasciatus* (Linnaeus 1758),

representing one of the most complete reference genomes (*rPleFas1.1*) to date of any species of

Scincidae, and the first for the subfamily Scincinae. We find that macrochromosome structure is

conserved across the family, but there are common rearrangements of the microchromosomes,

including a likely fusion in *P. fasciatus*, which has fewer microchromosomes than many other

skink species. We also present insight into the content and evolution of transposable elements in

skinks, which show remarkable conservation between species over the last ~115Ma.

Data availability

The data presented in the paper is available on NCBI (BioProject PRJNA1278702) and Figshare

(https://figshare.com/projects/Telomere-to-telomere_reference_genome_of_the_common_five-

lined_skink_Plestiodon_fasciatus_Squamata_Scincidae_/254723). The code for analyses can be

found at https://github.com/jomhoff/Chromosome-level_genome_assembly and

https://github.com/jomhoff/Genome-Annotation.

Acknowledgments

The authors thank Megan Wallace of the American Museum of Natural History's Institute of

Comparative Genomics for her laboratory expertise, as well as Mac Mahacek, Steven Price,

William Taylor, and Elijah Wessel for field assistance. The authors also thank the Michigan

Department of Natural Resources and Kentucky Department of Fish and Wildlife.

Funding

This work was funded in part by the American Museum of Natural History's Richard Gilder

Graduate School, the Theodore Roosevelt Memorial Fund, and the Society for the Study of

Amphibians and Reptiles' Roger Conant Grants-in-Herpetology. This research was supported in

part by the U.S. National Science Foundation: DBI 2029955 awarded to CJR and DEB 2323125

to FTB.

Figure Captions:

Figure 1: Ideogram for *Plestiodon fasciatus*. **Row A** represents chromosome number
and relative size. **Row B** is a heatmap of gene density for 50kb windows, where darker
colors represent a higher density of genes. **Row C** is a heatmap of repeat elements for
50kb windows, where darker colors represent a higher density of repeat elements. **Row
D** is a line chart of GC content across the genome. **E** is the the region of the genome
that is syntenic with the X chromosome of *Bassiani duperreyi*. **F** is a photograph of an
adult male and juvenile *Plestiodon fasciatus* (photo via Herps of Arkansas). Plot made
with Circos (Krzywinski et al., 2009).

278
279 **Figure 2:** Divergence estimates of TEs shown with Kimura 2-Parameter Distance for all
280 five chromosome-level skink genomes where the higher the Kimura 2-Parameter
281 Distance, the more diverged the sequences. The proportion of each transposable
282 element subdivision in the genome for each species is displayed with a pie chart.
283
284 **Figure 3:** Synteny of BUSCO genes between *Plestiodon fasciatus* and other lizards.
285 **From top to bottom**: *P. fasciatus* and *Tiliqua scincoides; P. fasciatus* and *Spondylurus*
286 *nitidus; P. fasciatus* and *Carinascincus ocellatus; P. fasciatus* and *Bassiana duperreyi*.
287

References

Alam, S.M.I., Sarre, S.D., Gleeson, D., Georges, A., Ezaz, T., 2018. Did Lizards Follow
Unique Pathways in Sex Chromosome Evolution? Genes (Basel) 9, 239.
https://doi.org/10.3390/genes9050239
Baril, T., Galbraith, J., Hayward, A., 2024. Earl Grey: A Fully Automated User-Friendly
Transposable Element Annotation and Analysis Pipeline. Molecular Biology and
Evolution 41, msae068. https://doi.org/10.1093/molbev/msae068
Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170
Bowen, N.J., Jordan, I.K., 2002. Transposable elements and the evolution of eukaryotic
complexity. Curr Issues Mol Biol 4, 65–76.
Brandies, P., Peel, E., Hogg, C.J., Belov, K., 2019. The Value of Reference Genomes in
the Conservation of Threatened Species. Genes 10, 846.
https://doi.org/10.3390/genes10110846
Brazeau, D., Freitag, R., Hecnar, S.J., Hecnar, D.R., 2015. Comparing Common Five-
lined Skink (Plestiodon fasciatus) Diet Among Populations and Time. |
EBSCOhost [WWW Document]. URL
https://openurl.ebsco.com/contentitem/gcd:115178012?sid=ebsco:plink:crawler&i
d=ebsco:gcd:115178012 (accessed 6.11.25).
Brown, J., Pirrung, M., McCue, L.A., 2017. FQC Dashboard: integrates FastQC results
into a web-based, interactive, and extensible FASTQ quality control tool.
Bioinformatics 33, 3137–3139. https://doi.org/10.1093/bioinformatics/btx373
Brown, M.R., Manuel Gonzalez de La Rosa, P., Blaxter, M., 2025. tidk: a toolkit to
rapidly identify telomeric repeats from genomic datasets. Bioinformatics 41,
btaf049. https://doi.org/10.1093/bioinformatics/btaf049
Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., Huerta-Cepas, J., 2021.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Molecular Biology and Evolution 38, 5825–
5829. https://doi.org/10.1093/molbev/msab293
Catlin, N.S., Josephs, E.B., 2022. The important contribution of transposable elements
to phenotypic variation and evolution. Current Opinion in Plant Biology 65,
102140. https://doi.org/10.1016/j.pbi.2021.102140

Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., Li, H., 2021. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. Nat Methods 18, 170–175. https://doi.org/10.1038/s41592-020-01056-5

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and BCFtools. GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008

Deweese, J.E., Wright, J.W., 1970. A preliminary karyological analysis of scincid lizards. Mammalian Chromosomes Newsletter 11, 95–96.

Dissanayake, D.S.B., Holleley, C.E., Hill, L.K., O'Meally, D., Deakin, J.E., Georges, A., 2020. Identification of Y chromosome markers in the eastern three-lined skink (Bassiana duperreyi) using in silico whole genome subtraction. BMC Genomics 21, 667. https://doi.org/10.1186/s12864-020-07071-2

Dodge, T.O., Farquharson, K.A., Ford, C., Cavanagh, L., Schubert, K., Schumer, M., Belov, K., Hogg, C.J., n.d. Genomes of two Extinct-in-the-Wild reptiles from Christmas Island reveal distinct evolutionary histories and conservation insights. Molecular Ecology Resources n/a. https://doi.org/10.1111/1755-0998.13780

Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S., Aiden, E.L., 2016. Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst 3, 95–98. https://doi.org/10.1016/j.cels.2016.07.002

Ezaz, T., Sarre, S.D., O'Meally, D., Graves, J.A.M., Georges, A., 2009. Sex chromosome evolution in lizards: independent origins and rapid transitions. Cytogenet Genome Res 127, 249–260. https://doi.org/10.1159/000300507

Fitch, H.S., Fitch, H.S., 1954. Life history and ecology of the five-lined skink, Eumeces fasciatus. University of Kansas publications, Museum of Natural History 8, 1--156.

Gable, S.M., Mendez, J.M., Bushroe, N.A., Wilson, A., Byars, M.I., Tollis, M., 2023. The State of Squamate Genomics: Past, Present, and Future of Genome Research in the Most Speciose Terrestrial Vertebrate Order. Genes 14, 1387. https://doi.org/10.3390/genes14071387

Gamble, T., Coryell, J., Ezaz, T., Lynch, J., Scantlebury, D.P., Zarkower, D., 2015. Restriction Site-Associated DNA Sequencing (RAD-seq) Reveals an Extraordinary Number of Transitions among Gecko Sex-Determining Systems. Molecular Biology and Evolution 32, 1296–1309. https://doi.org/10.1093/molbev/msv023

Giovannotti, M., Caputo, V., O'Brien, P.C.M., Lovell, F.L., Trifonov, V., Cerioni, P.N., Olmo, E., Ferguson-Smith, M.A., Rens, W., 2009. Skinks (Reptilia: Scincidae) have highly conserved karyotypes as revealed by chromosome painting. Cytogenet Genome Res 127, 224–231. https://doi.org/10.1159/000295002

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A., 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol 29, 644–652. https://doi.org/10.1038/nbt.1883

Haas, B.J., Delcher, A.L., Mount, S.M., Wortman, J.R., Smith, R.K., Hannick, L.I., Maiti, R., Ronning, C.M., Rusch, D.B., Town, C.D., Salzberg, S.L., White, O., 2003. Improving the Arabidopsis genome annotation using maximal transcript alignment assemblies. Nucleic Acids Res 31, 5654–5666. https://doi.org/10.1093/nar/gkg770

Haas, B.J., Salzberg, S.L., Zhu, W., Pertea, M., Allen, J.E., Orvis, J., White, O., Buell, C.R., Wortman, J.R., 2008. Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. Genome Biology 9, R7. https://doi.org/10.1186/gb-2008-9-1-r7

Hanrahan, B.J., Alreja, K., Reis, A.L.M., Chang, J.K., Dissanayake, D.S.B., Edwards, R.J., Bertozzi, T., Hammond, J.M., O'Meally, D., Deveson, I.W., Georges, A., Waters, P., Patel, H.R., 2025. A genome assembly and annotation for the Australian alpine skink Bassiana duperreyi using long-read technologies. G3 Genes|Genomes|Genetics 15, jkaf046. https://doi.org/10.1093/g3journal/jkaf046

Hao, Z., Lv, D., Ge, Y., Shi, J., Weijers, D., Yu, G., Chen, J., 2020. RIdeogram: drawing SVG graphics to visualize and map genome-wide data on the idiograms. PeerJ Comput. Sci. 6, e251. https://doi.org/10.7717/peerj-cs.251

Harding, J.H., 1997. Amphibians and Reptiles of the Great Lakes Region. University of Michigan Press.

Hardy, L.M., Raymond, L.R., Harris, S., 2017. The Karyotype of Plestiodon anthracinus (Baird, 1850) (Sauria: Scincidae): A Step Toward Solving an Enigma. sena 16, 326–330. https://doi.org/10.1656/058.016.0318

Howes, B.J., Lougheed, S.C., 2008. Genetic diversity across the range of a temperate lizard. Journal of Biogeography 35, 1269–1278. https://doi.org/10.1111/j.1365-2699.2007.01867.x

Huang, N., Li, H., 2023. compleasm: a faster and more accurate reimplementation of BUSCO. Bioinformatics 39, btad595. https://doi.org/10.1093/bioinformatics/btad595

Janzen, F.J., Phillips, P.C., 2006. Exploring the evolution of environmental sex determination, especially in reptiles. J Evol Biol 19, 1775–1784. https://doi.org/10.1111/j.1420-9101.2006.01138.x

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., Hunter, S., 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics 30, 1236–1240. https://doi.org/10.1093/bioinformatics/btu031

Kostmann, A., Kratochvíl, L., Rovatsos, M., 2021. Poorly differentiated XX/XY sex chromosomes are widely shared across skink radiation. Proceedings of the Royal Society B: Biological Sciences 288, 20202139. https://doi.org/10.1098/rspb.2020.2139

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM [WWW Document]. arXiv.org. URL https://arxiv.org/abs/1303.3997v2 (accessed 6.11.25).

Mezzasalma, M., Guarino, F.M., Odierna, G., 2021. Lizards as Model Organisms of Sex Chromosome Evolution: What We Really Know from a Systematic Distribution of Available Data? Genes 12, 1341. https://doi.org/10.3390/genes12091341

Mezzasalma, M., Macirella, R., Odierna, G., Brunelli, E., 2024. Karyotype Diversification
and Chromosome Rearrangements in Squamate Reptiles. Genes (Basel) 15,
371. https://doi.org/10.3390/genes15030371
Oliver, K.R., Greene, W.K., 2009. Transposable elements: powerful facilitators of
evolution. BioEssays 31, 703–714. https://doi.org/10.1002/bies.200800219
Palmer, J.M., Stajich, J., 2020. Funannotate v1.8.1: Eukaryotic genome annotation.
https://doi.org/10.5281/zenodo.4054262
Pasquesi, G.I.M., Adams, R.H., Card, D.C., Schield, D.R., Corbin, A.B., Perry, B.W.,
Reyes-Velasco, J., Ruggiero, R.P., Vandewege, M.W., Shortt, J.A., Castoe, T.A.,
2018. Squamate reptiles challenge paradigms of genomic repeat element
evolution set by birds and mammals. Nat Commun 9, 2774.
https://doi.org/10.1038/s41467-018-05279-1
Pellegrino, K.C., Rodrigues, M.T., Yonenaga-Yassuda, Y., 1999. Chromosomal
polymorphisms due to supernumerary chromosomes and pericentric inversions
in the eyelidless microteiid lizard Nothobachia ablephara (Squamata,
Gymnophthalmidae). Chromosome Res 7, 247–254.
https://doi.org/10.1023/a:1009218628942
Pinto, B.J., Gamble, T., Smith, C.H., Wilson, M.A., 2023. A lizard is never late:
Squamate genomics as a recent catalyst for understanding sex chromosome and
microchromosome evolution. Journal of Heredity 114, 445–458.
https://doi.org/10.1093/jhered/esad023
Platt, R.N., Vandewege, M.W., Ray, D.A., 2018. Mammalian transposable elements and
their impacts on genome evolution. Chromosome Res 26, 25–43.
https://doi.org/10.1007/s10577-017-9570-z
Powell, R., Conant, R., Collins, J.T., 2016. A Field Guide to Reptiles & Amphibians:
Eastern and Central North America. Houghton Mifflin Harcourt.
Robinson, J.T., Turner, D., Durand, N.C., Thorvaldsdóttir, H., Mesirov, J.P., Aiden, E.L.,
2018. Juicebox.js Provides a Cloud-Based Visualization System for Hi-C Data.
Cell Syst 6, 256-258.e1. https://doi.org/10.1016/j.cels.2018.01.001
Rovatsos, M., Altmanová, M., Johnson Pokorná, M., Velenský, P., Sánchez Baca, A.,
Kratochvíl, L., 2017. Evolution of Karyotypes in Chameleons. Genes (Basel) 8,
382. https://doi.org/10.3390/genes8120382
Schmid, M., Feichtinger, W., Nanda, I., Schakowski, R., Visbal Garcia, R., Manzanilla
Puppo, J., Fernández Badillo, A., 1994. An extraordinarily low diploid
chromosome number in the reptile Gonatodes taniae (Squamata, Gekkonidae). J
Hered 85, 255–260. https://doi.org/10.1093/oxfordjournals.jhered.a111452
Schrader, L., Schmitz, J., 2019. The impact of transposable elements in adaptive
evolution. Molecular Ecology 28, 1537–1549. https://doi.org/10.1111/mec.14794
Seburn, C.N.L., 1993. Spatial distribution and microhabitat use in the five-lined skink
(Eumeces fasciatus). Can. J. Zool. 71, 445–450. https://doi.org/10.1139/z93-064
Soltis, D.E., Morris, A.B., McLACHLAN, J.S., Manos, P.S., Soltis, P.S., n.d. Comparative
phylogeography of unglaciated eastern North America.
Stanke, M., Diekhans, M., Baertsch, R., Haussler, D., 2008. Using native and
syntenically mapped cDNA alignments to improve de novo gene finding.
Bioinformatics 24, 637–644. https://doi.org/10.1093/bioinformatics/btn013

Sun, C., Shepard, D.B., Chong, R.A., López Arriaza, J., Hall, K., Castoe, T.A.,
Feschotte, C., Pollock, D.D., Mueller, R.L., 2012. LTR retrotransposons
contribute to genomic gigantism in plethodontid salamanders. Genome Biol Evol
4, 168–183. https://doi.org/10.1093/gbe/evr139
Title, P.O., Singhal, S., Grundler, M.C., Costa, G.C., Pyron, R.A., Colston, T.J., Grundler,
M.R., Prates, I., Stepanova, N., Jones, M.E.H., Cavalcanti, L.B.Q., Colli, G.R., Di-
Poï, N., Donnellan, S.C., Moritz, C., Mesquita, D.O., Pianka, E.R., Smith, S.A.,
Vitt, L.J., Rabosky, D.L., 2024. The macroevolutionary singularity of snakes.
Science 383, 918–923. https://doi.org/10.1126/science.adh2449
Worley, K.C., Richards, S., Rogers, J., 2017. The value of new genome references.
Experimental Cell Research 358, 433–438.
https://doi.org/10.1016/j.yexcr.2016.12.014
Xu, W., Li, T., Li, J., Liu, S., Yu, X., Tang, M., Dong, J., Liu, J., Bu, X., Xia, X., Zhou, H.,
Nie, L., 2024. The First Identification of Homomorphic XY Sex Chromosomes by
Integrating Cytogenetic and Transcriptomic Approaches in Plestiodon elegans
(Scincidae). Genes 15, 664. https://doi.org/10.3390/genes15060664
Xu, W., Zhu, W., 2024. First karyological analysis of Plestiodon capito (Squamata:
Scincidae). Cytologia 89, 147–151. https://doi.org/10.1508/cytologia.89.147
Zhou, C., McCarthy, S.A., Durbin, R., 2023. YaHS: yet another Hi-C scaffolding tool.
Bioinformatics 39, btac808. https://doi.org/10.1093/bioinformatics/btac808
478