Telomere-to-telomere reference genome of the common five-lined skink, *Plestiodon*

***fasciatus* (Squamata: Scincidae)**

Jon J. Hoffman[1,2*]

Frank T. Burbrink[1]

R. Alexander Pyron[1,3]

Christopher J. Raxworthy[1]

[1]Department of Herpetology, American Museum of Natural History, New York, NY 10024, USA;

[2]Richard Gilder Graduate School, American Museum of Natural History, New York, NY 10024,

USA

[3]Department of Biological Sciences, The George Washington University, Washington, DC

20052, USA

*Corresponding author: jhoffman1@amnh.org

Keywords: Hi-C; Whole Genome Sequencing; Plestiodon fasciatus; Scincidae; Scincinae

Abstract

Although the publication of high-quality reference genomes is steadily increasing, many

clades remain chronically neglected. Skinks (order: Squamata; family: Scincidae) are one of the

most diverse lizard families (1,785 species), yet there are currently just six published

chromosome-level skink genomes. Here, we present the first telomere-to-telomere, chromosome-

level reference genome for one of the most abundant lizards in the eastern United States, the

common five-lined skink (*Plestiodon fasciatus*). Through the sequencing of RNA, long-read

DNA, and Hi-C chromatin interactions, we produced an annotated reference genome (N50 =

227MB, L50 = 3) consisting of 6 macrochromosome pairs and 7 microchromosome pairs with

98% of BUSCO genes represented (lineage: Sauropsida; 7480 BUSCO markers) represented,

providing one of the most complete skink genomes to date: *rPleFas1.1*. Functional annotation

predicts 32,520 protein-coding genes (16,100 unique, named genes) with an average gene length

of 9,372bp. Repeat annotations estimate that transposable elements comprise 46.7% of the

genome, for which we show the amount and content is remarkably conserved across Scincidae.

Introduction

With the advent of whole-genome sequencing, numerous genomic resources are now

available for many clades across the vertebrate tree of life, with birds and mammals comprising

most of these genomes. Despite the abundance of ecological and evolutionary research on

squamate reptiles, the availability of high-quality genomes has, until recently, lagged far behind

other amniote groups. There has been a recent drastic push to increase the production of long-

read genomes for many squamate groups (Gable et al., 2023). As a result, the number of

reference-level genomes of NCBI has risen from 115 in 2023 to 292 in Spring of 2025, a ~150%

increase over the last year and a half. Still, there is an underrepresentation of diverse families

such as skinks, geckos, chameleons, and amphisbaenids (Pinto et al., 2023). Generating high-

quality reference genomes across squamates is paramount, as broad sampling is an important

aspect of understanding and preserving a genetic record of biodiversity, hypothesizing species

relationships, and conservation of endangered taxa (Brandies et al., 2019; Worley et al., 2017)

For example, there is a large variation of chromosomal architecture within squamates,

with karyotypes ranging from 2n=16 in the gecko *Gonatodes taniae* to 2n = 62 in the microteiid

Notobrachia ablephara and the chameleon *Rieppeleon brevicaudatus* (Mezzasalma et al., 2024;

Pellegrino et al., 1999; Rovatsos et al., 2017; Schmid et al., 1994). In skinks, the number of

chromosomes ranges from 2n=22 to 2n=32, and genomic architecture is relatively conserved

(Deweese and Wright, 1970; Giovannotti et al., 2009). Most *Plestiodon* have 13 chromosome

pairs, where 2n=26 with 6 macrochromosome pairs and 7 microchromosome pairs (Xu and Zhu,

2024), though *P. anthracinus* has 12 chromosome pairs (Hardy et al., 2017).

There is also great diversity of sex-determination systems in squamates (Alam et al.,

2018; Ezaz et al., 2009; Janzen and Phillips, 2006), ranging from temperature-dependent sex

determination to genotypic sex determination with XY/XX (male heterogamy) and ZZ/ZW

(female heterogamy). Sex-determining systems are extremely labile in squamates (Ezaz et al.,

2009; Mezzasalma et al., 2021). For example, it is estimated that there have been 17–25 in sex-

determination transitions in geckos, with multiple sex-determining systems in some genera

(Gamble et al., 2015). Unlike geckos but similar to iguanas, skinks have a conserved XY sex

determining system (male heterogamy) with homomorphic sex chromosomes that are difficult to

distinguish (Kostmann et al., 2021; Xu et al., 2024). There is little apparent variation in the sex-

determining system of skinks, but this may be an artifact of limited genomic resources (Dodge et

al., 2023). It has been hypothesized that the XY sex-determination system may have evolved

independently from that of other squamates, such as *Podarcis mualris* (Lacertidae) and *Anolis*

carolinensis (Dactyloidae), which are commonly used in comparative studies (Kostmann et al.,

2021).

Skinks represent 4% of amniote diversity, yet there are currently only six chromosome-

level skink genomes, four of which are Australian. To increase both the phylogenetic and

biogeographic diversity of genomic resources for skinks, we present a telomere-to-telomere

annotated reference genome for *Plestiodon fasciatus*. The common five-lined skink, *P. fasciatus*

(Linnaeus 1758), is one of the most abundant lizards of the eastern U.S and southeastern Canada

(Powell et al., 2016). These generalist skinks are small (total length: 12.5–22.2 cm, maximum

snout–vent length: 8.6 cm), found in mesic wooded areas, and reside in cover materials, such as

rock crags, logs, and tree bark, but will emerge to thermoregulate or search for invertebrate prey

(Brazeau et al., 2015; Fitch and Fitch, 1954; Powell et al., 2016). This genome is the seventh

chromosome-level in the family Scincidae, which currently includes *Bassiana duperreyi*

(Hanrahan et al., 2025), *Spondylurus nitidus* (Rivera et al., 2024)*, Carinascincus ocellatus,*

Tiliqua scincoides, Cryptoblepharus egeriae (Dodge et al., 2025), and the congener *P. gilberti*

(Richmond et al., 2025)*.* Barring *P. gilberti*, all of the other species represented which shares a

common ancestor with *P. fasciatus* ~115Ma (Title et al., 2024), while *P. gilberti* and *P. fasciatus*

diverged ~17Ma (Brandley et al., 2011)

Materials & Methods

Sample Acquisition

We collected an individual female *Plestiodon fasciatus* in Allegan County, Michigan

(Lat: 42.5394; Long: -85.9949), and euthanized with MS-222 prior to tissue collection following

Michigan DNR permits and approved IACUC protocols at the American Museum of Natural

History (AMNH). We sampled liver, lung, heart, skeletal muscle, kidney, and skin from the

individual and stored them in NAP buffer (Camacho-Sanchez et al., 2013) to preserve the RNA.

Due to a suboptimal PacBio Revio whole-genome sequencing effort of the NAP preserved liver

tissue from this individual, we collected a female *P. fasciatus* in McCracken County, Kentucky

(Lat: 37.1501; Long: -88.7953), with Kentucky Department of Fish and Wildlife permits to

provide a blood sample (stored in EDTA) for genomic sequencing. Despite the 650 km between

these localities, there are no major biogeographical barriers (Soltis et al., 2006). Furthermore,

there is little mitochondrial phylogeographic structure within the population (Howes and

Lougheed, 2008) and therefore likely little genomic variation, though this has not been tested.

Both individuals are vouchered specimens, catalogued in the Herpetology Collections at the

American Museum of Natural History (AMNH) as AMNH-179334 (Allegan County, MI) and

AMNH-179327 (McCracken County, KY).

RNA extraction and sequencing

For RNA sequencing, we sent six tissues (liver, lung, heart, skeletal muscle, kidney, and

skin) to Azenta/Genewiz for extraction, library preparation with poly(A) selection to target

eukaryotic strand-specific mRNA, and sequencing on an Illumina NovaSeq 2x150bp, generating

~100M paired-end reads. All RNA sequences passed the initial quality check run with FastQC

(Brown et al., 2017). Sequencing adapters in the resulting sequences were filtered and trimmed

with trimmomatic using the default settings (Bolger et al., 2014).

Genomic DNA extraction and sequencing

Genomic DNA was extracted from blood stored in EDTA from AMNH-179327 using the Qiagen MagAttract High Molecular Weight DNA kit following their 'Manual Purification of High-Molecular Weight Genomic DNA from Whole Blood' protocol from the MagAttract HMW DNA Handbook in the AMNH ICG. The extracted DNA was then sent to Azenta/Genewiz, where it was sequenced with PacBio Revio HiFi sequencing on 1 SMRT cell, which typically results in ~15 million reads and ~100GB of data, depending on the quality of the input sample. We expected a coverage of ~66x based on an estimated 1.5GB genome size. Sequencing generated ~6.3 million read pairs and ~79GB of data, with an average read length of ~12,000bp, resulting in ~53x coverage.

Finally, we generated Hi-C data with Phase Genomics using the Proximo® Hi-C animal genome scaffolding platform from a collected blood sample. Proximity-ligated fragments were sequenced on an Illumina NovaSeq to produce 2x150bp paired-end reads. The sequencing generated 300 million read-pairs, of which 56% were high quality, yielding 1,721,863 read-pairs per contig that were 'usable', indicating that they mapped to different >5kb contigs.

Draft Genome assembly

A draft genome assembly was made from the HiFi long-read sequencing using Hifiasm v0.25.0, a haplotype-resolved de-novo assembly tool for PacBio HiFi reads (Cheng et al., 2021). Hifiasm was run without the Hi-C sequences as its inclusion led to a drastic increase of contigs, from 34 without Hi-C to ~6,000 with Hi-C. Due to the high content of low-divergence repeats,

137   we soft-masked the draft genome with the *Earlgrey* v4.1.1pipeline (Baril et al., 2024) prior to Hi-

138   C mapping, which also provide transposable element (TEs) annotations (see below).

140   *Hi-C incorporation*

141   To incorporate the Hi-C sequencing with the draft genome assembly, we initially aligned

the Hi-C sequences to the HiFi draft genome with a Burrow-Wheeler Alignment (*BWA* v 0.7.19)

(Li, 2013). Then, we processed the resulting alignments with *SAMtools* v1.22.1 (Danecek et al.,

2021) to remove duplicate sequences. We then scaffolded the assembly using the *de novo YAHS*

v1.2.2 assembly pipeline (Zhou et al., 2023). From there, we used Juicer Tools to generate a Hi-

C contact map (Durand et al., 2016). Finally, we visualized the scaffolding of the chromosome-

level assembly with *Juicebox Assembly Tools* v2.20.0 (Robinson et al., 2018).

TE and gene annotation

150   We annotated the final assembly by first modelling and quantifying TEs by soft-masking

the genome with *EarlGrey* v4.1.1, a fully automated TE annotation pipeline (Baril et al., 2024)

with the default settings, which includes a 100bp minimum length and 10 iterations of the

BLAST, extract, and extend process. Next, we functionally annotated the predicted gene regions

using the train, predict, update, fix, and annotate steps of the *funannotate* pipeline (Palmer and

Stajich, 2020). The 'train' step aligns RNA-seq data, assembles it with *Trinity* (Grabherr et al.,

2011), and runs *PASA*, which models gene structures based on alignments of expressed

transcripts (Haas et al., 2003). The 'predict' step uses *PASA* gene models to train *Augustus*, a *de*

novo gene finder (Stanke et al., 2008), prior to running *EvidenceModeler* (Haas et al., 2008). We

included RNA-seq data from the liver, lung, heart, skeletal muscle, kidney, and skin as evidence

for *EvidenceModeler*. We used the Tetrapoda BUSCO database with *Taeniopygia guttata* as the

seed species for the 'predict' step and kept the default options for each step. The 'update' step

fixes gene models that disagree with RNA-seq data, which are corrected in the 'fix' step. Prior to

running the 'annotate' step, we ran *InterProScan* v5.74-105.0 (Jones et al., 2014) to run predicted

genes against the InterPro database for gene families and downloaded *Eggnog-mapper* v2.1.12

(Cantalapiedra et al., 2021) locally to be run during the 'annotate' step. The 'annotate' step

incorporates the generated data into an annotated genome. We used the default functional

annotation databases in the annotation step. To compare across existing chromosome-level skink

genomes, we ran the above pipeline on the assemblies of *Tiliqua scinciodes, Spondylurus nitidus,*

Bassiana duperreyi, and Carinascincus occelatus.

Synteny analysis

To assess genomic synteny of the *Plestiodon fasciatus* genome and other chromosome-

level squamate assemblies, we created a custom pipeline called *Synk*

(https://github.com/jomhoff/Synk) that uses the output files from *compleasm* (Huang and Li,

2023) and isolates the BUSCO genes to create comparative text files, and uses *RIdeogram* (Hao

et al., 2020) to plot the syntenic chromosomal regions from BUSCO genes between species in

one script. Compared to methods for calculating and visualizing whole genome synteny, *Synk*

runs much faster. In addition, limiting the dataset to BUSCO genes minimizes paralogy issues, as

BUSCO genes are highly conserved, single copy orthologs. This is especially effective in

ensuring appropriate estimates of synteny in cross-species and cross-genera comparisons. Here,

we show synteny between *P. fasciatus*, *Bassiana duperreyi*, *Spondylurus nitidus, Carinascincus*

ocellatus, Tiliqua scincoides, and *P. gilberti*.

Results and Discussion

Assembly of the P. fasciatus genome

After completing *Hifiasm* with the long-read PacBio sequences, the draft assembly of the

genome was close to complete, with an L50 of 3 and a total of 34 contigs, 9 of which represented

near telomere-to-telomere chromosomes (**Table 1**). With the Hi-C data incorporated, we

successfully generated a chromosome-level assembly of *Plestiodon fasciatus* with 18 unplaced

scaffolds that we refer to as *rPleFas1.1* after Vertebrate Genome Project naming rules (Rhie et al.

2021) (**Table 1; Figure 1**). One of the 18 unplaced scaffolds includes the complete

mitochondrial genome. The other 17 unplaced scaffolds range in size from 8,262bp to

1,241,752bp and consist of unplaced TEs and mRNA sequences. For the complete autosomal

genome, the presence of telomeres was estimated with the characteristic rise in GC content at the

ends of chromosomes (**Figure 1**) and confirmed with *tidk* v0.2.65, a toolkit for identifying

telomeres that outputs a count of telomeric repeats in windows across the genome (Brown et al.,

2025). There is also a large spike in GC content in the middle of the second chromosome, which

corresponds to a repeat-dense region consisting of Long Interspersed Nuclear Elements (LINE),

simple, and unclassified repeats.

	Metric	Draft Genome	Hi-C Informed Genome
Basic Stats	Total assembly size (bp)	1,537,706,373	1,537,706,373
	Number of contigs	34	34
	Number of scaffolds	34	31
	GC content (%)	45.6	45.6
Contiguity	Contigs N50 (bp)	166,278,287	166,278,287
	Contigs L50 (bp)	3	3
	Largest Contig (bp)	244,253,184	244,253,184
	Scaffolds N50 (bp)	166,278,287	227,050,938

	Scaffolds L50	3	2
	Largest Scaffold (bp)	244,253,184	304,269,845
Completeness	BUSCO Complete (%)	98.17	98.17
	Missing BUSCOs (%)	1.24	1.24
	Fragmented BUSCOs (%)	0.19	0.19
	Sequencing coverage (X)	53	116
	Telomere to Telomere Chromosomes	6	13

Table 1: Assembly statistics of the genome before and after Hi-C incorporation

Annotation of genes and TEs

The resulting annotations from the *funannotate* pipeline consists of an estimated 32,520

genes. Filtering for named genes and removing isoforms resulted in 16,100 unique genes with

common names (**Table 2**). Like other squamates, the genome of *P. fasciatus* has a high

proportion of repetitive regions which account for 46.7% of the genome (Pasquesi et al., 2018).

Unclassified repeats account for 23.3% of the genome, while DNA repeats and LINEs

respectively make up 7.0% and 9.3% of the genome (**Figure 2**). The Kimura 2-parameter

distance between the repeat sequences produces an approximately bimodal distribution, with a

peak indicating a large degree of moderately diverged repeats and a peak indicating recently

diverged repeats (**Figure 2**). The more diverged peak is mainly driven by unclassified repeats.

Interestingly, this a pattern that is seen in all five skinks tested. Furthermore, the more recently

diverged peak is driven by a large proportion of both DNA and LINE repeats and is also

conserved across the species analyzed (**Figure 2**).

Although the impact of TEs on the diversification and adaptation of squamates has yet to

be studied in detail, it has been hypothesized that variation in TEs can impact phenotypic

adaptation through TE domestication, exaptation, host-gene regulation, formation of retrogenes,

and genomic plasticity (reviewed in Schrader and Schmitz, 2019), and that TEs contribute

significantly to genomic variation (Catlin and Josephs, 2022). Since TEs are highly mobile

across the genome, are abundant, labile, and have been major players in the evolution of

eukaryotic genomes (Bowen and Jordan, 2002; Oliver and Greene, 2009), we expect variation of

TEs among distantly diverged clades. Despite originating around ~115mya, the skinks in this

study display remarkable conservation of TE composition, especially when considering the

variation seen in younger clades, such as extant mammals (Platt et al., 2018) and plethodontid

salamanders (Sun et al., 2012).

Category	Metric	Value
Gene Content	Number of protein-coding genes	32,520
	Unique genes with common names	16,100
Gene Structure	Average gene length (bp)	9,372.34
	Total number of exons	252,300
	Average exon length (bp)	255.27
	Multiple-exon transcripts	25,572
	Single-exon transcripts	7,372
Functional Annotation	Genes with GO term annotations	21,578
	Genes with InterProScan hits	24,316
	Genes with eggNOG annotations	26,007
	Genes with Pfam domains	19,449
	CAZyme-annotated genes	319

MEROPS-annotated genes	1,143

Table 2: Annotation Statistics of the Genome after the *Funannotate* pipeline

Synteny of skink genomes

The genomes of *Plestiodon gilberti* and *P. fasciatus* appear highly syntenic, with the only

rearrangement being an inversion in the eighth chromosome. Despite the ~150ma divergence

(Title et al., 2024), the macrochromosomes are largely conserved across the skink species

analyzed, aside from a large inversion on chromosome 1 between *Tiliqua scincoides* and all other

species (**Figure 3**). Compared to the other scincids, *P. fasciatus* and *P. gilberti* have fewer

microchromosomes, where chromosomes 14, 15, and 16 in *S. nitidus* and *T. scincoides* are

syntenic with chromosome 8 in *P. fasciatus* and *P. gilberti*. There is also a rearrangement creating

syntenic blocks relating chromosomes 6 and 7 in *S. nitidus* and *T. scincoides* to chromosome 5 in

P. fasciatus and *P. gilberti*.

Due to the cryptic, homogametic XY sex-determining system in many skinks, little is

known about the position of sex-determining regions on the chromosomes; however,

hypothetical sex chromosomes have been identified in *Bassiana duperryi* (Dissanayake et al.,

2020; Hanrahan et al., 2025). Here, we show that a block of chromosome 5 is syntenic with the

X chromosome in *B. duperreyi*, indicating a potential location for sex-determining regions in *P.*

fasciatus (**Figure 3**). The same block of the fifth chromosome of *Plestiodon gilberti* is syntenic

with the X chromosome of *B. duperreryi* (Richmond et al., 2025). In an attempt to further

identify sex-linked regions, we used the *FindZX* pipeline (Sigeman et al., 2022) with two male

and two female *P. fasciatus*; however, the results were inconclusive. Despite some preliminary

work, more research is required to further classify sex-determination in skinks, ideally with

population-level sampling of populations with numerous representatives from both sexes across

Scincidae.

Conclusion

We present a high-quality, telomere-to-telomere, chromosome-level annotated reference

assembly of the North American common five-lined skink *Plestiodon fasciatus* (Linnaeus 1758),

representing one of the most complete reference genomes (*rPleFas1.1*) to date of any species of

Scincidae. We find that macrochromosome structure is conserved across the family, but there are

common rearrangements of the microchromosomes, including a likely fusion in *P. fasciatus*,

which has fewer microchromosomes than many other skink species. We also present insight into

the content and evolution of transposable elements in skinks, which show remarkable

conservation between species over the last ~115Ma.

Data availability

The data presented in the paper is available on NCBI (BioProject PRJNA1278702) and Figshare

(https://figshare.com/projects/Telomere-to-telomere_reference_genome_of_the_common_five-

lined_skink_Plestiodon_fasciatus_Squamata_Scincidae_/254723). The code for analyses can be

found at https://github.com/jomhoff/Chromosome-level_genome_assembly and

https://github.com/jomhoff/Genome-Annotation.

Acknowledgments

The authors thank Megan Wallace of the American Museum of Natural History's Institute of

Comparative Genomics for her laboratory expertise, as well as Mac Mahacek, Steven Price,

William Taylor, and Elijah Wessel for field assistance. The authors also thank the Michigan

Department of Natural Resources and Kentucky Department of Fish and Wildlife. Finally, we

thank two anonymous reviewers that improved the quality of this manuscript.

Funding

This work was funded in part by the American Museum of Natural History's Richard Gilder

Graduate School, the Theodore Roosevelt Memorial Fund, and the Society for the Study of

Amphibians and Reptiles' Roger Conant Grants-in-Herpetology. This research was supported in

part by the U.S. National Science Foundation: DBI 2029955 awarded to CJR and DEB 2323125

to FTB.

Figure Captions:

Figure 1: Ideogram for *Plestiodon fasciatus*. **Row A** represents chromosome number
and relative size. **Row B** is a heatmap of gene density for 50kb windows, where darker
colors represent a higher density of genes. **Row C** is a heatmap of repeat elements for
50kb windows, where darker colors represent a higher density of repeat elements. **Row
D** is a line chart of GC content across the genome. **E** is the the region of the genome
that is syntenic with the X chromosome of *Bassiani duperreyi*. **F** is a photograph of an
adult male and juvenile *Plestiodon fasciatus* (photo via Herps of Arkansas). Plot made
with Circos (Krzywinski et al., 2009).
292
293   **Figure 2:** Divergence estimates of TEs shown with Kimura 2-Parameter Distance for
294   five chromosome-level skink genomes where the higher the Kimura 2-Parameter
295   Distance, the more diverged the sequences. The proportion of each transposable
296   element subdivision in the genome for each species is displayed with a pie chart.
297
298   **Figure 3:** Synteny of BUSCO genes between *Plestiodon fasciatus* and other skinks. **A**.
299   *P.* gilberti; **B**. *Tiliqua scincoides;* **C**. *Spondylurus nitidus*; **D**. *Carinascincus ocellatus;* **E**.
300   *Bassiana duperreyi*.
301

References

Alam, S.M.I., Sarre, S.D., Gleeson, D., Georges, A., Ezaz, T., 2018. Did Lizards Follow
Unique Pathways in Sex Chromosome Evolution? Genes (Basel) 9, 239.
https://doi.org/10.3390/genes9050239

Baril, T., Galbraith, J., Hayward, A., 2024. Earl Grey: A Fully Automated User-Friendly
Transposable Element Annotation and Analysis Pipeline. Molecular Biology and
Evolution 41, msae068. https://doi.org/10.1093/molbev/msae068
Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170
Bowen, N.J., Jordan, I.K., 2002. Transposable elements and the evolution of eukaryotic
complexity. Curr Issues Mol Biol 4, 65–76.
Brandies, P., Peel, E., Hogg, C.J., Belov, K., 2019. The Value of Reference Genomes in
the Conservation of Threatened Species. Genes 10, 846.
https://doi.org/10.3390/genes10110846
Brandley, M.C., Wang, Y., Guo, X., Montes de Oca, A.N., Fería-Ortíz, M., Hikida, T., Ota,
H., 2011. Accommodating Heterogenous Rates of Evolution in Molecular
Divergence Dating Methods: An Example Using Intercontinental Dispersal of
Plestiodon (Eumeces) Lizards. Systematic Biology 60, 3–15.
https://doi.org/10.1093/sysbio/syq045
Brazeau, D., Freitag, R., Hecnar, S.J., Hecnar, D.R., 2015. Comparing Common Five-
lined Skink (Plestiodon fasciatus) Diet Among Populations and Time. |
EBSCOhost [WWW Document]. URL
https://openurl.ebsco.com/contentitem/gcd:115178012?sid=ebsco:plink:crawler&i
d=ebsco:gcd:115178012 (accessed 6.11.25).
Brown, J., Pirrung, M., McCue, L.A., 2017. FQC Dashboard: integrates FastQC results
into a web-based, interactive, and extensible FASTQ quality control tool.
Bioinformatics 33, 3137–3139. https://doi.org/10.1093/bioinformatics/btx373
Brown, M.R., Manuel Gonzalez de La Rosa, P., Blaxter, M., 2025. tidk: a toolkit to
rapidly identify telomeric repeats from genomic datasets. Bioinformatics 41,
btaf049. https://doi.org/10.1093/bioinformatics/btaf049
Camacho-Sanchez, M., Burraco, P., Gomez-Mestre, I., Leonard, J.A., 2013.
Preservation of RNA and DNA from mammal samples under field conditions. Mol
Ecol Resour 13, 663–673. https://doi.org/10.1111/1755-0998.12108
Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., Huerta-Cepas, J., 2021.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Molecular Biology and Evolution 38, 5825–
5829. https://doi.org/10.1093/molbev/msab293
Catlin, N.S., Josephs, E.B., 2022. The important contribution of transposable elements
to phenotypic variation and evolution. Current Opinion in Plant Biology 65,
102140. https://doi.org/10.1016/j.pbi.2021.102140
Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., Li, H., 2021. Haplotype-resolved de
novo assembly using phased assembly graphs with hifiasm. Nat Methods 18,
170–175. https://doi.org/10.1038/s41592-020-01056-5
Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham,
A., Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of
SAMtools and BCFtools. GigaScience 10, giab008.
https://doi.org/10.1093/gigascience/giab008
Deweese, J.E., Wright, J.W., 1970. A preliminary karyological analysis of scincid lizards.
Mammalian Chromosomes Newsletter 11, 95–96.

Dissanayake, D.S.B., Holleley, C.E., Hill, L.K., O'Meally, D., Deakin, J.E., Georges, A.,
2020. Identification of Y chromosome markers in the eastern three-lined skink
(Bassiana duperreyi) using in silico whole genome subtraction. BMC Genomics
21, 667. https://doi.org/10.1186/s12864-020-07071-2
Dodge, T.O., Farquharson, K.A., Ford, C., Cavanagh, L., Schubert, K., Schumer, M.,
Belov, K., Hogg, C.J., 2025. Genomes of two Extinct-in-the-Wild reptiles from
Christmas Island reveal distinct evolutionary histories and conservation insights.
Mol Ecol Resour 25, e13780. https://doi.org/10.1111/1755-0998.13780
Dodge, T.O., Farquharson, K.A., Ford, C., Cavanagh, L., Schubert, K., Schumer, M.,
Belov, K., Hogg, C.J., n.d. Genomes of two Extinct-in-the-Wild reptiles from
Christmas Island reveal distinct evolutionary histories and conservation insights.
Molecular Ecology Resources n/a. https://doi.org/10.1111/1755-0998.13780
Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S.,
Aiden, E.L., 2016. Juicer Provides a One-Click System for Analyzing Loop-
Resolution Hi-C Experiments. Cell Syst 3, 95–98.
https://doi.org/10.1016/j.cels.2016.07.002
Ezaz, T., Sarre, S.D., O'Meally, D., Graves, J.A.M., Georges, A., 2009. Sex
chromosome evolution in lizards: independent origins and rapid transitions.
Cytogenet Genome Res 127, 249–260. https://doi.org/10.1159/000300507
Fitch, H.S., Fitch, H.S., 1954. Life history and ecology of the five-lined skink, Eumeces
fasciatus. University of Kansas publications, Museum of Natural History 8, 1--
156.
Gable, S.M., Mendez, J.M., Bushroe, N.A., Wilson, A., Byars, M.I., Tollis, M., 2023. The
State of Squamate Genomics: Past, Present, and Future of Genome Research in
the Most Speciose Terrestrial Vertebrate Order. Genes 14, 1387.
https://doi.org/10.3390/genes14071387
Gamble, T., Coryell, J., Ezaz, T., Lynch, J., Scantlebury, D.P., Zarkower, D., 2015.
Restriction Site-Associated DNA Sequencing (RAD-seq) Reveals an
Extraordinary Number of Transitions among Gecko Sex-Determining Systems.
Molecular Biology and Evolution 32, 1296–1309.
https://doi.org/10.1093/molbev/msv023
Giovannotti, M., Caputo, V., O'Brien, P.C.M., Lovell, F.L., Trifonov, V., Cerioni, P.N.,
Olmo, E., Ferguson-Smith, M.A., Rens, W., 2009. Skinks (Reptilia: Scincidae)
have highly conserved karyotypes as revealed by chromosome painting.
Cytogenet Genome Res 127, 224–231. https://doi.org/10.1159/000295002
Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis,
X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N.,
Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K.,
Friedman, N., Regev, A., 2011. Full-length transcriptome assembly from RNA-
Seq data without a reference genome. Nat Biotechnol 29, 644–652.
https://doi.org/10.1038/nbt.1883
Haas, B.J., Delcher, A.L., Mount, S.M., Wortman, J.R., Smith, R.K., Hannick, L.I., Maiti,
R., Ronning, C.M., Rusch, D.B., Town, C.D., Salzberg, S.L., White, O., 2003.
Improving the Arabidopsis genome annotation using maximal transcript
alignment assemblies. Nucleic Acids Res 31, 5654–5666.
https://doi.org/10.1093/nar/gkg770

Haas, B.J., Salzberg, S.L., Zhu, W., Pertea, M., Allen, J.E., Orvis, J., White, O., Buell, C.R., Wortman, J.R., 2008. Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. Genome Biology 9, R7. https://doi.org/10.1186/gb-2008-9-1-r7

Hanrahan, B.J., Alreja, K., Reis, A.L.M., Chang, J.K., Dissanayake, D.S.B., Edwards, R.J., Bertozzi, T., Hammond, J.M., O'Meally, D., Deveson, I.W., Georges, A., Waters, P., Patel, H.R., 2025. A genome assembly and annotation for the Australian alpine skink Bassiana duperreyi using long-read technologies. G3 Genes|Genomes|Genetics 15, jkaf046. https://doi.org/10.1093/g3journal/jkaf046

Hao, Z., Lv, D., Ge, Y., Shi, J., Weijers, D., Yu, G., Chen, J., 2020. RIdeogram: drawing SVG graphics to visualize and map genome-wide data on the idiograms. PeerJ Comput. Sci. 6, e251. https://doi.org/10.7717/peerj-cs.251

Harding, J.H., 1997. Amphibians and Reptiles of the Great Lakes Region. University of Michigan Press.

Hardy, L.M., Raymond, L.R., Harris, S., 2017. The Karyotype of Plestiodon anthracinus (Baird, 1850) (Sauria: Scincidae): A Step Toward Solving an Enigma. sena 16, 326–330. https://doi.org/10.1656/058.016.0318

Howes, B.J., Lougheed, S.C., 2008. Genetic diversity across the range of a temperate lizard. Journal of Biogeography 35, 1269–1278. https://doi.org/10.1111/j.1365-2699.2007.01867.x

Huang, N., Li, H., 2023. compleasm: a faster and more accurate reimplementation of BUSCO. Bioinformatics 39, btad595. https://doi.org/10.1093/bioinformatics/btad595

Janzen, F.J., Phillips, P.C., 2006. Exploring the evolution of environmental sex determination, especially in reptiles. J Evol Biol 19, 1775–1784. https://doi.org/10.1111/j.1420-9101.2006.01138.x

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., Hunter, S., 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics 30, 1236–1240. https://doi.org/10.1093/bioinformatics/btu031

Kostmann, A., Kratochvíl, L., Rovatsos, M., 2021. Poorly differentiated XX/XY sex chromosomes are widely shared across skink radiation. Proceedings of the Royal Society B: Biological Sciences 288, 20202139. https://doi.org/10.1098/rspb.2020.2139

Krzywinski, M., Schein, J., Birol, İ., Connors, J., Gascoyne, R., Horsman, D., Jones, S.J., Marra, M.A., 2009. Circos: An information aesthetic for comparative genomics. Genome Res 19, 1639–1645. https://doi.org/10.1101/gr.092759.109

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM [WWW Document]. arXiv.org. URL https://arxiv.org/abs/1303.3997v2 (accessed 6.11.25).

Mezzasalma, M., Guarino, F.M., Odierna, G., 2021. Lizards as Model Organisms of Sex Chromosome Evolution: What We Really Know from a Systematic Distribution of Available Data? Genes 12, 1341. https://doi.org/10.3390/genes12091341

Mezzasalma, M., Macirella, R., Odierna, G., Brunelli, E., 2024. Karyotype Diversification
and Chromosome Rearrangements in Squamate Reptiles. Genes (Basel) 15,
371. https://doi.org/10.3390/genes15030371
Oliver, K.R., Greene, W.K., 2009. Transposable elements: powerful facilitators of
evolution. BioEssays 31, 703–714. https://doi.org/10.1002/bies.200800219
Palmer, J.M., Stajich, J., 2020. Funannotate v1.8.1: Eukaryotic genome annotation.
https://doi.org/10.5281/zenodo.4054262
Pasquesi, G.I.M., Adams, R.H., Card, D.C., Schield, D.R., Corbin, A.B., Perry, B.W.,
Reyes-Velasco, J., Ruggiero, R.P., Vandewege, M.W., Shortt, J.A., Castoe, T.A.,
2018. Squamate reptiles challenge paradigms of genomic repeat element
evolution set by birds and mammals. Nat Commun 9, 2774.
https://doi.org/10.1038/s41467-018-05279-1
Pellegrino, K.C., Rodrigues, M.T., Yonenaga-Yassuda, Y., 1999. Chromosomal
polymorphisms due to supernumerary chromosomes and pericentric inversions
in the eyelidless microteiid lizard Nothobachia ablephara (Squamata,
Gymnophthalmidae). Chromosome Res 7, 247–254.
https://doi.org/10.1023/a:1009218628942
Pinto, B.J., Gamble, T., Smith, C.H., Wilson, M.A., 2023. A lizard is never late:
Squamate genomics as a recent catalyst for understanding sex chromosome and
microchromosome evolution. Journal of Heredity 114, 445–458.
https://doi.org/10.1093/jhered/esad023
Platt, R.N., Vandewege, M.W., Ray, D.A., 2018. Mammalian transposable elements and
their impacts on genome evolution. Chromosome Res 26, 25–43.
https://doi.org/10.1007/s10577-017-9570-z
Powell, R., Conant, R., Collins, J.T., 2016. A Field Guide to Reptiles & Amphibians:
Eastern and Central North America. Houghton Mifflin Harcourt.
Richmond, J.Q., Escalona, M., Marimuthu, M.P.A., Nguyen, O., Sacco, S., Beraut, E.,
Toffelmier, E., Karin, B.R., Cooper, R.D., Fisher, R.N., Wang, I.J., Shaffer, H.B.,
2025. A chromosome-level reference genome assembly for Gilbert's skink
Plestiodon gilberti. J Hered esaf040. https://doi.org/10.1093/jhered/esaf040
Rivera, D., Henderson, J.B., Lam, A.W., Hostetter, N.J., Collazo, J.A., Bell, R.C., 2024.
High-Quality, Chromosome-Level Reference Genomes of the Viviparous
Caribbean Skinks Spondylurus nitidus and S. culebrae. Genome Biology and
Evolution 16, evae079. https://doi.org/10.1093/gbe/evae079
Robinson, J.T., Turner, D., Durand, N.C., Thorvaldsdóttir, H., Mesirov, J.P., Aiden, E.L.,
2018. Juicebox.js Provides a Cloud-Based Visualization System for Hi-C Data.
Cell Syst 6, 256-258.e1. https://doi.org/10.1016/j.cels.2018.01.001
Rovatsos, M., Altmanová, M., Johnson Pokorná, M., Velenský, P., Sánchez Baca, A.,
Kratochvíl, L., 2017. Evolution of Karyotypes in Chameleons. Genes (Basel) 8,
382. https://doi.org/10.3390/genes8120382
Schmid, M., Feichtinger, W., Nanda, I., Schakowski, R., Visbal Garcia, R., Manzanilla
Puppo, J., Fernández Badillo, A., 1994. An extraordinarily low diploid
chromosome number in the reptile Gonatodes taniae (Squamata, Gekkonidae). J
Hered 85, 255–260. https://doi.org/10.1093/oxfordjournals.jhered.a111452
Schrader, L., Schmitz, J., 2019. The impact of transposable elements in adaptive
evolution. Molecular Ecology 28, 1537–1549. https://doi.org/10.1111/mec.14794

Seburn, C.N.L., 1993. Spatial distribution and microhabitat use in the five-lined skink
(Eumeces fasciatus). Can. J. Zool. 71, 445–450. https://doi.org/10.1139/z93-064
Sigeman, H., Sinclair, B., Hansson, B., 2022. Findzx: an automated pipeline for
detecting and visualising sex chromosomes using whole-genome sequencing
data. BMC Genomics 23, 328. https://doi.org/10.1186/s12864-022-08432-9
Soltis, D.E., Morris, A.B., McLACHLAN, J.S., Manos, P.S., Soltis, P.S., n.d. Comparative
phylogeography of unglaciated eastern North America.
Stanke, M., Diekhans, M., Baertsch, R., Haussler, D., 2008. Using native and
syntenically mapped cDNA alignments to improve de novo gene finding.
Bioinformatics 24, 637–644. https://doi.org/10.1093/bioinformatics/btn013
Sun, C., Shepard, D.B., Chong, R.A., López Arriaza, J., Hall, K., Castoe, T.A.,
Feschotte, C., Pollock, D.D., Mueller, R.L., 2012. LTR retrotransposons
contribute to genomic gigantism in plethodontid salamanders. Genome Biol Evol
4, 168–183. https://doi.org/10.1093/gbe/evr139
Title, P.O., Singhal, S., Grundler, M.C., Costa, G.C., Pyron, R.A., Colston, T.J., Grundler,
M.R., Prates, I., Stepanova, N., Jones, M.E.H., Cavalcanti, L.B.Q., Colli, G.R., Di-
Poï, N., Donnellan, S.C., Moritz, C., Mesquita, D.O., Pianka, E.R., Smith, S.A.,
Vitt, L.J., Rabosky, D.L., 2024. The macroevolutionary singularity of snakes.
Science 383, 918–923. https://doi.org/10.1126/science.adh2449
Worley, K.C., Richards, S., Rogers, J., 2017. The value of new genome references.
Experimental Cell Research 358, 433–438.
https://doi.org/10.1016/j.yexcr.2016.12.014
Xu, W., Li, T., Li, J., Liu, S., Yu, X., Tang, M., Dong, J., Liu, J., Bu, X., Xia, X., Zhou, H.,
Nie, L., 2024. The First Identification of Homomorphic XY Sex Chromosomes by
Integrating Cytogenetic and Transcriptomic Approaches in Plestiodon elegans
(Scincidae). Genes 15, 664. https://doi.org/10.3390/genes15060664
Xu, W., Zhu, W., 2024. First karyological analysis of Plestiodon capito (Squamata:
Scincidae). Cytologia 89, 147–151. https://doi.org/10.1508/cytologia.89.147
Zhou, C., McCarthy, S.A., Durbin, R., 2023. YaHS: yet another Hi-C scaffolding tool.
Bioinformatics 39, btac808. https://doi.org/10.1093/bioinformatics/btac808
518

Figure 1: Ideogram for *Plestiodon fasciatus*. **Row A** represents chromosome number and relative size. **Row B** is a heatmap of gene density for 50kb windows, where darker colors represent a higher density of genes. **Row C** is a heatmap of repeat elements for 50kb windows, where darker colors represent a higher density of repeat elements. **Row D** is a line chart of GC content across the genome. **E** is the the region of the genome that is syntenic with the X chromosome of *Bassiani duperreyi*. **F** is a photograph of an adult male and juvenile *Plestiodon fasciatus* (photo via Herps of Arkansas). Plot made with Circos (Krzywinski et al., 2009).

Figure 2: Divergence estimates of TEs shown with Kimura 2-Parameter Distance for five chromosome-level skink genomes where the higher the Kimura 2-Parameter Distance, the more diverged the sequences. The proportion of each transposable element subdivision in the genome for each species is displayed with a pie chart.

Figure 3: Synteny of BUSCO genes between *Plestiodon fasciatus* and other skinks. **A**. *P*. gilberti; **B**. *Tiliqua scincoides;* **C**. *Spondylurus nitidus*; **D**. *Carinascincus ocellatus;* **E**. *Bassiana duperreyi.*