Differential impacts of drought and esca expression on Ascomycota fungi in the trunks and young organs of mature grapevines

Pierre GASTOU[1,2], Giovanni BORTOLAMI[1,3], Nathalie FERRER[1], Gregory A. GAMBETTA[4], Samuele MORETTI[1], Jessica VALLANCE[1], Chloé E.L. DELMAS[1,*]

[1] SAVE, INRAE, Bordeaux Sciences Agro, ISVV, F-33882 Villenave d'Ornon, France

[2] Département Sciences de l'Environnement, Univ. Bordeaux, F-33405 Talence, France

[3] Present address: Plant Ecology Research Laboratory PERL, School of Architecture, Civil and Environmental Engineering ENAC, EPFL, CH-1015 Lausanne, Switzerland.

[4] EGFV, Bordeaux Sciences Agro, Univ. Bordeaux, INRAE, ISVV, F-33882 Villenave d'Ornon, France

*Corresponding author: chloe.delmas@inrae.fr

Author contributions: C.E.L.D., G.A.G. and G.B. conceived the experimental drought x esca study; G.B. and N.F. sampled the tissues and ground the samples; G.B. analyzed the trunk section images; N.F. extracted DNA and performed PCR; P.G and J.V processed the bioinformatics pipeline; P.G. analyzed the data; S.M. contributed to the discussion section and to data interpretation; P.G wrote the first draft of the paper under the supervision of C.E.L.D.; All authors critically revised and approved the final version of the manuscript for submission.

Abstract

Perennial plant decline is increasingly threatening the profitability and sustainability of agriculture and forestry worldwide. It results from intricate interactions between microbial communities, the plant host, and abiotic stressors. We investigated the effects of drought and esca disease on mature grapevine phytobiomes. Grapevines display no esca leaf symptoms during droughts, but the impacts of drought and esca expression on fungal communities and wood health in mature plants remain poorly understood.

We studied 43 uprooted 30-year-old naturally infected vines in three experimental conditions: well-watered asymptomatic (Control) vines, vines with esca symptoms (Esca), and vines subjected to water deficit (WD) over two consecutive summers. We profiled trunk, cane, stem and petiole Ascomycota communities by DNA metabarcoding with primers specifically designed for grapevine trunk-associated Ascomycota, and quantified wood necrosis.

The Ascomycota communities of trunks and younger organs clearly differed, and drought and esca had different impacts on the Ascomycota communities of perennial and young organs. In the trunk, drought significantly decreased fungal diversity in healthy wood and increased the abundance of wood pathogens (e.g. *Phaeomoniella chlamydospora, Botryosphaeria dothidea*). In young organs, esca expression decreased the species richness and diversity of the Ascomycota community to a greater extent than drought. We also found that the relative proportion of healthy wood was smaller in plants with esca symptoms than in asymptomatic plants.

50 Thus, drought increased Ascomycota pathogen abundance in the trunk but did not

51 increase wood degradation and esca expression, highlighting the need to investigate

52 the molecular basis of plant-microbiome interactions under multi-stress conditions.

54 **Keywords:** Ascomycota communities, drought, esca, grapevine, metabarcoding,

55 perennial plant decline, wood integrity

Introduction

A balanced plant microbiota is essential to preserve plant health and productivity in a context of global changes and emerging biotic pressures (Compant et al. 2019; Trivedi et al. 2020, 2022). The disequilibrium or dysbiosis of endophytic microbiota has been associated with plant dieback, although the causal mechanism is generally unknown (Arnault et al. 2023, Bettenfeld et al. 2020, Broberg et al. 2018, Griffiths et al. 2020). Pathogenic and saprotrophic species that are present in a latent endophytic phase within healthy plants (Hallmann et al. 1997, Marsberg et al. 2017) are often overabundant or more active in symptomatic plants, potentially worsening the decline of these plants (Bettenfeld et al. 2020, Jiao et al. 2023).

Plant microbial communities are also affected by abiotic factors, and these interactions are associated with plant dieback events worldwide (Ciesla and Donaubauer 1994, Desprez-Loustau et al. 2006, Manion 1981). In a context of climate change, the interactions of drought with plant physiology, plant pathogens and the whole microbial community (i.e. holobiont) are a major issue that cannot be reduced to unequivocal relationships. Depending on plant host species and physiological status, pathogen lifestyle, and the intensity and timing of the stress, drought may exacerbate, inhibit or have no effect on plant-pathogen interactions (Desprez-Loustau et al. 2006, Jactel et al. 2012, Luna et al. 2025, Ramegowda and Senthil-Kumar 2015, Torres-Ruiz et al. 2024). The impact of drought stress on phytobiomes has been documented principally for the rhizosphere and the phyllosphere (Bechtold et al. 2021, Fitzpatrick et al. 2018, Jaeger et al. 2024, Xu et al. 2018) but remains largely unknown for other compartments (e.g. endosphere or woody perennial organs). A comprehensive understanding of the interactions between perennial plant phytobiomes, abiotic and

biotic stresses would contribute to deciphering the relative roles of the various factors

in plant decline.

Grapevine (*Vitis vinifera* L.), a major cultivated perennial crop, is particularly severely

affected by complex dieback processes, mostly involving interactions between trunk

diseases and environmental factors (Claverie et al. 2020, Etienne et al. 2024). Esca is

a trunk disease causing grapevine dieback worldwide, with harmful consequences for

the productivity and perenniality of the vineyard (Gramaje et al. 2018). Esca triggers

physiological modifications in the leaf, stem and whole plant, impairing growth,

hydraulic conductivity, gas exchange, and metabolism within perennial and vegetative

organs (Bortolami et al. 2019, 2021a,b, 2023, Dell'Acqua et al. 2024a, Magnin-Robert

et al. 2016, 2017). This disease is detected in the summer on plant canopies displaying

typical symptoms of scorching between the veins of the leaf (so-called "tiger-stripe"

symptoms) and apoplexy, which may be observed on some or all of the stems of plants

randomly distributed over the vineyard (Lecomte et al. 2024). Internal wood necrosis,

with black necrotic tissue and white rot, is observed in both symptomatic and

asymptomatic plants (Bertsch et al. 2013). However, white rot tends to be more

frequent in symptomatic plants (Bruez et al. 2020, Maher et al. 2012). The trunk

microbiota differs between necrotic and non-necrotic (healthy) wood tissues (Bruez et

al. 2020, Elena et al. 2018). In particular, black necroses are characterized by a high

abundance of Ascomycota whereas white rot necrosis is caused by Basidiomycota

fungi, such as *Fomitiporia mediterranea* (Bruez et al. 2020; Moretti et al. 2021).

Conversely, there is no clear difference between the microbial communities of healthy

wood collected from plants with and without esca leaf symptoms (Del Frari et al. 2019,

Dell'Acqua et al. 2025, Hofstetter et al. 2012, Monod et al. 2024). Several

environmental factors are typically involved in perennial plant dieback but drought has

a particularly strong impact in viticulture, and this impact is growing even stronger with the acceleration of climate change (van Leeuwen et al. 2024). In this context, the impact of drought on esca pathogens is a growing matter of concern, as both drought and esca have deleterious effects on grapevine hydraulic function.

Recent advances have improved our understanding of the environmental drivers of esca symptom expression, highlighting the key role of drought through its impact on plant water status. Contrasting with the many observations of synergistic drought-pathogen interactions in trees (Desprez-Loustau et al. 2006, Gomez-Gallego et al. 2022, Torres-Ruiz et al. 2024), esca and drought are antagonist in grapevine, inhibiting each other. Bortolami et al. (2021b) demonstrated that uprooted naturally infected plants subjected to moderate drought (predawn water potential around -1 MPa) did not express esca leaf symptoms at any time in the course of a two-year experiment. Dell'Acqua et al. (in press) showed that vines with esca symptoms are less affected by intense drought stress and recover more quickly after such stress. The characteristics of fungal communities probably play a key role in drought-esca interactions, but this role has never been elucidated in mature plants.

Drought modifies the structure and functioning of plant microbial communities across compartments (see for example Carbone et al. 2021, Debray et al. 2022; Lin et al. 2023, Trivedi et al. 2022). A recent pluriannual study on young potted vines revealed a significant shift in wood fungal communities during drought, with a decrease in the complexity of microbiota structure and interactions, and an increase in the abundance of *P. chlamydospora*, a pathogenic Ascomycota involved in Petri disease that has been implicated in the esca pathosystem (Leal et al. 2024). These results seem to contradict the antagonism between drought and esca. There is therefore a need to

validate these findings on mature grapevines and to consider again which Ascomycota

pathogens are associated with the different disease symptoms.

Here, we studied the effects of abiotic (drought) and biotic (esca) stresses on

Ascomycota communities in the various organs of grapevine. We addressed the

following questions: (i) What impact do seasonal drought and esca leaf symptom

expression have on the Ascomycota communities in trunk and younger vegetative

organs? (ii) Is the antagonistic relationship between esca and drought associated with

shifts in the functional properties of Ascomycota communities? (iii) How do drought

and esca expression affect internal wood necrosis and the corresponding Ascomycota

communities? We used the same 43 uprooted vines with different esca histories and

exposure to contrasting water conditions as described by Bortolami et al. (2021b). We

sampled the trunk (separating the necrotic and healthy wood) and young organs (two-

year-old canes, current-year stems, and petioles), and used a metabarcoding

approach to study grapevine trunk-associated ascomycete fungi.

Materials and methods

Plant material and sampling. Sampling was conducted between October 7 and 14

2019 on a subset of uprooted vines (*V. vinifera* L. cv. Sauvignon blanc; *n*=43) from the

experimental design described in detail by Bortolami et al. (2021b). This experimental

design included three factors: (i) two levels of watering regime in two consecutive

summer seasons in 2018 and 2019: WW (well-watered for two years, Ψ_{PD} close to 0

MPa), and WD (water-deficit, presenting $\Psi_{PD} \approx$ -1 MPa from July to October in 2018

and 2019); (ii) two contrasting histories of esca symptoms since 2012: pA (previously

asymptomatic, i.e. plants that had never expressed esca leaf symptoms at any time

since 2012), and pS (previously symptomatic, i.e. plants that had expressed esca leaf

symptoms at least once since 2012); iii) two canopy phenotypes in 2019: presence of

esca leaf symptoms (typical tiger-stripe symptoms) and asymptomatic plants

(hereafter referred to as control plants). Plants subjected to a WD never expressed

esca leaf symptoms (see Bortolami et al. 2021b). We therefore sampled vines from

four distinct groups (Table 1): Water Deficit-pS (WD, n=12 asymptomatic plants in

2019); Esca-pS (Esca, n=10 well-watered plants expressing tiger-stripe symptoms in

2019); Control-pA (n=10, well-watered plants asymptomatic since 2012); Control-pS

(n=11, well-watered plants asymptomatic in 2019). Esca history had no significant

effect (see the Results section), so the Control-pA and Control-pS groups were

merged into a single control category (n=21, well-watered plants asymptomatic in

2019) for some analyses. During the growing seasons, the plants were regularly

treated for powdery mildew (with 80% micronized sulfur-based multisite products).

We sampled four different organs from each plant for DNA metabarcoding

(Table 1; Fig. S1): petioles (several petioles pooled together), stems from the current

year (one debarked internode from the middle of the stem), two-year-old canes (one

debarked internode), and trunks (with the separation of necrotic and healthy wood).

Consequently, both endophytic and epiphytic Ascomycota communities were included

in petiole analyses but only endophytic Ascomycota communities were included in the

analyses of woody organs (i.e. stems, canes, trunks). For plants expressing esca

symptoms, we separated petioles, stems and canes according to their disease status:

asymptomatic (AS) or displaying tiger-stripe symptoms on the leaves (S) (see pictures

of leaves in Fig. S1).

We used sterilized tools to remove the bark from the trunk of all plants. We

searched for the presence of a brown stripe, a colored band of xylem band appearing

under the bark concomitantly with esca leaf symptoms (Lecomte et al. 2012, 2024). A

brown stripe was observed in all the plants presenting leaf symptoms (Fig S1) but

none of the asymptomatic plants. We sampled chips from the brown stripe at different

heights along the trunk to obtain sufficient material. We then cut a 2 cm-thick cross-

section of the trunk 15 cm from the top of the plant and photographed it for further

analyses (Fig. S1). We collected tissue of different wood types from this cross-section

(after removal of the bark) with sterile shears in a sterile field. This tissue was

immediately immersed in liquid nitrogen and stored at -80°C. We collected the

following wood types (Table 1): apparently healthy wood, black necrotic wood, white

rot, and brown stripe. For white rot, if insufficient necrotic tissue was collected for

pathogen detection, we sampled additional white rot tissues from just above or below

the cut section. Necrotic tissue was sampled only when present, whereas healthy

wood was sampled from every plant. We obtained a total of 251 samples (Table 1):

134 young organs (48 petioles, 47 stems, 39 canes) and 117 trunks (43 healthy wood,

36 black necrosis, 29 white rot, and 9 brown stripes).

DNA extraction and sequencing. Samples (n=251) were ground in liquid nitrogen

with a one-ball mill TissueLyser II (Qiagen, Hilden, Germany), taking care to sterilize

the balls and sample holders with ethanol 70% between samples. Genomic DNA was

extracted from 60 mg wood tissue samples with the Indvisorb Spin Plant mini Kit

(Invitek Diagnostics, Berlin, Germany) according to the manufacturer's instructions.

Empty tubes — containing only the extraction reagents, without sample — were

included as negative controls for DNA extraction (n = 5). DNA was then quantified with

a DS-11 spectrophotometer (DeNovix, Wilmington, USA) and adjusted to a uniform

concentration of 15 ng.µl^{-1}. The ITS2 region of the nuclear ribosomal repeat unit was

then amplified with the GTAAf/GTTAr primers

(AAAACTTTCAACAACGGATC/GACCTCGGATCAGGTAGGRA), which were

specifically designed and optimized for the profiling of pathogenic Ascomycota fungal

communities associated with trunk diseases (Morales-Cruz et al. 2018b). DNA was

amplified by PCR with an Epgradient Mastercycler (Eppendorf, Hambourg, Germany)

in a reaction mixture (25 µl final volume) consisting of 1 µl DNA template (15 ng.µl-1),

10 µl of 2X Platinum Hot Start PCR Master Mix (Invitrogen,Waltham, USA), 0.5 µl of

each primer (10 µM), 1.25 µl 20X bovine serum albumin (New England BioLabs,

Ipswich, USA), and 11.75 µl DNAse/RNAse-free sterile water. The cycling parameters

were as follows: enzyme activation at 95°C for 2 min; 35 cycles of denaturation at 95°C

for 45 s, 60°C for 1 min, 72°C for 1.5 minutes; and a final extension at 72°C for 10 min.

Empty wells on the PCR plate — i.e. those containing the PCR mix but without

template — were used as negative controls (n = 9). The PCR products were visualized

by electrophoresis in a 2% agarose gel in TBE (molecular weight ≈350 bp). They were

then sent to the PGTB sequencing facility (Genome Transcriptome Facility of

Bordeaux, Pierroton, France) for sequencing on an Illumina MiSeq platform (v3

chemistry, 2x300 bp). PCR product purification, the addition of multiplex identifiers and

sequencing adapters, library sequencing and sequence demultiplexing (with exact

index search) were performed by the sequencing service, who included another five

negative PCR control samples at this stage.

Sequence analysis and OTU table construction. Metabarcoding data were

processed with the FROGS pipeline (Find Rapidly OTUs with Galaxy Solution)

implemented on a Galaxy instance (https://vm-galaxy-prod.toulouse.inra.fr/galaxy/;

Escudié et al. 2018). In brief, VSearch was used to merge paired reads with a

maximum of mismatch of 10% in the overlapping region. After denoising (i.e. reads

with no expected length - between 150 and 590 bp - and containing ambiguous bases,

N, were discarded) and the removal of primers/adapters removal with CUTADAPT

(Martin 2011), clustering was performed with SWARM (Mahé et al. 2014), with an

aggregation distance d = 1. Chimeras were removed and low-abundance sequences

were filtered out at 0.00005% (i.e. we retained OTUs accounting for at least 0.00005%

of all sequences, adapted from Bokulich et al. 2013), to discard singletons from the

datasets. The ITS2 region was extracted with ITSx (Bengtsson-Palme et al. 2013)

before determining the taxonomic affiliation of the fungal OTUs by BLAST queries

against the UNITE Fungi v8.2 (implemented in FROGS), trunkdiseaseID.org

(http://www. grapeipm.org/), and NCBI (nr/nt) nucleotide collection

(https://blast.ncbi.nlm.nih.gov/Blast. cgi) databases.

We performed a decontamination step, based on 19 DNA extraction and PCR

blanks, with the *metabaR v.1.0.0* package (Zinger et al. 2021) implemented in R

v.4.2.1 software. Graphical and statistical methods were used to flag and selectively

remove non-Ascomycota, putatively contaminating or artifactual OTUs, and potentially

unusable samples (i.e. highly contaminated or low sequencing depth).

Metabarcoding statistical analysis. We compared the diversity, structure and

composition of Ascomycota communities (see metrics below) between different

categories (Table 1): (i) organs (trunk, cane, stem, petiole; n = 251 samples),

considering plant status as a covariate; (ii) plant statuses in healthy trunk wood

samples; n = 43); (iii) plant statuses and organ statuses in young organs (n = 134); (iv)

trunk wood types (i.e. healthy wood, black necrosis, white rot, brown stripe; n = 117),

considering plant status as a covariate; (v) esca symptom histories (i.e. previously

asymptomatic *vs.* previously symptomatic) in control plants, considering plant status

as a covariate. The (v) analysis was performed separately on trunk (n = 51) and young

organ (n = 58) samples. For all analyses, we checked the normality of the residuals

(QQ-plot) and the homogeneity of the residual variance (residuals vs. fitted, residuals

vs. predictors) graphically. Non-parametric analyses were performed if these

assumptions were not satisfied. All analyses were performed with R v.4.2.1 and with

an alpha risk of 5%.

To estimate the diversity, structure and composition of Ascomycota

communities we calculated three alpha-diversity indices (i.e. observed richness,

Shannon index, Simpson index) with the "estimate_richness" function in *phyloseq*

v.1.42.0 package (McMurdie and Holmes 2013), and compared them across

categories by ANOVA procedures followed by Tukey's post-hoc tests. For Simpson

index, the normality assumptions were not fulfilled, so we performed non-parametric

alternatives (i.e. Kruskal-Wallis test followed by Dunn's test). We subjected the

Ascomycota community structure of the samples to principal component analysis

(PCA) based on a redundancy dependence analysis (RDA) ordination on centered

log-ratio (CLR, recommended for compositional data; Gloor et al. 2017)-transformed

data, with the *phyloseq* "ordinate" function. We performed a permutational analysis of

variance (PERMANOVA) on CLR-transformed data, with the "adonis2" function in the

vegan v.2.6-4 package (Oksanen et al. 2022), to assess the effect of organ, plant

statuses, or trunk wood type on Ascomycota community structure, with plant status

included as a covariate. The multivariate homogeneity of group dispersions was

checked *a posteriori*, and multilevel pairwise comparisons were calculated with the

"pairwise.adonis2" function in *pairwiseAdonis v.0.4.1* (Arbizu 2025). A pseudocount of

one was added to all OTU counts to handle zero values before CLR transformation,

thereby ensuring numerical stability during log-ratio calculations.

For some analyses, we classified Ascomycota OTUs into three groups

according to function: putative grapevine wood pathogens, putative antagonists of

grapevine wood pathogens (tested with success *in vitro*, in the greenhouse or in the

vineyard), and other taxa. This taxonomic classification to genus level was based on

the available scientific literature (Supplementary Table S1). The effect of plant status,

or wood type on these functional classes was evaluated by ANOVA followed by

Tukey's post-hoc tests.

We performed differential analysis with linear discriminant analysis effect size

(LEfSe) to identify the Ascomycota species most likely to explain differences between

wood types, with the "run_lefse" function in the *microbiomeMarker v.1.4.0* package

(Cao et al. 2022) and the following parameters: kw_cutoff = 0.05 and lda_cutoff = 2.

We performed a differential analysis with the Aldex2 procedure to identify the

Ascomycota species most likely to explain differences between control and Esca

plants, and between control and WD plants. In this case, the "run_aldex" function was

applied to RLE-transformed data, with a *p*-value cutoff of 0.05.

Analysis of trunk necrosis. We used ImageJ software (Schneider et al. 2012) to

analyze each trunk cross-section image. We measured the area of apparently healthy

wood, black necrosis, and white rot (in mm²) relative to the entire cross section. The

brown stripe was not visible on the cross sections because it affects only the outermost

xylem rings and was not, therefore, quantified as for the other tissues.

We assessed the effect of plant status (control, esca, WD) on the relative

proportions of healthy wood, black necrosis and white rot in trunk cross sections (n =

43). The effect of plant status on the statistical distribution of wood type proportions

was assessed with Fisher's exact test (simulated *p*-value, 2,000 replicates). The effect

of plant status on the proportion of each type of necrosis was assessed by ANOVA

followed by Tukey's post-hoc tests with "HSD.test" in the *agricolae v.1.3-7* package

(de Mendiburu 2023). We then plotted the statistical distribution of healthy wood and

compared it across plant statuses, using the minimal proportion of healthy wood in

control plants as a threshold. Finally, we assessed the effect of esca symptom history

on the proportion of each type of necrosis in control plants (n = 21).

Results

General description of the Ascomycota community and differences between

organs. In total, 10,765,305 reads were generated and assigned to 320 operational

taxonomic units (OTUs) (Supplementary Tables S2 and S3). The final decontaminated

dataset contained 9,641,320 reads (i.e. 10.4% of reads were removed) from 251 plant

samples that were affiliated to 297 OTUs (i.e. 7.2% were removed). Ascomycota

communities were composed of four main classes: Dothideomycetes (59.5%),

Eurotiomycetes (19.0%), Sordariomycetes (18.4%) and Leotiomycetes (2.2%). Less

than 1% of OTUs were affiliated to multiple classes or to no class. Global metrics for

both the raw and decontaminated datasets are provided in Supplementary Table S2.

The decontaminated OTUs are presented in Supplementary Table S3. The number of

reads per sample after decontamination and sample metadata are provided in

Supplementary Table S4.

We compared the diversity and structure of Ascomycota communities between

plant organs (i.e. trunk, cane, stem, petiole) across all samples. Shannon and Simpson

indices differed significantly between organs, whereas observed richness did not ($P =$

0.005, $P = 10^{-4}$ and $P = 0.10$, respectively). Ascomycota richness and diversity were

slightly lower in trunk samples than in other organs (Supplementary Fig. S2A). PCA

and PERMANOVA analyses revealed that organ accounted for 21% of the variance

of the total Ascomycota community ($P = 0.001$). Trunk samples were clearly separated

from canes, stems and petioles (Supplementary Fig. S2B). We therefore decided to

compare trunk samples and young organ samples (canes, stems and petioles,

corresponding to organs that were no more than two years old) in subsequent

analyses.

Effects of drought and esca on the Ascomycota community in healthy trunk

wood samples. Three species predominated in healthy trunk wood samples:

Phaeomoniella chlamydospora (28.2%), *Cladosporium sphaerospermum* (13.0%) and

Phaeoacremonium minimum (8.1%; Fig. 1A). Plant status significantly influenced the

Shannon and Simpson indices of healthy wood but not observed richness ($P = 0.01$,

$P = 0.04$ and $P = 0.17$, respectively). Ascomycota diversity was significantly lower in

the healthy wood of WD plants than in that of control plants, whereas it did not differ

significantly between esca and the other two experimental conditions (Fig. 1B). PCA

and PERMANOVA analyses revealed that plant status accounted for 6.9% of the

variance of the Ascomycota community in healthy wood ($P = 0.001$). WD plants

differed significantly from control plants, whereas esca plants were intermediate (Fig.

1C).

Plant status significantly affected the relative abundance of putative grapevine

wood pathogens and antagonists (Supplementary Table S1) in healthy wood, but did

not affect the abundance of other taxa (P = 0.006, P = 0.006 and P = 0.38, respectively;

Fig. 1D). The relative abundance of putative pathogens was significantly higher in WD

plants than in control plants (Fig. 1D), whereas the abundance of putative antagonists

was significantly lower in WD plants than in control plants (Fig. 1D).

Finally, Aldex2 differential analyses to identify Ascomycota species highlighted

differences in microbial composition between plants with different disease statuses.

One phylum-affiliated Ascomycota OTU was significantly less abundant in plants with

esca symptoms than in control plants. Four fungal species, including the pathogens

P. chlamydospora and *Botryosphaeria dothidea*, were significantly more abundant in

WD plants than in control plants. Conversely, four fungal species, including putative

antagonist *Cladosporium* spp. and *E. nigrum*, were significantly less abundant in WD

plants than in control plants (Fig. 1A; Supplementary Table S5).

Effects of drought and esca on Ascomycota communities in young organs. We

first looked for differences in the Ascomycota community between asymptomatic and

esca-symptomatic young organs from plants with esca symptoms. We found no

significant difference between these two groups of organs for either alpha-diversity or

beta-diversity metrics (Supplementary Table S6).

We then compared the Ascomycota communities in young organs between

control, esca (pooling samples from asymptomatic and symptomatic canes, stems and

petioles) and WD plants. Three species predominated in young organs: *Cladosporium*

sphaerospermum (44.5%), *Cladosporium halotolerans* (20.0%), and *Diaporthe*

ampelina (5.1%; Fig. 2A). The genus *Cladosporium* accounted for 69.1% of the OTUs

retrieved from young organs.

Plant status significantly influenced observed richness, and the Shannon and

Simpson indices ($P = 10^{-6}$, $P = 0.003$ and $P = 0.02$, respectively). Ascomycota diversity

was lower in the young organs of esca plants than in those of control and WD plants

(Fig. 2B). PCA and PERMANOVA analyses revealed that plant status accounted for

5.5% of the variance of the Ascomycota community in young organs ($P = 0.001$). In

pairwise comparisons, there were slight differences between plant statuses (Fig. 2C).

We performed Aldex2 differential analyses to identify the Ascomycota species

underlying the differences in microbial composition between the young organs of

plants with different disease statuses. *Erysiphe necator* was significantly more

abundant in esca plants than in control plants. Conversely, 12 fungal species,

including several putative antagonist *Cladosporium* spp. and *Aureobasidium pullulans*,

were significantly less abundant in esca plants than in control plants. Two fungal

species, *Exophiala* sp. and *B. dothidea*, were found to be significantly more abundant

in young organs from WD plants than in those of control plants. Conversely, six fungal

species, including several *Cladosporium* spp. and *Aureobasidium pullulans*, were

found to be significantly less abundant in WD plants than in control plants (Fig. 2A;

Supplementary Table S5). Overall, putative wood pathogens accounted for 11.8 ±

0.2% of the OTUs in young organs. These pathogens were slightly more abundant in

esca and WD plants than in control plants ($P = 0.05$). Putative antagonistic genera

accounted for 72.9 ± 0.2% of OTUs in young organs, and were less abundant in esca

and WD plants than in control plants ($P = 10^{-4}$).

Effects of drought and esca on wood integrity. We quantified the areas occupied

by apparently healthy wood, black necrotic wood and white rot in trunk cross sections

(Supplementary Table S7). The relative proportions of these three types of wood

differed significantly between plant statuses (i.e. control, esca and WD; $P = 10^{-4}$). The

proportion of healthy wood was significantly lower in plants with esca symptoms than

in control plants, but did not differ significantly between WD plants and the other two

categories (Fig. 3A). The proportions of black necrosis and white rot did not differ

significantly between plant disease statuses ($P = 0.10$ and $P = 0.12$, respectively), but

there was a trend towards a higher proportion of white rot in plants with esca symptoms

and a higher proportion of black necrosis in WD plants (Fig. 3A). In 60% of plants with

esca symptoms, the proportion of healthy wood was lower than the minimum value

recorded in control plants (34.3%).

Ascomycota communities in trunk necrotic wood tissues. We compared trunk

Ascomycota communities between apparently healthy wood, brown stripe, black

necrosis and white rot samples collected in October, several weeks after symptom

onset. Brown stripe samples were collected only from plants displaying esca

symptoms (Table 1) and the sample size is lower for this tissue type. Wood type

significantly affected alpha-diversity metrics, i.e. observed richness, and the Shannon

and Simpson indices ($P < 10^{-16}$, $P = 10^{-12}$ and $P = 10^{-6}$, respectively). Ascomycota

richness and diversity were lower in necrotic wood (i.e. black necrosis or white rot),

than in healthy wood and brown stripe (Fig. 3B). PCA and PERMANOVA analysis

revealed that wood type accounted for 11.9% of the variance of the Ascomycota

community in a model including plant status and wood type * plant status as covariates

($P = 0.001$). All pairwise comparisons between wood types were significant, but the

clearest difference was that between necrotic and non-necrotic areas (Fig. 3C). The

interaction between wood type and plant status was not significant: the differences in

Ascomycota communities between non-necrotic and necrotic wood were observed in

control, esca or WD plants (Supplementary Fig. S3).

Classifying Ascomycota OTUs into putative wood pathogens, antagonists and

others (Supplementary Table S1), we found that wood type significantly modified the

relative abundance of putative wood pathogens and putative antagonists, but not of

other taxa ($P = 10^{-15}$, $P = 10^{-10}$ and $P = 0.16$, respectively; Fig. 4A). The abundance of

putative pathogens was higher within the areas of black and white rot than in healthy

wood (Fig. 4A). The abundance of putative antagonists was high in healthy wood and

brown stripe, whereas these fungi were almost entirely absent from necrotic areas

(Fig. 4A). Finally, we performed a linear discriminant analysis to identify the

Ascomycota species driving the differences in microbial composition between wood

types. Six fungal species, including the putative antagonist of trunk pathogens

Epicoccum nigrum, were identified as biomarkers of healthy wood. Nine fungal species

were identified as biomarkers of brown stripe samples, including several pathogens

belonging to the Botryosphaeriaceae and putative antagonist *Cladosporium* spp.

Three fungal species were significantly associated with black necrosis, including the

pathogenic species *Phaeoacremonium minimum*. Four fungal species were

significantly associated with white rot, including the pathogenic species *Phaeomoniella*

chlamydospora and several *Penicillium* spp. (Fig. 4B).

Effects of esca history on wood necrosis and Ascomycota communities. We

assessed the effect of esca history on the distribution of wood types and the properties

of Ascomycota communities in control plants. Esca history (comparing pS and pA

plants with plant status as a covariable) had no significant impact on the distribution

of wood types (healthy wood, black necrosis and white rot) in trunk cross sections

(Supplementary Table S8).

In addition, the Ascomycota community in healthy trunk wood and young organ

samples did not differ significantly between pA and pS plants in terms of either alpha-

diversity or beta-diversity metrics (Supplementary Table S8).

Discussion

Disentangling the complex interplay between plants, microbial communities and

abiotic stresses is crucial for understanding the processes underlying perennial plant

decline. In this study, we used a unique experimental design to characterize plant and

Ascomycota community responses to esca disease and drought stress in mature

grapevines. We extended the work of Bortolami et al. (2021b) showing that the

expression of esca symptoms is inhibited by drought under controlled conditions, using

naturally infected plants. We sampled the plants at the end of the experiment and

demonstrated that the impact of drought on Ascomycota communities was greatest in

the trunk, whereas the expression of esca symptoms affected mostly younger organs.

The esca-drought antagonism revealed by Bortolami et al. (2021b) is not related to a

decrease in the abundance of grapevine wood pathogens. Finally, we demonstrated

that the incidence of esca symptoms on leaves was associated with a significant

decrease in the relative proportion of apparently healthy (i.e. non-necrotic) wood in the

trunk, with a corresponding increase in the proportion of necrotic tissues with a low

diversity of Ascomycota and a high abundance of wood pathogens.

Drought favors the pathogenic component of the Ascomycota community in trunk healthy wood whereas esca does not. In this study, the endophytic Ascomycota community in the healthy wood of the trunk was particularly strongly affected by intense drought stresses applied during two consecutive summers, whereas the seasonal expression of esca leaf symptoms affected the Ascomycota within young organs. The impact of drought has been studied in belowground communities in particular (e.g. Carbone et al. 2021, Jaeger et al. 2024), but drought can also shape endophytic microbial communities, by (i) directly affecting the living environment of fungi, by decreasing xylem water potential and humidity (Aung et al. 2018, Luard and Griffin 1981, Dell'Acqua et al., in press) and (ii) modulating microbe-microbe and plant-microbe interactions (Trivedi et al. 2020). We found a low diversity of Ascomycota in healthy trunk wood under drought conditions. This finding is consistent with the results of Leal et al. (2024), who used one-year-old grapevine cuttings to demonstrate a substantial decrease in microbial interactions, thereby providing evidence in support of a drought-induced microbial imbalance. The effects of drought stress on plant and microbial communities include osmotic shock and the generation of reactive oxygen species (ROS; Schimel et al., 2007; Miller et al., 2010, Singh et al., 2011). Under drought conditions, the Ascomycota community in the trunk may shift towards specific taxa with more efficient intracellular ROS scavenging mechanisms and an ability to produce compatible osmolytes. The relative abundance of Ascomycota wood pathogens, including *Phaeomoniella chlamydospora* in particular, increased in the trunk under water deficit conditions, even in the absence of foliar symptoms on water-stressed plants (Bortolami et al. 2021b). The increase in *P. chlamydospora* levels under drought conditions has been reported before (Hrycan et al. 2025, Leal et al. 2024). It may be due to a change in fungal interactions, with an

increase in the abundance of competitive species and their activities under drought,

and a greater adaptation of *P. chlamydospora* to a constrained environment (i.e. in the

presence of activated plant defenses and a low water potential) potentially favoring its

development at the expense of its usual antagonists and competitors. Interestingly, *P.*

chlamydospora seems to cope well with the oxidative stress induced by plant cells due

to its efficient ROS scavenging arsenal (Fischer et al., 2016). However, this finding is

more surprising given that the foliar expression of esca symptoms was inhibited under

drought conditions.

Interestingly, during the years following rewatering, the WD plants kept alive

had a probability of expressing esca leaf symptoms similar to that of control plants (C.

Delmas, personal communication). The increase in *P. chlamydospora* abundance did

not lead to a higher incidence of symptoms in subsequent years. The total wood

pathogen load appeared to be higher in plants expressing esca, due to the presence

of larger necrotic areas rich in pathogens. However, the observation that these

pathogens are not more abundant in the healthy wood of symptomatic plants, where

the process of degradation leading to necrosis presumably occurs, raises questions

about the exact role of Ascomycota in esca pathogenesis. Ascomycota pathogens are

generally considered to play an important, if not essential role in the expression of

esca symptoms (e.g. Bertsch et al. 2013, Gramaje et al. 2018). However, relative

abundance, studied here through metabarcoding approaches, may not be the most

relevant trait. Instead, the activity of Ascomycota pathogens (e.g. *Phaeomoniella*

chlamydospora, *Botryosphaeriaceae* spp.), including their expression of virulence

factors, may decrease in water deficit conditions, or increase before the expression of

esca symptoms (Desprez-Loustau et al. 2006, Morales-Cruz et al. 2018a, Nerva et al.

2022). There is also growing scientific evidence to suggest that the contribution of

Ascomycota to esca expression is overestimated (Hofstetter et al. 2012, Nerva et al.

2022, Moretti et al. 2023). These results call for refining our knowledge of gene

expression and metabolic changes in both plants and pathogens in the context of

multiple stress experiments.

Drought did not significantly modify the proportion of necrotic wood, but it did

slightly decrease the white rot/black necrosis ratio. This suggests that drought may

interfere slightly with the activity of white-rot basidiomycetes that degrade wood

through both enzymatic and non-enzymatic decay pathways (Moretti et al. 2021,

2023). Drought has already been identified as a variable that may affect wood decay

(Jia et al. 2024). Indeed, to grow through the lignocellulose matrix and release their

arsenal of decay-mediating molecules (i.e. enzymes and low-molecular weight

compounds) into the wood cell wall, wood decay fungi (Basidiomycota included)

require the moisture content in the wood to be at least above the fiber saturation point

(Goodell 2020, Zaber and Morrell 2012). This highlights the importance of studying the

abundance and activity of Basidiomycota, but also of bacteria that can act in synergy

in the wood degradation process (Del Frari et al. 2021, Haidar et al. 2021).

The expression of esca symptoms did not significantly modify plant-microbiome

interactions in healthy trunk tissues (despite a trend towards lower species richness

and an increase in the abundance of wood pathogens) as reported in other studies

(Del Frari et al. 2019, Monod et al. 2024; Dell'Acqua et al. 2025). Very few studies

have quantified the impact of dieback diseases on endophytic communities in other

plants (e.g. Raghavendra et al. 2017, Steinrucken et al. 2016), although modifications

in belowground microbial diversity were highlighted (Darriaut et al. 2024, Farooq et al.

2022, Han et al. 2022, Lasa et al. 2024). Such modifications may be accompanied by

an ecological disequilibrium, with an increase in the pathogenic component and a loss

of beneficial functions, such as symbioses between plants and fungi and the promotion

of plant growth (Gómez-Aparicio et al. 2022, Ma et al. 2024, Solís-García et al. 2021).

Our findings confirm that esca is a complex pathosystem, in which plant physiology,

interacting with the abiotic environment, plays a key role in leaf symptom onset, without

any major change in fungal communities in healthy wood tissue.

Esca induces a more significant shift in Ascomycota communities in young

organs than drought. Two consecutive summer seasons under drought conditions

had a greater impact on the trunk than esca, but seasonal esca expression had a

greater effect on Ascomycota communities in younger organs, which undergo

significant physical, physiological and metabolic changes during the expression of

esca symptoms (Bortolami et al. 2021a,b, Dell'Acqua et al. 2024, 2025). The

Ascomycota communities inhabiting young organs (mainly endophytes, except for the

petioles, from which we sampled both endophytes and epiphytes) on plants displaying

esca symptoms were less diverse than those of control plants and were largely

dominated by *Cladosporium* species. This finding suggests that the presence of leaf

symptoms influences, albeit slightly, microbial communities distal to the infection zone

characterized by necrosis and abundant trunk pathogens (i.e. perennial organs). Leaf

endophytic fungal communities have been shown to be modified by esca expression

(Del Frari et al. 2025). Accumulation and qualitative changes have been reported for

primary and secondary metabolites (e.g. sugars, terpenes, flavonoids; Bortolami et al.

2021b, Del Frari et al. 2025, Weiller et al. 2024) in leaves and stems expressing esca

symptoms, and these effects may be greater than those observed in the trunk,

explaining the stronger response of fungal communities to esca expression in younger

organs. Moreover, biotic stresses (such as esca disease) also induce ROS generation,

and fungi have developed strategies for scavenging ROS (Lehmann et al. 2015).

Therefore, as discussed above for the trunk, Ascomycota communities in young

organs may shift towards specific taxa with more efficient ROS scavenging

mechanisms.

However, esca had only a small effect in our experiment and other studies

have revealed the stem microbial communities to be similar in asymptomatic plants

and plants with esca symptoms (Del Frari et al. 2019). Water deficit had an even

weaker effect on the Ascomycota communities of young organs, despite the highly

significant physiological impact of drought during the experiment (Bortolami et al.

2021b). The main physiological difference between the plants subjected to water

deficit and those expressing esca symptoms was that water potential was strongly

affected by drought but not by esca expression, despite both conditions decreasing

plant transpiration. Photosynthesis and the balance of non-structural carbohydrates

were more strongly affected by esca expression, with a decrease in leaf chlorophyll

content and an accumulation of monosaccharides in leaves and stems, at the expense

of sucrose production and starch storage (Bortolami et al. 2021b). The Ascomycota

communities in young organs may have been more sensitive to carbon resource

availability and the concentration of non-structural carbohydrates, whereas wood

water status may be ecologically more selective for trunk communities. Conversely,

these results demonstrate the resilience of the microbial communities in young aerial

organs to decreases in water potential during drought, with no significant decrease in

diversity and no increase in the abundance of specific ecological guilds. Finally, the

relative abundance of *E. necator*, the causal agent of grapevine powdery mildew, was

higher in asymptomatic vegetative organs on plants displaying esca symptoms,

potentially due to a slight increase in gas exchange in these specific organs (Bortolami

et al. 2021b). *E. necator* was less prevalent in organs expressing esca symptoms,

perhaps due to the impairment of carbon assimilation (Guilpart et al. 2017).

On the relationship between wood necrosis, pathogens and the expression of

esca foliar symptoms. The causal links between the internal consequences of plant-

pathogen interactions (i.e. wood necrosis) and the external expression of disease (i.e.

foliar symptoms) remain poorly understood in perennial plant decline (Claverie et al.

2020, Gosling et al. 2024, Skovsgaard et al. 2010). Esca foliar symptoms have been

reported to be associated with a higher proportion of wood necrosis, especially white

rot, in *Vitis vinifera* L. cv. 'Cabernet-Sauvignon' (Maher et al. 2012, Ouadi et al. 2019).

Consistent with these findings, we found that plants displaying esca symptoms had

significantly lower proportions of healthy wood in their trunks than control plants.

However, our results demonstrate that the proportion of healthy wood explained esca

symptom expression in *V. vinifera* cv. 'Sauvignon blanc' better than the type of

necrosis. This may reflect a cumulative loss of functional xylem area (i.e. of the active

tissues located in the outer xylem; McElrone et al. 2021) in symptomatic plants,

regardless of the type of degradation. The mean proportion of white rot was about 20%

in control plants (i.e. well-watered and asymptomatic in the year of sampling), higher

than in previous reports. As plant age and genotype are significant drivers of esca leaf

symptom expression (Etienne et al. 2024, Gastou et al. 2024), we can hypothesize

that these factors modulate the proportion of necrotic wood (Claverie et al. 2023). In

old plants (30-y-o in our study), control plants with an asymptomatic history had similar

necrosis extent and Ascomycota community in the trunk than previously symptomatic

plants. In addition to a threshold of necrotic wood proportion in the trunk, the area of

contact between active xylem tissue (efficiently transporting sap) and specific necrotic

tissues, or the activity of microbial communities (e.g. virulence factors) may be

involved in the onset of foliar symptoms and should be explored further*.*

In the classical model of plant-microorganism interactions underpinning esca

expression, Ascomycota fungi are considered to be responsible for black necrosis,

black spots, wood streaks and discoloration, whereas white rot is associated with

Basidiomycota wood-degrading species, probably in interaction with bacteria (Bertsch

et al. 2013, Bruez et al. 2020, Haidar et al. 2021, 2024, Moretti et al. 2021). However,

less attention has been paid to the relationship between the total microbiome and the

type and extent of necrosis and to the role of Ascomycota fungi in white rot

propagation. We show here that, whatever the health status of the plant (i.e. control,

Esca or water deficit), the necrotic areas in the trunk harbored less diverse

Ascomycota fungal communities than adjacent apparently healthy areas, confirming

the results obtained for cordons (Bruez et al. 2020). These areas were dominated by

Ascomycota wood pathogens, the proportion of which increased at the expense of

other taxa, including putative antagonists, as previously highlighted in grapevine trunk

diseases (Martín et al. 2022), but not in other pathosystems (Bakys et al. 2009,

Denman et al. 2016). We found that *Phaeoacremonium minimum* was clearly

overrepresented in black necrotic areas, whereas *Phaeomoniella chlamydospora* was

not. *P. chlamydospora* was significantly overrepresented in white rot tissues, as

previously reported by Bruez et al. (2020) and Pacetti et al. (2021). The previously

favored hypothesis of a pioneering role of Ascomycota fungi before Basidiomycota

species in the sequence of events occurring during the colonization of healthy

grapevine wood has recently been revisited in light of new findings on potential intra-

and extracellular machinery for the detoxification of wood phenolic compounds in

Fomitiporia mediterranea (Moretti et al. 2021, Schilling et al. 2022, Garcia et al. 2024).

For *P. chlamydospora* a hypothetical role in wood decay through the opportunistic

colonization of highly degraded areas or co-degradation, possibly alongside

Basidiomycota, therefore appears more plausible than the pioneer hypothesis.

Moreover, *P. chlamydospora* may be less closely associated with black necrosis, as

previously suggested (e.g. Bertsch et al. 2013, Surico et al. 2006). It is, in any case, a

very abundant endophyte in healthy tissues (Del Frari et al. 2019, Hofstetter et al.

2012; Dell'Acqua et al. 2025). Finally, we resequenced healthy wood samples, using

ITS markers to quantify Basidiomycota and determine their relative abundance in

healthy wood samples; we confirmed that *Fomitiporia mediterranea* was present at

high abundance in the wood of plants expressing esca symptoms (Chambard et al.,

submitted).

We also characterized the Ascomycota community associated with the brown

stripe xylem observed in plants with symptomatic esca about three months after the

onset of leaf symptoms. We showed that the Ascomycota community in this

symptomatic tissue is diverse and similar to that retrieved from healthy woody tissues.

This finding is consistent with the fact that the brown stripe has not yet become a

necrotic area. This tissue no longer conducts water, as shown by the use of a dye to

follow sapwood transport (Lecomte et al. 2024), and in other vascular pathosystems

(Agrios 2005, Bibuci and Cirulli, 2012, Jiménez-Díaz et al. 2012). The brown color

probably results from the translocation of oxidizing phytotoxins and degradation

products and is associated with non-gaseous vascular embolism (e.g. tyloses and

gels; Agrios 2005, Bubici and Cirulli, 2012, Lecomte et al. 2024). We confirmed the

very high abundance of Botryosphaeriaceae spp. in brown stripe tissues late in the

season (Larignon et al. 2001), alongside *Cladosporium* spp. However, these

observations were performed three months after symptom onset. We cannot therefore

draw any firm conclusions about the changes in plant-fungus interactions potentially

leading to the co-occurrence of brown stripe and foliar symptoms. More detailed

studies of the microbial and physiological features related to the brown stripe

symptom, and of the temporal dynamics linking internal and external symptoms, are

now required.

Conclusion. Overall, this experimental study provides a new perspective on the ways

in which abiotic and biotic stresses can shape microbial communities in perennial

plants and contribute to plant decline. The trunk and younger grapevine organs had

different endophytic fungal communities with different responses to drought and esca.

Only esca significantly altered wood integrity. However, we were unable to investigate

the dual-stress situation, as water deficit inhibited the expression of esca foliar

symptoms. Based on these findings, assessments of the responses of the grapevine

holobiont to combinations of stresses at the physiological and microbiome levels

across various cultivars appear relevant to identify promising genotypes in the context

of climate change and perennial plant decline. Indeed, wood degradation (Arnold et

al. 2025), the incidence of esca foliar symptoms (Gastou et al. 2024), drought

tolerance (Dayer et al. 2022) and endophytic microbial communities (Bekris et al.

2021, Hamaoka et al. 2022) differ between grapevine genotypes. Finally, plant dieback

is related not only to endophytic communities, but also to microbiomes inhabiting the

soil compartment (Darriaut et al. 2021, 2023, Gómez-Aparicio et al. 2022, Lasa et al.

2024). Future holistic *in natura* microbiome studies (i.e. multidisciplinary,

multicompartments and/or multitaxa) would be particularly helpful, providing precise

and generalizable results on the response of the holobiont to stressful conditions.

Data availability. DNA sequencing datasets will be made available in the ENA

database under the accession number PRJEB87388.

Acknowledgments

We would like to thank Elena Farolfi and Jérôme Pouzoulet for help with plant

sectioning and wood sampling, and Corinne Vacher for advice on metabarcoding

analyses.

Funding

This work was supported by (1) the French Ministry of Agriculture and Food,

FranceAgriMer, and the *Comité National des Interprofessions des Vins à appellation*

d'origine et à indication géographique within the PHYSIOPATH Project (22001150-

1506; Program *Plan National Dépérissement du Vignoble*) and (2) the French

government in the framework of the IdEX Bordeaux University "Investments for the

Future" program / GPR Bordeaux Plant Sciences. PG was awarded a PhD by the

French *Ministère de l'Enseignement Supérieur et de la Recherche*.

Competing Interests statement

The authors have no competing interests to declare.

List of supplementary material

Supplementary Fig. S1: Pictures of the sampling

Supplementary Fig. S2: Properties of Ascomycota communities in all samples (i.e.

trunk, cane, stem and petiole samples) across organs.

Supplementary Fig. S3: Ascomycota community structure across wood types for

each plant status.

Supplementary Fig.S2: Properties of Ascomycota communities in all samples (i.e.

trunk, cane, stem and petiole samples) across organs.

Supplementary Table S1: Ascomycota genera classified as putative wood pathogens

and putative wood antagonists.

Supplementary Table S2: Global metrics describing the GTAA metabarcoding

dataset before and after the decontamination process.

Supplementary Table S3: OTU table corresponding to the entire decontaminated

GTAA metabarcoding dataset.

Supplementary Table S4: List of experimental factors and Ascomycota reads for

each sample included in the GTAA metabarcoding study.

Supplementary Table S5: Summary of the results obtained in Aldex2 differential

analyses in healthy trunk samples and young samples.

Supplementary Table S6: Statistical modelling of organ health status (S-AS vs S-S)

effects on alpha-diversity metrics and Ascomycota community structure computed for

young organs.

Supplementary Table S7: List of experimental factors and wood type percentages

for each plant

Supplementary Table S8: Statistical modelling of esca history (pA vs pS) effects on

the distribution of wood types and alpha-diversity metrics and Ascomycota community

structure computed for healthy trunk and young samples.

Tables and figures

Table 1. Number of *Vitis vinifera* cv. Sauvignon blanc plants and samples included in this study, classified by experimental plant status, organ and sample types. For esca plants, samples of petioles, current-year stems, and two-year-old canes were classified by organ status: asymptomatic (AS) or presenting tiger-stripe symptoms on the leaves (S). Plants were classified by esca history: previously asymptomatic (pA) or previously symptomatic (pS).

Plant status – esca history	Nb of plants	Organ	Nb of samples (per sample type or organ status)
Control – pA	$N = 11$	Trunk	$n = 22$ (10 healthy; 6 black necrosis; 6 white rot)
		Cane	$n = 9$
		Stem	$n = 9$
		Petiole	$n = 9$
Control – pS	$N = 10$	Trunk	$n = 29$ (11 healthy; 10 black necrosis; 8 white rot)
		Cane	$n = 10$
		Stem	$n = 11$
		Petiole	$n = 10$
Esca – pS	$N = 10$	Trunk	$n = 35$ (10 healthy; 9 brown stripe; 8 black necrosis; 8 white rot)
		Cane	$n = 9$ (8 AS; 1 S)
		Stem	$n = 14$ (4 AS; 10 S)
		Petiole	$n = 18$ (10 AS; 8 S)
Water deficit – pS	$N = 12$	Trunk	$n = 31$ (12 healthy; 12 black necrosis; 7 white rot)
		Cane	$n = 11$
		Stem	$n = 13$
		Petiole	$n = 11$

Fig. 1. Properties of endophytic Ascomycota communities in healthy trunk samples according to plant status

 A, Distribution of the 20 most abundant Ascomycota OTUs in trunk samples, by plant status. The mean relative abundance was calculated for each OTU, by plant status. *Stacked bars are colored according to the OTU. Less abundant OTUs are grouped in the category "Other" and are colored in light gray.* **B**, Effect of plant status on alpha-diversity metrics (i.e. observed richness, Shannon and Simpson indices) associated

with Ascomycota communities. **B**, PCA summarizing the structure of Ascomycota

communities according to an RDA on CLR-transformed data. Ellipses correspond to

the 95% interval for each group. **C**, Relative abundance of putative wood pathogens,

putative antagonists and other taxa, by plant status. All graphs are colored according

to plant status.

Fig. 2. Properties of endophytic and epiphytic Ascomycota communities in

young organs (i.e. petioles, stems and canes), by plant status.

A, Distribution of the 20 most abundant Ascomycota OTUs, by plant status, in young

samples. Mean relative abundances were calculated for each OTU, by plant status.

Stacked bars are colored according to the OTU. Less abundant OTUs are grouped

into the "Other" category and colored in light gray. **B**, Effect of plant status on alpha-

diversity metrics (i.e. observed richness, Shannon and Simpson indices) associated

with Ascomycota communities. **C**, PCA summarizing the structure of Ascomycota

communities according to an RDA on CLR-transformed data. Ellipses correspond to

the 95% interval for each group. Panels B and C are colored according to plant status.

Fig. 3. Relative proportions of wood symptoms in trunk cross-sections

according to plant status (control, esca expression, water deficit), and the

properties of the endophytic Ascomycota communities associated with each

wood type.

A, Mean proportions of healthy wood, black necrosis and white rot for each set of

experimental conditions. **B**, Effect of wood type on the alpha-diversity metrics (i.e.

observed richness, Shannon and Simpson indices) associated with Ascomycota

communities. **C**, PCA summarizing the structure of Ascomycota communities

according to an RDA on CLR-transformed data. Ellipses correspond to the 95%

interval for each group. **D**, OTUs displaying significant differences in abundance

between wood types according to a LeFse analysis. Features with an LDA score > 2

and a Kruskal-Wallis *p*-value < 0.05 are shown, and each bar is colored according to

the group in which the OTU is overrepresented. **E**, Relative abundance of putative

wood pathogens, putative antagonists and other taxa across wood types. All graphs

are colored according to wood type.

Fig. 4. Functional and taxonomic characterization of Ascomycota communities across wood types. A, Relative abundance of putative wood pathogens, putative antagonists and other taxa, by wood type. All graphs are colored according to wood type. **B**, OTUs with significantly different abundances between wood types according to a LeFse analysis. Features with an LDA score > 2 and a Kruskal-Wallis *p*-value < 0.05 are shown, and each bar is colored according to the group in which the OTU is overrepresented.

Literature cited

Agrios, G. N. 2005. *Plant Pathology, Fifth Edition*. Academic Press.

Arnault, G., Mony, C., and Vandenkoornhuyse, P. 2023. Plant microbiota dysbiosis and the Anna Karenina Principle. Trends in Plant Science 28:18–30. https://doi.org/10.1016/j.tplants.2022.08.012.

Arbizu PM. 2025. pmartinezarbizu/pairwiseAdonis.

Arnold, G., Prado, E., Dumas, V., Butterlin, G., Duchêne, É., Merdinoglu, D., and Avia, K. 2025. Discovery of a locus associated with susceptibility to esca in grapevine. Plant Disease. https://doi.org/10.1094/PDIS-06-24-1258-RE.

Bakys, R., Vasaitis, R., Barklund, P., Thomsen, I. M., and Stenlid, J. 2009. Occurrence and pathogenicity of fungi in necrotic and non-symptomatic shoots of declining common ash (*Fraxinus excelsior*) in Sweden. Eur J Forest Res 128:51–60. https://doi.org/10.1007/s10342-008-0238-2.

Bechtold, E. K., Ryan, S., Moughan, S. E., Ranjan, R., and Nüsslein, K. 2021. Phyllosphere Community Assembly and Response to Drought Stress on Common Tropical and Temperate Forage Grasses. Applied and Environmental Microbiology 87:e00895-21. https://doi.org/10.1128/AEM.00895-21.

Bekris, F., Vasileiadis, S., Papadopoulou, E., Samaras, A., Testempasis, S., Gkizi, D., Tavlaki, G., Tzima, A., Paplomatas, E., Markakis, E., Karaoglanidis, G., Papadopoulou, K. K., and Karpouzas, D. G. 2021. Grapevine wood microbiome analysis identifies key fungal pathogens and potential interactions with the bacterial community implicated in grapevine trunk disease appearance. Environmental Microbiome 16:23. https://doi.org/10.1186/s40793-021-00390-1.

Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., De Wit, P., Sánchez-García, M., Ebersberger, I., de Sousa, F., Amend, A., Jumpponen, A., Unterseher, M., Kristiansson, E., Abarenkov, K., Bertrand, Y. J. K., Sanli, K., Eriksson, K. M., Vik, U., Veldre, V., and Nilsson, R. H. 2013. Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of

environmental sequencing data. Methods in Ecology and Evolution 4:914–919.

https://doi.org/10.1111/2041-210X.12073.

Bertsch, C., Ramirez-Suero, M., Magnin-Robert, M., Larignon, P., Chong, J., Abou-Mansour,

E., Spagnolo, A., Clément, C., and Fontaine, F. 2013. Grapevine trunk diseases: complex and

still poorly understood. Plant Pathology 62:243–265. https://doi.org/10.1111/j.1365-

3059.2012.02674.x.

Bettenfeld, P., Fontaine, F., Trouvelot, S., Fernandez, O., and Courty, P.-E. 2020. Woody

Plant Declines. What's Wrong with the Microbiome? Trends in Plant Science 25:381–394.

https://doi.org/10.1016/j.tplants.2019.12.024.

Bokulich, N. A., Subramanian, S., Faith, J. J., Gevers, D., Gordon, J. I., Knight, R., Mills, D.

A., and Caporaso, J. G. 2013. Quality-filtering vastly improves diversity estimates from Illumina

amplicon sequencing. Nat Methods 10:57–59. https://doi.org/10.1038/nmeth.2276.

Bortolami, G., Farolfi, E., Badel, E., Burlett, R., Cochard, H., Ferrer, N., King, A., Lamarque,

L. J., Lecomte, P., Marchesseau-Marchal, M., Pouzoulet, J., Torres-Ruiz, J. M., Trueba, S.,

Delzon, S., Gambetta, G. A., and Delmas, C. E. L. 2021a. Seasonal and long-term

consequences of esca grapevine disease on stem xylem integrity. Journal of Experimental

Botany 72:3914–3928. https://doi.org/10.1093/jxb/erab117.

Bortolami, G., Ferrer, N., Baumgartner, K., Delzon, S., Gramaje, D., Lamarque, L. J.,

Romanazzi, G., Gambetta, G. A., and Delmas, C. E. L. 2023. Esca grapevine disease involves

leaf hydraulic failure and represents a unique premature senescence process. Tree

Physiology 43:441–451. https://doi.org/10.1093/treephys/tpac133.

Bortolami, G., Gambetta, G. A., Cassan, C., Dayer, S., Farolfi, E., Ferrer, N., Gibon, Y., Jolivet,

J., Lecomte, P., and Delmas, C. E. L. 2021b. Grapevines under drought do not express esca

leaf symptoms. Proceedings of the National Academy of Sciences 118:e2112825118.

https://doi.org/10.1073/pnas.2112825118.

Bortolami, G., Gambetta, G. A., Delzon, S., Lamarque, L. J., Pouzoulet, J., Badel, E., Burlett,

R., Charrier, G., Cochard, H., Dayer, S., Jansen, S., King, A., Lecomte, P., Lens, F., Torres-

Ruiz, J. M., and Delmas, C. E. L. 2019. Exploring the Hydraulic Failure Hypothesis of Esca

Leaf Symptom Formation. Plant Physiology 181:1163–1174.

https://doi.org/10.1104/pp.19.00591.

Broberg, M., Doonan, J., Mundt, F., Denman, S., and McDonald, J. E. 2018. Integrated multi-

omic analysis of host-microbiota interactions in acute oak decline. Microbiome 6:21.

https://doi.org/10.1186/s40168-018-0408-5.

Bruez, E., Vallance, J., Gautier, A., Laval, V., Compant, S., Maurer, W., Sessitsch, A., Lebrun,

M.-H., and Rey, P. 2020. Major changes in grapevine wood microbiota are associated with

the onset of esca, a devastating trunk disease. Environ Microbiol 22:5189–5206.

https://doi.org/10.1111/1462-2920.15180.

Bubici, G., and Cirulli, M. 2012. Control of *Verticillium* wilt of olive by resistant rootstocks. Plant

Soil 352:363–376. https://doi.org/10.1007/s11104-011-1002-9.

Cao Y, Dong Q, Wang D, Zhang P, Liu Y, Niu C. 2022. microbiomeMarker: an R/Bioconductor

package for microbiome marker identification and visualization. Bioinformatics 38: 4027–4029.

Carbone, M. J., Alaniz, S., Mondino, P., Gelabert, M., Eichmeier, A., Tekielska, D., Bujanda,

R., and Gramaje, D. 2021. Drought Influences Fungal Community Dynamics in the Grapevine

Rhizosphere and Root Microbiome. Journal of Fungi 7:686.

https://doi.org/10.3390/jof7090686.

Ciesla, W. M., and Donaubauer, E. 1994. *Decline and Dieback of Trees and Forests: A Global*

Overview. Food & Agriculture Org.

Claverie, M., Lecomte, P., Delorme, G., Dumot, V., Jacquet, O., and Cochard, H. 2023. Xylem

water transport is influenced by age and winter pruning characteristics in grapevine (*Vitis*

vinifera). OENO One 57:53–68. https://doi.org/10.20870/oeno-one.2023.57.3.7452.

Claverie, M., Notaro, M., Fontaine, F., and Wery, J. 2020. Current knowledge on Grapevine

Trunk Diseases with complex etiology: a systemic approach. Phytopathologia Mediterranea

59:29–53. https://doi.org/10.36253/phyto-11150.

Compant, S., Samad, A., Faist, H., and Sessitsch, A. 2019. A review on the plant microbiome:

Ecology, functions, and emerging trends in microbial application. Journal of Advanced

Research 19:29–37. https://doi.org/10.1016/j.jare.2019.03.004.

Darriaut, R. C., Marzari, T., Lailheugue, V., Tran, J., Martins, G., Marguerit, E., Isabelle, M.-

P., and Lauvergeat, V. 2024. Microbial dysbiosis in roots and rhizosphere of grapevines

experiencing decline is associated with active metabolic functions. Front. Plant Sci. 15.

https://doi.org/10.3389/fpls.2024.1358213.

Darriaut, R., Martins, G., Dewasme, C., Mary, S., Darrieutort, G., Ballestra, P., Marguerit, E.,

Vivin, P., Ollat, N., Masneuf-Pomarède, I., and Lauvergeat, V. 2021. Grapevine decline is

associated with difference in soil microbial composition and activity. OENO One 55:67.

https://doi.org/10.20870/oeno-one.2021.55.3.4626.

Darriaut, R., Tran, J., Martins, G., Ollat, N., Masneuf-Pomarède, I., and Lauvergeat, V. 2023.

In grapevine decline, microbiomes are affected differently in symptomatic and asymptomatic

soils. Applied Soil Ecology 183:104767. https://doi.org/10.1016/j.apsoil.2022.104767.

Dayer, S., Lamarque, L. J., Burlett, R., Bortolami, G., Delzon, S., Herrera, J. C., Cochard, H.,

and Gambetta, G. A. 2022. Model-assisted ideotyping reveals trait syndromes to adapt

viticulture   to   a   drier   climate.   Plant   Physiology   190:1673–1686.

https://doi.org/10.1093/plphys/kiac361.

Debray, R., Socolar, Y., Kaulbach, G., Guzman, A., Hernandez, C. A., Curley, R., Dhond, A.,

Bowles, T., and Koskella, B. 2022. Water stress and disruption of mycorrhizas induce parallel

shifts   in   phyllosphere   microbiome   composition.   New   Phytologist   234:2018–2031.

https://doi.org/10.1111/nph.17817.

de Mendiburu F. 2023. agricolae: Statistical Procedures for Agricultural Research.

Del Frari, G., Gobbi, A., Aggerbeck, M. R., Oliveira, H., Hansen, L. H., and Ferreira, R. B.

2019. Characterization of the Wood Mycobiome of *Vitis vinifera* in a Vineyard Affected by

Esca. Spatial Distribution of Fungal Communities and Their Putative Relation With Leaf

Symptoms. Frontiers in Plant Science 10.

Del Frari, G., Ingrà, C., Rønne Aggerbeck, M., Gobbi, A., Nascimento, T., Cabral, A., Oliveira,

H., Hestbjerg Hansen, L., Ferrandino, A., and Boavida Ferreira, R. 2025. Fungal community

dynamics and anthocyanin profiling of grapevine leaves in a vineyard affected by esca. Plant

Stress 15:100793. https://doi.org/10.1016/j.stress.2025.100793.

Del Frari, G., Oliveira, H., and Boavida Ferreira, R. 2021. White Rot Fungi (Hymenochaetales)

and Esca of Grapevine: Insights from Recent Microbiome Studies. J Fungi (Basel) 7:770.

https://doi.org/10.3390/jof7090770.

Dell'Acqua, N., Gambetta, G. A., Delzon, S., Ferrer, N., Lamarque, L. J., Saurin, N., Theodore,

P., and Delmas, C. E. L. 2024. Mechanisms of grapevine resilience to a vascular disease:

investigating stem radial growth, xylem development and physiological acclimation. Annals of

Botany 133:321–336. https://doi.org/10.1093/aob/mcad188.

Dell'Acqua N., Gambetta G.A., Comont G., Ferrer N., Rochepeau A., Petriacq P., Delmas

CEL. (2025) Nitrogen nutrition impacts grapevine esca leaf symptom incidence, physiology

and metabolism. Journal of Experimental Botany, eraf172 https://doi.org/10.1093/jxb/eraf172

Denman, S., Plummer, S., Kirk, S., Peace, A., and McDonald, J. E. 2016. Isolation studies

reveal a shift in the cultivable microbiome of oak affected with Acute Oak Decline. Systematic

and Applied Microbiology 39:484–490. https://doi.org/10.1016/j.syapm.2016.07.002.

Desprez-Loustau, M.-L., Marçais, B., Nageleisen, L.-M., Piou, D., and Vannini, A. 2006.

Interactive effects of drought and pathogens in forest trees. Ann. For. Sci. 63:597–612.

https://doi.org/10.1051/forest:2006040.

Elena, G., Bruez, E., Rey, P., and Luque, J. 2018. Microbiota of grapevine woody tissues with

or without esca-foliar symptoms in northeast Spain. Phytopathologia Mediterranea 57:425–

438.

Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S.,

Hernandez-Raquet, G., Combes, S., and Pascal, G. 2018. FROGS: Find, Rapidly, OTUs with

Galaxy Solution. Bioinformatics 34:1287–1294. https://doi.org/10.1093/bioinformatics/btx791.

Etienne, L., Fabre, F., Martinetti, D., Frank, E., Michel, L., Bonnardot, V., Guérin-Dubrana, L.,

and Delmas, C. E. L. 2024. Exploring the role of cultivar, year and plot age in the incidence of

esca and Eutypa dieback: Insights from 20 years of regional surveys in France. Plant

Pathology 73:2344–2358. https://doi.org/10.1111/ppa.13975.

Farooq, Q. U. A., Hardy, G. E. St. J., McComb, J. A., Thomson, P. C., and Burgess, T. I. 2022.

Changes to the Bacterial Microbiome in the Rhizosphere and Root Endosphere of *Persea*

americana (Avocado) Treated With Organic Mulch and a Silicate-Based Mulch or Phosphite,

and Infested With *Phytophthora cinnamomi*. Front Microbiol 13:870900.

https://doi.org/10.3389/fmicb.2022.870900.

Garcia, J. F., Figueroa-Balderas, R., Comont, G., Delmas, C. E. L., Baumgartner, K., and

Cantu, D. 2024. Genome analysis of the esca-associated Basidiomycetes *Fomitiporia*

mediterranea, *Fomitiporia polymorpha*, *Inonotus vitis*, and *Tropicoporus texanus* reveals

virulence factor repertoires characteristic of white-rot fungi. G3 Genes|Genomes|Genetics

14:jkae189. https://doi.org/10.1093/g3journal/jkae189.

Gastou, P., Irvine, A. D., Arcens, C., Courchinoux, E., This, P., Leeuwen, C. van, and Delmas,

C. 2024. Large gradient of susceptibility to esca disease revealed by long-term monitoring of

46 grapevine cultivars in a common garden vineyard. OENO One 58.

https://doi.org/10.20870/oeno-one.2024.58.2.8043.

Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., and Egozcue, J. J. 2017. Microbiome

Datasets Are Compositional: And This Is Not Optional. Front. Microbiol. 8.

https://doi.org/10.3389/fmicb.2017.02224.

Gómez-Aparicio, L., Domínguez-Begines, J., Villa-Sanabria, E., García, L. V., and Muñoz-

Pajares, A. J. 2022. Tree decline and mortality following pathogen invasion alters the diversity,

composition and network structure of the soil microbiome. Soil Biology and Biochemistry

166:108560. https://doi.org/10.1016/j.soilbio.2022.108560.

Gomez-Gallego, M., Galiano, L., Martínez-Vilalta, J., Stenlid, J., Capador-Barreto, H. D.,

Elfstrand, M., Camarero, J. J., and Oliva, J. 2022. Interaction of drought- and pathogen-

induced mortality in Norway spruce and Scots pine. Plant, Cell & Environment 45:2292–2305.

https://doi.org/10.1111/pce.14360.

Goodell, B. 2020. Fungi Involved in the Biodeterioration and Bioconversion of Lignocellulose

Substrates. In *Genetics and Biotechnology*, eds. J. Philipp Benz and Kerstin Schipper. Cham:

Springer International Publishing, pp. 369–397. https://doi.org/10.1007/978-3-030-49924-

2_15.

Gosling, R. H., Jackson, R. W., Elliot, M., and Nichols, C. P. 2024. Oak declines: Reviewing

the evidence for causes, management implications and research gaps. Ecological Solutions

and Evidence 5:e12395. https://doi.org/10.1002/2688-8319.12395.

Gramaje, D., Úrbez-Torres, J. R., and Sosnowski, M. R. 2018. Managing Grapevine Trunk

Diseases With Respect to Etiology and Epidemiology: Current Strategies and Future

Prospects. Plant Disease 102:12–39. https://doi.org/10.1094/PDIS-04-17-0512-FE.

Griffiths, S. M., Galambao, M., Rowntree, J., Goodhead, I., Hall, J., O'Brien, D., Atkinson, N.,

and Antwis, R. E. 2020. Complex associations between cross-kingdom microbial endophytes

and host genotype in ash dieback disease dynamics. Journal of Ecology 108:291–309.

https://doi.org/10.1111/1365-2745.13302.

Guilpart, N., Roux, S., Gary, C., and Metay, A. 2017. The trade-off between grape yield and

grapevine susceptibility to powdery mildew and grey mould depends on inter-annual variations

in water stress. Agricultural and Forest Meteorology 234–235:203–211.

https://doi.org/10.1016/j.agrformet.2016.12.023.

Haidar, R., Compant, S., Robert, C., Antonielli, L., Yacoub, A., Grélard, A., Loquet, A., Brader,

G., Guyoneaud, R., Attard, E., and Rey, P. 2024. Two *Paenibacillus* spp. strains promote

grapevine wood degradation by the fungus *Fomitiporia mediterranea*: from degradation

experiments to genome analyses. Sci Rep 14:15779. https://doi.org/10.1038/s41598-024-

66620-x.

Haidar, R., Yacoub, A., Vallance, J., Compant, S., Antonielli, L., Saad, A., Habenstein, B.,

Kauffmann, B., Grélard, A., Loquet, A., Attard, E., Guyoneaud, R., and Rey, P. 2021. Bacteria

associated with wood tissues of Esca-diseased grapevines: functional diversity and synergy

with *Fomitiporia mediterranea* to degrade wood components. Environ Microbiol 23:6104–

6121. https://doi.org/10.1111/1462-2920.15676.

Hallmann, J., Quadt-Hallmann, A., Mahaffee, W. F., and Kloepper, J. W. 1997. Bacterial

endophytes in agricultural crops. Can. J. Microbiol. 43:895–914. https://doi.org/10.1139/m97-

131.

Hamaoka, K., Aoki, Y., Takahashi, S., Enoki, S., Yamamoto, K., Tanaka, K., and Suzuki, S.

2022. Diversity of endophytic bacterial microbiota in grapevine shoot xylems varies depending

on wine grape-growing region, cultivar, and shoot growth stage. Sci Rep 12:15772.

https://doi.org/10.1038/s41598-022-20221-8.

Han, G., Mannaa, M., Jeon, H., Jung, H., Kim, J.-C., Park, A. R., and Seo, Y.-S. 2022.

Dysbiosis in the Rhizosphere Microbiome of Standing Dead Korean Fir (*Abies koreana*).

Plants 11:990. https://doi.org/10.3390/plants11070990.

Hofstetter, V., Buyck, B., Croll, D., Viret, O., Couloux, A., and Gindro, K. 2012. What if esca

disease of grapevine were not a fungal disease? Fungal Diversity 54:51–67.

https://doi.org/10.1007/s13225-012-0171-z.

Hrycan, J., Bowen, P., Forge, T., Hart, M., and Úrbez-Torres, J. R. 2025. Impact of water

stress on *Phaeomoniella chlamydospora* abundance and Petri disease symptom development

in young grapevines. OENO One 59. https://doi.org/10.20870/oeno-one.2025.59.1.8317.

Jactel, H., Petit, J., Desprez-Loustau, M.-L., Delzon, S., Piou, D., Battisti, A., and Koricheva,

J. 2012. Drought effects on damage by forest insects and pathogens: a meta-analysis. Global

Change Biology 18:267–276. https://doi.org/10.1111/j.1365-2486.2011.02512.x.

Jaeger, A. C. H., Hartmann, M., Conz, R. F., Six, J., and Solly, E. F. 2024. Drought-induced

tree mortality in Scots pine mesocosms promotes changes in soil microbial communities and

trophic groups. Applied Soil Ecology 194:105198.

https://doi.org/10.1016/j.apsoil.2023.105198.

Jia, S., Yuan, T., Fu, Y., Penuelas, J., Zhou, G., Zhou, L., Liu, D., He, Y., Liu, R., Wang, X.,

Song, B., Jiang, Z., and Zhou, X. 2024. Fungi and bacteria trade-off mediates drought-induced

reduction in wood decomposition. CATENA 243:108169.

https://doi.org/10.1016/j.catena.2024.108169.

Jiao, Z., Gao, Z., Liao, Y., Liu, Y., Dong, L., and Sun, H. 2023. Effects of Pine Wilt Disease on

Rhizosphere Microbiota and Fine Root Fungi: Insights into Enzyme Activity, Ectomycorrhizal

Infection and Microbial Composition. Forests 14:1884. https://doi.org/10.3390/f14091884.

Jiménez-Díaz, R. M., Cirulli, M., Bubici, G., del Mar Jiménez-Gasco, M., Antoniou, P. P., and

Tjamos, E. C. 2012. Verticillium Wilt, A Major Threat to Olive Production: Current Status and

Future Prospects for its Management. Plant Disease 96:304–329.

https://doi.org/10.1094/PDIS-06-11-0496.

Larignon, P., and Dubos, B. 1997. Fungi associated with esca disease in grapevine. European

Journal of Plant Pathology 103:147–157. https://doi.org/10.1023/A:1008638409410.

Larignon, P., Fulchic, R., Cere, L., and Dubos, B. 2001. Observation on black dead arm in

French vineyards. Phytopathologia Mediterranea 40:S336–S342.

Lasa, A. V., Fernández-González, A. J., Villadas, P. J., Mercado-Blanco, J., Pérez-Luque, A.

J., and Fernández-López, M. 2024. Mediterranean pine forest decline: A matter of root-

associated microbiota and climate change. Science of The Total Environment 926:171858.

https://doi.org/10.1016/j.scitotenv.2024.171858.

Leal, C., Bujanda, R., Carbone, M. J., Kiss, T., Eichmeier, A., Gramaje, D., and Maldonado-

González, M. M. 2024. Drought Influences the Structure, Diversity, and Functionality of the

Fungal Community Inhabiting the Grapevine Xylem and Enhances the Abundance of

Phaeomoniella chlamydospora. Phytobiomes Journal PBIOMES-02-24-0027-R.

https://doi.org/10.1094/PBIOMES-02-24-0027-R.

Lecomte, P., Bénétreau, C., Diarra, B., Meziani, Y., Delmas, C., and Fermaud, M. 2024.

Logistic modeling of summer expression of esca symptoms in tolerant and susceptible

cultivars in Bordeaux vineyards. OENO One 58. https://doi.org/10.20870/oeno-

one.2024.58.1.7571.

Lecomte, P., Darrieutort, G., Liminana, J.-M., Comont, G., Muruamendiaraz, A., Legorburu,

F.-J., Choueiri, E., Jreijiri, F., El Amil, R., and Fermaud, M. 2012. New Insights into Esca of

Grapevine: The Development of Foliar Symptoms and Their Association with Xylem

Discoloration. Plant Disease 96:924–934. https://doi.org/10.1094/PDIS-09-11-0776-RE.

van Leeuwen, C., Sgubin, G., Bois, B., Ollat, N., Swingedouw, D., Zito, S., and Gambetta, G.

A. 2024. Climate change impacts and adaptations of wine production. Nat Rev Earth Environ

5:258–275. https://doi.org/10.1038/s43017-024-00521-5.

Lin, T., Tang, J., Li, S., Li, S., Han, S., Liu, Y., Yang, C., Chen, G., Chen, L., and Zhu, T. 2023.

Drought stress-mediated differences in phyllosphere microbiome and associated pathogen

resistance between male and female poplars. The Plant Journal 115:1100–1113.

https://doi.org/10.1111/tpj.16283.

Luard, E. J., and Griffin, D. M. 1981. Effect of water potential on fungal growth and turgor.

Transactions of the British Mycological Society 76:33–40. https://doi.org/10.1016/S0007-

1536(81)80006-X.

Luna, I. J., Santiago, L., Ezcurra, E., Xi, M., Ashworth, V. E. T. M., Nothnagel, E., and

Rolshausen, P. E. 2025. Physiological symptoms induced by drought stress outweigh vascular

pathogen infection in walnut. Tree Physiology 45:tpaf034.

https://doi.org/10.1093/treephys/tpaf034.

Ma, Y., Shen, Y., Zhou, X., Ma, H., Lan, J., Fu, B., and Xue, Q. 2024. Biological Decline of

Alfalfa Is Accompanied by Negative Succession of Rhizosphere Soil Microbial Communities.

Plants 13:2589. https://doi.org/10.3390/plants13182589.

Magnin-Robert, M., Adrian, M., Trouvelot, S., Spagnolo, A., Jacquens, L., Letousey, P.,

Rabenoelina, F., Harir, M., Roullier-Gall, C., Clément, C., Schmitt-Kopplin, P., Vallat, A., Abou-

Mansour, E., and Fontaine, F. 2017. Alterations in Grapevine Leaf Metabolism Occur Prior to

Esca Apoplexy Appearance. MPMI 30:946–959. https://doi.org/10.1094/MPMI-02-17-0036-R.

Magnin-Robert, M., Spagnolo, A., Boulanger, A., Joyeux, C., Clément, C., Abou-Mansour, E.,

and Fontaine, F. 2016. Changes in Plant Metabolism and Accumulation of Fungal Metabolites

in Response to Esca Proper and Apoplexy Expression in the Whole Grapevine.

Phytopathology® 106:541–553. https://doi.org/10.1094/PHYTO-09-15-0207-R.

Mahé, F., Rognes, T., Quince, C., Vargas, C. de, and Dunthorn, M. 2014. Swarm: robust and

fast clustering method for amplicon-based studies. PeerJ 2:e593.

https://doi.org/10.7717/peerj.593.

Maher, N., Piot, J., Bastien, S., Vallance, J., Rey, P., and Guérin-Dubrana, L. 2012. Wood

necrosis in esca-affected vines: types, relationships and possible links with foliar symptom

expression. OENO One 46:15–27. https://doi.org/10.20870/oeno-one.2012.46.1.1507.

Manion, P. D. 1981. *Tree disease concepts*. Englewood Cliffs, N.J: Prentice-Hall.

Marsberg, A., Kemler, M., Jami, F., Nagel, J. H., Postma-Smidt, A., Naidoo, S., Wingfield, M.

J., Crous, P. W., Spatafora, J. W., Hesse, C. N., Robbertse, B., and Slippers, B. 2017.

Botryosphaeria dothidea: a latent pathogen of global importance to woody plant health.

Molecular Plant Pathology 18:477–488. https://doi.org/10.1111/mpp.12495.

Martín, L., García-García, B., and Alguacil, M. del M. 2022. Interactions of the Fungal

Community in the Complex Patho-System of Esca, a Grapevine Trunk Disease. International

Journal of Molecular Sciences 23:14726. https://doi.org/10.3390/ijms232314726.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet.journal 17:10–12. https://doi.org/10.14806/ej.17.1.200.

McElrone, A. J., Manuck, C. M., Brodersen, C. R., Patakas, A., Pearsall, K. R., and Williams,

L. E. 2021. Functional hydraulic sectoring in grapevines as evidenced by sap flow, dye

infusion, leaf removal and micro-computed tomography. AoB PLANTS 13:plab003.

https://doi.org/10.1093/aobpla/plab003.

McMurdie, P. J., and Holmes, S. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLOS ONE 8:e61217.

https://doi.org/10.1371/journal.pone.0061217.

Monod, V., Hofstetter, V., Viret, O., Zufferey, V., Gindro, K., and Croll, D. 2024. Landscape-

scale endophytic community analyses in replicated grapevine stands reveal that dieback

disease is not caused by specific fungal communities. 2024.04.05.588363.

https://doi.org/10.1101/2024.04.05.588363.

Morales-Cruz, A., Allenbeck, G., Figueroa-Balderas, R., Ashworth, V. E., Lawrence, D. P.,

Travadon, R., Smith, R. J., Baumgartner, K., Rolshausen, P. E., and Cantu, D. 2018a. Closed-

reference metatranscriptomics enables in planta profiling of putative virulence activities in the

grapevine trunk disease complex. Molecular Plant Pathology 19:490–503.

https://doi.org/10.1111/mpp.12544.

Morales-Cruz, A., Figueroa-Balderas, R., García, J. F., Tran, E., Rolshausen, P. E.,

Baumgartner, K., and Cantu, D. 2018b. Profiling grapevine trunk pathogens in planta: a case

for community-targeted DNA metabarcoding. BMC Microbiology 18:214.

https://doi.org/10.1186/s12866-018-1343-0.

Moretti, S., Goddard, M.-L., Puca, A., Lalevée, J., Di Marco, S., Mugnai, L., Gelhaye, E.,

Goodell, B., Bertsch, C., and Farine, S. 2023. First Description of Non-Enzymatic Radical-

Generating Mechanisms Adopted by *Fomitiporia mediterranea*: An Unexplored Pathway of the

White Rot Agent of the Esca Complex of Diseases. Journal of Fungi 9:498.

https://doi.org/10.3390/jof9040498.

Moretti, S., Pacetti, A., Pierron, R., Kassemeyer, H.-H., Fischer, M., Péros, J.-P., Perez-

Gonzalez, G., Bieler, E., Schilling, M., Marco, S. D., Gelhaye, E., Mugnai, L., Bertsch, C., and

Farine, S. 2021. Fomitiporia mediterranea M. Fisch., the historical Esca agent: a

comprehensive review on the main grapevine wood rot agent in Europe. Phytopathologia

Mediterranea 60:351–379. https://doi.org/10.36253/phyto-13021.

Nerva, L., Garcia, J. F., Favaretto, F., Giudice, G., Moffa, L., Sandrini, M., Cantu, D., Zanzotto,

A., Gardiman, M., Velasco, R., Gambino, G., and Chitarra, W. 2022. The hidden world within

plants: metatranscriptomics unveils the complexity of wood microbiomes. J Exp Bot 73:2682–

2697. https://doi.org/10.1093/jxb/erac032.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos

P, Stevens MHH, Szoecs E, et al. 2022. vegan: Community Ecology Package.

Ouadi, L., Bruez, E., Bastien, S., Vallance, J., Lecomte, P., Domec, J.-C., and Rey, P. 2019.

Ecophysiological impacts of Esca, a devastating grapevine trunk disease, on *Vitis vinifera* L.

PLOS ONE 14:e0222586. https://doi.org/10.1371/journal.pone.0222586.

Pacetti, A., Moretti, S., Pinto, C., Compant, S., Farine, S., Bertsch, C., and Mugnai, L. 2021.

Trunk Surgery as a Tool to Reduce Foliar Symptoms in Diseases of the Esca Complex and

Its Influence on Vine Wood Microbiota. Journal of Fungi 7:521.

https://doi.org/10.3390/jof7070521.

Raghavendra, A. K. H., Bissett, A. B., Thrall, P. H., Morin, L., Steinrucken, T. V., Galea, V. J.,

Goulter, K. C., and van Klinken, R. D. 2017. Characterisation of above-ground endophytic and

soil fungal communities associated with dieback-affected and healthy plants in five exotic

invasive species. Fungal Ecology 26:114–124. https://doi.org/10.1016/j.funeco.2017.01.003.

Ramegowda, V., and Senthil-Kumar, M. 2015. The interactive effects of simultaneous biotic

and abiotic stresses on plants: mechanistic understanding from drought and pathogen

combination. J Plant Physiol 176:47–54. https://doi.org/10.1016/j.jplph.2014.11.008.

Riou, C., Agostini, D., Aigrain, P., Barthe, M., Robert, M.-L. des, Gervais, J.-P., Jobard, E.,

Lurton, L., Moncomble, D., and Prêtet-Lataste, C. 2016. Action plan against declining

vineyards: An innovative approach. BIO Web Conf. 7:01040.

https://doi.org/10.1051/bioconf/20160701040.

Schilling, M., Maia-Grondard, A., Baltenweck, R., Robert, E., Hugueney, P., Bertsch, C.,

Farine, S., and Gelhaye, E. 2022. Wood degradation by *Fomitiporia mediterranea* M. Fischer:

Physiologic, metabolomic and proteomic approaches. Front Plant Sci 13:988709.

https://doi.org/10.3389/fpls.2022.988709.

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. 2012. NIH Image to ImageJ: 25 years of

image analysis. Nat Methods 9:671–675. https://doi.org/10.1038/nmeth.2089.

Skovsgaard, J. P., Thomsen, I. M., Skovgaard, I. M., and Martinussen, T. 2010. Associations

among symptoms of dieback in even-aged stands of ash (*Fraxinus excelsior* L.). Forest

Pathology 40:7–18. https://doi.org/10.1111/j.1439-0329.2009.00599.x.

Solís-García, I. A., Ceballos-Luna, O., Cortazar-Murillo, E. M., Desgarennes, D., Garay-

Serrano, E., Patiño-Conde, V., Guevara-Avendaño, E., Méndez-Bravo, A., and Reverchon, F.

2021. Phytophthora Root Rot Modifies the Composition of the Avocado Rhizosphere

Microbiome and Increases the Abundance of Opportunistic Fungal Pathogens. Front.

Microbiol. 11. https://doi.org/10.3389/fmicb.2020.574110.

Steinrucken, T. V., Bissett, A., Powell, J. R., Raghavendra, Anil. K. H., and van Klinken, R. D.

2016. Endophyte community composition is associated with dieback occurrence in an invasive

tree. Plant Soil 405:311–323. https://doi.org/10.1007/s11104-015-2529-y.

Surico, G., Mugnai, L., and Marchi, G. 2006. Older and More Recent Observations on Esca :

A   Critical   Overview.   Phytopathologia   Mediterranea   68–86.

https://doi.org/10.14601/Phytopathol_Mediterr-1847.

Torres-Ruiz, J. M., Cochard, H., Delzon, S., Boivin, T., Burlett, R., Cailleret, M., Corso, D.,

Delmas, C. E. L., De Caceres, M., Diaz-Espejo, A., Fernández-Conradi, P., Guillemot, J.,

Lamarque, L. J., Limousin, J.-M., Mantova, M., Mencuccini, M., Morin, X., Pimont, F., De Dios,

V. R., Ruffault, J., Trueba, S., and Martin-StPaul, N. K. 2024. Plant hydraulics at the heart of

plant, crops and ecosystem functions in the face of climate change. New Phytologist 241:984–

999. https://doi.org/10.1111/nph.19463.

Trivedi, P., Batista, B. D., Bazany, K. E., and Singh, B. K. 2022. Plant–microbiome interactions

under a changing world: responses, consequences and perspectives. New Phytologist

234:1951–1959. https://doi.org/10.1111/nph.18016.

Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., and Singh, B. K. 2020. Plant–microbiome

interactions: from community assembly to plant health. Nat Rev Microbiol 18:607–621.

https://doi.org/10.1038/s41579-020-0412-1.

Weiller, F., Diniz, I., Pimentel, D., Erban, A., Reis, P., Soares, F., Rego, C., Kopka, J., and

Fortes, A. M. 2024. Metabolomic analysis of grapes and leaves from symptomatic and

asymptomatic *Vitis vinifera* grapevines with Esca disease. Current Plant Biology 40:100378.

https://doi.org/10.1016/j.cpb.2024.100378.

Xu, L., Naylor, D., Dong, Z., Simmons, T., Pierroz, G., Hixson, K. K., Kim, Y.-M., Zink, E. M.,

Engbrecht, K. M., Wang, Y., Gao, C., DeGraaf, S., Madera, M. A., Sievert, J. A., Hollingsworth,

J., Birdseye, D., Scheller, H. V., Hutmacher, R., Dahlberg, J., Jansson, C., Taylor, J. W.,

Lemaux, P. G., and Coleman-Derr, D. 2018. Drought delays development of the sorghum root

microbiome and enriches for monoderm bacteria. Proceedings of the National Academy of

Sciences 115:E4284–E4293. https://doi.org/10.1073/pnas.1717308115.

Zabel, R. A., and Morrell, J. J. 2012. *Wood Microbiology: Decay and Its Prevention*. Academic

Press.

Zinger, L., Lionnet, C., Benoiston, A.-S., Donald, J., Mercier, C., and Boyer, F. 2021. metabaR:

An r package for the evaluation and improvement of DNA metabarcoding data quality.

Methods in Ecology and Evolution 12:586–592. https://doi.org/10.1111/2041-210X.13552.