E. coli Nissle 1917 occupies previously undocumented host niches in the insect-

parasitic nematode *Steinernema hermaphroditum*

Victoria Chen[1,2#], John P. Marken[2#], Richard M. Murray[2], Mengyi Cao[2,3*]

[1] Department of Biological Engineering, Massachusetts Institute of Technology,

Cambridge, MA 02139

[2] Division of Biology and Biological Engineering, California Institute of Technology,

Pasadena, CA 91125

[3] Division of Biosphere Sciences and Engineering, Carnegie Institution for Science,

Pasadena, CA 91125

Running title: *Steinernema* nematode and *E. coli* Nissle interaction

Keywords: *E. coli* Nissle 1917, entomopathogenic, nematode, *Steinernema*

hermaphroditum, coelomocyte, intestine, cuticle

[#]Denotes equal contributions

[*]To whom correspondence may be addressed:

Mengyi Cao: mcao@carnegiescience.edu, mengyi.cao77@gmail.com

Co-authors:

Victoria Chen: vichen@mit.edu

John P. Marken: jmarken@caltech.edu

Richard M. Murray: murray@cds.caltech.edu

Data Availability Statement: all data supporting the findings of this study are available

within the manuscript and its supplementary materials.

Ethics Statement: experiments involving animals were conducted in accordance with

institutional and national guidelines under approved protocols.

Conflict of Interest: the authors declare no competing interests.

Authors contributions:

MC and JPM oversaw and conceptualized the project. VC, JPM, and MC acquired

preliminary data on *E. coli* colonization in *Steinernema* nematode cuticle that led to the

initiation of the project. VC, JPM, and MC designed the experiments. VC and MC

performed experiments, analyzed the data, and wrote the manuscript. JPM and RMM

edited the manuscript. MC and RMM supervised the project. RMM, JPM, and MC

acquired funding.

Acknowledgements

We thank Tara Chari for sharing the *E. coli* Nissle strain with constitutive mScarlet-I

reporter used in this research. We thank Heidi Goodrich-Blair, Jennifer Heppert, and

Omar Alani for sharing the *Xenorhabdus* bacterial strains used in this research. We

thank Paul Sternberg for providing space and equipment support.

Funding source:

MC, VC and JPM were supported in part by the Resnick Sustainability Institute. MC was

also supported by Carnegie Institution for Science. JPM and RMM were also supported

in part by the Institute for Collaborative Biotechnologies through contract W911NF-19-

D-0001 from the U.S. Army Research Office. The content of the information on this

page does not necessarily reflect the position or the policy of the Government, and no

official endorsement should be inferred.

Abstract

Steinernema species are soil-dwelling and insect-parasitic nematodes that associate

with symbiotic *Xenorhabdus* bacteria. During the infective juvenile (IJ) stage,

Steinernema nematodes package species-specific *Xenorhabdus* bacteria in the anterior

intestinal pockets. The nematodes can survive in the soil for months as they seek insect

prey. The mechanisms of how these nematodes associate with environmental microbes

other than their *Xenorhabdus* symbionts is barely known. Here, we report a new

mechanism of *E. coli* Nissle (EcN) association with the nematode

Steinernema hermaphroditum. We show that EcN cells are enclosed and lysed in at

least four pairs of coelomocytes, suggesting these immune cells respond to bacterial

invasion. During the IJ stage of nematode development, EcN cells localize to posterior

intestinal vacuoles and enter the inter-cuticular space, where they proliferate,

aggregate, then lyse. EcN cells expressed proteins in the cell lysates were maintained

in the nematode cuticle over eight weeks in non-sterile soil. These observations suggest

sequential steps of EcN colonization in the host nematodes involving an immune

response that is distinctive from interactions with mutualistic symbiont. Our work

establishes a novel framework of nematode-bacteria interaction with potential

applications in environmental bioengineering.

Introduction

Animal tissues create complex, compartmentalized spaces that provide abiotic

and biotic environments capable of selecting and maintaining mutualistic

microorganisms while antagonizing pathogenic ones. Reciprocally, microbes develop

within these host-derived environments, shaping the identity of the niches they occupy.

This process of microbial modulation of the local environment, known as niche

construction, can drive long-term coevolution. In contrast, pathogens can exploit

unoccupied niches, leading to opportunistic infections and niche deconstruction

(Baquero *et al.*, 2021). Identifying and characterizing novel niches within animal hosts,

as well as the dynamic interactions of microbes within these microenvironments, are

crucial for understanding the mechanisms and consequences of host-microbe crosstalk.

Binary symbioses, involving a single core symbiont and a single animal species, provide

efficient models for revealing the temporal and spatial dynamics of symbiont

colonization within specific host cells and tissues. The niche-specific interactions

observed in diverse binary symbiosis models are critical for host development and

ecosystem health (Nyholm and McFall-Ngai, 1998; Stilwell *et al.*, 2018; Maire *et al.*,

2020; Tivey *et al.*, 2022). While binary symbiosis often involves colonization by a single

microbial species within a specialized organ, additional genera or species can

sometimes associate with distinct tissues, maintaining spatial separation from primary

symbionts (Dale and Moran, 2006; Nyholm and McFall-Ngai, 2021). The relatively

simple composition of these natural microbiomes makes such animal models

advantageous for studying multiple types of host-microbe interactions simultaneously.

The entomopathogenic (EPN, insect-parasitic) nematode *Steinernema spp*

maintains a stable association with mutualistic *Xenorhabdus* bacteria in a binary

symbiosis. Each generation of nematodes acquires their symbiotic bacteria from the

external environment (semi-horizontal) while a maternal factor can influence the

bacterial colonization of the next generation (semi-vertical). The intestine of

Steinernema nematodes contains compartmentalized niches where multiple

microenvironments enable the attachment, persistence, and proliferation of

Xenorhabdus bacteria (Chaston *et al.*, 2013) (Fig. 1A and Fig. 1B). *Xenorhabdus*

bacteria first attach to the anterior intestinal caecum (AIC) in juveniles, while nematodes

feed on bacteria, molt through four juvenile stages (J1-J4) and become adults to

reproduce. Signals from overcrowding and lack of food source induce the J2 stage of

juveniles to molt into IJs through an alternative developmental pathway. During IJ

development, the symbiotic bacteria first localize to the pharyngeal intestinal valves

(PIV) in the pre-IJs. These narrow pouches restrict bacterial colonization to a few cells

per animal. During the infective juvenile (IJ) stage, symbiotic bacteria migrate into the

anterior intestinal pocket, termed the receptacle (Fig. 1B). The multiple intestinal tissues

involved in the colonization events, including AIC, PIV, and the receptacle, are

proposed to provide specific nutrients, signaling molecules, and physical restrictions to

select the specific species or strain of *Xenorhabdus* bacteria.

Steinernema are soil-dwelling nematodes that also encounter other species of

microbes from their natural environment. Bacteria from multiple genera, termed the

'pathobiome', are found to frequently associate with certain *Steinernema* spp and

potentially contribute to entomopathogenic (insect-killing) activities (Ogier *et al.*, 2020).

Despite its generalization of pathobiomes among multiple *Steinernema* species, the

frequency, localization, and molecular mechanisms of pathobiome association is

unknown. Little else is known about other environmental microbes -- either pathogenic,

mutualistic, or commensal -- that interact with EPNs. This knowledge gap limits the full

application of *Steinernema* as a symbiosis model and its applications in bioengineering.

Therefore, understanding the spatial and temporal interactions of pathogens and

commensals in *Steinernema* provides a foundational framework for novel modes of

symbiotic interactions.

Escherichia coli is the most established bacterial species for molecular biology

and is a leading candidate for bioengineering applications across diverse niches,

including biosensing in complex microenvironments (Del Valle *et al.*, 2021) and

modifying gene expression in multicellular organisms (Gao and Sun, 2021). *E. coli* is

commonly associated with the animal intestinal tract (host-dependent) and circulates

through the soil and water environment (host-independent) via a biphasic lifestyle (Fig.

1C). Certain *E. coli* strains can persist in soil environments, where they influence

indigenous microflora encountered by insects and soil-dwelling nematodes (Elsas *et al.*,

2010). The *E. coli* Nissle 1917 (EcN) strain, originally isolated from a soldier who

survived a *Shigella* endemic, is known to interact with the mammalian intestinal

epithelium and mucosal immune system in a non-pathogenic manner, antagonize

pathogenic enterobacteria, and confer probiotic benefits in humans (Sonnenborn,

2016). In model nematode *Caenorhabditis elegans* adults, EcN as food source was

observed to show both probiotic and neurodegenerative effects (Kim and Moon, 2019;

Redweik and Xue, 2025) . Engineered EcN were used to carry biosensors that monitor

the intestinal environment of the animal (Rutter *et al.*, 2019). However, the interactions

of EcN with nematode species beyond *C. elegans* is not explored. Understanding the

interactions between EcN and *Steinernema* are therefore useful for establishing a

foundation for potential synthetic biology solutions to challenges in host-microbe

interactions and application in the soil environment (Jones *et al.*, 2023) .

In this study, we investigated the interactions between *E. coli* Nissle and the

entomopathogenic nematode *Steinernema hermaphroditum*. We discovered that EcN

localizes within previously undocumented cell and tissue types in *S. hermaphroditum*,

including immune cells (coelomocytes), posterior intestinal tissues, and inter-cuticular

spaces. These snapshots of EcN localization in nematodes suggest a sequential

process of cell- and tissue-specific microbe-host interactions synchronized with infective

juvenile (IJ) development. Notably, EcN colonization occurs in spatially distinct tissues

from those occupied by *Xenorhabdus*, the nematode's native symbiont. Our findings

established *E. coli* Nissle–*S. hermaphroditum* interactions as a model to monitor niche-

specific microbial interactions. As an emerging genetic model organism,

S. hermaphroditum offers a promising delivery system for engineered biosensors due to

its transparency, short life cycle in vitro, and bacterivorous nature (Bracho *et al.*, 2016;

Hwang *et al.*, 2017; Rutter *et al.*, 2019).

Experimental Procedures

Construction of fluorescent *E. coli* Nissle strains with mScarlet-I. Genetic

constructs for constitutive expression of red fluorescent protein (mScarlet-I) was

assembled using the 3G assembly method (Halleran *et al.*, 2018) with standard parts

from the MoClo modular cloning toolkit (Iverson *et al.*, 2016). The resulting constructs

were cloned into the "KL" genome integration vector from the pOSIP one-step genome

integration system (St-Pierre *et al.*, 2013) , which facilitates integration at the lambda

integrase attachment site in the *E. coli* genome. Plasmids were transformed into

chemically competent *E. coli* strains (Nissle 1917, CSH50 (Sokurenko *et al.*, 1992) , and

BW25113 (Grenier *et al.*, 2014)) using a standard heat shock protocol. Briefly, 50 µL of

competent cells were mixed with 100 ng of plasmid DNA, incubated on ice for 30

minutes, heat shocked at 42 °C for 45 seconds, and then placed back on ice for 2

minutes. Cells were allowed to recover in 950 µL of SOC medium at 37 °C with

shaking for 1 hour before being plated on LB agar plates containing kanamycin (50

µg/mL) to select for successful integrants. Plates were incubated at 37 °C overnight.

Resulting transformants were screened via colony PCR on the integration locus using

the standard screening primers from (St-Pierre *et al.*, 2013) and Sanger sequenced to

validate the successful integration.

Liver-kidney agar plates. The recipe for Liver-kidney agar was adapted from (Flores-

Lara *et al.*, 2007): 50 g of pork liver and 50 g of pork kidney were blended in 250 mL

distilled water until smooth. 2.5 g of NaCl, 7.5 g agar, 250 mL of water, and the liver-

kidney solution was transferred to a flask, then autoclaved and cooled in a 55°C water

bath. To avoid contamination, the liver-kidney agar was supplemented with kanamycin

(50 µg/mL) and poured into 60 x 15mm Petri plates under a fume hood. The agar was

continuously swirled before pouring to avoid pouring any larger chunks.

***S. hermaphroditum* axenic egg extraction and colonization assay.** Conventional

S. hermaphroditum nematodes were maintained by propagating through

Galleria mellonela 5[th] instar larvae (Cao *et al.*, 2021). The method for axenic egg

extraction for *S. hermaphroditum* was adapted from (Murfin *et al.*, 2012). Liquid cultures

of symbiotic bacterium *X. griffiniae* (HGB2511) were grown overnight in LB media that

were kept in the dark and added to lipid agar plates (Vivas and Goodrich-Blair, 2001;

Cao *et al.*, 2021). Plates were cultured for 48 hours to form a bacterial lawn. To prepare

axenic (germ-free) eggs, conventional IJs (exclusively propagated through insects) were

added to the lawns of *X. griffiniae* (HGB2511) and incubated for 2-3 days to grow to

gravid adults. Nematodes were washed off the bacterial lawn with water and then

transferred to 50 mL conical tubes (Falcon, Corning, NY) and centrifuged at 3000 rpm

for 2 minutes. The supernatant was removed using a vacuum, and 2 mL of a lysis

solution (2.4 mL 8% bleach, 500 µL 5M KOH, 7.1 mL water) was added to lyse the

gravid adult hermaphrodites. Nematodes were incubated in the lysis solution for 5

minutes at room temperature and mixed several times per minute by inverting. The

solution was then centrifuged at 3000 rpm for two minutes, and the lysis solution was

removed using a vacuum. Nematode eggs were washed thrice with 10 mL of LB

medium. Their density was assessed by viewing the number of IJs in 2 µL spots under a

stereoscope.

For colonization assays, approximately 500 eggs were added to the prepared liver-

kidney or lipid agar plates, with or without bacterial lawn of *X. griffiniae* expressing GFP

(Thomas *et al.*, 2024) or *E. coli* expressing mScarlet-I and incubated at 25℃ for 1 week.

Plates were then moved to 100 x 25mm Petri dishes filled with 10-15 mL of water to trap

infective juveniles. Three biological replicates of each colonization condition were

screened to identify colonization phenotypes based on the presence of EcN in different

tissues. Colonization phenotypes and frequencies were assessed using fluorescence

microscopy (See Microscopy and Image Acquisition for details).

Calculation of CFU per IJ. The CFU per IJ assessment was adapted from (Murfin *et*

al., 2019). IJ suspension from water-trap was added to a sterile 1.5 mL microfuge tubes

(Eppendorf, Germany) and centrifuged at 634 g for two minutes to concentrate the

nematodes. The supernatant was removed using a vacuum, and 1 mL of a 1% bleach

solution was added to the pellet, which was incubated for 2 min at room temperature to

surface sterilize IJs. IJs were washed thrice with and resuspended in 1 mL of LB

medium, and their concentration was assessed by viewing 2 μL spots under a

stereoscope. 200 IJs were then resuspended in 250 μL of LB broth. IJs were

homogenized for 2 minutes using a hand-held motor-driven grinder and Kontes

polypropylene microtube pellet pestle. Serial dilutions of the homogenate were then

plated on LB agar plates with 50 μg/mL kanamycin.

Soil Challenge of IJ colonization. IJ suspensions were transferred to sterile 15 mL

conical tubes and centrifuged at 3000 rpm for 2 minutes to form a pellet. IJs were then

resuspended in 10 mL of M9 buffer (Stiernagle, 2006) and split into two 50 mL culture

flasks, 5 mL each. The concentrations of IJs in each flask were assessed under a

stereoscope and recorded. For each plate, a 1:5 dilution of M9-soaked soil (5 g Miracle-

Gro Potting Mix in 45 mL of M9 incubated at room temperature overnight, approximately

8000 CFU/µL) was filtered through a Whatman paper and added to one of the flasks to

simulate microbial challenge. Weekly, the colonization frequency for each flask was

measured via fluorescence microscopy by counting the number of colonized and

uncolonized worms. 200 IJs were removed from the flask, paralyzed with 2 mM

levamisole in 10 µL of sterile water, and imaged on 5% agar pads.

Microscopy and Image Acquisition. IJ nematodes were concentrated from water-

traps by centrifugation at 1761 g for one minute and immobilized by addition of

levamisole to a final concentration of 5 mM. The nematodes were mounted onto 5%

agarose in water and covered with a cover glass (Cao *et al.*, 2021) . Photomicrographs

and videos were acquired using Zeiss Imager Z2 microscope equipped with Apotome 2

and Axiocam 506 mono using Zen 2 Blue software.

Results

E. coli Nissle cells interact with coelomocytes of _S. hermaphroditum_ IJ.

Using fluorescence microscopy, we observed that mScarlet-I-expressing _E. coli_ Nissle

cells localize to at least four 'spots' on the lateral sides of _S. hermaphroditum_ IJs, with

two spots on the posterior dorsal side and two on the anterior ventral side

(approximately 5-10 μm in diameter) (Fig. 2A and 2B). Z-stack imaging for these four

'spots' showed that each consists of a pair of large ovoid cells (Fig. 2D, Video S1).

Within each cell, the cytosol, containing lysates from fluorescent bacteria, surrounds a

dark center which features the nuclease (Fig. 2D). Based on the morphology and

localization of these cells, we identify them as the coelomocytes, a cell type known to

serve defense and immune functions in invertebrates including some nematode species

(Engelmann _et al._, 2005; Tahseen, 2009; Smith _et al._, 2010). Nematode coelomocytes

are known to have diverse morphologies, sizes, and numbers. The morphology of

coelomocytes ranges from oval-shaped to star-shaped; the sizes of these cells correlate

with the size of the animals (from less than 1 mm to a few meters long). They are

proposed to participate in the up-take and scavenging of proteins circulating in the

pseudocoelomic space. Whether or not the coelomocytes could engulf bacterial cells

depends on the species of nematode (Tahseen, 2009). For instance, _C. elegans_

coelomocytes were found to only have endocytosed toxins, such as nematode-

expressed green fluorescent proteins that circulate in the body cavity, but not pathogens

(Fares and Greenwald, 2001). In contrast, the mammalian parasitic nematode _Ascaris_

suum was observed to phagocytose living bacterial cells in their coelomocytes

(Tahseen, 2009). Neither the morphology nor the function of *Steinernema* coelomocytes

has been previously documented. As expected, we observed that *S. hermaphroditum*

coelomocytes are oval-shaped, like other *Rhabditis* nematodes; rod-shaped bacterial

cells were visible in the coelomocytes (Fig. 2C, 2E, and 2F) consistent with the model

that bacteria are trapped and lysed in the lysosomes. These observations suggest the

phagocytotic and endocytic functions of *Steinernema* coelomocytes directly towards

invading *E. coli* Nissle which causes an active immune response in the host nematode.

E. coli Nissle cells localize and proliferate in the IJ intestine.

As typical of bacterivorous nematodes, the intestine of *Steinernema* serves as the

largest organ facilitating microbial interactions where bacteria can be engulfed,

digested, or colonized (McGhee, 2007) . Prior to IJ development, EcN cells are engulfed

and digested in the intestinal lumen of *S. hermaphroditum* juveniles (Fig. S1). When the

intestinal lumen collapses and closes during IJ development, we observe EcN cells

attaching to the posterior end of the IJ intestine within multiple small 'pockets'. We term

these structures, which were not previously identified, as 'intestinal vacuoles' (Fig. 3A

and 3B). Within the same IJ, EcN cells localize and proliferate in one or more intestinal

vacuoles (Fig. 3B), showing their role as reservoirs for invading microbes. Individual

bacterial cells were also observed to 'escape' from the vacuoles, pass through the anus,

and eventually enter the space in between the internal cuticle and a layer of growing

exterior cuticle (Fig 3B, panel c). This process may be caused by either bacterial motility

(such as swimming), or by the movement of the nematode intestine during IJ

development as it collapses and squeezes out the bacteria accumulated inside the

intestinal lumen. Intestinal colonization was not observed for two other tested laboratory

strains of *E. coli*: strain BW25113 and strain CSH50, suggesting that *E. coli* Nissle

contains genetic factors that allow colonization not found in these other laboratory

strains (Fig. 3C). Intestinal colonization was specific to liver-kidney agar condition

lacking *X. griffiniae* symbiotic bacteria (Fig. 3C), therefore, nutritional profile of liver-

kidney agar and the specific features of the *E. coli* Nissle strain are essential. The

addition of the *X. griffiniae* symbiont was also shown to inhibit intestinal colonization of

EcN (Fig. 3C), possibly due to its ability to antagonize *E. coli* growth by producing

antimicrobial molecules (Dreyer *et al.*, 2018). Overall, EcN is likely a mild and

opportunistic pathogen of *S. hermaphroditum,* and some EcN cells can gain resistance

to the host immune response by invading and persisting in the IJ intestine.

***E. coli* Nissle cells aggregate and migrate in the IJ inter-cuticular space**.

S. hermaphroditum IJs can have up to two cuticle layers: an inner collagenous cuticle

similar to that seen in other developmental stages (Wolkow and Hall, 2011), and

sometimes an outer cuticle that is known to protect the animal from pathogenic

microbes (Timper and Kaya, 1989). We observed aggregates of bacterial cells trapped

in the posterior end of the inter-cuticular space (Fig. 4A and 4B; Fig. S3). Cuticular

association of EcN is either exclusively localized to the posterior side or co-localized at

both the posterior and anterior sides (Fig. S3). No IJ was only associated at the anterior

cuticle without posterior colonization, confirming the directionality of colonization from

the posterior to the anterior; thus, this event must follow intestinal colonization. Like

colonization of the posterior intestinal vacuoles, inter-cuticular colonization was also

specific to growth on liver-kidney agar without the presence of any *X. griffiniae*

symbionts, and was not observed for the laboratory *E. coli* strains BW25113 or CSH50

(Fig. 4C). This is consistent with our hypothesis that these two colonization phenotypes

are correlated (Fig. 3C; Fig. 4C). Along this line of evidence, we also observed the co-

occurrence of EcN cells during intestinal and cuticular colonization (Fig. S3). We

attempted to assess the CFU per cuticle by surface-sterilizing and grinding IJs, and

growing EcN cells on the LB agar plates. However, bleaching process removed the

outer cuticle and killed the EcN cells in the inter-cuticular space, causing nearly 0 CFU

per IJ.

Proteins from *E. coli* cell lysates persist in the IJ inter-cuticular space.

We observed live EcN cells that were temporarily maintained in the IJ cuticular space

for approximately two weeks before they were observed to lyse. Lysis is marked by the

presence of both individual bacterial cells and a smear of bright fluorescence from the

mScarlet-I reporter (Fig. 5A). Eventually, no individual living mScarlet-I-tagged bacteria

are observed in the intercuticular space, suggesting all the observed fluorescence

comes from residual mScarlet-I proteins persisting in the bacterial lysate (Fig. 5B).

Surprisingly, diffused mScarlet-I fluorescence is observed in the inter-cuticular in IJs

grown under all conditions (Fig. 5C), including those where cuticular proliferation of

living bacteria was not observed (Fig. 4C). This observation suggests that in addition to

living bacterial colonization as described above, other mechanisms could cause

bacterial proteins to be expelled and trapped in the inter-cuticular space.

To test the robustness of this lysate's fluorescence to external challenges associated

with the nematode's natural environment, we investigated if fluorescent proteins

contained in the EcN lysates in the nematode cuticle can persist long-term when the IJs

were challenged by external soil microbes (Fig. 6; Fig. S4). Although cuticle growth and

colonization percentages varied among different replicates and experiments within

populations of IJs, mScarlet-I proteins in the lysates remained fluorescent in the cuticle

over eight weeks of IJ aging (Fig. 6; Fig. S4). Overall, soil challenge did not significantly

affect mScarlet-I persistence in the cuticle.

Discussion

In this research, we found EcN cells interacting with four pairs of *S. hermaphroditum* coelomocytes. In model organism *C. elegans* that has the most characterized anatomy and immune system among nematode species, three pairs of coelomocytes are attached to the pseudocoelomic space with three layers of defense preventing bacterial access: the grinding of bacteria by the pharynx, multiple layers of cuticles, and an active immune system defending against the invading microbes (Altun and Hall, 2005). Here, we show that *E. coli* Nissle can access *S. hermaphroditum* coelomocytes, suggesting that at least one of the three barriers fails to defend against EcN. Invertebrate immune cells are known to interact with both mutualistic and pathogenic symbionts. For instance, in the Hawaiian bobtail squid *Euprymna scolopes,* symbiotic *Vibrio fischeri* bacteria adhere to a single blood-cell type, the haemocytes, with less affinity in comparison to pathogen adherence (Nyholm *et al.*, 2009). In *S. hermaphroditum*, the mutualistic symbiont *X. griffiniae* has not been observed to interact with host coelomocytes, suggesting that the role of these immune cells may be specific to pathogens.

EcN cells were also found enclosed in the posterior intestinal pockets of IJs. These pockets, which we term 'vacuoles', are shown in both conventional IJs (those caught from nature and grown only through the natural insect infection process) and axenic IJs (those that are gnotobiotic). This suggests that neither the presence of EcN nor the absence of *Xenorhabdus* symbionts causes the formation of these structures. Based on the size and morphology of the debris present in the intestinal vacuoles, these void

spaces are likely to have formed via cell death of the intestinal cells or organelles (Fig.

3B, panel a; Fig. S1; Video S2 and S3). Our data suggest that the posterior end, similar

to the anterior end of the IJ intestine (Chaston *et al.*, 2013) , features compartments that

are specific to the colonization and proliferation of certain environmental microbes.

In previous studies, environmental microbes were found in *S. scapterisci* in the 'inter-

cuticular space', defined as the space in between the inner cuticle of J2 and J3 larvae

(Bonifassi *et al.*, 1999) . Those rod-shaped bacteria, reported as 'contaminants', were

thought to be wrapped into the J3 cuticle during nematode growth and molting (Fig. 7A).

Note that in this research, EcN cells are localized to the space in between the

collagenous inner cuticle and the outer cuticle. Based on our observations, we propose

that the bacteria enter the inter-cuticular space after being expelled from the intestine

(Fig. 4A and 4B). Within the inter-cuticular space, EcN cells were observed to migrate

from the posterior to the anterior side where they form aggregates, proving that these

bacterial cells are indeed living. What causes the lysis of EcN cells in the cuticle is

unknown, some possibilities include host nematode immune response, or an exhaustion

of nutrients in the inter-cuticular space.

Despite the potential of binary symbiosis systems in environmental engineering, they

remain underexplored. Engineering vertically transmitted symbionts or endosymbionts,

which reside in host cell cytosol and are maternally passed through oocytes (Bright and

Bulgheresi, 2010) , is technically challenging. In contrast, horizontally transmitted

microbiomes—such as the association of *S. hermaphroditum* with *Xenorhabdus*

griffiniae and EcN—are acquired extracellularly from the environment in each

generation. This mode of symbiont acquisition offers key advantages for microbial

bioengineering: symbiotic bacteria (including mutualistic and pathogenic) can be

cultured and genetically modified *ex vivo*, then easily re-associated with the host. This

enables the use of engineered microbes to monitor microenvironments within host

tissues or to assess external environmental conditions within relevant ecosystems.

Conclusion

In this work, we report previously unidentified cells and tissues within *Steinernema*

nematodes where *E. coli* Nissle (EcN) can localize, proliferate, and lyse. We

characterized sequential EcN localization within infective juveniles (IJs), including

immune cells (coelomocytes), posterior intestinal vacuoles, and the inter-cuticular space

between the interior collagenous cuticle and the exterior cuticle (Fig. 7B). Remarkably,

mScarlet-I proteins from EcN lysates persisted within the inter-cuticular space for over

eight weeks under non-sterile soil conditions. This observation challenges the paradigm

that environmental microbes are passively "wrapped" within molting nematode cuticles

(Fig. 7A). Instead, our findings demonstrate that EcN can actively associate with soil-

dwelling *Steinernema* in a tissue-specific manner in a niche that is separate from the

nematode's natural symbiont. During EcN invasion, the immune response of

Steinernema hermaphroditum engages the pseudocoelom, intestinal structures, and

cuticle. Based on our data in this research, we propose EcN to be a mild pathogen of

S. hermaphroditum, which opens a new avenue of establishing *S. hermaphroditum* as a

pathogenesis model. EcN cells and bacterial proteins have the potential for engineering

biocircuits to monitor host-microbe interactions and soil ecosystem.

Figure 1: *Steinernema hermaphroditum* intestine provides specific niches for host-

microbe interactions. (A). *S. hermaphroditum* life cycle and symbiont colonization in

specific tissues at the anterior intestine. (B). *S. hermaphroditum* IJ that has the potential

to interact with both mutualistic and pathogenic microbes. (C). Biphasic lifestyle of

E. coli that oscillates between associating with mammalian gut and persisting in the soil,

which provides an opportunity to interact with soil-dwelling *Steinernema* nematodes. Bb:

basal bulb; AIC: anterior intestinal caecum; PIV: pharyngeal intestinal valve.

Figure 2: *E. coli* Nissle localizes in the *Steinernema hermaphroditum* IJ coelomocytes.

(A). A schematic diagram showing *S. hermaphroditum* IJs has at least four pairs of

coelomocytes. (B) and (C). A representative IJ showing *E. coli* Nissle localized and

lysed within coelomocytes. (D). Each ovoid coelomocyte structure contains two cells.

(E) and (F). Rod-shaped bacterial cells are observed inside of the coelomocyte cytosol.

Scale bars in panel D, E, and F equals to 20 µm. (Also see Video S1).

Figure 3: *E. coli* Nissle localizes in the *S. hermaphroditum* intestinal lumen and

vacuoles. (A). A schematic diagram showing *E. coli* Nissle invading nematode intestine

and proliferating in the intestinal vacuoles. (B). A representative IJ showing intestinal

colonization of bacteria: an empty intestinal vacuole containing cell or organelle debris-

looking substance (a); bacterial colonization in the IJ intestinal vacuoles (b); bacterial

cells escape from the posterior intestinal opening (c). (C). Quantification of intestinal

colonization frequency of *E. coli* strains: Nissle (Nis), *E. coli* CSH50 (CSH), BW25113

(BW) under nutritional conditions of either liver-kidney (LK) or lipid agar (LA).

Treatments were either with (+) or without (-) symbiont *X. griffiniae.* Numbers in the

parenthesis indicate the number of animals screened in each treatment.

Figure 4: Inter-cuticular colonization of *E. coli* Nissle. (A). A schematic diagram showing

E. coli Nissle colonizes in the inter-cuticular space in the IJs. (B). A representative IJ

showing inter-cuticular proliferation of bacteria: the head region (a) and the tail region

(b). Arrow indicates the intestinal vacuole colonization observed in the same IJ with

cuticular colonization. (C). Quantification of cuticular proliferation frequency of *E. coli*

strains: Nissle (Nis), *E. coli* CSH50 (CSH), BW25113 (BW) under nutritional conditions

of either liver-kidney (LK) or lipid agar (LA). Treatments were either with (+) or without (-

) symbiont *X. griffiniae.* Numbers in the parenthesis indicates the number of animals

screened in each treatment.

Figure 5: *E. coli* Nissle cells lyse in the IJ inter-cuticular space. (A). A representative IJ

(2 weeks post trapping) showing transitional stage in which bacterial cells and mScarlet-

I contained in cell lysate co-localize in the inter-cuticular space. (B). Diffused

fluorescence from mScarlet-I expressing *E. coli* cells is found in symbiont-colonized IJs.

(C). Quantification of cuticular diffusion of mScarlet-I expressing *E. coli* strains: Nissle

(Nis), *E. coli* CSH50 (CSH), BW25113 (BW) under nutritional conditions of either liver-

kidney (LK) or lipid agar (LA). Treatments were either with (+) or without (-) symbiont *X.*

griffiniae. Numbers in the parenthesis indicates the number of animals screened in each

treatment.

Figure 6: Proportion of developed outer cuticles and cuticularly colonized IJs as a

fraction of total IJs with and without soil challenge. Developed cuticles (noted as

'developed') include those with fully developed, less developed, or broken outer

cuticles. The presence of *E. coli* Nissle produced mScarlet-I proteins in IJ cuticles

(noted as 'colonized') was assessed by fluorescence microscopy. (A). IJ cuticle

development and colonization of mScarlet-I protein in the inter-cuticular space over

three weeks. (I), (II), and (III) represents three biological replicates. (B). Cuticle

development and cuticular mScarlet-I colonization frequency over ten weeks. (IV), (V),

and (VI) represent three biological replicates. (C). A comparison of IJ cuticle

development and cuticular colonization by mScarlet-I in week 8 and week 9 with and

without challenge by non-sterile soil.

Figure 7: Hypothetical models of *E. coli* Nissle interaction in *S. hermaphroditum*. (A). a

hypothetical model based on a previous publication of *S. scapterisci* cuticular microbes

(Bonifassi *et al.*, 1999) : environmental microbes attaching to the inner cuticle during IJ

formation. (B). Current hypothetical model based on data from this research: *E. coli*

Nissle possibly causes mild infection in *S. hermaphroditum*.

References

Altun, Z.F. and Hall, D.H. (2005) WormAtlas Hermaphrodite Handbook - Coelomocyte System. *WormAtlas*.

Baquero, F., Coque, T.M., Galán, J.C., and Martinez, J.L. (2021) The Origin of Niches and Species in the Bacterial World. *Front Microbiol* **12**: 657986.

Bonifassi, E., Saux, M.F.-L., Boemare, N., Lanois, A., Laumond, C., and Smart, G. (1999) Gnotobiological Study of Infective Juveniles and Symbionts of Steinernema scapterisci: A Model to Clarify the Concept of the Natural Occurrence of Monoxenic Associations in Entomopathogenic Nematodes. *J Invertebr Pathol* **74**: 164–172.

Bracho, O.R., Manchery, C., Haskell, E.C., Blanar, C.A., and Smith, R.P. (2016) Circumvention of Learning Increases Intoxication Efficacy of Nematicidal Engineered Bacteria. *Acs Synth Biol* **5**: 241–249.

Bright, M. and Bulgheresi, S. (2010) A complex journey: transmission of microbial symbionts. *Nat Rev Microbiol* **8**: 218–230.

Cao, M., Schwartz, H.T., Tan, C.-H., and Sternberg, P.W. (2021) The entomopathogenic nematode Steinernema hermaphroditum is a self-fertilizing hermaphrodite and a genetically tractable system for the study of parasitic and mutualistic symbiosis. *Genetics* **220**:.

Chaston, J.M., Murfin, K.E., Heath-Heckman, E.A., and Goodrich-Blair, H. (2013) Previously unrecognized stages of species-specific colonization in the mutualism between Xenorhabdus bacteria and Steinernema nematodes: Species-specific host colonization by Xenorhabdus. *Cell Microbiol* **15**: 1545–1559.

Dale, C. and Moran, N.A. (2006) Molecular Interactions between Bacterial Symbionts and Their Hosts. *Cell* **126**: 453–465.

Dreyer, J., Malan, A.P., and Dicks, L.M.T. (2018) Bacteria of the Genus Xenorhabdus, a Novel Source of Bioactive Compounds. *Front Microbiol* **9**: 3177.

Elsas, J.D. van, Semenov, A.V., Costa, R., and Trevors, J.T. (2010) Survival of Escherichia coli in the environment: fundamental and public health aspects. *ISME J* **5**: 173–183.

Engelmann, P., Cooper, E.L., and Németh, P. (2005) Anticipating innate immunity without a Toll. *Mol Immunol* **42**: 931–942.

Fares, H. and Greenwald, I. (2001) Genetic Analysis of Endocytosis in Caenorhabditis elegans: Coelomocyte Uptake Defective Mutants. *Genetics* **159**: 133–145.

Flores-Lara, Y., Renneckar, D., Forst, S., Goodrich-Blair, H., and Stock, P. (2007) Influence of
nematode age and culture conditions on morphological and physiological parameters in the
bacterial vesicle of Steinernema carpocapsae (Nematoda: Steinernematidae). *J Invertebr*
Pathol **95**: 110–118.

Gao, B. and Sun, Q. (2021) Programming gene expression in multicellular organisms for
physiology modulation through engineered bacteria. *Nat Commun* **12**: 2689.

Grenier, F., Matteau, D., Baby, V., and Rodrigue, S. (2014) Complete Genome Sequence of
Escherichia coli BW25113. *Genome Announc* **2**: e01038-14.

Halleran, A.D., Swaminathan, A., and Murray, R.M. (2018) Single Day Construction of
Multigene Circuits with 3G Assembly. *Acs Synth Biol* **7**: 1477–1480.

Hwang, I.Y., Koh, E., Wong, A., March, J.C., Bentley, W.E., Lee, Y.S., and Chang, M.W.
(2017) Engineered probiotic Escherichia coli can eliminate and prevent Pseudomonas
aeruginosa gut infection in animal models. *Nat Commun* **8**: 15028.

Iverson, S.V., Haddock, T.L., Beal, J., and Densmore, D.M. (2016) CIDAR MoClo: Improved
MoClo Assembly Standard and New E. coli Part Library Enable Rapid Combinatorial Design
for Synthetic and Traditional Biology. *ACS Synth Biol* **5**: 99–103.

Jones, E.M., Marken, J.P., and Silver, P.A. (2023) Synthetic microbiology in sustainability
applications. *Nat Rev Microbiol* **22**: 345–359.

Kim, J. and Moon, Y. (2019) Worm-Based Alternate Assessment of Probiotic Intervention
against Gut Barrier Infection. *Nutrients* **11**: 2146.

Maire, J., Parisot, N., Ferrarini, M.G., Vallier, A., Gillet, B., Hughes, S., et al. (2020) Spatial and
morphological reorganization of endosymbiosis during metamorphosis accommodates adult
metabolic requirements in a weevil. *Proc Natl Acad Sci* **117**: 19347–19358.

McGhee, J. (2007) The C. elegans intestine. *WormBook* 1–36.

Murfin, K.E., Chaston, J., and Goodrich-Blair, H. (2012) Visualizing bacteria in nematodes
using fluorescent microscopy. *J Vis Exp Jove*.

Murfin, K.E., Ginete, D.R., Bashey, F., and Goodrich☐Blair, H. (2019) Symbiont☐mediated
competition: Xenorhabdus bovienii confer an advantage to their nematode host Steinernema
affine by killing competitor Steinernema feltiae. *Environ Microbiol* **21**: 3229–3243.

Nyholm, S.V. and McFall-Ngai, M.J. (2021) A lasting symbiosis: how the Hawaiian bobtail
squid finds and keeps its bioluminescent bacterial partner. *Nat Rev Microbiol* **19**: 666–679.

Nyholm, S.V. and McFall-Ngai, M.J. (1998) Sampling the light-organ microenvironment of
Euprymna scolopes: description of a population of host cells in association with the bacterial
symbiont Vibrio fischeri. *Biological Bulletin* **195**: 89–97.

Nyholm, S.V., Stewart, J.J., Ruby, E.G., and McFall☐Ngai, M.J. (2009) Recognition between
symbiotic Vibrio fischeri and the haemocytes of Euprymna scolopes. *Environ Microbiol* **11**:
483–493.

Ogier, J.-C., Pagès, S., Frayssinet, M., and Gaudriault, S. (2020) Entomopathogenic nematode-
associated microbiota: from monoxenic paradigm to pathobiome. *Microbiome* **8**: 25.

Redweik, G. and Xue, D. (2025) Secreted autotransporter toxin produced by probiotic
Escherichia coli Nissle 1917 enhances neurodegeneration in Caenorhabditis elegans.
microPublication Biol **2025**: 10.17912/micropub.biology.001366.

Rutter, J.W., Ozdemir, T., Quintaneiro, L.M., Thomas, G., Cabreiro, F., and Barnes, C.P. (2019)
Detecting changes in the Caenorhabditis elegans intestinal environment using an engineered
bacterial biosensor. *Biorxiv* 717215.

Smith, L.C., Ghosh, J., Buckley, K.M., Clow, L.A., Dheilly, N.M., Haug, T., et al. (2010)
Echinoderm immunity. *Adv Exp Med Biol* **708**: 260–301.

Sokurenko, E.V., Courtney, H.S., Abraham, S.N., Klemm, P., and Hasty, D.L. (1992) Functional
heterogeneity of type 1 fimbriae of Escherichia coli. *Infect Immun* **60**: 4709–4719.

Sonnenborn, U. (2016) Escherichia coli strain Nissle 1917—from bench to bedside and back:
history of a special Escherichia coli strain with probiotic properties. *FEMS Microbiol Lett*
363: fnw212.

Stiernagle, T. (2006) Maintenance of C. elegans. *Wormbook* 1–11.

Stilwell, M.D., Cao, M., Goodrich-Blair, H., and Weibel, D.B. (2018) Studying the Symbiotic
Bacterium Xenorhabdus nematophila in Individual, Living Steinernema carpocapsae
Nematodes Using Microfluidic Systems. *Msphere* **3**: e00530-17.

St-Pierre, F., Cui, L., Priest, D.G., Endy, D., Dodd, I.B., and Shearwin, K.E. (2013) One-Step
Cloning and Chromosomal Integration of DNA. *Acs Synth Biol* **2**: 537–541.

Tahseen, Q. (2009) Coelomocytes: Biology and Possible Immune Functions in Invertebrates
with Special Remarks on Nematodes. *Int J Zoology* **2009**: 1–13.

Thomas, N.M.St., Myers, T.G., Alani, O.S., Goodrich-Blair, H., and Heppert, J.K. (2024) Green
and red fluorescent strains of Xenorhabdus griffiniae HGB2511, the bacterial symbiont of the
nematode Steinernema hermaphroditum (India). *microPublication Biol* **2024**:
10.17912/micropub.biology.001064.

Timper, P. and Kaya, H.K. (1989) Role of the second-stage cuticle of entomogenous nematodes
in preventing infection by nematophagous fungi. *J Invertebr Pathol* **54**: 314–321.

Tivey, T.R., Coleman, T.J., and Weis, V.M. (2022) Spatial and Temporal Patterns of Symbiont
Colonization and Loss During Bleaching in the Model Sea Anemone Aiptasia. *Front Mar Sci*
9: 808696.

Del Valle, I.D., Fulk, E.M., Kalvapalle, P., Silberg, J.J., Masiello, C.A., and Stadler, L.B. (2021)
Translating New Synthetic Biology Advances for Biosensing Into the Earth and
Environmental Sciences. *Front Microbiol* **11**: 618373.

Vivas, E.I. and Goodrich-Blair, H. (2001) Xenorhabdus nematophilus as a model for host-
bacterium interactions: rpoS is necessary for mutualism with nematodes. *J Bacteriol* **183**:
4687–4693.

Wolkow, C. and Hall, D.H. (2011) WormAtlas Dauer Handbook - The Dauer Cuticle.
WormAtlas.

604
605
606

607 **Supplemental Materials**

608 **Figure S1**: cell or organelle debris in the posterior intestinal vacuoles in a

609 representative infective juvenile. Black arrows: gut granules; white arrows: cell or

610 organelle debris in the intestinal vacuoles. Scale bar = 20 µm.

613
614

Figure S2: A representative *S. hermaphroditum* juvenile fed on *E. coli* Nissle. mScarlet-

I expressing EcN cells are engulfed and digested in the intestinal lumen of J2 stage of

S. hermaphroditum. 'P' denotes posterior end of the nematode; 'int. lumen' denotes

'intestinal lumen'; 'b.b' denotes 'basal bulb'; 'AIC' denotes 'anterior intestinal caecum';

'PIV' denotes 'pharyngeal intestinal valve'.

Figure S3: A profile of twenty-nine IJs with EcN colonization in the intestinal vacuoles

and the inter-cuticualr space. 'P' denotes the posterior side of the IJ.

Figure S4: outer cuticle growth and bacterial protein (mScarlet-I) colonization in

Steinernema hermaphroditum IJs. (A): Fractions of IJs with developed outer cuticle over

9 week's time course of experiment. (B): Fraction of developed IJs with outer cuticle that

has cuticular colonization of mScarlet from EcN lysate over the 9 weeks time course of

experiment.

Video S1: bacterial lysis within the *S. hermaphroditum* coelomocytes. *E. coli* Nissle

expressing mScarlet is localized in the coelomocytes.

Video S2: intestinal cell or organelle debris in a representative intestinal vacuole.

Video S3: a representative IJ with two empty intestinal vacuoles housing nematode
organelle debris, and two vacuoles housing EcN cells expressing mScarlet-I.
640