Title: Chromosomal rearrangements and segmental deletions contribute to gene loss in

squamates

Buddhabhushan Girish Salve, Sonal H, Nagarjun Vijay

buddhabhushan21@iiserb.ac.in, sonal24@iiserb.ac.in, nagarjun@iiserb.ac.in

Computational Evolutionary Genomics Lab, Department of Biological Sciences, IISER

Bhopal, Bhauri, Madhya Pradesh, India

Author for Correspondence: Nagarjun Vijay, nagarjun@iiserb.ac.in

Abstract

Background

Genomic rearrangements, including segmental deletions, duplications, translocations, and

inversions of DNA segments, can contribute to gene losses, thereby reshaping genome

architecture and potentially resulting in functional consequences. In squamates, karyotypic

evolution mainly involves chromosome number reduction through fusions and

microchromosome to macrochromosome translocations, although fissions have also

contributed to diversification in several lineages. Despite these dynamics, the evolutionary

processes and underlying genetic mechanisms driving chromosomal rearrangements and

associated gene losses in squamates remain poorly understood.

Results

In this study, we analysed chromosome/scaffold-level assemblies of 261 squamates,

corroborated by short-read, long-read, and transcriptomic data. We found multiple lines of

evidence for the putative loss of 53 genes in the squamate lineage. Synteny and phylogenetic

analysis revealed that, among the 53 unretrieved orthologs, 14 are lost in squamates with no retained paralog, 15 show ortholog loss with retained paralogs, and 24 remain as unretrieved orthologs. Furthermore, we find that many of the genes lost from squamates are organised in syntenic clusters and are involved in essential immune functions—raising important questions about the role of paralogs in compensating for the function of lost genes, strengthening the "less is more" hypothesis in the squamate lineage.

Conclusions

Together, our comparative genomic analyses highlight that the loss of crucial genes in squamate lineages has occurred primarily through inter- and intrachromosomal rearrangements, including segmental deletions. These findings offer insights into the evolutionary loss of genes involved in macrophage differentiation and inform the development of novel pharmaceutical approaches for modulating immune responses.

Keywords

segmental deletions, macrophage polarisation, gene loss, karyotype evolution, squamates

Background

Genomic rearrangements, including deletions, inversions, translocations, and duplications, play a major role in shaping genome architecture and can potentially contribute to gene loss [1–3]. Such structural changes often disrupt gene integrity by breaking coding regions, altering regulatory landscapes, or relocating genes into heterochromatic or transcriptionally silent regions [4, 5]. For example, large-scale deletions can physically remove entire genes, whereas rearrangements such as translocations or inversions may separate essential gene components or place them under incompatible regulatory control, rendering them non-functional [3, 6, 7]. Moreover, rearrangements mediated by repetitive elements can create unstable genomic regions prone to further structural variation, thereby accelerating gene

erosion [8–12]. These processes are particularly evident in rapidly evolving lineages, such as

birds and mammals, where gene loss contributes to lineage-specific adaptation and

diversification [13–16].

The recent dramatic increase in the number of high-quality genome assemblies of diverse

organisms has facilitated studies on lineage-specific gene losses [17–20]. The NCBI RefSeq

annotation of orthologues for multiple vertebrate species and the availability of short- and

long-read sequencing data allow the confirmation of gene losses [21, 22]. However, most

studies linking gene loss and genomic rearrangements in vertebrate evolution have focused

on birds [3, 23], mammals [11] and fishes [24], and the association of such chromosomal

rearrangement and gene loss have not been systematically performed in squamates.

Squamate reptiles are a diverse monophyletic group of vertebrates classified into lizard and

snake families, which diverged from tuatara approximately 250 million years ago (MYA) [25,

26]. With more than 11,000 described species, they are more species-rich than birds [27].

Squamates have emerged as excellent systems for studying genomic rearrangement [28],

karyotype [29], genome size evolution [30], and unique biology such as tissue regeneration

and limb degeneration [31, 32]. Yet, the dearth of genome assemblies has been a bottleneck

for large-scale comparative genomic studies [33]. The recent availability of high-quality

genomes across the squamate tree of life has opened new avenues to look at lineage-specific

genome dynamics, such as sex chromosome evolution [34], changes in the repeat landscape

[35], and GC content heterogeneity [36]. Notably, the genome size of squamates is

heterogenous, with a trend toward genome size reduction [37]. Similar reductions in birds

and bats have been associated with the loss of gene families and entire syntenic blocks,

suggesting that the contraction of the gene repertoire may accompany genome size reduction

[37, 38]. Although gene and whole-genome duplications drive evolutionary innovation,

emerging evidence now reinforces the "less-is-more" hypothesis, highlighting gene loss as an

equally potent force in shaping genomic and phenotypic diversity [39–41]. The growing

availability of chromosome-scale assemblies from initiatives such as the Vertebrate Genomes

Project (VGP) is now enabling the systematic identification of such gene loss events,

providing unprecedented opportunities to explore how gene reduction contributes to lineage-

specific adaptation and evolutionary novelty [42–44]. In this context, investigating patterns of

gene loss across the squamate radiation can therefore yield critical insights into how genomic

reduction and restructuring contribute to phenotypic innovation, evolutionary diversification,

and an alternative to phyletic gradualism [45].

To address this gap, we focused on gene loss in squamates and the role of genomic

rearrangements. Our preliminary genome-wide screening of vertebrates revealed that 53 key

genes with conserved functions are not annotated in NCBI RefSeq in squamates. Among

these genes, previous studies have reported the loss of *PLAAT1*, *SLC24A1,* and *IL34* in

squamates and associated them with adaptation to low-light environments [46], changes in

the visual system [47], and myeloid cell-type evolution [48], respectively. These studies

suggest the evolutionary loss of these genes based on a search of the genome assemblies and

failure to recover any remnants. Here, we leverage high-quality genomes of squamates

coupled with a comprehensive search of genomic short- and long-read and transcriptomic

data, assembly validation, syntenic and phylogenetic analysis to rule out inadequacies in the

genome assembly posing as gene loss [42, 49, 50]. We began our search by looking for genes

annotated in the five outgroup species (i.e., Western clawed frog (*Xenopus tropicalis*),

American alligator (*Alligator mississippiensis*), chicken (*Gallus gallus*), green sea turtle

(*Chelonia mydas*), and human (*Homo sapiens*)) and not annotated in all annotated squamate

genomes. Furthermore, we looked for direct (search of genomes, raw read datasets, statistical

and phylogenetic signal of gene loss) and indirect evidence (co-evolution of the receptor)

supporting the loss of these genes. We found that gene loss in squamates occurred in

conserved syntenic clusters, similar to the pattern observed in birds [38]. The putative gene

loss we identified needs to be evaluated with better-quality genome assemblies, and its

functional impact should be assessed through comparative functional assays.

Results

First, as a pilot analysis to identify unretrieved orthologs specific to the entire squamate

lineage, we focused on genes consistently identified as not annotated in NCBI RefSeq across

all 30 annotated squamate genomes, while being present in outgroup species (Additional file

1: Table S1-S2). To ensure robust inference, we used high-quality genome assemblies from

multiple vertebrate species, including 261 squamates. Of these, 77 assemblies meet the

quality standards recommended by global initiatives such as the Vertebrate Genome Project

and the Earth BioGenome Project, with contig N50 values exceeding 1 Mb and scaffold N50

values exceeding 10 Mb (Additional file 1: Table S3). The 53 unretrieved orthologs were

systematically classified into distinct categories based on multiple lines of genomic and

syntenic evidence. These included: (i) assembly verification using long-read data through the

Klumpy tool, (ii) local synteny assessment, (iii) evidence for segmental deletion or

chromosomal rearrangement, (iv) ortholog sequence properties such as GC-content

distribution and sequence identity with human orthologs, (v) local assembly using HybPiper

followed by phylogenetic analysis, (vi) pairwise genome alignment, (vii) paralog annotation,

(viii) transcriptional activity at the syntenic locus and cross-species expression mapping in

central bearded dragon and chicken, (ix) BLASTn and tBLASTx searches in the near T2T

phased genome assembly of central bearded dragon, and (x) BLASTn search against 261

squamate genomes. Based on the integration of syntenic and phylogenetic evidence,

unretrieved orthologs were further subdivided into three categories: (i) ortholog loss with no

retained paralog, supported by evidence for segmental deletion and/or syntenic and

phylogenetic analysis and GC-content < 65% and sequence identity > 55% with no sequence

paralog found, (ii) putative ortholog loss with retained paralog, syntenic and phylogenetic

analysis found paralog along with loss of orthologs, and (iii) unclear cases, where high

sequence divergence for highly diverged ortholog or translocated ortholog with high

sequence divergence or germline-restricted genes or hard to sequence genomic regions

containing unretrieved orthologs present in dark matter or non-B-form DNA structures or

proximity to telomeric and centromeric regions, prevented confident classification

(Additional file 1: Table S1 and Additional file 2: Gene-wise evidences).

Unretrieved orthologs occur in conserved syntenic blocks and chromosomal

rearrangements

We examined the genomic locations of the unretrieved orthologs in the chicken genome and

found that they are distributed across both macro- and microchromosomes (**Fig. 1**). This

contrasts with previous studies on gene loss in birds, where unretrieved orthologs were

predominantly located on microchromosomes [38, 42]. We found clusters of unretrieved

orthologs on chromosomes 1 (*LCP1*, *RUBCNL*, *SIAH3*, *LRCH1*, *ARL11*, *DLEU7*, *STOML3*,

INTS6 and *FREM2*) and 4 (*TBC1D14*, *MAN2B2*, *GPR78*, *GRK4*, *TNIP2*, *KDM3A*) of

chicken. We also noted 7 clusters of two or more genes within ~ 2 Mb of each other in the

chicken genome. The cross-species comparison of syntenic loci revealed that these regions

are prone to both intra- and inter-chromosomal rearrangements in squamate species such as

the central bearded dragon (*Pogona vitticeps*), brown anole (*Anolis sagrei*), and terrestrial

garter snake (*Thamnophis elegans*) (**Fig. 1**; Additional file 1: Table S4). In several cases,

these loci are also mapped to telomeric regions. In contrast, comparable rearrangements were

not observed in other reptiles, including the American alligator (*Alligator mississippiensis*)

and the green sea turtle (*Chelonia mydas*), suggesting squamate-specific genomic

rearrangements.

Integrated multi-evidence stratified framework for putative gene loss validation

Based on the syntenic analysis along with local genome assembly validation, the 53 unretrieved orthologs could be classified as putatively lost due to segmental deletions (n=36) and intra/inter chromosomal rearrangements (n=17) (**Fig. 2A**). In the case of segmental deletions, the evidence for the loss of the gene consists of squamate-specific phylogenetic signal of reduction in intergenic distance, genome blast coverage, and phylogenetic assessment of sequences assembled by HybPiper or retrieved by blast search. While in the case of chromosomal rearrangements, the evidence relied largely on syntenic and phylogenetic analysis. An evidence score for the loss of each gene based on these integrated multiple layers of evidence was used to rank the unretrieved orthologs.

The BLASTn search of 261 squamate genomes revealed that most unretrieved orthologs are absent from squamates. Of the 53 genes, 21 have query coverage (search on squamate genomes using human orthologues as a query) of ≤ 25%, 23 genes between 25 - 74%, and 9 genes have coverage of ≥ 75% (*SUV39H2*, *FREM2*, *DNM3*, *RBBP7*, *MOB1A*, *SSTR4*, *WNT2*, *LCP1* and *CAV3*; **Fig. 2B**; Additional file 1: Table S5). Our bioinformatic analysis (using BioMart human paralog annotation) revealed two or more paralogs for each gene having query coverage >75% (**Fig. 3**; Additional file 1: Table S6). Furthermore, we assembled gene sequences using raw whole-genome sequencing datasets. The phylogenetic analysis of the retrieved sequences revealed that they cluster with closely related paralogues rather than the expected orthologues (Additional file 1: Table S7 and Additional file 2: Gene-wise evidences).

It has been previously reported that unretrieved orthologs or "missing" or "hidden" genes in genome assemblies are often associated with high GC content and high sequence divergence [42, 49, 51, 52]. However, in our analysis, the mean GC content of the unretrieved ortholog

(calculated across vertebrate orthologues) ranged from 39.56% to 64.72% (**Fig. 2B**). Notably,

nineteen genes—*CAV3* (55.21%), *NOTCH2* (55.30%), *CYYR1* (55.43%), *DIP2A* (55.47%),

UNKL (55.80%), *RIPPLY3* (56.06%), *ADRA1B* (56.48%), *IL34* (56.76%), *WNT2* (56.90%),

TNIP2 (57.83%), *SLC9A5* (58.12%), *MATK* (58.23%), *SIAH3* (58.52%), *DLEU7* (58.67%),

ARL11 (58.92%), *HSD17B1* (61.35%), *YBX3* (61.65%), *SH2D2A* (64.47%), and *SSTR4*

(64.72%)—had GC content exceeding 55% (Additional file 1: Table S1).

Immune genes often evolve under strong selective pressure due to the ongoing evolutionary

arms race between the host immune system and invading pathogens. This rapid evolution

results in low amino acid conservation among orthologous proteins across species, limiting

the effectiveness of sequence similarity-based tools such as BLASTn for detecting these

genes [52–55]. Therefore, some of these genes may not be truly lost but may be present as

highly diverged orthologs. We observed that 24 genes including *ARL11, GPR78, HHLA2,*

HPGDS, HSD17B1, IFTAP, IL13RA2, IL26, MATK, MOB1A, OLAH, PLAAT1, RBBP7,

SH2D2A, SLC24A1, TMEM273, UTS2B, ZNF438, DLEU7, DNM3, UNKL, MAN2B2,

RUBCNL, and TNIP2 share less than 65% sequence identity with their human orthologs (**Fig.**

2B; Additional file 1: Table S1), suggesting that moderate to high sequence divergence may

contribute to their loss or lack of annotation in squamate genomes, therefore remained

classified as unretrieved orthologs. Comparative genome alignment between human and

tuatara reveals that several genes were intact in the ancestral lineage with conserved gene

synteny, highlighting their loss as specific to squamates (Additional File 1: Table S8).

Based on syntenic analysis, 36 genes were found to have a conserved gene order similar to

that of the outgroup species, while eight genes showed evidence of intrachromosomal

rearrangements and nine genes indicated interchromosomal rearrangements (**Fig. 2A**;

Additional file 1: Table S1). Most of the 53 unretrieved orthologs lacked evidence of

transcription, with some genes having spurious mapping (Additional File 3: Cross species

RNA-seq mapping). We found evidence for segmental deletion in the case of 19 out of 36 genes, namely, *ADAP2*, *ELOVL3*, *GRK4*, *HHLA2*, *HSD17B1*, *IL34*, *IL5RA*, *KDM3A*, *LAPTM5*, *MOB1A*, *RBBP7*, *RIPPLY3*, *SLC24A1*, *SLC9A5, SSTR4, STAP1, SUV39H2, TBC1D14, TMEM273* and *WNT2* (Additional file 1: Table S1).

Paralogs were not found for 14 of the 53 unretrieved orthologs. Among these 14 genes, (i) *IL34*, *IL5RA*, and *RIPPLY3* exhibit high sequence divergence while retaining conserved synteny, with additional evidence of segmental deletion, (ii) A similar pattern (except for high sequence divergence) was observed for *GRK4*, *LAPTM5*, *SLC9A5*, *SSTR4*, *STAP1*, and *WNT2*, (iii) Despite weak evidence of segmental deletion, *CYYR1* (with frame-disrupting changes in *Anolis sagrei*, partial exon remnants in a genome assembly validated syntenic locus in *Pogona vitticeps* and lack of transcription in *Pogona vitticeps* and *Rhineura floridana*) and *GPR82* (with no blast hits even to paralogs despite low sequence divergence among orthologs) have conserved synteny (Additional file 4: Figure S1), (iv) The phylogenetic analysis suggests the lack of paralogs among the chromosomal rearrangement genes, *DIP2A*, *NOTCH2*, and *SH2D1A*. Collectively, these 14 genes were classified as cases of ortholog loss with no retained paralog (Additional file 1: Table S1, and Additional file 2: Gene-wise evidences).

Multi-gene families are often prone to inactivation or deletion, consistent with the birth-and-death model of gene family evolution [13, 24, 56, 57]. We found that 39 of the 53 unretrieved orthologs belong to multi-gene families and have at least one paralogue (**Fig. 2** and **Fig. 3**; Additional file 1: Table S1, S6 and S7). Among these 39 genes, (i) the evidence for conserved synteny and segmental deletion with limited sequence divergence was found for *ADAP2*, *CAV3*, *EVOVL3*, *INTS6*, *KDM3A*, *SUV39H2*, and *TBC1D14*, (ii) chromosomal rearrangement was identified in the case of *ADRA1B*, *KREMEN1*, *FREM2*, *LCP1*, *LRCH1*, *SIAH3*, *STOML3*, and *YBX3* with limited sequence divergence. Collectively, these 15 genes were

classified as cases of ortholog loss with retained paralog (Additional file 1: Table S1, and

Additional file 2: Gene-wise evidences).

Of the remaining 24 of the 39 genes with paralogs, *ARL11*, *GPR78*, *HHLA2*, *HSD17B1*,

IFTAP, *IL13RA2*, *IL26, OLAH*, *PLAAT1*, *SH2D2A*, *SLC24A1*, *TMEM273*, *UTS2B*, *DLEU7*,

UNKL, *FREM2*, *LCP1*, *LRCH1*, *MAN2B2*, *RUBCNL*, *SIAH3*, *STOML3*, and *TNIP2* had high

sequence divergence and remain as unretrieved orthologs. In summary, synteny and

phylogenetic analyses suggest that, among the 53 unretrieved orthologs, 14 genes are lost in

squamates with no retained paralog, 15 show ortholog loss with retained paralogs, and 24

remain as unretrieved orthologs (Additional file 1: Table S1, and Additional file 2: Gene-wise

evidences).

Furthermore, pair-wise genome alignments (at these 53 unretrieved orthologs loci) between

chicken (*Gallus gallus*) and 34 other vertebrate species revealed that the squamate (~280

MYA) genomes consistently having lower aligned exonic regions compared to tuatara

(*Sphenodon punctatus*; ~280 MYA), Testudines (~261 MYA), and Crocodilians (~245 MYA),

despite these groups having comparable evolutionary divergence from chicken (**Fig. 4A**;

Additional file 1: Table S9 and Additional file 4: Figure S2; pair-wise divergence time

obtained from Timetree website [58]). The flanking genes to focal genes show significantly

higher levels of alignments at exonic regions (**Fig. 4B;** Additional file 1: Table S9, and

Additional file 2: Gene-wise evidences). The ENSEMBL available and chains files (pair-wise

genome-wide LASTZ alignment) created by this study do not show any marked difference

(Wilcoxon, p-value = 0.16, **Fig. 4C**). We also found that the aligned regions at exonic

positions of focal genes are significantly lower in squamates compared to Birds,

Coelacanthiformes, Testudines, Crocodylia, and Sphenodontia (**Fig. 4D**, Additional file 1:

Table S9). The occurrence of unaligned regions in non-squamate species for some genes

could be due to assembly gaps and/or high sequence divergence. The aligned region could

also be from the paralogous regions.

Comparative genomic evidence for *IL34* loss via segmental deletion in squamates

As reported previously [48], our comparative genomic analysis confirmed the absence of the

IL34 gene in squamates despite its widespread conservation across other vertebrates. Synteny

analysis revealed that *COG4* and *SF3B3* typically flank *IL34* on the left and *MTSS2* and

VAC14 on the right (**Fig. 5A**; Additional file 1: Table S4). Notable exceptions are observed in

species such as the zebrafish (*Danio rerio*), green sea turtle (*Chelonia mydas*), American

alligator (*Alligator mississippiensis*), and tiger rattlesnake (*Crotalus tigris*), which have

IONP2, *PARD6A*, *BBS2*, and *CNOT1,* respectively, on the right flank of the *IL34* locus. The

lack of annotation for the *IL34* gene in squamates could potentially result from assembly

artefacts. To rule out this possibility, we examined the syntenic locus of *IL34* in squamate

species such as the common wall lizard (*Podarcis muralis*) and the Indian cobra (*Naja naja*)

using PacBio and Oxford Nanopore long-read data. In both cases, overlapping reads span the

IL34 gene locus, indicating that the assemblies are correct at this locus (**Fig. 6**). BLASTn

analysis of squamate genome assemblies and raw read datasets failed to recover any sequence

corresponding to the *IL34* gene, further supporting its absence in this lineage and ruling out

the possibility of its translocation to another genomic region (Additional file 1: Table S5 and

S7). Next, we compared intergenic distances by examining squamate species for evidence of

segmental deletions compared to non-squamate species (see Methods). Wilcoxon rank-sum

tests (alternative = "less") consistently yielded adjusted *p*-values < 0.001 for all combinations

of gene distance ratios involving *SF3B3–MTSS2*, indicating significant contraction in this

region in squamates (**Fig. 5B**; Additional file 1: Table S1; Additional file 4: Fig. S3). In

contrast, other gene combinations did not significantly differ across all comparisons.

Additionally, evidence for segmental deletion was supported by phylogenetic logistic

regression analyses, with significant results from both logistic_MPLE (slope = 0.2; *p*-value =

0.0022) and logistic_IG10 (slope = 0.2; *p*-value = 0.0018) (**Fig. 5C**; Additional file 1: Table

S1). Additionally, we found intrachromosomal rearrangement of the *IL34* locus in some of

the Viperidae snake species (Additional file 4: Fig. S4).

Our bioinformatic analysis, integrating synteny, long-read mapping-based genome assembly

verification, BLASTn searches, transcriptome searches, intergenic distance comparisons, and

phylogenetic regression, robustly confirms the loss of the *IL34* gene in squamates.

Co-evolution of *CSF1R* with *IL34* gene loss in the squamate lineage

Our previous results from multifaceted approaches concurrently infer loss of the *IL34* gene

from squamates. Next, we investigated the evolutionary consequences of *IL34* gene loss on

its receptor —*CSF1R,* in squamates. Previous studies have suggested that *IL34* remains

relatively unchanged, with detectable co-variation with *CSF1R*, whereas no such correlation

is observed for *CSF1* [Garceau et al. [59], reviewed in [60]]. However, this hypothesis

remains untested in lineages where *IL34* is lost. The availability of high-quality genome

assemblies of squamates and *IL34* gene loss opens the opportunity to address this interesting

evolutionary question─how two ligands evolve with a single receptor? Using selection

analysis tools, we detected signatures of relaxed selection acting on *CSF1R* in several

squamate species. We found squamate species such as Western terrestrial garter snake

(*Thamnophis elegans*; *p*-values = 0.0003, k-value = 0.63), komodo dragon (*Varanus*

komodoensis; *p*-value = 0.0053, k-value = 0.62), and central bearded dragon (*Pogona*

vitticeps; *p*-value = 0.0047, k-value = 0.59) show signatures of relaxed selection by RELAX

(with *p*-values <0.05 and k-value <0) and/or codeml (**Fig. 7**; Additional file 1: Table S10).

Notably, signatures of relaxed selection were not pervasive across the screened vertebrate

lineages, suggesting lineage-specific evolutionary changes. Several sites within the

extracellular domain of *CSF1R* were identified as positively selected by both MEME and

FEL analyses, with squamate species exhibiting a relatively higher number of positively

selected sites (Additional file 1: Table S10). Together, the results of comparative sequence

analyses revealed lineage-specific modifications in *CSF1R* in squamates, potentially

reflecting compensatory evolution following *IL34* gene loss. These findings provide indirect

but compelling evidence for *IL34* gene loss in squamates and support the hypothesis of

receptor adaptation following the loss of one of its ligands.

Discussion

Our study identifies large-scale gene losses across squamates, with 53 genes consistently

unretrievable in all 261 squamate genomes screened. These genes are dispersed across macro-

and microchromosomes in the chicken genome and are typically lost individually, while some

are unretrievable in clusters. Based on synteny conservation, the unretrieved orthologs are

classified as putatively lost due to segmental deletions (n=36) and intra/inter chromosomal

rearrangements (n=17). Among the 53 unretrieved orthologs, 21 had < 25% blastn query

coverage, 23 had intermediate coverage of 25–74%, and 9 had ≥75% coverage in the

genomes of squamates. The mean GC content of the unretrieved orthologs ranges from 39.56

% to 64.72 %, is within a GC-neutral range and suggests that GC bias is unlikely to cause

them to be unretrievable. Additionally, many of these genes exhibit less than 70% sequence

identity to their human orthologues, indicating substantial evolutionary sequence divergence.

Despite extensive efforts, we could not recover these genes from available genomic or

transcriptomic resources, including searches in high-quality genome assemblies, analyses of

raw genomic sequencing reads, and examination of RNA-seq data across diverse tissues in

the central bearded dragon (*Pogona vitticeps*). The synteny and phylogenetic analyses of the

53 unretrieved orthologs revealed that 14 genes are lost in squamates with no retained

paralog, 15 show ortholog loss with retained paralogs, and 24 remain unretrieved orthologs.

Based on the evidence score, *IL34* is the strongest empirical case of evolutionary gene loss.

Our comparative genomic analyses confirmed the complete loss of *IL34* in squamates via

segmental deletion and intrachromosomal rearrangements at the *IL34* locus in some

Viperidae snakes. We also identified squamate-specific modifications in *CSF1R*, suggesting

possible compensatory changes following *IL34* loss. Similarly, *CAV3* and *KDM3A* have the

highest evidence score among the 14 orthologs lost with retained paralogs. While the cross-

species mapping found noisy RNA-seq support for both these genes, the phylogenetic

analysis and synteny confirm the loss of the ortholog with the paralogs retained in line with

the "less is more" hypothesis.

Evolutionary loss of conserved genes in squamates: patterns and limitations

In squamates, we discovered that segmental deletions and intra- and inter-chromosomal

genomic rearrangements cause the putative loss of these 53 unretrieved orthologs. In contrast

with previously documented gene losses in birds, which are primarily found on

microchromosomes and are found to be due to assembly errors [42], these genes are

dispersed throughout different macro- and microchromosomes rather than clustered on

microchromosomes in the chicken genome [38, 61]. We observed that some gene losses

occurred in clusters, with two or more adjacent genes unretrievable from the same genomic

region. For *IL34*, both segmental deletions and intra-chromosomal rearrangements were

observed, suggesting that these loci lie in structurally unstable regions. The loss of *IL34* in

squamates, along with the signatures of loss on the receptor and presence of a structural

paralog [48, 59, 62], makes the case for functional compensation by distant structural

homologs [63, 64]. Sequence paralogs from ancient whole-genome or segmental duplications

are more widely studied than structural homologs. The example of ortholog loss with retained

paralog, such as *CAV3*, *KDM3A*, *ADAP2*, *SUV39H2*, and *TBC1D14,* could be compensated

for by their paralogs, such as *CAV3*-like, *KDM3B*, *ADAP1L*, *SUV39H1* and *TBC1D12*.

Furthermore, the pattern of *TNIP2* presence in the vicinity of the telomeric region of chicken

chromosome 4, compared to interchromosomal rearrangement in squamates (Additional file

4: Fig. S5), suggests that gene loss may be facilitated by the prevalence of the gene in highly

variable and rapidly evolving regions such as telomeric and sub-telomeric regions [65, 66].

Genes such as *TNIP2* are retained as unretrieved orthologs due to the challenges of

distinguishing evolutionary loss of genes from methodological limitations.

Previous studies [38, 52, 53, 61] have shown that some genes initially reported as "missing"

in birds were later recovered using transcriptomic data [50], with their absence attributed to

high GC content and limitations in earlier genome assemblies [49]. We incorporated GC

content analysis to avoid similar errors, searched short- and long-read datasets, and examined

RNA-seq data to minimise the risk of false gene loss inference in squamates. The GC content

of unretrieved orthologs in squamates is less than that reported for the "missing" GC-rich

genes in birds, which were later found to be "dark matter" missing in prior genome

assemblies [42, 49, 67–69]. While lineage-specific GC content shifts in squamates cannot be

ruled out, the presence of most unretrieved orthologs in tuatara weakens the likelihood of a

widespread increase in GC content in squamates. Furthermore, we assembled gene sequences

by performing BLASTn searches using raw whole-genome sequencing datasets. Phylogenetic

analysis of the retrieved sequences revealed that they clustered with closely related

paralogues rather than the expected orthologues. We verified genome assemblies for

conserved synteny and ruled out assembly artefacts as the cause of unretrieved orthologs,

suggesting that structural rearrangements, rather than assembly artefacts, may play a role in

gene loss. Many of the unretrieved orthologs, including several interleukins, show less than

70% identity to their human orthologues and are primarily linked to immune functions (for

gene-specific detailed references, see Additional file 1: Table S11). A plausible explanation is

the involvement of these genes in host-pathogen arms races, where rapid evolution driven by

changing selective pressures can lead to high sequence divergence and positive selection [70–

73]. Consequently, these genes may still exist in squamates but in highly diverged forms,

making them difficult to detect using standard sequence homology-based approaches.

Furthermore, we could not accurately estimate deletion sizes, as the genes are absent in all

the examined squamates, and the nearest outgroup with intact genes, tuatara, diverged ~250

MYA [26]. This deep divergence limits the precise delineation of deletion boundaries, unlike

in previous studies, where segmental deletions could be more clearly defined [6, 54].

The discovery of squamate-specific gene losses offers insights into their evolutionary

importance, supporting the idea that gene loss can be adaptive and extending the "less-is-

more" hypothesis to squamates vis-à-vis providing an alternative to phyletic gradualism [39,

65, 74–77]. Many unretrieved orthologs have multiple paralogs; in some cases, their apparent

absence may result from assembly limitations. However, if the gene loss is true, its function

is likely compensated by paralogous genes. Additionally, most unretrieved orthologs form

conserved syntenic clusters in non-squamate vertebrates, suggesting their loss in squamates

likely involved block deletions, potentially contributing to genome size reduction [37]. Future

genome assemblies should aim to assess the presence or absence of these genes to understand

the mechanisms and evolutionary consequences of their loss.

Functional implications of unretrieved orthologs

Earlier studies reported the absence of *PLAAT1*, *SLC24A1*, and *IL34* in squamates and

associated their loss with adaptations to low-light environments, modifications in the visual

system, and the evolution of myeloid cell types, respectively [46–48]. Our methodology has

extended the search genome-wide and identified these genes as unretrieved orthologs,

reinforcing the reliability of our gene loss detection approach and highlighting the need for

detailed functional investigations of unretrieved orthologs. A literature survey revealed that

the 53 unretrieved orthologs from squamates are linked to diverse biological functions,

including immunity and inflammation, cell growth, apoptosis, cancer, development,

cardiovascular and metabolic processes, musculoskeletal formation, sensory function, and

core cellular mechanisms like autophagy, protein degradation, and transcriptional regulation

(for gene-specific detailed references, see Additional file 1: Table S11). The involvement of

these genes in fundamental physiological and immune functions raises critical questions

about how these pathways operate in their absence, offering a valuable opportunity to explore

compensatory mechanisms and lineage-specific adaptations. The loss of these genes may play

a pivotal role in evolutionary genetics by influencing genome size and complexity, affecting

the rate of evolution and adaptation, offering insights into gene and pathway essentiality, and

contributing to phenotypic diversity in squamates [13]. Furthermore, we found that the genes

KDM3A, *SUV39H2*, and *RBBP7*, which are unretrieved in squamates, are associated with the

Gene Ontology (GO) term "chromatin remodeling", and are involved in DNA replication,

recombination, and repair. The absence of these genes suggests a potential impact on

chromatin architecture, with possible consequences for genome accessibility and stability in

squamates [3]. The genes *FREM2*, *WNT2* and *NOTCH2* involved in eye development are

unretrieved and may be associated with changes in the visual system in squamates [31, 46,

47].

The present study reports the loss of four genes— *IL34*, *STAP1*, *LAPTM5* and *TNIP2*-

involved in macrophage activation and polarisation. *IL34* expression is observed in tissues

such as the skin, brain, kidneys, and testes and is modulated in diseases like Alzheimer's,

cancer and HBV viral infection [78–80]. The macrophages derived from *IL34* exhibit more

anti-viral activity than those from *CSF1* [81–83]. Despite having a normal phenotype, $IL34^{-/-}$

mice exhibit impaired responses to skin antigens and viral infections, highlighting its

physiological importance [84, 85]. These findings suggest that squamates may be

functionally compromised owing to the loss of *IL34*, potentially rendering them more

susceptible to pathogenic infections. In mice, *STAP1* deficiency impairs TCR-mediated T cell

activation [86], *TNIP2* acts as a regulator of NF-κB signalling [87, 88], and *LAPTM5* inhibits

HIV-1 progeny infectivity by transporting viral Env to the lysosome for degradation in

macrophages [89]. However, the role of these genes in reptiles remains largely unexplored

and needs functional validation. The limited availability of tools for assessing immune

function in free-living reptiles presents a challenge in functionally testing this scenario. These

findings offer new insights into the evolutionary reduction in the gene repertoire regulating

macrophage differentiation, which may help develop novel pharmaceutical strategies for the

immune modulation of monocytes and macrophages.

Conclusions

This study systematically identified and confirmed that 53 genes are putatively lost in the

squamate lineage but retained in outgroup species, suggesting lineage-specific gene loss

events primarily associated with large-scale segmental deletions and chromosomal

rearrangements. Among the 53 unretrieved orthologs, 14 genes are lost in squamates with no

retained paralog, 15 show ortholog loss with retained paralogs, and 24 remain unretrieved

orthologs. Notably, four unretrieved orthologs are involved in macrophage activation and

polarisation—*IL34*, *STAP1*, *LAPTM5* and *TNIP2*, while others play key roles in immunity,

development, metabolism, cellular processes, and neuronal and musculoskeletal functions.

The unretrieved orthologs are GC-neutral and are members of multi-gene families with

several paralogues. Our study provides direct and indirect evidence of potential gene losses in

the squamate lineage. Despite comprehensive searches across genome assemblies, raw reads,

and transcriptomes, none could be recovered, supporting their possible loss in squamates.

Further studies are needed to generate high-quality squamate genomes to identify the

causative forces behind these gene losses and their final confirmation.

Methods

Identification of unretrieved orthologs from squamates

We retrieved the list of all 20,595 protein-coding genes annotated in the human genome from NCBI (Additional file 1: Table S1-S2). The orthologues for each human gene across 679 annotated vertebrate genomes were obtained using the datasets command-line utility of NCBI (parameters: --ortholog all) [90]. First, as a pilot analysis to identify unretrieved orthologs specific to squamates, we focused on genes consistently identified as not annotated in NCBI RefSeq across all 30 annotated squamate genomes. As a filtering criterion to ensure that these genes were not universally absent or pseudogenised or annotation artefact, we retained only those genes that were classified as intact (annotated without the "Low quality protein" tag of NCBI) in multiple distantly related species, namely Western clawed frog (*Xenopus tropicalis*), American alligator (*Alligator mississippiensis*), chicken (*Gallus gallus*), green sea turtle (*Chelonia mydas*), and human (*Homo sapiens*) allowing to confidently infer squamate-specific unretrieved orthologs by comparing across a broad phylogenetic tree of vertebrates (Additional file 1: Table S1-S2).

Analysis of evolutionary rearrangement and chromosomal mapping of unretrieved orthologs

To investigate the chromosomal distribution and rearranged genomic regions of unretrieved orthologs across squamates, we mapped them to chicken karyotypes. We used GENESPACE v1.3.1 [91] to understand patterns of synteny and orthology across chicken (*Gallus gallus*), American alligator (*Alligator mississippiensis*), green sea turtle (*Chelonia mydas*), central bearded dragon *(Pogona vitticeps)*, brown anole (*Anolis sagrei*), and terrestrial garter snake (*Thamnophis elegans*) genomes. A comparative framework enabled the visualisation of conserved syntenic blocks and rearranged genomic regions linked to these 53 unretrieved

orthologs in squamates. We used BEDTools nuc to calculate GC content across 100 bp

sliding windows along the length of the chromosome. To identify sequences associated with

non-B DNA-forming motifs, we used the GFA v2 program [92]. The resulting outputs were

used to plot the GC content and the presence of non-B DNA motifs across chicken

chromosomes. Additionally, the unretrieved orthologs loci were compared between the

genome assemblies of chicken (*Gallus gallus*) and green anole (*Anolis carolinensis*) using the

NCBI Comparative Genome Viewer [93].

BLASTn search of 53 unretrieved orthologs in squamate genomes and raw read

datasets

To increase the power of screening for unretrieved orthologs in squamate genomes, we

performed BLASTn [94] searches (parameters: -evalue 0.05 -outfmt '17 SQ', 1, 6, and 7)

across 261 squamate genome assemblies, including both chromosome-level and scaffold-

level assemblies, using human (*Homo sapiens*) and chicken (*Gallus gallus*) gene sequences

as queries (Additional file 1: Table S5). The BLASTn output in format '17 SQ' was used to

generate IGV reports [95], which were then screened to assess the presence of gene

sequences in the squamate assemblies. Furthermore, we screened publicly available short-

and long-read DNA and RNA sequencing data from PacBio (HiFi and Revio), Oxford

Nanopore, and Illumina sequencing technologies—from squamate species (Additional file 1:

Table S12). We employed BLASTn (parameters: -evalue 0.05 -max_target_seqs 10000) with

query sequences of unretrieved orthologs of the five outgroup species. To distinguish between

orthologs and paralogs, the sequence obtained from HybPiper [96] was analysed using IQ-

TREE v2.3.6 [97], along with coding DNA sequences from 39 species that produced

BLASTn hits to the extracted sequence (parameters: -evalue 0.001)(Additional file: Table

S12). We used the 'ape' package to identify the closest phylogenetic branches to the

HybPiper recovered gene sequence [98]. Similar phylogenetic analysis was performed using

newly available near-complete T2T genomes of the Australian central bearded dragon

(*Pogona vitticeps*) [99, 100]. To identify orthologous genes across 28 species (19 squamates

and 9 outgroups), protein sequences of all annotated genes in FASTA format were used as

input to OrthoFinder 2 [101]. The analysis involved an all-vs-all sequence similarity search

using DIAMOND, followed by clustering of genes into orthogroups, each representing a set

of genes derived from a single ancestral gene in the last common ancestor. From this dataset,

we focused on 53 genes of particular interest and retrieved their corresponding orthogroup

IDs. The gene IDs within these orthogroups were then extracted. To confirm whether these

sequences represented true orthologs, or paralogs, or gene duplicates, we conducted synteny

analyses and phylogenetic inference using IQ-TREE 2 (Additional file 1: Table S13) [97].

Furthermore, to assess the paralog prevalence, we used Ensembl BioMart to obtain annotated

paralogs, GC content, sequence identity, and transcript count data for human and chicken

orthologs of squamate-specific unretrieved orthologs.

To explore potential transcriptional evidence for unretrieved orthologs, we mapped RNA-seq

data from the central bearded dragon (*Pogona vitticeps*) of the testis, muscle, lung, liver,

kidney, eye, heart, and brain tissues to the chicken genome using minimap2 v2.29-r1283

[102] (parameters: -ax splice -uf -k14) and FLAIR v2.2.0 [103] (Additional file 1: Table

S12). We then screened for the presence of transcripts corresponding to unretrieved orthologs

and visualised the read coverage and splice junctions using IGV-report, allowing us to assess

transcriptional signals for these genes in the central bearded dragon (*Pogona vitticeps*)

despite their apparent genomic absence (i.e., lack of annotation and failure to be picked up by

blastn search).

Identification of 1-to-1 orthologs, assembly verification, GC content, paralog annotation

of unretrieved orthologs

We confirmed gene synteny of unretrieved orthologs and identified 1-to-1 orthologs across

eight vertebrate species, namely Western clawed frog (*Xenopus tropicalis*), American

alligator (*Alligator mississippiensis*), chicken (*Gallus gallus*), green sea turtle (*Chelonia*

mydas), green anole (*Anolis carolinensis*), Common wall lizard (*Podarcis muralis*), Central

bearded dragon (*Pogona vitticeps*), Eastern brown snake (*Pseudonaja textilis*), tiger

rattlesnake (*Crotalus tigris*), tuatara (*Sphenodon punctatus*), platypus (*Ornithorhynchus*

anatinus), house mouse (*Mus musculus*) and human (*Homo sapiens*) using NCBI genome

viewer [22] and ENSEMBL [104]. To assess the integrity of the unretrieved orthologs loci in

the chromosome/scaffold-level assembly of Indian cobra (*Naja naja*), tiger rattlesnake

(*Crotalus tigris*), Central bearded dragon (*Pogona vitticeps*) and common wall lizard

(*Podarcis muralis*). We used the *scan_alignment* and *alignment_plot* options in the Klumpy

v1.0.10 tool [105], the required BAM file was generated by mapping long-read sequencing

data from squamates species (Additional file 1: Table S12) with Minimap2 v2.17-r941 [102]

(parameters: -ax map-pb or -ax map-hifi or -ax map-ont). The resulting alignment plots were

examined for the presence of continuous and overlapping reads spanning the unretrieved

orthologs loci.

The lack of annotations and/or sequence for the unretrieved ortholog from the genome

assembly of a species could be due to its high GC content and/or high sequence divergence

[42, 49, 106]. Therefore, we calculated GC content across the coding sequence of vertebrate

orthologs using seqkit (parameters: fx2tab --name –gc) and sequence identity with respect to

the human ortholog using EMBOSS-needle [107].

Squamate lineage-specific evidence of segmental deletion

a. Dataset preparation

We developed an integrated analysis pipeline encompassing orthologue retrieval, synteny mapping, intergenic distance measurement, and gap detection to investigate segmental deletions underlying unretrieved orthologs. Firstly, four neighbouring syntenic genes (two upstream and two downstream of the focal gene) were identified with synteny analysis, and their orthologs across vertebrate genomes were retrieved from NCBI datasets (parameters: --include product-report --ortholog all). The longest isoform was selected for each ortholog based on protein length and species, where orthologs of all four syntenic genes were annotated. We calculated pairwise intergenic distances using BEDTools closest. Gene pairs on different chromosomes were flagged as potential Evolutionary Breakpoint Regions (EBRs). A comprehensive gene distance matrix with complete orthologous information was assembled for all species. Next, to detect gaps in the locus, genomic intervals spanning the outermost flanking genes were merged with BEDTools merge -d 100000000, and the corresponding sequences were retrieved using NCBI efetch. These regions were scanned for assembly gaps using the Klumpy v1.0.10 find_gaps subprogram, and the total number of gaps was recorded per species. Each species' taxonomy classification (class and order) was obtained using NCBI Taxonomy resources (esearch and efetch). Finally, all data—including intergenic distances, gap counts, and taxonomic info—were consolidated into a unified summary table. A subset of species with complete data was used for downstream visualisation (heatmap) and phylogenetic logistic regression analysis. Based on the presence of intact synteny and absence of gaps, species were classified as having either an intact locus (non-squamates) or a potentially deleted locus (squamates).

b. Statistical analysis of gene distance and phylogenetic signal

i. Ratio-Based Comparative Analysis

To evaluate whether changes in intergenic distances between the focal gene (genes which are unretrievable in squamates) and its flanking genes are associated with segmental

deletion, we first excluded species exhibiting large-scale genomic disruptions such as gaps

and EBRs. We calculated six pairwise ratios representing the remaining species' upstream,

downstream, and across-gene intergenic distances. We conducted Wilcoxon rank-sum tests

for each ratio with the alternative hypothesis set to "less" to compare distributions between

species with intact versus deleted gene loci. The resulting p-values were adjusted for

multiple testing using the FDR method, and significant comparisons were visualised as a

heatmap using the pheatmap package in R [108].

ii. Phylogenetic Logistic Regression

To quantify the phylogenetic association between relative gene distance and gene loss, we

performed phylogenetic logistic regression using the "phylolm" R package [109]. Squamates

(0) and non-squamates (1) were used as the binary variable. The independent variable was the

relative intergenic distance for the focal gene, computed as:

$$\overset{G_{L2}}{\longleftarrow} \qquad \overset{G_{L1}}{\longrightarrow} \qquad \overset{focal\ gene}{\Longrightarrow} \qquad \overset{G_{R1}}{\longleftarrow} \qquad \overset{G_{R2}}{\longrightarrow}$$

576   Where

G_{L1}: Gene on the left flank of a lost gene on 1^{st} position

G_{L2}: Gene on the left flank of a lost gene on 2^{nd} position

G_{R1}: Gene on the right flank of a lost gene on 1^{st} position

G_{R2}: Gene on the right flank of a lost gene on 2^{nd} position

$focal\ gene$: unretrieved orthologs in squamates

The direction of the arrow indicates gene orientation

$$Relative\ distance\ in\ percentage\ = \left(\frac{d_{focal}}{d_{upstream}\ +\ d_{focal}\ +\ d_{downstream}} \right) \times 100$$

Where

$$d_{focal} = \text{distance between } G_{L1} \text{ and } G_{R1}$$

$$d_{downstream} = \text{distance between } G_{L1} \text{ and } G_{L2}$$

$$d_{upstream} = \text{distance between } G_{R1} \text{ and } G_{R2}$$

Two methods were used for model fitting: logistic_MPLE and logistic_IG10 (Additional file

1: Table S1).

An UpSet plot was used to visualise multiple lines of evidence supporting gene loss,

including synteny-based indicators of segmental deletion or chromosomal rearrangement,

genome blast coverage, validation of genome assembly integrity at the gene locus using

klumpy, phylogenetic analyses for ortholog and paralog detection using whole-genome

sequencing data, hybpiper & IQ-TREE, evidence of segmental deletion inferred from

intergenic distances, paralog annotations from Ensembl, and orthologous sequence-based

metrics such as GC content and sequence identity.

Signatures of receptor-ligand co-evolution as indirect evidence of *IL34* loss in squamates

Structurally similar ligands can functionally compensate for the loss of each other [110].

Therefore, to explore the evolutionary impact of *IL34* loss in squamates, we investigated the

IL34–CSF1–CSF1R signalling axis, focusing on the functional interplay between these

ligands (*IL34*, *CSF1*) and their shared receptor *CSF1R*. *CSF1R* is a type III receptor tyrosine

kinase critical for developing, surviving, and functioning myeloid lineage cells [111].

Previous studies have shown evolutionary co-variation between *IL34* and *CSF1R*, but not

between *CSF1* and *CSF1R* [Garceau et al. [59] and reviewed in [60]]. Based on this, we

hypothesised that *IL34* loss may influence the evolution and function of *CSF1R* in squamates.

We leveraged ENSEMBL annotation for *CSF1R* across vertebrates and retrieved the coding

DNA sequence (CDS) of the *CSF1R* with a conserved exon phase (Additional file: Table

S10). Multiple sequence alignments (MSAs) of intact *CSF1R* from Squamates and vertebrate

clades were generated using PRANK v.170427 in the Guidance2 suite [112, 113]. Nucleotide

sequence alignments were trimmed using trimAl v1.4.1 (parameters: *-resoverlap 0.05 -*

seqoverlap 90), removing sequences with >10% gaps and positions with gaps in >5% of

sequences [114] and used to assess the strength of selection. Further, the alignments and

phylogenetic tree obtained from the Timetree website (https://timetree.org/) were used to

investigate the strength of selection [58]. We relied on the dN/dS-based method of Hyphy

v2.5.48 and PAML v4.9f [115, 116]. Our selection suite consists of RELAX, BUSTED,

aBSERL, MEME, and FEL frameworks of branch-site and site models of Hyphy, along with

the null (M0) and branch (branch-free and branch-neutral) models of codeml [117–123]. To

investigate whether the gene evolves under relaxed selection, we specified one branch as the

foreground and all other branches as the background at a time in the RELAX model of the

HyPhy program [68] and codeML of the PAML program [69]. A significant result of k > 1

indicates intensified selection along test branches, while k < 1 suggests relaxed selection,

with significance determined at $p < 0.05$. We also implemented aBSREL and BUSTED,

which are branch and gene-wide tests, respectively, to test for signatures of selection on

branches of the genes. Site-specific models, such as FEL and MEME, were used to identify

sites evolving under positive and negative selection, and ggplot2 was used for visualisation in

the R program [124].

Quantifying cross-species alignment conservation

To validate the absence of unretrieved orthologs in squamates, we utilised pair-wise genome

alignment to quantify the alignment coverage of diverse vertebrates and compared squamates

vs non-squamate orders. For this, we analysed pairwise whole-genome alignments between

chicken (GRCg7b) and 27 other vertebrate species, including squamates, using precomputed

LASTZ alignments from the Ensembl Comparative database

(https://ftp.ensembl.org/pub/current_maf/ensembl-compara/pairwise_alignments/) [104].

Additionally, we created seven new pairwise whole genome alignments using newly available

squamates genome assemblies with BUSCO > 96% (Additional file 1: Table S12). The

multiple alignment format (MAF) files for each species were downloaded and processed

using MafFilter v1.3.1 [125] with the OutputCoordinates filter enabled (output_src_size=yes)

to extract conserved syntenic blocks between chicken and the focal species. Next, we used

BEDTools intersect with the -wao option to compute overlap between each exon of 53 genes

and the aligned regions for each focal species. Finally, to assess alignment for each

unretrieved ortholog, we normalised the aligned region with CDS length. The resulting

matrix (gene × species) of normalised aligned values was visualised as a heatmap using the

pheatmaps v1.0.13 R package (Additional file 1: Table S9). Pairwise Wilcoxon rank-sum

tests (alternative = "greater") were performed to compare the aligned regions between

squamates and other orders. Multiple testing correction was applied using the "BH" method,

and adjusted p-values were used to assess significance (Additional file 1: Table S9).

Comparisons of alignments across syntenic genes within squamates were visualised using

alluvial plots and heatmaps followed by statistical evaluation using Wilcoxon rank-sum tests

(alternative = "less") to compare the left and right neighbouring genes of each focal

unretrieved ortholog. Genome alignments of squamate species were assessed using both the

assemblies available in Ensembl and those generated in this study.

Data availability

643 The data and relevant code for this study are available on GitHub:

644 https://github.com/CEGLAB-Buddhabhushan/Missing_genes_in_squamata.git

Abbreviations

EBR: Evolutionary Breakpoint Region

HBV: Hepatitis B Virus

MYA: Million Years Ago

TCR: T Cell Receptor

PacBio: Pacific Biosciences

ONT: Oxford Nanopore Technologies

FDR: False Discovery Rate

BH: Benjamini–Hochberg correction

NCBI: National Center for Biotechnology Information

BLASTn: Basic Local Alignment Search Tool (nucleotide)

MAF: Multiple Alignment Format

IGV: Integrative Genomics Viewer

RNA-seq: RNA sequencing

Mb: Megabase

Kb: Kilobase

References

1. Burssed B, Zamariolli M, Bellucco FT, Melaragno MI. Mechanisms of structural
chromosomal rearrangement formation. Mol Cytogenet. 2022;15.

2. Claude SJ, Park S, Park SJ. Gene loss, genome rearrangement, and accelerated substitution
rates in plastid genome of Hypericum ascyron (Hypericaceae). BMC Plant Biol. 2022;22.

3. Huang Z, De O. Furo I, Liu J, Peona V, Gomes AJB, Cen W, et al. Recurrent chromosome
reshuffling and the evolution of neo-sex chromosomes in parrots. Nat Commun. 2022;13.

4. Harewood L, Fraser P. The impact of chromosomal rearrangements on regulation of gene
expression. Hum Mol Genet. 2014;23:R76–82.

5. Stewart NB, Rogers RL. Chromosomal rearrangements as a source of new gene formation
in Drosophila yakuba. PLOS Genet. 2019;15:e1008314.

6. Salve BG, Kurian AM, Vijay N. Concurrent loss of ciliary genes WDR93 and CFAP46 in
phylogenetically distant birds. R Soc Open Sci. 2023;10.

7. Ungrová L, Geryk J, Kohn M, Kučerová D, Krchlíková V, Hron T, et al. Avian interferon
regulatory factor (IRF) family reunion: IRF3 and IRF9 found. BMC Biol. 2025;23:1–19.

8. Stankiewicz P, Shaw CJ, Withers M, Inoue K, Lupski JR. Serial segmental duplications
during primate evolution result in complex human genome architecture. Genome Res.
2004;14:2209–20.

9. Hinsch H, Hannenhalli S. Recurring genomic breaks in independent lineages support
genomic fragility. BMC Evol Biol. 2006;6:1–12.

10. Emanuel BS, Shaikh TH. Segmental duplications: an "expanding" role in genomic
instability and disease. Nat Rev Genet. 2001;2:791–800.

11. Shinde SS, Sharma S, Teekas L, Sharma A, Vijay N. Recurrent erosion of
COA1/MITRAC15 exemplifies conditional gene dispensability in oxidative phosphorylation.
Sci Reports 2021 111. 2021;11:1–16.

12. Meredith RW, Gatesy J, Cheng J, Springer MS. Pseudogenization of the tooth gene
enamelysin (MMP20) in the common ancestor of extant baleen whales. Proc R Soc B Biol
Sci. 2011;278:993–1002.

13. Albalat R, Cañestro C. Evolution by gene loss. Nat Rev Genet 2016 177. 2016;17:379–
91.

14. Blumer M, Brown T, Freitas MB, Destro AL, Oliveira JA, Morales AE, et al. Gene losses
in the common vampire bat illuminate molecular adaptations to blood feeding. Sci Adv.
2022;8:6494.

15. Sharma V, Hecker N, Roscito JG, Foerster L, Langer BE, Hiller M. A genomics approach
reveals insights into the importance of gene losses for mammalian adaptations. Nat Commun
2018 91. 2018;9:1–9.

16. Guijarro-Clarke C, Holland PWH, Paps J. Widespread patterns of gene loss in the
evolution of the animal kingdom. Nat Ecol Evol 2020 44. 2020;4:519–23.

17. Koepfli KP, Paten B, O'brien SJ, Antunes A, Belov K, Bustamante C, et al. The Genome
10K Project: a way forward. Annu Rev Anim Biosci. 2015;3:57–111.

18. Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, et al. Phylogenomic analyses data
of the avian phylogenomics project. Gigascience. 2015;4.

19. Jebb D, Huang Z, Pippel M, Hughes GM, Lavrichenko K, Devanna P, et al. Six
reference-quality genomes reveal evolution of bat adaptations. Nature. 2020;583:578–84.

20. Stiller J, Feng S, Chowdhury AA, Rivas-González I, Duchêne DA, Fang Q, et al.
Complexity of avian evolution revealed by family-level genomes. Nat 2024 6298013.
2024;629:851–60.

21. Wheeler DL, Barrett T, Benson DA, Bryant SH, Canese K, Chetvernin V, et al. Database
resources of the National Center for Biotechnology Information. Nucleic Acids Res. 2007;35
SUPPL. 1.

22. Rangwala SH, Kuznetsov A, Ananiev V, Asztalos A, Borodin E, Evgeniev V, et al.
Accessing NCBI data using the NCBI Sequence Viewer and Genome Data Viewer (GDV).
Genome Res. 2021;31:159–69.

23. Friocourt F, Lafont AG, Kress C, Pain B, Manceau M, Dufour S, et al. Recurrent DCC

gene losses during bird evolution. Sci Reports 2017 71. 2017;7:1–11.

24. Kato A, Pipil S, Ota C, Kusakabe M, Watanabe T, Nagashima A, et al. Convergent gene
losses and pseudogenizations in multiple lineages of stomachless fishes. Commun Biol.
2024;7:408.

25. Simões TR, Caldwell MW, Tałanda M, Bernardi M, Palci A, Vernygora O, et al. The
origin of squamates revealed by a Middle Triassic lizard from the Italian Alps. Nature.
2018;557:706–9.

26. Gemmell NJ, Rutherford K, Prost S, Tollis M, Winter D, Macey JR, et al. The tuatara
genome reveals ancient features of amniote evolution. Nat 2020 5847821. 2020;584:403–9.

27. Simões TR, Pyron RA. THE SQUAMATE TREE OF LIFE. Bull Museum Comp Zool.
2021;163:47–95.

28. Mezzasalma M, Macirella R, Odierna G, Brunelli E. Karyotype Diversification and
Chromosome Rearrangements in Squamate Reptiles. Genes 2024, Vol 15, Page 371.
2024;15:371.

29. Waters PD, Patel HR, Ruiz-Herrera A, Alvarez-Gonzalez L, Lister NC, Simakov O, et al.
Microchromosomes are building blocks of bird, reptile, and mammal chromosomes. Proc
Natl Acad Sci U S A. 2021;118:e2112494118.

30. Pinto BJ, Gamble T, Smith CH, Wilson MA. A lizard is never late: Squamate genomics
as a recent catalyst for understanding sex chromosome and microchromosome evolution. J
Hered. 2023;114:445–58.

31. Emerling CA. Genomic regression of claw keratin, taste receptor and light-associated
genes provides insights into biology and evolutionary origins of snakes. Mol Phylogenet
Evol. 2017;115:40–9.

32. Roscito JG, Sameith K, Kirilenko BM, Hecker N, Winkler S, Dahl A, et al. Convergent
and lineage-specific genomic differences in limb regulatory elements in limbless reptile
lineages. Cell Rep. 2022;38:110280.

33. Gable SM, Mendez JM, Bushroe NA, Wilson A, Byars MI, Tollis M. The State of
Squamate Genomics: Past, Present, and Future of Genome Research in the Most Speciose
Terrestrial Vertebrate Order. Genes (Basel). 2023;14:1387.

34. Modi WS, Crews D. Sex chromosomes and sex determination in reptiles: Commentary.
Curr Opin Genet Dev. 2005;15:660–5.

35. Gable SM, Bushroe NA, Mendez JM, Wilson A, Pinto BJ, Gamble T, et al. Differential
Conservation and Loss of Chicken Repeat 1 (CR1) Retrotransposons in Squamates Reveal
Lineage-Specific Genome Dynamics Across Reptiles. Genome Biol Evol. 2024;16.

36. Alföldi J, Di Palma F, Grabherr M, Williams C, Kong L, Mauceli E, et al. The genome of
the green anole lizard and a comparative analysis with birds and mammals. Nature.
2011;477:587–91.

37. Kapusta A, Suh A, Feschotte C. Dynamics of genome size evolution in birds and
mammals. Proc Natl Acad Sci U S A. 2017;114:E1460–9.

38. Lovell P V, Wirthlin M, Wilhelm L, Minx P, Lazar NH, Carbone L, et al. Conserved
syntenic clusters of protein coding genes are missing in birds. Genome Biol. 2014;15:565.

39. Olson M V. When Less Is More: Gene Loss as an Engine of Evolutionary Change. Am J
Hum Genet. 1999;64:18–23.

40. Sánchez-Serna G, Badia-Ramentol J, Bujosa P, Ferrández-Roldán A, Torres-Águila NP,
Fabregà-Torrus M, et al. Less, but More: New Insights From Appendicularians on Chordate
Fgf Evolution and the Divergence of Tunicate Lifestyles. Mol Biol Evol. 2025;42.

41. Hoffmann FG, Opazo JC, Storz JF. Whole-Genome Duplications Spurred the Functional
Diversification of the Globin Gene Superfamily in Vertebrates. Mol Biol Evol. 2012;29:303–
12.

42. Kim J, Lee C, Ko BJ, Yoo DA, Won S, Phillippy AM, et al. False gene and chromosome
losses in genome assemblies caused by GC content variation and repeats. Genome Biol.
2022;23:1–29.

43. Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, et al. Towards
complete and error-free genome assemblies of all vertebrate species. 2021;592:737–46.

44. Zhou Y, Shearwin-Whyatt L, Li J, Song Z, Hayakawa T, Stevens D, et al. Platypus and
echidna genomes reveal mammalian biology and evolution. Nat 2020 5927856.
2021;592:756–62.

45. Gould SJ. Is uniformitarianism necessary? Am J Sci. 1965;263:223–8.

46. Drabeck DH, Wiese J, Gilbertson E, Arroyave J, Stiassny MLJ, Alter SE, et al. Gene loss
and relaxed selection of plaat1 in vertebrates adapted to low-light environments. Proc R Soc
B. 2024;291.

47. Gower DJ, Fleming JF, Pisani D, Vonk FJ, Kerkkamp HMI, Peichl L, et al. Eye-
Transcriptome and Genome-Wide Sequencing for Scolecophidia: Implications for Inferring
the Visual System of the Ancestral Snake. Genome Biol Evol. 2021;13.

48. Pinheiro D, Mawhin MA, Prendecki M, Woollard KJ. In-silico analysis of myeloid cells
across the animal kingdom reveals neutrophil evolution by colony-stimulating factors. Elife.
2020;9:1–20.

49. Hron T, Pajer P, Pačes J, Bartůněk P, Elleder D. Hidden genes in birds. Genome Biol.
2015;16:164.

50. Bornelöv S, Seroussi E, Yosefi S, Pendavis K, Burgess SC, Grabherr M, et al.
Correspondence on Lovell et al.: identification of chicken genes previously assumed to be
evolutionarily lost. Genome Biol. 2017;18:112.

51. Hara Y, Kuraku S. The impact of local genomic properties on the evolutionary fate of
genes. Elife. 2023;12.

52. Rohde F, Schusser B, Hron T, Farkašová H, Plachý J, Härtle S, et al. Characterization of
chicken tumor necrosis factor-α, a long missed cytokine in birds. Front Immunol. 2018;9
APR:605.

53. Janusova S, Krchlikova V, Hron T, Elleder D, Stepanek O. Identification of GC-rich LAT
genes in birds. PLoS One. 2023;18 4 April:11–4.

54. Salve BG, Sharma S, Vijay N. Evolutionary diversity of CXCL16-CXCR6: Convergent
substitutions and recurrent gene loss in sauropsids. Immunogenet 2024 765. 2024;76:397–
415.

55. Brocker C, Thompson D, Matsumoto A, Nebert DW, Vasiliou V. Evolutionary
divergence and functions of the human interleukin (IL) gene family. Hum Genomics.
2010;5:30.

56. Nei M, Gu X, Sitnikova T. Evolution by the birth-and-death process in multigene families
of the vertebrate immune system. Proc Natl Acad Sci U S A. 1997;94:7799–806.

57. Nei M, Rooney AP. Concerted and birth-and-death evolution of multigene families. Annu
Rev Genet. 2005;39 Volume 39, 2005:121–52.

58. Kumar S, Stecher G, Suleski M, Hedges SB. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. Mol Biol Evol. 2017;34:1812–9.

59. Garceau V, Smith J, Paton IR, Davey M, Fares MA, Sester DP, et al. Pivotal Advance:
Avian colony-stimulating factor 1 (CSF-1), interleukin-34 (IL-34), and CSF-1 receptor genes
and gene products. J Leukoc Biol. 2010;87:753–64.

60. Baghdadi M, Umeyama Y, Hama N, Kobayashi T, Han N, Wada H, et al. Interleukin-34,
a comprehensive review. J Leukoc Biol. 2018;104:931–51.

61. Ko BJ, Lee C, Kim J, Rhie A, Yoo DA, Howe K, et al. Widespread false gene gains
caused by duplication errors in genome assemblies. Genome Biol. 2022;23:1–26.

62. Hume DA, Gutowska-Ding MW, Garcia-Morales C, Kebede A, Bamidele O, Trujillo
AV, et al. Functional evolution of the colony-stimulating factor 1 receptor (CSF1R) and its
ligands in birds. J Leukoc Biol. 2020;107:237–50.

63. Bayly-Jones C, Whisstock JC. Mining folded proteomes in the era of accurate structure
prediction. PLOS Comput Biol. 2022;18:e1009930.

64. Deem KD, Brisson JA. Problems with Paralogs: The Promise and Challenges of Gene
Duplicates in Evo-Devo Research. Integr Comp Biol. 2024;64:556.

65. Neves F, Marques JP, Areal H, Pinto-Pinho P, Colaço B, Melo-Ferreira J, et al. TLR7 and
TLR8 evolution in lagomorphs: different patterns in the different lineages. Immunogenetics.
2022;74:475–85.

66. Baird DM. Telomeres and genomic evolution. Philos Trans R Soc B Biol Sci. 2018;373.

67. Sedlazeck FJ, Lee H, Darby CA, Schatz MC. Piercing the dark matter: bioinformatics of
long-range sequencing and mapping. Nat Rev Genet 2018 196. 2018;19:329–46.

68. Li M, Sun C, Xu N, Bian P, Tian X, Wang X, et al. De Novo Assembly of 20 Chicken
Genomes Reveals the Undetectable Phenomenon for Thousands of Core Genes on
Microchromosomes and Subtelomeric Regions. Mol Biol Evol. 2022;39.

69. Seroussi E, Cinnamon Y, Yosefi S, Genin O, Smith JG, Rafati N, et al. Identification of
the Long-Sought Leptin in Chicken and Duck: Expression Pattern of the Highly GC-Rich
Avian leptin Fits an Autocrine/Paracrine Rather Than Endocrine Function. Endocrinology.
2016;157:737–51.

70. Neves F, Abrantes J, Steinke JW, Esteves PJ. Maximum-likelihood approaches reveal
signatures of positive selection in IL genes in mammals. Innate Immun. 2014;20:184–91.

71. Neves F, Abrantes J, Almeida T, De Matos AL, Costa PP, Esteves PJ. Genetic
characterization of interleukins (IL-1α, IL-1β, IL-2, IL-4, IL-8, IL-10, IL-12A, IL-12B, IL-15
and IL-18) with relevant biological roles in lagomorphs. Innate Immun. 2015;21:787–801.

72. Sironi M, Cagliani R, Forni D, Clerici M. Evolutionary insights into host-pathogen
interactions from mammalian sequence data. Nat Rev Genet. 2015;16:224–36.

73. Tenthorey JL, Emerman M, Malik HS. Evolutionary Landscapes of Host-Virus Arms
Races. Annu Rev Immunol. 2022;40 Volume 40, 2022:271–94.

74. Wang X, Grus WE, Zhang J. Gene Losses during Human Origins. PLOS Biol.
2006;4:e52.

75. Helsen J, Voordeckers K, Vanderwaeren L, Santermans T, Tsontaki M, Verstrepen KJ, et
al. Gene Loss Predictably Drives Evolutionary Adaptation. Mol Biol Evol. 2020;37:2989–
3002.

76. Brandis G, Hughes D. The SNAP hypothesis: Chromosomal rearrangements could
emerge from positive Selection during Niche Adaptation. PLoS Genet. 2020;16.

77. van der Loo W, Magalhaes MJ, de Matos AL, Abrantes J, Yamada F, Esteves PJ.
Adaptive Gene Loss? Tracing Back the Pseudogenization of the Rabbit CCL8 Chemokine. J
Mol Evol. 2016;83:12–25.

78. Baghdadi M, Endo H, Tanaka Y, Wada H, Seino K ichiro. Interleukin 34, from
pathogenesis to clinical applications. Cytokine. 2017;99:139–47.

79. Walker DG, Tang TM, Lue LF. Studies on colony stimulating factor receptor-1 and
ligands colony stimulating factor-1 and interleukin-34 in Alzheimer's disease brains and
human microglia. Front Aging Neurosci. 2017;9 AUG:271250.

80. Cheng ST, Tang H, Ren JH, Chen X, Huang AL, Chen J. Interleukin-34 inhibits hepatitis
B virus replication in vitro and in vivo. PLoS One. 2017;12:e0179605.

81. Hossainey MRH, Hauser KA, Garvey CN, Kalia N, Garvey JM, Grayfer L. A perspective
into the relationships between amphibian (Xenopus laevis) myeloid cell subsets. Philos Trans
R Soc B Biol Sci. 2023;378.

82. Lopez Ruiz V, Robert J. The amphibian immune system. Philos Trans R Soc B.
2023;378.

83. Lin H, Lee E, Hestir K, Leo C, Huang M, Bosch E, et al. Discovery of a Cytokine and Its
Receptor by Functional Screening of the Extracellular Proteome. Science (80-).
2008;320:807–11.

84. Wang Y, Szretter KJ, Vermi W, Gilfillan S, Rossini C, Cella M, et al. IL-34 is a tissue-
restricted ligand of CSF1R required for the development of Langerhans cells and microglia.
Nat Immunol. 2012;13:753–60.

85. Greter M, Lelios I, Pelczar P, Hoeffel G, Price J, Leboeuf M, et al. Stroma-Derived
Interleukin-34 Controls the Development and Maintenance of Langerhans Cells and the
Maintenance of Microglia. Immunity. 2012;37:1050–60.

86. Kagohashi K, Sasaki Y, Ozawa K, Tsuchiya T, Kawahara S, Saitoh K, et al. Role of
Signal-Transducing Adaptor Protein-1 for T Cell Activation and Pathogenesis of
Autoimmune Demyelination and Airway Inflammation. J Immunol. 2024;212:951–61.

87. Banks CAS, Boanca G, Lee ZT, Eubanks CG, Hattem GL, Peak A, et al. TNIP2 is a Hub
Protein in the NF-κB Network with Both Protein and RNA Mediated Interactions. Mol Cell
Proteomics. 2016;15:3435–49.

88. Van Huffel S, Delaei F, Heyninck K, De Valck D, Beyaert R. Identification of a Novel
A20-binding Inhibitor of Nuclear Factor-κB Activation Termed ABIN-2. J Biol Chem.
2001;276:30216–23.

89. Zhao L, Wang S, Xu M, He Y, Zhang X, Xiong Y, et al. Vpr counteracts the restriction of
LAPTM5 to promote HIV-1 infection in macrophages. Nat Commun 2021 121. 2021;12:1–
14.

90. O'Leary NA, Cox E, Holmes JB, Anderson WR, Falk R, Hem V, et al. Exploring and
retrieving sequence and metadata for species across the tree of life with NCBI Datasets. Sci
Data 2024 111. 2024;11:1–10.

91. Lovell JT, Sreedasyam A, Schranz ME, Wilson M, Carlson JW, Harkess A, et al.
GENESPACE tracks regions of interest and gene copy number variation across multiple
genomes. Elife. 2022;11.

92. Cer RZ, Donohue DE, Mudunuri US, Temiz NA, Loss MA, Starner NJ, et al. Non-B DB
v2.0: A database of predicted non-B DNA-forming motifs and its associated tools. Nucleic
Acids Res. 2013;41.

93. Rangwala SH, Rudnev D V., Ananiev V V., Oh DH, Asztalos A, Benica B, et al. The
NCBI Comparative Genome Viewer (CGV) is an interactive visualization tool for the
analysis of whole-genome eukaryotic alignments. PLOS Biol. 2024;22:e3002405.

94. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. J Mol Biol. 1990;215:403–10.

95. Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP. igv.js: an embeddable JavaScript
implementation of the Integrative Genomics Viewer (IGV). Bioinformatics. 2023;39.

96. Johnson MG, Gardner EM, Liu Y, Medina R, Goffinet B, Shaw AJ, et al. HybPiper:
Extracting coding sequence and introns for phylogenetics from high-throughput sequencing
reads using target enrichment. Appl Plant Sci. 2016;4.

97. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, et
al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
Genomic Era. Mol Biol Evol. 2020;37:1530–4.

98. Paradis E, Blomberg S, Bolker B, Brown J, Claramunt S, Claude J, et al. Analyses of
Phylogenetics and Evolution [R package ape version 5.8-1]. CRAN Contrib Packag. 2024.
https://doi.org/10.32614/CRAN.PACKAGE.APE.

99. Guo Q, Pan Y, Dai W, Guo F, Zeng T, Chen W, et al. A near-complete genome assembly
of the bearded dragon Pogona vitticeps provides insights into the origin of Pogona sex
chromosomes. Gigascience. 2025;14.

100. Patel HR, Alreja K, Reis ALM, Chang JK, Chew ZA, Jung H, et al. A near telomere-to-
telomere phased genome assembly and annotation for the Australian central bearded dragon
Pogona vitticeps. Gigascience. 2025;14.

101. Emms DM, Kelly S. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019;20:1–14.

102. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34:3094–100.

103. Tang AD, Soulette CM, van Baren MJ, Hart K, Hrabeta-Robinson E, Wu CJ, et al. Full-length transcript characterization of SF3B1 mutation in chronic lymphocytic leukemia reveals downregulation of retained introns. Nat Commun. 2020;11.

104. Martin FJ, Amode MR, Aneja A, Austine-Orimoloye O, Azov AG, Barnes I, et al. Ensembl 2023. Nucleic Acids Res. 2023;51:D933–41.

105. Madrigal G, Minhas BF, Catchen J. Klumpy: A tool to evaluate the integrity of long-read genome assemblies and illusive sequence motifs. Mol Ecol Resour. 2024;:e13982.

106. Beauclair L, Ramé C, Arensburger P, Piégu B, Guillou F, Dupont J, et al. Sequence properties of certain GC rich avian genes, their origins and absence from genome assemblies: Case studies. BMC Genomics. 2019;20:734.

107. Rice P, Longden L, Bleasby A. EMBOSS: The European Molecular Biology Open Software Suite. Trends Genet. 2000;16:276–7.

108. Kolde R. pheatmap: Pretty Heatmaps. CRAN Contrib Packag. 2010. https://doi.org/10.32614/CRAN.PACKAGE.PHEATMAP.

109. Ives AR, Garland T. Phylogenetic Logistic Regression for Binary Dependent Variables. Syst Biol. 2010;59:9–26.

110. Krchlíková V, Hron T, Těšický M, Li T, Ungrová L, Hejnar J, et al. Dynamic Evolution of Avian RNA Virus Sensors: Repeated Loss of RIG-I and RIPLET. Viruses. 2023;15.

111. Stanley ER, Chitu V. CSF-1 receptor signaling in myeloid cells. Cold Spring Harb Perspect Biol. 2014;6.

112. Sela I, Ashkenazy H, Katoh K, acids research TP-N, undefined 2015. GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. academic.oup.com. 2015;43:7–14.

113. Löytynoja A. Phylogeny-aware alignment with PRANK. Methods Mol Biol. 2014;1079:155–70.

114. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25:1972–3.

115. Kosakovsky Pond SL, Frost SDW, Muse S V. HyPhy: Hypothesis testing using phylogenies. Bioinformatics. 2005.

116. Yang Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Mol Biol Evol. 2007;24:1586–91.

117. Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, Kosakovsky Pond SL. Less is more: an adaptive branch-site random effects model for efficient detection of episodic diversifying selection. Mol Biol Evol. 2015;32:1342–53.

118. Murrell B, Weaver S, Smith MD, Wertheim JO, Murrell S, Aylward A, et al. Gene-wide identification of episodic selection. Mol Biol Evol. 2015;32:1365–71.

119. Kosakovsky Pond SL, Frost SDW. Not so different after all: A comparison of methods for detecting amino acid sites under selection. Mol Biol Evol. 2005;22:1208–22.

120. Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL. Detecting Individual Sites Subject to Episodic Diversifying Selection. PLOS Genet.

2012;8:e1002764.

121. Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. RELAX:
Detecting Relaxed Selection in a Phylogenetic Framework. Mol Biol Evol. 2015;32:820–32.

122. Poon AFY, Lewis FI, Kosakovsky Pond SL, Frost SDW. An Evolutionary-Network
Model Reveals Stratified Interactions in the V3 Loop of the HIV-1 Envelope. PLOS Comput
Biol. 2007;3:e231.

123. Yang Z, Nielsent R. Codon-Substitution Models for Detecting Molecular Adaptation at
Individual Sites Along Specific Lineages. Mol Biol Evol. 2002;19:908–17.

124. Wickham H. ggplot2. 2016. https://doi.org/10.1007/978-3-319-24277-4.

125. Dutheil JY, Gaillard S, Stukenbrock EH. MafFilter: A highly flexible and extensible
multiple genome alignment files processor. BMC Genomics. 2014;15:1–10.

Acknowledgements

We gratefully acknowledge the Vertebrate Genomes Project (VGP;

https://vertebrategenomesproject.org) for generating and making available the high-quality

genome assemblies that greatly facilitated this work. We thank the Council of Scientific &

Industrial Research (CSIR) and the University Grant Commission (UGC) for a fellowship to

BGS and SH, respectively. We used BioRender (https://biorender.com) to arrange figures and

images of species.

Funding

980   This article was funded by the Department of Biotechnology, Ministry of Science and

981   Technology, India (grant no. BT/11/IYBA/2018/03) and Science and Engineering Research

982   Board (grant no. ECR/2017/001430) provided funds for procuring computational resources

983   (i.e. Har Gobind Khorana Computational Biology cluster) used.

Author information

Authors and Affiliations

Buddhabhushan Girish Salve (buddhabhushan21@iiserb.ac.in), Sonal H

(sonal24@iiserb.ac.in), Nagarjun Vijay (nagarjun@iiserb.ac.in)

Computational Evolutionary Genomics Lab, Department of Biological Sciences, IISER

Bhopal, Bhauri, Madhya Pradesh, India

Contributions

BGS and NV. conceived and designed the study; BGS, SH, and NV performed research and

analysed data; and BGS, SH and NV wrote the manuscript. All authors have read and

approved the final version of the manuscript and agree with the order of presentation of the

authors.

Corresponding author

Correspondence to Nagarjun Vijay (nagarjun@iiserb.ac.in)

Ethics declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

1006

Figure legends

Fig. 1

Unretrieved orthologs are localised to discrete chromosomal regions, occur within syntenic

blocks, and are associated with chromosomal rearrangements. The left panel shows the

genomic locations of 53 unretrieved orthologs mapped across 21 chromosomes of the

Chicken karyotype (black horizontal bars scaled by genomic position on the x-axis, and

chromosome numbers are shown on the left side). Unretrieved orthologs are indicated by

light green dots labelled with gene names. GC content, calculated in 100 bp windows, is

represented by a gradient from light grey (low GC) to red (high GC). Below the GC track, the

presence of predicted non-B DNA motifs is indicated by colour-coded bars. The right panel

depicts chromosomal rearrangements across representative sauropsid species (*Gallus gallus*,

Alligator mississippiensis, *Chelonia mydas*, *Pogona vitticeps*, *Anolis sagrei*, and *Thamnophis*

elegans). The chromosome numbers are mentioned above each grey-coloured rectangle, and

coloured ribbons (as defined in the left panel) represent conserved synteny blocks connecting

orthologous regions, with seven regions of interest (ROI_1 to ROI_7) highlighted. These

regions are implicated in the loss of two or more genes**.** The chicken orthologues of the

human genes *IFTAP*, *PLAAT1*, and *TMEM273* correspond to *C11orf74*, *HRASLS*, and

C10orf128, respectively.

Fig. 2

Multiple lines of evidence support gene loss in the Squamate lineage. **A.** Evidence types used

to validate candidate gene losses/unretrieved orthologs. The left panels summarise evidence

of segmental deletions (n = 36 genes) and chromosomal rearrangements (n = 17 genes).

Upset plots display intersections of evidence categories, including synteny, genome blast

coverage, GC content, sequence identity, paralog status (Ensembl), closest paralog placement

(Hybpiper and IQ-TREE-based phylogenetic analysis), phylogenetic signal for segmental

deletion, intergenic distance-based evidence of segmental deletion, and assembly gaps found

using klumpy. Horizontal bar plots show total set size for each evidence type, while vertical

bars represent intersection sizes and gene names are mentioned on bars (the font size varies

with increase in evidence). The stacked bar plots below indicate cumulative evidence scores

for each gene, with contributions color-coded by evidence type. **B.** Quantitative evaluation of

evidence metrics across 53 candidate gene losses/unretrieved orthologs. Violin plots display

distributions of (i) maximum genome coverage obtained using BLASTn search for each

squamate genome, (ii) sequence identity (%), (iii) GC content (%), and (iv) GC stretches.

Each gene is represented by its own distribution profile, with dashed lines marking thresholds

used to support gene classification as candidate gene losses or unretrieved orthologs.

Fig. 3

Unretrieved orthologs often belong to multi-gene families of paralogues. Human genes

corresponding to squamate- unretrieved orthologs are plotted based on their GC-content (x-

axis) and paralogs (y-axis) as annotated in Ensembl. Each point represents a paralogue, with

the size of the light-coloured filled circle indicating the percentage sequence identity between

the unretrieved orthologs and their paralog. Dashed lines indicate an approximate GC content

of 55% for both axes (vertical red for the unretrieved orthologs and horizontal blue for the

paralogues). This representation highlights that genes lost in squamates often belong to multi-

gene families with moderate GC content.

Fig. 4

LASTZ alignment-based evidence for candidate gene loss in squamates. **A.** Heatmap of

normalised aligned regions at the exonic region of Chicken (alignment length/CDS length)

for 53 genes across 34 amniote species. Squamates show consistently reduced alignment at

focal loci compared to other clades. In some cases, where alignment is detected, it may

originate from paralogous sequences rather than true orthologs, while unaligned regions may

reflect highly divergent orthologs. **B.** Boxplots of normalised aligned regions across clades

for left, focal, and right gene positions. Significant reductions (wilcox.test; alternative =

"greater") are observed at focal loci in squamates, which is not the case for other orders. **C.**

Comparison of Ensembl LASTZ chains (n = 7 species) and manually created LASTZ chains

in this study (n = 6 species with two genomes of *Pogona vitticeps*, of squamate species

BUSCO >97) shows no significant difference (Wilcoxon test, p = 0.16). **D.** Clade-level

comparison highlights significantly lower alignment coverage in squamates relative to birds,

crocodilians, testudines, and other lineages (wilcox.test; alternative = "greater"). Statistically

significant pairwise differences are indicated by bars above the boxplots (*□p□≤□0.05,

□p□≤□0.01, *□p□≤□0.001, ****□p□≤□0.0001, ns□=□non-significant). Boxplots

are generated using ggplot [124].

Fig. 5

Evidence for *IL34* gene loss via segmental deletion. **A.** Micro-gene synteny of *IL34* in 24

vertebrate representative species. The phylogenetic branches of species with *IL34* lost are

depicted with dashed red, while blue phylogenetic branches represent the 3rd round of whole

genome duplication. The green-coloured thunderbolt marks the gene loss event in squamates.

Arrows represent genes, with their direction indicating gene orientation and gene names

labelled inside each arrow. **B.** The heatmap shows the reduction in gene distance at the *IL34*

locus (between *SF3B3-MTSS2*) in squamates. The values inside the box represent adjusted p-

values (significant differences between squamates and non-squamates, calculated using

pairwise Wilcox one-tailed test (alternative = "less"). **C.** Phylogenetic logistic regression

analyses. The relative genomic distance between flanking genes (x-axis, in %) and the status

of the *IL34* gene (y-axis) are correlated using two methods (logistic_IG10 and

logistic_MPLE) to evaluate the segmental deletion of the *IL34* gene across vertebrates

(n□=□179). The species tree was obtained from the Timetree website (https://timetree.org/)

and annotated in iTOL (https://itol.embl.de/). Species images are from https://BioRender.com.

Fig. 6

Validation of genome assembly at *IL34* gene syntenic locus in common wall lizard and Indian

cobra genomes using long-read sequencing. UCSC Genome Browser snapshots of the *IL34*

syntenic region are shown for the common wall lizard (*Podarcis muralis*; panel A,

chr8:39,120,000–39,125,000) and the Indian cobra (*Naja naja*; panel B, chr4:770,000–780,000). The top portion of each panel displays GC content (black line), annotated syntenic genes (*MTSS2*/*SF3B3* in *P. muralis*, E2320_000953/958 in *N. naja*), and RepeatMasker annotations (including SINEs, LINEs, and low-complexity elements). The bottom portion of both panels shows long-reads from PacBio and Nanopore spanning the entire region. Grey and yellow segments indicate forward- and reverse-strand alignments, respectively. Transparent red and blue boxes mark exons of flanking syntenic genes. SRA (Short Read Archive) accession numbers for the datasets used to generate the BAM files are indicated next to the plots.

Fig. 7

Indirect evidence for *IL34* loss: Co-evolution of receptor *CSF1R*. Selection analysis of the *CSF1R* coding sequence was conducted across 32 vertebrate species using both branch-based (RELAX, aBSREL, and CodeML) and site (FEL and MEME) models. The phylogenetic tree (left) illustrates the evolutionary relationships among species, with squamate lineages highlighted in red text and red box. Selection inferences such as relaxed/intensified and positive selection, based on RELAX, CodeML, and aBSREL analyses, are shown for each species. The central panel displays the distribution of positively and negatively selected codon sites along the gene, with codon positions on the x-axis and selection signals from focal species (foreground) against the rest of the species in the background. The right panel summarises the number of positively selected sites detected by FEL and MEME for each species, visualised using ggplot2 [124].

Fig. 8

Evolutionary relationships and genomic context of caveolin (CAV) genes in Squamates.

(A) Maximum-likelihood phylogeny of *CAV1*, *CAV2*, *CAV3*, and *CAV3*-like genes across representative vertebrates. *CAV1* and *CAV2* form distinct clades, while *CAV3* and *CAV3*-like (blue branch) sequences cluster separately. The phylogenetic tree made using IQ-TREE and rooted to *CAV2*. The bootstrap support values are indicated at nodes. (B) Synteny of the *CAV3*-like locus in Pogona vitticeps. (C) Transcriptomic evidence for the *CAV3*-like gene in central bearded dragon (*Pogona vitticeps*). Long-read (Nanopore) and short-read RNA-seq data confirm exon–intron structure and transcriptionally active status of *CAV3*-like (*LOC110078681*) gene in *Pogona vitticeps*.

Danio rerio
Oryzias melastigma
Poecilia formosa
Poecilia reticulata
Latimeria chalumnae
Xenopus tropicalis
Rana temporaria
Hyla sarda
Bufo bufo
Ornithorhynchus anatinus
Phascolarctos cinereus
Loxodonta africana
Homo sapiens
Mus musculus
Rhinolophus ferrumequinum
Orcinus orca
Equus caballus
Canis lupus
Sphenodon punctatus
Rhineura floridana
Thamnophis elegans
Varanus komodoensis
Pogona vitticeps
Anolis carolinensis
Dermochelys coriacea
Caretta caretta
Alligator mississippiensis
Gallus gallus
Cygnus olor
Calypte anna
Taeniopygia guttata
Rissa tridactyla

0   250   500   750   1000
Codon

-60 -40 -20 0 20 40
No. of Sites

Relaxed selection (RELAX)   Relaxed selection (codeml)   Positive selection (aBSREL)
Intensified selection (RELAX)   Intensified selection (codeml)   non-significant
↑ Positive sites (FEL)   Negative sites (FEL)   ■ Positive sites (MEME)   ● No. of Positive sites (FEL)   ● No. of Positive sites (MEME)