1 **Transgenerational transmission of an environmental modification in quails:**

2 **changes in phenotypic variance components across three generations**

S. Rousse [a], S. Leroux [a], D. Gourichon [b], L. Calandreau [c], I. David [a], A. Rau [d], S.

Lagarrigue [e], T. Zerjal [d], F. Pitel [a], S.E. Eynard [a]

[a] *GenPhySE, Université de Toulouse, INRAE, ENVT, 31326, Castanet-Tolosan,*

France

[b] *PEAT, UE1295 - INRAE Centre Val de Loire, 37380, Nouzilly,* France

[c] *INRAE, CNRS, IFCE, Université de Tours, Centre Val de Loire UMR Physiologie de*

la Reproduction et des Comportements, 37380 Nouzilly, France

[d] *GABI, Université Paris Saclay, INRAE, AgroParisTech, 78350 Jouy-en-Josas,*

France

[e] *PEGASE, INRAE, Institut Agro, 35590 Saint Gilles,* France

Corresponding author:

Stacy Rousse. Email: stacy.rousse@inrae.fr

Sonia E. Eynard. Email: sonia.eynard@inrae.fr

17 **Abstract**

While epigenetic variations can contribute to shaping phenotypic diversity, it can be

challenging to isolate and quantify the portion of trait variability under epigenetic

influence. The present study compares the phenotypic response for different

phenotypes of two epilines of Japanese quails (*Coturnix japonica*), with similar

genetic structure across three generations following an initial genistein ingestion in

the ancestors' diet. Linear and linear mixed models were fitted to extract the fraction

of variance allocated to multiple factors such as dam, family, sex, reproductive status

and epiline. The latter was found to be significantly associated with body weight and

with heart, abdominal fat and testicle weight adjusted by body weight. The proportion

of phenotypic variance explained by the epiline progressively increased from the first

generation (G0) to the last (G2), leading - for example in body weight at slaughter - to

an average difference for adult males and females in G2 of 9 grams and 14 grams

respectively, between genistein treated offspring (epi+) and controls (epi-). These

findings suggest a possible transgenerational effect of the initial diet disruption. In

tissue weights, production and behavioural traits, the effect of the epiline accounted

for smaller fractions of phenotypic variability and overall large residual variances

were observed, underlining the complex nature of the analysed traits. This innovative

experimental design allowed for a characterization of the impact of transgenerational

effects on a variety of phenotypic traits, which will be useful for prioritizing traits for

future genetic and epigenetic association studies.

Keywords

Epigenetic, Phenotype, Environmental modification, Transgenerational, *Coturnix*

japonica

Implications

Nowadays, changes in the environment occur more frequently and livestock species

can struggle to adapt to these rapid shifts. It is therefore important to understand the

impact of the environment on animals. In this study, we evaluate how an initial

modification in prior generations can also affect subsequent ones. By collecting

different traits of interest (growth, tissues, production, and behaviour), we compare

the response to a change in the ancestors diet across three successive generations.

Our findings highlight the importance of considering the potential multi-generational

effects of changes to the environment. This could provide guidelines for better poultry

management in order to ensure animal welfare while maintaining sustainable

production yields.

Introduction

The evolution of farming practices as well as the increased frequency of

environmental constraints due to climate change can be sources of stress for farm

animals. Heat waves, variability of food accessibility and quality are examples of

environmental changes livestock species must deal with. One of the key drivers of

adaptation is the emergence of new phenotypes, which arise from the combined

effects of genetic influence of parental alleles, non-genetic factors and the interaction

between them. Among non-genetic factors, epigenetic phenomena, such as DNA

methylation or histone modifications, can be involved in the construction of

phenotypes (Duncan et al., 2014). Fluctuations in the parental environment can

trigger epigenetic modifications, resulting in changes in gene regulatory mechanisms

and thus gene expression, even in the progeny (Guerrero-Bosagna et al., 2018).

Understanding the impact and transmission of environmental effects to subsequent

generations is crucial not only to comprehend how animals adapt to changing

environments but also to respond to production needs while ensuring animal

welfare.

Recently, the impact of the parental epigenome on offspring's developmental

phenotypes has been a research topic of growing interest in different livestock

species like cattle (Kiefer et al., 2021), pig (Wu et al., 2024), sheep (Fonseca et al.,

2023) and poultry (Hu et al., 2020). Even if most studies tackle the direct impact of

the environment on the traits of exposed animals and on the performances of the

subsequent generation, a handful have managed to evaluate the possible

transgenerational inheritance of epigenetic phenotypes as defined by Skinner 2008.

In this study, we align with this definition and consider an environmental effect to be

transgenerationally transmitted when the individuals exhibiting trait variations were

not exposed to the environmental stressor, even as germ cells. Transgenerational

studies in farm animal are challenging due to the need to raise multiple generations

and the large number of animals involved. To this end, quail is an ideal model for

transgenerational analysis, thanks to its short generation interval, the wide range of

measurable phenotypes, and the controlled breeding practices and limited space in

which it can be reared. In addition, despite the predominance of chickens, Japanese

quails (*Coturnix japonica*) were estimated to account for around 11% of the global

share in poultry meat and egg production worldwide based on 2018 FAO data

(Lukanov 2019). Previous studies on a variety of traits such as body weight, egg

production and behavioural characteristics using Japanese quails as model

organisms have reported multi and transgenerational phenotypic responses of the

progeny after a thermal increase in the parental embryonic environment (Vitorino

Carvalho et al., 2023), after in-ovo injection of genistein, an endocrine disruptor

(Leroux et al., 2017), and after exposure to maternal stress (Charrier et al., 2024).

These studies suggest a possible transmission of environmental effects through

changes occurring in the prenatal embryonic environment. However, to our

knowledge, no study in birds has specifically examined the impact of a less intrusive

environmental modification close to current rearing conditions, such as a shift in the

dam diet on complex phenotypes of offspring. Some evidence of the importance of

grandmother's diet was documented in mule and Muscovy ducks, with significant

variations in ducklings and adult weights after methionine deprivation in the diets of

the grandmothers (Brun et al., 2015). This suggests that diet disruptions can indeed

be a cause of phenotypic response to an unknown yet memorised environment,

through a specific inheritance mechanism.

To investigate the existing link between the environment, the epigenome and the

regulation of phenotypic expression at a transgenerational level, the present study

used genistein as the environmental stressor in the ancestor diets of Japanese

quails. Genistein is a phytoestrogen compound naturally present in soy, and was

notably associated with significant shifts in methylation profiles in the promoter region

of the Agouti gene in mice (Dolinoy et al., 2006). In human, genistein is known for its

beneficial effects on health (reducing the risk of cancer, cardiovascular disease,

diabetes, etc.), and today, its use in human medicine offers promising results for

cancer treatment regarding its methylome regulation capacities (Rasheed et al.,

2022; Casari et al., 2025). In certain regions of the world, a societal shift towards

diets with lighter daily intake of animal proteins leads consumers to compensate with

soy rich diets. However, numerous studies have demonstrated harmful effects of soy

overconsumption on reproductive functions, in both sexes (Liu et al., 2020; Lozi et

al., 2021; Rashid et al., 2022; Toktay et al., 2020) or cognitive functions (Svensson et

al., 2021). The effects of endocrine disruptors on health seem to be persistent over

several generations, although this remains relatively unexplored, despite a few

specific studies (Brulport et al., 2021). In addition, the use of genistein as

supplementation within the feed of Japanese quails has been shown to lead to better

adaptation of birds to heat stress by stimulating feed efficiency and growth rate

(Onderci et al., 2011). Genistein supplementation is thus a promising candidate

method with potential to trigger phenotypic variation through changes on the

methylome at a transgenerational level.

While epigenetic variations can contribute to shaping phenotypic diversity in

offspring, isolating and quantifying the portion of trait variation influenced by

epigenetics remains challenging. To date, although genetic contribution has largely

been documented for a wide range of traits in many poultry species, no direct

quantification of the epigenetic influence on these traits has been reported.

Regarding body weights, the genetic heritabilities described in the literature differ

largely according to studies (Gaya et al., 2006, Moreira et al., 2019, Prince et al.,

2020, Dadousis et al., 2021, Chomchuen et al., 2022, Wang et al., 2024) providing a

compelling reason to investigate how much phenotypic variability is explained by

epigenetic factors. The same conclusion can be drawn concerning behavioural traits,

for which the environment, the emotional state and personalities account for large

parts of the variabilities observed in individuals' response to stress (van Oers et al.,

2003, Wright and Henriksen, 2020, Lormant et al., 2020) without quantifying the role

of the epigenetic component. Overall, the objectives of this study are to 1) investigate

the transgenerational transmission of the effects of an initial diet disruption across

three generations of Japanese quails; 2) quantify the portion of phenotypic variability

linked to epigenetic modifications and its evolution throughout generations; and 3)

establish a starting point for further analyses using methylomes and genomes to

establish on the transgenerational inheritance mechanisms underlying trait variability.

Material and methods

Quail mating design

All animals in the experiment are from a quail experimental line (HSR, for High Social

Reinstatement) arising from a divergent selection on social motivation (Mills and

Faure, 1991). From this common genetic background, twenty initial founding families

were crossed, producing generation G-1. In each founding family, two full sib females

were mated at maturity with the same male, originating from another family. One of

the females was administered a genistein capsule (100 mg, Molekula group) for thirty

days after mating while the other one was only provided with empty capsules, leading

to the development of G0 individuals in two different embryonic environments. The

females that received the genistein are the dams of the treated epiline (thereafter

called epi+), in opposition to the control epiline (epi-). After this initial diet disruption,

the quails were mated following a mirror design, with males and females from the

same founding families mated in each line, to maximize the genetic balance between

the two epilines. Groups were then maintained under standard rearing conditions for

the three subsequent generations (G0, G1 and G2), inside a facility with water and

feed *ad libitum* (Fig.1). From egg hatching to 49 days-of-age, individuals were floor-

raised on litter altogether, in climate-controlled housing. Animals were then placed

within cages of two individuals of same sex, until the age of twenty weeks, when they

were placed in couples for mating. All diet, temperature and light conditions were

standard and the same for all animals, all reared in the same barn.

Phenotypic traits

Phenotypic records were collected over four generations for body weights, tissue

weights, egg production and behavioural response to isolation. The summary

statistics of traits collected from G0 onwards are available in Table 1. The summary

statistics of traits collected in G-1 are included in Supplementary Table S1.

Body weights (BW)

Body weights were measured at different developmental stages: one week, four

weeks, seven weeks and slaughter for all quails. A unique scale was used for all

measurements and weights were collected at the same age when possible to ensure

consistency between generations. One-, four- and seven-week body weights were

collected respectively at eight or nine days, thirty or thirty-two days and fifty or fifty-

two days. The generation G-1 was slaughtered at thirty-one weeks (217 days), G0

and G2 at thirty weeks (210 or 211 days) and G1 at twenty-eight weeks (196 or 197

days). Slaughter was done through electronarcosis immediately following the

recording of the final body measure.

Tissue weights (TW)

Several tissues were weighed: abdominal fat (for all generations), heart, liver (for G0

onwards), and testicle (for G1 onwards). Samples from those tissues were collected

from a subset of the animals used to produce the next generation, immediately

frozen in liquid nitrogen and stored for further analysis. All weights were adjusted by

correcting for the weight of the quail at slaughter.

Egg production (PROD)

At twenty weeks of age, quails were placed in cages for reproduction. Egg production

data were recorded from the first day in the cage for 49 days (total egg number

during the laying period, precocity). The laying rate was estimated as the number of

eggs laid over the laying period. The mean egg weight per quail was calculated on a

basis of four eggs collected and weighed per quail.

Behaviour at isolation (BHV)

Before slaughter, the quails underwent an open field behavioural test in which their

reactivity to a new environment was assessed. Each quail was placed in a cubic

arena (80 cm) of white wood with a floor made of beige linoleum, in an unknown

experimental room surrounded by an unknown white curtain. Each quail was

individually placed at the centre of the arena and allowed to freely explore it for 300

seconds. For each trial, the time spent in the inner area (a square of 20 cm × 20 cm

at the centre of the arena), the frequency of time spent at the periphery (outside the

inner area) as well as the total distance covered by the quail was recorded by a

camera placed above the arena. The open field test was performed for all

generations.

Statistical analysis : pairwise differences between epilines

One-way ANOVA was applied to compare the effect of epilines on the raw

phenotypes. Given the strong sexual dimorphism in quails, the test was applied to all

raw phenotypes in each sex and generation separately. The null hypothesis of H0

"No significant effect of the epiline on the phenotype" was rejected at the 5% level of

significance using a F-test. A Tukey HSD (Honestly Significant Difference) test was

then conducted to estimate pairwise differences between epilines for each phenotype

with a statistical significant epiline effect using the R 'rstatix' package (version 0.7.2).

Statistical analysis : variance components estimation

To estimate the percentage of variance explained by influential factors such as the

family, dam, sex and epiline, linear and linear mixed models were fitted to normalized

trait variables.

Data normalization

To minimize skewness and meet the assumptions of linear modeling, phenotypes

were normalized to minimize skewness using the R 'moments' package (version

0.14.1). Skewness of Log_{10}, square root and inverse transformed data were

compared to that of the raw data and the transformation associated with the lowest

skewness value was subsequently retained for each trait.

Linear models (LM)

For each phenotype, the optimal set of parameters was first determined with a linear

model and stepwise model selection using the stepAIC function of the R 'MASS'

package (version 7.3-65). A linear model was fitted for each trait as follows:

$$Y = \mu + X\beta + \varepsilon$$

where Y is the phenotype, μ is the overall mean, X is the design matrix for the fixed

effects, β is the coefficient vector of fixed effects and ε is the residual. Fixed effects

included in the model were sex (male, female), epiline (epi-, epi+), reproductive

status (0,1), family (origin from founders) and their pairwise interactions. Effects were

iteratively added or deleted until the model with the lowest Akaike information

criterion (AIC) was identified for each trait.

Linear mixed models (LMM)

In poultry, it is known that dams have a major effect on offspring growth and

development through egg size and composition (Halbersleben DL and Mussekl FE,

1922; Widowski et al., 2022). Therefore, for some traits it is important to take into

consideration the variability attributable to dam. To this end, a linear mixed model

(LMM) with family, dam and interaction parameters as random effects using the R

'lme4' package (version 1.1-37) was fit as follows, where fixed effects corresponded

to the ones obtained after LM selection :

$$Y = \mu + X\beta + Z\alpha + \varepsilon$$

where Y is the phenotype, μ is the overall mean, X is the design matrix for the fixed

effects, β is the coefficient vector of fixed effects, Z is the associate matrix design for

the random effects, α is the vector of random effects and ε is the residual.To select

the best model for each trait (LM or LMM), ANOVA was used to determine whether

the random effect of dam was significant. When it was the case, LMM was kept,

otherwise we kept LM.

Cross Validations

After selecting the optimal models for each trait on the whole dataset, we evaluated

the fit of the model using 100 random fold cross validations (80% of data in the Train

set, 20% in the Test set). For each fold, the Train set was used to fit the model, which

was then used to predict the phenotypic response of the Test set. Correlations

between observed and predicted Test phenotypes were then assessed. The

coefficient of determination R^2, which represents the overall variance of the trait

captured by the model, was computed for each fold using the r.squaredGLMM

function from the R package 'MuMIn' (version 1.48.11).

Variance components

The proportion of variance explained by each factor for a given trait was estimated in

one of two ways, according to whether the optimal model identified was a LM or a

LMM. In the case of a LM, we performed an analysis of variance (ANOVA). The total

sum of squares SS_{Total} is decomposed into the sum of squares for each factor j

SS_{Factor_j} and residual variance SS_{res}. The proportion of variance explained (PVE) of

factor j may be calculated as:

$$PVE_j = \frac{SS_{Factor\,j}}{SS_{Total}}$$

260 Estimations were performed using the R package 'car' (version 3.1-3) with Type II

261 sum of squares.

262 In LMM, variance components for the fixed effects were computed using the post hoc

263 calculation described by Hoffman and Schadt, 2016 where:

264
$$\hat{\sigma}^2_{\beta_j} = var(X_j\widehat{\beta_j}),$$

265 where $\widehat{\beta_j}$ is the estimated effects for factor j with incidence matrix X_j. The total

266 variance is thus :

$$\hat{\sigma}^2_{Total} = \sum_j \hat{\sigma}^2_{\beta_j} + \sum_k \hat{\sigma}^2_{\alpha_k} + \sigma^2_\varepsilon$$

267 so that the proportion of variance explained by the j^{th} fixed factor is:

$$PVE_j = \frac{\hat{\sigma}^2_{\beta_j}}{\hat{\sigma}^2_{Total}}$$

268 and the proportion of variance explained by the k^{th} random factor is:

269
$$PVE_k = \frac{\hat{\sigma}^2_{\alpha_k}}{\hat{\sigma}^2_{Total}} .$$

All statistical analyses were performed using R 4.5.0 (R core team).

Results

Impact of the epiline on phenotypes and evolution across generations

Significant differences between epilines were observed across several traits.

Body weights (BW)

For BW traits, females showed significant differences starting from generation G1 at

all ages with epi+ quails systematically weighing more on average than controls. The

same pattern was found in males where the groups began to consistently diverge in

G2 at all ages; however, no significant difference was observed in earlier

generations, with the exception of G0 at four weeks of age (Fig. 2).

Differences in average BW at all ages became increasingly more pronounced over

the generations. In G0, the difference between epi+ and epi- at slaughter was on

average 4 grams, which was not found to be significant. In G1, the difference

between female epilines was on average 7 grams and reached 14 grams in G2. In

males, a similar pattern was observed in G2, with epi+ quails weighing 9 grams more

on average at slaughter whereas a maximum difference of 2 grams was observed in

earlier generations (Fig. 3).

Tissue weights (TW)

For tissues, after adjustment on full body weight, significant weight differences

between epilines were found in G1 for heart weight in both males and females, as

well as in G2 males, although this latter result appears to be likely driven by the

results of a single individual (Fig. 4).

Egg production (PROD)

For PROD traits, despite the absence of significant difference in laying rate, the

difference between epilines seemed to increase on average in later generations, with

epi+ quails showing signs of slightly better productivity, even though not significant

than epi- individuals. For precocity and egg weight, results depended on the

generation. Epi+ individuals were found to start laying eggs slightly later than controls

in G1, although no difference was revealed in other generations. Egg weight was

slightly heavier for epi+ in G0, but this tendency was erased in later generations (Fig.

5).

Behaviour at isolation (BHV)

Finally, in BHV traits, females of both epilines did not show any difference in

behaviour for any of the traits, including in late generations. Epi- males in G2

exhibited a significantly stronger agitation (frequency at centre and at periphery

higher on average) than their epi+ counterparts, most probably due to the exploratory

behaviour of a few individuals only, but this observation did not translate into higher

total distance or velocity. In G1, epi- quails were found to spend significantly more

time at the centre of the arena though this tendency was not observed in other

generations (Fig. 6). Only the total distance, the frequency at periphery and the

duration at centre were considered in the next steps.

Phenotypic variance explained by parameters: evolution across generations

Effect of the epiline

Model fitting revealed that epiline as well as other parameters such as dam, sex,

family and reproductive status explained a significant portion of the variability

observed in the traits considered. For BW, epiline was found significant at all ages,

from G1 onwards. In addition, it was found to account for more variability in later

generations, with an increase in the percentage of variance explained going

respectively from 0 (G0) to 5.5% (G2) at one week of age, 1.3 (G0) to 9.4% (G2) at

four weeks, 0.5 (G0) to 6.7% (G2) at seven weeks and from 0.02 (G0) to 2.7% (G2)

at slaughter. A similar pattern across generations was observed for other traits,

including total weight adjusted abdominal fat (0.06% in G0 to 0.8% in G2) and testicle

(0 to 2.6% in G2) and frequency at the periphery (0% in G0 to 0.9% in G2). Other

phenotypes like egg weight and duration at centre behaved differently, with the

portion of variation explained by epiline decreasing between G0 and G2 (resp. 2.4%

in G0 to 0.5% in G2 for egg weight and 0.8% to 0% in G2 for the duration at centre).

Lastly, adjusted heart weight, precocity and total distance showed a higher

percentage of variance explained by the epiline in G2 compared to G0, but lower

than in G1. The laying rate and liver weight did not show a significant part of

variability related to epiline (Table 2). In a fair share of traits (BW, abdominal fat

weight, testicle weight, egg weight, precocity, total distance), a significant interaction

parameter involving the epiline was found, adding to the overall effect of this factor

(Fig. 7). Details on the estimation of the percentage of phenotypic variance related to

the interaction with epiline (among other parameters) are available in Supplementary

Table S2.

Effect of other factors

Concerning other parameters, expectedly the dam effect had a stronger influence on

BW variability in younger individuals, and was influential in two of the three PROD

traits (Fig. 7). In G0, it accounted for 7.7% of the BW at one week of age and

decreased to 3.9% at slaughter (Fig. 7A). This pattern intensified in G1 and G2, with

overall higher effect of the dam (from 16.1% at one week to 3.1% at slaughter in G1

and resp. 37.2% to 5.2% in G2). In PROD traits, the effect of the dam was somewhat

similar for egg weight with an increase in G2 compared to G1 (25.9% compared to

19.5%) but no variation was observed for precocity between G0 and G2, where it

accounted for almost a third of the total variance (Fig. 7B).

Contrary to the increase in dam effect across generations for BW, the share of

variance explained by the family declined, starting from 42.3% at one week in G0 to

16.8% in G1 and decreasing again to 3.2% in G2 (Fig. 7A). The family effect was

also stronger in young animals than adults. On the other hand, the influence of family

slightly strengthened across generations for adjusted abdominal fat, doubling in G2

from explaining 6% in G0. In other phenotypes like BHV traits, the family was found

significant for total distance at all generations and for the frequency at periphery in

G0 and G1, but with no striking change of its influence. For these traits, family was

found to explain between 10 to 15% of the overall variance (Fig. 7B).

As expected, sex had relevant effects on the variability of adult quail traits. In BW, the

proportion of variability explained by sex increased with age, from nearly 0% to 1% at

one week of age to over 50% at slaughter. (Fig. 7A). In TW and BHV, the amount of

variance explained by sex was trait dependent, yet variable across generations, with

the exception of duration at centre where the model struggled to capture the

variability. It was strongest for liver weight where it ranged from 41% to 70%, lighter

for heart weight (from 7.7 to 29%) and total distance (8 to 20%); the sex component

was least important for abdominal fat (1.3 to 8.9%) and frequency at periphery (5 to

10%) (Fig. 7B).

The reproductive status of quails was found to be significantly involved in two traits

(testicle weight in G1 and in laying rate in G0 and G1), where it explained between 3

to 9% of the variability.

Cross validations

The predictive performance of the models, evaluated using cross validations, varied

considerably across traits and generations. Traits related to BW, such as BW at

slaughter, consistently exhibited the highest prediction accuracy across all

generations ($R^2 \geq 0.79$, Corr ≥ 0.86), highlighting good model performance for

growth-related traits. In contrast, behavioural traits like the duration at centre showed

the lowest coefficients of determination and predictive values ($R^2 \leq 0.04$, Corr ≤ 0.16),

suggesting limited predictive power of our models for these traits; these results are in

line with the large residual variance components estimated for behavioural traits. For

PROD traits, adjusted egg weight had moderate R^2 values in G0 (0.56) but

decreased in G1 (0.23), recovering somewhat in G2 (0.37); the corresponding

correlation values followed a similar trend. Interestingly, the models struggled to fit

the laying rate phenotype with an extremely low R^2 (0.06) in G0 to a moderate value

in G2 (0.32) and, overall low correlation values, suggesting the limited capacity of the

models for this trait. Among TW traits, liver weight demonstrated high predictive

power in G0 and G2 (R^2= 0.73 and 0.68, Corr = 0.76 and 0.82, respectively). These

results underline the differential predictability of traits, with body composition and

weight traits generally yielding higher R^2 and correlation values compared to

behaviour or some production-related traits (Table 3).

Discussion

The aim of this study was to analyse phenotypic variations across three generations

of quails following supplementation with genistein, an endocrine disruptor.

Genistein ingestion is associated with different phenotypic responses across

generations

In both sexes, body weight (BW) was increased in the treated epiline (epi+)

compared to the control epiline (epi-), with the difference between epilines becoming

progressively more pronounced from generation G0 to G2. In a pilot study (Leroux et

al 2017), a similar pattern was observed in G2 females: genistein injection into the

egg was associated with increased BW at three weeks of age. However, in that

study, the effect was reversed in males (genistein injection associated with

decreased BW). In contrast, we did not observe such sex-specific interactions in the

present study, regardless of the generation or age considered. On the other hand,

the results obtained concur with other studies concerning the significant impact of

genistein on BW, even though the scientific community remains divided on the

impact of the phytoestrogen on the organism (Xiang et al., 2023; Rockwood et al.,

2019), with most studies tackling the direct effect of ingestion on the concerned

individual.

Additionally, production (PROD) traits did not exhibit the strong epiline-related effects

seen previously in the pilot study, where epi+ females began laying eggs more than

eight days later than control females on average, even though epiline was

systematically kept as a factor in the model, explaining a marginal yet significant

portion of the precocity and the egg weight variances. Likewise, abdominal fat weight

and behaviour (BHV) traits did not demonstrate the previously reported significant

differences between epilines. Here, they only explained a limited fraction of the

variability depending on the generation considered. The discrepancies between the

two studies, despite using the same quail genetic line and the same endocrine

disruptor, are likely attributable to differences in the method of genistein

administration: in the pilot study, 500 µg of genistein were directly injected into the

egg before incubation, whereas in the present study genistein was administered

orally to the G-1 mother to mimic more realistic breeding conditions, resulting in

substantially lower concentrations of the molecule in the egg (Lin et al., 2004).

Overall, results reveal different sensitivities of the phenotypes towards genistein

ingestion. Although this molecule is already used as a supplement to increase BW in

broilers (Lv et al., 2018), it has been observed that it affects other phenotypes as

well, but not as strongly (Navarro-Guillén et al., 2025). These different sensitivities

may explain to a certain extent the contrasts observed in potential transmission

patterns, with more pronounced effect of the epiline in BW traits in later generations

compared to PROD or BHV traits.

Missing variability highlights the complexity of phenotypes

The proportion of phenotypic variability explained by epigenetic (epiline) or genetic

(dam, family) factors displayed a clear trend with age, decreasing over the lifetime

(with the exception of week 4 in G1). This may reflect the growing influence of other

factors on BW variability, particularly the sex of the individual. Regardless, residual

variance was substantial across all traits. This is highlighted by the generally weak

values of coefficient of determination (R^2), suggesting the difficulty of the models to

accurately reflect the overall variability of traits. A possible explanation resides in the

complex determinism of phenotypic traits collected, influenced by various

environmental factors, some of which the design may have failed to eliminate despite

the controlled management. Regarding PROD traits, residual variance can be

overestimated due to the remaining skewness of the traits, even after applying a

transformation to normalize data. Heritabilities reported in the literature for these

phenotypes are thus impacted and can be very low, even in selected lines of laying

hens (Wolc et al., 2012). Additionally, limited sample sizes in each generation can

complicate the accurate estimation of genetic parameters and phenotypic scores, as

revealed by the correlation between observed and predicted values. Another

example on a New Zealand bird species found that home-range sharing and genetic

relatedness among individuals contributed to phenotypic similarities for several

morphological traits, yet residual variance remained high (Rutschmann et al., 2020).

Similarly, Beziers et al 2019 reported high residual variance for stress-induced

corticosterone levels in a barn owl population, hypothesizing an additional factor

beyond genetics or nest environment: the experienced environment. Since all

individuals from the same generation in our study were reared under identical

conditions, a plausible explanation for this residual variance would be the

transmission of the experienced environment from the first generation, possibly

involving slight variations in genistein dosage among eggs. The embryonic

environment is known to influence the individual phenotype later in life, especially in

birds (Frésard et al., 2013).

Possible multi and transgenerational transmission of genistein effect

The availability of phenotypic records for all individuals of the design highlighted a

progressive increase of the epiline effect from G0 to G2 in a handful of traits (TW,

BW, precocity). Interestingly, this phenomenon was strongly emphasized in BW traits

at all ages, where the percentage of variance explained by the epiline largely

amplified from G0 to G2. Adjusted liver, abdominal fat and testicle weights displayed

a similar trend as BW, despite the overall modest percentage of variability associated

with the epiline for those phenotypes. The progressive impact of the epiline across

generations might be explained by genetic drift (Falconer and Mackay., 1996,

Hospital and Chevalet, 1996), even though, the "mirror" mating design was

implemented to minimize genetic effects, ensuring a balanced origin of founders

between epilines in each generation.

Alternatively, an increasing environmental influence across generations has been

reported in several species, including quail (Vitorino Carvalho et al., 2023), but such

effects were generally associated with repeated environmental challenges in each

generation. Here, genistein was only administered once and individuals were not

selected, suggesting a possible transmission of the environmental effects to the

progeny. This hypothesis concurs with results found by Karami et al., 2025

(submitted). Indeed, the authors identified a significant difference between expected

and estimated sire and dams path coefficients, marking a likely environmental effect

on BW traits from the same dataset. Their result supports the possible transmission

of environmental effects via non-genetic mechanisms.

While the phenotypic effects of genistein exposure varied in nature and intensity

across generations and traits, our findings demonstrate that environmental exposure

during development can induce persistent differences between epilines, particularly

on BW. The increasing proportion of phenotypic variance explained by the epiline

across generations suggests the potential involvement of non-genetic inheritance

mechanisms, although the contribution of genetic drift cannot be excluded despite

the controlled mating design and the absence of progeny selection (Supplementary

Figure S1). To better disentangle these effects, future studies could focus on

integrating genome-wide epigenetic profiling (e.g., DNA methylation) across

generations. For the traits showing epiline effect, running EWAS and metQTL

analyses to identify genomic regions where epigenetic marks govern phenotypic

changes or where genetic variation is driving epigenetic changes will help decipher

the variance due to genetic and epigenetic mechanisms.

Ethics approval

Animals were bred at the UE1295 PEAT (Nouzilly, France, doi:

10.15454/1.5572326250887292E12) with official authorization for the animals

(APAFIS#29977-2021010717073072 v5), and PEAT agreement (D371751).

Data and model availability statement

All data are available on Zenodo, 10.5281/zenodo.15796146

(https://zenodo.org/records/15796146). The code used to run the analysis is

available on GitLab:

https://forge.inrae.fr/stacy.rousse/transgenerational_pheno_quails

Declaration of generative AI and AI-assisted technologies in the writing

process

The authors did not resort to using AI or AI-assisted technologies for this work.

Author ORCIDs

S.Rousse: https://orcid.org/0009-0007-2894-9356

S.Leroux: https://orcid.org/0000-0003-2840-5109

D.Gourichon: https://orcid.org/0000-0002-4631-0149

L.Calandreau: https://orcid.org/0000-0002-7535-5733

I.David: https://orcid.org/0000-0002-2514-6693

A.Rau: https://orcid.org/0000-0001-6469-488X

S.Lagarrigue: https://orcid.org/0000-0002-4887-7245

T.Zerjal: https://orcid.org/0000-0002-0666-5420

F.Pitel: https://orcid.org/0000-0002-1477-7633

S.Eynard: https://orcid.org/0000-0002-8609-5869

Declaration of interest

The authors declare that they do not have financial conflicts of interest in relation to

the content of the article.

Acknowledgements

The authors would like to thank Mary Ann Ottinger and Catherine Bennetau-

Pelissero for their advice and help concerning genistein. Many thanks to all the

collaborators and partners, in particular the colleagues from PEAT, especially David

Gourichon, Sandrine Rivière and Michael Troquet who carefully reared the quails and

kindly helped for all experiments. Finally, special thanks to the Genobioinfo platform

for data management.

Financial support statement

523   This project has received funding from the European Union's Horizon 2020 research

524   and innovation program under grant agreement N°101000236 (GEroNIMO). This

525   project is part of EuroFAANG (https://eurofaang.eu). Stacy Rousse is co-funded by

526   the INRAE Animal Genetics Division and the French Occitanie Region.

References

Bartoń, K., 2025. MuMIn: Multi-Model Inference. doi:10.32614/CRAN.package.MuMIn

<https://doi.org/10.32614/CRAN.package.MuMIn>, R package version 1.48.11,

<https://CRAN.R-project.org/package=MuMIn>.

Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models

Using lme4. Journal of Statistical Software, 67(1),1-48. doi:10.18637/jss.v067.i01.

Béziers, P., San-Jose, L.M., Almasi, B., Jenni, L., Roulin, A., 2019. Baseline and stress-

induced corticosterone levels are heritable and genetically correlated in a barn owl

population. Heredity 123, 337–348. https://doi.org/10.1038/s41437-019-0203-5

Brulport, A., Le Corre, L., Maquart, G., Barbet, V., Dastugue, A., Severin, I., Vaiman, D.,

Chagnon, M.-C., 2021. Multigenerational study of the obesogen effects of bisphenol S

after a perinatal exposure in C57BL6/J mice fed a high fat diet. Environ. Pollut. 270,

116243. https://doi.org/10.1016/j.envpol.2020.116243

Brun, J.-M., Bernadet, M.-D., Cornuez, A., Leroux, S., Bodin, L., Basso, B., Davail, S., Jaglin,

M., Lessire, M., Martin, X., Sellier, N., Morisson, M., Pitel, F., 2015. Influence of grand-

mother diet on offspring performances through the male line in Muscovy duck. BMC

Genet. 16, 145. https://doi.org/10.1186/s12863-015-0303-z

Casari, G., Romaldi, B., Scirè, A., Minnelli, C., Marzioni, D., Ferretti, G., Armeni, T., 2025.

Epigenetic Properties of Compounds Contained in Functional Foods Against Cancer.

Biomolecules 15, 15. https://doi.org/10.3390/biom15010015

Charrier, M., Lumineau, S., George, I., Meurisse, M., Georgelin, M., Palme, R., Angelier, F.,

Coustham, V., Nicolle, C., Bertin, A., Darmaillacq, A.-S., Dickel, L., Guémené, D.,

Calandreau, L., Houdelier, C., 2024. Maternal stress effects across generations in a

precocial bird. R. Soc. Open Sci. 11, 231826. https://doi.org/10.1098/rsos.231826

Chomchuen, K., Tuntiyasawasdikul, V., Chankitisakul, V., Boonkum, W., 2022. Genetic

Evaluation of Body Weights and Egg Production Traits Using a Multi-Trait Animal

Model and Selection Index in Thai Native Synthetic Chickens (Kaimook e-san2). Anim.

Open Access J. MDPI 12, 335. https://doi.org/10.3390/ani12030335

Dadousis, C., Somavilla, A., Ilska, J.J., Johnsson, M., Batista, L., Mellanby, R.J., Headon, D.,

Gottardo, P., Whalen, A., Wilson, D., Dunn, I.C., Gorjanc, G., Kranis, A., Hickey, J.M.,

2021. A genome-wide association analysis for body weight at 35 days measured on

137,343 broiler chickens. Genet. Sel. Evol. 53, 70. https://doi.org/10.1186/s12711-021-

00663-w

Dolinoy, D.C., Weidman, J.R., Waterland, R.A., Jirtle, R.L., 2006. Maternal Genistein Alters

Coat Color and Protects Avy Mouse Offspring from Obesity by Modifying the Fetal

Epigenome. Environ. Health Perspect. 114, 567–572. https://doi.org/10.1289/ehp.8700

Duncan, E.J., Gluckman, P.D., Dearden, P.K., 2014. Epigenetics, plasticity, and evolution:

How do we link epigenetic change to phenotype? J. Exp. Zoolog. B Mol. Dev. Evol.

322, 208–220. https://doi.org/10.1002/jez.b.22571

Falconer, D.S. and Mackay, T.F.C. (1996). Introduction to quantitative genetics. 4 . Ed.

Harlow,UK, Longman Group, 464 p. ISBN 978-0-582-24302-6

Fonseca, P.A.S., Suárez-Vega, A., Pelayo, R., Marina, H., Alonso-García, M., Gutiérrez-Gil,

B., Arranz, J.-J., 2023. Intergenerational impact of dietary protein restriction in dairy

ewes on epigenetic marks in the perirenal fat of their suckling lambs. Sci. Rep. 13,

4351. https://doi.org/10.1038/s41598-023-31546-3

Fox, J., Weisberg, S., 2019. An R Companion to Applied Regression, Third edition. Sage,

Thousand Oaks CA. <https://www.john-fox.ca/Companion/>.

Frésard, L., Morisson, M., Brun, J.-M., Collin, A., Pain, B., Minvielle, F., Pitel, F., 2013.

Epigenetics and phenotypic variability: some interesting insights from birds. Genet. Sel.

Evol. 45, 16. https://doi.org/10.1186/1297-9686-45-16

Gaya, L.G., Ferraz, J.B.S., Rezende, F.M., Mourão, G.B., Mattos, E.C., Eler, J.P., Michelan

Filho, T., 2006. Heritability and Genetic Correlation Estimates for Performance and

Carcass and Body Composition Traits in a Male Broiler Line. Poult. Sci. 85, 837–843.

https://doi.org/10.1093/ps/85.5.837

Guerrero-Bosagna, C., Morisson, M., Liaubet, L., Rodenburg, T.B., de Haas, E.N., Košťál, Ľ.,

Pitel, F., 2018. Transgenerational epigenetic inheritance in birds. Environ. Epigenetics

4, dvy008. https://doi.org/10.1093/eep/dvy008

Halbersleben DL and Mussekl FE. Relationship of egg weight to chickweight at hatching.

Poult Sci 1: 143–144, 1922

Hoffman, G.E., Schadt, E.E., 2016. variancePartition: interpreting drivers of variation in

complex gene expression studies. BMC Bioinformatics 17, 483.

https://doi.org/10.1186/s12859-016-1323-z

Hospital, F., Chevalet, C., 1996. Interactions of selection, linkage and drift in the dynamics of

polygenic characters. Genet. Res. 67, 77–87.

https://doi.org/10.1017/s0016672300033498

Hu, Y., Feng, Y., Ding, Z., Lv, L., Sui, Y., Sun, Q., Abobaker, H., Cai, D., Zhao, R., 2020.

Maternal betaine supplementation decreases hepatic cholesterol deposition in chicken

offspring with epigenetic modulation of SREBP2 and CYP7A1 genes. Poult. Sci. 99,

3111–3120. https://doi.org/10.1016/j.psj.2019.12.058

Kassambara, A., 2023. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.

doi:10.32614/CRAN.package.rstatix <https://doi.org/10.32614/CRAN.package.rstatix>,

R package version 0.7.2, <https://CRAN.R-project.org/package=rstatix>.

Kiefer, H., Sellem, E., Bonnet-Garnier, A., Pannetier, M., Costes, V., Schibler, L., Jammes,

H., 2021. The epigenome of male germ cells and the programming of phenotypes in

cattle. Anim. Front. 11, 28–38. https://doi.org/10.1093/af/vfab062

Komsta, L., Novomestky, F., 2022. moments: Moments, Cumulants, Skewness, Kurtosis and

Related Tests. doi:10.32614/CRAN.package.moments

<https://doi.org/10.32614/CRAN.package.moments>, R package version 0.14.1,

<https://CRAN.R-project.org/package=moments>.

Leroux, S., Gourichon, D., Leterrier, C., Labrune, Y., Coustham, V., Rivière, S., Zerjal, T.,

Coville, J.-L., Morisson, M., Minvielle, F., Pitel, F., 2017. Embryonic environment and

transgenerational effects in quail. Genet. Sel. Evol. 49, 14.

https://doi.org/10.1186/s12711-017-0292-7

Lin, F., Wu, J., Abdelnabi, M.A., Ottinger, M.A., Giusti, M.M., 2004. Effects of Dose and

Glycosylation on the Transfer of Genistein into the Eggs of the Japanese Quail

(Coturnix japonica). J. Agric. Food Chem. 52, 2397–2403.

https://doi.org/10.1021/jf034921f

Liu, X., Li, F., Xie, J., Huang, D., Xie, M., 2020. Fetal and neonatal genistein exposure

aggravates to interfere with ovarian follicle development of obese female mice induced

by high-fat diet. Food Chem. Toxicol. 135, 110982.

https://doi.org/10.1016/j.fct.2019.110982

Lormant, F., Ferreira, V.H.B., Meurisse, M., Lemarchand, J., Constantin, P., Morisse, M.,

Cornilleau, F., Parias, C., Chaillou, E., Bertin, A., Lansade, L., Leterrier, C., Lévy, F.,

Calandreau, L., 2020. Emotionality modulates the impact of chronic stress on memory

and neurogenesis in birds. Sci. Rep. 10, 14620. https://doi.org/10.1038/s41598-020-

71680-w

Lozi, A.A., Pinto da Matta, S.L., Sarandy, M.M., Silveira Alves de Melo, F.C., Araujo, D.C.,

Novaes, R.D., Gonçalves, R.V., 2021. Relevance of the Isoflavone Absorption and

Testicular Function: A Systematic Review of Preclinical Evidence. Evid. Based

Complement. Alternat. Med. 2021, 8853172. https://doi.org/10.1155/2021/8853172

Lukanov, H., 2019. Domestic quail (Coturnix japonica domestica), is there such farm animal?

Worlds Poult. Sci. J. 75, 547–558. https://doi.org/10.1017/S0043933919000631

Lv, Z., Fan, H., Zhang, B., Xing, K., Guo, Y., 2018. Dietary genistein supplementation for

breeders and their offspring improves the growth performance and immune function of

broilers. Sci. Rep. 8, 5161. https://doi.org/10.1038/s41598-018-23530-z

Mills, A.D., Faure, J.-M., 1991. Divergent selection for duration of tonic immobility and social

reinstatement behavior in Japanese quail (Coturnix coturnix japonica) chicks. J. Comp.

Psychol. 105, 25–38. https://doi.org/10.1037/0735-7036.105.1.25

Moreira, G.C.M., Poleti, M.D., Pértille, F., Boschiero, C., Cesar, A.S.M., Godoy, T.F., Ledur,

M.C., Reecy, J.M., Garrick, D.J., Coutinho, L.L., 2019. Unraveling genomic

associations with feed efficiency and body weight traits in chickens through an

integrative approach. BMC Genet. 20, 83. https://doi.org/10.1186/s12863-019-0783-3

Morris, K.M., Hindle, M.M., Boitard, S., Burt, D.W., Danner, A.F., Eory, L., Forrest, H.L.,

Gourichon, D., Gros, J., Hillier, L.W., Jaffredo, T., Khoury, H., Lansford, R., Leterrier,

C., Loudon, A., Mason, A.S., Meddle, S.L., Minvielle, F., Minx, P., Pitel, F., Seiler, J.P.,

Shimmura, T., Tomlinson, C., Vignal, A., Webster, R.G., Yoshimura, T., Warren, W.C.,

Smith, J., 2020. The quail genome: insights into social behaviour, seasonal biology and

infectious disease response. BMC Biol. 18, 14. https://doi.org/10.1186/s12915-020-

0743-4

Navarro-Guillén, C., Huesa-Cerdán, R., Hidalgo-Pérez, J.A., Simó-Mirabet, P., Rodríguez-

Viera, L., Martos-Sitcha, J.A., Martínez-Rodríguez, G., Perera, E., 2025. One‐carbon

nutrients and genistein as nutritional programming effectors in juvenile gilthead

seabream (Sparus aurata): Contrasting effects on phenotypic traits. Aquaculture 598,

742063. https://doi.org/10.1016/j.aquaculture.2024.742063

Onderci, M., Sahin, K., Sahin, N., Gursu, M.F., Doerge, D., Sarkar, F.H., Kucuk, O., 2004.

The effect of genistein supplementation on performance and antioxidant status of

Japanese quail under heat stress. Arch. Anim. Nutr. 58, 463–471.

https://doi.org/10.1080/00039420400020017

Prince, L.L.L., Rajaravindra, K.S., Rajkumar, U., Reddy, B.L.N., Paswan, C., Haunshi, S.,

Chatterjee, R.N., 2020. Genetic analysis of growth and egg production traits in

synthetic colored broiler female line using animal model. Trop. Anim. Health Prod. 52,

3153–3163. https://doi.org/10.1007/s11250-020-02340-4

R Core Team (2025). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rasheed, S., Rehman, K., Shahid, M., Suhail, S., Akash, M.S.H., 2022. Therapeutic

potentials of genistein: New insights and perspectives. J. Food Biochem. 46, e14228.

https://doi.org/10.1111/jfbc.14228

Rashid, R., Kumari, A., Chattopadhyay, N., Jha, R., Rajender, S., 2022. Genistein lowers

fertility with pronounced effect in males: Meta-analyses on pre-clinical studies.

Andrologia 54, e14511. https://doi.org/10.1111/and.14511

Rockwood, S., Mason, D., Lord, R., Lamar, P., Prozialeck, W., Al-Nakkash, L., 2019.

Genistein diet improves body weight, serum glucose and triglyceride levels in both

male and female ob/ob mice. Diabetes Metab. Syndr. Obes. Targets Ther. 12, 2011–

2021. https://doi.org/10.2147/DMSO.S216312

Rutschmann, A., de Villemereuil, P., Brekke, P., Ewen, J.G., Anderson, N., Santure, A.W.,

2020. Consequences of space sharing on individual phenotypes in the New Zealand

hihi. Evol. Ecol. 34, 821–839. https://doi.org/10.1007/s10682-020-10063-z

Skinner, M.K., 2008. What is an epigenetic transgenerational phenotype? Reprod. Toxicol.

25, 2–6. https://doi.org/10.1016/j.reprotox.2007.09.001

Svensson, T., Sawada, N., Mimura, M., Nozaki, S., Shikimoto, R., Tsugane, S., 2023. Midlife

Intakes of the Isoflavone Genistein and Soy and the Risk of Late-life Cognitive

Impairment: The JPHC Saku Mental Health Study. J. Epidemiol. 33, 342–349.

https://doi.org/10.2188/jea.JE20210199

Toktay, E., Selli, J., Gurbuz, M.A., Tastan, T.B., Ugan, R.A., Un, H., Halici, Z., 2020. Effects

of soy isoflavonoids (genistein and daidzein) on endometrial receptivity. Iran. J. Basic

Med. Sci. 23, 1603–1609. https://doi.org/10.22038/ijbms.2020.48294.11089

van Oers, K., Drent, P.J., de Goede, P., van Noordwijk, A.J., 2004. Realized heritability and

repeatability of risk-taking behaviour in relation to avian personalities. Proc. R. Soc.

Lond. B Biol. Sci. 271, 65–73. https://doi.org/10.1098/rspb.2003.2518

Venables, W. N. & Ripley, B. D. (2002) Modern Applied Statistics with S. Fourth Edition.

Springer, New York. ISBN 0-387-95457-0

Vitorino Carvalho, A., Hennequet-Antier, C., Rouger, R., Delaveau, J., Bordeau, T., Crochet,

S., Couroussé, N., Pitel, F., Collin, A., Coustham, V., 2023. Thermal conditioning of

quail embryos has transgenerational and reversible long-term effects. J. Anim. Sci.

Biotechnol. 14, 124. https://doi.org/10.1186/s40104-023-00924-2

Wang, J., Liu, J., Lei, Q., Liu, Z., Han, H., Zhang, S., Qi, C., Liu, W., Li, D., Li, F., Cao, D.,

Zhou, Y., 2024. Elucidation of the genetic determination of body weight and size in

Chinese local chicken breeds by large-scale genomic analyses. BMC Genomics 25,

296. https://doi.org/10.1186/s12864-024-10185-6

Widowski, T.M., Cooley, L., Hendriksen, S., Peixoto, M.R.L.V., 2022. Maternal age and

maternal environment affect egg composition, yolk testosterone, offspring growth and

behaviour in laying hens. Sci. Rep. 12, 1828. https://doi.org/10.1038/s41598-022-

05491-6

Wolc, A., Arango, J., Settar, P., O'Sullivan, N.P., Olori, V.E., White, I.M.S., Hill, W.G.,

Dekkers, J.C.M., 2012. Genetic parameters of egg defects and egg quality in layer

chickens. Poult. Sci. 91, 1292–1298. https://doi.org/10.3382/ps.2011-02130

Wright, D., Henriksen, R., 2020. Advances in understanding the genetics of poultry

behaviour. In Understanding the behaviour and improving the welfare of chickens.

Burleigh Dodds Series In Agriculture Science Number 91. (ed. Prof. Nicol, C. Royal

Veterinary College – University of London, UK). Burleigh Dodds Science Publishing.

DOI 10.19103/AS.2020.0078. pp 3-217

Wu, P., Wang, Junge, Ji, X., Chai, J., Chen, L., Zhang, T., Long, X., Tu, Z., Chen, S., Zhang,

Lijuan, Wang, K., Zhang, Liang, Guo, Z., Wang, Jinyong, 2024. Maternal

Hypermethylated Genes Contribute to Intrauterine Growth Retardation of Piglets in

Rongchang Pigs. Int. J. Mol. Sci. 25, 6462. https://doi.org/10.3390/ijms25126462

Xiang, J., Mlambo, R., Dube, P., Machona, O., Shaw, I., Seid, Y., He, Y., Luo, M., Hong, T.,

He, B., Zhou, W., Tan, S., 2023. The obesogenic side of Genistein. Front. Endocrinol.

14. https://doi.org/10.3389/fendo.2023.1308341

Table 1

Summary statistics for all phenotypic traits in different generations from G0 onwards.

Trait		G0						G1						G2					
		Nobs	mean	sd	CV	min	max	Nobs	mean	sd	CV	min	max	Nobs	mean	sd	CV	min	max
BW[1]	1 week	432	28.26	3.06	0.11	17	38	392	25.79	2.87	0.11	16	32	519	25.62	2.86	0.11	17	34
	4 weeks	432	120.94	9.71	0.08	93	152	393	120.93	10.29	0.09	81	162	514	119.63	10.94	0.09	87	153
	7 weeks	431	157.26	10.55	0.07	129	187	392	153.55	12.28	0.08	125	195	508	157.05	12.46	0.08	128	192
	slaughter	398	204.53	28.13	0.14	145	279	373	205.1	32.02	0.16	128	281	414	202.92	31.37	0.15	142	284
TW[2]	heart	135	0.01	0	0.18	0	0.02	370	0.01	0	0.15	0	0.01	408	0.01	0	0.16	0.01	0.02
	liver	161	0.03	0.01	0.23	0.01	0.05	371	0.02	0.01	0.24	0.01	0.04	410	0.02	0.01	0.31	0.01	0.04
	abdo. fat[5]	390	0.01	0	0.75	0	0.02	371	0.01	0.01	0.91	0	0.05	410	0.01	0	0.86	0	0.04
	testicle	-	-	-	-	-	-	173	0.02	0	0.23	0	0.04	207	0.02	0	0.19	0.01	0.03
PROD[3]	precocity	212	74.72	7.26	0.1	65	113	202	81.01	11.15	0.14	69	116	216	76.99	6.65	0.09	67	129
	laying rate	212	0.81	0.19	0.24	0.02	1	202	0.79	0.19	0.24	0.06	1	215	0.82	0.18	0.21	0.04	1
	egg weight	188	0.05	0	0.09	0.03	0.06	189	0.05	0	0.09	0.03	0.07	199	0.05	0	0.09	0.04	0.06
BHV[4]	total distance	404	3362.43	2309.15	0.69	68.59	14968.27	386	2691.85	1769.25	0.66	227.87	8320.07	408	3708.42	2081.7	0.56	334.19	14295.31
	duration at centre	404	125.58	83.38	0.66	0.08	300.08	374	113.08	78.67	0.7	0.28	300.04	393	117.13	76.16	0.65	0.04	300.04
	freq. at periphery[6]	404	26.23	21.06	0.8	0	159	386	25.76	19.21	0.75	0	99	408	34.76	28.15	0.81	0	242

Nobs = number of phenotypic records; mean; sd = standard deviation; CV = coefficient of variation; min = minimum; max = maximum.

[1] BW = body weights

[2] TW = tissue weights. All tissue weights are adjusted (corrected by the weight at slaughter).

[3] PROD = production traits. Egg weight is adjusted (corrected by the weight at slaughter).

[4] BHV = behavioural traits. Results to an isolation test.

[5] abdominal fat

[6] frequency at periphery

Table 2

Percentage of phenotypic variance explained by the epiline in each generation.

Type	Trait	G0	G1	G2
BW	1 week	0	2.846	5.487
			[2.139;3.552]	[4.688;6.285]
	4 weeks	1.304	5.329	9.375
		[0.806;1.802]	[4.232;6.425]	[7.779;10.97]
	7 weeks	0.471	3.624	6.718
		[0.204;0.738]	[2.733;4.515]	[5.544;7.892]
	slaughter	0.023	0.792	2.693
		[-0.01;0.055]	[0.565;1.019]	[2.267;3.118]
TW	abdo. fat[1]	0.063	0.168	0.815
		[-0.021;0.146]	[-0.018;0.354]	[0.398;1.233]
	heart	0.21	4.12	0.42
		[-0.099;0.519]	[3.231;5.009]	[0.113;0.727]
	liver	0	0	0.018
				[-0.004;0.039]
	testicle	-	0.136	2.622
			[-0.033;0.305]	[1.453;3.79]
PROD	laying rate	0	0	0
	precocity	0.267	3.205	0.166
		[-0.077;0.611]	[2.152;4.259]	[-0.083;0.415]
	egg weight	2.438	0.38	0.514
		[1.464;3.412]	[-0.003;0.764]	[0.016;1.012]
BHV	total distance	0	0.228	0.079
			[0.024;0.432]	[-0.028;0.186]
	freq. at	0	0.598	0.958
	periphery[2]		[0.22;0.977]	[0.51;1.406]
	duration at	0.792	0.227	0
	centre	[0.384;1.2]	[-0.016;0.47]	

All figures are percentages and represent the average share of phenotypic variability explained by the

epiline. The confidence interval is in the brackets and was calculated using the standard deviations

across the 100 folds of cross validations.

[1] Abdominal fat weight

[2] Frequency at periphery

Table 3

Percentage of phenotypic variance explained by the epiline in each generation.

Type	Trait	Generation	R² av	R² sd	Corr av	Corr sd
BW	1 week	G0	0.48	0.03	0.63	0.06
		G1	0.37	0.03	0.48	0.08
		G2	0.47	0.02	0.59	0.06
	4 weeks	G0	0.4	0.03	0.54	0.07
		G1	0.36	0.03	0.47	0.07
		G2	0.53	0.02	0.65	0.05
	7 weeks	G0	0.48	0.03	0.59	0.07
		G1	0.48	0.03	0.6	0.06
		G2	0.57	0.02	0.68	0.05
	slaughter	G0	0.81	0.01	0.88	0.02
		G1	0.79	0.01	0.88	0.02
		G2	0.79	0.02	0.86	0.03
TW	abdo. fat	G0	0.49	0.04	0.47	0.09
		G1	0.48	0.02	0.52	0.07
		G2	0.35	0.02	0.34	0.09
	heart	G0	0.11	0.04	0.25	0.22
		G1	0.43	0.02	0.6	0.06
		G2	0.28	0.02	0.34	0.09
	liver	G0	0.73	0.02	0.76	0.07
		G1	0.45	0.02	0.66	0.07
		G2	0.68	0.01	0.82	0.03
	testicle	G1	0.31	0.06	0.35	0.11
		G2	0.42	0.05	0.38	0.12
PROD	laying rate	G0	0.06	0.01	0.25	0.09
		G1	0.29	0.03	0.23	0.12
		G2	0.32	0.03	0.2	0.13
	precocity	G0	0.37	0.07	0.36	0.15
		G1	0.03	0.01	0.16	0.15
		G2	0.35	0.09	0.33	0.13
	egg weight	G0	0.56	0.03	0.48	0.11
		G1	0.23	0.07	0.18	0.15
		G2	0.37	0.05	0.35	0.13
BHV	total	G0	0.32	0.02	0.46	0.07
		G1	0.32	0.02	0.43	0.08
	distance	G2	0.32	0.02	0.26	0.1
	freq. at	G0	0.21	0.02	0.29	0.09
		G1	0.25	0.02	0.36	0.09
	periphery	G2	0.06	0.01	0.22	0.1
	duration at	G0	0.04	0.01	0.16	0.09
		G1	0.04	0.01	0.11	0.12
	centre	G2	0.01	0	0.14	0.03

R² av: average R²; R² sd: R² standard deviation; Corr av: average correlation; Corr sd: correlation

standard deviation. The results were obtained by estimating the model performance across cross

validations. between observed (real) and predicted data

All figures are percentages and represent the average share of phenotypic variability explained by the

epiline. The confidence interval is in the brackets and was calculated using the standard deviations

across the 100 folds of cross validations.

Figure captions

Fig. 1. Quail mirror mating design. Genistein capsules were administered to females in G-1 as their full-sib female counterparts were provided empty capsules. Both dams were then mated with the same male coming from another genetic background. The G0 progeny was divided in 2 epilines (epi+ deriving from the dams having ingested the genistein and epi- deriving from the control dams) of similar genetic background. The following matings were controlled to respect the balance of the genetic contribution from founders by ensuring a mirror between epilines (solid lines: epi- and dashed lines: epi+). The genetic contribution of founders is represented by the colors.

Fig. 2. Analysis of variance on body weights at one week, four weeks, seven weeks and at slaughter (thirty weeks) between epilines. The study was conducted for each combination of sex and generation independently. The asterisk indicates the level of significance of the Tukey-HSD test results.

Fig. 3. Average body weight differences between epilines throughout quails' lifetime across generations.

Fig. 4. Analysis of variance on adjusted tissue weights (heart, liver, abdominal fat and testicle) between epilines. The study was conducted by sex for abdominal fat, heart and liver weights and by generation for all tissues. All weights were adjusted by the body weight at slaughter. The asterisk indicates the level of significance of the Tukey-HSD test results.

Fig. 5. Analysis of variance on production traits (laying rate, precocity and egg

weight) between epilines. The study was conducted by generation. The red dotted

line in the distribution of laying rate corresponds to the mean value of each epiline.

Egg weight was corrected by the body weight at slaughter. The asterisk indicates the

level of significance of the Tukey-HSD test results.

Fig. 6. Analysis of variance on behavioural traits (total distance, velocity, frequency

at centre, frequency at periphery, duration at centre and duration at periphery)

between epilines. The study was conducted by sex and generation. The asterisk

indicates the level of significance of the Tukey-HSD test results.

Fig. 7. Phenotypic variance partition between all factors (dam, sex, family, epiline,

reproductive status and interaction) per generation. The interaction is split into 2

factors: the interaction parameter involving epiline and the interaction parameter

involving other factors. **A**: Body weights at one, four, seven and thirty (slaughter)

weeks of age in G0, G1 and G2; **B**: Tissue weights (abdominal fat, liver, heart and

testicle weights), Production traits (laying rate, precocity, and egg weight) and

Behavioural traits (total distance, duration at centre and frequency at the periphery)

in G0, G1 and G2.

pwc: **Tukey HSD**; p.adjust: **Tukey**

adj. TW

pwc: **Tukey HSD**; p.adjust: **Tukey**

20 founding families

genistein

genistein

genistein

genistein

G-1

G0

G1

G2

Legend:

founding family combination

♀

♂

epi -

epi +

BW

pwc: **Tukey HSD**; p.adjust: **Tukey**

PROD