Immune Responses to Salmonella Typhi Antigens among Typhoid Recovered Individuals

in an Endemic Region

Atharv Athavale[1], Adarsh Subramaniam[1], Aftab Hussain[1], Indu Rani[1], Deepak Kumar Rathore[1],

Santosh Kumar Upadhyay[2], Anil Kumar Pandey[3], Praveen Kumar Malik[3], Amit Awasthi[1,*]

Ramesh Chandra Rai[1,*]

[1]BRIC-Translational Health Science and Technology Institute, Faridabad-121001, Haryana,

India

[2]Department of Biotechnology, Kumaun University, Nainital- 263001, Uttarakhand, India

[3]ESIC Medical College and Hospital, Faridabad-121012, Haryana, India

Corresponding Authors:

* Amit Awasthi

BRIC-Translational Health Science and Technology Institute, Faridabad- 121001, Haryana, India

aawasthi@thsti.res.in

Telephone: (+91)1292876482

* Ramesh Chandra Rai

BRIC-Translational Health Science and Technology Institute, Faridabad- 121001, Haryana, India

rairamesh@gmail.com

Telephone: (+91)1292876218

orcid.org/0000-0002-8008-3904

Abstract

Background: Typhoid fever remains a major public health concern in endemic regions, yet the

nature of antibody responses to individual Salmonella Typhi antigens and their potential role in

modulating cellular immune functions are not well understood. This gap also applies to the long-

term immune response after the vaccination against the disease.

Objectives: To evaluate the humoral and cellular immune response against the *Salmonella Typhi*

antigens among the Typhoid recovered individuals.

Methods: ELISA was used to measure antibody responses against five *S. Typhi* antigens in

plasma samples from typhoid-recovered individuals and healthy participants. Vi polysaccharide-

specific plasma antibodies were tested for their ability to activate NK cells isolated from healthy

donors. Flow cytometry-based assays were performed to analyze T cell and B cell profiles in the

peripheral circulation of these individuals.

Results: Polysaccharide antigens mounted better IgA responses among typhoid-recovered

individuals compared with healthy participants. Anti-*S. Typhi* IgM was significantly elevated in

the typhoid group for all five antigens (p<0.05), while no significant differences were observed

in IgG responses. NK cell activation assays showed comparable levels of activation markers of

the NK cells isolated from healthy individuals and treated with the plasma of Typhoid recovered

individuals. Analysis of B cell subsets revealed largely comparable distributions between the two

groups; however, transitional B cells were elevated among the Typhoid recovered individuals.

HlyE-specific memory B cells were detectable in both the groups. CD4□ T cell subsets were

similar between typhoid-recovered and healthy individuals, whereas differences were observed

in CD8□ T cell populations, with higher TEMRA cells among recovered individuals and

relatively higher central memory CD8□ T cells in healthy participants.

Conclusion: Polysaccharide antigens elicited significantly better IgA response in recovered

typhoid participants, possibly reflecting mucosal immune priming from the gut. Persistent IgM

antibodies after recovery suggest durable IgM responses following typhoid infection. Similar

IgG levels between groups and the presence of antigen-specific memory cells in healthy

individuals highlight the influence of repeated exposure in endemic settings. Elevated

transitional B cells suggest the reinforcement of the B cells to the peripheral circulation during the disease which remained in the circulation of these individuals. However, elevated TEMRA population of the CD8 T cells has been observed in several other infectious diseases including this study. These findings demonstrate the complexity of humoral and cellular immune responses to *S. Typhi* in endemic regions.

Keywords

1. Introduction

Typhoid and paratyphoid fevers, collectively known as enteric fevers, are caused by *Salmonella enterica* serovars Typhi and Paratyphi. The morbidity and mortality because of Typhoid fever is a major concern in developing countries, especially in those with poor healthcare systems. According to the World Health Organization (WHO), there are 9 million cases of typhoid annually, resulting in 1,10,000 deaths per year (1). *S.* Typhi is transmitted via the fecal-oral route, predominantly through ingestion of contaminated food and water (2). Humans are the only natural reservoirs, and chronic carriers of the bacteria and play a critical role in sustaining transmission (3). Variations in disease presentation across endemic regions and the rise of multidrug☐resistant strains pose challenges for timely diagnosis and effective control (2,4)

Isolating *S.* Typhi using blood culture is the gold standard method of typhoid diagnosis but it has limited sensitivity and requires adequate laboratory infrastructure (5,6). Serological tests such as Widal agglutination assay and rapid immunoassays (e.g., Typhidot, Tubex) are used in resource limited areas and typically measure antibodies against O (lipopolysaccharide/LPS) and H (flagellin) antigens. These tests suffer from low sensitivity and specificity, where the obtained results are often unreliable and show cross-reactivity with other species of Salmonella (7,8). There is no reliable method for the detection of chronic typhoid and serological tests may not distinguish between current and past infection (1).

There is a gap in our understanding about *S.* Typhi infection and its pathogenesis, where the human-restricted nature of this pathogen has limited our ability to model host-pathogen interactions effectively. Adaptive immunity against *S.* Typhi includes both cellular and humoral arms (9). Humoral responses involve antibodies to diverse antigens such as surface polysaccharides (LPS [O antigen], Vi capsule) and bacterial proteins (e.g., flagellin [H antigen], HlyE toxin, outer membrane proteins). The role of antibodies in protection against *S.* Typhi is not fully understood, however anti-Vi-polysaccharide responses correlate with vaccine-induced immunity (9,10). Recent studies have identified IgA against HlyE and LPS as strong markers of acute typhoid infection (11). However, little is known about how long these antibodies persist post-infection or how they differ among individuals of endemic area without clinical illness. In addition to antibody responses, the phenotypic landscape of circulating B cell and T cell subsets

among typhoid-recovered individuals remains poorly characterized, particularly in endemic
regions. Understanding how infection shapes memory B cell and T cell compartments is
important for interpreting long-term immune responses and may provide insights into vaccine-
induced immunity against S. Typhi.

Key virulence determinants of *S*. Typhi include the Vi capsular polysaccharide, Type III
secretion systems (T3SS-1 and T3SS-2), and typhoid toxin, which facilitate intestinal invasion,
intracellular survival within macrophages, and systemic dissemination (12). Protective immunity
against typhoid fever involves antibodies directed against the Vi antigen, robust Th1 responses
characterized by IL-12 and IFN-γ production, and cytotoxic T cell lymphocyte activity that
promotes intracellular killing of bacteria (13,14). A detailed understanding of these correlates of
protection underpins current vaccine strategies (15). Both B cells and T cells play important roles
in shaping protective immunity against intracellular bacterial pathogens. B cells contribute to the
generation of antigen-specific antibody responses and memory B cell populations, while T cells
mediate cellular immunity through helper and cytotoxic responses that facilitate bacterial
clearance. However, information regarding the distribution of circulating B cell and T cell
subsets following recovery from typhoid infection remains limited.

S. Typhi capsule consists of Vi polysaccharide and is encoded by the genes of Salmonella
Pathogenicity Islands (SPI) (3,16). It has been noted in a study from India, that about 10% of the
fresh isolates lack Vi-polysaccharide antigens (17). However, strains of *S*. Typhi with Vi
polysaccharide are more infectious and virulent than Vi-negative strains (2). H-protein, a
flagellar antigen recognized by Toll-like receptor 5 (TLR5), triggers strong antibody responses,
while the Vi capsular polysaccharide antigen acts as a virulence factor by masking O antigens
and protecting the bacteria from complement-mediated lysis and phagocytosis (18,19). LPS,
composed of lipid A, core oligosaccharide, and O-specific polysaccharide chains, is recognized
by TLR4 and induces robust inflammatory responses (20,21). Interactions among these antigens
are complex; for instance, Vi expression can suppress immune recognition of LPS, altering host-
pathogen interactions (22–24). In addition to these antigens, Outer Membrane Proteins (OMPs)
represent significant immunogenic component involved with host-pathogen interactions during
S. Typhi infection. These proteins play essential roles in the pathogen's adaptation to
environmental conditions, motility, adherence, and host cell colonization. *S*. typhi OMPs have

demonstrated potent immunogenic properties, eliciting long-lasting and protective immunity,
which makes them a potential vaccine candidate, and diagnostic antigen.

In endemic regions, where typhoid fever is widespread, immune responses to *S.* Typhi are
complicated by frequent exposure to the pathogen. How different antigenic components
contribute to long-term immunity in endemic settings, where individuals may experience
repeated exposure and subclinical infections, remains poorly understood.

This study focuses on evaluating the antibody responses- IgA, IgG, and IgM; against five critical
S. Typhi antigens: Vi polysaccharide, H-protein, LPS, Hemolysin E and Outer Membrane
Protein and examining the distribution of circulating B cell and T cell subsets to understand the
long-term immune landscape shaped by *S.* Typhi exposure in endemic regions. Plasma samples
were collected three months after typhoid to assess the dynamics of the humoral immunity, with
a focus on identifying immunological patterns that could inform vaccine response studies. To
further improve our understanding about cellular innate immunity, we tested whether Vi-
polysaccharide specific antibodies can activate the NK cells, owing to the role of the NK cells in
eliminating cells infected by an intracellular pathogen. In addition, we characterized circulating
B cell and T cell subsets in order to understand the broader immune landscape following
recovery from typhoid infection. The findings of this study reveal heterogeneous humoral
responses among typhoid-recovered individuals against these antigens. Notably, the substantial
immune responses observed in healthy participants also highlight the *S.* Typhi exposure in this
region, emphasizing the challenges posed by the pathogen endemicity.

2. Materials and Methods

2.1 Ethics statement

This study was conducted at the BRIC-Translational Health Science and Technology Institute,
Faridabad, India. Participant enrolment and sample collection were done at ESIC Medical
College and Hospital, Faridabad. Written informed assent/consent was obtained from each of the
study participants. This study was approved by the Institutional Ethics Committees from both
institutes [*ESIC Hospital and Medical College, Faridabad File no. 134 X/11/13/2022- IEC/110*
and THSTI Faridabad Ref No: THS 1.8.1/ (152)].

2.2 Participant details

Blood samples were collected from a total of 80 participants **(Table 1)**. Their age, gender, infection date (in the case of typhoid-recovered patients) and blood collection date were recorded. The study population consisted primarily of rural residents and healthcare workers.. Among the 80 participants we recruited for the study, 33 had recovered from typhoid fever at least for 3 months, while 47 were healthy participants and had no recorded history of the disease.

2.3 Inclusion and Exclusion criteria

The study included participants ages 18–59 years tested for Typhoid fever. Eligible individuals were either healthy or were recovered from typhoid fever at least three months prior to blood collection. A positive Widal test was not considered as a confirmatory test; a positive Typhidot IgM/IgG or blood culture tests were used instead. Pregnant women and people with immunocompromising comorbidities were excluded from the study.

2.4 Blood collection and sample processing

Eight ml of blood was collected from each participant in the Greiner Bio-One tubes (Cat. No. 22-040-134) coated with sodium heparin. Blood samples were processed immediately to isolate plasma and peripheral blood mononuclear cells (PBMCs) using Lymphoprep density gradient centrifugation method. Plasma was stored at -80 °C and PBMCs were resuspended in fetal bovine serum (FBS) supplemented with 10% dimethyl sulfoxide (DMSO) and then stored in liquid nitrogen (25).

2.5 ELISA

2.5.1   Polysaccharide antigens

Anti- *S*. Typhi IgA, IgG and IgM antibody responses are measured against Vi polysaccharide and Lipopolysaccharide after the three-months from infection of the typhoid patients by using indirect ELISA. Nunc Maxisorp 96-well plate was coated with 100 µL of poly-l-lysine (2 µg/ml) and incubated at 37°C for 2 hours. The poly-l-lysine coating was not done in case of the H antigen. All antigens were diluted in sterile PBS (2 µg/ml for Vi polysaccharide; 1 µg/ml for

Lipopolysaccharide and H-protein). Antigens were coated and incubated overnight (37°C for Vi

polysaccharide and lipopolysaccharide; 4°C for H-protein). Coated plates were blocked with a

200 µl block buffer (1% BSA in PBS) and incubated for 1 hour at 37°C. Plasma samples were

diluted 1:100 in a diluent (1% BSA, 0.05% Tween-20 in PBS) and 100 µl/well was added and

incubated for one hour at 37°C. Then, 1/10000 secondary antibody was diluted in a buffer

containing 1% BSA, and 0.05% Tween-20 in PBS and 100 µl was added to each well and

incubated. After an hour 100 µl pNPP (p-Nitrophenyl Phosphate) substrate is added to each well

which is diluted in Tris-MgCl$_2$ and incubated for an hour at room temperature in dark conditions.

At last, 50 µl 3M NaOH is added to stop the reaction. After every step of ELISA, 4 to 5 washing

is given with 0.05% Tween-20 in PBS except after the addition of substrate. Then, optical

density is measured using a plate reader at 405 nm and 490 nm. The readings at 490 nm were

subtracted from those at 405 nm to correct for optical imperfections.

2.5.2   Protein Antigens

For the protein antigens (H-protein, Hemolysin E and OMP), similar ELISA was performed with

some modifications. HRP-TMB was used as the conjugate-substrate combination instead of

alkaline phosphatase-pNPP; because it was found to give better and consistent results during the

optimization experiments. Hence, the stop solution was changed to 2N H$_2$SO$_4$.

2.6 Avidity assay

Antibody avidity against S. Typhi Vi polysaccharide and LPS was measured using a modified

indirect ELISA protocol as described above. Briefly, plates were coated with poly-L-lysine (10

µg/mL) and incubated at 37 °C for 2 hours, followed by coating with Vi antigen or LPS (0.5

µg/mL) overnight at 37 °C. Plates were blocked with 1% BSA in PBS and plasma samples

diluted in sample diluent (1% BSA + 0.1% Brij-35 in PBS) were added and incubated for 1 hour

at 37 °C.

After sample incubation, 8 M urea (100 µL/well) was added to half of the replicate wells, while

wash buffer was added to the remaining wells as control, and the plates were incubated for 10

minutes at 37 °C. Plates were then washed and incubated with alkaline phosphatase–conjugated

secondary antibody (1:10,000 dilution) for 30 minutes at 37 °C. The reaction was developed

using pNPP substrate and stopped with 3 M NaOH. Optical density was measured at 405 nm and

490 nm using a microplate reader, and background correction was performed by subtracting the

OD at 490 nm from the OD at 405 nm.

2.7 Antibody-dependent NK cell activation

Nunc MaxiSorp® flat-bottom plate was coated with streptavidin (5 µg/ml) and incubated

overnight at 37°C. After 3 washes with PBS (200 µl/well), biotinylated Vi-polysaccharide

(26,27) (3 µg/ml) was added, and the plate was incubated at 37°C for 3 hours. After 5 washes

with PBS, diluted plasma (plasma: PBS; 1:10) from typhoid-recovered patients was added and

the plate was incubated at 37°C for 2 hours. NK cells obtained using Miltenyi MACS NK cell

isolation kit (cat no. 130-092-657) were added to the plates after 5 PBS washes. 200 µl complete

RPMI media was added, and the plate was incubated at 37°C for 12 hours. The cells from the

microplate were carefully scraped off with 100 µl/well FACS buffer (PBS + 1% FBS) and

transferred to a 96-well round bottom plate using a multichannel pipette. The step was repeated if

required, to get most of the residual cells. The cells were washed twice by adding a FACS buffer

and centrifuged at 3000 rpm for 5 minutes. Excess FACS buffer was decanted carefully, and a

cocktail of antibodies was added (40 µl/well) **(Table 2)**. Samples were acquired at

BDFACSCanto II and analyzed using FlowJo software (version v10). The gating strategy of the

data analysis has been shown in **Supplementary Figure 1**.

2.8 Labelling of HlyE Protein with the Alexa Fluor-488

219   Labelling of the HlyE protein was performed using the Alexa Fluor Microscale Protein Labelling

220   kit (Cat No. #A30006; Thermo Fisher Scientific) as per the manufacturer's instructions.

221   Fluorophore-labelled HlyE was aliquoted and stored at −20 °C for further use in testing antigen-

222   specific memory B cell response.

2.9 T cell and B cell phenotyping using flow cytometry

224   Peripheral blood mononuclear cells (PBMCs) isolated from study participants were used for

225   immunophenotyping of B cell and T cell subsets (Table 3). Cryopreserved PBMCs were thawed,

226   washed with RPMI medium supplemented with fetal bovine serum and resuspended in FACS

buffer (PBS + 1% FBS). Cells were stained with fluorochrome-conjugated antibodies against CD3, CD4, CD8, CD19, CD27, CD38, IgD, CD24, CD45RA and CCR7. Surface staining was performed for 30 minutes at 4°C in the dark. After washing with FACS buffer, cells were acquired on a BD FACSymphony A3 flow cytometer and analyzed using FlowJo software (version 10.10.0).

B cells were identified as CD3☐CD19☐ cells. Naïve (CD27☐IgD☐), switched memory (CD27☐IgD☐), unswitched memory (CD27☐IgD☐) and double negative (CD27☐IgD☐) B cell populations were analyzed within the B cell compartment. Classic memory B cells were defined as CD27☐CD38☐, and transitional B cells were defined as CD27☐CD24☐CD38^low.

T cells were gated as CD3☐ lymphocytes, followed by separation into CD4☐ and CD8☐ populations. T cell subsets were defined based on CD45RA and CCR7 expression as naïve (CD45RA☐CCR7☐), central memory (CD45RA☐CCR7☐), effector memory (CD45RA☐CCR7☐) and TEMRA (CD45RA☐CCR7☐) cells.

2.10   Statistics

All the data visualization and statistical analyses were done using GraphPad Prism 10.4.2. Outliers were removed using the ROUT method (Q=1%). All the datasets were checked for normal distribution first. The Mann-Whitney test was used for statistical analysis to compare the two datasets. p value <0.05 was considered statistically significant.

3. Results

3.1 Polysaccharide antigens lead to a better IgA response

Anti-polysaccharide IgA (to LPS and Vi) was significantly higher in recovered participants than in healthy ones. For anti-LPS IgA, median OD was ~0.5852 in typhoid-recovered vs 0.4064 in healthy participants (p=0.0087). For anti-Vi IgA, median OD was ~0.406 vs 0.273 (p=0.0140) **[Fig. 1]**. IgA responses to H antigen, and OMP did not differ significantly; however, IgA response against HlyE showed significant difference among Typhoid recovered and healthy individuals **[Fig. 1]**.

3.2 Sustained IgM response against *S.* Typhi antigen among study participants

We observed sustained IgM response against the S. Typhi antigens among the study participants. However, IgM levels were significantly elevated in the typhoid-recovered group for all five antigens **[Fig. 2]**. IgM to H antigen, HlyE, and OMP were all roughly twofold higher in the typhoid group (all p<0.01). This elevated IgM level among typhoid recovered individuals suggests durability of IgM response after typhoid, complicating the diagnosis in endemic regions. Notably, healthy participants samples showed detectable IgA and IgM levels against all the antigens, indicating that they may have been exposed to the pathogen and remained asymptomatic.

3.3 IgG antibody responses and IgG antibody avidity (against Vi PS) did not differ between typhoid-recovered and healthy individuals

No significant differences were observed in antigen-specific IgG levels between the groups. Mean IgG ODs were similar in recovered vs healthy participants for all antigens (p>0.1). This indicates that by >3 months post-infection, IgG levels against these antigens had declined to the background levels of the endemic population **[Fig. 3]**. The chaotropic agent-based modified ELISA assay for testing the avidity of the IgG antibodies yielded interesting results. The median IgG antibody avidity index for both participant groups exceeded 70%, substantiating the presence of high-affinity antibodies targeting the Vi polysaccharide antigen **[Fig. 3(i)(b)]**. A summary of the serological data has been provided in **Table 4**.

3.4 No sex-specific disparities observed in serological immune responses

The study cohort demonstrated a balanced representation of both sexes, as evident from **Table 1**. Specifically, the participant groups included approximately 55% females among typhoid-recovered individuals and 43% females in the healthy group. Sex-disaggregated analysis of the immune responses was performed to evaluate any potential disparities between immune responses in males and females against *S.* Typhi. Our analysis revealed that there is no statistically significant difference in serological immune responses between these participants **[Supplementary Figures 2-6]**. This data indicates that sex does not appear to influence the immune response at recovery to typhoid infection.

3.5 NK cell response to antibody-dependent activation

We investigated the potential of anti-Vi polysaccharide (Vi-PS) antibodies present in the plasma samples of typhoid-recovered individuals to be able to activate natural killer (NK) cells isolated from the PBMCs of the healthy participants. The primary hypothesis was that the NK cells isolated from healthy participants would exhibit distinct immune responses when exposed to anti-Vi PS antibodies. To test this, we measured NK cell activation and proliferation using molecular markers CD16, CD25 and CD69 after incubating NK cells with typhoid-recovered participant's plasma and culturing the cells for 12 hours. Cells incubated with plasma showed similar levels of activation and proliferation markers as those cultured without any plasma **[Fig. 4]**. Probably, due to the endemicity of the disease in India, the control samples also behaved like the plasma treated ones. Another possibility is that the interaction of the antigen with antibodies present in the plasma of the typhoid participants is needed to activate the NK cells.

3.6 Distribution of circulating B cell and T cell subsets

To further understand the immune landscape following typhoid infection, circulating B cell and T cell subsets were analyzed using flow cytometry.

Within the B cell compartment, the proportions of naïve, switched memory, unswitched memory and double negative B cells were comparable between typhoid-recovered and healthy individuals **(Fig. 6)**. Similarly, the frequency of classic memory B cells (CD27☐CD38☐) did not differ substantially between the two groups. However, a modest but statistically significant difference was observed in transitional B cells (CD27☐CD24☐CD38^low), where typhoid-recovered individuals showed slightly higher frequencies compared with healthy participants.

We also evaluated antigen-specific memory B cells in the two study groups; using HlyE as it elicited better humoral response among protein antigens. HlyE-specific memory B cells were detectable in both typhoid-recovered and healthy participants, indicating prior exposure to the antigen in this endemic setting. The overall frequency of HlyE-specific memory B cells and the proportion of IgG-producing or IgM-producing cells within this population did not differ significantly between the two groups **(Fig. 7)**.

Analysis of T cell subsets revealed that the distribution of CD4▯ T cell populations (naïve, central memory, effector memory and TEMRA cells) was largely similar between typhoid-recovered and healthy individuals **(Fig. 8)**. On the other hand, differences were observed within the CD8▯ T cell compartment. Typhoid-recovered individuals showed higher frequencies of CD8▯ TEMRA cells, while healthy participants showed relatively higher levels of CD8▯ central memory T cells **(Fig. 9)**. These observations suggest alterations in CD8▯ T cell differentiation and or distribution following typhoid infection.

4. Discussion

The findings of this study provide insight into the humoral response against different classes of antigens and cellular responses against *S.* Typhi in individuals from an endemic region. We observed a robust IgM response three months post-infection against all the antigens we tested. In general, the presence of IgM antibodies suggests a recent infection while IgG antibodies imply long-term exposure. However, in certain viral and bacterial infections, long-lasting IgM response has been observed (28–30). Another possibility for a prolonged IgM antibody response is possibly due to a continued presence of the pathogen or re-exposure which is common in endemic areas. The persistence of IgM antibodies further complicates the interpretation of immune status in endemic populations (31). Interestingly, while the IgM antibody levels were elevated in case of the typhoid-recovered individuals as compared to the healthy group, IgG levels were not substantially different. These results imply that IgG mediated humoral immunity against *S.* Typhi returned to its baseline level during the post three-month recovery period. Alternatively, healthy participants likely have background IgG from asymptomatic exposures which is leading to the underestimation of the IgG levels present in recovered individuals.

Another key finding of this study is the better IgA response among typhoid-recovered individuals against polysaccharide antigens (Vi-PS and LPS). Vi capsule and LPS are the antigens that are heavily expressed on *S.* Typhi and engage with immune cells. IgA antibodies are present in the mucosal secretions and are known to contribute to the local immunity in the gut, potentially preventing *S.* Typhi from establishing the infection (9,10). IgA response against LPS was also found to be an accurate diagnostic test in a human challenge study (32). These results suggest that we may need to test the immune responses against such other antigens of *S.*

Typhi to expand the repertoire of the options available to be used for multigenic screening for
Typhoid.

Polysaccharides are classic T cell independent antigens and primarily trigger IgM and IgG2
responses but generate little immunological memory (33). In regions with frequent exposure,
repeated encounters with antigens tend to sustain IgM levels. In contrast, immune response
against protein antigens are T cell dependent (34). They usually induce IgG antibodies and
promote long-term memory. In this study, IgG levels to HlyE, H protein and OMP showed no
significant difference between groups after three months of disease. This could reflect the
waning of IgG over time towards the basal level which is significantly high in endemic regions.

Another aspect explored in this study was the distribution of circulating B cell and T cell subsets
among typhoid-recovered and healthy individuals. The overall composition of major B cell
subsets was largely comparable between the two groups. However, transitional B cells showed
elevated frequencies among typhoid-recovered individuals compared with healthy participants,
indicating possible alterations in early B cell maturation following infection. The presence of
HlyE-specific memory B cells in both groups further suggests that individuals living in endemic
regions may encounter the pathogen or related antigens even in the absence of clinically
diagnosed disease. Interestingly, differences were more prominent in the CD8□ T cell
compartment. Typhoid-recovered individuals showed higher frequencies of terminally
differentiated TEMRA cells, whereas healthy individuals exhibited relatively higher proportions
of central memory CD8□ T cells. Elevated levels of TEMRA cells may be a result of repeated
antigen exposure and are associated with cytotoxic effector function (35). Therefore, the
increased frequency of this subset in recovered individuals may reflect antigen-driven
differentiation during infection. These observations support the idea that immune responses to S.
Typhi in endemic regions are shaped not only by visible infection but also by repeated exposure
to the pathogen.

The antibody-dependent NK cell activation (ADNKA) assay showed that the plasma from
typhoid-recovered individuals did not significantly enhance NK cell activation and proliferation
markers (CD16, CD25 & CD69 respectively). These findings are at variance with a previous
study (36), probably because of the modified protocol we followed for this study or due to the

possibly exposed healthy individuals from whom PBMCs were used for this assay. The mature, $CD56^{dim}CD16+$ NK cells are identified as a highly differentiated subset known for their potent cytotoxicity and capacity to induce antibody-dependent cellular cytotoxicity (ADCC); on the other hand, less mature $CD56^{bright}CD16-$ NK cells are major cytokine producers with regulatory functions (37). The enigmatic behavior of NK cell responses in this study could be attributed to several factors. One possibility is that the plasma from typhoid-recovered individuals possibly contains immunomodulatory factors that suppress NK cell activity, similar to what is observed in other chronic infections like Hepatitis C and HIV (38,39).

A limitation of this study is the lack of acute-phase samples, which might have given clear differentiation of the immune parameters tested between healthy and infected individuals. Another limitation is the age group of our participants which are mainly in the adult category and hence even healthy participants are probably exposed to the pathogen due to its endemicity in the region. Therefore, understanding the impact of endemicity on the immune response to typhoid with a larger sample size is essential for developing effective prevention and control strategies. Despite these limitations, this study provides critical insights into immune responses to *S*. Typhi antigens in endemic regions, offering an understanding of the immune correlates which could be protective. Moreover, the healthy population at a larger sample size may need to be screened against different antigens and antibodies to draw a cut-off to differentiate the symptomatic from exposed individuals, particularly for endemic regions. Additionally, the pre-existing immunity in healthy individuals, as reflected by their antibody levels, could be a result of prior, subclinical infections, contributing to the overall immune landscape of the population (40). Further, many of the recent vaccines against typhoid are having Vi-polysaccharide as a component, this study could provide a hindsight for the kind of immune response one can expect in an endemic region against the Vi-antigen. Together, these results emphasize the need for tailored vaccine strategies that considers pre-existing immunity and endemic exposure.

5. Acknowledgements

The authors wish to acknowledge BRIC☐THSTI, Faridabad for intramural funding to support this work (THSTI Core/EX-DBT-55-2022/P366). We are thankful to Jishnu Sankar and Dr. Dinesh Mahajan for help with biotinylation of the Vi-PS; and to Harshit Singh, Nafees Ahmed,

395 Anmol Gaur, Sakshi Gaur, Manas Ranjan Tripathy and Jitender Chandilla, Immunobiology and
396 Immunotherapy laboratory, BRIC☐THSTI, Faridabad for the suggestions and help. The authors
397 also thank Mr. Nitesh Dhull and Dr. Sankalp for their help with the samples and ethical approval.
398 We are particularly thankful to the participants for donating their blood samples, which made
399 this study feasible.

6. Author's contribution

Atharv Athavale☐ collected samples, performed experiments, made tables and figures, wrote,
reviewed and edited the different versions of the manuscript; Adarsh Subramaniam and Aftab
Hussain- collected samples and performed experiments; Indu Rani- performed experiments;
Deepak Kumar Rathore and Santosh K. Upadhyay- edited and reviewed the different versions of
the manuscript; Anil Kumar Pandey and Praveen Kumar Malik ☐ provided samples for the
study, reviewed the different versions of the manuscript; Amit Awasthi☐ conceptualized the
study, and reviewed the different versions of the manuscript; Ramesh Chandra Rai☐
conceptualized the study, wrote the manuscript, reviewed and edited the different versions of the
manuscript, tables and figures. All authors read and approved the final version of the manuscript.

7. Declaration of interest statement

411 The authors declare no conflict of interest.

8. Data Availability

413 Data related to the article has been presented in the form of figures and tables. Keeping
414 confidentiality of the participants the details could be shared as per the journal's policy.

9. References

1. CDC. Typhoid and Paratyphoid Fever | Yellow Book | CDC [Internet]. 2025 [cited 2025
May 13]. Available from: https://www.cdc.gov/yellow-book/hcp/travel-associated-
infections-diseases/typhoid-and-paratyphoid-fever.html

2. Parry CM, Hien TT, Dougan G, White NJ, Farrar JJ. Typhoid Fever. N Engl J Med. 2002
Nov 28;347(22):22. doi:10.1056/NEJMra020201

3. Wain J, House D, Zafar A, Baker S, Nair S, Kidgell C, et al. Vi Antigen Expression in
Salmonella enterica Serovar Typhi Clinical Isolates from Pakistan. J Clin Microbiol. 2005
Mar;43(3):3. doi:10.1128/JCM.43.3.1158-1165.2005

4. Wain J, Hendriksen RS, Mikoleit ML, Keddy KH, Ochiai RL. Typhoid fever. Lancet. 2015
Mar 21;385(9973):1136–45. doi:10.1016/S0140-6736(13)62708-7 PubMed PMID:
25458731; PubMed Central PMCID: PMC11567078.

5. Antillon M, Saad NJ, Baker S, Pollard AJ, Pitzer VE. The Relationship Between Blood
Sample Volume and Diagnostic Sensitivity of Blood Culture for Typhoid and Paratyphoid
Fever: A Systematic Review and Meta-Analysis. The Journal of Infectious Diseases. 2018
Nov 10;218(suppl_4):suppl_4. doi:10.1093/infdis/jiy471

6. Kumar P, Kumar R. Enteric Fever. Indian J Pediatr. 2017 Mar;84(3):3. doi:10.1007/s12098-
016-2246-4

7. Moses AE, Atting IA, Inyang OS. Evidence of Overlapping Infections of Dengue, Malaria
and Typhoid in Febrile Patients Attending a Tertiary Health Facility in Uyo, South-South
Nigeria. Journal of Advances in Medicine and Medical Research. 2016 Aug 3;1–9.
doi:10.9734/BJMMR/2016/27668

8. Tuuminen T, Lounamo K, Leirisalo-Repo M. A Review of Serological Tests to Assist
Diagnosis of Reactive Arthritis: Critical Appraisal on Methodologies. Front Immunol. 2013
Dec 4;4. doi:10.3389/fimmu.2013.00418

9. Sztein MB, Salerno-Goncalves R, McArthur MA. Complex adaptive immunity to enteric
fevers in humans: lessons learned and the path forward. Front Immunol. 2014;5:516.
doi:10.3389/fimmu.2014.00516 PubMed PMID: 25386175; PubMed Central PMCID:
PMC4209864.

10. Dahora LC, Jin C, Spreng RL, Feely F, Mathura R, Seaton KE, et al. IgA and IgG1 Specific
to Vi Polysaccharide of Salmonella Typhi Correlate With Protection Status in a Typhoid
Fever Controlled Human Infection Model. Frontiers in Immunology [Internet]. 2019 [cited
2024 Feb 15];10. Available from:
https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2019.02582

11. Kumar S, Nodoushani A, Khanam F, DeCruz AT, Lambotte P, Scott R, et al. Evaluation of a
Rapid Point-of-Care Multiplex Immunochromatographic Assay for the Diagnosis of Enteric
Fever. Papasian CJ, editor. mSphere. 2020 Jun 24;5(3):3. doi:10.1128/mSphere.00253-20

12. Song J, Gao X, Galán JE. Conferring Virulence: Structure and Function of the chimeric
A2B5 Typhoid Toxin. Nature. 2013 Jul 18;499(7458):7458. doi:10.1038/nature12377
PubMed PMID: 23842500; PubMed Central PMCID: PMC4144355.

13. Hornick RB. Immunological control of typhoid fever. J Egypt Public Health Assoc.
1970;45(1):236–46. PubMed PMID: 5484828.

14. Sztein MB. Cell-mediated immunity and antibody responses elicited by attenuated
Salmonella enterica Serovar Typhi strains used as live oral vaccines in humans. Clin Infect
Dis. 2007 Jul 15;45 Suppl 1:S15-19. doi:10.1086/518140 PubMed PMID: 17582562.

15. Patel PD, Patel P, Liang Y, Meiring JE, Misiri T, Mwakiseghile F, et al. Safety and Efficacy
of a Typhoid Conjugate Vaccine in Malawian Children. N Engl J Med. 2021 Sep
16;385(12):12. doi:10.1056/NEJMoa2035916 PubMed PMID: 34525285; PubMed Central
PMCID: PMC8202713.

16. Marcus SL, Brumell JH, Pfeifer CG, Finlay BB. Salmonella pathogenicity islands: big
virulence in small packages. Microbes and Infection. 2000 Feb 1;2(2):2. doi:10.1016/S1286-
4579(00)00273-2

17. Maurya P, Gulati AK, Nath G. Status of Vi gene, its expression and Salmonella
Pathogenicity Island (SPI-7) in Salmonella Typhi in India. Southeast Asian J Trop Med
Public Health. 2010 Jul;41(4):913–9. PubMed PMID: 21073066.

18. Chaban B, Hughes HV, Beeby M. The flagellum in bacterial pathogens: For motility and a
whole lot more. Seminars in Cell & Developmental Biology. 2015 Oct;46:91–103.
doi:10.1016/j.semcdb.2015.10.032

19. Dufresne K, Saulnier-Bellemare J, Daigle F. Functional Analysis of the Chaperone-Usher
Fimbrial Gene Clusters of Salmonella enterica serovar Typhi. Front Cell Infect Microbiol.
2018 Feb 8;8:26. doi:10.3389/fcimb.2018.00026

20. Chessa D, Spiga L, De Riu N, Delaconi P, Mazzarello V, Ganau G, et al.
Lipopolysaccharides Belonging to Different Salmonella Serovars Are Differentially Capable
of Activating Toll-Like Receptor 4. McCormick BA, editor. Infect Immun. 2014
Nov;82(11):11. doi:10.1128/IAI.02297-14

21. Hoare A, Bittner M, Carter J, Alvarez S, Zaldívar M, Bravo D, et al. The Outer Core
Lipopolysaccharide of *Salmonella enterica* Serovar Typhi Is Required for Bacterial Entry
into Epithelial Cells. Infect Immun. 2006 Mar;74(3):3. doi:10.1128/IAI.74.3.1555-
1564.2006

22. De Jong HK, Parry CM, Van Der Poll T, Wiersinga WJ. Host–Pathogen Interaction in
Invasive Salmonellosis. Chitnis CE, editor. PLoS Pathog. 2012 Oct 4;8(10):e1002933.
doi:10.1371/journal.ppat.1002933

23. Wilson RP, Raffatellu M, Chessa D, Winter SE, Tükel Ç, Bäumler AJ. The Vi-capsule
prevents Toll-like receptor 4 recognition of Salmonella. Cell Microbiol. 2008 Apr;10(4):4.
doi:10.1111/j.1462-5822.2007.01090.x

24. Isibasi A, Ortiz V, Vargas M, Paniagua J, González C, Moreno J, et al. Protection against
Salmonella typhi infection in mice after immunization with outer membrane proteins isolated
from Salmonella typhi 9,12,d, Vi. Infect Immun. 1988 Nov;56(11):2953–9.
doi:10.1128/iai.56.11.2953-2959.1988 PubMed PMID: 2844676; PubMed Central PMCID:
PMC259676.

25. Crotty S, Aubert RD, Glidewell J, Ahmed R. Tracking human antigen-specific memory B
cells: a sensitive and generalized ELISPOT system. Journal of Immunological Methods.
2004 Mar;286(1–2):1–2. doi:10.1016/j.jim.2003.12.015

26. Deshpande NU, Jayakannan M. Biotin-Tagged Polysaccharide Vesicular Nanocarriers for
Receptor-Mediated Anticancer Drug Delivery in Cancer Cells. Biomacromolecules. 2018
Aug 13;19(8):8. doi:10.1021/acs.biomac.8b00833

27. Dj O, Mj D, Rh Q. A novel procedure for labeling immunoglobulins by conjugation to
oligosaccharide moieties. Immunology letters. 1984;8(5):5. doi:10.1016/0165-
2478(84)90008-7 PubMed PMID: 6394473.

28. Athavale A, Gaur A, Ahmed N, Subramaniam A, Dandotiya J, Raj S, et al. Receptor Binding
Domain-Specific B Cell Memory Responses Among Individuals Vaccinated Against SARS-
CoV-2. Vaccines. 2024 Dec 12;12(12):1396. doi:10.3390/vaccines12121396

29. Costa DM do N, Coêlho MRCD, Gouveia PA da C, Bezerra LA, Marques CDL, Duarte
ALBP, et al. Long-Term Persistence of Serum-Specific Anti-Chikungunya IgM Antibody -
A Case Series of Brazilian Patients. Rev Soc Bras Med Trop. 2021;54:e0855.
doi:10.1590/0037-8682-0855-2020 PubMed PMID: 33886823; PubMed Central PMCID:
PMC8047711.

30. Racine R, Winslow GM. IgM in microbial infections: Taken for granted? Immunology
Letters. 2009 Aug 15;125(2):2. doi:10.1016/j.imlet.2009.06.003

31. House D, Wain J, Ho VA, Diep TS, Chinh NT, Bay PV, et al. Serology of typhoid fever in
an area of endemicity and its relevance to diagnosis. J Clin Microbiol. 2001 Mar;39(3):3.
doi:10.1128/JCM.39.3.1002-1007.2001 PubMed PMID: 11230418; PubMed Central
PMCID: PMC87864.

32. Darton TC, Baker S, Randall A, Dongol S, Karkey A, Voysey M, et al. Identification of
Novel Serodiagnostic Signatures of Typhoid Fever Using a Salmonella Proteome Array.
Front Microbiol. 2017;8:1794. doi:10.3389/fmicb.2017.01794 PubMed PMID: 28970824;
PubMed Central PMCID: PMC5609549.

33. Mond JJ, Vos Q, Lees A, Snapper CM. T cell independent antigens. Curr Opin Immunol.
1995 Jun;7(3):349–54. doi:10.1016/0952-7915(95)80109-x PubMed PMID: 7546399.

34. El Shikh MEM, El Sayed RM, Szakal AK, Tew JG. T-Independent Antibody Responses to
T-Dependent Antigens: A Novel Follicular Dendritic Cell-Dependent Activity12. The
Journal of Immunology. 2009 Mar 15;182(6):3482–91. doi:10.4049/jimmunol.0802317

35. Martin MD, Badovinac VP. Defining Memory CD8 T Cell. Front Immunol. 2018 Nov 20;9.
doi:10.3389/fimmu.2018.02692

36. Jin C, Hill J, Gunn BM, Yu WH, Dahora LC, Jones E, et al. Vi-specific serological
correlates of protection for typhoid fever. Journal of Experimental Medicine. 2021 Feb
1;218(2):2. doi:10.1084/jem.20201116

37. Petrov S, Taskov H, Murdjeva M. Guardians of immunity: NK cell-mediated defense in
COVID-19 and post-COVID scenarios. FM. 2024 Feb 29;66(1):1.
doi:10.3897/folmed.66.e113356

38. Lang B, Wang M, Zhang Z, Fu Y, Han X, Hu Q, et al. Inhibitory receptor CD47 binding to
plasma TSP1 suppresses NK-cell IFN-γ production via activating the JAK/STAT3 pathway
during HIV infection. J Transl Med. 2023 Nov 30;21(1):1. doi:10.1186/s12967-023-04667-6
PubMed PMID: 38037074; PubMed Central PMCID: PMC10688093.

39. Lanier LL. Up on the tightrope: natural killer cell activation and inhibition. Nat Immunol.
2008 May;9(5):5. doi:10.1038/ni1581 PubMed PMID: 18425106; PubMed Central PMCID:
PMC2669298.

40. Bharmoria A. Typhoid Fever as a Challenge for Developing Countries and Elusive
Diagnostic Approaches Available for the Enteric Fever. IJVR. 2017 Jun 2;2(2):2.
doi:10.15226/2473-2176/2/2/00118

Figure legends

Fig. 1 IgA ELISA against Vi PS, LPS, H protein, HlyE and OMP

Fig. 2 IgM ELISA against Vi PS, LPS, H protein, HlyE and OMP

Fig. 3 IgG ELISA against Vi PS, LPS, H protein, HlyE and OMP; IgG Avidity against Vi PS

Fig. 4 Percentage of NK cell proliferation and activation markers after antibody-dependent
activation

Fig. 5 Distribution of B cell subsets within CD3□CD19□ B cells showing naïve (CD27□IgD□),
switched memory (CD27□IgD□), unswitched memory (CD27□IgD□) and double negative
(CD27□IgD□) populations.

Fig. 6 Frequency of classic memory B cells (CD27□CD38□) and transitional B cells
(CD27□CD24□CD38^low) among circulating lymphocytes.

Fig. 7 Frequency of HlyE-specific memory B cells and the proportion of IgG□ and IgM□ cells
within this population.

Fig. 8 Distribution of CD4□ T cell subsets (naïve, central memory, effector memory and
TEMRA) among typhoid-recovered and healthy participants.

Fig. 9 Distribution of CD8□ T cell subsets (naïve, central memory, effector memory and
TEMRA) among typhoid-recovered and healthy participants.

564
565

566   **Table 1:** Participant demographics

		Typhoid Recovered	**Healthy**
Sex	Male	15	27
	Female	18	20
Age (years)	Range	13-53	12-58
	Median	25	24
Months since infection	Range	3-10	-
	Median	3	-

569   **Table 2:** Molecular markers used for estimation of NK cell proliferation and activation

Sr. No.	Molecular marker	Fluorophore	Make and Catalogue number
1	Fixable viability dye	Violet Dye	Invitrogen #L34964
2	CD3	PerCP	BioLegend #344814
3	CD19	PE	BioLegend #302208
4	CD25	APC	BioLegend #302610
5	CD56	FITC	BioLegend #318304
6	CD69	BV510	BioLegend #310936
7	CD14	APC-Cy7	BioLegend #301820

571   **Table 3:** Molecular markers used for analyzing B cell and T cell phenotyping

Sr. No.	Molecular marker	Fluorophore	Catalogue number
1	Fixable viability dye	Violet Dye	Invitrogen #L34964
2	CD3	RB705	344814
3	CD19	PE	302208
4	CD21	RY703	551064
5	CD27	PE-Cy7	350526

6	CD24	APC	311118

Antigens	Vi polysaccharide	Lipopolysaccharide

7	CD38	BV650	356620
8	CD69	BV510	310936
9	CD138 (Syndecan-1)	BV785	356538
10	IgG Fc	BV711	410740
11	IgM	BV750	314566
12	IgD	APC-Cy7	348218
13	CD25	AF700	302622
14	CD11c	BV605	563929
15	Bcl-6	PE/Dazzle^{TM} 594	358510

Antibody class	IgA	IgG	IgM	IgA	IgG	IgM
Relative Antibody response among Typhoid-recovered individuals over the healthy participants	* p=0.0140 (p<0.05)	ns	* p=0.0107 (p<0.05)	* p=0.0227 (p<0.05)	ns	* p=0.0222 (p<0.05)

Antigens	H Protein			Hemolysin E			OMP		
Antibody class	IgA	IgG	IgM	IgA	IgG	IgM	IgA	IgG	IgM
Relative Antibody response among Typhoid-recovered individuals over the healthy participants	ns	ns	**** (p<0.0001)	*** p=0.0009 (p<0.001)	ns	** p=0.0022 (p<0.01)	ns	ns	** p=0.0025 (p<0.01)

Table 4: Antigen-wise seropositivity among individuals after 3 months of recovery from typhoid (ELISA)

Polysaccharide antigens

(i) Vi PS

(ii) LPS

Protein antigens

(iii) H protein

(iv) HlyE

(v) OMP

Polysaccharide antigens

(i) Vi PS

(ii) LPS

Protein antigens

(iii) H protein

(iv) HlyE

(v) OMP

(a) CD16

(b) CD69

(c) CD25

Naïve
$CD27^-IgD^+$

Switched
$CD27^+IgD^-$

Subsets of
memory cells
within B cells
$(CD3^-CD19^+)$

Unwitched
$CD27^+IgD^+$

Double Negative
$CD27^-IgD^-$

Figure 5

Classic Memory cells
$CD27^+CD38^-$

Transitional B cells (T1/T2)
$CD27^-CD24^+CD38^{low}$

Figure 6

Figure 7

CD4

Figure 8

CD8

Figure 9