Title:

Flexible foraging effort shapes Adélie penguin reproductive success across Antarctica and sea ice conditions

Authors:

Téo Barracho [1,2,3*], Gaël Bardon [1,3], Aymeric Houstin [1,3,4], Michaël Beaulieu [5], Pierrick Blanchard [6], Robin Cristofari [7], Lana Lenourry [8], Thierry Raclot [1], Clarisse Tabillon [3], Daniel P. Zitterbart [4,9], Nicolas Lecomte [2*†] and Céline Le Bohec [3,8*†]

[†] Co-last authors

[*] tbarracho@centrescientifique.mc

[*] nicolas.lecomte@umoncton.ca

[*] celine.le-bohec@cnrs.fr

Affiliations:

[1] Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[2] Canada Research Chair in Polar and Boreal Ecology and Centre d'Études Nordiques, Department of Biology, University of Moncton, Moncton, NB, Canada

[3] Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of Monaco

[4] Friedrich-Alexander-University Erlangen-Nürnberg, Department of Physics, Erlangen, Germany

[5] Wild Animal Initiative 5123 W 98th St, 1204 Minneapolis, MN 55437

[6] Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), Université de Toulouse, Toulouse INP, CNRS, IRD, Toulouse, France

[7] Institute of Biotechnology, University of Helsinki, Helsinki, Finland

[8] CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France

[9] Woods Hole Oceanographic Institution, Applied Ocean Physics and Engineering Department, Woods Hole, Massachusetts, USA

Abstract

Understanding species responses across a wide spectrum of environmental conditions is essential for predicting demographic responses to climate change. In Antarctica, changing sea ice dynamics threaten ice-dependent species. Behavioural adjustments could help mitigate these threats, but how they link to demographic rates, as well as potential thresholds, remains elusive. Long-term studies and circumpolar approaches could address this challenge. Here, we used a 15-year dataset (2010-2024) comprising over 23,000 foraging trips from Adélie penguins *(Pygoscelis adeliae)* in East Antarctica, to investigate how the species responds to variable sea ice conditions during reproduction via flexible foraging effort (trip duration). We identified a non-linear relationship between sea ice concentration (SIC) and foraging effort in the absence of landfast ice, a feature predicted to define the species' future habitat during chick-rearing. Minimal effort occurred around 10-20% SIC and increased more than two-fold in the presence of landfast ice, or of heavy pack-ice (>76% SIC). This relationship was sex-specific, with females showing greater environmental sensitivity. Through a circumpolar synthesis, we also showed that foraging effort during chick-guard predicted reproductive success across ten populations spread across the Adélie penguin's range (marginal $R^2 = 0.74$). Threshold analysis revealed that reproductive success remains comparatively high until chick-guard foraging trip duration exceeds ~29 hours (approximately three times minimum values), beyond which demographic performance declines at a stronger rate. Our findings provide a comprehensive assessment of Adélie penguin's responses across a large environmental gradient, revealing that behavioural flexibility can buffer moderate changes but has clear limits. As it mediates the relationship between environmental variability and reproductive success, our results also establish trip duration as a valuable early warning indicator for population responses to environmental change.

Keywords:

Antarctica | environmental variation | circumpolar synthesis | sea ice | Adélie penguin | foraging effort | reproductive success

Introduction

Global change is exerting an increasing pressure on ecosystems, challenging their resilience, and threatening species with extinction (Urban, 2015; Wiens & Zelinka, 2024). As changes in species abundance stem from changes in vital rates such as survival and reproduction (Sæther et al., 2016), adopting a demographic framework is a powerful way towards understanding and forecasting the impacts of global change on biodiversity (Capdevila et al., 2020; Jenouvrier, 2013; Paniw et al., 2019, 2023). For example, widespread declines in bird offspring production (e.g., Halupka et al., 2023) may foreshadow future population declines (Cerini et al., 2023).

Nevertheless, such declines are not homogeneous, and some species are able to withstand, or even benefit from, the modifications in their environment associated with global change (Finn et al., 2023; Rosenberg et al., 2019). While compensation among vital rates (e.g., increases in one vital rate proportional to decreases in another (Iles et al., 2019; Reed et al., 2013) may explain part of this resilience, a major and rapid mechanism through which species can respond to changing environments is by modifying their behaviour (Beever et al., 2017; Milles et al., 2023; Muthukrishnan et al., 2025; Wong & Candolin, 2015). Behavioural modifications observed at the population-level in response to changing environments (adjustments) can arise through a variety of processes. These involve reversible changes within the same individuals (contextual plasticity), or a turnover of individuals with similar (developmental plasticity) or distinct (genetic evolution) genotypes (Andrade et al., 2025; Charmantier & Gienapp, 2014; Piersma & Drent, 2003; Stager et al., 2024). Regardless of the origins of behavioural adjustments (here defined at the population level), species or population are often able to adjust key behaviours such as breeding phenology or activity budgets, which can help them mitigate or even entirely escape the demographic consequences of mismatched food resources (Both et al., 2006; Charmantier et al., 2008; Møller et al., 2008; Visser & Gienapp, 2019), unpredictable food shortages (Burger & Piatt, 1990; Grémillet et al., 2012; Harding et al., 2007) or rising temperatures (Kearney et al., 2009; Mason et al., 2014).

While behaviour is considered an early response to stressors (Cerini et al., 2023), the limitations and potential thresholds of behavioural adjustments in allowing species to cope with changing environments are poorly known (Beever et al., 2017). Specifically, examples charting the efficacy of behavioural adjustments for buffering demography declines with increasing environmental constraint are few (e.g., Fuentes et al., 2023; Wanless et al., 2023). Shorter-term studies can provide a binary response to the efficacy of behavioural adjustments (e.g., sufficient: Dehnhard et al., 2016; Grémillet et al., 2012; Hamer et al., 2007; Senner et al., 2015 ; insufficient: Fayet et al., 2017; Horswill et al., 2017). However, a

limited cover of environmental conditions will hide the full suite of possible responses and flexibility (Grémillet & Charmantier, 2010). In this context, long term studies and cross-populations syntheses (e.g., space for time substitutions, Şen et al., 2025) can help resolve this challenge (Orgeret et al., 2022; Viblanc & Muller-Landau, 2025).

Antarctica presents an ideal natural laboratory for exploring these questions, with rising temperatures, shifts in wind and precipitation patterns, changing sea ice dynamics, and fisheries development all combining to affect food webs across the continent (e.g., Chown et al., 2022; Convey & Peck, 2019; Purich & Doddridge, 2023; Raphael et al., 2025; Rintoul et al., 2018; Trathan et al., 2025; Vignon et al., 2021; Watters et al., 2020). Many Antarctic species are long lived and as such are likely to rely heavily on plasticity (physiological or behavioural) for responding to these changes (Convey & Peck, 2019; Peck, 2018). Among them, the Adélie penguin (*Pygoscelis adeliae*) is the focus of several long-term research programs around the continent (Ainley, 2002; Cimino et al., 2023; Clarke et al., 2003; Hinke et al., 2014; Olmastroni et al., 2000; Watanabe et al., 2020) making this species suitable for continent-wide approaches.

Adélie penguins are reliant on sea ice (Ainley, 2002; Emmerson & Southwell, 2008; Iles et al., 2020), a dominant feature of the Southern Ocean (e.g., Fraser et al., 2023; Fricker et al., 2025). Sea ice can impact Adélie penguin food availability in two ways: it controls the population dynamics (abundance) of its main prey species (i.e. Antarctic and ice krill, Antarctic silverfish, *Euphausia superba & E. crystallorophias, Pleuragramma antarctica*, respectively, (Ainley, 2002; Corso et al., 2022; Kawaguchi et al., 2023), and can also constrain penguins' accessibility to these prey. While non-breeding Adélie penguins can escape local food constraints by travelling thousands of kilometers (Oosthuizen et al., 2022; Thiebot et al., 2019), they are central place foragers bound to their breeding site during the breeding season (Orians & Pearson, 1979).

During the chick-guard stage of the breeding cycle, foraging penguins are particularly exposed to local changes in food availability, with the regular provisioning of young and thermally dependent chicks requiring them to forage over smaller areas compared to other breeding and non-breeding stages (Takahashi et al., 2018; Thiebot et al., 2019; Widmann et al., 2015). The availability of food during chick-guard can be affected by sea ice at different timescales, the main of which appears to differ across Antarctic regions. In areas where sea ice persists near colonies during chick-guard, such as the Ross Sea or East Antarctica, concurrent sea ice conditions are considered a reliable predictor of local food availability (Ainley et al., 2006; Ballard et al., 2010; Kokubun et al., 2021; Le Guen et al., 2018;

Watanabe et al., 2020). However, in areas where sea ice seldom persists after incubation (e.g., the western Antarctic Peninsula, (Oosthuizen et al., 2022; Stammerjohn et al., 2008) variations in chick-guard food availability are thought to be mostly driven by sea ice conditions prior to chick-rearing, such as in the previous winter or in early spring (Cimino et al., 2023; Gallagher et al., 2023).

Regardless of its drivers, chick-guarding penguins can respond to variable food availability by adjusting their foraging effort, for example by performing longer trips when food availability is reduced (Ballance et al., 2009; Nicol et al., 2008; Watanabe et al., 2020). While such adjustments may lead to higher energy expenditure (Ballance et al., 2009), they could allow foraging parents to find sufficient food to sustain their chicks. Nevertheless, the extent to which foraging adjustments allow penguins to buffer reproductive success against variable food availability is poorly known (Emmerson et al., 2015). For populations where chick-guard food availability is driven by concurrent sea ice conditions, the type of ice encountered by penguins likely affects this outcome (Emmerson et al., 2015; Emmerson & Southwell, 2008). For instance, the persistence of landfast ice (consolidated sea ice anchored to the coastline) in front of breeding colonies during chick-guard leads to much longer foraging trips because penguins have to travel by walking rather than swimming (Lescroël et al., 2014; Watanabe et al., 2020). In such cases, reproductive success is often severely reduced (Emmerson & Southwell, 2008; Ropert-Coudert et al., 2018). In the absence of landfast ice, foraging penguins can still face varying concentrations of pack ice, ranging from open water (0% Sea Ice Concentration, SIC) to heavy pack ice (up to 80% SIC). In this scenario, the foraging response of penguins and the consequences on reproductive success are not fully resolved, although foraging effort was found to be minimal at intermediate 10-20% SIC when measured over a narrow range of SIC (0-30% SIC, Ballard et al., 2010; Le Guen et al., 2018). Because most Adélie penguin populations typically have direct access to open water during chick-rearing (i.e. with an absence of landfast ice; Ainley, 2002; Santora et al., 2020), and because landfast ice is expected to recede further under climate change (Fraser et al., 2023), understanding penguin responses to the variable pack ice conditions they predominantly encounter is crucial for projecting their demographic responses to future environmental changes.

Here, we combine a unique 15-year-long time series of Adélie penguin's foraging effort (trip duration) and reproductive success from Adélie Land, Antarctica, with an additional circumpolar analysis of previously published data spanning the species range to address two main questions. First, we ask how Adélie penguins at our Adélie Land study site adjust their foraging effort to variable food availability, as proxied by sea ice conditions in their chick-guard foraging area. Specifically, we investigate how penguins respond to a wide range of sea ice concentrations (0-99%) while contrasting both ice types (landfast vs. pack ice) to match the expected icescape the species will face in the coming decades.

Second, we explore the consequences of these foraging effort adjustments on reproductive success. We predict that increased foraging effort, as reflected by longer foraging trips, will lead to lower reproductive success (Emmerson et al., 2015; Watanabe et al., 2020), even in the absence of landfast ice. By scaling-up the analysis to include other colonies, we aim at quantifying the relationship between foraging effort and reproductive success across the species distribution. Such data integration over a wide range of conditions allows us to understand the relevance of foraging effort as a fundamental mechanism linking environmental variability to reproductive success, providing a robust baseline for projecting the species responses to changing sea ice dynamics.

Material and methods

Study site and monitoring design

The Adélie penguin's breeding cycle spans the austral summer (see Ainley (2002) for a general description). In Adélie Land, following arrival at the breeding colonies (October) and the laying of two eggs (November), Adélie penguin partners alternate incubation shifts and long foraging trips (5-20 days), before switching to shorter trips upon hatching (guard stage: December-January). Broods (1 or 2 chicks) grow rapidly, and as chicks acquire self-thermoregulation abilities, parents then start to forage simultaneously to maximize food deliveries (crèche stage: January-February). Depending on the region, chicks finally fledge in January-February, with fledging mass being an important predictor of their post fledging survival (Barracho et al., 2025).

We carried out fieldwork on Île des Pétrels (66°40′S, 140°01′E), Pointe Géologie archipelago, Adélie Land, Antarctica, adjacent to the Dumont d'Urville research station. This archipelago has hosted 30,000-50,000 breeding pairs of Adélie penguins annually for the last two decades (Barbraud et al., 2020), making it a medium-sized colony for this species (Ainley, 2002; Santora et al., 2020). From the 2006-2007 to 2023-2024 breeding seasons, we established a comprehensive monitoring system for a breeding patch comprising approximately 250 annual breeding pairs and situated within a natural canyon (hereafter study colony, Barracho et al., 2025).

Starting in 2006-2007 (breeding seasons will be referred to by the chick-fledging year, 2007 in this case), we implanted Radio Frequency IDentification (RFID) tags in both chicks and breeding adults (n = 3,248 and n = 313 as of 2024, respectively). From the 2010 breeding season onwards, the study colony was enclosed with directional RFID gateways, strategically positioned to capture all colony entries and exits.

This setup enables the precise tracking of individual colony attendance patterns over their entire breeding cycle (see Barracho et al., 2025 and supplementary material F in Bardon et al., 2023 for a description of the system, as well as similar systems in Adélie (K. Kerry et al., 1993; Lescroël et al., 2014; Olmastroni et al., 2000) and other penguins species (Afanasyev et al., 2015; Bardon et al., 2023; Dehnhard et al., 2013; Robinson et al., 2005)). The RFID record produced by this system was used to determine individual trip durations between the 2010 and 2024 breeding seasons (see trip duration section below).

Each season, this electronic monitoring was complemented by daily observations of a subset of nests spread over the study colony (n = 0 to 60 depending on the season), starting in 2012. We identified breeding partners on their nest using either hand-held RFID readers or repeated visual observations coupled with the automatic RFID system. We monitored both the breeding outcome and phenology (e.g., hatching and crèching dates) of each of these nests throughout the breeding season. Because nests were monitored from a distance using binoculars to minimize disturbance, the hatching date could not be measured precisely for all nests. Nests for which hatching dates could not be measured with a precision of ± 4 days (i.e. more than 4 days between the last observation of an egg and the first observation of a chick) were thus excluded from the analyses. Crèching dates were observed with a 1-day precision. For a given nest, hatching dates were taken as the hatching of the first egg, and crèching dates as the first time the brood was seen unattended by parents. Finally, we conducted weekly photo counts of both adults and chicks present in the study colony from the 2011 season onwards. These counts were used to determine the reproductive success of the study colony (chicks fledged per breeding pair), which closely represented the broader archipelago population (r = 0.99, 2011-2017, Barracho et al., 2025, see also (Méheust et al., 2024)).

Focus on the chick-guard stage

The chick-guard stage of the Adélie penguin starts at hatching and ends when chicks are left unattended at the colony while both parents forage simultaneously (crèching stage). We decided to focus our analyses on the chick-guard stage of the Adélie penguin breeding cycle for three reasons. First, foraging parents are most constrained in time and space during chick-guard. Compared to incubation and crèching, they forage over a smaller area (Widmann et al., 2015) and provision chicks more frequently (Ainley, 2002; Ballard et al., 2010; Riaz et al., 2020). These constraints on parental effort during chick-guard are further illustrated by the body mass loss incurred by breeders, whereas this pattern is typically reversed during incubation and crèching (Clarke et al., 2006; Emmerson et al., 2019). Second, chicks have a higher probability of mortality during chick-guard compared to crèche (Davis & McCaffrey,

1986; Jennings et al., 2023), making this stage critical for chick survival and reproductive success. Finally, focusing on chick-guard facilitates comparisons of foraging trips duration with other study sites, as data is more abundant for chick-guard than for crèching, in part because of the practical difficulties associated with retrieving electronic data loggers during crèching (another common methodology for acquiring trip duration data, e.g., Kim et al., 2024; Machado-Gaye et al., 2025; Park et al., 2022).

In Adélie penguins, the phenology of reproductive stages varies among years and populations (Ainley, 2002; Youngflesh et al., 2017). In Adélie Land, peak hatching and crèching typically occur in late December and mid-to-late January, respectively (Beaulieu et al., 2009). Here, we defined the chick-guard stage on an annual basis according to season-specific median hatching and crèching dates. Nest-level hatching dates were obtained from the RFID record by adding 31 days (i.e. the duration of incubation minus the average two days taken by the female to leave the colony after laying (Ainley, 2002) to the departure date of females for their first incubation foraging trip (average annual n = 77 ± 41 nests). Crèching dates were obtained from both daily observations at the colony and the RFID record, with crèching defined as the first time two known partners were away from the colony (average annual n = 25 ± 12 nests). Both hatching and crèching dates derived from the RFID record were validated using the visual observation data for the subset of nests for which both were available. As the field observations of crèching dates started in 2013, crèching phenology was unavailable in 2010, 2011, and 2012. For those years, we defined the median crèching date as the median hatching date plus the average duration of brooding in other years (25 days).

Trip duration, sea ice, and individual parameters

Individual trip durations were extracted from the RFID record and defined as the duration (hours, h) between colony departure and return. We followed the data cleansing protocol of Bardon et al. (2023) to process missed detections (e.g., individuals detected with only one of the two directional antennas). Foraging trip data were also filtered to exclude trips shorter than 6 h (likely non-foraging, Ballard et al., 2010; Lescroël et al., 2021) and trips exceeding 5 days, as they likely reflect late incubation trips of late breeders, or unresolvable missing RFID detections. We ensured that we focused on trips from individuals engaged in chick-guarding activities by only using data from successful breeders, or from failed breeders prior to their breeding failure. To identify successful breeders, we used the *RFIDeep* workflow (Bardon et al., 2023), which identifies breeding success from individual RFID records with high accuracy (>95%). For failed breeders, we retained only individuals for which the daily observation protocol (see above) allowed us to determine the date of breeding failure with a precision of ± 2 days or less.

We considered two sea ice variables to investigate how penguin foraging effort responded to sea ice variability and associated food availability. First, we assessed the binary presence/absence of landfast ice (sea ice anchored to the coastline, also called landfast ice) in front of the breeding colony during the chick-guard stage. This was achieved by visual inspection of MODIS and VIIRS satellite imagery using NASA Worldview (https://worldview.earthdata.nasa.gov). Second, we measured the sea ice concentration (SIC) in the chick-guard foraging area for each trip. A single SIC value per trip was calculated by averaging daily concentrations from one day pre-departure to one day post-return. Adélie penguins' foraging area during chick-guard at our study site was defined using previously published tracking data (Michelot et al., 2021; Ropert-Coudert et al., 2018; Widmann et al., 2015) and ranged from 65.5°S to 66.7°S and from 138°E to 141°E (Fig. S1). This spatial window is also consistent with Barreau et al. (2019) findings that January SIC in that area was related to reproductive success for the Pointe Géologie archipelago Adélie penguin population. Trip specific SIC was calculated using daily SIC data from Advanced Microwave Scanning Radiometer AMSR-E and AMSR-2 (6.25×6.25 km grid datasets), downloaded from the University of Bremen website (https://data.seaice.uni-bremen.de/amsr2/asi_daygrid_swath/) for all breeding seasons except 2012. AMSR data was unavailable for 2012 due to satellite replacement, and we instead used data from the National Snow and Ice Data Center (https://nsidc.org/data/g02202/versions/4, 25×25 km grid, Meier et al., 2021 in that season, as SIC from both datasets were strongly correlated for the 2009-2024 period ($R^2 = 0.96$, $\beta = 0.96 \pm 0.005$). The possible range for sea ice concentration values is between 0% (open water), and 100% (total ice cover).

In addition to landfast ice presence and SIC variability, we also included sex in our trip duration models, since female Adélie penguins usually forage for longer (Beaulieu, Dervaux, et al., 2010; Clarke et al., 1998; Dugger et al., 2006; Emmerson et al., 2015), travel farther offshore (Widmann et al., 2015), and target different food sources (Beaulieu, Dervaux, et al., 2010; Michelot, 2020) compared to males. Sex assignment was done by analyzing sex-specific colony attendance patterns using RFIDeep (Bardon et al., 2023). Sex was determined by calculating the average percentage of support across multiple breeding seasons. Individuals for which the average percentage of support was below 75% (i.e. individuals with too little stereotyped colony-attendance patterns to be sexed confidently) were excluded from subsequent analyses (mean support for sex-assignment in the final dataset = 89%).

Mean trip duration as an indicator of reproductive success

While our first hypothesis dealt with the effects of sea ice on foraging effort at the trip-scale, our second hypothesis focused on linking colony-scale measures of foraging effort to reproductive success. We

tested our prediction that increased trip duration during chick-guard leads to lower reproductive success

at two spatial scales. First, we averaged trip durations for each year at our Adélie Land study site and

investigated the relationship between mean trip duration and reproductive success using both our entire

dataset (n = 15 years) and a dataset only containing years when landfast ice was absent from the front

of the colony. In both cases, we modelled reproductive success as a function of mean trip duration

calculated over either all individuals, males, or females, and compared the three models using Akaike

Information Criterion (AIC). As reproductive success data for our study colony was unavailable in 2010,

we extracted it at the population level from Barreau et al. (2019) using PlotDigitizer

(https://plotdigitizer.com/).

Second, we surveyed the literature using Google Scholar with the keywords 'Adélie penguin' and 'trip

duration'. For each identified study with trip duration data, we then looked for concurrent data on

reproductive success, either from the same study or from a different study (see supplementary text). We

extracted the reported trip duration data only when concurrent reproductive success data was available

(i.e. same colony and same season). As there were slight differences among the 14 identified studies in

how trip duration data was reported, we used our own data to ensure comparability of methods (see

supplementary text for details).

Statistical analyses

Trip duration and sea ice

We used Generalized Linear Mixed Models (GLMM) and Generalized Additive Mixed Models

(GAMM) as implemented in the {*gamm*} function of R package {*mgcv*} (Wood, 2011) to investigate

linear and non-linear relationships between trip duration and SIC. We fitted linear and quadratic GLMM

as well as GAMMs, and compared these models to assess the support for non-linearity in the response

of trip duration to SIC, using an AIC-based model selection framework (Burnham & Anderson, 2002).

As a conservative way to avoid model overfitting and to optimize biological interpretation, the

maximum number of basis functions used for smoothing the relationship between trip duration and SIC

in GAMMs was set to $k = 4$. In all models, we used a Gamma error distribution with a log link function

and included individual ID as a random effect to account for individual-level repeated measurements.

Additional fixed effects of sex, as well as the interaction of sex with SIC were then included, and their

support was investigated using AIC. For GAMMs, we employed a tensor product smoother (t2, (Wood

et al., 2013) to allow sex-specific smoothing of the relationship between trip duration and SIC (i.e.

interaction, (Pedersen et al., 2019). Data from each landfast ice scenario (present/absent) was analyzed

separately because there were large differences in both sample size and range of SIC values covered between scenarios.

Reproductive success and mean trip duration: study site

To model the relationship between colony-scale mean trip duration and reproductive success (number of chicks fledged per pair) for our study site, we first averaged all trip duration values by year, without accounting for repeated measurements at the individual level (i.e. a slightly different approach to that of Emmerson et al., 2015 but similar to Ainley et al., 2004). We then fitted Generalized Linear Models (GLM) with a log link function as implemented in package {*glmmTMB*} (Brooks et al., 2017) with reproductive success and mean trip duration as response and explanatory variables, respectively. We fitted three models, first using annual mean trip duration for all individuals, then for males and females separately. Reproductive success was modeled using a Tweedie distribution to accommodate the presence of 0 (2 values out of 15) and the lower bound at 0 (Foster & Bravington, 2013). We weighted observations using the inverse standard error of each annual trip duration mean, thus giving greater weight to more precise estimates of mean trip duration (i.e. measured over larger sample sizes, Table S1). We assessed the validity of the unweighted models by checking residuals for dispersion, uniformity, and outliers using package {*DHARMa*} (Hartig, 2024). We used a variance-based pseudo-R^2 (squared correlation coefficient between predicted and observed data) to provide an indication of the predictive performance of our models.

Reproductive success and mean trip duration: across colonies

We used the same approach to upscale the analysis using data from 9 other colonies. We fit Tweedie GLMM with a colony random effect to account for possible differences in mean reproductive success among colonies. We compared a linear and a quadratic model using AIC to test for non-linear effect of mean trip duration on reproductive success. Because standard errors for each annual mean trip duration were only available for <30% of the data entries, we weighted observations using a log-transformation of the number of trips used to calculate each annual mean (supplementary text). As mean trip duration estimates made over a larger number of trips are likely to be more precise (although asymptotically, (Southwell et al., 2006), this allowed to put more weight on more precise estimates. Model fit was again assessed using the package {*DHARMa*}. Because initial checks revealed significant zero-inflation, we incorporated a zero-inflation component dependent on the explanatory variable in the Tweedie GLMMs. We used package {*performance*} to compute conditional and marginal R^2 (Lüdecke et al., 2021; Nakagawa et al., 2017).

All statistical analyses were performed using R version 4.4.2 (*R Core Team*, 2024). Mean values are presented ± SE unless specified otherwise.

Results

Our 15-year dataset (2010-2024) of foraging trip duration during the chick-guard period (inter-annual average: Dec. 23rd to Jan. 19th) encompassed 23,459 trips from 565 individual penguins that performed a total of 1,779 breeding cycles. On average, each individual was included in our dataset in 3 different breeding seasons (range: 1-12). The annual number of trips averaged 782 trips per year-sex combination (range: 10-3,013, Table S1). Landfast ice persisted in front of the colony through the chick-guard stage in five years (2012, 2014, 2015, 2017, 2018), when minimal distances to open water ranged from 4 km in 2012 to 60 km in 2017. For these five years, foraging penguins faced elevated sea ice concentrations (SIC) ranging from 43 to 99% (mean = 58 ± 0.33%; Fig. 1). For all the other years, landfast ice was absent in front of the colony during the chick-guard stage and penguins experienced a lower average (20 ± 0.14%) but a wider range (0-84%) of SIC, which even included near-total open water conditions (Fig. 1).

Trip duration response to the sea ice scape

In line with this sea ice variability, foraging trip duration during the chick-guard period also varied considerably (mean = 23.4 ± 0.08 h). The presence of landfast ice in front of the breeding colony was the main driver of foraging trip duration. Trips were twice as long in years of landfast ice presence (mean = 43.9 ± 0.46, n = 1,621 trips) compared to when landfast ice was absent (mean = 21.9 ± 0.06 h, n = 21,838 trips, Welch's t-test p < 0.001). Over the range of SIC values shared by both landfast ice presence/absence scenarios (i.e. 43-84%, Fig. 2), trips were, on average, 1.4 times longer when landfast ice was present than when it was absent (Welch's t-test p < 0.001). In the absence of landfast ice, we found strong support for a non-linear effect of SIC on trip duration (GLMM vs. GAMM ΔAIC = 450; Table S2; Fig. 2), with the model-predicted 10% shortest trips (range: 19-20 h) occurring within 8-18% SIC (Fig. 2). Foraging trip duration lengthened rapidly between 20 and 40% SIC, but increased more slowly at ca. 70-80% SIC, with the model-predicted 10% longest trip duration (range: 29-37 h) occurring above 76% SIC. Comparison of raw data at optimal (8-18%) versus high SIC (>76%) indicates that this represents an average 2.5-fold increase in trip duration. At the other end of the SIC range (<8% SIC), we also observed an increase in trip duration, but of much lower magnitude (1.04-fold increase compared to optimal 8-18% SIC; Fig. 2). Similar to the landfast ice absent scenario, we found that foraging trip duration lengthened with increasing SIC when landfast ice was present.

The relationship between trip duration and SIC is mediated by sex

Chick-guard foraging trip durations differed between male and female Adélie penguins in both landfast ice scenarios (Table S2, Fig. 1B). Overall, females exhibited foraging trips almost 2 hours longer on average compared to males (females: mean = 24.4 ± 0.11 h, n = 12,135 trips *vs.* males: mean = 22.4 ± 0.10 h, n = 11,324 trips, Welch's t-test $p < 0.001$). Regardless of the landfast ice scenario, sex-differences were exacerbated under increasing SIC, as shown by the overall relationship between SIC and the annual male-female trip duration difference (LM, $\beta = 0.22 \pm 0.03$, $p < 0.001$, $R^2 = 0.79$; Fig. S2). Specifically, while the difference in trip duration between sexes was under 2 hours below 20% SIC, it increased beyond 14 hours above 70% SIC (Fig. S2). This relationship was maintained when landfast ice was absent (LM, $\beta = 0.15 \pm 0.04$, $p = 0.002$, $R^2 = 0.70$; Fig. S2).

Consequences on reproduction

Our 15-year study encompassed substantial interannual variation in reproductive success, ranging from 0 (2014, 2017) to a maximum of 1.29 chicks/pair (2022), with an overall mean of 0.66 ± 0.43 chicks/pair. Reproductive success was negatively related to mean SIC in the chick-guard foraging area (Tweedie GLM: $\beta = -0.020 \pm 0.005$, $p < 0.001$, pseudo-$R^2 = 0.58$, Fig. 3A) and was lower in years when landfast ice was present in front of the colony (Welch's t-test, $p < 0.001$, Fig. 1C). We found that longer foraging trips during chick-guard were associated with a linear reduction in reproductive success, with a model-predicted 4.3% decrease in reproductive success every 1-h increase in mean trip duration (Tweedie GLM: $\beta = -0.044 \pm 0.006$, $p < 0.001$, pseudo-$R^2 = 0.73$, Fig. 3B). This relationship was maintained when considering only years without landfast ice (Tweedie GLM: $\beta = -0.035 \pm 0.007$, $p < 0.001$, pseudo-$R^2 = 0.48$). Overall, the mean trip duration of females was a better predictor of reproductive success (pseudo-$R^2 = 0.78$) than that of males (pseudo-$R^2 = 0.57$), including in years when landfast ice was absent (Table S3).

The circumpolar analysis including data from 9 other Adélie penguin populations spread across the species range (see supplementary text) yielded an additional 49 concurrent measures of trip duration and reproductive success. Over the total 64 observations, covering East Antarctica, the Ross Sea, and the Western Antarctic Peninsula (Fig. 4), reproductive success averaged 0.88 ± 0.37 chicks/pair, while mean trip duration averaged 29.7 h (range 9.6-72.2 h). Similar to our study site, we found a significant relationship between reproductive success and trip duration (Table S4, $p < 0.001$, marginal $R^2 = 0.74$), but with non-linear effects of trip duration on reproductive success (linear vs. quadratic GLMM ΔAIC = 33, Table S4). Model-predicted reproductive success remained well above average (>1 chicks/pair) below approximately 30 h of trip duration, before declining more sharply (Fig. 4). To investigate this

potential threshold, we carried-out a piecewise regression analysis *a posteriori* and estimated the breakpoint at 29.2 h (bootstrapped 95% confidence interval: 17-40 h, Fig. S3).

Discussion

Behavioural adjustments to fluctuating environments are likely to be a central mechanism through which species can respond to climate change and other human pressures (Van Buskirk, 2012; Wong & Candolin, 2015). As such, capturing these behavioural adjustments over their largest possible extent and understanding their effectiveness for buffering demographic rates against environmental variability is crucial for identifying potential perturbation thresholds (e.g., Cury et al., 2011) and species tipping points (Beltran et al., 2023; Gil et al., 2020), as well as for designing reactive conservation policies (Beever et al., 2017; Cerini et al., 2023). Here, we showed how Adélie penguins responded to an unprecedentedly wide range of sea ice variability through flexible foraging effort, and identified a non-linear linkage between foraging effort and reproductive success across the species pan-Antarctic range. Our results provide a powerful template to assess Adélie penguins' response to anticipated changes in Antarctica.

The sea ice scape constrains foraging effort

The strong, positive relationship between landfast ice presence (sea ice anchored to the coastline) and trip duration found in our 15-year study corroborates previous findings (Ballard et al., 2010; Emmerson et al., 2015; Ropert-Coudert et al., 2018), confirming that the need to walk over ice rather than swim significantly increases foraging effort (Watanabe et al., 2020). Importantly, a remarkable contribution of our long term study is the characterization of the species response over a wide spectrum of sea ice concentration (SIC) in the absence of landfast ice, a condition that predominates for most Adélie penguin populations (Ainley, 2002; Santora et al., 2020) and is likely to become increasingly common as landfast ice recedes over the coming century (Fraser et al., 2023).

The non-linear relationship we observed between foraging effort and SIC (Fig. 2) aligns with previous studies in the Ross Sea (Ballard et al., 2010; Lescroël et al., 2014) and East Antarctica (Le Guen et al., 2018) showing optimal foraging at intermediate sea ice concentrations (8-18% SIC). However, by extending the range of SIC covered from 0-30% (Ballard et al., 2010) to 0-84%, our study revealed that foraging effort increases substantially at high SIC (>76%), even in the absence of landfast ice. Although such high SIC values are rarely attained in the absence of landfast ice (Fig. 2), dense pack-ice still

presents a significant challenge to foraging penguins, with trip duration increased up to 2.5-fold compared to optimal conditions. This value is comparable with the average 2.2-fold increase in trip duration between optimal conditions (8-18%SIC, no landfast ice) and landfast ice conditions.

Interestingly, male and female Adélie penguins differed in their response to challenging sea ice conditions. While females consistently undertook longer trips than males, matching previous observations locally (Beaulieu, Dervaux, et al., 2010; Widmann et al., 2015) and at other colonies around Antarctica (Clarke et al., 1998; Emmerson et al., 2015; Machado-Gaye et al., 2025), we showed that females experienced a steeper increase in trip duration with increasing SIC compared to males (Fig. S2). Such differences could be related to sex-specific diets, with males and females preferentially targeting fish and krill, respectively (Beaulieu, Dervaux, et al., 2010; Clarke et al., 1998), or to differences in body condition at the time of chick-guard, with females in poorer condition more prone to engage in longer self-feeding trips. Sex-specific sensitivity to foraging conditions could also signal the potential for sex specific threats (e.g., Gianuca et al., 2019; Schultz et al., 2021), although no difference in survival were found between male and female breeders in the Ross Sea (Morandini et al., 2024).

How does sea ice modulates foraging effort?

The increased effort at both very low (0% SIC) and high (>50% SIC) concentrations likely reflects different limitations on foraging efficiency. In open water (0% SIC), Adélie penguins may struggle to locate and exploit prey patches efficiently (Kokubun et al., 2021; Riaz et al., 2023). As Antarctic krill and silverfish aggregate near sea ice edges (Brierley et al., 2002; Corso et al., 2022), the complete absence of ice could remove these aggregating structures. The absence of sea ice during summer may also directly reduce the local abundance of prey species associated with sea ice year-round (e.g., *E. crystallorophias*), although the abundance of both *E. crystallorophias and E. superba* has been more convincingly linked to winter ice cover rather than summer SIC (Kawaguchi et al., 2023; Merkel et al., 2023; Ryabov et al., 2023). At the opposite end of the SIC spectrum, high ice concentrations physically impede access to prey, thereby reducing food accessibility through increased travelling costs (Emmerson et al., 2015; Watanabe et al., 2020). The optimal intermediate SIC (10-20%) therefore likely provides sufficient ice-edge habitat for prey aggregation, while maintaining adequate open water access for efficient travel and foraging (Le Guen et al., 2018).

Foraging within the future sea ice scape

Another key result is that, over the course of our study, very low SIC (<8%) did not significantly increase Adélie penguin foraging effort, which remained within 10% of their lowest observed effort (Fig. 2). Together with even shorter trip durations sometimes observed in the summer-ice-free western Antarctic peninsula (i.e. around 10 h, Machado-Gaye et al., 2025; Watters et al., 2020), this suggest that Adélie penguins are able to obtain sufficient food when SIC is severely reduced within their chick-rearing foraging area. In the context of ongoing and projected sea ice loss (Fraser et al., 2023; Purich & Doddridge, 2023; Rackow et al., 2022; Raphael et al., 2025), this is a crucial finding for understanding and anticipating the species' response under expected environmental shifts. Nonetheless, expected sea ice declines are not confined to the summer season, and will also likely affect winter ice dynamics. Because winter sea ice is a critical habitat and driver of krill abundance during summer (Kawaguchi et al., 2023; Meyer et al., 2017; Ryabov et al., 2023), the ability of Adélie penguins to mitigate lower ice concentrations during summer may not prove to be a sufficient response if waning winter sea ice further alters summer food abundance. Such lags could explain unexpected observations of lower reproductive success despite low summer SIC values at our study site (Barreau et al., 2019) or in the western Antarctic Peninsula (Fig. 4), but further concurrent data on food abundance, foraging effort and reproductive success in low SIC contexts will be key to test this hypothesis.

While counter-intuitive, predicted increase in air and water temperatures in Antarctica could also foster local increases in landfast ice and dense pack-ice persistence during the breeding season, which we both showed to significantly impact Adélie penguin foraging effort. Grounded icebergs can indeed act as anchor points for landfast ice, and glacier calvings are expected to increase under climate change-driven changes in temperatures, windstorms, and rain regimes (Fraser et al., 2023). While the location and duration of iceberg grounding events remain highly unpredictable (Fraser et al., 2023), such mechanisms could explain the rising landfast ice trends in certain regions, notably East Antarctica (Fraser et al., 2021).

Foraging effort is a reliable predictor of reproductive success

A major outcome of our study is the robust relationship between foraging trip duration during chick-guard and reproductive success, both at our study site (Fig. 3B) and across the species' range (Fig. 4). This relationship persisted even when considering only years without landfast ice, demonstrating that varying pack-ice concentrations alone can significantly impact reproductive outcomes. While substantial variability remains, the non-linear relationship we revealed over different Antarctic

regions—from the ice-heavy Ross Sea to the increasingly maritime West Antarctic Peninsula— highlights the limits of flexible foraging effort as a mechanism for coping with changes in food availability during a critical phase of the species' breeding cycle.

Why is chick-guard trip duration a good predictor of reproductive success?

Reproductive failure often peaks during early chick-rearing in Adélie penguins (Davis & McCaffrey, 1986; Jennings et al., 2023; Smiley & Emmerson, 2016) as well as other seabirds (e.g., Lewis et al., 2001; Mínguez & Oro, 2003; Ponchon et al., 2014) but see (Ollason & Dunnet, 1980). This makes the chick-guard period, and the ability of breeders to provision their chicks with food at a high rate by performing short trips, critical for reproductive success. While trip duration is constrained by foraging conditions encountered during chick-rearing, it also depends on the conditions encountered during incubation. Adélie penguins indeed use incubation foraging trips to regain the body condition that they subsequently lose during chick-guard, in order to maximize energy allocation towards the offspring (Clarke et al., 2006; Emmerson et al., 2019). Accordingly, failure to regain body conditions during incubation leads to longer foraging trips during chick-guard (Ballard et al., 2010; Ropert-Coudert et al., 2018). The integration of foraging conditions during both incubation and early chick-rearing thus likely makes chick-guard trip duration a reliable predictor of reproductive success.

The foraging strategy (Houston & McNamara, 2014; Schoener, 1971) employed by Adélie penguins is also important in explaining this relationship. Adélie penguins operate as time-minimizers during chick-guard, with foraging parents aiming at provisioning their offspring as quickly as possible, rather than maximizing food loads (Alonzo et al., 2003; Clarke et al., 2006). This is consistent with the limited stomach capacity of guard-stage chicks compared to maximal parental food loads (i.e. > 1 kg, (Emison, 1968) and with observations that chicks fed more often have faster growth rates and daily survival (Jennings et al., 2021, 2023). Importantly, the temporal constraints on foraging parents during chick-guard are not as strong during the incubation and crèching phases, explaining failed attempts at using trip duration during incubation as an index of food availability (Miller & Davis, 1993).

Trip duration as a tool for forecasting population reproductive performance

Seabirds are considered potent indicators of food supplies, with their behaviour more responsive to changes in food availability compared to demographic rates or abundance (Cairns, 1987; Lewis et al., 2006; Piatt et al., 2007). Consistently, behavioural changes sit at the forefront of the *timeline to collapse*, whereby the consequences of ramping environmental forcings should successively become observable on behaviour, morphological traits, demographic rates and ultimately abundance (Cerini et al., 2023). While this framework can improve forecasts of populations' fate under climate change, it requires well-defined baselines against which observed changes in (behavioural) traits can be contrasted, as well as an understanding of the linkage between behaviour and demographic rates (e.g., (Clements & Ozgul, 2016; Enquist et al., 2015; Sutherland & Norris, 2002; Wildermuth et al., 2013). Here, our results provide such a template for Adélie penguins, and the existence of similar data on foraging effort and reproductive success for other central-place foragers should prompt similar endeavours. Furthermore, as the foraging effort of females appears to be a better predictor of reproductive success than that of males (perhaps because female's longer trips can have more impact on the pair-level provisioning rate, see also (Emmerson et al., 2015), our results highlight the importance of considering and including sex-specific strategies in behaviour-based assessment of population responses (Nakazawa et al., 2023).

Although the relationship we observed between trip duration and reproductive success was consistent across conditions and space, extra variability remains (Fig. 3B, 4) and must be considered prior to using foraging effort as a proxy for population reproductive performance. First, extreme landfast ice extent, such as that observed at our study site in 2017 (over 60 km of landfast ice between the colony and open-water) may enable chick-rearing penguins to forage only temporarily, and only for a subset of highly performant individuals (Lescroël et al., 2020). Accordingly, chick-guard trip durations were particularly variable in 2017 in Pointe Géologie because some individuals were first able to forage in ice cracks, but as resources in these areas rapidly became depleted, individuals then engaged in very long trips (> 5 days, not considered in our analysis), which led to all chicks ultimately dying by mid-January (Ropert-Coudert et al., 2018). Second, land-based factors independent of food availability have been shown to cause significant reductions in reproductive success for polar seabirds, including Adélie penguins (Descamps et al., 2023; Fraser et al., 2013; Ropert-Coudert et al., 2018; Yannic et al., 2014). Unusual weather events, particularly snow accumulation around hatching and strong winds, can indeed cause significant eggs and chick losses. Accordingly, the 2023 breeding season at our study site showed lower reproductive success than predicted by trip duration alone (Fig. 3B), largely due to heavy snowfall around hatching (personal observation). Elsewhere in Antarctica, the weather regime of the western Antarctic Peninsula is currently shifting towards a more maritime and rainy state (Vignon et al., 2021).

In this region, the reproductive success of Adélie penguins appears to be particularly driven by such food-independent factors (Chapman et al., 2011; Cimino et al., 2023, 2025; Fraser et al., 2013), perhaps explaining the deviations from the general relationship between trip duration and reproductive success observed there (e.g., for Admiralty Bay, Fig. 4).

Limits to behavioural flexibility

Identifying thresholds of behavioural flexibility beyond which environmental variability impacts demographic rates would foster our ability to gauge the scale of perturbations and forecast future population trajectories (Beever et al., 2017). Our results, combining behavioural and demographic data from across Antarctica, suggest that the population-level threshold above which increased foraging effort ceases to buffer Adélie penguin reproductive success is about three times the minimum recorded foraging effort (i.e. around 29 h of trip duration, Fig. 3B). While a more precise breakpoint will likely be defined as more data become available, our results also suggest that behavioural flexibility does not break down immediately above this threshold. Instead, reproductive success appears to decline linearly with increasing foraging trip duration, only dropping to near zero when conditions are so unfavourable that trips exceed two times the average foraging effort (i.e. 60 h). Although this links to earlier findings that only some individuals within populations display the high foraging abilities needed for successful breeding under the harshest conditions (Lescroël et al., 2010, 2020), it remains unclear whether such foraging effort thresholds are consistent across species — a question that future research should address. Additionally, future research should also quantify the magnitude and frequency of poor reproductive events for which population persistence could be jeopardized (e.g., Emmerson & Southwell, 2022; Hinke et al., 2017).

Behavioural adjustments could entail carry-over effects

While behavioural adjustments may entirely or partially buffer short-term fitness components (i.e. current reproduction) against environmental variability, the potential carryover effects of such adjustments could entail negative consequences on longer-term fitness components (e.g., adult survival, (Cunningham et al., 2021). For example, increased foraging effort by common guillemot (*Uria aalge*) parents in response to low food availability led to decreased survival to the next breeding season (Wanless et al., 2023). Alternately, the foraging adjustments displayed by little auks (*Alle alle*) at several colonies with contrasted resource availability allowed them not only to maintain reproductive output, but also to preserve body condition and subsequent survival (Grémillet et al., 2012). In Adélie penguins, survival costs of reproduction linked to increased foraging effort have not been apparent at the

population-level (Emmerson et al., 2011; Lescroël et al., 2009). In long-lived species, we can expect the body condition threshold below which breeding individuals switch to self-feeding (Beaulieu, Spée, et al., 2010; Clarke et al., 2006; Tveraa et al., 1997; Watanuki et al., 2002) to be sufficiently fine tuned to prevent individuals from entering the post-breeding season with critical energy reserves. For Adélie penguins, there is considerable inter-annual variability in adult survival at some colonies (e.g., Hinke et al., 2014), and part of this variability has been linked to environmental conditions during the moult (Schmidt et al., 2023). This stage of the species life cycle takes place right after reproduction (Ainley, 2002). There is therefore a significant opportunity for investigating carry-over effects of breeding effort on adult survival in Adélie penguins, which would provide a more complete picture into the efficacy of behavioural adjustments in buffering fitness against environmental variability.

Conclusions and perspectives

To conclude, our study has addressed how Adélie penguins adjusted their foraging effort in response to an unprecedentedly wide range of sea ice conditions, especially in the absence of landfast ice, a feature predicted to define the future foraging and breeding habitats of Antarctic species (Fraser et al., 2023; Trathan et al., 2020). Importantly, we demonstrated that chick-guard foraging trip duration, a simple measure of foraging effort, represents a valuable and readily applicable metric for monitoring the reproductive performance of a seabird's colonies across its entire (continental) distribution, warranting its use as a potential early warning indicator of population-level declines (Cerini et al., 2023). Such ecological indicators are fundamental, and critical to the implementation of conservation and management plans in Antarctica (e.g., Agnew, 1997), particularly with the projected development of fisheries bolstered by sea ice losses.

Acknowledgements

We are deeply grateful to all the present and former members of Project 137, including the member of the MIBE team at IPHC in Strasbourg for their expertise and support, Yvon Le Maho (former PI of 137-ECOPHY), and all the wintering and summering field teams since the inception of this project in the field in 2005. We would also like to thank all the members of the missions in Dumont D'Urville since then, and the French Polar Institute-IPEV logistics team in Dumont d'Urville for their important and continuous support in the field. This study was approved by the French Polar Environmental Committee and authorizations for handling animals and accessing breeding sites were delivered by the Terres Australes et Antarctiques Françaises (TAAF).

Funding

This study was supported by the Institut Polaire Français Paul-Emile Victor (IPEV) within the framework of the Project 137-ANTAVIA-POLAROBS, by the Centre Scientifique de Monaco with additional support from the LIA-647 and RTPI-NUTRESS (CSM/CNRS-UNISTRA), by the Centre National de la Recherche Scientifique (CNRS) through the Programme Zone Atelier Antarctique et Terres Australes (ZATA), by the Canada Research Program, NSERC, and Université de Moncton, and by the Deutsche Forschungsgemeinschaft (DFG) grants FA336/5-1 and ZI1525/3-1 in the framework of the priority program ''Antarctic research with comparative investigations in Arctic ice areas''.

Authors contributions

Author contributions follow the CRediT guidelines. In each section, authors are listed by order of
appearance in the author list.

Conceptualization: TB, NL, CLB

Methodology: TB, AH, MB, RC, TR, NL, CLB

Software: TB, GB

Validation: TB, NL, CLB

Formal analysis: TB, GB, NL, CLB

Investigation: TB, AH, MB, PB, RC, TR, DPZ, NL, CLB

Resources: CLB

Data Curation: TB, GB, CT, LL, CLB

Writing – original draft preparation: TB, NL, CLB

Writing – review and editing: All authors

Visualization: TB

Supervision: NL, CLB

Project administration: CLB

Funding acquisition: DPZ, NL, CLB

References

Afanasyev, V., Buldyrev, S. V., Dunn, M. J., Robst, J., Preston, M., Bremner, S. F., Briggs, D. R., Brown, R., Adlard, S., & Peat, H. J. (2015). Increasing Accuracy: A New Design and Algorithm for Automatically Measuring Weights, Travel Direction and Radio Frequency Identification (RFID) of Penguins. *PLOS ONE*, *10*(4), e0126292. https://doi.org/10.1371/journal.pone.0126292

Agnew, D. J. (1997). Review—The CCAMLR Ecosystem Monitoring Programme. *Antarctic Science*, *9*(3), 235–242. https://doi.org/10.1017/S095410209700031X

Ainley, D. G. (2002). *The Adélie penguin: Bellwether of climate change*. Columbia University Press.

Ainley, D. G., Ballard, G., & Dugger, K. M. (2006). Competition Among Penguins and Cetaceans Reveals Trophic Cascades in the Western Ross Sea, Antarctica. *Ecology*, *87*(8), 2080–2093. https://doi.org/10.1890/0012-9658(2006)87[2080:CAPACR]2.0.CO;2

Ainley, D. G., Ribic, C. A., Ballard, G., Heath, S., Gaffney, I., Karl, B. J., Barton, K. J., Wilson, P. R., & Webb, S. (2004). Geographic Structure of Adélie Penguin Populations: Overlap in Colony-Specific Foraging Areas. *Ecological Monographs*, *74*(1), 159–178. https://doi.org/10.1890/02-4073

Alonzo, S. H., Switzer, P. V., & Mangel, M. (2003). Ecological games in space and time: The distribution and abundance of Antarctic krill and penguins. *Ecology*, *84*(6), 1598–1607. https://doi.org/10.1890/0012-9658(2003)084[1598:EGISAT]2.0.CO;2

Andrade, P., Franco, A. M. A., Acácio, M., Afonso, S., Marques, C. I., Moreira, F., Carneiro, M., & Catry, I. (2025). Mechanisms underlying the loss of migratory behaviour in a long-lived bird. *Journal of Animal Ecology*. https://doi.org/10.1111/1365-2656.70035

Ballance, L. T., Ainley, D. G., Ballard, G., & Barton, K. (2009). An energetic correlate between colony size and foraging effort in seabirds, an example of the Adélie penguin Pygoscelis adeliae. *Journal of Avian Biology*, *40*(3), 279–288. https://doi.org/10.1111/j.1600-048X.2008.04538.x

Ballard, G., Dugger, K., Nur, N., & Ainley, D. (2010). Foraging strategies of Adélie penguins: Adjusting body condition to cope with environmental variability. *Marine Ecology Progress Series*, *405*, 287–302. https://doi.org/10.3354/meps08514

Barbraud, C., Delord, K., Bost, C. A., Chaigne, A., Marteau, C., & Weimerskirch, H. (2020). Population trends of penguins in the French Southern Territories. *Polar Biology*, *43*(7), Article 7. https://doi.org/10.1007/s00300-020-02691-6

Bardon, G., Cristofari, R., Winterl, A., Barracho, T., Benoiste, M., Ceresa, C., Chatelain, N., Courtecuisse, J., Fernandes, F. A. N., Gauthier-Clerc, M., Gendner, J.-P., Handrich, Y., Houstin, A., Krellenstein, A., Lecomte, N., Salmon, C.-E., Trucchi, E., Vallas, B., Wong, E. M., … Le Bohec, C. (2023). RFIDeep: Unfolding the potential of deep learning for radio-frequency identification. *Methods in Ecology and Evolution*, *14*(11), 2814–2826. https://doi.org/10.1111/2041-210X.14187

Barracho, T., Bardon, G., Choquet, R., Houstin, A., Winterl, A., Beaulieu, M., Blanchard, P., Cornet, C. C., Cristofari, R., Raclot, T., Richter, S., Chatelain, N., Courtecuisse, J., Brucker, M., Zitterbart, D. P., Lecomte, N., & Le Bohec, C. (2025). Declining juvenile survival of Adélie penguins in Antarctica. *bioRxiv*, 2024.08.23.609289. https://doi.org/10.1101/2024.08.23.609289

Barreau, E., Ropert-Coudert, Y., Delord, K., Barbraud, C., & Kato-Ropert, A. (2019). Scale matters: Sea ice and breeding success of Adélie penguins. *Polar Biology*, *42*(7), Article 7. https://doi.org/10.1007/s00300-019-02531-2

Beaulieu, M., Dervaux, A., Thierry, A.-M., Lazin, D., Le Maho, Y., Ropert-Coudert, Y., Spée, M., Raclot, T., & Ancel, A. (2010). When sea-ice clock is ahead of Adélie penguins' clock. *Functional Ecology*, *24*(1), 93–102. https://doi.org/10.1111/j.1365-2435.2009.01638.x

Beaulieu, M., Spée, M., Lazin, D., Ropert-Coudert, Y., Le Maho, Y., Ancel, A., & Raclot, T. (2010). Ecophysiological response of Adelie penguins facing an experimental increase in breeding constraints. *The Journal of Experimental Biology*, *213*, 33–39. https://doi.org/10.1242/jeb.035378

Beaulieu, M., Thierry, A.-M., Raclot, T., Le Maho, Y., Ropert-Coudert, Y., Gachot-Neveu, H., & Ancel, A. (2009). Sex-specific parental strategies according to the sex of offspring in the Adélie Penguin. *Behavioral Ecology*, *20*, 870–883. https://doi.org/10.1093/beheco/arp076

Beever, E. A., Hall, L. E., Varner, J., Loosen, A. E., Dunham, J. B., Gahl, M. K., Smith, F. A., & Lawler, J. J. (2017). Behavioral flexibility as a mechanism for coping with climate change. *Frontiers in Ecology and the Environment*, *15*(6), 299–308. https://doi.org/10.1002/fee.1502

Beltran, R. S., Hernandez, K. M., Condit, R., Robinson, P. W., Crocker, D. E., Goetsch, C., Kilpatrick, A. M., & Costa, D. P. (2023). Physiological tipping points in the relationship between foraging success and lifetime fitness of a long-lived mammal. *Ecology Letters*, *26*(5), 706–716. https://doi.org/10.1111/ele.14193

Both, C., Bouwhuis, S., Lessells, C. M., & Visser, M. E. (2006). Climate change and population declines in a long-distance migratory bird. *Nature*, *441*(7089), 81–83. https://doi.org/10.1038/nature04539

Brierley, A. S., Fernandes, P. G., Brandon, M. A., Armstrong, F., Millard, N. W., McPhail, S. D., Stevenson, P., Pebody, M., Perrett, J., Squires, M., Bone, D. G., & Griffiths, G. (2002). Antarctic Krill Under Sea Ice: Elevated Abundance in a Narrow Band Just South of Ice Edge. *Science*, *295*(5561), 1890–1892. https://doi.org/10.1126/science.1068574

Brooks, M., E., Kristensen, K., Benthem, K., J.,van, Magnusson, A., Berg, C., W., Nielsen, A., Skaug, H., J., Mächler, M., & Bolker, B., M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378. https://doi.org/10.32614/RJ-2017-066

Burger, A. E., & Piatt, J. F. (1990). *Flexible time budgets in breeding common murres: Buffers against variable prey abundance*. *14*.

Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information-theoretic approach* (2. ed., [4. printing]). Springer.

Cairns, D. K. (1987). Seabirds as Indicators of Marine Food Supplies. *Biological Oceanography*, *5*, 261–

271.

Capdevila, P., Stott, I., Beger, M., & Salguero-Gómez, R. (2020). Towards a Comparative Framework of

Demographic Resilience. *Trends in Ecology & Evolution*, *35*(9), 776–786.

https://doi.org/10.1016/j.tree.2020.05.001

Cerini, F., Childs, D. Z., & Clements, C. F. (2023). A predictive timeline of wildlife population collapse.

Nature Ecology & Evolution, *7*(3), 320–331. https://doi.org/10.1038/s41559-023-01985-2

Chapman, E. W., Hofmann, E. E., Patterson, D. L., Ribic, C., & Fraser, W. R. (2011). Marine and terrestrial

factors affecting Adélie penguin Pygoscelis adeliae chick growth and recruitment off the

western Antarctic Peninsula. *Marine Ecology Progress Series*, *436*, 273–289.

https://doi.org/10.3354/meps09242

Charmantier, A., & Gienapp, P. (2014). Climate change and timing of avian breeding and migration:

Evolutionary versus plastic changes. *Evolutionary Applications*, *7*(1), 15–28.

https://doi.org/10.1111/eva.12126

Charmantier, A., McCleery, R. H., Cole, L. R., Perrins, C., Kruuk, L. E. B., & Sheldon, B. C. (2008). Adaptive

Phenotypic Plasticity in Response to Climate Change in a Wild Bird Population. *Science*,

320(5877), 800–803. https://doi.org/10.1126/science.1157174

Chown, S. L., Leihy, R. I., Naish, T. R., Brooks, C. M., Convey, P., Henley, B. J., Mackintosh, A. N., Phillips,

L. M., Ii, M. C. K., & Grant, S. M. (2022). *Antarctic Climate Change and the Environment: A*

Decadal Synopsis and Recommendations for Action. Scientific Committee on Antarctic

Research. www.scar.org

Cimino, M. A., Conroy, J. A., Connors, E., Bowman, J., Corso, A., Ducklow, H., Fraser, W., Friedlaender,

A., Kim, H. H., Larsen, G. D., Moffat, C., Nichols, R., Pallin, L., Patterson-Fraser, D., Roberts, D.,

Roberts, M., Steinberg, D. K., Thibodeau, P., Trinh, R., … Stammerjohn, S. (2023). Long-term

patterns in ecosystem phenology near Palmer Station, Antarctica, from the perspective of the

Adélie penguin. *Ecosphere*, *14*(2), e4417. https://doi.org/10.1002/ecs2.4417

Cimino, M. A., Larsen, G. D., Johnston, D. W., & Groff, D. V. (2025). Long-term, landscape- and wind-driven snow conditions influence Adélie penguin colony extinctions. *Landscape Ecology*, *40*(4), 78. https://doi.org/10.1007/s10980-025-02088-y

Clarke, J., Emmerson, L. M., Townsend, A., & Kerry, K. R. (2003). Demographic characteristics of the Adélie penguin population on Béchervaise Island after 12 years of study. *CCAMLR Science*, *10*, 53–74.

Clarke, J., Emmerson, L., & Otahal, P. (2006). Environmental conditions and life history constraints determine foraging range in breeding Adélie penguins. *Marine Ecology Progress Series*, *310*, 247–261. https://doi.org/10.3354/meps310247

Clarke, J., Manly, B., Kerry, K., Gardner, H., Franchi, E., Corsolini, S., & Focardi, S. (1998). Sex differences in Adélie penguin foraging strategies. *Polar Biology*, *20*(4), 248–258. https://doi.org/10.1007/s003000050301

Clements, C. F., & Ozgul, A. (2016). Including trait-based early warning signals helps predict population collapse. *Nature Communications*, *7*(1), 10984. https://doi.org/10.1038/ncomms10984

Convey, P., & Peck, L. S. (2019). Antarctic environmental change and biological responses. *Science Advances*, *5*(11), eaaz0888. https://doi.org/10.1126/sciadv.aaz0888

Corso, A. D., Steinberg, D. K., Stammerjohn, S. E., & Hilton, E. J. (2022). Climate drives long-term change in Antarctic Silverfish along the western Antarctic Peninsula. *Communications Biology*, *5*(1), 104. https://doi.org/10.1038/s42003-022-03042-3

Cunningham, S. J., Gardner, J. L., & Martin, R. O. (2021). Opportunity costs and the response of birds and mammals to climate warming. *Frontiers in Ecology and the Environment*, *19*(5), 300–307. https://doi.org/10.1002/fee.2324

Cury, P. M., Boyd, I. L., Bonhommeau, S., Anker-Nilssen, T., Crawford, R. J. M., Furness, R. W., Mills, J. A., Murphy, E. J., Österblom, H., Paleczny, M., Piatt, J. F., Roux, J.-P., Shannon, L., & Sydeman, W. J. (2011). Global Seabird Response to Forage Fish Depletion—One-Third for the Birds. *Science*, *334*(6063), 1703–1706. https://doi.org/10.1126/science.1212928

Davis, L. S., & McCaffrey, F. T. (1986). Survival Analysis of Eggs and Chicks of AdÉlie Penguins (Pygoscelis adeliae). *The Auk*, *103*(2), Article 2. https://doi.org/10.1093/auk/103.2.379

Dehnhard, N., Ludynia, K., Masello, J. F., Voigt, C. C., McGill, R. A. R., & Quillfeldt, P. (2016). Plasticity in foraging behaviour and diet buffers effects of inter-annual environmental differences on chick growth and survival in southern rockhopper penguins Eudyptes chrysocome chrysocome. *Polar Biology*, *39*(9), 1627–1641. https://doi.org/10.1007/s00300-015-1887-5

Dehnhard, N., Ludynia, K., Poisbleau, M., Demongin, L., & Quillfeldt, P. (2013). Good Days, Bad Days: Wind as a Driver of Foraging Success in a Flightless Seabird, the Southern Rockhopper Penguin. *PLOS ONE*, *8*(11), e79487. https://doi.org/10.1371/journal.pone.0079487

Descamps, S., Hudson, S., Sulich, J., Wakefield, E., Grémillet, D., Carravieri, A., Orskaug, S., & Steen, H. (2023). Extreme snowstorms lead to large-scale seabird breeding failures in Antarctica. *Current Biology*, *33*(5), R176–R177. https://doi.org/10.1016/j.cub.2022.12.055

Dugger, K. M., Ballard, G., Ainley, D. G., & Barton, K. J. (2006). Effects of Flipper Bands on Foraging Behavior and Survival of Adélie Penguins (Pygoscelis Adeliae). *The Auk*, *123*(3), Article 3. https://doi.org/10.1093/auk/123.3.858

Emison, W. B. (1968). Feeding preferences of the Adélie penguin at Cape Crozier, Ross Island. *Antarctic Research Series*, 191–212. https://doi.org/10.1029/AR012p0191

Emmerson, L., Pike, R., & Southwell, C. (2011). Reproductive consequences of environment-driven variation in Adélie penguin breeding phenology. *Marine Ecology Progress Series*, *440*, 203–216. https://doi.org/10.3354/meps09265

Emmerson, L., & Southwell, C. (2008). Sea Ice Cover and Its Influence on Adélie Penguin Reproductive Performance. *Ecology*, *89*(8), 2096–2102. https://doi.org/10.1890/08-0011.1

Emmerson, L., & Southwell, C. (2022). Environment-triggered demographic changes cascade and compound to propel a dramatic decline of an Antarctic seabird metapopulation. *Global Change Biology*, *28*(24). https://doi.org/10.1111/gcb.16437

Emmerson, L., Southwell, C., Clarke, J., Tierney, M., & Kerry, K. (2015). Adélie penguin response parameters signal reduced prey accessibility: Implications for predator–prey response curves. *Marine Biology*, *162*(6), 1187–1200. https://doi.org/10.1007/s00227-015-2661-5

Emmerson, L., Walsh, S., & Southwell, C. (2019). Nonbreeder birds at colonies display qualitatively similar seasonal mass change patterns as breeders. *Ecology and Evolution*, *9*(8), Article 8. https://doi.org/10.1002/ece3.5067

Enquist, B. J., Norberg, J., Bonser, S. P., Violle, C., Webb, C. T., Henderson, A., Sloat, L. L., & Savage, V. M. (2015). Scaling from Traits to Ecosystems. In *Advances in Ecological Research* (Vol. 52, pp. 249–318). Elsevier. https://doi.org/10.1016/bs.aecr.2015.02.001

Fayet, A., Shoji, A., Freeman, R., Perrins, C., & Guilford, T. (2017). Within-pair similarity in migration route and female winter foraging effort predict pair breeding performance in a monogamous seabird. *Marine Ecology Progress Series*, *569*, 243–252. https://doi.org/10.3354/meps12083

Finn, C., Grattarola, F., & Pincheira-Donoso, D. (2023). More losers than winners: Investigating Anthropocene defaunation through the diversity of population trends. *Biological Reviews*, *98*(5), 1732–1748. https://doi.org/10.1111/brv.12974

Foster, S. D., & Bravington, M. V. (2013). A Poisson–Gamma model for analysis of ecological non-negative continuous data. *Environmental and Ecological Statistics*, *20*(4), 533–552. https://doi.org/10.1007/s10651-012-0233-0

Fraser, A. D., Massom, R. A., Handcock, M. S., Reid, P., Ohshima, K. I., Raphael, M. N., Cartwright, J., Klekociuk, A. R., Wang, Z., & Porter-Smith, R. (2021). Eighteen-year record of circum-Antarctic landfast-sea-ice distribution allows detailed baseline characterisation and reveals trends and variability. *The Cryosphere*, *15*(11), 5061–5077. https://doi.org/10.5194/tc-15-5061-2021

Fraser, A. D., Wongpan, P., Langhorne, P. J., Klekociuk, A. R., Kusahara, K., Lannuzel, D., Massom, R. A., Meiners, K. M., Swadling, K. M., Atwater, D. P., Brett, G. M., Corkill, M., Dalman, L. A., Fiddes, S., Granata, A., Guglielmo, L., Heil, P., Leonard, G. H., Mahoney, A. R., … Wienecke, B. (2023).

Antarctic Landfast Sea Ice: A Review of Its Physics, Biogeochemistry and Ecology. *Reviews of Geophysics*, *61*(2), e2022RG000770. https://doi.org/10.1029/2022RG000770

Fraser, W., Patterson-Fraser, D., Ribic, C., Schofield, O., & Ducklow, H. (2013). A Nonmarine Source of Variability in Adélie Penguin Demography. *Oceanography*, *26*(3), Article 3. https://doi.org/10.5670/oceanog.2013.64

Fricker, H. A., Galton-Fenzi, B. K., Walker, C. C., Freer, B. I. D., Padman, L., & DeConto, R. (2025). Antarctica in 2025: Drivers of deep uncertainty in projected ice loss. *Science*, *387*(6734), 601–609. https://doi.org/10.1126/science.adt9619

Fuentes, M., Santos, A., Abreu, A., Briseno, R., Al-Khayat, J., Hamza, S., Saliba, S., Anderson, D., Rusenko, K., Mitchell, N., Gammon, M., Bentley, B., Beton, D., Booth, D., Broderick, A., Colman, L., Snape, R., Calderon-Campuzano, M., Cuevas, E., & Monsinjon, J. (2023). Adaptation of sea turtles to climate warming: Will phenological responses be sufficient to counteract changes in reproductive output? *Global Change Biology*, *30*. https://doi.org/10.1111/gcb.16991

Gallagher, K. L., Dinniman, M. S., & Lynch, H. J. (2023). Quantifying Antarctic krill connectivity across the West Antarctic Peninsula and its role in large-scale Pygoscelis penguin population dynamics. *Scientific Reports*, *13*(1), 12072. https://doi.org/10.1038/s41598-023-39105-6

Gianuca, D., Votier, S. C., Pardo, D., Wood, A. G., Sherley, R. B., Ireland, L., Choquet, R., Pradel, R., Townley, S., Forcada, J., Tuck, G. N., & Phillips, R. A. (2019). Sex-specific effects of fisheries and climate on the demography of sexually dimorphic seabirds. *Journal of Animal Ecology*, *88*(9), 1366–1378. https://doi.org/10.1111/1365-2656.13009

Gil, M. A., Baskett, M. L., Munch, S. B., & Hein, A. M. (2020). Fast behavioral feedbacks make ecosystems sensitive to pace and not just magnitude of anthropogenic environmental change. *Proceedings of the National Academy of Sciences*, *117*(41), 25580–25589. https://doi.org/10.1073/pnas.2003301117

Grémillet, D., & Charmantier, A. (2010). Shifts in phenotypic plasticity constrain the value of seabirds as ecological indicators of marine ecosystems. *Ecological Applications*, *20*(6), 1498–1503. https://doi.org/10.1890/09-1586.1

Grémillet, D., Welcker, J., Karnovsky, N., Walkusz, W., Hall, M., Fort, J., Brown, Z., Speakman, J., & Harding, A. (2012). Little auks buffer the impact of current Arctic climate change. *Marine Ecology Progress Series*, *454*, 197–206. https://doi.org/10.3354/meps09590

Halupka, L., Arlt, D., Tolvanen, J., Millon, A., Bize, P., Adamík, P., Albert, P., Arendt, W. J., Artemyev, A. V., Baglione, V., Bańbura, J., Bańbura, M., Barba, E., Barrett, R. T., Becker, P. H., Belskii, E., Bolton, M., Bowers, E. K., Bried, J., … Halupka, K. (2023). The effect of climate change on avian offspring production: A global meta-analysis. *Proceedings of the National Academy of Sciences*, *120*(19), e2208389120. https://doi.org/10.1073/pnas.2208389120

Hamer, K., Humphreys, E., Garthe, S., Hennicke, J., Peters, G., Grémillet, D., Phillips, R., Harris, M., & Wanless, S. (2007). Annual variation in diets, feeding locations and foraging behaviour of gannets in the North Sea: Flexibility, consistency and constraint. *Marine Ecology Progress Series*, *338*, 295–305. https://doi.org/10.3354/meps338295

Harding, A. M. A., Piatt, J. F., Schmutz, J. A., Shultz, M. T., Pelt, T. I. V., Kettle, A. B., & Speckman, S. G. (2007). Prey Density and the Behavioral Flexibility of a Marine Predator: The Common Murre (uria Aalge). *Ecology*, *88*(8), 2024–2033. https://doi.org/10.1890/06-1695.1

Hartig, F. (2024). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models* (Version R package version 0.4.7) [Computer software]. https://github.com/florianhartig/dharma

Hinke, J. T., Trivelpiece, S. G., & Trivelpiece, W. Z. (2014). Adélie penguin (Pygoscelis adeliae) survival rates and their relationship to environmental indices in the South Shetland Islands, Antarctica. *Polar Biology*, *37*(12), Article 12. https://doi.org/10.1007/s00300-014-1562-2

Hinke, J. T., Trivelpiece, S. G., & Trivelpiece, W. Z. (2017). Variable vital rates and the risk of population

declines in Adélie penguins from the Antarctic Peninsula region. *Ecosphere*, *8*(1), e01666.

https://doi.org/10.1002/ecs2.1666

Horswill, C., Trathan, P. N., & Ratcliffe, N. (2017). Linking extreme interannual changes in prey

availability to foraging behaviour and breeding investment in a marine predator, the macaroni

penguin. *PLOS ONE*, *12*(9), e0184114. https://doi.org/10.1371/journal.pone.0184114

Houston, A. I., & McNamara, J. M. (2014). Foraging currencies, metabolism and behavioural routines.

Journal of Animal Ecology, *83*(1), 30–40. https://doi.org/10.1111/1365-2656.12096

Iles, D. T., Lynch, H., Ji, R., Barbraud, C., Delord, K., & Jenouvrier, S. (2020). Sea ice predicts long-term

trends in Adélie penguin population growth, but not annual fluctuations: Results from a range-

wide multiscale analysis. *Global Change Biology*, *26*(7), Article 7.

https://doi.org/10.1111/gcb.15085

Iles, D. T., Rockwell, R. F., & Koons, D. N. (2019). Shifting Vital Rate Correlations Alter Predicted

Population Responses to Increasingly Variable Environments. *The American Naturalist*, *193*(3),

Article 3. https://doi.org/10.1086/701043

Jennings, S., Dugger, K. M., Ballard, G., & Ainley, D. G. (2021). Effects of Diet and Provisioning Behavior

on Chick Growth in Adélie Penguins (Pygoscelis adeliae). *Waterbirds*, *44*(1).

https://doi.org/10.1675/063.044.0105

Jennings, S., Dugger, K. M., Ballard, G., & Ainley, D. G. (2023). Faster growth and larger size at crèche

onset are associated with higher offspring survival in Adélie Penguins. *Ornithology*, ukad006.

https://doi.org/10.1093/ornithology/ukad006

Jenouvrier, S. (2013). Impacts of climate change on avian populations. *Global Change Biology*, *19*(7),

Article 7. https://doi.org/10.1111/gcb.12195

Kawaguchi, S., Atkinson, A., Bahlburg, D., Bernard, K. S., Cavan, E. L., Cox, M. J., Hill, S. L., Meyer, B., &

Veytia, D. (2023). Climate change impacts on Antarctic krill behaviour and population

dynamics. *Nature Reviews Earth & Environment*, *5*(1), 43–58. https://doi.org/10.1038/s43017-023-00504-y

Kearney, M., Shine, R., & Porter, W. P. (2009). The potential for behavioral thermoregulation to buffer "cold-blooded" animals against climate warming. *Proceedings of the National Academy of Sciences*, *106*(10), 3835–3840. https://doi.org/10.1073/pnas.0808913106

Kerry, K., Clarke, J., & Else, G. (1993). The use of an automated weighing and recording system for the study of the biology of Adélie penguins. *Proc. NIPR Symp. Polar Biol.*, *6*, 62–75.

Kim, Y., Kim, J.-U., Oh, Y., Park, J., Lee, K., Kim, H.-C., & Kim, J.-H. (2024). Adaptive foraging strategies of Adélie penguins in the Ross Sea Region: Balancing chick feeding and body condition in changing marine environments. *Marine Biology*, *172*(1), 15. https://doi.org/10.1007/s00227-024-04575-3

Kokubun, N., Emmerson, L., McInnes, J., Wienecke, B., & Southwell, C. (2021). Sea-ice and density-dependent factors affecting foraging habitat and behaviour of Adélie penguins throughout the breeding season. *Marine Biology*, *168*(7), 97. https://doi.org/10.1007/s00227-021-03899-8

Le Guen, C., Kato, A., Raymond, B., Barbraud, C., Beaulieu, M., Bost, C.-A., Delord, K., MacIntosh, A. J. J., Meyer, X., Raclot, T., Sumner, M., Takahashi, A., Thiebot, J.-B., & Ropert-Coudert, Y. (2018). Reproductive performance and diving behaviour share a common sea-ice concentration optimum in Adélie penguins (Pygoscelis adeliae). *Global Change Biology*, *24*(11), Article 11. https://doi.org/10.1111/gcb.14377

Lescroël, A., Ballard, G., Grémillet, D., Authier, M., & Ainley, D. G. (2014). Antarctic Climate Change: Extreme Events Disrupt Plastic Phenotypic Response in Adélie Penguins. *PLOS ONE*, *9*(1), Article 1. https://doi.org/10.1371/journal.pone.0085291

Lescroël, A., Ballard, G., Toniolo, V., Barton, K. J., Wilson, P. R., Lyver, P. O., & Ainley, D. G. (2010). Working less to gain more: When breeding quality relates to foraging efficiency. *Ecology*, *91*(7), Article 7. https://doi.org/10.1890/09-0766.1

Lescroël, A., Dugger, K. M., Ballard, G., & Ainley, D. G. (2009). Effects of individual quality, reproductive success and environmental variability on survival of a long-lived seabird. *Journal of Animal Ecology*, *78*(4), Article 4. https://doi.org/10.1111/j.1365-2656.2009.01542.x

Lescroël, A., Lyver, P., Jongsomjit, D., Veloz, S., Dugger, K., Kappes, P., Karl, B., Whitehead, A., Pech, R., Cole, T., & Ballard, G. (2020). Inter-individual differences in the foraging behavior of breeding Adélie penguins are driven by individual quality and sex. *Marine Ecology Progress Series*, *636*, 189–205. https://doi.org/10.3354/meps13208

Lescroël, A., Schmidt, A., Elrod, M., Ainley, D. G., & Ballard, G. (2021). Foraging dive frequency predicts body mass gain in the Adélie penguin. *Scientific Reports*, *11*(1), 22883. https://doi.org/10.1038/s41598-021-02451-4

Lewis, S., Wanless, S., Elston, D. A., Schultz, M. D., Mackley, E., Du Toit, M., Underhill, J. G., & Harris, M. P. (2006). Determinants of quality in a long-lived colonial species. *Journal of Animal Ecology*, *75*(6), 1304–1312. https://doi.org/10.1111/j.1365-2656.2006.01152.x

Lewis, S., Wanless, S., Wright, P., Harris, M., Bull, J., & Elston, D. A. (2001). Diet and breeding performance of Black-legged Kittiwakes Rissa tridactyla at a North Sea colony. *Marine Ecology-Progress Series - MAR ECOL-PROGR SER*, *221*, 277–284. https://doi.org/10.3354/meps221277

Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance: An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of Open Source Software*, *6*(60), 3139. https://doi.org/10.21105/joss.03139

Machado-Gaye, A. L., Chimienti, M., Kato, A., Ropert-Coudert, Y., Barbosa, A., Bustamante, P., Cosse, M., Menéndez-Blázquez, J., Zaldúa, N., & Soutullo, A. (2025). Energy expenditure of Adélie penguins during the breeding season: Females pay the cost in years of low food availability. *Animal Behaviour*, *222*, 123144. https://doi.org/10.1016/j.anbehav.2025.123144

Mason, T. H. E., Stephens, P. A., Apollonio, M., & Willis, S. G. (2014). Predicting potential responses to future climate in an alpine ungulate: Interspecific interactions exceed climate effects. *Global Change Biology*, *20*(12), 3872–3882. https://doi.org/10.1111/gcb.12641

Méheust, Y., Delord, K., Bonnet-Lebrun, A.-S., Raclot, T., Vasseur, J., Allain, J., Decourteillle, V., Bost, C.-A., & Barbraud, C. (2024). Human infrastructures correspond to higher Adélie penguin breeding success and growth rate. *Oecologia*, *204*(3), 675–688. https://doi.org/10.1007/s00442-024-05523-0

Meier, W., Fetterer, F., Windnagel, A., & Stewart, S. (2021). *NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 4* [Dataset]. NSIDC. https://doi.org/10.7265/EFMZ-2T65

Merkel, B., Trathan, P., Thorpe, S., Murphy, E. J., Pehlke, H., Teschke, K., & Griffith, G. P. (2023). Quantifying circumpolar summer habitat for Antarctic krill and Ice krill, two key species of the Antarctic marine ecosystem. *ICES Journal of Marine Science*, *80*(6), 1773–1786. https://doi.org/10.1093/icesjms/fsad110

Meyer, B., Freier, U., Grimm, V., Groeneveld, J., Hunt, B. P. V., Kerwath, S., King, R., Klaas, C., Pakhomov, E., Meiners, K. M., Melbourne-Thomas, J., Murphy, E. J., Thorpe, S. E., Stammerjohn, S., Wolf-Gladrow, D., Auerswald, L., Götz, A., Halbach, L., Jarman, S., … Yilmaz, N. I. (2017). The winter pack-ice zone provides a sheltered but food-poor habitat for larval Antarctic krill. *Nature Ecology & Evolution*, *1*(12), 1853–1861. https://doi.org/10.1038/s41559-017-0368-3

Michelot, C. (2020). *Le manchot Adélie, sentinelle de la glace de mer: Étude du comportement de recherche alimentaire en réponse aux variations environnementales en période de reproduction*.

Michelot, C., Kato, A., Raclot, T., & Ropert-Coudert, Y. (2021). Adélie penguins foraging consistency and site fidelity are conditioned by breeding status and environmental conditions. *PLOS ONE*, *16*(1), Article 1. https://doi.org/10.1371/journal.pone.0244298

Miller, G. D., & Davis, L. S. (1993). Foraging flexibility of Adélie penguins Pygoscelis adeliae: Consequences for an indicator species. *Biological Conservation*, *63*, 223–230.

Milles, A., Banitz, T., Bielcik, M., Frank, K., Gallagher, C. A., Jeltsch, F., Jepsen, J. U., Oro, D., Radchuk, V., & Grimm, V. (2023). Local buffer mechanisms for population persistence. *Trends in Ecology & Evolution*, *38*(11), 1051–1059. https://doi.org/10.1016/j.tree.2023.06.006

Mínguez, E., & Oro, D. (2003). *Variations in Nest Mortality in the European Storm Petrel Hydrobates Pelagicus*.

Møller, A. P., Rubolini, D., & Lehikoinen, E. (2008). Populations of migratory bird species that did not show a phenological response to climate change are declining. *Proceedings of the National Academy of Sciences*, *105*(42), 16195–16200. https://doi.org/10.1073/pnas.0803825105

Morandini, V., Dugger, K. M., Schmidt, A. E., Varsani, A., Lescroël, A., Ballard, G., Lyver, P. O., Barton, K., & Ainley, D. G. (2024). Sex-specific recruitment rates contribute to male-biased sex ratio in Adélie penguins. *Ecology and Evolution*, *14*(2), e10859. https://doi.org/10.1002/ece3.10859

Muthukrishnan, R., Smiley, T. M., Title, P. O., Fudickar, A. M., Jahn, A. E., & Lau, J. A. (2025). Chasing the Niche: Escaping Climate Change Threats in Place, Time, and Space. *Global Change Biology*, *31*(4), e70167. https://doi.org/10.1111/gcb.70167

Nakagawa, S., Johnson, P. C. D., & Schielzeth, H. (2017). The coefficient of determination R2 and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *Journal of The Royal Society Interface*, *14*(134), 20170213. https://doi.org/10.1098/rsif.2017.0213

Nakazawa, T., Hsu, Y., & Chen, I. (2023). Why sex matters in phenological research. *Oikos*, *2023*(8), e09808. https://doi.org/10.1111/oik.09808

Nicol, S., Clarke, J., Romaine, S. J., Kawaguchi, S., Williams, G., & Hosie, G. W. (2008). Krill (Euphausia superba) abundance and Adélie penguin (Pygoscelis adeliae) breeding performance in the waters off the Béchervaise Island colony, East Antarctica in 2 years with contrasting ecological conditions. *Deep Sea Research Part II: Topical Studies in Oceanography*, *55*(3–4), 540–557. https://doi.org/10.1016/j.dsr2.2007.11.013

Ollason, J., & Dunnet, G. M. (1980). Nest failures in the fulmar: The effect of observers. *Journal of Field*

Ornithology.

Olmastroni, S., Corsolini, S., Pezzo, F., Focardi, S., & Kerry, K. (2000). The first five years of the Italian-

Australian joint programme on the Adélie Penguin: An overview. *Italian Journal of Zoology*,

67(sup1), 141–145. https://doi.org/10.1080/11250000009356369

Oosthuizen, W. C., Pistorius, P. A., Korczak-Abshire, M., Hinke, J. T., Santos, M., & Lowther, A. D. (2022).

The foraging behavior of nonbreeding Adélie penguins in the western Antarctic Peninsula

during the breeding season. *Ecosphere*, *13*(5). https://doi.org/10.1002/ecs2.4090

Orgeret, F., Thiebault, A., Kovacs, K. M., Lydersen, C., Hindell, M. A., Thompson, S. A., Sydeman, W. J.,

& Pistorius, P. A. (2022). Climate change impacts on seabirds and marine mammals: The

importance of study duration, thermal tolerance and generation time. *Ecology Letters*, *25*(1),

218–239. https://doi.org/10.1111/ele.13920

Orians, G. H., & Pearson, N. E. (1979). On the theory of central place foraging. In *Analysis of Ecological*

Systems (D.J. Horn, R.D. Mitchell, G.R. Stairs, pp. 155–177). Ohio State University Press.

Paniw, M., García-Callejas, D., Lloret, F., Bassar, R. D., Travis, J., & Godoy, O. (2023). Pathways to global-

change effects on biodiversity: New opportunities for dynamically forecasting demography and

species interactions. *Proceedings of the Royal Society B: Biological Sciences*, *290*(1993),

20221494. https://doi.org/10.1098/rspb.2022.1494

Paniw, M., Maag, N., Cozzi, G., Clutton-Brock, T., & Ozgul, A. (2019). Life history responses of meerkats

to seasonal changes in extreme environments. *Science*, *363*(6427), 631–635.

https://doi.org/10.1126/science.aau5905

Park, S., Chung, H., & Lee, W. Y. (2022). Behavioral responses of Adélie penguins confronting a giant ice

floe. *Deep Sea Research Part II: Topical Studies in Oceanography*, *203*, 105152.

https://doi.org/10.1016/j.dsr2.2022.105152

Peck, Ll. (2018). *Antarctic Marine Biodiversity: Adaptations, Environments and Responses to Change:*

An Annual Review (pp. 105–236). https://doi.org/10.1201/9780429454455-3

Pedersen, E. J., Miller, D. L., Simpson, G. L., & Ross, N. (2019). Hierarchical generalized additive models in ecology: An introduction with mgcv. *PeerJ*, *7*, e6876. https://doi.org/10.7717/peerj.6876

Piatt, J. F., Harding, A. M. A., Shultz, M., Speckman, S. G., Van Pelt, T. I., Drew, G. S., & Kettle, A. B. (2007). Seabirds as indicators of marine food supplies: Cairns revisited. *Marine Ecology Progress Series*, *352*, 221–234.

Piersma, T., & Drent, J. (2003). Phenotypic flexibility and the evolution of organismal design. *Trends in Ecology & Evolution*, *18*(5), 228–233. https://doi.org/10.1016/S0169-5347(03)00036-3

Ponchon, A., Grémillet, D., Christensen-Dalsgaard, S., Erikstad, K. E., Barrett, R. T., Reiertsen, T. K., McCoy, K. D., Tveraa, T., & Boulinier, T. (2014). When things go wrong: Intra-season dynamics of breeding failure in a seabird. *Ecosphere*, *5*(1), Article 1. https://doi.org/10.1890/ES13-00233.1

Purich, A., & Doddridge, E. W. (2023). Record low Antarctic sea ice coverage indicates a new sea ice state. *Communications Earth & Environment*, *4*(1), Article 1. https://doi.org/10.1038/s43247-023-00961-9

R Core Team. (2024). [Computer software].

Rackow, T., Danilov, S., Goessling, H. F., Hellmer, H. H., Sein, D. V., Semmler, T., Sidorenko, D., & Jung, T. (2022). Delayed Antarctic sea-ice decline in high-resolution climate change simulations. *Nature Communications*, *13*(1), 637. https://doi.org/10.1038/s41467-022-28259-y

Raphael, M. N., Maierhofer, T. J., Fogt, R. L., Hobbs, W. R., & Handcock, M. S. (2025). A twenty-first century structural change in Antarctica's sea ice system. *Communications Earth & Environment*, *6*(1), 1–9. https://doi.org/10.1038/s43247-025-02107-5

Reed, T. E., Grøtan, V., Jenouvrier, S., Sæther, B.-E., & Visser, M. E. (2013). Population Growth in a Wild Bird Is Buffered Against Phenological Mismatch. *Science*, *340*(6131), 488–491. https://doi.org/10.1126/science.1232870

Riaz, J., Bestley, S., Wotherspoon, S., Cox, M. J., & Emmerson, L. (2023). Spatial link between Adélie penguin foraging effort and krill swarm abundance and distribution. *Frontiers in Marine Science*, *10*, 1060984. https://doi.org/10.3389/fmars.2023.1060984

Riaz, J., Bestley, S., Wotherspoon, S., Freyer, J., & Emmerson, L. (2020). From trips to bouts to dives: Temporal patterns in the diving behaviour of chick-rearing Adélie penguins, East Antarctica. *Marine Ecology Progress Series*, *654*, 177–194. https://doi.org/10.3354/meps13519

Rintoul, S. R., Chown, S. L., DeConto, R. M., England, M. H., Fricker, H. A., Masson-Delmotte, V., Naish, T. R., Siegert, M. J., & Xavier, J. C. (2018). Choosing the future of Antarctica. *Nature*, *558*(7709), Article 7709. https://doi.org/10.1038/s41586-018-0173-4

Robinson, S., Chiaradia, A., & Hindell, M. A. (2005). The effect of body condition on the timing and success of breeding in Little Penguins Eudyptula minor. *Ibis*, *147*(3), Article 3. https://doi.org/10.1111/j.1474-919x.2005.00431.x

Ropert-Coudert, Y., Kato, A., Shiomi, K., Barbraud, C., Angelier, F., Delord, K., Poupart, T., Koubbi, P., & Raclot, T. (2018). Two Recent Massive Breeding Failures in an Adélie Penguin Colony Call for the Creation of a Marine Protected Area in D'Urville Sea/Mertz. *Frontiers in Marine Science*, *5*. https://doi.org/10.3389/fmars.2018.00264

Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., Stanton, J. C., Panjabi, A., Helft, L., Parr, M., & Marra, P. P. (2019). Decline of the North American avifauna. *Science*, *366*(6461), 120–124. https://doi.org/10.1126/science.aaw1313

Ryabov, A., Berger, U., Blasius, B., & Meyer, B. (2023). Driving forces of Antarctic krill abundance. *Science Advances*, *9*(50), eadh4584. https://doi.org/10.1126/sciadv.adh4584

Saenz, B. T., Ainley, D. G., Daly, K. L., Ballard, G., Conlisk, E., Elrod, M. L., & Kim, S. L. (2020). Drivers of concentrated predation in an Antarctic marginal-ice-zone food web. *Scientific Reports*, *10*(1), 7282. https://doi.org/10.1038/s41598-020-63875-y

Sæther, B.-E., Grøtan, V., Engen, S., Coulson, T., Grant, P. R., Visser, M. E., Brommer, J. E., Rosemary Grant, B., Gustafsson, L., Hatchwell, B. J., Jerstad, K., Karell, P., Pietiäinen, H., Roulin, A., Røstad,

O. W., & Weimerskirch, H. (2016). Demographic routes to variability and regulation in bird populations. *Nature Communications*, *7*(1), 12001. https://doi.org/10.1038/ncomms12001

Santora, J. A., LaRue, M. A., & Ainley, D. G. (2020). Geographic structuring of Antarctic penguin populations. *Global Ecology and Biogeography*, *29*(10), 1716–1728. https://doi.org/10.1111/geb.13144

Schmidt, A. E., Ballard, G., Lescroël, A., Dugger, K. M., Jongsomjit, D., Elrod, M. L., & Ainley, D. G. (2021). The influence of subcolony-scale nesting habitat on the reproductive success of Adélie penguins. *Scientific Reports*, *11*(1), 15380. https://doi.org/10.1038/s41598-021-94861-7

Schmidt, A. E., Lescroël, A., Lisovski, S., Elrod, M., Jongsomjit, D., Dugger, K. M., & Ballard, G. (2023). Sea ice concentration decline in an important Adélie penguin molt area. *Proceedings of the National Academy of Sciences*, *120*(46), e2306840120. https://doi.org/10.1073/pnas.2306840120

Schoener, T. W. (1971). Theory of Feeding Strategies. *Annual Review of Ecology and Systematics*, *2*, 369–404.

Schultz, H., Chang, K., Bury, S. J., Gaskett, A. C., Dennis, T. E., Ismar-Rebitz, S. M. H., Southey, I., Hohnhold, R. J., & Millar, C. D. (2021). Sex-specific foraging of an apex predator puts females at risk of human–wildlife conflict. *Journal of Animal Ecology*, *90*(7), 1776–1786. https://doi.org/10.1111/1365-2656.13496

Şen, B., Che-Castaldo, C., LaRue, M. A., Krumhardt, K. M., Landrum, L., Holland, M. M., Lynch, H. J., Delord, K., Barbraud, C., & Jenouvrier, S. (2025). *Temporal and spatial equivalence in demographic responses of emperor penguins (Aptenodytes forsteri) to environmental change*. https://doi.org/10.1111/1365-2656.70025

Senner, N. R., Verhoeven, M. A., Abad-Gómez, J. M., Gutiérrez, J. S., Hooijmeijer, J. C. E. W., Kentie, R., Masero, J. A., Tibbitts, T. L., & Piersma, T. (2015). When Siberia came to the Netherlands: The response of continental black-tailed godwits to a rare spring weather event. *Journal of Animal Ecology*, *84*(5), 1164–1176. https://doi.org/10.1111/1365-2656.12381

Smiley, K., & Emmerson, L. (2016). A matter of timing: Adélie penguin reproductive success in a seasonally varying environment. *Marine Ecology Progress Series*, *542*, 235–249. https://doi.org/10.3354/meps11536

Southwell, C., Clarke, J., & Emmerson, L. M. (2006). *Assessing the power to detect systematic change in Adélie penguin foraging trip duration*.

Stager, M., Velotta, J. P., Cheviron, Z. A., & Senner, N. R. (2024). Assessing the evidence for treating developmental plasticity and phenotypic flexibility as different phenomena. *Functional Ecology*, 1365-2435.14590. https://doi.org/10.1111/1365-2435.14590

Stammerjohn, S. E., Martinson, D. G., Smith, R. C., & Iannuzzi, R. A. (2008). Sea ice in the western Antarctic Peninsula region: Spatio-temporal variability from ecological and climate change perspectives. *Deep Sea Research Part II: Topical Studies in Oceanography*, *55*(18–19), 2041–2058. https://doi.org/10.1016/j.dsr2.2008.04.026

Sutherland, W. J., & Norris, K. (2002). Behavioural models of population growth rates: Implications for conservation and prediction. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *357*(1425), 1273–1284. https://doi.org/10.1098/rstb.2002.1127

Takahashi, A., Ito, M., Nagai, K., Thiebot, J., Mitamura, H., Noda, T., Trathan, P., Tamura, T., & Watanabe, Y. (2018). Migratory movements and winter diving activity of Adélie penguins in East Antarctica. *Marine Ecology Progress Series*, *589*, 227–239. https://doi.org/10.3354/meps12438

Thiebot, J.-B., Ropert-Coudert, Y., Raclot, T., Poupart, T., Kato, A., & Takahashi, A. (2019). Adélie penguins' extensive seasonal migration supports dynamic Marine Protected Area planning in Antarctica. *Marine Policy*, *109*, 103692. https://doi.org/10.1016/j.marpol.2019.103692

Trathan, P., Friedlaender, A., Johnson, C., & Reisinger, R. (2025). The fishery for Antarctic krill – Conflicts between industrial production, protection of biodiversity, and legal governance. *Marine Policy*, *180*, 106787. https://doi.org/10.1016/j.marpol.2025.106787

Trathan, P. N., Wienecke, B., Barbraud, C., Jenouvrier, S., Kooyman, G., Le Bohec, C., Ainley, D. G., Ancel, A., Zitterbart, D. P., Chown, S. L., LaRue, M., Cristofari, R., Younger, J., Clucas, G., Bost, C.-A.,

Brown, J. A., Gillett, H. J., & Fretwell, P. T. (2020). The emperor penguin—Vulnerable to projected rates of warming and sea ice loss. *Biological Conservation*, *241*, 108216. https://doi.org/10.1016/j.biocon.2019.108216

Trivelpiece, W. Z., Trivelpiece, S. G., Geupel, G. R., Kjelmyr, J., & Volkman, N. J. (1990). Adélie and Chinstrap Penguins: Their Potential as Monitors of the Southern Ocean Marine Ecosystem. In K. R. Kerry & G. Hempel (Eds.), *Antarctic Ecosystems* (pp. 191–202). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-84074-6_20

Tveraa, T., Lorensten, S.-H., & Sæther, B.-E. (1997). Regulation of foraging trips and costs of incubation shifts in the Antarctic petrel (*Thalassoica antarctica*). *Behavioral Ecology*, *8*(5), 465–469. https://doi.org/10.1093/beheco/8.5.465

Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, *348*(6234), 571–573. https://doi.org/10.1126/science.aaa4984

Van Buskirk, J. (2012). Behavioural plasticity and environmental change. In *Behavioral responses to a changing world: Mechanisms and consequences* (Candolin U and Wong BBM). Oxford University Press.

Viblanc, V., & Muller-Landau, H. (2025). Insights Into Ecology, Evolution and Global Change Responses From Very Long-Term Studies. *Ecology Letters*, *28*. https://doi.org/10.1111/ele.70072

Vignon, É., Roussel, M.-L., Gorodetskaya, I. V., Genthon, C., & Berne, A. (2021). Present and Future of Rainfall in Antarctica. *Geophysical Research Letters*, *48*(8), e2020GL092281. https://doi.org/10.1029/2020GL092281

Visser, M. E., & Gienapp, P. (2019). Evolutionary and demographic consequences of phenological mismatches. *Nature Ecology & Evolution*, *3*(6), 879–885. https://doi.org/10.1038/s41559-019-0880-8

Wanless, S., Albon, S. D., Daunt, F., Sarzo, B., Newell, M. A., Gunn, C., Speakman, J. R., & Harris, M. P. (2023). Increased parental effort fails to buffer the cascading effects of warmer seas on

common guillemot demographic rates. *Journal of Animal Ecology*, *92*(8), 1622–1638. https://doi.org/10.1111/1365-2656.13944

Watanabe, Y. Y., Ito, K., Kokubun, N., & Takahashi, A. (2020). Foraging behavior links sea ice to breeding success in Antarctic penguins. *Science Advances*, *6*(26), Article 26. https://doi.org/10.1126/sciadv.aba4828

Watanuki, Y., Kato, A., Sato, K., Niizuma, Y., Bost, C. A., Le Maho, Y., & Naito, Y. (2002). Parental mass change and food provisioning in Adélie penguins rearing chicks in colonies with contrasting sea-ice conditions. *Polar Biology*, *25*(9), 672–681. https://doi.org/10.1007/s00300-002-0399-2

Watters, G. M., Hinke, J. T., & Reiss, C. S. (2020). Long-term observations from Antarctica demonstrate that mismatched scales of fisheries management and predator-prey interaction lead to erroneous conclusions about precaution. *Scientific Reports*, *10*(1), 2314. https://doi.org/10.1038/s41598-020-59223-9

Widmann, M., Kato, A., Raymond, B., Angelier, F., Arthur, B., Chastel, O., Pellé, M., Raclot, T., & Ropert-Coudert, Y. (2015). Habitat use and sex-specific foraging behaviour of Adélie penguins throughout the breeding season in Adélie Land, East Antarctica. *Movement Ecology*, *3*(1), Article 1. https://doi.org/10.1186/s40462-015-0052-7

Wienecke, B. C., Lawless, R., Rodary, D., Thomson, R., Pauly, T., Robertson, G., Kerry, K. R., & LeMaho, Y. (2000). *Adélie penguin foraging behaviour and krill abundance along the Wilkes and Adélie land coasts, Antarctica*.

Wiens, J. J., & Zelinka, J. (2024). How many species will Earth lose to climate change? *Global Change Biology*, *30*(1), e17125. https://doi.org/10.1111/gcb.17125

Wildermuth, R. P., Anadón, J. D., & Gerber, L. R. (2013). Monitoring behavior: Assessing population status with rapid behavioral assessment. *Conservation Letters*, *6*(2), 86–97. https://doi.org/10.1111/j.1755-263X.2012.00298.x

Wong, B. B. M., & Candolin, U. (2015). Behavioral responses to changing environments. *Behavioral Ecology*, *26*(3), Article 3. https://doi.org/10.1093/beheco/aru183

Wood, S. N. (2011). Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation of Semiparametric Generalized Linear Models. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *73*(1), 3–36. https://doi.org/10.1111/j.1467-9868.2010.00749.x

Wood, S. N., Scheipl, F., & Faraway, J. J. (2013). Straightforward intermediate rank tensor product smoothing in mixed models. *Statistics and Computing*, *23*(3), 341–360. https://doi.org/10.1007/s11222-012-9314-z

Yannic, G., Aebischer, A., Sabard, B., & Gilg, O. (2014). Complete breeding failures in ivory gull following unusual rainy storms in North Greenland. *Polar Research*. https://doi.org/10.3402/polar.v33.22749

Youngflesh, C., Jenouvrier, S., Li, Y., Ji, R., Ainley, D. G., Ballard, G., Barbraud, C., Delord, K., Dugger, K. M., Emmerson, L. M., Fraser, W. R., Hinke, J. T., Lyver, P. O., Olmastroni, S., Southwell, C. J., Trivelpiece, S. G., Trivelpiece, W. Z., & Lynch, H. J. (2017). Circumpolar analysis of the Adélie Penguin reveals the importance of environmental variability in phenological mismatch. *Ecology*, *98*(4), Article 4. https://doi.org/10.1002/ecy.1749

Figures and Tables

Figure 1: Cascading relationship between sea ice concentration (SIC), foraging trip duration and reproductive success for Adélie penguins (*Pygoscelis adeliae*) breeding in Pointe Géologie archipelago, Adélie Land, East Antarctica over the 2010-2024 period. Two contrasting scenarios of access to open water during chick-rearing are apparent: direct access to open water is either possible (*landfast ice absent*, blue) or precluded by the presence of solid landfast ice in front of the breeding colony (*landfast ice present*, yellow). **A)** Annual density distribution of SIC experienced by Adélie penguins during their foraging trips throughout the chick-guard stage. **B)** Annual density distribution of foraging trip duration (n = 23,459 trips) for chick-guarding male and female Adélie penguins from a ca. 250 pairs breeding patch (study colony) located on Île des Pétrels, Pointe Géologie archipelago. **C)** Reproductive success time series for the study colony (2010-2024 average ± SD = 0.66 ± 0.43 chicks per breeding pair). Reproductive success was calculated as the number of chicks fledged per breeding pair and can exceed 1 because Adélie penguins can raise two chicks (Ainley, 2002). **D)** Satellite imagery representative of the absence (top) and presence (bottom) of landfast ice in front of the study colony (red marker). MODIS satellite imagery was downloaded from the NASA Worldview application (https://worldview.earthdata.nasa.gov), part of the NASA Earth Observing System Data and Information System (EOSDIS, top and bottom images from Jan 10th, 2022, and Jan 1st, 2018, respectively).

Figure 2: Relationship between the foraging trip duration (log scale) of chick-guarding Adélie penguins (*Pygoscelis adeliae*) and sea ice concentrations (SIC) for two contrasting scenarios of access to open water: direct access to open water is either possible (*landfast ice absent*, blue) or precluded by the presence of solid landfast ice in front of the breeding colony (*landfast ice present*, yellow). Trip duration data (n = 23,459 trips) was collected using a RadioFrequency Identification System deployed around a breeding patch of ca. 250 pairs within the Pointe Géologie archipelago population (Adélie Land, Antarctica). Lines and associated confidence intervals represent the predictions of the Generalized Additive Mixed Models (GAMM) examined in Table S2. When landfast ice is absent, the 10% lowest trip durations predicted by the GAMM are attained for SIC ranging 8-18% (raw data trip duration mean ± SE = 19.4 ± 0.09 h). Conversely, the predicted 10% longest trip durations occur at high SIC (>76%, raw data trip duration mean ± SE: 47.5 ± 2.17 h), indicating an average 2.5-fold increase between optimal and high SIC. When direct access to open water is precluded by the presence of landfast ice in front of the breeding colony, trip durations are lengthened, averaging 43.9 ± 0.46 h (raw data). This represents an average 1.4-fold increase in trip duration compared to landfast ice absence conditions over the comparable range of SIC (43-84%), and an average 2.2-fold increase compared to the optimal 8-18% SIC in the absence of landfast ice.

Figure 3: Relationship between Adélie penguin (*Pygoscelis adeliae*) reproductive success (chicks/pair) and A) sea ice concentration (SIC) in the foraging area used during chick-guard (pseudo-R² = 0.58); and B) mean foraging trip duration (pseudo-R² = 0.71) of chick-guarding Adélie penguins breeding on Île des Pétrels, Pointe Géologie archipelago, Adélie Land, Antarctica (2010-2024). Black lines and associated confidence intervals are from generalized linear models with a Tweedie distribution. When only years without landfast ice are considered (blue), the relationship between SIC and reproductive success (A) is not statistically significant (p = 0.058, pseudo-R² = 0.25), while the relationship between mean trip duration and reproductive success (B) remains significant (p = 0.003, pseudo-R² = 0.46).

1212

Figure 4: Relationship between Adélie penguin (*Pygoscelis adeliae*) reproductive success and mean foraging trip duration across 10 colonies around Antarctica. The regression lines and associated confidence interval for the general relationship (black line) are from a Generalized Linear Mixed Model (GLMM) including colony as a random effect (marginal $R^2 = 0.74$). Numbers in parentheses after colony abbreviations are colony-specific sample sizes (number of breeding seasons with concurrent reproductive success and trip duration data). Data for Pointe Géologie mainly comes from the present study (n = 15 seasons) but also includes data retrieved from the literature (Barreau et al., 2019; Beaulieu, Dervaux, et al., 2010; Wienecke et al., 2000)Data for all other colonies was retrieved from the literature, and full references and raw data can be found in supplementary text. Abbreviations for colony names: ARDL, Ardley Island (Machado-Gaye et al., 2025); ADMI, Admiralty Bay (Trivelpiece et al., 1990; Watters et al., 2020); HUKU, Hukuro Cove (Watanabe et al., 2020); BECH, Béchervaise Island (Emmerson et al., 2015); PGEO, Pointe Géologie archipelago; HALL, Cape Hallett (Kim et al., 2024); EDMO, Edmonson Point (Clarke et al., 1998; Olmastroni et al., 2000); ROYD, Cape Royds (Ainley et al., 2004, 2006; Saenz et al., 2020; Schmidt et al., 2021); BIRD, Cape Bird (Ainley et al., 2004); CROZ, Cape Crozier (Ainley et al., 2004, 2006; Jennings et al., 2021).

Supporting information

SUPPORTING INFORMATION FOR:

Flexible foraging effort shapes Adélie penguin reproductive success across Antarctica and sea ice conditions

Barracho et al.

Supporting text:

Circumpolar analysis of the effect of trip duration on reproductive success using published data

To upscale our analysis relating chick-guard trip duration to reproductive success from our study to several populations around the species range, we surveyed the literature using Google Scholar with the keywords 'Adélie penguin' and 'trip duration'. For each study for which trip duration data was available, we then looked for concurrent data on reproductive success, either from the same study or from a different study (see Supplementary Table S5). We extracted the reported annual trip duration summary statistics only when the reproductive success data was available for same colony and for the same season. Overall, we obtained 14 different studies to use in our circumpolar analysis.

Because the reporting of trip duration data varied among these 14 studies, we used our own dataset to compare the different methodologies and ensure comparability between our approach and those used by other authors. The differences between our approaches and those found in the literature can be summarized in three main themes:

1. *Use of a different summary statistic*

In one study (Watanabe et al., 2020), the summary statistics used to report annual trip duration was the median. In contrast, and to remain consistent with most other studies, we used the mean. Using our own data to compare mean and median trip duration, we found that *annual medians* were highly correlated to *annual means* (linear model: $\beta = 0.98 \pm 0.03$, $p < 0.001$, $R^2 = 0.99$).

2. *Use of a different method to compute annual mean trip durations (means of means vs. grand mean)*

Mean of means-per-period: In two studies, mean trip durations were reported separately for four (Ainley et al., 2004) to six (Saenz et al., 2020) periods of 5-6 days spanning the entire chick-rearing period. In one study (Ainley et al., 2006), mean trip duration were reported daily. In all three cases, we considered only periods falling during the chick-guard stage and averaged these period-specific means in order to obtain a single annual mean trip duration value. Using our own dataset, we compared this method (i.e. *mean of means-per-period*) to using the *grand mean* over all annual trips. Both methods yielded comparable estimates of annual mean trip duration, with the *mean of means-per-period* being a strong predictor of the *grand mean* both when using 6 equal-sized and successive subsamples within season (linear model: $\beta = 0.99 \pm 0.01$, $p < 0.001$, $R^2 = 0.99$) and when using daily means (linear model: $\beta = 0.99 \pm 0.02$, $p < 0.001$, $R^2 = 0.99$).

Mean of means-per-individual: In one study (Emmerson et al., 2015), individual means were first calculated for each individual (yielding a single mean trip duration value per individual per year) before performing the averaging of individual means, as recommended by Southwell et al. (2006). In contrast, and to ensure direct comparability with most other studies, we calculated mean trip duration using all trips directly (i.e. *grand mean*, without prior within-individual averaging). Using our own dataset, we compared both methods. We found that mean trip duration calculated as the *mean of means-per-individual* was highly comparable to the *grand mean* (linear model: $\beta = 0.95 \pm 0.02$, $p < 0.001$, $R^2 = 0.99$).

Mean of means-per-sex: In three studies (Clarke et al., 1998; Emmerson et al., 2015; Machado-Gaye et al., 2025), mean trip durations were reported separately for males and females. We averaged these sex-specific estimates in order to obtain a single annual mean trip duration value to use in our comparative analysis. Using our own dataset, we compared this method (i.e. *mean of means-per-sex*) to using the *grand mean* over all trips. Both methods yielded comparable estimates of annual mean trip duration, with the *mean of means-per-sex* being a strong predictor of the *grand mean* (linear model: $\beta = 1.03 \pm 0.02$, $p < 0.001$, $R^2 = 0.99$).

3. Data not restricted to the chick-guard stage

In three studies (Jennings et al., 2021; Lescroël et al., 2020; Watanabe et al., 2020), annual trip duration metrics were reported for the entire chick-rearing season (i.e. over both the brooding stage and part of the crèching stage). These studies accounted for 7 out of the 64 total data entries. Because we could not isolate trip duration for the guard-stage (i.e. as we did for (Ainley et al., 2004, 2006; Saenz et al., 2020; Watters et al., 2020), we used these estimates directly in our comparative analysis. Using a version of our own dataset that also included trip duration data during crèching ($n = 25{,}304$ additional foraging trips), we compared annual mean trip duration calculated only for the chick-guard stage to annual mean trip duration calculated for the entire chick-rearing period (chick-guard and crèching). We found that both values were acceptably comparable (linear model: $\beta = 1.33 \pm 0.13$, $p < 0.001$, $R^2 = 0.88$).

When the number of trips used to compute mean trip duration was not reported by the authors, we used the minimal possible value (e.g., number of individuals for which trip duration data is reported, neglecting the possibility that an individual contributes more than one trip to the dataset).

Overall, despite some minor difference in the way trip duration data was reported across studies, we are confident that our approach is reasonable and should not impact the overall model, for which the predicted error is larger than the among-studies differences highlighted here.

Supporting figures and tables:

Figure S1: Foraging area (orange) of chick-guarding Adélie penguins from the Pointe Géologie archipelago, based on tracking data from Widmann et al. (2015), Michelot et al (2021), and Ropert-Coudert et al. (2018). The main map was produced using Quantarctica (Matsuoka et al. 2021) (https://www.npolar.no/quantarctica/).

Figure S2: Relationship between the male-female difference in mean trip duration and Sea Ice Concentration (SIC) for chick-guarding Adélie penguin (*Pygoscelis adeliae*) breeding on Île des Pétrels, Pointe Géologie archipelago, Adélie Land, Antarctica (2010-2024, Linear Model, $\beta = 0.22 \pm 0.03$, $p < 0.001$, $R^2 = 0.77$). The landfast ice categorization indicates whether or not landfast ice was present in front of the breeding colony during chick-rearing, precluding direct access to open water. Higher values of the trip duration difference (y-axis) indicate longer trips by females compared to males.

Figure S3: A) Akaike Information Criterion (AIC) profile of the piecewise regression analysis examining the non-linear relationship between trip duration and reproductive success of Adélie penguins (*Pygoscelis adeliae*) presented in Fig. 4, main text. The red vertical line denotes the most supported breakpoint (29.2 hours). **B) Distribution of bootstrapped breakpoint estimates** with the median (blue vertical line) and the 2.5th and 97.5th percentiles (red vertical lines). We conducted the piecewise regression analysis *a posteriori* to identify the trip duration threshold at which reproductive success begins to change at a different rate with increasing trip duration. Building on model 4 in Table S4, we evaluated 70 candidate breakpoints equally spaced between 10 and 40 hours of trip duration (i.e. ~0.43 hours increment). We discarded models with less than 10 datapoints on either side of the breakpoint. We compared the remaining models using AIC to identify the most strongly supported breakpoint. To estimate a 95% confidence interval around this breakpoint we conducted a nonparametric bootstrap procedure by resampling the full dataset with replacement 500 times. For each bootstrapped sample, we re-estimated the optimal breakpoint by fitting a series of segmented Tweedie GLMMs across 70 candidate breakpoints, as explained above. The breakpoint associated with the lowest AIC was retained for each sample. We then derived the 95% bootstrap confidence interval calculated the 2.5th and 97.5th percentiles of the resulting breakpoint distribution to derive a 95% bootstrap confidence interval.

Table S1: Sample sizes and summary statistics of the sex-specific trip duration data used to link trip duration during chick-guard to sea ice conditions and reproductive success for Adélie penguin (*Pygoscelis adeliae*) breeding on Île des Pétrels, Pointe Géologie archipelago, Adélie Land, Antarctica. N = *sample size*.

Breeding season	N trips	N trips ♂	N trips ♀	N individuals ♂	N individuals ♀	N individuals	Sex ratio trips (males prop.)	Sex ratio individuals (males prop.)	Mean trip duration ± SE	♀ mean trip duration ± SE	♂ mean trip duration ± SE
2010	295	215	80	20	8	28	72.9	71.4	29.2 ± 0.6	31.7 ± 1.3	28.3 ± 0.7
2011	316	220	96	19	9	28	69.6	67.9	27.3 ± 0.7	32.3 ± 1.2	25.1 ± 0.8
2012	87	49	38	8	8	16	56.3	50	45.4 ± 2.1	54.8 ± 3.3	38.2 ± 2.1
2013	473	254	219	37	34	71	53.7	52.1	39.4 ± 1	48 ± 1.4	32.1 ± 1.2
2014	28	10	18	5	10	15	35.7	33.3	72.2 ± 4.4	78.8 ± 5.5	60.5 ± 5.7
2015	274	125	149	24	30	54	45.6	44.4	53 ± 1.4	57.8 ± 1.9	47.3 ± 2
2016	5297	2284	3013	111	147	258	43.1	43	17.4 ± 0.1	18.1 ± 0.1	16.5 ± 0.1
2017	35	20	15	10	9	19	57.1	52.6	34 ± 3.5	47.8 ± 5.9	23.7 ± 2.4
2018	1197	609	588	73	71	144	50.9	50.7	41.3 ± 0.5	44.3 ± 0.7	38.4 ± 0.6
2019	2433	1174	1259	96	99	195	48.3	49.2	23.1 ± 0.2	24.1 ± 0.2	21.9 ± 0.3
2020	2122	1028	1094	111	116	227	48.4	48.9	29 ± 0.2	31.1 ± 0.3	26.8 ± 0.3
2021	4255	2052	2203	110	116	226	48.2	48.7	18.9 ± 0.1	19.6 ± 0.2	18.2 ± 0.1
2022	2899	1486	1413	111	103	214	51.3	51.9	21.6 ± 0.1	22.1 ± 0.2	21.1 ± 0.1
2023	1381	650	731	51	51	102	47.1	50	24.2 ± 0.2	23.9 ± 0.2	24.4 ± 0.4
2024	2367	1148	1219	88	94	182	48.5	48.4	23.6 ± 0.2	24.6 ± 0.2	22.6 ± 0.2
All (2010-2024)	23459	11324	12135	260	305	565	48.3	46	23.4 ± 0.1	24.4 ± 0.1	22.4 ± 0.1

Table S2: Model selection table for the effects of sea ice concentrations (SIC) and sex on the trip duration of chick-rearing Adélie penguins (*Pygoscelis adeliae*) during the guard stage in two scenarios of access to open water for breeding penguins: direct access to open water is either possible (*fast ice absent*) or precluded by the presence of solid landfast ice in front of the breeding colony (*fast ice present*). All models used trip duration as the response variable, included a random individual effect to account for repeated measurements on the same individuals, and used a Gamma error distribution with a log link. Trip duration data was collected at Île des Pétrels, Pointe Géologie archipelago, Adélie Land, Antarctica. Abbreviations: GAMM: Generalized Additive Mixed Model; GLMM: Generalized Linear Mixed Model; adj-R²: percentage of variance explained by the fixed effect component of the mode; AIC: Akaike Information Criterion; ΔAIC: AIC difference between candidate model and model with the lowest AIC within the model set.

Model index	Model type	Sex effect	SIC effect	Interaction SIC×SEX	adj-R^2	AIC	ΔAIC
FAST ICE : ABSENT (*n* = 21,838 trips)							
1	GAMM	yes	Nonlinear smooth	yes	0.16	17039.9	0
2	GAMM	yes	Nonlinear smooth	no	0.16	17050.7	10.8
3	GLMM	yes	Quadratic	yes	0.16	17103.6	63.8
4	GLMM	yes	Quadratic	no	0.16	17129.6	89.7
5	GLMM	yes	Linear	yes	0.15	17489.4	449.6
6	GLMM	yes	Linear	no	0.15	17514.9	475.0
7	GLMM	no	Linear	-	0.13	17521.1	481.2
8	GLMM	yes	-	-	0.01	21473.4	4433.6
9	GLMM	no	-	-	0	21475.6	4435.8
FAST ICE : PRESENT (*n* = 1,621 trips)							
10	GAMM	yes	Nonlinear smooth	yes	0.14	1375.2	0
11	GLMM	yes	Quadratic	yes	0.14	1382.0	6.8
12	GAMM	yes	Nonlinear smooth	no	0.15	1382.4	7.2
13	GLMM	yes	Quadratic	no	0.13	1387.5	12.2
14	GLMM	yes	Linear	no	0.13	1435.6	60.3
15	GLMM	yes	Linear	yes	0.13	1439.6	64.3
16	GLMM	no	Linear	-	0.07	1472.8	97.5
17	GLMM	yes	-	-	0.05	1628.6	253.4
18	GLMM	no	-	-	0	1665.4	290.1

Table S3: Summary of models relating mean trip duration during chick-guard to reproductive success for Adélie penguin (*Pygoscelis adeliae*) at Pointe Géologie archipelago, Adélie Land, Antarctica. The effect of trip duration on reproductive success was modeled using Generalized Linear Models (GLM) with a Tweedie distribution and a log link. Pseudo R^2 were calculated based on the percentage of observed variance in the data explained by the model. AIC: Akaike Information Criterion; ΔAIC: AIC difference between candidate model and model with the lowest AIC within the model set.

Dataset	Mean trip duration calculated over	Tweedie dispersion parameter	Effect size	SE effect size	P value	Pseudo R^2	AIC	ΔAIC
All seasons (n=15)	All individuals	0.046	-0.044	0.006	<0.001	0.73	-22.17	0
All seasons (n=15)	Females	0.049	-0.038	0.006	<0.001	0.78	-5.88	16.29
All seasons (n=15)	Males	0.052	-0.049	0.008	<0.001	0.57	3.83	26
Seasons without landfast ice (n=10)	All individuals	0.039	-0.035	0.007	<0.001	0.48	-15.1	0
Seasons without landfast ice (n=10)	Females	0.046	-0.03	0.008	<0.001	0.48	-9.71	5.39
Seasons without landfast ice (n=10)	Males	0.042	-0.044	0.010	<0.001	0.46	-9.51	5.59

Table S4: Summary of models investigating linear and non-linear effects of foraging trip duration (annual mean) on reproductive success for chick-rearing Adélie penguins (*Pygoscelis adeliae*). All models included a random population effect (n = 10 populations, Fig. 4, Table S5) and a fixed effect of trip duration. Abbreviations: AIC: Akaike Information Criterion. ΔAIC: AIC difference between candidate model and model with the lowest AIC within the model set.

Model index	Model	Fixed effect	Random effect	Fixed effect shape	Weighting	Tweedie dispersion parameter	Conditional R²	Marginal R²	AIC	ΔAIC
1	Reproductive success ~ Trip Duration + (1|Colony)	Mean trip duration	Colony	Quadratic	log(Ntrips)	0.049	0.87	0.74	-39.95	0
2	Reproductive success ~ Trip Duration + (1|Colony)	Mean trip duration	Colony	Linear	log(Ntrips)	0.049	0.83	0.56	-6.9	33.05
Model index	Model	Fixed effect	Random effect	Fixed effect shape	Weighting	Tweedie dispersion parameter	Conditional R²	Marginal R²	AIC	ΔAIC
3	Reproductive success ~ Trip Duration + (1|Colony)	Mean trip duration	Colony	Quadratic	none	0.049	0.83	0.75	26.48	0
4	Reproductive success ~ Trip Duration + (1|Colony)	Mean trip duration	Colony	Linear	none	0.065	0.65	0.49	36.89	10.41

1362 **Table S5: Literature review for concurrent data on Adélie penguin (*Pygoscelis adeliae*) foraging trip duration and reproductive success.**

1363 Full size table with extracted data and references available at: https://figshare.com/s/ef3fe5b9ce965fc6ca13

References (supporting information):

Ainley, D. G., Ballard, G., & Dugger, K. M. (2006). Competition Among Penguins and Cetaceans Reveals Trophic Cascades in the Western Ross Sea, Antarctica. Ecology, 87(8), 2080–2093. https://doi.org/10.1890/0012-9658(2006)87[2080:CAPACR]2.0.CO;2

Ainley, D. G., Ribic, C. A., Ballard, G., Heath, S., Gaffney, I., Karl, B. J., Barton, K. J., Wilson, P. R., & Webb, S. (2004). Geographic Structure of Adélie Penguin Populations: Overlap in Colony-Specific Foraging Areas. Ecological Monographs, 74(1), 159–178. https://doi.org/10.1890/02-4073

Clarke, J., Manly, B., Kerry, K., Gardner, H., Franchi, E., Corsolini, S., & Focardi, S. (1998). Sex differences in Adélie penguin foraging strategies. Polar Biology, 20(4), 248–258. https://doi.org/10.1007/s003000050301

Emmerson, L., Southwell, C., Clarke, J., Tierney, M., & Kerry, K. (2015). Adélie penguin response parameters signal reduced prey accessibility: Implications for predator–prey response curves. Marine Biology, 162(6), 1187–1200. https://doi.org/10.1007/s00227-015-2661-5

Jennings, S., Dugger, K. M., Ballard, G., & Ainley, D. G. (2021). Effects of Diet and Provisioning Behavior on Chick Growth in Adélie Penguins (Pygoscelis adeliae). Waterbirds, 44(1). https://doi.org/10.1675/063.044.0105

Lescroël, A., Lyver, P., Jongsomjit, D., Veloz, S., Dugger, K., Kappes, P., Karl, B., Whitehead, A., Pech, R., Cole, T., & Ballard, G. (2020). Inter-individual differences in the foraging behavior of breeding Adélie penguins are driven by individual quality and sex. Marine Ecology Progress Series, 636, 189–205. https://doi.org/10.3354/meps13208

Machado-Gaye, A. L., Chimienti, M., Kato, A., Ropert-Coudert, Y., Barbosa, A., Bustamante, P., Cosse, M., Menéndez-Blázquez, J., Zaldúa, N., & Soutullo, A. (2025). Energy expenditure of Adélie penguins during the breeding season: Females pay the cost in years of low food availability. Animal Behaviour, 222, 123144. https://doi.org/10.1016/j.anbehav.2025.123144

Matsuoka, K., Skoglund, A., Roth, G., de Pomereu, J., Griffiths, H., Headland, R., Herried, B., Katsumata, K., Le Brocq, A., Licht, K., Morgan, F., Neff, P. D., Ritz, C., Scheinert, M., Tamura, T., Van de Putte, A., van den Broeke, M., von Deschwanden, A., Deschamps-Berger, C., … Melvær, Y. (2021). Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic islands. Environmental Modelling & Software, 140, 105015. https://doi.org/10.1016/j.envsoft.2021.105015

Saenz, B. T., Ainley, D. G., Daly, K. L., Ballard, G., Conlisk, E., Elrod, M. L., & Kim, S. L. (2020). Drivers of concentrated predation in an Antarctic marginal-ice-zone food web. Scientific Reports, 10(1), 7282. https://doi.org/10.1038/s41598-020-63875-y

Southwell, C., Clarke, J., & Emmerson, L. M. (2006). Assessing the power to detect systematic change in Adélie penguin foraging trip duration.

Watanabe, Y. Y., Ito, K., Kokubun, N., & Takahashi, A. (2020). Foraging behavior links sea ice to breeding success in Antarctic penguins. Science Advances, 6(26), Article 26. https://doi.org/10.1126/sciadv.aba4828

Watters, G. M., Hinke, J. T., & Reiss, C. S. (2020). Long-term observations from Antarctica demonstrate that mismatched scales of fisheries management and predator-prey interaction lead to erroneous conclusions about precaution. Scientific Reports, 10(1), 2314. https://doi.org/10.1038/s41598-020-59223-9