Revalidation of *Manis aurita*, based on integrative genomic and morphological evidence

Narayan Prasad Koju[1,2,3,4]#, Zeling Zeng[1]#, Guihua Zhang[1]#, Xia Huang[1]#, Zhicheng Yao[1], Xiaoyun Wang[1], Melissa T.R. Hawkins[5], Arlo Hinckley[5,6], Mary Faith C. Flores[5], Ce Guo[7], Jun Li[7], Devendra Maharjan[8], Saraswoti Byanjankar[9], Lianghua Huang[10], Wenhua Yu[1], Liang Leng[11], Kai He[1*], Anderson Feijó [12 *], Yan Hua[7*]

#: equal contribution; *: corresponding authors

Address:

[1] South China Biodiversity Research Center, School of Life Sciences, School of Life Sciences, Guangzhou University, Guangzhou 510006, China

[2] Center for Postgraduate Studies, Nepal Engineering College, Pokhara University, Lalitpur, 44600, Nepal

[3] Key Laboratory of Genetic Evolution and Animal Models, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650201, China

[4] Department of Psychology, University of Washington, Seattle 98125, USA

[5] Division of Mammals, Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington DC 20024, USA

[6] Departamento de Zoología, Universidad de Sevilla, Seville, 41012, Spain

[7] Guangdong Provincial Key Laboratory of Silviculture, Protection and Utilization, Guangdong Academy of Forestry, Guangzhou, China

[8] Natural History Museum, Tribhuvan University, Kathmandu 44611, Nepal

[9] Aspect Ecology, West Court, Oxfordshire, OX16 2AF, UK

[10] Wildlife Forensic Science Service, Kunming, Yunnan 650201, China

[11] Chengdu University of Traditional Chinese Medicine, Chengdu 611137, China

[12] Negaunee Integrative Research Center, Field Museum of Natural History, Chicago 60605, USA

Details on Corresponding author

Kai He

South China Biodiversity Research Center, School of Life Sciences, Guangzhou University

#230 Waihuanxilu Road, Fanyu District, Guangzhou, Guangdong, China

Email: hekai@gzhu.edu.cn; Phone: 15925188783

ABSTRACT

Pangolins face severe conservation threats globally, necessitating accurate taxonomy for effective protection. Previous genomic research on the Chinese pangolin (*Manis pentadactyla*) identified two deeply divergent lineages (MPA and MPB), suggesting underestimated species diversity. The recent description of *M. indoburmanica* (corresponding to MPB), however, did not assess its relationship with historical names, particularly the subspecies *M. p. aurita*, leaving the group's taxonomic status uncertain. To resolve this issue, we employed an integrative framework, analyzing genomic and morphological data from museum specimens including the lectotype of *M. p. aurita* to clarify phylogenetic relationships and taxonomy within *M. pentadactyla* sensu lato. Our results demonstrate that the *MPB* lineage, which both *M. indoburmanica* and the *M. p. aurita* lectotype, is geographically restricted to the southern Himalayas and thus distinct from other *M. pentadactyla* populations. Genomic analyses indicate the two clades diverged ~1.8 Ma and have remained largely isolated since, with only minimal gene flow. Furthermore, morphometric analyses of both cranial and external features, reveal consistent and significant differentiation between the Himalayan lineage (MPB) and *M. pentadactyla* sensu stricto. Collectively, this congruent evidence unequivocally supports the revalidation of the Himalayan pangolin as a distinct species, *Manis aurita* Hodgson, 1836, representing the fifth extant species in the genus.

Keywords: *Manis aurita*，Integrative taxonomy, Phylogenomics, Morphometrics, Conservation genetics

INTRODUCTION

       Pangolins (Order Pholidota) are unique mammals characterized by their keratinized scales, an extraordinary trait among mammals [1]. These animals face severe threats including habitat loss and extensive illegal trade, which has resulted in dramatic population declines [2-4]. Consequently, all pangolin species are listed under Appendix I of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and are recognized as Evolutionarily Distinct and Globally Endangered (EDGE) species [5]. Accurate species identification, clarification of species distribution, and estimation of population sizes and genetic load are essential for effective conservation efforts [6-8].

       Until recently, eight extant pangolin species were recognized: four in Africa and four in Asia [9]. All Asian species belong to the genus *Manis*, including Chinese pangolin (*Manis pentadactyla*), Indian pangolin (*M. crassicaudata*), Sunda pangolin (*M. javanica*), and Philippine pangolin (*M. culionensis*). Recent genomic studies revealed cryptic genetic and taxonomic diversity within Asian pangolins. For instance, genome sequencing of confiscated scales uncovered a cryptic lineage, provisionally named "mysteria", for which no voucher specimens or precise locality data exist [10]. Moreover, analyses of broad sampling within the Sunda pangolin [11,12] and *M. pentadactyla* [10,13,14] have revealed distinct clades, indicating the existence of an undescribed species.

       The Chinese pangolin is distributed across a broad range, extending from Nepal in the west to the eastern coast of China, with insular populations on Hainan and Taiwan islands. Latitudinally, its range extends from central China southward to Southeast Asia [15]. This broad distribution is divided by geographic barriers, such as the mountains of southwestern China and southward-flowing river systems, which likely drive genetic divergence among geographic populations [16,17]. Accordingly, Hu, et al. [13] identified two major clades within *M. pentadactyla,* namely, MPA and MPB. While recent studies have found additional subclades closely related to MPA in China and Southeast Asia [14,18], and MPB remains the most genetically divergent lineage, and its geographic origin has been enigmatic. Initially, the MPB clade was known almost exclusively from specimens confiscated along the Sino-Myanmar border, suggesting a Myanmar origin. This inference was tentative, as the area is a major wildlife trafficking hub [19-23]. However, recent studies found occurrences of MPB in Nepal, South Tibet and northeastern India [24-27], strongly suggest that this clade represents the western populations inhabiting the South Himalayan region.

       Historically, three subspecies of *M. pentadactyla* were recognized: the nominate subspecies *M. p. pentadactyla* Linnaeus, 1758 described based on specimens from Taiwan; *M. p. pusilla* Allen, 1906 from the Hainan Island; and *M. p. aurita* Hodgson, 1836 from the central Nepal [28]. While the former two have been subject to extensive genetic studies [13,14,18,29,30], and their evolutionary relationships as well as distribution in China and SE Asia has been confirmed [13,14,31], but the taxonomy and evolutionary history of *M. p. aurita* remain unclear.

       Recently, Wangmo, et al. [27] describe a new species, *Manis indoburmanica*, from South Tibet, which highlights several challenges in modern taxonomy. The recognition of this new species is based solely on the mitochondrial genome, a data source susceptible to introgression,

incomplete lineage sorting, and maternal inheritance bias [32-34]. Furthermore, the work lacks a comprehensive morphological comparison with *M. pentadactyla*, thus failing to establish a clear morphological boundary for *M. indoburmanica*. Critically, as noted by Zijlstra [35], Wangmo, et al. [27] did not evaluate whether their samples correspond to existing historical names like *M. p. aurita* [36] or *M. assamensis* [37], leaving the novelty of the taxon in question. Such challenges underscore the necessity of integrative taxonomic approaches that incorporate genome-wide data, robust morphological comparisons with both *M. pentadactyla* and *M. p. aurita* [38-40].

To resolve this taxonomic uncertainty, we employed an integrative taxonomic framework combining genomic and morphological evidence [41], incorporating the type specimens of *M. p. aurita*. The objective of this study was to evaluate whether pangolins from South Himalaya correspond to the subspecies *aurita* and, if so, whether their degree of divergence supports elevation to full species status. Our analyses reveal clear and pronounced differentiation from *M. pentadactyla* across both genetic and morphological datasets, providing robust support for recognizing this lineage as the distinct species *Manis aurita* Hodgson, 1836.

RESULT

Genetic evidence for the distinctiveness of *M. aurita*

We generated seven new NGS datasets for *Manis aurita*, including the *M. aurita* lectotype (BMNH 43.1.12.85) (**Tables S1, S2**), and integrated them with previously published data. The final comprehensive dataset for genomic analysis comprised 55 whole-genome sequencing (WGS) datasets (representing 53 *Manis* individuals, one *Smutsia gigantea*, and one *Phataginus tricuspis;* (**Table S3**) and 70 mitogenomes (**Table S2**). Phylogenetic reconstruction based on the 70 mitogenomes yielded a robustly resolved Maximum Likelihood (ML) tree, with maximal bootstrap support (bootstrap support [BS] = 100) for all interspecific relationships (**Figure 1A**). The topology corroborates all previous findings [10,13,14,18,27], identifying *Manis pentadactyla* sensu lato (encompassing all three subspecies *pentadactyla*, *pusilla*, and *aurita*) as the sister clade to all other Asian congeners. The putative taxon "mysteria" emerges as the immediate sister to *M. javanica.* Within *M. pentadactyla* s. l. (n=44), samples clustered into two deeply diverged clades, corresponding to MPA and MPB identified in Hu, et al. [13]. Autosomal SNPs, sex-chromosome-derived SNPs, and single-copy orthologues (**Figure 1A**; **Supplementary Figures S1**, **S2**) yielded topologies identical to the mitochondrial tree, reinforcing these phylogenetic relationships.

When projected the SNP of *Manis* species onto principal-component (PC) space, every *M. pentadactyla* s. l. sample falls into a tight cluster that is clearly distinct from the other congeneric species (**Figures 1B**, **1C**). Restricting the analysis to *M. pentadactyla* s. l. reveals non-overlapping splitting between MPA and MPB along PC1 (Tracy-Widom test, $p = 7.49 \times 10^{-6}$; **Table S4**), underscoring consistent genomic differentiation despite their close relationship. The result of ADMIXTURE analysis echoes these patterns (**Figure 1D**). The lowest cross-validation error occurs at K = 2 (CV = 0.47), with marginal increases at K = 3 (CV = 0.50) and K = 4 (CV = 0.53; **Table S5**). At K = 2, the inferred clusters correspond closely to MPA and MPB, with three individuals displaying minor ancestry from the opposite cluster. Of important note, in all above phylogenetic and population genetic analyses, the lectotype of *M. aurita* fell in the MPB clade.

Taken together, the concordant mitochondrial and nuclear results robustly validate that MPB represents *M. aurita* and is genetically distinct from the *M. pentadactyla* sensu stricto (MPA) (which represent the other two subspecies).

Figure 1: Phylogenomic relationships and population structure of pangolins. (A) Maximum likelihood (ML) phylogenies based on complete mitochondrial genomes (left) and autosomal nuclear single-nucleotide polymorphisms (SNPs) (right). Numbers at nodes indicate bootstrap support values. Scale bars represent substitutions per site. (B) Principal Component Analysis (PCA) of genomic SNP data demonstrating the separation of four *Manis* species. (C) PCA restricted to the *M. pentadactyla* s.l. illustrating clear genetic differentiation between *M. pentadactyla* s.s. (MPA) and *M. aurita* (MPB). Values in parentheses indicate the percentage of total variance explained by each principal component. (D) Population structure of the *M. pentadactyla* species complex inferred by ADMIXTURE assuming K=2 to K=4. Each vertical bar represents an individual, with colored segments denoting the proportion of ancestry from each inferred cluster. The lectotype of *M. aurita* is marked with a red star.

To explore the geographic distribution of *M. aurita*, we augmented our mitogenome dataset with publicly available georeferenced mitochondrial *CYTB* and *COI* sequences representing *M. pentadactyla* s.l.. This expanded phylogenetic analysis recovered an identical backbone topology (**Figure 2A**). Critically, samples attributed to *Manis indoburmanica* (from South Tibet) were robustly embedded within MPB (BS=100). Furthermore, mapping the occurrences of MPB haplotypes revealed a distribution confined to Nepal, South Tibet, and Assam (**Figure 2B**). Collectively, these lines of evidence demonstrate that *M. indoburmanica* is conspecific with with *M. aurita* and therefore represents a junior synonym of this species, a species whose distribution is restricted to the southern Himalayas.

Figure 2:Mitochondrial phylogeny and geographic distribution of the *Manis pentadactyla* species complex. (A) Maximum likelihood phylogeny constructed from an expanded dataset of mitochondrial sequences (complete mitogenomes and partial *CYTB* and *COX1* fragments), integrating the new sequences generated in this study with publicly available data. Numbers at the nodes represent bootstrap support values. The scale bar indicates the number of substitutions per

site. *M. indoburmanica* specimens (MI, highlighted in orange color) and lectotype of *M. pentadactyla aurita* (BMNH 43.1.12.85) nest firmly within the *M. aurita* (MPB) clade. (B) Geographic distribution map with verified locality information (**see Table S1**). Circles represent data obtained from previous studies while triangles indicate samples collected in this study. Shaded areas denote the approximate ranges of Asian pangolin species according to IUCN Red List.

Morphometric evidence for the distinctiveness of *M. aurita*

To assess whether *M. aurita* is morphologically distinct from the *M. pentadactyla* sensu stricto (including subspecies *pentadactyla* and *pusilla*) and thus merits recognition as a separate species, we employed linear measurements-based and three-dimensional landmark-based geometric morphometric analyses. Principal Component Analysis (PCA) of 20 cranial measurements for 44 specimens positioned *M. pentadactyla* s. l. within a morphospace that showed partial overlap with the Indian pangolin (*M. crassicaudata*) but remained clearly separated from the Sunda (*M. javanica*) and Philippine (*M. culionensis*) pangolins (**Figure 3A**; **Supplementary Figures S3B**, **S4A**, **S4B**). Along PC1, which accounted for 39.4% of the shape variance and primarily reflected overall cranial size, specimens identified as *M. aurita* consistently occupied the positive extreme. This placement indicates that *M. aurita* possesses a significantly larger skull, and proportionally shorter palatine when compared to *M. pentadactyla* s.s. (**Tables S7A-6C**). Three-dimensional landmark-based geometric morphometric analyses of 75 cranial landmarks (**Table S6**) obtained from micro-CT scans also clearly distinguished *M. aurita* from *M. pentadactyla* s.s. along PC3 (**Figure 3B**; **Tables S8A**, **S8B**), but not PC1 or PC2 (**Supplementary Figures S4C**, **S4D**). Compared to *M. pentadactyla* s.s., *M. aurita* exhibits a relatively shorter yet broader nasal bone, a laterally expanded junction between the maxilla and palatine bones, a posteriorly positioned anterior point of the foramen magnum on the ventral occipital surface (landmark 25, **Figure S5**; **Supplementary Table S6**), and an anteriorly shifted M-shaped suture between the maxilla and palatine bones on the ventral view, culminating in a shorter ventral maxilla (**Supplementary Figure S10C**).

To further substantiate the morphological separation between *M. aurita* and *M. pentadactyla* s.s., we conducted a PCA based on fourteen external body measurements including *M. aurita* and *M. pentadactyla* s.s. and *M. javanica*. A crucial aspect of this analysis was the inclusion of the *M. aurita* lectotype, which notably lacks skull material. The results demonstrated a clear distinction: *M. aurita* separated unequivocally from *M. pentadactyla* s.s. along both PC1 (accounting for 52.1% of variance) and PC3 (6.9% of variance) (**Figure 3C**; **Supplementary Tables S9A-9C**, **Figures S4E**, **S4F**). The PCA-based separation is strongly supported by *t*-tests on individual cranial (**Figure 3D**; **Supplementary Figure S6A**; **Table S10**) and external measurements (**Figure 3E**; **Supplementary Figure S6B**; **Supplementary Table S11**), which show *M. aurita* having significantly longer body lengths (89.14±1.65 cm; n=6), back tail lengths (37.39±3.73 cm; n=6), larger hind feet (6.39±0.28 cm; n=6), an extended snout and markedly smaller ears (2.01± 0.05 cm; n=6) (**Supplementary Data S1**).

Finally, Canonical Variate Analyses (CVA) that using both cranial and external variables achieved 100% classification accuracy in separating *M. aurita* and *M. pentadactyla* s.s. (**Tables**

S12, **S13, S14**; **Supplementary Figure S7**; **Supplementary Data S2**). In contrast to these clear differences in skeletal and external body proportions, an analysis of scale characteristics using superimposed, scale-encoded heatmaps of 12 intact body skins (nine *M. pentadactyla* s. s. and *three M. aurita*) revealed no significant differences in either the total number of scale rows or their spatial distribution between the two taxa (**Supplementary Figure S8**; **Supplementary Table S15**; **Supplementary Data S3**, **S4**). Collectively, the consistent and pronounced size- and shape-based differences in both cranial and external morphology, strongly demonstrate that *M. aurita* is morphologically distinct from *M. pentadactyla* s.s. supporting its full species status.

Figure 3: Morphometric analyses based on cranial and external morphology. (A) Principal Component Analysis (PCA) based on 20 linear cranial measurements (PC1 *vs*. PC2). (B) PCA of cranial shape variation derived from 75 3D geometric morphometric landmarks (PC2 *vs*. PC3). (C) PCA based on 14 external body measurements (PC1 *vs*. PC2). For all PCA plots (A-C), values in parentheses on the axes indicate the percentage of variance explained. Note the inclusion of the *M. aurita* lectotype (BMNH: 43.1.12.85) in panel C, which clusters firmly with other *M. aurita* specimens. (D, E) Boxplots comparing key cranial (D) and external (E) measurements between the two taxa. Boxes depict the interquartile range (IQR) with the median indicated by the central line; whiskers extend to 1.5 times the IQR. Individual dots indicate outliers beyond 1.5 × IQR. Units are in millimeters (D) and centimeters (E). Statistical significance from Student's t-tests is denoted as: * *P* < 0.05; ** *P* < 0.01; *** *P* < 0.001. See Tables S5–S14 and Figures S4, S10, S11 for additional results.

Evolutionary and demographic history of *Manis aurita*

We estimated the divergence time within pangolins using 200 selected clock-like single-copy orthologous genes. The most recent common ancestor (MRCA) of the genus *Manis* was dated to approximately 29.4 million years ago (Ma) (95% confidence interval [CI] = 27.4 to 31.5 Ma; **Figure 4A**), a finding consistent with Gu et al. (2023). *M. aurita* (MA hereafter) and *M. pentadactyla* s. s. (MP hereafter) diverged approximately 1.8 Ma (95% CI = 1.3 to 2.5 Ma), during the early Pleistocene.

Pairwise Sequentially Markovian Coalescent (PSMC) analyses revealed distinct demographic trajectories for the two species post-divergence (Figure 4B). MP generally maintained a larger effective population size (N_e) than MA. Following an initial post-split decline, the MP population remained relatively stable from 1.0 Ma until a sharp reduction approximately 50 ka. In contrast, the MA population exhibited demographic stability until ~300 ka, after which it entered a sustained, long-term decline. To circumvent the limitations of PSMC in resolving recent demographic events, we employed GONE to estimate N_e fluctuations over the most recent 600 generations. For MP, a sharp population contraction was detected approximately 60–50 generations ago (corresponding to the 17th century), during which N_e plummeted to 2606 (ranging from 2387 to 2966). In contrast, MA experienced an earlier bottleneck approximately 100-110 generations ago (corresponding to the 15th century), with N_e declining to 1802 (1648-2033) (Supplementary Figure S9).

Given the inferred population declines and reduced contemporary N_e in both species, we evaluated whether these demographic changes were associated with increased inbreeding. (**Figure 4C**), indicating elevated levels of recent inbreeding. Consistent with this pattern, the mean inbreeding coefficient (F_{ROH}) was slightly higher in MP (~0.063) than contemporary MA population (~0.06), However, this difference was not statistically significant (Wilcoxon rank-sum test, $P > 0.05$). Notably several individuals from both species exhibited exceptionally high F_{ROH} (> 0.1), including contemporary MA individuals (Pan06 and PNG1) and MP individuals (GAEM0008, GAEM0009, and GAEM00019). This pattern points to pronounced inbreeding in certain populations such as *M aurita* around Kathmandu Valley.

We subsequently examined patterns of interspecific gene flow involving MA to assess the impact of current and historical range overlap with its congeners (**Figure 2B**). D-suite analyses revealed significant, albeit low-level, introgression from the Indian pangolin (*M. crassicaudata*) into both MP (f4-ratio ≈ 0.0025, p < 1x10-15) and MA (f4-ratio ≈ 0.0022, p < 1x10-15) (**Figure 4E**; **Supplementary Table S16**). Introgression from MA into the Malayan pangolin (*M. javanica*) was also detected using both Dsuite (f4-ratio ≈ 0.0017) and TreeMix (**Figure 4F**). In contrast, analysis of post-divergence gene flow between MP and MA using fastsimcoal2 strongly supported a model of no gene flow over alternative models (ΔAIC > 2.9; **Supplementary Table S17**). Collectively, these results indicate that although MA has experienced minimal introgression from other pangolin species, it has remained largely isolated from its sister species MP since their divergence.

Figure 4: Divergence time, demographic dynamics, and genomic divergence of *Manis* species.
(A) Divergence time estimates for pangolins based on 200 clock-like single-copy orthologous genes, inferred using *BEAST3. Numbers at nodes represent mean age estimates in millions of years (Ma), while the associated horizontal bars represent the 95% Confidence Intervals (CI). Asterisks indicate nodes used for calibration. (B) Historical effective population size (N_e) trajectories for three species inferred using PSMC. Shaded areas represent 100 bootstrap replicates. (C-D) Comparison of recent inbreeding levels between the contemporary population of *M. aurita* (MA contemporary) and *M. pentadactyla* s.s. (MP): (C) Distribution of Runs of Homozygosity (ROH) across individuals in MA contemporary and MP and (D) Genome-wide fraction of ROH (F_{ROH}). Differences between groups were assessed using a Wilcoxon rank-sum test; no statistically significant difference was detected ($P > 0.05$). (E-F) Interspecific gene flow analyses among four *Manis* species. MC: *M. crassicaudata*, MJ: *Manis javanica*: (E) Heatmap of Dsuite f-branch

statistics, where warmer colors indicate stronger signals of admixture on a given branch. (F) TreeMix population graph showing inferred drift and a migration event (arrow). (G-H) Genomic differentiation between *M. aurita* and *M. pentadactyla* s.s. (G) Manhattan plot of F_{ST} and D_{XY} values, with dashed lines indicating the 95% outlier thresholds used to define genomic islands of divergence. (H) Gene Ontology (GO) enrichment analysis of genes within the identified genomic islands, revealing significant enrichment for terms related to olfactory functions. Dot size corresponds to gene count and color to statistical significance (p.adjust).

Finally, we compared MA and MP genomes for signatures of local adaptation by calculating nucleotide diversity (π) and fixation index (F_{ST}), and absolute genetic differentiation (D_{XY}) (**Supplementary Figure S10**). Genomic islands of divergence were defined as windows in the top 5 % for both F_{ST} (>0.8575) and D_{XY} (>0.0373; **Figure 4G**). Gene Ontology (GO) enrichment analysis of genes within these identified islands revealed significant enrichment for olfactory-related functions, including "the detection of chemical stimuli and broader sensory perception of smell" (**Figure 4H**).

DISCUSSION
The Revalidation of *M. aurita*

Despite accumulating molecular evidence suggesting overlooked, or cryptic species [13,14,18], the taxonomy of *M. pentadactyla* s. l. lacked rigorous examination. Our mitochondrial and genome-wide phylogenomic analyses consistently reveal a divergence between *M. aurita* and *M. pentadactyla* s.s. at about 1.8 Ma (**Figure 4A**). The magnitude of this split surpasses commonly accepted thresholds for intraspecific variation [42-44]. Complementing this genetic divergence, morphometric analyses identify consistent and diagnostic differences that distinguish *M. aurita* from *M. pentadactyla* s.s. (**Figure 3**). Cranially, *M. aurita* possesses a significantly larger overall skull with a proportionally shorter and broader nasal region compared to *M. pentadactyla* s.s. (**Supplementary Figures S11**, **S12**; **Supplementary Tables S9**, **S10**). External morphological differences are particularly clear and facilitate differentiation. For instance, *M. aurita* has smaller ears than *M. pentadactyla*. However, Hodgson [36] chose the characteristic reflected in its original specific epithet (*aurita*, Latin for eared) because, compared with other pangolins he was familiar with (e.g., *M. crassicaudata*), this species possessed distinct and conspicuous external ears [35].

The distribution of *M. aurita* is associated with the Himalayan region (**Figure 2B**). It includes the type locality of *M. aurita* from central Nepal, as well as individuals from South Tibet recently described as *Manis indoburmanica* [27]. In contrast, *M. pentadactyla* s.s. is predominantly found in South and Eastern China and Southeast Asia. This geographical separation is likely influenced by biogeographic barriers, such as the rivers run in north-south directions (e.g., the Brahmaputra River) [6,45] and the Arakan Mountains [46].

Although pangolins are capable swimmers [47], major Himalayan landscape features likely impose strong ecological barriers to gene flow rather than purely physical ones. Broad river systems such as the Brahmaputra and steep elevational gradients across the Arakan and Himalayan ranges create discontinuities in forest cover and climatic conditions, limiting dispersal between

suitable habitats. Similar biogeographic filters have been shown to structure genetic divergence in other Himalayan mammals and herpetofauna [6,45,46,48-52]. These combined ecological and climatic constraints may explain the long-term isolation between *M. aurita* and *M. pentadactyla*. Future sampling across eastern Nepal, southern Tibet, and northern Myanmar is recommended to better define potential contact zones and refine the phylogeographic boundaries of *M. aurita*.

Collectively, the congruent evidence from our genetic, morphological, and distributional analyses unequivocally supports recognizing Hodgson's *aurita* as a distinct species. While Zijlstra (2025) previously argued for its revalidation, our study substantiates this view with broader and more integrative evidence. Consequently, in accordance with the Principle of Priority of the International Code of Zoological Nomenclature (ICZN), we reinstate *Manis aurita* Hodgson, 1836 as the valid name for this species. This action renders junior synonyms such as *Pholidotus assamensis* Fitzinger, 1872, and *Manis indoburmanica* Wangmo, Ghosh, Dolker, Joshi, Sharma and Thakur, 2025.

Evolutionary history of *M. aurita*

The population genetic and demographic analyses illuminate the evolutionary trajectory of *M. aurita* following its speciation. The ancestors of *M. aurita* and *M. pentadactyla* diverged in the early Pleistocene, period of, significant global climatic oscillations. Given that pangolins are generally warm-adapted species, with no historical fossil evidence suggesting their presence in northern China [53], the climatic shifts likely played a crucial role in driving the allopatric separation of the ancestor into western (Himalaya) vs. eastern (East/ Southeast Asia) refugium [48-52]. Pangolins are highly olfactory-reliant mammals, exhibiting significantly enlarged olfactory bulb [54,55], using smell to forage for ants and termites [56]. As the *M. aurita* and *M. pentadactyla* adapted to their respective environments, certain OR genes, certain OR genes likely diverged under selection as supported in our GO enrichment analysis (**Figure 4H**).

Our findings highlight the divergent evolutionary responses of MA and MP to Quaternary climatic oscillations. The maintenance of a larger N_e in MP until late in the Pleistocene (~50 ka) suggests that the extensive lowland forests of eastern and southern China served as critical refugia, providing substantial climatic buffering and habitat continuity [57]. Conversely, the demographic collapse of MA beginning around 300 ka coincides with the Penultimate Glaciation (~300–130 ka). During this period, intensified cooling and aridification likely triggered significant habitat contraction[58,59], This pattern aligns with the demographic histories of other warm-adapted mammals in southern China [60], such as the golden snub-nosed monkeys [61]. The result suggests that *M. aurita*, restricted to smaller distribution area, was more vulnerable to habitat fragmentation and aridification.

Interestingly, the recent demographic declines inferred by GONE for both species align temporally with the Little Ice Age (LIA). The 17th-century bottleneck observed in MP coincides with the "Late Ming Little Ice Age" in China. This era was characterized by severe cooling and megadrought [62,63] which has been linked to agricultural failure and the eventual collapse of the Ming Dynasty [64]. Similarly, the demographic contraction of MA around the 15th century

corresponds to the onset of the LIA in the Himalayan region [65,66]. Our findings underscore that even relatively recent, short-term climatic oscillations like the LIA can leave a discernible and lasting footprint on the genomic diversity of pangolins.

Our analyses further detected limited gene flow between *M. aurita* and two other Asian pangolins. The signal of introgression between *M. aurita* and *M. crassicaudata* (contributing to <0.25% of the genome) is unsurprising, given their sympatric ranges in southern Nepal and northeastern India. We also observed a similarly low level of gene flow between *M. aurita* and *M. javanica*, indicating that their distributions likely have overlapped at least historically. Despite these instances of historical admixture, the low level of gene flow indicates *M. aurita* has maintained its status as a distinct evolutionary entity.

It is noteworthy, in the admixture analysis, at both K=3 and K=4, *M. aurita* separated into two distinct genetic clusters: one comprising individuals from Nepal and the other comprising individuals from South Tibet and northeastern India, a pattern that indicates significant intraspecific subdivision (**Figure 1D**).

Conservation Implications

The formal recognition of *M. aurita* as a distinct species carries profound conservation significance. Its intrinsic vulnerability is underscored by a geographically restricted range, confined to the Himalayan–Indo-Burmese region, and a demographic history marked by a more pronounced and sustained population decline since the Middle Pleistocene compared to *M. pentadactyla*.

Despite the long-term population decline revealed by PSMC analyses, *M. aurita* on average exhibits lower levels of recent inbreeding than *M. pentadactyla* as evidenced by the results of ROH, F_{ROH} (**Figures 4C**, **4D**) and GONE (**Supplementary Figure S9**). This likely reflects differing histories of anthropogenic pressure nowadays; the less intensively developed landscapes of South Asia historically provided more contiguous refuge habitats compared to the range of *M. pentadactyla* in China and Southeast Asia, where has experienced intense human activity for centuries, leading to severe habitat fragmentation and the regional extinction of many medium-to-large mammal species [67-69]. However, it is noteworthy, *M. aurita* around Kathmandu valley exhibit extremely high inbreeding levels (i.e., $F_{ROH} > 0.1$), indicating that these populations are already facing severe inbreeding depression (**Figure 4D**). Conservation strategies must therefore prioritize the identification and management, and continue search for other locally vulnerable populations to prevent local extinctions.

Overarching these intrinsic vulnerabilities is the critical threat from illegal wildlife trade. Genetic analyses of confiscated scales have repeatedly confirmed that *M. aurita* is a target [13]. Alarmingly, our findings also reveal that products from this species have infiltrated regulated traditional medicine markets (data from Xie, et al. [70]; **Supplementary Figure S13**), demonstrating that illegally sourced materials are being laundered through formal supply chains. The comprehensive genomic dataset presented here, with its improved geographic coverage and inclusion of previously unsampled populations, provides an enhanced genetic baseline which will

bolster wildlife forensics by allowing for greater precision in determining the geographic provenance of confiscated scales, thereby helping to pinpoint poaching hotspots.

The recognition of *Manis aurita* as a distinct species also has important implications for its legal protection. As CITES and IUCN currently lists only the eight previously recognized pangolin species individually under Appendix I, and categorized as Endangered or Critically Endangered species. *M. aurita* remains temporarily unlisted. This creates a potential regulatory gap that may expose them to illegal trade risks. We therefore recommend that *M. aurita* be formally included under CITES Appendix I and listed as Critically Endangered on the IUCN Red List to ensure consistent international protection along with other pangolin species.

Taxonomy

Manis aurita Hodgson, 1836 stat. nov.

Manis auritus Hodgson, 1836. Original description.

Manis aurita Waterhouse,1838

Pholidotus assamensis Fitzinger, 1872

Manis pentadactyla auritus Thomas 1918

Manis pentadactyla Ellerman & Morrison-Scott, 1951 (part)

Manis indoburmanica Wangmo, Ghosh, Dolker, Joshi, Sharma and Thakur, 2025

Lectotype: BMNH 43.1.12.85, an adult (skin) collected by Brian Houghton Hodgson in 1836 from Central Nepal (**Supplementary Figure S12F**).

Emended diagnosis

Manis aurita is a medium-sized Asian pangolin from the South Himalaya; with a total length longer than *M. pentadactyla* but shorter than *M. crassicaudata.* The species is characterized by a flap-shaped external pinna, which is smaller than *M. pentadactyla* (**Supplementary Figures S11A**, **S11E-F**)*.* The rostral scales do not reach the tip of the snout, leaving a distinct patch of exposed, naked skin (typically connected to the nasal tip in *M. crassicaudata and M. culionensis*) (**Supplementary Figure S11A**). The dorsal and ventral scales of *M. aurita* are uniformly dark brown to deep chestnut (overall yellowish-brown in *M. crassicaudata* and pale yellow on the abdomen and distal limbs in *M. javanica*). The medial rows of dorsal scales are broadly trapezoidal with straight posterior margins (typically broad rhombic shape in *M. pentadactyla* and *M. crassicaudata*, and elongated rhombic and sharply pointed in *M. javanica* and *M. culionensis*) (**Supplementary Figure S11B**).

Cranially, *M. aurita* is distinguished by a shorter palatal process of the premaxilla (extending less than half the premaxillary length), in contrast to *M. crassicaudata*, and *M. culionensis*, where the process exceeds half the length (**Supplementary Fig. S11C, S12B**). The incisive foramen is oval-shaped and enclosed entirely within the premaxilla, neither extending to the posterior border nor contacting the ventral projection of the maxilla, whereas in *M. crassicaudata* and some specimens of *M. javanica*, the incisive foramen is formed by the premaxilla (anteriorly/laterally) and the maxilla's palatine processes (posteriorly). The maxillae in *M. aurita* are inflated, especially anteriorly where it meets the premaxilla, resulting in a more robust rostral profile; by contrast, in the other four *Manis* species, the rostral region appear

slenderer (**Supplementary Figures S11C**, **S12A**). The postglenoid foramen opens on the lateral aspect of the squamosal zygomatic process (whereas it opens on the posterior aspect in *M. javanica* and *M. culionensis*) (**Supplementary Figures S11D, S12D**). The jugular and hypoglossal foramina open within a single depressed fossa (whereas in the other four *Manis* species, these foramina are not recessed) (**Supplementary Figures S11C**, **S12C**).

Description

Manis aurita has an elongated head, small ears, short limbs, and a tail shorter than half the body length. Its external pinna is small and flap shaped. The head and body are covered in overlapping, exceptionally smooth keratin scales: dorsal scales are broadly trapezoidal with straight posterior margins, while lateral abdominal scales are broad and shield shaped. On the snout, the first row of head scales lies well behind the nasal tip, leaving a bare patch of skin. Scale coloration in adult's deep chestnut dark brown; juveniles are slate-grey or greyish brown, with more bristles between scales.

The skull presents a robust overall profile. The rostrum is notably thickened due to the inflated maxillae, particularly at their junction with the premaxilla. The ventral palatal process of the premaxilla is short, extending less than half the length of the premaxilla. The incisive foramen is oval and situated entirely within the premaxilla, without reaching the posterior border or contacting the ventral projection of the maxilla. In ventral view, the vomer is not visible. From a dorsal perspective, the nasal bones are broad and extend posteriorly to the parietal, terminating in an arch-like shape and sharing a broad suture with the maxilla. The braincase is dorsally inflated, resulting in a dome-shaped appearance.

The postglenoid foramen opens laterally on the zygomatic process of the squamosal, a condition recognized as an apomorphy for Pholidota [71]. In the zygomatic region, the maxillary process either connects to the squamosal via an ossified, rod-like structure or remains detached— an apomorphic trait in certain lineages [71]. The pterygoid does not extend beyond the posterior margin of the tympanic bulla. The jugular and hypoglossal foramina open within a single depressed fossa on the ventral surface of the cranium. External and cranial measurements are provided in Supplementary Data S1 and S2, respectively.

The total scale count ranges from 538 to 596. The trunk, spanning from the neck to the tail base, bears 254–274 scales arranged in 23–24 longitudinal and 15–17 transverse rows. The tail (caudal) scales total 140–153, 80–94 on the dorsal surface organized into five transverse by 16–20 longitudinal rows, and 50–63 on the ventral surface in three to five transverse by 18–20 longitudinal rows, with 17–19 scales along each lateral edge. Unilateral scale counts for the forelimbs and hindlimbs range from 37–55 and 32–41, respectively.

Comparisons

Manis aurita (MA) and *M. pentadactyla s.s.* (MP) exhibit a close overall morphological resemblance. Both species share a similar number of body scales, which are typically uniform dark brown to deep chestnut in coloration. Along the midline of the tail and the dorsal surface of the body, the scales are predominantly trapezoidal in shape, whereas those on the limbs and along the ventral margins are generally shield-shaped. This contrasts with *M. javanica*, which features pale

yellow scales the distal portion of the tail and ventral side and a unique reversed scale growth on the forelimbs. *M. crassicaudata* is easily distinguished by its significantly larger, earth-toned scales and sparse bristles, while *M. culionensis* exhibits more numerous, elongated, and leaf-shaped scales with sharper apices (**Supplementary Figures S11G-J**, **S12F**).

In MA and MP, the maxillary zygomatic process and the squamosal zygomatic process are sometimes connected by a slender, rod-like bony bridge, forming a complete zygomatic arch. This condition is also occasionally observed in *M. javanica*. By contrast, a complete zygomatic arch is consistently absent in *M. crassicaudata* and *M. culionensis* [30]. Additionally, the maxillary and squamosal zygomatic processes in MA and MP are generally positioned much closer to each other than in *M. javanica* and *M. culionensis*, in which the two processes are noticeably more widely separated (**Supplementary Figure S12D**). The position of the incisive foramen varies among species. In *MA*, *MP*, and *M. culionensis*, the incisive foramen is consistently enclosed entirely within the premaxilla and does not contact the ventral projection of the maxilla. By contrast, in *M. crassicaudata* the incisive foramen is formed anteriorly and laterally by the premaxilla and posteriorly by the palatine processes of the maxilla. *M. javanica* exhibits intraspecific variation and displays both conditions (**Supplementary Figure S12B**).

The palatal process of the premaxilla in MA and MP is notably short, extending less than half the total length of the premaxillary length. This condition contrasts with other *Manis* species, in which the palatal process is comparatively longer [30] (**Supplementary Figure S12B**). Additionally, the postglenoid foramen in MA and MP opens on the lateral aspect of the squamosal zygomatic process. This differs from *M. javanica* and *M. culionensis*, the foramen opens on the posterior aspect. In *M. crassicaudata*, the postglenoid foramen may open either posteriorly or laterally (**Supplementary Figure S12D**).

Among all species, only some specimens of *M. crassicaudata* show the vomer visible on the ventral surface of the skull. Most notably, the jugular and hypoglossal foramina of MA open within a single depressed fossa on the ventral surface of the cranium. Although both foramina open into the same recessed area, they remain clearly distinguishable from one another. In all other Asian pangolin species, these foramina are not recessed and open separately on the ventral surface of the skull (**Supplementary Figure S12C**).

Despite their overall similarity, MA can be unequivocally distinguished from MP through several robust morphometric and anatomical characters. Compared with MP, MA typically attains a larger adult body size, with a total length of 81.0–105.1 cm (mean = 95.2 cm) versus 60.5–80.6 cm in MP (mean = 71.2 cm). MA also possesses a longer tail (31.7–40.9 cm, mean = 37.4 cm) than MP (22.3–31.6 cm, mean = 27.5 cm) and has larger foot surfaces, with hindfoot lengths averaging 6.4 cm compared with 2.9 cm in MP. The external pinnae further differ between the two species, being distinctly smaller in MA than in MP (**Figure 3E**; **Supplementary Figures S6B**, **S11E–F**). The skull of MA is also larger and taller than that of MP, with cranial width ranging from 93.2 to 99.7 mm (mean = 95.6 mm) and height from 35.2 to 36.3 mm (mean = 35.8 mm), compared with MP, which shows a cranial width of 75.9–95.9 mm (mean = 84.6 mm) and a height of 23.8–31.5 mm (mean = 27.6 mm). This difference is further reflected in nasal proportions: MA

has a relatively smaller nasal length-to-width ratio (approximately 1.9:1), indicating broader nasal bones, whereas MP exhibits a higher ratio (approximately 2.5:1) and a comparatively more slender nasal region. In addition, the maximum width of the gap between the alveolar sulci is greater in MA (13.5–14.7 mm, mean = 13.8 mm) than in MP (10.1–12.0 mm, mean = 10.9 mm) (**Figure 3D**; **Supplementary Figure S6A**). At the anterior end of the skull, the maxilla in MA bulges laterally, producing a broader maxillary region in ventral view and a broader anterior skull profile in dorsal view, whereas this region appears distinctly narrower in MP (**Supplementary Figures S12A**, **S12E**). The palatine bone in MA is also positioned more anteriorly than in MP: the most posterior point of the palatine sutures in MA aligns approximately with the junction of the maxillary and squamosal zygomatic processes, whereas in MP this point lies posterior to that junction.

Distribution, habitat, and Conservation Statues

Manis aurita is distributed along the southern Himalayan foothills, with confirmed occurrences in Nepal, South Tibet and northeastern India (including Assam), southern Bhutan, and extending into southern Tibet. The species' altitudinal range spans from lowland areas to approximately 3,000 meters above sea level [72,73]. The eastern extent of its distribution remains undefined, but the Arakan Mountains appear to represent the eastern geographic limit for this species [46]. Further surveys are necessary to ascertain its potential presence in Myanmar.

Manis aurita predominantly inhabits mixed deciduous and evergreen forests. Key tree species in these habitats include *Shorea robusta, Schima wallichii, Castanopsis indica*, and *Alnus nepalensis* [74] [75]. *M. aurita* is also frequently observed in forested areas adjacent to human-modified landscapes, indicating some level of adaptability to disturbed environments [73-76].

Despite protected under national laws such as the Nepal's National Parks and Wildlife Conservation Act (1973), India's Wildlife Protection Act (Schedule I, 1972), and internationally under CITES Appendix I, *M. aurita* remains a significant target for illegal wildlife traffickers in South Asia. For example, Kathmandu functions as a major transit hub for smuggled pangolin scales [19,21]. Similar trafficking pressures are evident in neighboring India and Bhutan [20], where challenges such as limited enforcement capacity and porous international borders facilitate cross-border smuggling operations.

MATERIAL AND METHODS

Taxon sampling and sequencing

DNA was extracted from tissue samples (n=21) and purified using the Dneasy blood and tissue kit (Qiagen). Genomic DNA was sheared to ~350 bp fragments using a Covaris LE220R-plus system (Covaris, USA), followed by end-repair, A-tailing, and adapter ligation. Libraries were amplified by PCR, purified with AMPure XP beads.

Destructive sampling was conducted using skin specimens deposited in the National Museum of Natural History, the Smithsonian Institution (USNM), Field Museum of Natural History (FMNH); American Museum of Natural History (AMNH) and University of Michigan, Museum of Zoology (UMMZ) in the United States. Sterile practices were employed with routine bleaching, instrument sterilization, and changing of PPE to prevent any cross contamination

between specimens. Samples of 25 mg of material were placed in sterile cryovials or 1.5mL tubes and transferred to a dedicated Historic DNA processing facility at the Smithsonian Institution's Museum Support Center and Natural History Museum's sequencing laboratory where DNA extraction was performed via Qiagen MinElute or Investigator kits following the manufacturer's protocol. Single-stranded library preparation was performed with the SRSLY Pico Plus Kit (Claret Biosystems®) utilizing half reactions. Libraries had dual indexed iTru style adapters ligated to each sample and amplified for 18 cycles of PCR. Following amplification, samples were visualized on a High-Sensitivity Tape Station (Agilent) and pooled in equimolar ratios for sequencing. Sequencing was performed on an Illumina NovaSeq X platform to generate 150 bp paired end (PE150) reads. Raw FASTQ reads were processed with fastp v0.24.0 [77] to remove adapter contamination and poly-G tails, trimming bases with Phred score ≤ 5. Reads containing > 10 % ambiguous bases or > 50 % low-quality bases were discarded. We supplemented our dataset with 34 available genome re-sequencing data (**Supplementary Table S2**).

Mitochondrial assembly and phylogenetic analyses

Mitochondrial genomes were assembled from 65 raw sequencing datasets, including data from 31newly sequenced individuals and 34 previously published genomes of various pangolin species. For these datasets, clean reads were processed for *de novo* assembly and annotation in MITOZ v3.6 [78], which employed Megahit with k-mer values of 21, 33, and 55. This final set of assembled mitogenomes was then augmented with five additional mitogenomes obtained from GenBank (**Supplementary Table S2**). We aligned mitogenomes using MAFFT v7.5 [79], and trimmed poorly aligned regions using Gblocks. We retained 12S, 16S rRNA, and 12 coding genes on the heavy chain for phylogenetic analyses. We also downloaded available mitochondrial 18 *CYTB* and 12 *COI* genes identified as *M. pentadactyla* from GenBank. We used IQ-TREE v2.1.3 to estimate the best-scoring Maximum-likelihood (ML) trees. ModelFinder was used to automatically determine the optimal partitioning scheme and substitution models under the Bayesian Information Criterion (–m MFP+MERGE). Node support was evaluated with 1,000 ultrafast bootstrap replicates.

Genomic dataset and phylogenetic analyses

We aligned clean data to a *M. pentadactyla* reference genome (GCF_030020395.1) using BWA v0.7.17. The resulting SAM files were converted to coordinate-sorted BAMs and indexed with SAMtools v1.10 [80]. Duplicates were identified and flagged using MarkDuplicates tool of GATK v.4.6 [81]. Local realignment around indels was then performed using GATK's IndelRealigner to reduce false-positive variant calls. Per-sample gVCFs were generated with GATK's HaplotypeCaller and merged via CombineGVCFs. SNP calling was then executed with GenotypeGVCFs and SelectVariants to produce multisample VCF and isolate high-confidence SNP calls, respectively. We filtered SNP calls using GATK's VariantFiltration module with thresholds: QUAL < 30.0, QD < 2.0, MQ < 40.0, FS > 60.0, SOR > 3.0, MQRankSum < –12.5, ReadPosRankSum < –8.0, and SB ≥ –1.0. We applied VCFtools v0.1.16 to remove SNPs with minor-allele frequency < 2 % or depth outside the 2.5–97.5 percentile of the depth distribution [82].

Phylogenetic relationships were inferred using three datasets: high-quality single nucleotide polymorphisms (SNPs) for autosomal and sex chromosomes as well as single-copy orthologous genes. For the SNP-based analyses, autosomal and sex chromosome SNP data were separately extracted from the filtered, high-quality multi-sample VCF. These datasets were converted to PHYLIP format using vcf2phylip v2.8. ML trees were constructed for both datasets using IQ-TREE v2.1.4 as mentioned above. For the ortholog-based analysis, we first generated consensus sequences by applying bcftools consensus to high-quality SNP data using the reference genome. Based on the reference annotation file, AGAT was employed to extract the CDS sequences, which were then translated into protein sequences using EMBOSS transeq. We subsequently identified 23,752 single-copy orthologs using OrthoFinder v2.5.4 [83]. These orthologs underwent stringent filtering based on GC content, sequence variability, proportion of missing data, and ambiguous bases, resulting in 7,852 high-quality loci. Each loci was individually aligned and a concatenated ML tree was constructed using IQ-TREE as mentioned above.

Genetic structuring analyses

To investigate population genetic structure, we utilized linkage-disequilibrium (LD)–pruned autosomal SNPs. LD pruning was performed in PLINK v1.90b6.21 using the –indep-pairwise 50 5 0.2 parameters [84]. We then performed PCA on mean-imputed genotypes in PLINK to extract the top ten principal components. To assess the statistical significance of the principal components, we applied the Tracy-Widom test implemented in EIGENSOFT v8.0.0 [85]. We applied PCA to all *Manis* samples and to *M. pentadactyla* sensu lato (which encompasses individuals later identified as *M. aurita*). To infer individual ancestry proportions, we ran ADMIXTURE v1.3.0 on the *M. pentadactyla* s.l. samples. We evaluated models for K (number of ancestral populations) from 1 to 10, and selected the optimal K to minimize cross-validation error [86].

Molecular dating and population genetic analyses

Divergence times were estimated using *BEAST3 model [87] as implemented in BEAST v2.7.7 [88]. First, we identified the 2,000 most genetically divergent loci from an initial pool of 7,852 filtered orthologs. These loci were then assessed for their conformity to a strict molecular clock using SortaDate v2.0 [89]. The 200 orthologs that best fit the strict clock model were subsequently used for the divergence time estimation. The Bayesian inference employed a Birth–Death tree prior, a strict molecular clock, and the HKY+Γ substitution model. We employed two secondary calibrations derived from the molecular dating estimates of Gu, et al. [10]. The analysis was constrained by setting priors on i) the crown-group pangolin divergence, set to 33.19 million years ago (Ma) (95% CI: 29.8–30.3 Ma), and ii) the divergence of Asian pangolins, set to 16.9 Ma (95% CI: 12.9–22.3 Ma), and iii) the outgroup divergence, modeled as an exponential prior with an offset of 5 Ma [90] and a mean of 1.665. The Markov chain Monte Carlo (MCMC) analysis was run for 300 million generations. Convergence and parameter mixing were assessed in Tracer v1.7.2 [91], ensuring all key parameters achieved effective sample sizes (ESS) > 200.

Historical population dynamics were reconstructed using the Pairwise Sequentially Markovian Coalescent (PSMC) model [92]. According to the estimated divergence time between *M. pentadactyla* and *M. aurita* (see Results), we estimated the mutation rate per generation assuming

an average generation time of 7 years [93]. The analysis was run with the parameters -N25 -t15 -r5 -p "4+25*2+4+6", using the Manis *pentadactyla* genome [13] as the reference for mapping. To assess the variability in demographic inferences, 100 bootstrap replicates were performed.

Levels of inbreeding were quantified by calculating the average inbreeding coefficient (F_{ROH}) for each individual. Long runs of homozygosity (ROH) were identified using PLINK v2.0 [84] with the parameters --homozyg --homozyg-window-snp 20 --homozyg-kb 100. Individual F_{ROH} values were then calculated as the proportion of the autosomal genome covered by ROH, relative to the total autosomal SNP coverage, following McQuillan, et al. [94]. Because DNA from museum specimens may be degraded and highly fragmented, the detection of long ROHs may be reduced, potentially influencing F_{ROH} estimates.

To explore the long-term demographic reconstructions obtained from the PSMC analyses, we inferred recent demographic histories (last ~600 generations) of *M. aurita* and *M. pentadactyla* using the linkage-disequilibrium approach implemented in GONE [95]. For this analysis, we used the same whole-genome autosomal SNPs dataset filtered for high genotype quality and site callability. To assess robustness, we generated 100 independent replicate datasets through random subsampling, capping each chromosome at 100,000 SNPs. Each replicate was analyzed using identical GONE settings (cMMb = 1, maxNSP = 50,000, hc = 0.05).

Geneflow analyses

To infer the direction and magnitude of historical gene flow among *Manis* species, we analyzed a high-quality SNP dataset from 52 samples representing the four *Manis* species using Dsuite v0.5 [96] and TreeMix v1.13 (Pickrell & Pritchard, 2012). We first filtered the SNP dataset to retain sites with ≤20% missing data using VCFtools (--max-missing 0.8). To quantify admixture, we used the Dtrios program in Dsuite to calculate D-statistics and f4-ratio statistics for all relevant population trios. *P. tricuspis* and *S. gigantea* were used as outgroups. Results were visualized with the dtools.py script. To model population splits and migration events, the filtered SNP data was first pruned for linkage disequilibrium (PLINK; --indep-pairwise 50 5 0.6), and the output was converted to TreeMix format using plink2treemix.py. We then ran TreeMix, exploring models with 0 to 5 migration edges (five replicates each). The optimal model was selected by evaluating the explained variance and residual plot fits. The resulting population graph was visualized using the R script plotting_funcs.R provided with TreeMix.

We used FastSimCoal v2.8 [97] to test potential gene flow between *M. pentadactyla* (MP) and *M. aurita* (MA). First, we used VCF file through easySFS to generate folded observed joint site frequency spectrum (JSFS). We constructed four demographic models to test gene flow scenarios: (1) early gene flow (range from 120,000 to 130,000 generations); (2) recent gene flow (range from 1,000 to 20,000 generations); (3) continuous gene flow; and (4) no gene flow. For each tested demographic model, we performed 100 independent runs using 100,000 simulations, 40 expectation–maximization cycles, and a broad search range for parameters to determine the run with the best parameter estimates and maximum likelihood. Model comparisons were performed

using the Akaike Information Criterion (AIC) to identify the model that best explains the observed data.

Genomic Island of Differentiation

To identify genomic regions under divergent selection, we scanned the genomes of *M. pentadactyla* and *M. aurita*. We used Pixy v2.0 to calculate nucleotide diversity (π), fixation index (F_{ST}), and absolute genetic differentiation (D_{XY}) across 20 kb windows [98]. Candidate genomic islands of divergence were defined as outlier windows concurrently in the top 5th percentile for both F_{ST} and D_{XY}. We extracted all genes within these outlier regions. Genes annotated only with LOC identifiers were annotated based on their high-confidence human orthologs, identified via BLASTP [99]. This gene set was then tested for Gene Ontology (GO) term enrichment using the R package clusterProfiler [100].

Morphometric analyses

We accessed specimens deposited in FMNH, USNM, UMMZ, British Museum of Natural History, UK (BMNH), Natural History Museum, Tribhuvan University (NHM-TU) Institute of Zoology, Guangdong Academy of Sciences, China (IOZ-GDAS), and Guangdong Wildlife Monitoring and Rescue Center, China (GWMRC) (**Supplementary Table S1**). Fourteen external measurements were taken including full body length (abdominal) (FBLA), head body length (abdominal) (BLA), tail length (abdominal) (TLA), full body length (back) (FBLB), head body length (back) (HBL), tail length (back) (TLB), external pinna length (EL), external pinna thickness (ET), middle claw length of the back foot (MCLBF), middle claw length of the forefoot (MCLFF), the distance of the first scale from the tip of the nose (DFSTN), the maximum number of scales in the back of the body (MNS), the single-sided number of tail edge scales (SNTE), length of hind foot (LHF), using soft tape rulers (accurate to 0.1 mm) for flexible dimensions and MNT-150 standard-type zinc alloy digital caliper (Model: MNT952101; DEGUQMNT, accuracy to 0.01 mm) for more precise linear features (**Supplementary Figure S3A**). This data includes 26 skin specimens for *M. pentadactyla* (n=8), *M. javanica* (n=12), *M. aurita* (n=6) and also includes the lectotype of *M. aurita* (**Supplementary Data S1**).

Twenty cranial measurements were conducted with MNT-150 standard-type zinc alloy digital caliper (accuracy to 0.01 mm), following [30,101-105] including total skull length (TSL), nasal to foramen magnum base length (NFBL), length from premaxilla to palatine (PP), from Intersection between maxilla and palatine sutures to most posterior point of the palatine sutures(IMPS), tymapanic bulla (left) (TB), skull width (SW), skull height (SH), maximum width of nasal bone (WN), maximum length of nasal bone (LN), dorsal gap between maxillary (dorsal) (GM), maximum width of gap between alveolar sulci (MWG), left maximal length of orbit left) (MO), distance between tympanic bulla (DB), length of suture between parietal bone (DTEM), length of suture between frontal bone (DLOB), Length of the zygomatic process between maxilla and squamous bone (left) (ZMS), length of foremen magnun occipitals (FMO), mandible length (left) (ML), the distance between condylar process (DCP), length of palatal process of premaxilla(LPP) (**Supplementary Figure S3B**). This data includes 44 skulls for *M. pentadactyla* (n=25), *M. javanica* (n=7), *Manis crassicaudata* (n=2), *M. culionensis* (n=3) and *M. aurita* (n=7;

Supplementary Data S2). Sex was not considered in the analysis due to the absence of marked sexual dimorphism and the lack of information for some specimens. Specimen age was assessed based on body size, skull morphology, and available collection records. Juvenile and subadult individuals were excluded in the analysis. To ensure completeness of the data, we replaced missing values with the mean value calculated across specimens of the same species following Arbour and Brown [106]. We used the function prcomp () of stat to conduct principal component analysis (PCA) and Morpho to perform canonical variate analysis (CVA) on log_{10}-transformed variables. Finally, we conducted Levene's test for homogeneity of variances using the leveneTest () function and then performed Student's t-tests with t.test () to compare group means across taxa.

We downloaded 84 micro-CT scan data from Ferreira-Cardoso, et al. [107], and scan 13 specimens, obtaining high-resolution micro-CT scans of 97 skulls for *M. pentadactyla* (n = 42), *M. javanica* (n = 38), *M. crassicaudata* (n = 12), *M. culionensis* (n = 3), and *aurita* (n = 2; **Supplementary Table S2**). Raw DICOM stacks were imported into Materialise Mimics v19.0 for segmentation and construction of 3D cranial meshes. We placed 75 landmarks per mesh following Ferreira-Cardoso (2020; **Supplementary Figure S5**). After Procrustes alignment, we conduct principal component analysis (PCA) and canonical variate analysis (CVA) using MorphoJ v1.08 [108]. Landmark selection was based on prior studies addressing mammalian taxa [109,110]

Scale coding

We generated scale-encoded heatmaps from 12 intact skins (nine *M. pentadactyla,* three *M. aurita*) to visualize variation following Algewatta, et al. [111]; Guo, et al. [112] and Martin, et al. [113]. In brief, we recorded the x-y coordinates of every scale in Excel, exported standardized images to Photoshop v25, and overlaid all specimens within each species to generate species-specific composite maps (**Supplementary Data S3 and S4**).

DATA ACCESSIBILITY

All sequencing data were deposited in the National Center for Biotechnology Information (NCBI) database (BioProjectID PRJNA1332899). Sample metadata, including individual identifiers and accession numbers, are summarized in **Supplementary Table S2**. The analysis scripts and source data for the main figures are available on Dryad repository with the identifier Dryad http://datadryad.org/share/tUnpoAtBaYbC-LxLmopIeUXwwngHX96olu7F6k_uTBk[114]

SUPPLEMENTARY DATA
Supplementary data to this article can be found online.

COMPETING INTERESTS
The authors declare that they have no competing interests

AUTHOR CONTRIBUTIONS
NPK and KH: Conceptualization, Methodology, Investigation, Resources, Visualization, Validation, Supervision, Funding acquisition, Project administration, Writing – review & editing. All authors read and approved the final version of the manuscript. **ZZ, GZ, XH, AF, and YH**:

Methodology, Investigation, Formal analysis, Visualization, Validation, Writing – original draft, Writing – review & editing. **ZY, XW, JL, DM, SB, LH, WY, and LL** Methodology, Investigation, Formal analysis, Visualization, Validation. **MTRH, CG, AH, and MFCF,** Formal analysis, Validation, Writing – original draft, Writing – review & editing.

ACKNOWLEDGMENTS
We thank the following institutions for providing access to specimens for morphological and genetic analyses: Natural History Museum, UK, The National Museum of Natural History, Smithsonian Institution (USNM); Field Museum of Natural History (FMNH); American Museum of Natural History (AMNH); University of Michigan Museum of Zoology (UMMZ); the Institute of Zoology, Guangdong Academy of Sciences (IOZ-GDAS); and the Guangdong Wildlife Monitoring and Rescue Center (GWMRC), China. We are grateful to the Natural History Museum, Tribhuvan University, and the Department of National Parks and Wildlife Conservation (Nepal) for providing research permissions and institutional support. We sincerely thank the Ministry of Education, Science and Technology (MoEST), Nepal, and the Nepal Academy of Science and Technology (NAST) for their support. NPK's effort to this study was partially supported by MoEST, Nepal, 2019, an ANSO-CAS visiting Scientist fellowship at Kunming Institute of Zoology, Chinese Academy of Sciences, 2025 and South China Biodiversity Research Center, School of Life Sciences, Guangzhou University, Guangzhou, China, 2023. This study is supported by the ADCS Seed Grants program of the National Museum of Natural History, Smithsonian Institution (M.T.R.H. and A.H.), Guangdong Provincial Natural Science Foundation for Distinguished Young Scholars (2022B1515020033, K.H.) and the National Natural Science Foundation of China (32170452, K.H.).

REFERENCES
Challender, D. W., Nash, H. C. & Waterman, C. *Pangolins: science, society and conservation*. (Academic Press, 2019).
Zhang, F., Wu, S. & Cen, P. The past, present and future of the pangolin in Mainland China. *Global Ecology and Conservation* **33**, e01995, https://doi.org/:10.1016/j.gecco.2021.e01995 (2022).
Challender, D. W., Heinrich, S., Shepherd, C. R. & Katsis, L. K. in *pangolins*   259-276 (Elsevier, 2020).
Zhang, F. *et al.* Reducing pangolin demand by understanding motivations for human consumption in Guangdong, China. *Frontiers in Ecology and Evolution* **8**, 574161, https://doi.org/:10.3389/fevo.2020.574161 (2020).
Zoological Society of London. *Plight of the pangolin: all eight species to move up EDGE list. EDGE of Existence*, <https://www.edgeofexistence.org/blog/plight-of-the-pangolin-all-eight-species-move-up-edge-list/> (2014).
Hu, Y. *et al.* Genomic evidence for two phylogenetic species and long-term population bottlenecks in red pandas. *Science advances* **6**, eaax5751, https://doi.org/:10.1126/sciadv.aax575 (2020).

Ma, Y. *et al.* Demographic history and identification of threats revealed by population genomic analysis provide insights into conservation for an endangered maple. *Molecular Ecology* **31**, 767-779, https://doi.org/:10.1111/mec.16289 (2022).

Hoffmann, A. A., Sgrò, C. M. & Kristensen, T. N. Revisiting adaptive potential, population size, and conservation. *Trends in Ecology & Evolution* **32**, 506-517, https://doi.org/:10.1016/j.tree.2017.03.012 (2017).

Gaubert, P. *et al.* The complete phylogeny of pangolins: scaling up resources for the molecular tracing of the most trafficked mammals on earth. *Journal of Heredity* **109**, 347-359, https://doi.org/:10.1093/jhered/esx097 (2018).

Gu, T.-T. *et al.* Genomic analysis reveals a cryptic pangolin species. *Proceedings of the National Academy of Sciences* **120**, e2304096120, https://doi.org/:10.1073/pnas.2304096120 (2023).

Banterng, N., Ewart, K., Sitam, F. T. & Ogden, R. Mitogenomic analysis of Thai Sunda pangolins reveals regional phylogeography and informs conservation management. *Scientific Reports* **15**, 14067, https://doi.org/:10.1038/s41598-025-97182-1 (2025).

Hinckley, A., Flores, M. F. C., Inayah, N. & Hawkins, M. T. Uncovering new lineages in the Sunda pangolin (Manis javanica) with museum mitogenomics. *Biology Letters* **21**, 20250222, https://doi.org/:10.1098/rsbl.2025.0222 (2025).

Hu, J.-Y. *et al.* Genomic consequences of population decline in critically endangered pangolins and their demographic histories. *National Science Review* **7**, 798-814, https://doi.org/:10.1093/nsr/nwaa031 (2020).

Wei, S. *et al.* Conservation genomics of the critically endangered Chinese pangolin. *Science China Life Sciences*, 1-11, https://doi.org/:10.1007/s11427-023-2540-y (2024).

Challender, D. *et al.*   (Gland: IUCN, 2020).

He, K. & Jiang, X. Sky islands of southwest China. I: an overview of phylogeographic patterns. *Chinese Science Bulletin* **59**, 585-597, https://doi.org/:10.1007/s11434-013-0089-1 (2014).

Koju, N. P. *et al.* Multilocus approaches reveal underestimated species diversity and inter-specific gene flow in pikas (*Ochotona*) from southwestern China. *Molecular Phylogenetics and Evolution*, https://doi.org/:10.1016/j.ympev.2016.11.005 (2017).

Wang, Q. *et al.* Whole-genome resequencing of Chinese pangolins reveals a population structure and provides insights into their conservation. *Communications Biology* **5**, 821, https://doi.org/:10.1038/s42003-022-03757-3 (2022).

Bashyal, A., Shrestha, N., Dhakal, A., Khanal, S. N. & Shrestha, S. Illegal trade in pangolins in Nepal: Extent and network. *Global Ecology and Conservation* **32**, e01940 (2021).

Gomez, L., Joseph, T., Heinrich, S., Wright, B. & D'Cruze, N. Illegal trade of pangolins in India with international trade links: an analysis of seizures from 1991 to 2022. *European Journal of Wildlife Research* **69**, 85, https://doi.org/:10.1007/s10344-023-01708-9 (2023).

Koju, N. P. & Tamang, B. M. Illegal Wildlife Trade in Kathmandu Valley: A Hub of Biodiversity Crime. *Ecology and Evolution (unpublished)* (2025).

Zhang, M., Gouveia, A., Qin, T., Quan, R. & Nijman, V. Illegal pangolin trade in northernmost Myanmar and its links to India and China. *Global Ecology and Conservation* **10**, 23-31, https://doi.org/:10.1016/j.gecco.2017.01.006 (2017).

Nijman, V., Zhang, M. X. & Shepherd, C. R. Pangolin trade in the Mong La wildlife market and the role of Myanmar in the smuggling of pangolins into China. *Global Ecology and Conservation* **5**, 118-126, https://doi.org/:10.1016/j.gecco.2015.12.003 (2016).

Koju, N. P. Abundance and phylogenetic status of Chinese Pangolin (*Manis pentadactyla*) from Kathmandu valley, Nepal. 33 (Ministry of Education Science and Technology Government of Nepal (MoEST, GoN), Kathmandu, Nepal. https://doi.org/10.13140/RG.2.2.36784.12807/1, 2018).

Shrestha, S. *et al.* Mitochondrial DNA analysis of critically endangered Chinese Pangolins (*Manis pentadactyla*) from Nepal. *Mitochondrial DNA Part B* **5**, 3257-3261, https://doi.org/:10.1080/23802359.2020.1811174 (2020).

Hogan, F. *et al.* Genetic evidence reveals distinct lineage of Chinese pangolin in Nepal: insights from scat and blood samples for conservation and wildlife forensics. *Ecology and Evolution* **15**, e70982, https://doi.org/:10.1002/ece3.70982 (2025).

Wangmo, L. K. *et al.* Indo-Burmese pangolin (*Manis indoburmanica*): a novel phylogenetic species of pangolin evolved in Asia. *Mammalian Biology*, 1-8, https://doi.org/:10.1007/s42991-024-00475-7 (2025).

Wilson, D. E. & Mittermeier, R. A. *Handbook of the mammals of the world: vol. 8: insectivores, sloths and colugos*. 79 (Lynx Edicions, 2018).

Heighton, S. P. *et al.* Pangolin genomes offer key insights and resources for the world's most trafficked wild mammals. *Molecular Biology and Evolution* **40**, msad190 (2023).

Gaudin, T. J., Emry, R. J. & Wible, J. R. The phylogeny of living and extinct pangolins (Mammalia, Pholidota) and associated taxa: a morphology based analysis. *Journal of mammalian evolution* **16**, 235-305, https://doi.org/:10.1007/s10914-009-9119-9 (2009).

Corbet, G. B. & Hill, J. E. *The mammals of the Indomalayan region: a systematic review*. Vol. 488 (oxford university press Oxford, 1992).

Ballard, J. W. O. & Whitlock, M. C. The incomplete natural history of mitochondria. *Molecular ecology* **13**, 729-744, https://doi.org/:10.1046/j.1365-294X.2003.02063.x (2004).

Funk, D. J. & Omland, K. E. Species-Level Paraphyly and Polyphyly: Frequency, Causes, and Consequences, with Insights from Animal Mitochondrial DNA. *Annual Review of Ecology, Evolution, and Systematics* **34**, 397-423, https://doi.org/:10.1146/annurev.ecolsys.34.011802.132421 (2003).

Toews, D. P. & Brelsford, A. The biogeography of mitochondrial and nuclear discordance in animals. *Molecular ecology* **21**, 3907-3930, https://doi.org/:10.1111/j.1365-294X.2012.05664.x (2012).

Zijlstra, J. S. Manis aurita Hodgson 1837 as a valid species of pangolin: a comment to Wangmo et al.(2025). *Mammalian Biology*, 1-3, https://doi.org/:10.1007/s42991-025-00495-x (2025).

Hodgson, B. H. New Mammals in Nipdlese Catalogue. *The journal of the Asiatic Society of Bengal* **V**, 234-235 (1836).

Fitzinger, L. J. F. J. *Die natürliche Familie der Schuppenthiere (Manes).Sitzungsberichte der kaiserlichen* 65:9–83 (Akademie der Wissenschaften, 1872).

Padial, J. M., Miralles, A., De la Riva, I. & Vences, M. The integrative future of taxonomy. *Frontiers in zoology* **7**, 1-14, https://doi.org/:10.1186/1742-9994-7-16 (2010).

Schlick-Steiner, B. C. *et al.* Integrative taxonomy: a multisource approach to exploring biodiversity. *Annual review of entomology* **55**, 421-438, https://doi.org/:10.1146/annurev-ento-112408-085432 (2010).

Fujita, M. K., Leaché, A. D., Burbrink, F. T., McGuire, J. A. & Moritz, C. Coalescent-based species delimitation in an integrative taxonomy. *Trends in ecology & evolution* **27**, 480-488, https://doi.org/:10.1016/j.tree.2012.04.012 (2012).

Dayrat, B. Towards integrative taxonomy. *Biological journal of the Linnean society* **85**, 407-417, https://doi.org/:10.1111/j.1095-8312.2005.00503.x (2005).

Rieseberg, L. H. & Burke, J. M. The biological reality of species: gene flow, selection, and collective evolution. *Taxon* **50**, 47-67, https://doi.org/:10.2307/1224511 (2001).

Van Elst, T. *et al.* Integrative taxonomy clarifies the evolution of a cryptic primate clade. *Nature Ecology & Evolution* **9**, 57-72, https://doi.org/:10.1038/s41559-024-02547-w (2025).

Abreu, E. F. *et al.* Recent, Intricate Speciation in Amazonia Uncovered by a Multilayered Genomic Analysis of Tree Squirrels. *Systematic Biology*, syaf013, https://doi.org/:10.1093/sysbio/syaf013 (2025).

Lehman, S. M., Fleagle, J. G., Meijaard, E. & Groves, C. P. in *Primate Biogeography. Developments in Primatology: Progress and Prospects*   305-329 (Springer, 2006).

Hinckley, A., Maldonado, J. E., Tamura, N., Leonard, J. A. & Hawkins, M. T. Lost in synonymy: Integrative species delimitation reveals two unrecognized species of Southern Asian tree squirrels (Rodentia: *Sciuridae: Callosciurinae*). *Vertebrate Zoology* **74**, 683-707, https://doi.org/:10.3897/vz.74.e133467 (2024).

Sanyal, R. *A handbook of the management of wild animals in captivity in lower Bengal*. (Calcutta: Bengal Secretariat Press (reprinted by Central Zoo Authority in 1995), 1892).

Hinckley, A., Hawkins, M. T., Maldonado, J. E. & Leonard, J. A. Evolutionary history and patterns of divergence in three tropical east Asian squirrels across the Isthmus of Kra. *Journal of Biogeography* **50**, 1090-1102, https://doi.org/:10.1111/jbi.14598 (2023).

Louys, J. & Meijaard, E. Palaeoecology of Southeast Asian megafauna-bearing sites from the Pleistocene and a review of environmental changes in the region. *Journal of Biogeography* **37**, 1432-1449, https://doi.org/:10.1111/j.1365-2699.2010.02297.x (2010).

Camacho-Sanchez, M., Leonard, J. A., Fitriana, Y., Tilak, M.-K. & Fabre, P.-H. The generic status of Rattus annandalei (Bonhote, 1903)(Rodentia, Murinae) and its evolutionary implications. *Journal of Mammalogy* **98**, 1340-1355, https://doi.org/:10.1093/jmammal/gyx081 (2017).

Hawkins, M. T. *et al.* Evolutionary history of endemic Sulawesi squirrels constructed from UCEs and mitogenomes sequenced from museum specimens. *BMC Evolutionary Biology* **16**, 1-16, https://doi.org/:10.1186/s12862-016-0650-z (2016).

Leonard, J. A. *et al.* Phylogeography of vertebrates on the Sunda Shelf: a multi-species comparison. *Journal of Biogeography* **42**, 871-879, https://doi.org/:10.1111/jbi.12465 (2015).

Tong, H. Occurrences of warm-adapted mammals in north China over the Quaternary Period and their paleo-environmental significance. *Science in China Series D: Earth Sciences* **50**, 1327-1340, https://doi.org/:10.1007/s11430-007-0096-7 (2007).

Imam, A. *et al.* The brain of the tree pangolin (*Manis tricuspis*). II. The olfactory system. *Journal of Comparative Neurology* **526**, 2548-2569, https://doi.org/:10.1002/cne.24510 (2018).

Li, J. *et al.* Genomic signatures of sensory adaptation and evolution in pangolins. *BMC genomics* **25**, 1-13, https://doi.org/:10.1186/s12864-024-11063-x (2024).

DiPaola, J. D., Yindee, M. & Plotnik, J. M. Investigating the use of sensory information to detect and track prey by the Sunda pangolin (*Manis javanica*) with conservation in mind. *Scientific Reports* **10**, 9787, https://doi.org/:10.1038/s41598-020-65898-x (2020).

Zhai, T., Wang, J., Zhan, G., Hu, J. & Yu, L. Population genomic landscapes and insights for conservation of the critically endangered island-endemic Chinese pangolin in Taiwan. *Science China Life Sciences*, 1-16, https://doi.org/:10.1007/s11427-024-2904-6 (2025).

Ma, T. *et al.* Intensified climate drying and cooling during the last glacial culmination (20.8–17.5 cal ka BP) in the south-eastern Asian monsoon domain inferred from a high-resolution pollen record. *Quaternary Science Reviews* **278**, 107371, https://doi.org/:10.1016/j.quascirev.2022.107371 (2022).

Colleoni, F., Wekerle, C., Näslund, J.-O., Brandefelt, J. & Masina, S. Constraint on the penultimate glacial maximum Northern Hemisphere ice topography ($\approx$ 140 kyrs BP). *Quaternary Science Reviews* **137**, 97-112, https://doi.org/:10.1016/j.quascirev.2016.01.024 (2016).

Li, Z., Fan, H., Liao, Z., Wang, Y. & Wei, F. Global spatiotemporal patterns of demographic fluctuations in terrestrial vertebrates during the Late Pleistocene. *Science Advances* **11**, eadq3938, https://doi.org/:10.1126/sciadv.adq3938 (2025).

Dai, Y. *et al.* Historical changes in the distribution of the Sichuan golden snub‑nosed monkey (*Rhinopithecus roxellana*) in Sichuan Province, China. *Ecology and Evolution* **14**, e11270, https://doi.org/:https://doi.org/10.1002/ece3.11270 (2024).

Zheng, J. *et al.* How climate change impacted the collapse of the Ming dynasty. *Climatic change* **127**, 169-182, https://doi.org/:10.1007/s10584-014-1244-7 (2014).

Zhang, W. Q. & Huang, Z. G. The Little Ice Age and its environmental effects in the tropical zone of China. *Acta Geographica Sinica-Chinese Edition* **55**, 750-750 (2000).

Ren, M. *et al.* The collapse of the Ming Dynasty actually began with the Wanli megadrought: Insights from a hydroclimate reconstruction based on tree-ring δ18O over the past 460 years. *Palaeogeography, Palaeoclimatology, Palaeoecology* **655**, 112548, https://doi.org/:10.1016/j.palaeo.2024.112548 (2024).

Rowan, A. V. The 'Little Ice Age' in the Himalaya: A review of glacier advance driven by Northern Hemisphere temperature change. *The Holocene* **27**, 292-308, https://doi.org/:10.1177/0959683616658530 (2017).

Yadav, R. R., Braeuning, A. & Singh, J. Tree ring inferred summer temperature variations over the last millennium in western Himalaya, India. *Climate Dynamics* **36**, 1545-1554, https://doi.org/: 10.1007/s00382-009-0719-0 (2011).

Brook, S. M. *et al.* Lessons learned from the loss of a flagship: the extinction of the Javan rhinoceros *Rhinoceros sondaicus annamiticus* from Vietnam. *Biological Conservation* **174**, 21-29, https://doi.org/:10.1016/j.biocon.2014.03.014 (2014).

Goossens, B. *et al.* Habitat fragmentation and genetic diversity in natural populations of the Bornean elephant: Implications for conservation. *Biological Conservation* **196**, 80-92, https://doi.org/:10.1016/j.biocon.2016.02.008 (2016).

Boonratana, R. Asian primates in fragments: Understanding causes and consequences of fragmentation, and predicting primate population viability. *American Journal of Primatology* **82**, e23082, https://doi.org/:10.1002/ajp.23082 (2020).

Xie, X. *et al.* DNA mini-barcodes, a potential Weapon for Conservation and combating illegal trade of pangolin. *Tropical Conservation Science* **14**, 19400829211017361, https://doi.org/:10.1177/1940082921101736 (2021).

Gaudin, T. J. & Wible, J. R. The entotympanic of pangolins and the phylogeny of the Pholidota (Mammalia). *Journal of Mammalian Evolution* **6**, 39-65, https://doi.org/:10.1023/A:1020538313412 (1999).

Gurung, K. K. & Singh, R. *Field Guide to the Mammals of the Indian Subcontinent*. (Academic Press, 1996).

Sharma, H. P. *et al.* Potential distribution of the critically endangered Chinese Pangolin (*Manis pentadactyla*) in different land covers of Nepal: Implications for conservation. *Sustainability* **12**, 1282, https://doi.org/:10.3390/su12031282 (2020).

Shrestha, A., Bhattarai, S., Shrestha, B. & Koju, N. P. Factors influencing the habitat choice of pangolins (*Manis* spp.) in low land of Nepal. *Ecology and Evolution* **11**, 14689-14696, https://doi.org/:10.1002/ece3.8156 (2021).

Dhital, S. *et al.* Distribution of Chinese pangolin (*Manis pentadactyla*) in Nagarjun forest of Shivapuri Nagarjun National Park, Nepal. *Nepalese Journal of Zoology* **4**, 1-7, https://doi.org/:10.3126/njz.v4i1.30667 (2020).

Suwal, T. L. *et al.* Predicting the potential distribution and habitat variables associated with pangolins in Nepal. *Global Ecology and Conservation* **23**, e01049, https://doi.org/:10.1016/j.gecco.2020.e01049 (2020).

Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008, https://doi.org/:10.1093/gigascience/giab008 (2021).

Meng, G., Li, Y., Yang, C. & Liu, S. MitoZ: a toolkit for animal mitochondrial genome assembly, annotation and visualization. *Nucleic acids research* **47**, e63-e63, https://doi.org/:10.1093/nar/gkz173. (2019).

Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular biology and evolution* **30**, 772-780, https://doi.org/:10.1093/molbev/mst010 (2013).

Li, H. *et al.* The sequence alignment/map format and SAMtools. *bioinformatics* **25**, 2078-2079, https://doi.org/:10.1093/bioinformatics/btp352 (2009).

McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research* **20**, 1297-1303, https://doi.org/:10.1101/gr.107524.110 (2010).

Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158, https://doi.org/:10.1093/bioinformatics/btr330 (2011).

Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome biology* **16**, 1-14, https://doi.org/:10.1186/s13059-015-0721-2 (2015).

Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *The American journal of human genetics* **81**, 559-575, https://doi.org/:10.1086/519795 (2007).

Patterson, N., Price, A. L. & Reich, D. Population structure and eigenanalysis. *PLoS genetics* **2**, e190, https://doi.org/:10.1371/journal.pgen.0020190 (2006).

Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. *Genome research* **19**, 1655-1664, https://doi.org/:10.1101/gr.094052.109 (2009).

Douglas, J., Jiménez-Silva, C. L. & Bouckaert, R. StarBeast3: adaptive parallelized Bayesian inference under the multispecies coalescent. *Systematic Biology* **71**, 901-916, https://doi.org/:10.1093/sysbio/syac010 (2022).

Bouckaert, R. *et al.* BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS computational biology* **15**, e1006650, https://doi.org/:10.1371/journal.pcbi.1006650 (2019).

Smith, S. A., Brown, J. W. & Walker, J. F. So many genes, so little time: a practical approach to divergence-time estimation in the genomic era. *PloS one* **13**, e0197433, https://doi.org/:10.1371/journal.pone.0197433 (2018).

Botha, J. & Gaudin, T. An early pliocene pangolin (mammalia; pholidota) from langebaanweg, south africa. *Journal of Vertebrate Paleontology* **27**, 484-491, https://doi.org/:0.1671/0272-4634(2007)27[484:AEPPPF]2.0.CO;2 (2007).

Rambaut, A., Suchard, M. & Xie, D.   (2017).

Li, H. & Durbin, R. Inference of human population history from individual whole-genome sequences. *Nature* **475**, 493-496, https://doi.org/:10.1038/nature10231 (2011).

IUCN SSC Pangolin Specialist Group. *Regional Sunda Pangolin (Manis javanica) Conservation Strategy 2018-2028. IUCN SSC*. 37 (Pangolin Specialist Group, 2018).

McQuillan, R. *et al.* Runs of homozygosity in European populations. *The American Journal of Human Genetics* **83**, 359-372, https://doi.org/:10.1016/j.ajhg.2008.08.007 (2008).

Santiago, E. *et al.* Recent demographic history inferred by high-resolution analysis of linkage disequilibrium. *Molecular Biology and Evolution* **37**, 3642-3653, https://doi.org/:10.1093/molbev/msaa169 (2020).

Malinsky, M., Matschiner, M. & Svardal, H. Dsuite‐Fast D‐statistics and related admixture evidence from VCF files. *Molecular ecology resources* **21**, 584-595, https://doi.org/:10.1111/1755-0998.13265 (2021).

Excoffier, L. *et al.* fastsimcoal2: demographic inference under complex evolutionary scenarios. *Bioinformatics* **37**, 4882-4885, https://doi.org/:10.1093/bioinformatics/btab468 (2021).

Korunes, K. L. & Samuk, K. pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. *Molecular ecology resources* **21**, 1359-1368, https://doi.org/:10.1111/1755-0998.13326 (2021).

Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic acids research* **25**, 3389-3402, https://doi.org/:10.1093/nar/25.17.3389 (1997).

Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics: a journal of integrative biology* **16**, 284-287, https://doi.org/:10.1089/omi.2011.0118 (2012).

Wu, S., Naifa, L., Zhang Yingmei, Zhi, O. & Hai, C. Measurement and Comparison of Skull Variables in Chinese Pangolin and Malayan Pangolin. *Acta Theriologica Sinica* **24**, 211, https://doi.org/:10.16829/j.slxb (2005).

Sulaiman, M. H., Azmi, W. A., Hassan, M. & Chong, J. L. Current updates on the morphological measurements of the Malayan pangolin (*Manis javanica*). *Folia Zoologica* **66**, 262-266, https://doi.org/:10.25225/fozo.v66.i4.a8.2017 (2017).

Igado, O., Adekanmbi, J., Adebayo, M., Sulaimon, O. & Osundina, O. Facial and Body Morphometric Indices of the Arboreal African Pangolin (*Manis tricuspis*):

Anthropometric Tools in Describing the Pangolin: Facial and biometric indices of the African pangolin. *Sahel Journal of Veterinary Sciences* **17**, 28-33, https://doi.org/:10.54058/saheljvs.v17i3.122 (2020).

Morenikeji, O. *et al.* Morphometric characterisation of white-bellied pangolins rescued from trade for conservation in Nigeria. *Emerging Animal Species* **1**, 100018, https://doi.org/:10.1016/j.eas.2022.100018 (2022).

Gaubert, P. & Antunes, A. Assessing the taxonomic status of the Palawan pangolin *Manis culionensis* (Pholidota) using discrete morphological characters. *Journal of Mammalogy* **86**, 1068-1074, https://doi.org/:10.1644/1545-1542(2005)86[1068:ATTSOT]2.0.CO;2 (2005).

Arbour, J. H. & Brown, C. M. Incomplete specimens in geometric morphometric analyses. *Methods in Ecology and Evolution* **5**, 16-26, https://doi.org/:10.1111/2041-210X.12128 (2014).

Ferreira-Cardoso, S., Billet, G., Gaubert, P., Delsuc, F. & Hautier, L. Skull shape variation in extant pangolins (Pholidota: Manidae): allometric patterns and systematic implications. *Zoological Journal of the Linnean Society* **188**, 255-275, https://doi.org/:10.1093/zoolinnean/zlz096 (2019).

Dryden, I. L. & Mardia, K. V. Multivariate shape analysis. *Sankhyā: The Indian Journal of Statistics, Series A*, 460-480 (1993).

Goswami, A. Cranial modularity shifts during mammalian evolution. *The American Naturalist* **168**, 270-280, https://doi.org/:10.1086/505758 (2006).

Hautier, L., Billet, G., De Thoisy, B. & Delsuc, F. Beyond the carapace: skull shape variation and morphological systematics of long-nosed armadillos (genus *Dasypus*). *PeerJ* **5**, e3650, https://doi.org/:10.7717/peerj.3650 (2017).

Algewatta, H. R. *et al.* Updates on the morphometric characterization of Indian pangolin (*Manis crassicaudata*) in Sri Lanka. *Animals* **11**, 25, https://doi.org/:10.3390/ani11010025 (2020).

Guo, C. *et al.* Coding System and Comparisons of Scales Between Malayan Pangolin(*Manis javanica*) and Chinese Pangolin(*M. pentadactyla*). *Chinese Journal of Wildlife* **43**, 024 − 031, https://doi.org/:10.19711/j.cnki.issn2310-1490.2022.01.003 (2022).

Martin, J. M. *et al.* The Pangolin Universal Notching System: a scale-marking methodology for pangolins. *Oryx*, 1-7, https://doi.org/:1017/S0030605324000656 (2024).

Koju, N. P. *et al.* in *Dryad*   (Dryad. http://datadryad.org/share/tUnpoAtBaYbC-LxLmopIeUXwwngHX96olu7F6k_uTBk, 2025).