Revalidation of *Manis aurita*, based on integrative morphological and genomic evidence

Narayan Prasad Koju[1,2,3,4]#, Zeling Zeng[1]#, Guihua Zhang[1]#, Xia Huang[1]#, Zhicheng Yao[1],

Xiaoyun Wang[1], Melissa T.R. Hawkins[5], Arlo Hinckley[5,6], Mary Faith C. Flores[5], Ce Guo[7], Jun

Li[7], Devendra Maharjan[8], Saraswoti Byanjankar[9], Lianghua Huang[10], Wenhua Yu[1], Liang

Leng[11], Kai He[1]*, Anderson Feijó [12, *], Yan Hua[7]*

#: equal contribution; *: corresponding authors

Address:

[1] South China Biodiversity Research Center, School of Life Sciences, School of Life Sciences,
Guangzhou University, Guangzhou 510006, China
[2] Center for Postgraduate Studies, Nepal Engineering College, Pokhara University, Lalitpur,
44600, Nepal
[3] Key Laboratory of Genetic Evolution and Animal Models, Kunming Institute of Zoology,
Chinese Academy of Sciences, Kunming, Yunnan 650201, China
[4] Department of Psychology, University of Washington, Seattle 98125, USA
[5] Division of Mammals, Department of Vertebrate Zoology, National Museum of Natural
History, Smithsonian Institution, Washington DC 20024, USA
[6] Departamento de Zoología, Universidad de Sevilla, Seville, 41012, Spain
[7] Guangdong Provincial Key Laboratory of Silviculture, Protection and Utilization, Guangdong
Academy of Forestry, Guangzhou, China
[8] Natural History Museum, Tribhuvan University, Kathmandu 44611, Nepal
[9] Aspect Ecology, West Court, Oxfordshire, OX16 2AF, UK
[10] Wildlife Forensic Science Service, Kunming, Yunnan 650201, China
[11] Chengdu University of Traditional Chinese Medicine, Chengdu 611137, China
[12] Negaunee Integrative Research Center, Field Museum of Natural History, Chicago 60605,
USA

Details on Corresponding author

Kai He

South China Biodiversity Research Center, School of Life Sciences, Guangzhou University

#230 Waihuanxilu Road, Fanyu District, Guangzhou, Guangdong, China

Email: hekai@gzhu.edu.cn; Phone: 15925188783

Anderson Feijó: Email: afeijo@fieldmuseum.org ;

Yan Hua: Email: wildlife530@163.com

ABSTRACT

Pangolins face severe conservation threats globally, making accurate taxonomy essential for their

protection. Previous genomic research on the Chinese pangolin (*M. pentadactyla*) identified two

deeply divergent lineages (MPA and MPB), suggesting underestimated species diversity. The

recent description of *M. indoburmanica* for the MPB lineage, however, did not assess its

relationship with historical names, particularly the subspecies *M. p. aurita*, leaving the group's

taxonomic status uncertain. To resolve this issue, we employed an integrative framework,

analyzing genomic and morphological data from museum specimens to clarify the validity and

relationship of these taxa. Our results demonstrate that the MPB lineage, which encompasses *M.*

indoburmanica, is geographically restricted to the southern Himalayas, a range congruent with

the type locality of *M. aurita* and distinct from other *M. pentadactyla* populations. Genomic

analyses confirm this Himalayan lineage diverged ~1.2 Ma and has remained largely isolated

since, with only minimal gene flow. Furthermore, morphometric analyses of both cranial and

external features, including the *aurita* holotype, reveal consistent and significant differentiation

between the Himalayan lineage and *M. pentadactyla sensu stricto*. Collectively, this congruent

evidence unequivocally supports the revalidation of the Himalayan pangolin as a distinct species,

Manis aurita Hodgson, 1836, representing the fifth extant species in the genus. This taxonomic

revision provides a critical foundation for species-specific conservation strategies and enhances

wildlife forensic capabilities by creating a robust genetic baseline to trace the provenance of

illegally traded products.

Keywords: *Manis aurita*, Integrative taxonomy, Phylogenomics, Morphometrics, Conservation

genetics

INTRODUCTION

Pangolins (Order Pholidota) are unique mammals characterized by their keratinized scales, an extraordinary trait among mammals (Challender et al., 2019). These animals face severe threats including habitat loss and extensive illegal trade, which has resulted in dramatic population declines (Challender et al., 2020b; Zhang et al., 2020; Zhang et al., 2022). Consequently, all pangolin species are listed under Appendix I of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and are recognized as Evolutionarily Distinct and Globally Endangered (EDGE) species (Zoological Society of London, 2014). Accurate species identification, clarification of species distribution, and estimation of population sizes and genetic load are essential for effective conservation efforts.

Until recently, eight extant pangolin species were recognized: four in Africa and four in Asia (Gaubert et al., 2018). All Asian species belong to the genus *Manis*, including Chinese pangolin (*M. pentadactyla*), Indian pangolin (*M. crassicaudata*), Sunda pangolin (*M. javanica*), and Philippine pangolin (*M. culionensis*). Recent genomic studies revealed cryptic genetic and taxonomic diversity within Asian pangolins. For instance, genome sequencing of confiscated scales uncovered a cryptic lineage, provisionally named "mysteria", for which no voucher specimens or precise locality data exist (Gu et al., 2023). Moreover, analyses of broad sampling within *M. pentadactyla* have revealed distinct clades, indicating the existence of an undescribed species.

The Chinese pangolin is distributed across a broad range, extending from Nepal in the west to the eastern coast of China, with insular populations on Hainan and Taiwan islands. Latitudinally, its range extends from central China southward to Southeast Asia (Challender et al., 2020a). This broad distribution is divided by geographic barriers, such as the mountains of southwestern

China and southward-flowing river systems, which likely drive genetic divergence among

geographic populations (He & Jiang, 2014; Koju et al., 2017). Accordingly, Hu et al. (2020a)

identified two major clades within *M. pentadactyla,* namely, MPA and MPB. While recent

studies have found additional subclades closely related to MPA in China and Southeast Asia

(Wang et al., 2022; Wei et al., 2024), MPB remains the most genetically divergent lineage, and

its geographic origin has been enigmatic. Initially, the MPB clade was known almost

exclusively from specimens confiscated along the Sino-Myanmar border, suggesting a Myanmar

origin. This inference was tentative, as the area is a major wildlife trafficking hub (Nijman et al.,

2016; Zhang et al., 2017; Bashyal et al., 2021; Gomez et al., 2023; Koju & Tamang, 2025).

However, recent studies found occurrences of MPB in Nepal, South Tibet and northeastern India

(Koju, 2018; Shrestha et al., 2020; Hogan et al., 2025; Wangmo et al., 2025).

Historically, three subspecies of *M. pentadactyla* were recognized: the nominate subspecies *M. p.*

pentadactyla Linnaeus, 1758 described based on specimens from Taiwan (Corbet & Hill, 1992);

M. p. pusilla Allen, 1906 from the Hainan Island; and *M. p. aurita* Hodgson, 1836 from the

central Nepal (Wilson & Mittermeier, 2018). While the former two have been subject to

extensive genetic studies (Gaudin et al., 2009; Hu et al., 2020a; Wang et al., 2022; Heighton et

al., 2023; Wei et al., 2024), and their evolutionary relationships as well as distribution in China

and SE Asia has been confirmed (Hu et al., 2020a; Wei et al., 2024), the taxonomy and

evolutionary history of *M. p. aurita* remains unclear.

Recently, Wangmo et al. (2025) describe a new species, *Manis indoburmanica*, from South

Tibet, which highlights several challenges in modern taxonomy. The recognition of this new

species is based solely on the mitochondrial genome, a data source susceptible to introgression,

incomplete lineage sorting, and maternal inheritance bias (Funk & Omland, 2003; Ballard &

Whitlock, 2004; Toews & Brelsford, 2012). Furthermore, the work lacks a comprehensive

morphological comparison with *M. pentadactyla*, thus failing to establish a clear morphological

boundary for *M. indoburmanica*. Critically, as noted by Zijlstra (2025), the authors did not

evaluate whether their samples correspond to existing historical names like *M. p. aurita*

(Hodgson, 1836) or *M. assamensis* (Fitzinger, 1872), leaving the novelty of the taxon in

question. Such challenges underscore the necessity of integrative taxonomic approaches that

incorporate genome-wide data, robust morphological comparisons with both *M. pentadactyla*

and *M. p. aurita* (Padial et al., 2010; Schlick-Steiner et al., 2010; Fujita et al., 2012).

To resolve this taxonomic uncertainty, we employed an integrative taxonomic framework

combining genomic and morphological evidence. Our genomic analyses incorporated topotypical

specimens of *M. p. aurita*, while our morphological comparisons included an examination of the

external characteristics of its type specimen. The objective of this study was to evaluate whether

pangolins from South Asia correspond to the subspecies *aurita* and, if so, whether their degree of

divergence supports elevation to full species status. Our analyses reveal clear and pronounced

differentiation from *M. pentadactyla* across both genetic and morphological datasets, providing

robust support for recognizing this lineage as the distinct species *Manis aurita* Hodgson, 1836.

MATERIAL AND METHODS

Taxon sampling and sequencing

DNA was extracted from tissue samples (n=18) and purified using the Dneasy blood and tissue

kit (Qiagen). Genomic DNA was sheared to ~350 bp fragments using a Covaris LE220R-plus

system (Covaris, USA), followed by end-repair, A-tailing, and adapter ligation. Libraries were

amplified by PCR, purified with AMPure XP beads.

Destructive sampling was conducted using skin specimens deposited in the National Museum of

Natural History, the Smithsonian Institution (USNM), Field Museum of Natural History

(FMNH); American Museum of Natural History (AMNH) and University of Michigan, Museum

of Zoology (UMMZ) in the United States. Sterile practices were employed with routine

bleaching, instrument sterilization, and changing of PPE to prevent any cross contamination

between specimens. Samples of 25 mg of material were placed in sterile cryovials or 1.5mL

tubes and transferred to a dedicated Historic DNA processing facility at the Smithsonian

Institution's Museum Support Center where DNA extraction was performed via Qiagen

MinElute or Investigator kits following the manufacturer's protocol. Single-stranded library

preparation was performed with the SRSLY Pico Plus Kit (Claret Biosystems®) utilizing half

reactions. Libraries had dual indexed iTru style adapters ligated to each sample and amplified for

18 cycles of PCR. Following amplification, samples were visualized on a High-Sensitivity Tape

Station (Agilent), and pooled in equimolar ratios for sequencing.

Sequencing was performed on an Illumina NovaSeq X platform to generate 150 bp paired-end

(PE150) reads. Raw FASTQ reads were processed with fastp v0.24.0 (Danecek et al., 2021) to

remove adapter contamination and poly-G tails, trimming bases with Phred score ≤ 5. Reads

containing > 10 % ambiguous bases or > 50 % low-quality bases were discarded. We

supplemented our dataset with 48 available genome re-sequencing data including 24 *M.*

pentadactyla, 19 *M. javanica*, 2 *M. crassicaudata*, 1 *Phataginus tricuspis*, and 1 *Smutsia*

gigantea (**Table S1**).

Mitochondrial assembly and phylogenetic analyses

Mitochondrial genomes were assembled from 64 raw sequencing datasets, including data from

30 newly sequenced individuals and 34 previously published genomes of various pangolin

species. For these datasets, clean reads were processed for *de novo* assembly and annotation in

MITOZ v3.6 (Meng et al., 2019), which employed Megahit with k-mer values of 21, 33, and 55.

This final set of assembled mitogenomes was then augmented with five additional mitogenomes

obtained from GenBank (**Table S2**). We aligned mitogenomes using MAFFT v7.5 (Katoh &

Standley, 2013), and trimmed poorly aligned regions using Gblocks. We retained 12S, 16S

rRNA, and 12 coding genes on the heavy chain for phylogenetic analyses. We also downloaded

available mitochondrial 18 *CYTB* and 12 *COI* genes identified as *M. pentadactyla* from

GenBank. We used IQ-TREE v2.1.3 to estimate the best☐scoring Maximum☐likelihood (ML)

trees. ModelFinder was used to automatically determine the optimal partitioning scheme and

substitution models under the Bayesian Information Criterion (–m MFP+MERGE). Node

support was evaluated with 1,000 ultrafast bootstrap replicates.

Genome assembly and phylogenetic analyses

We aligned clean data to a *M. pentadactyla* reference genome (GCF_030020395.1) using BWA

v0.7.17. The resulting SAM files were converted to coordinate☐sorted BAMs and indexed with

SAMtools v1.10 (Li et al., 2009). Duplicates were identified and flagged using MarkDuplicates

tool of GATK v.4.6 (McKenna et al., 2010). Local realignment around indels was then

performed using GATK's IndelRealigner to reduce false☐positive variant calls. Per☐sample

gVCFs were generated with GATK's HaplotypeCaller and merged via CombineGVCFs. SNP

calling was then executed with GenotypeGVCFs and SelectVariants to produce multisample

VCF and isolate high-confidence SNP calls, respectively. We filtered SNP calls using GATK's

VariantFiltration module with thresholds: QUAL < 30.0, QD < 2.0, MQ < 40.0, FS > 60.0, SOR

> 3.0, MQRankSum < –12.5, ReadPosRankSum < –8.0, and SB ≥ –1.0. We applied VCFtools

v0.1.16 to remove SNPs with minor☐allele frequency < 2 % or depth outside the 2.5–97.5

percentile of the depth distribution (Danecek et al., 2011).

Phylogenetic relationships were inferred using three datasets: high-quality single

nucleotide polymorphisms (SNPs) for autosomal and sex chromosomes as well as single-copy

orthologous genes. For the SNP-based analyses, autosomal and sex chromosome SNP data were

separately extracted from the filtered, high-quality multi-sample VCF. These datasets were

converted to PHYLIP format using vcf2phylip v2.8. ML trees were constructed for both datasets

using IQ-TREE v2.1.4 as mentioned above. For the ortholog-based analysis, we first identified

23,752 single-copy orthologs using OrthoFinder v2.5.4 (Emms & Kelly, 2015). These orthologs

underwent stringent filtering based on GC content, sequence variability, proportion of missing

data, and ambiguous bases, resulting in 7,852 high-quality loci. Each loci was individually

aligned and a concatenated ML tree was constructed using IQ-TREE as mentioned above.

Genetic structuring analyses

To investigate population genetic structure, we utilized linkage-disequilibrium (LD)–pruned

autosomal SNPs. LD pruning was performed in PLINK v1.90b6.21 using the –indep-pairwise 50

5 0.2 parameters (Purcell et al., 2007). We then performed PCA on mean-imputed genotypes in

PLINK to extract the top ten principal components. To assess the statistical significance of the

principal components, we applied the Tracy-Widom test implemented in EIGENSOFT v8.0.0

(Patterson et al., 2006). We applied PCA to all *Manis* samples and to *M. pentadactyla* sensu lato

(which encompasses individuals later identified as *M. aurita*). To infer individual ancestry

proportions, we ran ADMIXTURE v1.3.0 on the *M. pentadactyla* s.l. samples. We evaluated

models for K (number of ancestral populations) from 1 to 10, and selected the optimal K to

minimize cross-validation error (Alexander et al., 2009).

Molecular dating and population genetic analyses

Divergence times were estimated using *BEAST3 model (Douglas et al., 2022) as implemented in BEAST v2.7.7 (Bouckaert et al., 2019). First, we identified the 2,000 most genetically divergent loci from an initial pool of 7,852 filtered orthologs. These loci were then assessed for their conformity to a strict molecular clock using SortaDate v2.0 (Smith et al., 2018). The 200 orthologs that best fit the strict clock model were subsequently used for the divergence time estimation. The Bayesian inference employed a Birth–Death tree prior, a strict molecular clock, and the HKY+Γ substitution model. We employed two secondary calibrations derived from the molecular dating estimates of Gu et al. (2023). The analysis was constrained by setting priors on i) the crown-group pangolin divergence, set to 33.19 million years ago (Ma) (95% CI: 29.8–30.3 Ma), and ii) the divergence of Asian pangolins, set to 16.9 Ma (95% CI: 12.9–22.3 Ma). The Markov chain Monte Carlo (MCMC) analysis was run for 200 million generations. Convergence and parameter mixing were assessed in Tracer v1.7.2 (Rambaut et al., 2017), ensuring all key parameters achieved effective sample sizes (ESS) > 200.

Historical population dynamics were reconstructed using the Pairwise Sequentially Markovian Coalescent (PSMC) model (Li & Durbin, 2011). According to the estimated divergence time between *M. pentadactyla* and *M. aurita* (see Results), we estimated the mutation rate per generation assuming an average generation time of 7 years (IUCN SSC Pangolin Specialist Group, 2018). The analysis was run with the parameters -N25 -t15 -r5 -p "4+25*2+4+6", using the Manis *pentadactyla* genome (Hu et al., 2020a) as the reference for mapping. To assess the variability in demographic inferences, 100 bootstrap replicates were performed.

Levels of inbreeding were quantified by calculating the average inbreeding coefficient (FROH) for each individual. Long runs of homozygosity (ROH) were identified using PLINK v2.0

(Purcell et al., 2007) with the parameters --homozyg --homozyg-window-snp 20 --homozyg-kb

100. Individual FROH values were then calculated as the proportion of the autosomal genome

covered by ROH, relative to the total autosomal SNP coverage, following McQuillan et al.

(2008).

Geneflow analyses

To infer the direction and magnitude of historical gene flow among *Manis* species, we analyzed a

high-quality SNP dataset from 52 samples representing the four *Manis* species using Dsuite v0.5

(Malinsky et al., 2021) and TreeMix v1.13 (Pickrell & Pritchard, 2012). We first filtered the

SNP dataset to retain sites with ≤20% missing data using VCFtools (--max-missing 0.8). To

quantify admixture, we used the Dtrios program in Dsuite to calculate D-statistics and f4-ratio

statistics for all relevant population trios. *P. tricuspis* and *S. gigantea* were used as outgroups.

Results were visualized with the dtools.py script. To model population splits and migration

events, the filtered SNP data was first pruned for linkage disequilibrium (PLINK; --indep-

pairwise 50 5 0.6), and the output was converted to TreeMix format using plink2treemix.py. We

then ran TreeMix, exploring models with 0 to 5 migration edges (five replicates each). The

optimal model was selected by evaluating the explained variance and residual plot fits. The

resulting population graph was visualized using the R script plotting_funcs.R provided with

TreeMix.

We used FastSimCoal v2.8 (Excoffier et al., 2021) to test potential gene flow between *M.*

pentadactyla (MP) and *M. aurita* (MA). First, we used VCF file through easySFS to generated

folded observed joint site frequency spectrum (JSFS). We constructed four demographic models

to test gene flow scenarios: (1) early gene flow (range from 120,000 to 130,000 generations); (2)

recent gene flow (range from 1,000 to 20,000 generations); (3) continuous gene flow; and (4) no

gene flow. For each tested demographic model, we performed 100 independent runs using

100,000 simulations, 40 expectation–maximization cycles, and a broad search range for

parameters to determine the run with the best parameter estimates and maximum likelihood.

Model comparisons were performed using the Akaike Information Criterion (AIC) to identify the

model that best explains the observed data.

Genomic Island of Differentiation

To identify genomic regions under divergent selection, we scanned the genomes of *M.*

pentadactyla and *M. aurita*. We used Pixy v2.0 to calculate nucleotide diversity (π), fixation

index (F_{ST}), and absolute genetic differentiation (d_{XY}) across 20 kb windows (Korunes & Samuk,

2021). Candidate genomic islands of divergence were defined as outlier windows concurrently in

the top 5th percentile for both F_{ST} and d_{XY}. We extracted all genes within these outlier regions.

Genes annotated only with LOC identifiers were annotated based on their high-confidence

human orthologs, identified via BLASTP (Altschul et al., 1997). This gene set was then tested

for Gene Ontology (GO) term enrichment using the R package clusterProfiler (Yu et al., 2012).

Morphometric analyses

We accessed specimens deposited in FMNH, USNM, UMMZ, British Museum of Natural

History, UK (BMNH), Natural History Museum, Tribhuvan University (NHM-TU) Institute of

Zoology, Guangdong Academy of Sciences, China (IOZ-GDAS), and Guangdong Wildlife

Monitoring and Rescue Center, China (GWMRC) (**Table S1**). Fourteen external measurements

were taken including full body length (abdominal) (FBLA), head body length (abdominal)

(BLA), tail length (abdominal) (TLA), full body length (back) (FBLB), head body length (back)

(HBL), tail length (back) (TLB), ear length (EL), ear thickness (ET), middle claw length of the

back foot (MCLBF), middle claw length of the forefoot (MCLFF), the distance of the first scale

from the tip of the nose (DFSTN), the maximum number of scales in the back of the body

(MNS), the single-sided number of tail edge scales (SNTE), length of hind foot (LHF), using soft

tape rulers (accurate to 0.1 mm) for flexible dimensions and MNT-150 standard-type zinc alloy

digital caliper (Model: MNT952101; DEGUQMNT, accuracy to 0.01 mm) for more precise

linear features (**Supplementary Figure S3A**). This data includes 26 skin specimens for *M.*

pentadactyla (n=8), *M. javanica* (n=12), *M. aurita* (n=6) and also includes the holotype of *M.*

aurita (**Data S1**).

Twenty cranial measurements were conducted with MNT-150 standard-type zinc alloy digital

caliper (accuracy to 0.01 mm), following (Gaubert & Antunes, 2005; Wu et al., 2005; Gaudin et

al., 2009; Sulaiman et al., 2017; Igado et al., 2020; Morenikeji et al., 2022) including total skull

length (TSL), nasal to foramen magnum base length (NFBL), premaxilla to palatine (PP), from

Intersection between maxilla and palatine sutures to most posterior point of the palatine

sutures(IMPS), tymapanic bulla (left) (TB), maximum width between lateral edge of auditory

bulla (WSLB), skull height (H), width of Nasal Bone (WN), length of Nasal Bone (LN), dorsal

gap between maxillary (dorsal) (GM), maximum width of Gap between maxillary (MWG), left

maximal length of orbit left） (MO), distance between auditory bulla (DA), length of suture

between parietal bone (DTEM), length of suture between frontal bone (DLOB), zygomatic

process between maxilla and squamous bone left） (ZMS), length of foremen magnun occipitals

(FMO), mandible length (left) (ML), the distance between condylar process (DCP), length of

palatal process of premaxilla(LPP) (**Supplementary Figure S3B**). This data includes 44 skulls

for *M. pentadactyla* (n=25), *M. javanica* (n=7), *Manis crassicaudata* (n=2), *M. culionensis* (n=3)

and *M. auritus* (n=7; **Data S2**). Sex was not considered in the analysis due to the absence of

marked sexual dimorphism and the lack of information for some specimens. Specimen age was

assessed based on body size, skull morphology, and available collection records. Juvenile and

subadult individuals were excluded in the analysis. To ensure completeness of the data, we

replaced missing values with the mean value calculated across specimens of the same species.

We used the function prcomp () of stat to conduct principal component analysis (PCA) and

Morpho to perform canonical variate analysis (CVA) on log□□-transformed variables. Finally,

we conducted Levene's test for homogeneity of variances using the leveneTest () function and

then performed Student's t-tests with t.test () to compare group means across taxa.

We downloaded 84 micro□CT scan data from Ferreira-Cardoso et al. (2019), and scan 13

specimens, obtaining high□resolution micro□CT scans of 97 skulls for *M. pentadactyla* (n = 42),

M. javanica (n = 38), *M. crassicaudata* (n = 12), *M. culionensis* (n = 3), and *aurita* (n = 2; **Table**

S2). Raw DICOM stacks were imported into Materialise Mimics v19.0 for segmentation and

construction of 3D cranial meshes. We placed 75 landmarks per mesh following Ferreira-

Cardoso (2020; **Supplementary Figure S5**). After Procrustes alignment, we conduct principal

component analysis (PCA) and canonical variate analysis (CVA) using MorphoJ v1.08 (Dryden

& Mardia, 1993). Landmark selection was based on prior studies addressing mammalian taxa

(Goswami, 2006; Hautier et al., 2017)

Scale coding

We generated scale-encoded heatmaps from 12 intact skins (nine *M. pentadactyla,* three *M.*

aurita) to visualize variation following Algewatta et al. (2020); Guo et al. (2022) and Martin et

al. (2024). In brief, we digitized the x-y coordinates of every scale in Excel, exported

standardized images into Photoshop v25, and overlaid all specimens per species to produce

species-specific composite maps (**Data S3, S4**).

RESULT

Phylogenetic relationships and distribution of *Manis pentadactyla*

We firstly estimated the Asian pangolins' phylogeny using 68 mitogenomes (**Table S2**). The

maximum likelihood (ML) tree is well-resolved, with every interspecific relationship strongly

supported (i.e., bootstrap support [BS] = 100; **Figure 1A**). The topology aligns well with

previous studies (Hu et al., 2020a; Wang et al., 2022; Gu et al., 2023; Wei et al., 2024; Wangmo

et al., 2025), reaffirming *Manis pentadactyla* sensu lato (including all three subspecies

pentadactyla, *pusilla*, and *aurita*) as the sister clade to all other Asian congeners. The putative

taxon "mysteria" emerges as the immediate sister to *M. javanica. M. pentadactyla* s. l., is

represented by 43 samples, clustered into two deeply diverged clades, corresponding to MPA

and MPB identified in Hu et al. (2020a).

We further estimated maximum-likelihood trees using autosomal and sex-chromosome derived

SNP and single-copy orthologues for a subset of samples (**Table S3**). The phylogenies derived

from all three datasets are topologically identical (**Figure 1A, Supplementary Figures S1, S2**),

mirroring the mitochondrial phylogeny. When projected the SNP of *Manis* species onto

principal-component (PC) space, every *M. pentadactyla* s. l. sample falls into a tight cluster that

is clearly distinct from the other congeneric species (**Figure 1B, 1C**). Restricting the analysis to

M. pentadactyla s. l. reveals non-overlapping splitting between MPA and MPB along PC1

(Tracy-Widom test, p = 1.09×10^{-5}; **Table S4**), underscoring consistent genomic differentiation

despite their close relationship. The result of ADMIXTURE analysis echoes these patterns

(**Figure 1D**). The lowest cross-validation error occurs at K = 2 (CV = 0.50), with marginal

increases at K = 3 and K = 4 (CV = 0.53 in both cases; **Table S5**). At K = 2, the inferred clusters

correspond closely to MPA and MPB, with three individuals displaying minor ancestry from the

opposite cluster. Taken together, the concordant mitochondrial and nuclear results support a

robust validation of the split within *M. pentadactyla* sensu lato.

Figure 1: Phylogenomic relationships and population structure of pangolins. (A) Maximum likelihood (ML) phylogenies based on complete mitochondrial genomes (left) and autosomal nuclear single-nucleotide polymorphisms (SNPs) (right). Numbers at the nodes represent bootstrap support values. The scale bars indicate the number of substitutions per site. (B)

Principal Component Analysis (PCA) of genomic SNP data showing the separation of four *Manis* species. (C) PCA focused on the *M. pentadactyla* s.l. illustrating the clear genetic differentiation between *M. pentadactyla* s.s. (MPA) and *M. aurita* (MPB). Values in parentheses on each axis indicate the percentage of total variance explained by that principal component. (D) Population structure of the *M. pentadactyla* species complex as inferred by ADMIXTURE, for K=2 to K=4 assumed ancestral populations. Each vertical bar represents an individual, and its colored segments show the proportion of that individual's ancestry from each of the K inferred clusters.

To define the geographic distribution of the MPB clade, we augmented our dataset with publicly

available mitochondrial *CYTB* and *COI* sequences representing *M. pentadactyla* s.l. A

phylogenetic analysis of this expanded dataset confirmed the same backbone topology as our

initial trees (**Figure 2A**). Critically, this analysis placed sequences from specimens described as

Manis indoburmanica (from South Tibet) firmly within the MPB clade. Furthermore, every MPB

haplotype with a reliable geographic origin was traced to either Assam, South Tibet, China or

Nepal, a region that includes the collection locality of Hodgson's *aurita* topotype from central

Nepal (**Figure 2B**). This direct association of the MPB clade with the *aurita* topotype provides

evidence to assign this lineage, geographically rooted in Himalaya, to *M. aurita*.

Figure 2:Mitochondrial phylogeny and geographic distribution of the *Manis pentadactyla* species complex. (A) Maximum likelihood phylogeny constructed from an expanded dataset of

mitochondrial sequences (complete mitogenomes and partial CYTB and COX1 fragments),
incorporating publicly available data. Numbers at the nodes represent bootstrap support values.
The scale bar indicates the number of substitutions per site. Note that specimens described as *M.*
indoburmanica (MI, highlighted in green) nest firmly within the *M. aurita* (MPB) clade. (B)
Geographic distribution map for specimens with reliable locality information (see Table S1).
Circles represent data obtained from previous studies while triangles data collected in this study.
Shaded areas denote the approximate ranges of Asian pangolin species based on distribution data
from the IUCN Red List.

Morphometric evidence for the distinctiveness of *M. aurita*

To assess whether *M. aurita* is morphologically distinct from the *M. pentadactyla* sensu stricto

(including subspecies *pentadactyla* and *pusilla*) and thus merits recognition as a separate species,

we employed linear measurements-based and three-dimensional landmark-based geometric

morphometric analyses (**Table S6**). Principal Component Analysis (PCA) of 20 cranial

measurements for 44 specimens positioned *M. pentadactyla* s. l. within a morphospace that

showed partial overlap with the Indian pangolin (*M. crassicaudata*) but remained clearly

separated from the Sunda (*M. javanica*) and Philippine (*M. culionensis*) pangolins (**Figure 3A**;

Supplementary Figures S3; S4A, S4B). Along PC1, which accounted for 39.4% of the shape

variance and primarily reflected overall cranial size, specimens identified as *M. aurita*

consistently occupied the positive extreme. This placement indicates that *M. aurita* possesses a

significantly larger skull, and proportionally shorter palatine when compared to *M. pentadactyla*

s.s. (**Tables S7A-6C**).

Three-dimensional landmark-based geometric morphometric analyses of 75 cranial landmarks

(**Table S6**) obtained from micro-CT scans also clearly distinguished *M. aurita* from *M.*

pentadactyla s.s. along PC3 (**Figure 3B**, **Table S8A-8B**), but not PC1 or PC2 (**Supplementary**

Figures S4C, S4D). Compared to *M. pentadactyla* s.s., *M. aurita* exhibits a relatively shorter yet

broader nasal bone, a laterally expanded junction between the maxilla and palatine bones, a

posteriorly positioned anterior point of the foramen magnum on the ventral occipital surface

(landmark 25, see **Table S6**; **Supplementary Figure S5**), and an anteriorly shifted M-shaped

suture between the maxilla and palatine bones in ventral view, culminating in a shorter ventral

maxilla (**Supplementary Figure S10C**).

To further substantiate the morphological separation between *M. aurita* and *M. pentadactyla* s.s.,

we conducted a PCA based on fourteen external body measurements including *M. aurita* and *M.*

pentadactyla s.s. and *M. javanica.* A crucial aspect of this analysis was the inclusion of the *M.*

aurita holotype (Natural History Museum, BMNH: 43.1.12.85), which notably lacks skull

material. The results demonstrated a clear distinction: all specimens from South Tibet,

Northeastern India and Nepal, including the *M. aurita* holotype, separated unequivocally from

M. pentadactyla s.s. along both PC1 (accounting for 52.1% of variance) and PC3 (6.9% of

variance) (**Figure 3C**; **Table S9A-9C**, **Supplementary Figures S4E, S4F**). The PCA-based

separation is strongly supported by *t*-tests on individual cranial (**Figure 3D**, **Supplementary**

Figure S6A; Table S10) and external measurements (**Figure 3E, Supplementary Figure S6B,**

Table S11), which show *M. aurita* having significantly longer body lengths, tail lengths, larger

hind feet, an extended snout and markedly smaller ears.

Finally, Canonical Variate Analyses (CVA) that using both cranial and external variables

achieved 100% classification accuracy in separating *M. aurita* and *M. pentadactyla* s.s. (**Tables**

S12, S13, 14; Supplementary Figure. S7). In contrast to these clear differences in skeletal and

external body proportions, an analysis of scale characteristics using superimposed, scale-encoded

heatmaps of 12 intact body skins (nine *M. pentadactyla* s. s. and *three M. aurita*) revealed no

significant differences in either the total number of scale rows or their spatial distribution

between the two taxa (**Supplementary Figure S8; Table S15**). Collectively, the consistent and

pronounced size- and shape-based differences in both cranial and external morphology, strongly

demonstrate that *M. aurita* is morphologically distinct from *M. pentadactyla* s.s. supporting its

full species status.

Figure 3: Morphometric analyses based on cranial and external morphology. (A) Principal
Component Analysis (PCA) based on 20 linear cranial measurements (PC1 vs. PC2). (B) PCA of
cranial shape variation derived from 75 3D geometric morphometric landmarks (PC2 vs. PC3).
(C) PCA based on 14 external body measurements (PC1 vs. PC2). For all PCA plots (A-C),
values in parentheses on the axes indicate the percentage of variance explained. Note the
inclusion of the *M. aurita* holotype (BMNH: 43.1.12.85) in panel C, which clusters firmly with
other *M. aurita* specimens. (D, E) Boxplots comparing key cranial (D) and external (E)
measurements between the two taxa. Boxes depict the interquartile range (IQR) with the median
indicated by the central line; whiskers extend to 1.5 times the IQR. Units are in millimeters (D)
and centimeters (E). Statistical significance from Student's t-tests is denoted as: * $P < 0.05$; ** $P
< 0.01$; *** $P < 0.001$. See Tables S5–S14 and Figures S4, S9, S10 for additional results.

Evolutionary and demographic history of *Manis aurita*

We estimated divergence time of pangolins using 200 selected clock-like single-copy orthologous genes. The most recent common ancestor (MRCA) of the genus *Manis* was dated to approximately 20.1 million years ago (Ma) (95% confidence interval [CI] = 17.5 to 22.6 Ma; **Figure 4A**), a finding consistent with Gu et al. (2023). *M. aurita* (MA hereafter) and *M. pentadactyla* s. s. (MP hereafter) diverged approximately 1.2 Ma (95% CI = 0.9 to 1.7 Ma), during the early Pleistocene. Pairwise Sequentially Markovian Coalescent (PSMC) analyses revealed MP maintained a larger effective population size (N□) than MA following their divergence (**Figure 4B**). After their split, the trajectory for MP was characterized by an initial decline, a subsequent period of prolonged stabilization since 1.0 Ma and a final, sudden drop around 50 ka. The trajectory for MA was broadly similar, showing stability until approximately 300 ka, at which point it entered a sustained decline.

As both species experienced population declines and possess small effective population sizes (N_e), we evaluated whether population decline resulted from inbreeding. The histogram of ROH length (**Figure 4C**) shows that individuals of MP carry more medium- and long-tract ROHs (>100 kb) than those of MA, indicating a higher degree of recent inbreeding. Consistent with this, the median F_{ROH} in MP is ~0.0447, twice that of MA (~0.0228) (**Figure 4D**). This finding confirms that MP has undergone more recent inbreeding which may be related to the abrupt Ne drop in the recent past. It is also noteworthy that some individuals in both species exhibit very high FROH (> 0.1), suggesting pronounced inbreeding in certain populations such as *M aurita* near Kathmandu Valley (e.g., Dhading).

Figure 4: Divergence time, demographic dynamics, and genomic divergence of *Manis* species. (A) Divergence time estimates for pangolins based on 200 clock-like single-copy orthologous genes, inferred using *BEAST3. Numbers at nodes represent mean age estimates in millions of years (Ma), while the associated horizontal bars represent the 95% Confidence Intervals (CI). Asterisks indicate nodes used for calibration. (B) Historical effective population size (N_e) trajectories for three species inferred using PSMC. Shaded areas represent 100 bootstrap replicates. (C-D) Comparison of recent inbreeding levels between *M. aurita* (MA) and *M. pentadactyla* s.s. (MP), showing (C) the distribution of Runs of Homozygosity (ROH) lengths and (D) the genome-wide fraction of ROH (F_{ROH}). (E-F) Interspecific gene flow analyses among four *Manis* species. MC: *M. crassicaudata*, MJ: *Manis javanica*. (E) Heatmap of Dsuite f-branch statistics, where warmer colors indicate stronger signals of admixture on a given branch. (F)

TreeMix population graph showing inferred drift and a migration event (arrow). (G-H) Genomic
differentiation between *M. aurita* and *M. pentadactyla* s.s. (G) Manhattan plot of FST and dXY
values, with dashed lines indicating the 95% outlier thresholds used to define genomic islands of
divergence. (H) Gene Ontology (GO) enrichment analysis of genes within the identified genomic
islands, revealing significant enrichment for terms related to olfactory functions. Dot size
corresponds to gene count and color to statistical significance (p.adjust).

We subsequently investigated patterns of interspecific gene flow involving MA to assess the

impact of current and historical range overlap with its congeners (**Figure 2B**). D-suite analyses

indicated significant, albeit low-level, introgression from the Indian pangolin (*M. crassicaudata*)

into both MP (f4-ratio ≈ 0.0025, p < 1x10-15) and MA (f4-ratio ≈ 0.0022, p < 1x10-15) (**Figure**

4E; Table S16). Evidence of introgression from MA into the Malayan pangolin (*M. javanica*)

was also detected by Dsuite (f4-ratio ≈ 0.0017) and TreeMix (**Figure 4F**). In contrast, when we

specifically examined gene flow between MP and MA post-divergence using fastsimcoal2, the

analysis strongly favored a scenario of no gene flow over alternative models (ΔAIC > 2.9; **Table**

S17). Collectively, these results indicate that while MA has experienced minimal introgression

from other pangolin species, it has remained largely isolated from its sister species MP since

their divergence.

Finally, we compared MA and MP genomes for signatures of local adaptation by calculating

nucleotide diversity (π) and fixation index (F_{ST}), and absolute genetic differentiation (D_{XY})

(**Supplementary Figure S9**). Genomic islands of divergence were defined as windows in the top

5 % for both F_{ST} (>0.8575) and D_{XY} (>0.0373; **Figure 4G**). Gene Ontology (GO) enrichment

analysis of genes within these identified islands revealed significant enrichment for terms related

to olfactory functions, notably "detection of chemical stimulus involved in sensory perception of

smell" and "sensory perception of smell" (**Figure 4H**).

DISCUSSION

The Revalidation of *M. aurita*

Despite accumulating molecular evidence suggesting overlooked, or cryptic species (Hu et al., 2020a; Wang et al., 2022; Wei et al., 2024), the taxonomy of *M. pentadactyla* s. l. lacked rigorous examination. Our mitochondrial and genome-wide phylogenomic analyses consistently reveal a divergence between *aurita* and *M. pentadactyla* s.s. at about 1.3 Ma (**Figure 4A**). The magnitude of this split surpasses commonly accepted thresholds for intraspecific variation (Rieseberg & Burke, 2001; Abreu et al., 2025; Van Elst et al., 2025). Complementing this genetic divergence, morphometric analyses identify consistent and diagnostic differences that distinguish *aurita* from *M. pentadactyla* s.s. (**Figure 3**). Cranially, *M. aurita* possesses a significantly larger overall skull with a proportionally shorter and broader nasal region compared to *M. pentadactyla* s.s. (**Supplementary Figures S10, S11, Table S9, Table S10**). External morphological differences are particularly clear and facilitate differentiation. For instance, *aurita* has obviously smaller ears than *M. pentadactyla,* a characteristic reflected in its original specific epithet (*aurita*, Latin for eared). On the other hand, our phylogenetic analysis fails to resolve *M. p. pentadactyla* and *M. p. pusilla* as two distinct evolutionary entities (**Figure 1A**).

The distribution of *M. aurita* is associated with the Himalayan region (**Figure 2B**). It includes the type locality of *M. aurita* from central Nepal, as well as individuals from South Tibet recently described as *Manis indoburmanica* (Wangmo et al., 2025). In contrast, *M. pentadactyla* s.s. is predominantly found in South and Eastern China and Southeast Asia. This geographical separation is likely influenced by biogeographic barriers, such as the rivers run in north-south directions (e.g., the Brahmaputra River) (Lehman et al., 2006; Hu et al., 2020b) and the Arakan Mountains (Hinckley et al., 2024).

Collectively, the congruent evidence from our genetic, morphological, and distributional

analyses unequivocally supports recognizing Hodgson's *aurita* as a distinct species. While

Zijlstra (2025) previously argued for its revalidation, our study substantiates this view with

broader and more integrative evidence. Consequently, in accordance with the Principle of

Priority of the International Code of Zoological Nomenclature (ICZN), we reinstate *Manis aurita*

Hodgson, 1836 as the valid name for this species. This action renders junior synonyms such as

Pholidotus assamensis Fitzinger, 1872, and *Manis indoburmanica* Wangmo, Ghosh, Dolker,

Joshi, Sharma and Thakur, 2025.

Evolutionary history of *M*. aurita

The population genetic and demographic analyses illuminate the evolutionary trajectory of *M*.

aurita following its speciation. The ancestors of *M. aurita* and *M. pentadactyla* diverged in the

early Pleistocene, period of, significant global climatic oscillations. Given that pangolins are

generally warm-adapted species, with no historical fossil evidence suggesting their presence in

northern China (Tong, 2007), the climatic shifts likely played a crucial role in driving the

allopatric separation of the ancestor into western (Himalaya) vs. eastern (East/ South East Asia)

refugium (Louys & Meijaard, 2010; Leonard et al., 2015; Hawkins et al., 2016; Camacho-

Sanchez et al., 2017; Hinckley et al., 2023). Pangolins are highly olfactory-reliant mammals,

exhibiting significantly enlarged olfactory bulb (Imam et al., 2018; Li et al., 2024), using smell

to forage for ants and termites (DiPaola et al., 2020). As the *M. aurita* and *M. pentadactyla*

adapted to their respective environments, certain OR genes, certain OR genes likely diverged

under selection as supported in our GO enrichment analysis (**Figure 4H**).

The population of *M. aurita* decline started around 300 Ka and is consistent with the Penultimate

Glaciation (~300–130 Ka), during which the climate cooled and dried (Colleoni et al., 2016; Ma

et al., 2022), likely leading pangolin habitats to contract. Notably, similar patterns are seen in

other warm-adapted mammals in southern China (Li et al., 2025), such as the golden snub-nosed

monkeys (Dai et al., 2024).

Our analyses further detected limited gene flow between *M. aurita* and two other Asian

pangolins. The signal of introgression between *M. aurita* and *M. crassicaudata* (contributing to

<0.25% of the genome) is unsurprising, given their sympatric ranges in southern Nepal and

northeastern India. We also observed a similarly low level of gene flow between *M. aurita* and

M. javanica, indicating that their distributions likely have overlapped at least historically.

Despite these instances of historical admixture, the low level of gene flow indicates *M. aurita*

has maintained its status as a distinct evolutionary entity.

It is noteworthy, in the admixture analysis, at both K=3 and K=4, *M. aurita* separated into two

distinct genetic clusters: one comprising individuals from Nepal and the other comprising

individuals from South Tibet and northeastern India, a pattern that indicates significant

intraspecific subdivision (**Figure 1D**).

Conservation Implications

The formal recognition of *M. aurita* as a distinct species carries profound conservation

significance. Its intrinsic vulnerability is underscored by a geographically restricted range,

confined to the Himalayan–Indo-Burmese region, and a demographic history marked by a more

pronounced and sustained population decline since the Middle Pleistocene compared to *M.*

pentadactyla.

Despite the long-term population decline revealed by PSMC analyses, *M. aurita* on average

exhibits lower levels of recent inbreeding than *M. pentadactyla* as evidenced by the results of

ROH and FROH (**Figures 4C, 4D**). This likely reflects differing histories of anthropogenic

pressure nowadays; the less intensively developed landscapes of South Asia historically provided

more contiguous refuge habitats compared to the range of *M. pentadactyla* in China and

Southeast Asia, where has experienced intense human activity for centuries, leading to severe

habitat fragmentation and the regional extinction of many medium-to-large mammal species

(Brook et al., 2014; Goossens et al., 2016; Boonratana, 2020). However, it is noteworthy, *M.*

aurita around Kathmandu valley exhibit extremely high inbreeding levels (i.e., FROH > 0.1),

indicating that these populations are already facing severe inbreeding depression (**Figure**. **4D**).

Conservation strategies must therefore prioritize the identification and management of these

highly vulnerable populations to prevent local extinctions.

Overarching these intrinsic vulnerabilities is the critical threat from illegal wildlife trade. Genetic

analyses of confiscated scales have repeatedly confirmed that *M. aurita* is a target (Hu et al.,

2020a). Alarmingly, our findings also reveal that products from this species have infiltrated

regulated traditional medicine markets (data from Xie et al. (2021); **Supplementary Figure S13**),

demonstrating that illegally sourced materials are being laundered through formal supply chains.

The comprehensive genomic dataset presented here, with its improved geographic coverage and

inclusion of previously unsampled populations, provides an enhanced genetic baseline which

will bolster wildlife forensics by allowing for greater precision in determining the geographic

provenance of confiscated scales, thereby helping to pinpoint poaching hotspots.

Taxonomy

Manis aurita Hodgson, 1836 stat. nov.

Manis auritus Hodgson, 1836. Original description.

Manis aurita Waterhouse,1838

Phatages bengalensis Fitzinger, 1872

Pholidotus assamensis Fitzinger, 1872

Manis pentadactyla auritus Thomas 1918

Manis pentadactyla Ellerman & Morrison-Scott, 1951 (part)

Manis indoburmanica Wangmo, Ghosh, Dolker, Joshi, Sharma and Thakur, 2025

Holotype: BMNH 43.1.12.85, an adult (skin) collected by Brian Houghton Hodgson in 1836

from Central Nepal (**Supplementary Figure S11B**).

Emended diagnosis

Manis aurita can be diagnosed by the following combination of characters that distinguish it

from all other Asian pangolins: the external pinna is small and distinctively flake-shaped. On the

snout, the first row of head scales lies well behind the nasal tip, exposing a bare patch of naked

skin (**Supplementary Figure S10A**); dorsal scales are smooth and trapezoidal with straight

posterior margins, whereas the lateral scales near the abdomen are broad and shield-shaped

(**Supplementary Figure S10B**). Cranially, the palatal process of the premaxilla extends

posteriorly but measures less than half of the premaxillary midline length, and the incisive

foramen does not extend to the premaxilla's posterior border. At its anterior end, the maxilla

bulges laterally, creating nearly parallel lateral margins (**Supplementary Figure S10C**). The

postglenoid foramen opens laterally within the central region of the postglenoid fossa.

(**Supplementary Figure S10D**). Both the jugular and hypoglossal foramina are recessed yet

remain distinct. Finally, the ventral surface of the occipital condyle is flat rather than convex

(**Supplementary Figure S10C**).

Description

Manis aurita is an Asian pangolin with an elongated head, small ears, short limbs, and a tail shorter than half the body length. Its external pinna is small and flake-shaped. The head and body are covered in overlapping, exceptionally smooth keratin scales: dorsal scales are trapezoidal with straight posterior edges, while lateral abdominal scales are broad and shield-shaped. On the snout, the first row of head scales lies well behind the nasal tip, leaving a bare patch of skin. The ventral surface is pale brown and densely fringed with short, stiff hair bristles. Claws are moderately curved: the third (middle) claw on each forefoot is the longest, the first and fifth claws are noticeably shorter, and all hind-foot claws are shorter than those on the forefeet. Scale coloration in adults ranges from dark brown to chestnut; juveniles are slate-grey or grey-brown, with more bristles between scales.

The skull exhibits a relatively elevated and broad profile. The ventral palatal process of the premaxilla is short, measuring less than half the length of the premaxilla itself. The incisive foramen of the premaxilla does not contact the ventral projection of the maxilla. The vomer does not protrude at the midline on the ventral surface of the maxilla. The nasal bones are broad and terminate in a semicircular shape. The postglenoid foramen is positioned laterally. The pterygoid hamulus is short and does not extend beyond the ectotympanic bulla, which is rounded in shape. The palatine sutures form an M-shaped structure. The zygomatic process of the maxilla may either be connected to the zygomatic process of the squamosal via a bony rod-like structure or remain unconnected. The external and skull measurements are presented in **Supplementary Data 1, 2**, respectively.

The total number of body scales ranges from 538 to 596. From the neck to the tail base there are 254–274 scales arranged in 23–24 longitudinal and 15–17 transverse rows. The tail bears 140–153 scales: 80–94 on the dorsal surface organized into five transverse by 16–20 longitudinal

rows, and 50–63 on the ventral surface in three to five transverse by 18–20 longitudinal rows,

with 17–19 scales along each lateral edge. The forelimbs carry 37–55 scales per side, while the

hindlimbs have 32–41 scales per side.

Comparisons

Manis aurita (MA) and *M. pentadactyla* s.s. (MP) share a close morphological resemblance,

exhibiting a similar number of body scales, typically uniform dark brown to deep chestnut. These

scales are generally shield-shaped or trapezoidal and are interspersed with sparse hair bristles

(**Supplementary Figure S11A**). This contrasts with *M. javanica*, whose scales on the tail tip, tail

base, and ventral side are pale yellow, and which uniquely features a reversed growth direction

in the last longitudinal scale row on its forelimbs. *M. crassicaudata* is set apart by its larger,

earth-toned scales and the absence of interspersing hair bristles. *M. culionensis* is characterized

by more numerous, elongated, and sharper-tipped scales that are dark brown and either leaf-

shaped or shield-shaped. In MA and MP, the zygomatic process of the maxilla and the

squamosal is sometimes connected via a bony rod to form a zygomatic arch. This contrasts with

all other species, that lack a complete zygomatic arch. The palatal process of the premaxilla in

MA and MP is notably short, not exceeding half the premaxilla's midline length, unlike in other

species, where this process is longer (**Supplementary Figure S11B**). Additionally, the

postglenoid foramen in MA and MP opens on the lateral aspect of the zygomatic process,

differing from *M. javanica* and *M. culionensis*, where it opens posteriorly, and from *M.*

crassicaudata, where its opening can be either posterior or lateral (**Supplementary Figure**

S11B).

Compared to MP, MA typically attains a larger adult body size, with a total body length ranging

from 81.0 to 105.1 cm (mean: 95.2 cm) compared to 60.5 to 80.6 cm in MP (mean: 71.2 cm).

MA also possesses a longer tail (31.7–40.9 cm, mean: 37.4 cm) than MP (22.3–31.6 cm, mean:

27.5 cm), and has larger foot surfaces, with hindfoot lengths averaging 6.4 cm versus 2.9 cm in

MP. Furthermore, the external pinnae differ, with those of MA being distinctly smaller than in

MP (**Figure**. 3E; **Supplementary Figure S6B**). The skull of MA is larger and taller than that of

MP, with cranial width ranging from 93.2 to 99.7 mm (mean: 95.6 mm) and height from 35.2 to

36.3 mm (mean: 35.8 mm), compared to MP's cranial width of 75.9 to 95.9 mm (mean: 84.6 mm)

and height of 23.8 to 31.5 mm (mean: 27.6 mm). Additionally, the maximum width of the gap

between the maxillary bones is wider in MA (13.5–14.7 mm, mean: 13.8 mm) than in MP (10.1–

12.0 mm, mean: 10.9 mm). The skull of MA is larger and taller than that of MP. This is reflected

in a relatively smaller nasal length-to-width ratio in MA (approximately 1.9:1), indicating a

broader nasal bone, whereas MP has a narrower nasal bone with a larger ratio (approximately

2.5:1), resulting in a comparatively slenderer nasal region (**Figure. 3D, Supplementary Figure**

S6A). The anterior portion of the skull in MA is also broader and deeper; its maxillary region

expands laterally, with the posterior lateral margins of this anterior expansion running nearly

parallel. In contrast, the anterior part of the skull in MP is more elongated, contributing to a more

triangular overall shape of the ventral skull surface. The palatine bone in MA is positioned more

anteriorly than in MP. Specifically, the most posterior point of the palatine sutures in MA aligns

approximately with the maxilla zygomatic process and squamosal zygomatic process junction. In

MP, however, this point lies posterior to this junction. The occipital condyle also shows clear

differences: in MP, it is distinctly prominent, whereas MA presents a relatively flat ventral

condylar articular surface that extends posteriorly. The condylar articular facet is comparatively

smaller in MP than in MA (**Supplementary Figure S11A**).

Distribution, habitat, and Conservation Statues

Manis aurita is distributed along the southern Himalayan foothills, with confirmed occurrences in Nepal, South Tibet and northeastern India (including Assam), southern Bhutan, and extending into southern Tibet. The species' altitudinal range spans from lowland areas to approximately 3,000 meters above sea level. The eastern extent of its distribution remains undefined, but the Arakan Mountains appear to represent the eastern geographic limit for this species (Hinckley et al., 2024). Further surveys are necessary to ascertain its potential presence in Myanmar.

Manis aurita predominantly inhabits mixed deciduous and evergreen forests. Key tree species in these habitats include *Shorea robusta, Schima wallichii, Castanopsis indica*, and *Alnus nepalensis. M. aurita* is also frequently observed in forested areas adjacent to human-modified landscapes, indicating some level of adaptability to disturbed environments (Dhital et al., 2020; Shrestha et al., 2021).

Despite protected under national laws such as the Nepal's National Parks and Wildlife Conservation Act (1973), India's Wildlife Protection Act (Schedule I, 1972), and internationally under CITES Appendix I, *M. aurita* remains a significant target for illegal wildlife traffickers in South Asia. For example, Kathmandu functions as a major transit hub for smuggled pangolin scales (Bashyal et al., 2021; Koju & Tamang, 2025). Similar trafficking pressures are evident in neighboring India and Bhutan, where challenges such as limited enforcement capacity and porous international borders facilitate cross-border smuggling operations.

DATA AVAILABILITY

All sequencing data were deposited in the National Center for Biotechnology Information (NCBI) database (BioProjectID …………………), and Genome Sequence Archive (GSA) database (https://ngdc.cncb.ac.cn/gsa/) (accession number ………………….).

SUPPLEMENTARY DATA

Supplementary data to this article can be found online.

COMPETING INTERESTS

The authors declare that they have no competing interests

AUTHOR CONTRIBUTIONS

AUTHOR CONTRIBUTIONS

NPK: Conceptualization, Methodology, Investigation, Formal analysis, Visualization,

Validation, Writing – original draft, Writing – review & editing. **ZZ**: Methodology,

Investigation, Formal analysis, Visualization, Validation, Writing – original draft, Writing –

review & editing. **GZ**: Methodology, Investigation, Formal analysis, Visualization, Validation,

Writing – original draft, Writing – review & editing. **XH**: Methodology, Investigation, Formal

analysis, Visualization, Validation. **ZY**: Methodology, Investigation, Formal analysis,

Visualization, Validation. **XW**: Methodology, Investigation, Formal analysis, Visualization,

Validation. **MTRH**: Formal analysis, Validation, Writing – original draft, Writing – review &

editing. **CG**: Formal analysis, Validation, Writing – original draft, Writing – review & editing.

AH: Formal analysis, Validation, Writing – original draft, Writing – review & editing. **MFCF**:

Formal analysis, Validation, Writing – original draft, Writing – review & editing. **JL**: Formal

analysis, Validation, Writing – original draft, Writing – review & editing. **DM**: Methodology,

Investigation, Formal analysis, Visualization. **SB:** Methodology, Investigation, Formal analysis,

Visualization **LH**: Formal analysis, Validation, Writing – original draft, Writing – review &

editing. **WY**: Formal analysis, Validation, Writing – original draft, Writing – review & editing.

LL: Formal analysis, Validation, Writing – original draft, Writing – review & editing. **AF**:

Methodology, Investigation, Resources, Visualization, Validation, Writing – review & editing.

All authors read and approved the final version of the manuscript. **YH**: Methodology,

Investigation, Resources, Visualization, Validation, Supervision, Funding acquisition, Project

administration, Writing – review & editing. All authors read and approved the final version of

the manuscript. **KH**: Conceptualization, Methodology, Investigation, Resources, Visualization,

Validation, Supervision, Funding acquisition, Project administration, Writing – review & editing.

All authors read and approved the final version of the manuscript.

ACKNOWLEDGMENTS

We thank the following institutions for providing access to specimens for morphological and

genetic analyses: The National Museum of Natural History, Smithsonian Institution (USNM);

Field Museum of Natural History (FMNH); American Museum of Natural History (AMNH);

University of Michigan Museum of Zoology (UMMZ); the Institute of Zoology, Guangdong

Academy of Sciences (IOZ-GDAS); and the Guangdong Wildlife Monitoring and Rescue Center

(GWMRC), China. We are grateful to the Natural History Museum, Tribhuvan University, and

the Department of National Parks and Wildlife Conservation (Nepal) for providing research

permissions and institutional support. We sincerely thank the Ministry of Education, Science and

Technology, Nepal, and the Nepal Academy of Science and Technology (NAST) for their

support. NPK's effort to this study was partially supported by an ANSO-CAS visiting Scientist

fellowship at Kunming Institute of Zoology, Chinese Academy of Sciences, 2025 and South

China Biodiversity Research Center, School of Life Sciences, Guangzhou University,

Guangzhou, China, 2023. This study is supported by the ADCS Seed Grants program of the

National Museum of Natural History, Smithsonian Institution (M.T.R.H. and A.H.), Guangdong

Provincial Natural Science Foundation for Distinguished Young Scholars (2022B1515020033,

K.H.) and the National Natural Science Foundation of China (32170452, K.H.).

REFERENCES

Abreu, EF, Prado, JR, Maldonado, JE, Wilson, DE, Percequillo, AR, and Pavan, SE. (2025).
Recent, Intricate Speciation in Amazonia Uncovered by a Multilayered Genomic
Analysis of Tree Squirrels. Systematic biology, syaf013.
https://doi.org/10.1093/sysbio/syaf013
Alexander, DH, Novembre, J, and Lange, K. (2009). Fast model-based estimation of ancestry in
unrelated individuals. Genome research, 19(9), 1655-1664.
https://doi.org/10.1101/gr.094052.109
Algewatta, HR, Perera, P, Karawita, H, Dayawansa, N, Manawadu, D, and Liyanage, M. (2020).
Updates on the morphometric characterization of Indian pangolin (*Manis crassicaudata*)
in Sri Lanka. Animals, 11(1), 25. https://doi.org/10.3390/ani11010025
Altschul, SF, Madden, TL, Schäffer, AA, Zhang, J, Zhang, Z, Miller, W, and Lipman, DJ.
(1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs. Nucleic acids research, 25(17), 3389-3402.
https://doi.org/10.1093/nar/25.17.3389
Ballard, JWO, and Whitlock, MC. (2004). The incomplete natural history of mitochondria.
Molecular ecology, 13(4), 729-744. https://doi.org/10.1046/j.1365-294X.2003.02063.x
Bashyal, A, Shrestha, N, Dhakal, A, Khanal, SN, and Shrestha, S. (2021). Illegal trade in
pangolins in Nepal: Extent and network. Global Ecology and Conservation, 32, e01940.
Boonratana, R. (2020). Asian primates in fragments: Understanding causes and consequences of
fragmentation, and predicting primate population viability. American journal of
primatology, 82(4), e23082. https://doi.org/10.1002/ajp.23082
Bouckaert, R, Vaughan, TG, Barido-Sottani, J, Duchêne, S, Fourment, M, Gavryushkina, A,
Heled, J, Jones, G, Kühnert, D, and De Maio, N. (2019). BEAST 2.5: An advanced
software platform for Bayesian evolutionary analysis. PLoS computational biology,
15(4), e1006650. https://doi.org/10.1371/journal.pcbi.1006650
Brook, SM, Dudley, N, Mahood, SP, Polet, G, Williams, AC, Duckworth, J, Van Ngoc, T, and
Long, B. (2014). Lessons learned from the loss of a flagship: the extinction of the Javan
rhinoceros *Rhinoceros sondaicus annamiticus* from Vietnam. Biological Conservation,
174, 21-29. https://doi.org/10.1016/j.biocon.2014.03.014
Camacho-Sanchez, M, Leonard, JA, Fitriana, Y, Tilak, M-K, and Fabre, P-H. (2017). The
generic status of Rattus annandalei (Bonhote, 1903)(Rodentia, Murinae) and its
evolutionary implications. Journal of Mammalogy, 98(5), 1340-1355.
https://doi.org/10.1093/jmammal/gyx081
Challender, D, Wu, S, Kaspal, P, Khatiwada, A, Ghose, A, Ching-Min Su, N, and Suwal, TL.
(2020a). *Manis pentadactyla*. The IUCN Red List of Threatened Species 2019:
e.T12764A168392151. In: Gland: IUCN.
Challender, DW, Heinrich, S, Shepherd, CR, and Katsis, LK. (2020b). International trade and
trafficking in pangolins, 1900–2019. In Pangolins (pp. 259-276). Elsevier.
https://doi.org/10.1016/B978-0-12-815507-3.00016-2

Challender, DW, Nash, HC, and Waterman, C. (2019). *Pangolins: science, society and*
conservation. Academic Press.

Colleoni, F, Wekerle, C, Näslund, J-O, Brandefelt, J, and Masina, S. (2016). Constraint on the
penultimate glacial maximum Northern Hemisphere ice topography ($\approx$ 140 kyrs BP).
Quaternary Science Reviews, 137, 97-112.
https://doi.org/10.1016/j.quascirev.2016.01.024

Corbet, GB, and Hill, JE. (1992). *The mammals of the Indomalayan region: a systematic review*
(Vol. 488). oxford university press Oxford.

Dai, Y, Xia, W, Zhu, Y, Hacker, C, Wang, X, and Li, D. (2024). Historical changes in the
distribution of the Sichuan golden snub‐nosed monkey (*Rhinopithecus roxellana*) in
Sichuan Province, China. Ecology and Evolution, 14(4), e11270.
https://doi.org/https://doi.org/10.1002/ece3.11270

Danecek, P, Auton, A, Abecasis, G, Albers, CA, Banks, E, DePristo, MA, Handsaker, RE,
Lunter, G, Marth, GT, and Sherry, ST. (2011). The variant call format and VCFtools.
bioinformatics, 27(15), 2156-2158. https://doi.org/10.1093/bioinformatics/btr330

Danecek, P, Bonfield, JK, Liddle, J, Marshall, J, Ohan, V, Pollard, MO, Whitwham, A, Keane,
T, McCarthy, SA, and Davies, RM. (2021). Twelve years of SAMtools and BCFtools.
Gigascience, 10(2), giab008. https://doi.org/10.1093/gigascience/giab008

Dhital, S, Paudel, SM, Thapa, S, Bleisch, WV, Shrestha, A, and Koju, NP. (2020). Distribution
of Chinese pangolin (*Manis pentadactyla*) in Nagarjun forest of Shivapuri Nagarjun
National Park, Nepal. Nepalese Journal of Zoology, 4(1), 1-7.
https://doi.org/10.3126/njz.v4i1.30667

DiPaola, JD, Yindee, M, and Plotnik, JM. (2020). Investigating the use of sensory information to
detect and track prey by the Sunda pangolin (*Manis javanica*) with conservation in mind.
Scientific reports, 10(1), 9787. https://doi.org/10.1038/s41598-020-65898-x

Douglas, J, Jiménez-Silva, CL, and Bouckaert, R. (2022). StarBeast3: adaptive parallelized
Bayesian inference under the multispecies coalescent. Systematic biology, 71(4), 901-
916. https://doi.org/10.1093/sysbio/syac010

Dryden, IL, and Mardia, KV. (1993). Multivariate shape analysis. Sankhyā: The Indian Journal
of Statistics, Series A, 460-480. https://www.jstor.org/stable/25050954

Emms, DM, and Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome biology, 16,
1-14. https://doi.org/10.1186/s13059-015-0721-2

Excoffier, L, Marchi, N, Marques, DA, Matthey-Doret, R, Gouy, A, and Sousa, VC. (2021).
fastsimcoal2: demographic inference under complex evolutionary scenarios.
bioinformatics, 37(24), 4882-4885. https://doi.org/10.1093/bioinformatics/btab468

Ferreira-Cardoso, S, Billet, G, Gaubert, P, Delsuc, F, and Hautier, L. (2019). Skull shape
variation in extant pangolins (Pholidota: Manidae): allometric patterns and systematic
implications. Zoological Journal of the Linnean Society, 188(1), 255-275.
https://doi.org/10.1093/zoolinnean/zlz096

Fitzinger, LJFJ. (1872). *Die natürliche Familie der Schuppenthiere (Manes).Sitzungsberichte der*
kaiserlichen Akademie der Wissenschaften.
https://www.biodiversitylibrary.org/part_231767

Fujita, MK, Leaché, AD, Burbrink, FT, McGuire, JA, and Moritz, C. (2012). Coalescent-based
species delimitation in an integrative taxonomy. Trends in ecology & evolution, 27(9),
480-488. https://doi.org/10.1016/j.tree.2012.04.012

Funk, DJ, and Omland, KE. (2003). Species-Level Paraphyly and Polyphyly: Frequency, Causes,
and Consequences, with Insights from Animal Mitochondrial DNA. Annual review of
ecology, evolution, and systematics, 34(1), 397-423.
https://doi.org/10.1146/annurev.ecolsys.34.011802.132421
Gaubert, P, and Antunes, A. (2005). Assessing the taxonomic status of the Palawan pangolin
Manis culionensis (Pholidota) using discrete morphological characters. Journal of
Mammalogy, 86(6), 1068-1074. https://doi.org/10.1644/1545-
1542(2005)86[1068:ATTSOT]2.0.CO;2
Gaubert, P, Antunes, A, Meng, H, Miao, L, Peigné, S, Justy, F, Njiokou, F, Dufour, S, Danquah,
E, and Alahakoon, J. (2018). The complete phylogeny of pangolins: scaling up resources
for the molecular tracing of the most trafficked mammals on earth. Journal of Heredity,
109(4), 347-359. https://doi.org/10.1093/jhered/esx097
Gaudin, TJ, Emry, RJ, and Wible, JR. (2009). The phylogeny of living and extinct pangolins
(Mammalia, Pholidota) and associated taxa: a morphology based analysis. Journal of
mammalian evolution, 16, 235-305. https://doi.org/10.1007/s10914-009-9119-9
Gomez, L, Joseph, T, Heinrich, S, Wright, B, and D'Cruze, N. (2023). Illegal trade of pangolins
in India with international trade links: an analysis of seizures from 1991 to 2022.
European Journal of Wildlife Research, 69(4), 85. https://doi.org/10.1007/s10344-023-
01708-9
Goossens, B, Sharma, R, Othman, N, Kun-Rodrigues, C, Sakong, R, Ancrenaz, M, Ambu, LN,
Jue, NK, O'Neill, RJ, and Bruford, MW. (2016). Habitat fragmentation and genetic
diversity in natural populations of the Bornean elephant: Implications for conservation.
Biological Conservation, 196, 80-92. https://doi.org/10.1016/j.biocon.2016.02.008
Goswami, A. (2006). Cranial modularity shifts during mammalian evolution. The American
Naturalist, 168(2), 270-280. https://doi.org/10.1086/505758
Gu, T-T, Wu, H, Yang, F, Gaubert, P, Heighton, SP, Fu, Y, Liu, K, Luo, S-J, Zhang, H-R, and
Hu, J-Y. (2023). Genomic analysis reveals a cryptic pangolin species. Proceedings of the
National Academy of Sciences, 120(40), e2304096120.
https://doi.org/10.1073/pnas.2304096120
Guo, C, Jinyao, L, Siyuan, L, Yuze, J, Chunfang, L, Yan, H, Bo, L, Li, L, Songyuan, T, and
Yanchun, X. (2022). Coding System and Comparisons of Scales Between Malayan
Pangolin(*Manis javanica*) and Chinese Pangolin(*M. pentadactyla*). Chinese Journal of
Wildlife, 43(1), 024－031. https://doi.org/10.19711/j.cnki.issn2310-1490.2022.01.003
Hautier, L, Billet, G, De Thoisy, B, and Delsuc, F. (2017). Beyond the carapace: skull shape
variation and morphological systematics of long-nosed armadillos (genus *Dasypus*).
PeerJ, 5, e3650. https://doi.org/10.7717/peerj.3650
Hawkins, MT, Leonard, JA, Helgen, KM, McDonough, MM, Rockwood, LL, and Maldonado,
JE. (2016). Evolutionary history of endemic Sulawesi squirrels constructed from UCEs
and mitogenomes sequenced from museum specimens. BMC evolutionary biology, 16, 1-
16. https://doi.org/10.1186/s12862-016-0650-z
He, K, and Jiang, X. (2014). Sky islands of southwest China. I: an overview of phylogeographic
patterns. Chinese Science Bulletin, 59, 585-597. https://doi.org/10.1007/s11434-013-
0089-1
Heighton, SP, Allio, R, Murienne, J, Salmona, J, Meng, H, Scornavacca, C, Bastos, AD,
Njiokou, F, Pietersen, DW, and Tilak, M-K. (2023). Pangolin genomes offer key insights

and resources for the world's most trafficked wild mammals. Molecular biology and
evolution, 40(10), msad190.
Hinckley, A, Hawkins, MT, Maldonado, JE, and Leonard, JA. (2023). Evolutionary history and
patterns of divergence in three tropical east Asian squirrels across the Isthmus of Kra.
Journal of Biogeography, 50(6), 1090-1102. https://doi.org/10.1111/jbi.14598
Hinckley, A, Maldonado, JE, Tamura, N, Leonard, JA, and Hawkins, MT. (2024). Lost in
synonymy: Integrative species delimitation reveals two unrecognized species of Southern
Asian tree squirrels (Rodentia: *Sciuridae: Callosciurinae*). Vertebrate Zoology, 74, 683-
707. https://doi.org/10.3897/vz.74.e133467
Hodgson, BH. (1836). New Mammals in Nipdlese Catalogue. The journal of the Asiatic Society
of Bengal, V, 234-235. https://www.biodiversitylibrary.org/bibliography/51678
Hogan, F, Wedrowicz, F, Pd. Khatiwada, A, Dev Joshi, J, Wasser, S, and Wright, W. (2025).
Genetic evidence reveals distinct lineage of Chinese pangolin in Nepal: insights from scat
and blood samples for conservation and wildlife forensics. Ecology and Evolution, 15(2),
e70982. https://doi.org/10.1002/ece3.70982
Hu, J-Y, Hao, Z-Q, Frantz, L, Wu, S-F, Chen, W, Jiang, Y-F, Wu, H, Kuang, W-M, Li, H, and
Zhang, Y-P. (2020a). Genomic consequences of population decline in critically
endangered pangolins and their demographic histories. National Science Review, 7(4),
798-814. https://doi.org/10.1093/nsr/nwaa031
Hu, Y, Thapa, A, Fan, H, Ma, T, Wu, Q, Ma, S, Zhang, D, Wang, B, Li, M, and Yan, L. (2020b).
Genomic evidence for two phylogenetic species and long-term population bottlenecks in
red pandas. Science Advances, 6(9), eaax5751. https://doi.org/10.1126/sciadv.aax575
Igado, O, Adekanmbi, J, Adebayo, M, Sulaimon, O, and Osundina, O. (2020). Facial and Body
Morphometric Indices of the Arboreal African Pangolin (*Manis tricuspis*):
Anthropometric Tools in Describing the Pangolin: Facial and biometric indices of the
African pangolin. Sahel Journal of Veterinary Sciences, 17(3), 28-33.
https://doi.org/10.54058/saheljvs.v17i3.122
Imam, A, Bhagwandin, A, Ajao, MS, Spocter, MA, Ihunwo, AO, and Manger, PR. (2018). The
brain of the tree pangolin (*Manis tricuspis*). II. The olfactory system. Journal of
Comparative Neurology, 526(16), 2548-2569. https://doi.org/10.1002/cne.24510
IUCN SSC Pangolin Specialist Group. (2018). *Regional Sunda Pangolin (Manis javanica)*
Conservation Strategy 2018-2028. IUCN SSC. Pangolin Specialist Group.
https://pangolinsg.org/wp-content/uploads/sites/4/2020/01/Pangolin-SG-et-
al_2018_Regional-Sunda-pangolin-conservation-strategy-2018-2028.pdf
Katoh, K, and Standley, DM. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Molecular biology and evolution, 30(4),
772-780. https://doi.org/10.1093/molbev/mst010
Koju, NP. (2018). *Abundance and phylogenetic status of Chinese Pangolin (Manis pentadactyla)*
from Kathmandu valley, Nepal.
Koju, NP, He, K, Chalise, MK, Ray, C, Chen, Z, Zhang, B, Wan, T, Chen, S, and Jiang, X.
(2017). Multilocus approaches reveal underestimated species diversity and inter-specific
gene flow in pikas (*Ochotona*) from southwestern China. Molecular Phylogenetics and
Evolution. https://doi.org/https://doi.org/10.1016/j.ympev.2016.11.005
Koju, NP, and Tamang, BM. (2025). Illegal Wildlife Trade in Kathmandu Valley: A Hub of
Biodiversity Crime. Ecology and Evolution (unpublished).

Korunes, KL, and Samuk, K. (2021). pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. Molecular ecology resources, 21(4), 1359-1368. https://doi.org/10.1111/1755-0998.13326

Lehman, SM, Fleagle, JG, Meijaard, E, and Groves, CP. (2006). The geography of mammals and rivers in mainland Southeast Asia. In Primate Biogeography. Developments in Primatology: Progress and Prospects (pp. 305-329). Springer. https://doi.org/10.1007/0-387-31710-4_11

Leonard, JA, den Tex, RJ, Hawkins, MT, Muñoz‑Fuentes, V, Thorington, R, and Maldonado, JE. (2015). Phylogeography of vertebrates on the Sunda Shelf: a multi‑species comparison. Journal of Biogeography, 42(5), 871-879. https://doi.org/10.1111/jbi.12465

Li, H, and Durbin, R. (2011). Inference of human population history from individual whole-genome sequences. Nature, 475(7357), 493-496. https://doi.org/10.1038/nature10231

Li, H, Handsaker, B, Wysoker, A, Fennell, T, Ruan, J, Homer, N, Marth, G, Abecasis, G, Durbin, R, and Subgroup, GPDP. (2009). The sequence alignment/map format and SAMtools. bioinformatics, 25(16), 2078-2079. https://doi.org/10.1093/bioinformatics/btp352

Li, J, Guo, C, Xie, M, Wang, K, Wang, X, Zou, B, Hou, F, Ran, C, Bi, S, and Xu, Y. (2024). Genomic signatures of sensory adaptation and evolution in pangolins. BMC genomics, 25(1), 1-13. https://doi.org/10.1186/s12864-024-11063-x

Li, Z, Fan, H, Liao, Z, Wang, Y, and Wei, F. (2025). Global spatiotemporal patterns of demographic fluctuations in terrestrial vertebrates during the Late Pleistocene. Science Advances, 11(21), eadq3938. https://doi.org/10.1126/sciadv.adq3938

Louys, J, and Meijaard, E. (2010). Palaeoecology of Southeast Asian megafauna‑bearing sites from the Pleistocene and a review of environmental changes in the region. Journal of Biogeography, 37(8), 1432-1449. https://doi.org/10.1111/j.1365-2699.2010.02297.x

Ma, T, Tarasov, PE, Huang, K, Leipe, C, Man, M, and Zheng, Z. (2022). Intensified climate drying and cooling during the last glacial culmination (20.8–17.5 cal ka BP) in the south-eastern Asian monsoon domain inferred from a high-resolution pollen record. Quaternary Science Reviews, 278, 107371. https://doi.org/10.1016/j.quascirev.2022.107371

Malinsky, M, Matschiner, M, and Svardal, H. (2021). Dsuite‑Fast D‑statistics and related admixture evidence from VCF files. Molecular ecology resources, 21(2), 584-595. https://doi.org/10.1111/1755-0998.13265

Martin, JM, Buckley, JY, Connelly, E, Hywood, L, Lacey, LM, Ruden, RM, Ruhagazi, D, and Wearn, A. (2024). The Pangolin Universal Notching System: a scale-marking methodology for pangolins. Oryx, 1-7. https://doi.org/1017/S0030605324000656

McKenna, A, Hanna, M, Banks, E, Sivachenko, A, Cibulskis, K, Kernytsky, A, Garimella, K, Altshuler, D, Gabriel, S, and Daly, M. (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome research, 20(9), 1297-1303. https://doi.org/10.1101/gr.107524.110

McQuillan, R, Leutenegger, A-L, Abdel-Rahman, R, Franklin, CS, Pericic, M, Barac-Lauc, L, Smolej-Narancic, N, Janicijevic, B, Polasek, O, and Tenesa, A. (2008). Runs of homozygosity in European populations. The American journal of human genetics, 83(3), 359-372. https://doi.org/10.1016/j.ajhg.2008.08.007

Meng, G, Li, Y, Yang, C, and Liu, S. (2019). MitoZ: a toolkit for animal mitochondrial genome assembly, annotation and visualization. Nucleic acids research, 47(11), e63-e63. https://doi.org/10.1093/nar/gkz173.

Morenikeji, O, Omotosho, O, Jarikre, T, Emikpe, B, Fawole, E, Adeyefa, A, and Olapade, B.
(2022). Morphometric characterisation of white-bellied pangolins rescued from trade for
conservation in Nigeria. Emerging Animal Species, 1, 100018.
https://doi.org/10.1016/j.eas.2022.100018
Nijman, V, Zhang, MX, and Shepherd, CR. (2016). Pangolin trade in the Mong La wildlife
market and the role of Myanmar in the smuggling of pangolins into China. Global
Ecology and Conservation, 5, 118-126. https://doi.org/10.1016/j.gecco.2015.12.003
Padial, JM, Miralles, A, De la Riva, I, and Vences, M. (2010). The integrative future of
taxonomy. Frontiers in zoology, 7, 1-14. https://doi.org/10.1186/1742-9994-7-16
Patterson, N, Price, AL, and Reich, D. (2006). Population structure and eigenanalysis. PLoS
genetics, 2(12), e190. https://doi.org/10.1371/journal.pgen.0020190
Purcell, S, Neale, B, Todd-Brown, K, Thomas, L, Ferreira, MA, Bender, D, Maller, J, Sklar, P,
De Bakker, PI, and Daly, MJ. (2007). PLINK: a tool set for whole-genome association
and population-based linkage analyses. The American journal of human genetics, 81(3),
559-575. https://doi.org/10.1086/519795
Rambaut, A, Suchard, M, and Xie, D. (2017). Drummond AJ. 2014. Tracer v1. 6. In.
Rieseberg, LH, and Burke, JM. (2001). The biological reality of species: gene flow, selection,
and collective evolution. Taxon, 50(1), 47-67. https://doi.org/10.2307/1224511
Schlick-Steiner, BC, Steiner, FM, Seifert, B, Stauffer, C, Christian, E, and Crozier, RH. (2010).
Integrative taxonomy: a multisource approach to exploring biodiversity. Annual review
of entomology, 55(1), 421-438. https://doi.org/10.1146/annurev-ento-112408-085432
Shrestha, A, Bhattarai, S, Shrestha, B, and Koju, NP. (2021). Factors influencing the habitat
choice of pangolins (*Manis* spp.) in low land of Nepal. Ecology and Evolution, 11(21),
14689-14696. https://doi.org/10.1002/ece3.8156
Shrestha, S, Bashyal, A, Dhakal, A, McGreevy Jr, TJ, Buffum, B, Joshi, J, Chaudhary, HK, and
Khanal, SN. (2020). Mitochondrial DNA analysis of critically endangered Chinese
Pangolins (*Manis pentadactyla*) from Nepal. Mitochondrial DNA Part B, 5(3), 3257-
3261. https://doi.org/10.1080/23802359.2020.1811174
Smith, SA, Brown, JW, and Walker, JF. (2018). So many genes, so little time: a practical
approach to divergence-time estimation in the genomic era. Plos one, 13(5), e0197433.
https://doi.org/10.1371/journal.pone.0197433
Sulaiman, MH, Azmi, WA, Hassan, M, and Chong, JL. (2017). Current updates on the
morphological measurements of the Malayan pangolin (*Manis javanica*). Folia
Zoologica, 66(4), 262-266. https://doi.org/10.25225/fozo.v66.i4.a8.2017
Toews, DP, and Brelsford, A. (2012). The biogeography of mitochondrial and nuclear
discordance in animals. Molecular ecology, 21(16), 3907-3930.
https://doi.org/10.1111/j.1365-294X.2012.05664.x
Van Elst, T, Sgarlata, GM, Schüßler, D, Tiley, GP, Poelstra, JW, Scheumann, M, Blanco, MB,
Aleixo-Pais, IG, Rina Evasoa, M, and Ganzhorn, JU. (2025). Integrative taxonomy
clarifies the evolution of a cryptic primate clade. Nature Ecology & Evolution, 9(1), 57-
72. https://doi.org/10.1038/s41559-024-02547-w
Wang, Q, Lan, T, Li, H, Sahu, SK, Shi, M, Zhu, Y, Han, L, Yang, S, Li, Q, and Zhang, L.
(2022). Whole-genome resequencing of Chinese pangolins reveals a population structure
and provides insights into their conservation. Communications Biology, 5(1), 821.
https://doi.org/10.1038/s42003-022-03757-3

Wangmo, LK, Ghosh, A, Dolker, S, Joshi, BD, Sharma, LK, and Thakur, M. (2025). Indo-
997   Burmese pangolin (*Manis indoburmanica*): a novel phylogenetic species of pangolin
998   evolved in Asia. Mammalian biology, 1-8. https://doi.org/10.1007/s42991-024-00475-7
999   Wei, S, Fan, H, Zhou, W, Huang, G, Hua, Y, Wu, S, Wei, X, Chen, Y, Tan, X, and Wei, F.
1000   (2024). Conservation genomics of the critically endangered Chinese pangolin. Science
1001   China Life Sciences, 1-11. https://doi.org/10.1007/s11427-023-2540-y
1002   Wilson, DE, and Mittermeier, RA. (2018). *Handbook of the mammals of the world: vol. 8:*
1003   *insectivores, sloths and colugos*. Lynx Edicions.
1004   Wu, S, Naifa, L, Zhang Yingmei, Zhi, O, and Hai, C. (2005). Measurement and Comparison of
1005   Skull Variables in Chinese Pangolin and Malayan Pangolin. ACTA THERIOLOGICA
1006   SINICA, 24(3), 211. https://doi.org/10.16829/j.slxb
1007   Xie, X, Ye, H, Cai, X, Li, C, Li, F, Tian, E, and Chao, Z. (2021). DNA mini-barcodes, a
1008   potential Weapon for Conservation and combating illegal trade of pangolin. Tropical
1009   Conservation Science, 14, 19400829211017361.
1010   https://doi.org/10.1177/1940082921101736
1011   Yu, G, Wang, L-G, Han, Y, and He, Q-Y. (2012). clusterProfiler: an R package for comparing
1012   biological themes among gene clusters. Omics: a journal of integrative biology, 16(5),
1013   284-287. https://doi.org/10.1089/omi.2011.0118
1014   Zhang, F, Wu, S, and Cen, P. (2022). The past, present and future of the pangolin in Mainland
1015   China. Global Ecology and Conservation, 33, e01995.
1016   https://doi.org/10.1016/j.gecco.2021.e01995
1017   Zhang, F, Yu, Y, Wu, S, Mahmood, A, Yu, J, and Min, Y. (2020). Reducing pangolin demand
1018   by understanding motivations for human consumption in Guangdong, China. Frontiers in
1019   Ecology and Evolution, 8, 574161. https://doi.org/10.3389/fevo.2020.574161
1020   Zhang, M, Gouveia, A, Qin, T, Quan, R, and Nijman, V. (2017). Illegal pangolin trade in
1021   northernmost Myanmar and its links to India and China. Global Ecology and
1022   Conservation, 10, 23-31. https://doi.org/10.1016/j.gecco.2017.01.006
1023   Zijlstra, JS. (2025). Manis aurita Hodgson 1837 as a valid species of pangolin: a comment to
1024   Wangmo et al.(2025). Mammalian biology, 1-3. https://doi.org/10.1007/s42991-025-
1025   00495-x
1026   Zoological Society of London. (2014). *Plight of the pangolin: all eight species to move up*
1027   *EDGE list. EDGE of Existence*. Retrieved September 12 from
1028   https://www.edgeofexistence.org/blog/plight-of-the-pangolin-all-eight-species-move-up-
1029   edge-list/

1030