Taurine is a Natural Host Cytoadhesion Inhibitor in Asymptomatic

Malaria Cases

Authors: Gretchen Diffendall[1*], Perine Millot[2], Patty Chen[1], Dominique Dorin-Semblat[2], Fanny

Aprahamian[3], Sylvère Durand[3], Benoît Gamain[2], Antoine Claessens[4,5*] and Artur Scherf [1*]

Affiliations: *[1]Institut Pasteur, Biology of Host-Parasite Interactions Unit, INSERM U1201,*

CNRS EMR9195 ; Paris, France.

[2]Sorbonne Université, INSERM, CNRS, Centre d'Immunologie et des Maladies Infectieuses,

CIMI F-75013 ; Paris, France.

[3]Metabolomic platform of Gustave Roussy, U1138/UMS AMMICa (US 23 INSERM / UMS

3655 CNRS); Paris, France

[4]LPHI, MIVEGEC, CNRS, INSERM, University of Montpellier, France

[5]Medical Research Council Unit The Gambia at the London School of Hygiene and Tropical

Medicine, Banjul, The Gambia

*Corresponding authors: artur.scherf@pasteur.fr, antoine.claessens@umontpellier.fr,

gdiffendall@hsph.harvard.edu

One Sentence Summary:

Metabolomics uncovers taurine as a natural host inhibitor of *Plasmodium falciparum*

cytoadhesion in asymptomatic malaria carriers, offering new insights for disease control.

Abstract:

The prolonged dry season in malaria-endemic regions of sub-Saharan Africa can be divided

into periods of high and low transmission. The majority of symptomatic malaria cases are

restricted to the short high transmission period that follows the rainy season. Shortly after,

persistent asymptomatic malaria cases are more prevalent throughout the prolonged low

transmission period. It is still unclear whether host metabolic alterations play a role in

asymptomatic infections during seasonal malaria. In this study, we analyzed the blood plasma

metabolome (n=199) of individuals in The Gambia, West Africa, capturing data from both

high and low malaria transmission periods. Plasma samples from individuals (n=16) were

collected monthly throughout the low transmission season, enabling a longitudinal analysis of

metabolic alternations over six months. Our findings reveal that significant changes in host

plasma metabolite composition are associated with seasonality and malaria pathogenicity.

Notably, we observed elevated levels of taurine in asymptomatic malaria infections,

especially during periods of low transmission. *In vitro*, this naturally occurring host molecule

inhibits the cytoadhesion of malaria-infected red blood cells (iRBCs), which is key to malaria

disease severity and mortality. Exogenous taurine can significantly reduce or reverse binding

of iRBCs to the common adhesion receptor CD36 and the endothelial protein C receptor

(EPCR), the later being associated with cerebral malaria. This study uncovers a mechanism

by which elevated taurine plasma levels in subclinical infections could reduce cytoadhesion

and lead to increased splenic clearance, thereby strengthening host resistance to symptomatic

infections. In the absence of health strategies targeting dry season parasite reservoirs, our

findings highlight taurine as a potential prophylactic or therapeutic agent to reduce

symptomatic malaria in sub-Saharan Africa.

Main Text:

INTRODUCTION

Malaria, a mosquito-borne disease, remains a significant global health issue, causing nearly

600,000 deaths annually, with the majority occurring among children in sub-Saharan Africa [1].

Most fatalities are due to infection by *Plasmodium falciparum*, which produces adhesive

proteins known as PfEMP1 on the surface of infected red blood cells (iRBCs) [2]. These

proteins enable the parasite to bind to endothelial cells in various organs in a process called

cytoadhesion, helping it to evade clearance by the spleen. The accumulation of parasites in

organs is associated with symptomatic malaria, and when this occurs in the brain, it can lead

to cerebral malaria, a more severe form of the disease [3]. Due to its central role in virulence,

PfEMP1 is one of the most extensively studied surface protein families, consisting of 60

members per genome. By switching expression between these family members, iRBCs can

exhibit different adhesive properties and antigenic variants, enabling them to evade the

immune system and to sequester in different organs [4,5].

The extracellular domain of PfEMP1 proteins is comprised of 2-10 Duffy-binding-like (DBL)

domains and cysteine-rich interdomain regions (CIDR) [6-9]. The majority of PfEMP1

members contain a DBL domain at the N-terminus followed by a CIDR domain (called head

structure) followed by a variable number of domains that form the adhesive portion exposed

on the surface of iRBCs. Individual domains commonly act as specific receptor-binding

regions of endothelial surface receptors [10]. Despite the significant sequence diversity among

PfEMP1 family members, the N-terminal CIDRα domains retain a structural configuration

that allows binding to either CD36 (CIDRα2-6) [11,12] or EPCR (CIDRα1) [10,13,14]. CD36 is

predicted to be the most common adhesive phenotype in the PfEMP1 family, since

approximately 84% of PfEMP1 proteins bind CD36 [10]. This scavenger receptor, commonly

expressed on endothelial and innate immune cells, interacts with iRBCs from laboratory

strains and field isolates. Apart from preventing splenic clearance, binding to CD36

expressed on dendritic cells may also inhibit the host innate immune response to infection

[15,16]. A subset of 5-7 PfEMP1 members are predicted to bind to EPCR. The CIDRα1

domain's interaction with EPCR on brain endothelial cells [17,18] has been associated with

severe childhood malaria [13,19] and notably cerebral malaria [20,21]. This interaction is a key

target for major vaccine efforts aimed at preventing severe malaria.

The analysis of *P. falciparum* gene expression in cultured parasites uncovered a pre-

programmed, just-in-time transcription cascade that orchestrates asexual blood stage

development [22,23]. Superimposed on this regulatory framework is the heterochromatin-driven

stochastic derepression of clonally variant gene families, which underpins the mechanism of

PfEMP1-mediated antigenic variation [24,25]. The transcriptional program can be influenced by

the addition or depletion of external metabolites in controlled *in vitro* culture conditions or

using an experimental mouse malaria model. Several studies have demonstrated that

environmental sensing is a crucial aspect of the pathogen, modulating its proliferation,

virulence gene expression, and developmental pathways [26–31]. In the context of environmental

sensing, a phenomenon known as seasonal malaria in Sub-Saharan countries is of particular

interest [32,33]. In these malaria-endemic regions, asymptomatic subclinical *P. falciparum*

infections during periods of low transmission alternate with symptomatic infections during

periods of high transmission. The influence of environmental changes on parasite virulence

regulation and clinical manifestations remains a compelling area of investigation. Several

epidemiological studies investigating factors that decrease symptomatic infections have been

performed in Mali. The antibody response in asymptomatic individuals in Mali did not

provide evidence that the immune response plays a major role in subclinical infections [33].

However, the same authors observed an extended blood circulation time of infected red blood

cells (iRBCs) in asymptomatic individuals, suggesting that enhanced clearance by the spleen

reduces parasite load and clinical symptoms. The factors contributing to this phenotype

remain unknown [33–36].

Over the course of a year, The Gambia can be divided into periods of high transmission (late

September through mid-December), when mosquitoes are more prevalent to transmit *P.*

falciparum parasites to new human hosts, and periods of low transmission (late December

through mid-September), when transmission is reduced (Fig. 1A) [37]. Both high and low

transmission periods occur during the dry season (mid-September through mid-June) whereas

the wet season is limited to late June through early September. After the high rain levels

cease in September, symptomatic infections are limited to the high transmission months. A

sizeable fraction of infections remain chronic, resulting in asymptomatic infections being

detected throughout the year, not limited to a specific period [38].

Although metabolomic analyses of human plasma from *P. falciparum*-infected individuals,

have been conducted, most have focused on the effects of disease severity [39,39–43] rather than

on seasonal variations in malaria. To our knowledge, only one recent study [36] has examined

the impact of persistent asymptomatic infections on host plasma metabolites in Mali.

However, its small sample size may have limited the ability to detect significant metabolic

alterations.

In this study, we explored whether the dry season affects the host plasma environment of

infected individuals, with a particular focus on identifying metabolites associated with

disease severity and seasonal impact on asymptomatic infections. Metabolite profiling was

performed on human plasma samples (n = 199) collected from individuals in The Gambia,

including those infected with *P. falciparum*—both symptomatic and asymptomatic—as well

as non-infected volunteers, as part of a previously conducted two-part study[37,44].

Our results reveal significant changes in metabolite levels associated with malaria severity—

comparing symptomatic and asymptomatic infections during the high transmission season—

as well as with seasonality, by comparing asymptomatic infections across high and low

transmission periods. Notably, one of the most highly elevated metabolites in subclinical

infections, consistently increased in subclinical infections, was taurine, an amino sulfonic

acid. While taurine is synthesized in the liver from methionine and cysteine, it is not a

component of proteins but plays a role in various physiological host functions [reviewed in

[45]. Although taurine showed no effect on iRBCs development of cultured parasites, it
strongly inhibited iRBCs adhesion to CD36 and EPCR receptors as well as by a human cell
line expressing CD36 at concentrations observed in dry season-infected individuals.
Additionally, taurine was able to reverse iRBCs binding to these receptors. The identification
of a natural host molecule that inhibits iRBC binding to CD36 and EPCR offers a novel
therapeutic and prophylactic strategy to target symptomatic malaria and reduce dry-season
reservoirs.
141

RESULTS
Study design and metabolomics analysis
To investigate host metabolic changes during malaria infection, we analyzed a total of 199
plasma samples collected from individuals living in neighboring villages in The Gambia (Fig.
1A) as part of a previously published longitudinal study [37,44]. The cohort design included two
complementary arms: (1) During the high transmission season (October–November), active
case detection yielded 46 blood samples from asymptomatic *P. falciparum* carriers, while
individuals presenting at local health posts yielded 38 samples from symptomatic malaria
case. (2) During the low transmission season (December–May), a subset of 16 asymptomatic
individuals with persistent *P. falciparum* infections was followed monthly for up to 6 months.
Also included were 9 unique asymptomatic individuals that had samples taken in December.
Additionally, 11 non-infected individuals were sampled between December to February.
154

As expected, parasite loads were significantly higher in symptomatic individuals (Fig. S1C).
Although participants were not age-matched across groups, the overall age distribution was
representative of the local population (median age: 19 years, Fig. S1A–B, [37]). Metabolomics
was performed on a total 199 plasma samples using gas chromatography coupled to triple
quadrupole mass spectrometry (GC–QqQ/MS), targeting a broad panel of compounds. After
quality filtering, 78 metabolites were retained for analysis (Fig. 1B), spanning diverse
chemical classes including amino acids, fatty acids, and carbohydrates (Fig. 1C). These
metabolites play a role in various metabolic pathways, determined by pathway enrichment
analysis (Fig. 1D, SM3). Initial Principal component analysis (PCA) revealed clustering of
samples according to infection status and season (Fig. 1E).
165

Metabolites associated with seasonality

We assessed the influence of seasonality on host metabolic responses by comparing asymptomatic infections from the peak transmission period (October) to those from the end of the dry season (May), when malaria transmission is nearly absent. Ten metabolites differed significantly between these timepoints (SM5). Specifically, allantoin ($p = .0019$), palmitoleic acid ($p = 0.0095$), glycerol-3-phosphate ($p = 0.0100$) , 4-hydroxymandelic acid ($p = 0.0108$) were elevated in asymptomatic infections during the high transmission period while taurine ($p = 0.0002$), tryptophan ($p = 0.0018$), guanosine ($p = 0.0038$), glyceric acid ($p = 0.0082$), maltose-trehalose ($p = .0264$) and o-phosphoethanolamine ($p = 0.0391$) were significantly increased in May. The results for these metabolites listed in Fig. 2A, indicate that the dry season influences host plasma metabolites, possibly through changes in diet or microbiota composition.

Metabolites associated with symptomatic malaria

To identify metabolic signatures associated with malaria symptoms, we compared 38 individuals with symptomatic infections to 46 asymptomatic infections collected during the same high transmission months (October–November). Three metabolites were significantly elevated in symptomatic individuals: (1) 2-oxoglutaric acid (more commonly known as alpha-ketoglutarate) that is part of the tricarboxylic acid (TCA) cycle, previously identified as a biomarker of *P. falciparum* infection [40]. (2) Palmitoleic acid, a monounsaturated fatty acid that may accumulate during *P. falciparum* infection [46]. (3) 3-hydroxybutyric acid, also a fatty acid, was found to accumulate in host plasma as a result of acidosis induced by the parasite [47] (see Fig. 2A). These findings reveal a distinct metabolic host signature associated with symptomatic malaria, highlighting disruptions in energy metabolism and fatty acid pathways during acute *P. falciparum* infection.

Metabolite dynamics during chronic asymptomatic infection in the dry season

We took advantage of the longitudinal monthly sampling from 16 individuals with persistent
asymptomatic infections over the dry season (December to May) to examine the temporal
evolution of plasma metabolites. Five metabolites: guanosine, maltose;trehalose, taurine,
threonic acid, and glyceric acid showed consistent increase across the dry season, particularly
from March onward (Fig. S2A). Allantoin showed the opposite pattern, with high levels early
in the dry season and lower levels later (Fig. S2A). These patterns likely reflect a
combination of dietary shifts and host physiological adaptation. For instance, allantoin, an
oxidative product of urate, is a known marker of oxidative stress; maltose;trehalose and
threonic acid are diet-derived; guanosine and taurine can be either diet-derived or
endogenously synthesized, with taurine playing a protective role against oxidative stress [48].

Notably, while guanosine, maltose;trehalose, and threonic acid were not significantly
elevated in asymptomatic individuals compared to uninfected controls (Fig. 2B), taurine (Fig.
2C), and to a lesser extent, glyceric acid (Fig. 2C) were increased during the dry season,
suggesting infection-specific upregulation. To assess whether the host metabolome
differentiates asymptomatic carriers from uninfected individuals, we analyzed samples
collected in January (the month with the most available data) comparing 16 asymptomatic
individuals to 11 uninfected controls. Thirteen metabolites were significantly altered in the
asymptomatic group: nine were elevated (including taurine) and four were reduced (Fig.
S2B). These findings show a dynamic shift in specific plasma metabolites during chronic
asymptomatic infection across the dry season likely reflecting dietary changes and host
physiological adaptations, including responses to oxidative stress.

This dataset motivated us to identify metabolites potentially involved in modulating *P.*
falciparum persistence and virulence. We prioritized metabolites meeting three progressively
less stringent criteria: (1) elevated levels in asymptomatic individuals compared to uninfected
controls, (2) lower levels than in symptomatic cases, and (3) a consistent increase across the
dry season. Taurine emerged as the strongest candidate, fulfilling all three conditions. These
findings led us to further investigate the functional role of taurine in malaria pathogenesis, a
direction supported by a recent study showing that taurine supplementation in a *P.*
berghei mouse model conferred host benefits through immune modulation [49].

Taurine inhibits ability of iRBCs to cytoadhere to CD36 and EPCR

Given the key role of iRBC adhesion in malaria pathogenesis, our goal was to identify
candidate metabolites significantly elevated in asymptomatic infections during the low
transmission period that may influence *P. falciparum* sequestration to endothelial receptors.
Additionally to taurine, threonic acid, a breakdown product of ascorbic acid, was selected due
to previous findings that ascorbic acid significantly inhibits *P. falciparum* growth; however,
its potential effect on iRBC cytoadhesion has not yet been studied [50]. We next investigated
whether taurine and threonic acid could reduce iRBC cytoadhesion in vitro. To establish
physiologically relevant high and low taurine concentrations for the cultures, we measured
the absolute taurine quantity in plasma samples exhibiting the highest and lowest values from
our mass spectrometry analysis (Fig. 3A). These taurine concentrations aligned with levels
reported in human plasma [51] and confirmed the mass spectrometry results. Due to the absence
of an assay kit for threonic acid, a reported physiological concentration of 0.27 mM [52] was
used for all analyses. Growth curve analysis of wild-type *P. falciparum* cultures
supplemented with or without various taurine concentrations, or threonic acid [0.27mM] (Fig.
S3A) revealed no effect of taurine nor threonic acid on growth over the course of five days.
These data are consistent with what was observed in another study [53] at even higher
concentrations of taurine. Physiological taurine levels ranging from 0.64 to 0.93 mM have
been observed in clinical volunteer studies following oral administration of 4 g of taurine [54].

To determine if cytoadhesion of iRBCs is affected by taurine or threonic acid, we performed
cytoadhesion assays with RBCs infected with late-stage wild-type 3D7 parasites grown in
media supplemented with a range of taurine concentrations for up to 40 hours (Fig. 3B, S3B).
The lowest taurine concentration of [0.02mM] already significantly decreased cytoadhesion
of iRBCs to CD36 ($p < 0.0001$) by up to 60% (Fig. 3C, D). With even higher taurine
concentrations, binding was drastically decreased by > 97%. Thus, physiological levels of
plasma taurine found in asymptomatic individuals almost completely blocks binding of
iRBCs to recombinant CD36 receptor *in vitro*. Threonic acid, at physiological levels of
[0.27mM] [52], had no significant effect on cytoadhesion of iRBCs to CD36 (Fig. S3C). As
metabolomic MS analysis revealed no significant differences in methionine levels between
conditions (SM4), a physiological concentration of 0.04 mM was used as a control, which
also had no significant effect on cytoadhesion (Fig. 3B)[55]. Incubation of iRBCs with
[0.16mM] of taurine for one hour prior to the cytoadhesion assays was sufficient to decrease
cytoadhesion by > 97% (Fig. 3B), suggesting that taurine influences cytoadhesion through

direct interaction with the iRBC surface, rather than by altering parasite transcription or
protein expression on the surface of iRBC.

The surface PfEMP1 domains that bind to CD36 and EPCR are associated with a single
CIDR domain [56]. We next set out to determine if taurine affects iRBC binding to the
endothelial cell protein C receptor (EPCR), which is important in cerebral malaria [13]. We
used a parasite strain (FCR3) that predominantly express the Itvar19 PfEMP1 previously
found to bind to EPCR [57]. After a one-hour incubation with [0.16mM] or [0.64mM] taurine,
we observed a significant decrease in cytoadhesion of FCR3 iRBCs to recombinant EPCR of
almost 50% and >70%, respectively (Fig. 3D). This parasite line, which also expresses a
distinct PfEMP1 member was additionally used for cytoadhesion assays with CD36 (Fig.
S3D). Taurine supplementation similarly reduced receptor binding across strains, indicating
that taurine inhibitory effect is not strain specific and is effective despite differences in *var*
gene family repertoires.

In contrast to adhesion of infected red blood cells to CD36 and EPCR, which involves diverse
PfEMP1 variants containing CIDR domains, adhesion to placental chondroitin sulfate A
(CSA) is mediated by a unique PfEMP1 encoded by the *var2CSA* gene, which comprises
multiple DBL domains but lacks a CIDR domain [56]. We performed cytoadhesion assays with
the NF54 parasite strain that expresses VAR2CSA [58] in the presence or not of [0.64mM]
taurine (Fig. S3E). A small but significant decrease in iRBC binding was observed (p =
0.0288). Finally we assessed the capacity of malaria naïve human serum to inhibit iRBC
binding to CD36. Human serum caused a slight reduction in our iRBC binding assay (Fig.
S3F), comparable to the effect of 0.02 mM taurine supplementation (Fig. 3B), likely due to
the taurine content in the serum used for parasite culture, which reached up to 0.02 mM. Our
findings show that physiological taurine levels inhibit iRBC adhesion to CD36 and EPCR,
pointing to a novel host-mediated mechanism contributing to reduced *P. falciparum* virulence
in subclinical infections.

284 Taurine reverses adhesion of iRBCs to CD36 and EPCR
To assess its therapeutic potential, we set out to determine if taurine could reverse, rather than
merely prevent, iRBC binding to CD36 (Fig. 4A) and EPCR (Fig. 4B), using the same
strains from (Fig. 3B, D). When iRBCs already adhering to CD36 or EPCR were incubated
without or with [0.64mM] taurine for 45 minutes, cytoadhesion decreased by over 77% (p <

0.0001) to CD36 and nearly 30% (p = 0.0017) to EPCR (Fig. 4A, B). These demonstrate that

taurine can disrupt existing interactions of iRBCs with endothelial receptors.

To explore the mechanisms by which taurine inhibits and reverses iRBC binding to

endothelial receptors, we examined PfEMP1 expression. Since a one-hour taurine incubation

was sufficient to nearly abolish completely cytoadhesion, it is unlikely that transcriptional

changes could account for such rapid effects on protein expression. Indeed, FACS analysis of

FCR3 Itvar19 iRBCs supplemented with taurine showed no significant changes in surface

PfEMP1 protein levels (Fig. S4A, S4E, F). We additionally tested whether taurine interacts

directly with CD36 to prevent iRBC binding. Cytoadhesion assays were performed with

CD36 receptors that were pre-treated with or without [0.16mM] taurine and no significant

difference in cytoadhesion of iRBCs was observed between the two conditions (Fig. S4B).

The observed rapid action of taurine suggests it may directly target the iRBC surface,

inducing conformational changes in parasite adhesins and thereby altering binding properties.

Taurine reduces binding of iRBCs to CD36 on human C32 melanoma cells

To investigate taurine inhibition in a cellular context, we employed the human C32

amelanotic melanoma cell line, which has been used extensively in previous *P. falciparum*

CD36 adhesion studies [59]. This cell line shares adhesion receptors with human endothelial

cells, and the mechanisms of binding are comparable, making it a suitable surrogate for

endothelial cells in adhesion assays [60,61]. Parasitized erythrocytes primarily adhere to these

melanoma cells via the glycoprotein CD36, which is also widely expressed on vascular

endothelium [62]. In three biological replicates we observed > 60% inhibition of FCR3 strain

adhesion to C32 at taurine concentration of 0.93 mM. This inhibition was dose-dependent

(Fig. 4C, D). Similar C32 inhibition results were obtained using a genetically distinct strain

(3D7) (Fig. S4C, D). Taurine inhibits iRBC adhesion at the cellular level, albeit at higher

concentrations than with recombinant receptors. Its effectiveness in reducing iRBC binding

by multiple methods, highlights its potential as a natural supplement to lessen disease severity

in endemic areas.

DISCUSSION

The mechanisms by which *P. falciparum* persists through the rainy season without causing

symptoms remain unclear. In this study, we investigated if environmental factors contribute

to subclinical and persistent *P. falciparum* infections during the dry season in sub-Saharan

Africa. To do so, we conducted a metabolomics analysis on plasma samples collected from

infected and non-infected individuals in The Gambia, West Africa, from both high and low

transmission times during the dry season (Fig. 1). Of the statistically validated metabolites

identified (n=78), nine showed distinct profiles in the cohort (Fig. 2C). We identified

metabolites that are dependent on disease severity or seasonality. Metabolites associated with

severity include fatty acids, which have been shown to increase in *P. falciparum* infections,

as well as a previously identified biomarker involved in the TCA cycle [40]. Metabolites linked

to seasonality include those potentially derived from the diet, as well as metabolites produced

endogenously by the host to combat oxidative stress.

Since the development and proliferation of *P. falciparum* rely on nutrient uptake from the

host serum [63], the observed changes in the host metabolome may directly affect the parasite.

However, it remains unclear whether the altered metabolite levels—either elevated or

reduced—are taken up by infected red blood cells (iRBCs) and subsequently influence

parasite development. Of note, a recent report suggests that external lactate levels, a major

cause of parasite-induced acidosis, may serve as metabolic sensors in *P. falciparum* iRBCs [64].

An increase in environmental lactate was linked to elevated levels of epigenetic regulators,

sirtuins [65], and histone lysine lactylation [64] in the parasite. In our study, lactate levels were

significantly lower in uninfected individuals and asymptomatic infections compared to those

with symptomatic infections during the high transmission period. Another major contributor

to parasite-induced acidosis, 3-hydroxybutyric acid, was found significantly increased in our

symptomatic high transmission infections [47].

One of the most striking metabolic changes observed was in taurine levels, which were as

low as 0.02 mM during periods of high transmission and rose to 0.16 mM by the end of the

low transmission season in infected individuals (Fig. 3A). For comparison, oral

administration of taurine (4 g) in eight healthy volunteers increased plasma levels from 0.03–

0.06 mM to peak concentrations of 0.53–0.93 mM for several hours after taurine

administration [54]. The cause of elevated taurine levels in individuals with asymptomatic

infections remains unclear. Although the human liver synthesizes taurine, dietary sources

such as meat, fish, poultry, and dairy also contribute to plasma levels. In rural Gambia, diets

mainly consist of rice, wheat-flour bread, and groundnuts, with seasonal availability of leafy

vegetables, millet, and dried fish. Since there's no evidence of increased protein intake during

the dry season in this cohort, we hypothesize that elevated plasma taurine levels are due to

increased endogenous bile secretion. Notably, persistent high host-derived taurine levels have

been reported for extended periods following the resolution of enteric infections in mice [66]. It

would be important to investigate whether changes in the gut microbiome play a role in

stimulating increased bile taurine production in this cohort. A recent study using a mouse

malaria model (*P. berghei*) suggested that taurine, when combined with artemisinin, may

enhance host defenses against infection. Although *P. berghei* lacks *var* genes, the findings

indicate a potential role for taurine in strengthening the immune response to help control the

infection [67].

Building on our initial observation, we explored whether taurine could play a direct role in

the development of subclinical infections. To investigate this, we focused on two key factors

implicated in asymptomatic infections: the growth of iRBCs and their binding to host

endothelial receptors. External taurine concentrations were selected based on levels reported

in our cohort study and in human volunteers after a 4 g taurine oral dose [54]. Although taurine

treatment did not affect parasite growth (Fig. S3A), it strongly inhibited the binding of iRBCs

to recombinant receptor CD36 (Fig. 3B). Taurine also inhibited the binding of iRBCs to the

brain endothelial receptor EPCR (Fig. 3D), which is closely associated with severe malaria

[10,13,14], although the inhibition levels were slightly lower than those observed for CD36.

Conversely, taurine had a very mild effect on cytoadhesion of RBCs infected with parasites

expressing an atypical PfEMP1 family member (VAR2CSA) with a unique sequence that has

no CIDRα domain (Fig. S3E). Importantly, we demonstrate that taurine also inhibits iRBC

adhesion at the cellular level, though at higher concentrations than using recombinant

receptors. We used the human C32 amelanotic melanoma cell line, a well-established

surrogate for endothelial cells in *P. falciparum* adhesion assays due to its shared adhesion

receptors and comparable binding mechanisms.

Since preincubation of CD36 with taurine did not impair iRBC binding (Fig. S4B), we

hypothesize that taurine disrupts parasite adhesion by interfering with PfEMP1-mediated

receptor interactions. Three key findings support this hypothesis. First, a one-hour incubation

of adhesive iRBCs with taurine prior to the CD36 adhesion assay markedly reduced binding

(Fig. 3B). Second, taurine pretreatment did not alter PfEMP1 surface expression (Fig. S4A),

indicating that its effect is not due to changes in protein trafficking. Finally, the distinct

inhibitory effect of taurine on CD36 and EPCR binding supports a mechanism that
specifically disrupts PfEMP1 interactions, rather than broadly affecting iRBC membrane
components shared across all PfEMP1 variants.

Moreover, we observed the capacity of taurine to revert adhesion to CD36 (>77%) and EPCR
(>29%) (Fig. 4A, B). Thus, taurine shows blocking activity to different PfEMP1s. In the case
of CD36, we have tested different CD36-binding strains (3D7 and FCR3) and have not
observed differences in the binding inhibition to this receptor despite the sequence variability
that occurs in the CIDRα domain between different family members and different strains [11].

A study conducted in Mali[33] provided evidence that increased circulation time of infected red
blood cells (iRBCs) is associated with asymptomatic infections during the dry season,
although the underlying mechanism was not elucidated. Our findings from The Gambia offer
mechanistic support for these observations, suggesting that elevated plasma taurine levels in
persistent asymptomatic infections may reduce or reverse iRBC binding, thereby promoting
the circulation of mature parasite forms and their clearance by the spleen. In a recent small-
scale plasma metabolomics study involving 35 samples from 20 individuals with
symptomatic and subclinical infections in Mali, taurine levels were not reported [36]. Further
studies are needed to explore whether elevated plasma taurine levels are a consistent feature
of asymptomatic *P. falciparum* infections during the dry season across sub-Saharan Africa. It
cannot be excluded that, in addition to taurine, other metabolites associated with plasma
alterations in asymptomatic malaria contribute to subclinical malaria by influencing the iRBC
proliferation or by increasing their susceptibility to immune-mediated clearance.

Other host receptors such as ICAM-1 have been linked to severe malaria and aggregation of
iRBCs and uninfected RBCs (known as rosettes) [reviewed in [2]]. Both adhesion phenotypes
are significantly rarer than CD36 adhesion and it remains to be shown if these phenotypes
play an important role in symptomatic infections during seasonal malaria. A sulfated
polysaccharide derived from heparin was shown to interfere with rosette formation [68]. While
this negatively charged carbohydrate compound shows promise, further studies are needed to
fully understand its role in subclinical infections.

Binding of *P. falciparum* iRBCs to human receptors is central to the development of
symptomatic and severe malaria. Yet, no adjunctive therapies or broad-spectrum vaccines

currently target CD36-binding parasites, and efforts to block EPCR-binding variants
implicated in severe and cerebral malaria remain ongoing [69]. Clinically, taurine is already
used as a dietary supplement with documented benefits for conditions like hypertension and
hypercholesterolemia [reviewed in [70]]. Our identification of taurine, a natural host molecule,
as an effective inhibitor of iRBC binding to both CD36 and EPCR reveals a novel public
health strategy to reduce or eliminate dry-season reservoirs and prevent symptomatic malaria
at the start of the transmission season.

MATERIALS AND METHODS
Study individuals and ethical approval
Study design and sample collection was alredy reported in [37,38,44]. The first part of the study
was a cross-sectional study implemented in October and November 2016 to estimate the
prevalence of *P. falciparum* infections in four neighboring villages (Madina Samako: K,
Njayel: J, Sendebu: P, and Karandaba: N) within a 5km radius in Upper River Region (URR),
eastern Gambia. *P. falciparum* asymptomatic infections were detected by varATS qPCR and
individuals were invited to provide a venous blood sample before receiving anti-malarial
treatment. Symptomatic cases presented themselves at the local health facility with fever.
Malaria cases were defined as anxially temperature >37.5C, positive RDT and later
confirmation of *P. falciparum* infection by varATS qPCR. For the second part of the study, in
December 2016, asymptomatic *P. falciparum* carriers were recruited for a longitudinal cohort
with monthly samplings until May 2017, or until the infection naturally cleared [Ref
Collins2022].

The high transmission season months include October and November and the low
transmission season months include late December, January, February, March, April, and
May. The months pertaining to the two transmission periods was directly determined by the
reports of symptomatic malaria infections for that specific year for all villages used in the
study [37]. For additional information about each sample, see (SM1).

The study protocol was reviewed and approved by the Gambia Government/MRC Joint
Ethics Committee (SCC 1476, SCC 1318, L2015.50) and by the London School of Hygiene
& Tropical Medicine ethics committee (Ref 10982). The field studies were also approved by
local administrative representatives, the village chiefs. Written informed consent was

obtained from participants over 18 years of age and from parents/guardians for participants
under 18 years. Written assent was obtained from all individuals aged 12–17 years.

Human plasma sample collection

From *P. falciparum* positive individuals (by varATS qPCR), 5 to 8mL venous blood was
collected in a EDTA vacutainer. Collected venous blood samples were immediately separated
in plasma, PBMCs and red blood cells, by centrifugation on a layer of Lymphoprep™
according to manufacturer's instructions. Plasma was directly stored at -80 degrees Celsius
and never thawed until use in this study.

Metabolomics of human plasma sample preparation and widely targeted detection by
GC-MS
50 µl of collected plasma were mixed with 500 µl of ice-cold extraction mixture
(methanol/water, 9/1, -20°C, with labelled internal standard). To facilitate endogenous
metabolites extraction, samples were then completely homogenized (vortexed 5 minutes at
2500 rpm) and then centrifuged (15 min at 10000 g, 4°C). 150 µl of supernatants were
collected to be analyzed by gas chromatography coupled with mass spectrometer (GC/MS) [71].

Widely targeted analysis by GC-MS/MS was performed on a 7890A gas chromatography
(Agilent Technologies) coupled with Triple Quadrupole 7000C (Agilent Technologies) and
was previously described in [72]. All targeted treated data were merged and cleaned with a
dedicated R (version 4.0) package (@Github/Kroemerlab/GRMeta).

Plasma taurine concentration assay
Plasma taurine levels were determined using a commercial taurine assay kit (Cell Biolabs
MET-5071). Complete information on tested individual samples can be found in (SM1). The
protocol was followed as described in the kit and read on a Synergy 2 microplate reader for
spectrophotometric reading.

Parasite culture and synchronization
Asexual blood stage *P. falciparum* parasites were cultured as previously described in [24].
Parasites were cultured in human RBCs (obtained from the Etablissement Français du Sang
with approval number HS 2019-24803) in RPMI-1640 culture medium (Thermo Fisher

11875) supplemented with 10% v/v Albumax I (Thermo Fisher 11020), hypoxanthine (0.1

mM final concentration, C.C.Pro Z-41-M) and 10 mg gentamicin (Sigma G1397) at 4%

hematocrit and under 5% O_2, 3% CO_2 at 37 °C. Static parasite development was monitored

by Giemsa staining. Parasites were synchronized by sorbitol (5%, Sigma S6021) lysis at ring

stage, plasmagel (Plasmion, Fresenius Kabi) enrichment of late stages 24 hours later, and an

additional sorbitol lysis 3 hours after plasmagel enrichment. The 0-hour time point was

considered to be 1.5 hours after plasmagel enrichment.

Parasite growth assay

Parasite growth was measured as described previously [73]. Wild type 3D7 parasites were

tightly synchronized and diluted to 0.2% parasitemia (5% hematocrit) at the ring stage using

the blood of three different donors separately (n=3). Each culture was split into control and

parasites supplemented with [0.16mM] taurine. The growth curve was performed with three

technical replicates per condition per blood. Parasitemia was measured every 24 hours by

counting ten randomly selected different fields on Giemsa-stained slides each day for a total

of five days.

Static cytoadhesion binding assays

3D7-CD36 binding assay: Mature stage iRBCs for 3D7 control, supplementation of taurine

for increasing concentrations [0.02, 0.04, 0.08, and 0.16mM] from a [200mM] stock, and

supplementation of methionine [0.04mM] after synchronization were used for cytoadhesion

assays as described earlier [74]. Additionally, supplementation of [0.16mM] taurine and

[0.27mM] threonic acid for one hour prior to the assays were used. Receptors diluted in

Dulbecco's phosphate-buffered saline (DPBS, Thermo Fisher 14190) (10 µg/mL target CD36

or negative control 1% BSA) were spotted on labeled petri dishes overnight at 4°C in

duplicate. Dishes were blocked for 30 minutes at 37°C with 1% BSA/DPBS while

parasitemia was determined after trophozoite and young schizont iRBCs were isolated by

plasmagel (Plasmion, Fresenius Kabi) enrichment. Adjusted amounts of iRBCs were

resuspended in DPBS for a concentration of 2.2×10^8 iRBCs/mL. iRBCs/DPBS were added

to each petri dish and incubated for one hour at 37°C. Unbound cells were removed, and the

dishes were washed five times with DPBS by carefully tilting from side to side. Adherent

iRBCs were counted in duplicate spots using 40x lens with a Nikon ECLIPSE TE200 in

seven randomly selected fields (each with 0.2 mm^2) using ImageJ2 (version 2.14.0/1.54i). A

total of three biological replicates were performed in duplicate and the results were expressed

as the percentage of bound iRBCs compared to the control per 0.2 mm^2 of target receptor

monolayer.

<u>FCR3-*var19*-EPCR and FCR3-CD36 binding assay:</u> For EPCR binding, the protocol listed

above was followed with the following changes: Receptors were diluted in in DPBS (50

µg/mL EPCR, 50 µg/mL CD36 or negative control 1% BSA) were spotted on labeled petri

dishes overnight at 4°C in duplicate. Late stage iRBCs from FCR3 *var19*-expressing parasites

[57] were purified by VARIOMACS (Miltenyi Biotec France) and counted. The iRBC pellet

was split into control and addition of taurine [0.64mM] for one hour in RPMI-1640 culture

medium (Thermo Fisher 11875) supplemented with 10% v/v Albumax I (Thermo Fisher

11020), hypoxanthine (0.1 mM final concentration, C.C.Pro Z-41-M) and 10 mg gentamicin

(Sigma G1397) at 4% hematocrit and under 5% O_2, 3% CO_2 at 37□°C. The spots were

washed twice with DPBS and blocked for 30 minutes at room temperature with 1%

BSA/DPBS, 3.75 x 10^6 parasites/mL diluted in 10uL DPBS were added to each spot and

incubated for one hour at room temperature. Plates were gently washed with 25mL DPBS on

an orbital shaker for 2 minutes and repeated four times before visualization and imaging

using EVOS 5000. Adherent iRBCs were counted in duplicate spots using 20x lens with a

Nikon ECLIPSE TE200 in seven randomly selected fields (each with 0.3 mm^2).

<u>NF54-*var2CSA*-CSA binding assay:</u> For CSA binding, the protocol listed above (for 3D7-

CD36 binding) was followed for control and supplementation of [0.64mM] Taurine using

CSA-panned NF54 parasites [58,75].

<u>Evaluation of taurine's ability to reverse iRBC binding to CD36 and EPCR</u>: A two-step

protocol was used to evaluate the reversal of binding of 3D7 and FCR3-*var19*-expressing

parasites to CD36 or EPCR. In Step 1, binding was conducted as previously described for

3D7 to CD36 and FCR3-*var19* to EPCR. In Step 2, DPBS or [0.64mM] taurine + DPBS was

added for 45 minutes, followed by DPBS washes. Adherent iRBCs were counted in duplicate

spots using a 10x objective lens to count and assess the percent reduction in bound iRBCs

after taurine incubation compared to the control.

<u>Taurine-CD36 incubation prior to iRBC adhesion assay</u>: For CD36-Taurine binding, the

protocol was followed the same as listed above (for 3D7-CD36 binding) for mature stage

iRBCs from 3D7 parasites with the following changes: Control and supplementation of
[0.16mM] taurine was added during blocking stage with 1% BSA/DPBS.

Human serum- and human serum+taurine-CD36 binding assay: For human serum-CD36
binding and human serum + taurine-CD36 binding, the protocol was followed the same as
listed above (for 3D7-CD36 binding) for mature stage iRBCs from 3D7 parasites with the
following changes: 10% human serum without and with supplementation of [0.64mM]
taurine from three different donors (n=3) was added for one hour prior to the assay.

C32 melanoma cell binding assay
C32 melanoma binding assays were performed as previously described [76]. To prepare iRBCs
for the cellular inhibition assay, mature-stage iRBCs (3D7 or FCR3) were prepared as for the
receptor binding assays above. Plasma gel-enriched FCR3 iRBCs were overlaid onto
confluent C32 cell monolayers in 6-well plastic dishes for 45 minutes in binding medium.
After a 1-hour incubation, unbound iRBCs were removed by gentle washes with RPMI
medium adjusted to pH 6.8 until only adhering parasites were observed. For each taurine
concentration, ten independent 1□mm² microscopic fields were counted across three
biological replicates. iRBC adhesion was quantified by counting bound iRBC per mm^2 of
confluent C32 cells in 10 random fields. Representative micrographs and raw data are
provided in the supplementary materials (Fig. 4D, S4D, SM6).

Flow cytometry assessment of *var19* surface expression
Control and supplementation of [0.64mM] taurine in FCR3 *var19*-expressing parasites iRBCs
at mid/late trophozoites stages were resuspended in DPBS 1% BSA after VARIOMACS
purification and counted. For each assay, 3×10^5 iRBCs were washed in DPBS and incubated
with 50µl of purified rabbit anti-*var19* antibodies [57] diluted 1:100 in PBS 1% BSA for 1-hour
at room temperature. iRBCs were washed twice with DPBS and resuspended in 100µl of PE
conjugated goat anti-rabbit antibody diluted 1: 100 in DPBS for 30 min at RT. iRBCs were
washed twice in DPBS 1% BSA and resuspended in paraformaldehyde 4% in DPBS and kept
at 4°C overnight in darkness. Cells were washed twice with DPBS and analyzed by flow
cytometry using a BD LSR Fortessa flow cytometer. The results were analyzed using the
FlowJo 10.0 software. Parasite nuclei were stained by TO-PRO-3 (1:10000 dilution). The
results are expressed as the percentage of *var19* surface expression. A total of three biological
replicates (n=3) were performed.

Statistical analysis

For metabolomics MS analysis, GRmeta (https://github.com/kroemerlab/GRMeta) was used

for the data shape QC correction (based on global pool for normalization between 3 total

batches).

Fold change (FC) was determined when comparing each condition of each month to all other

conditions in each month using the following code:

#Fold Change averages

mm = mergedata$Data$AreaCorrLog2Cen2

mm = cbind(as.data.frame(mergedata$Grp2),mm)

mm = apply(mm[,-1], 2, function(x) tapply(x, mm$`mergedata$Grp2`, mean)) %>% t()

FC = combn(1:ncol(mm),2,function(x) mm[,x[2]] - mm[,x[1]])

p values were determined when comparing each condition of each month to all other

conditions in each month using Kruskall wallis with BH adjustment with the following code:

KruskalWallis$data[[mergedata$Annot$MetName[i]]]=as.data.frame(dunn.test(mergedata$

Data$AreaCorrLog2Cen2[,i],mergedata$Grp2, method="BH", kw=TRUE,

label=TRUE,wrap=FALSE, table=TRUE, list=FALSE, rmc=FALSE, alpha=0.05,

altp=FALSE)$P.adjusted)

All other statistical analyses were performed using GraphPad Prism version 9.1.0 (216) for

Mac. To test for a normal distribution of the data, the Shapiro-Wilk normality test was used.

To test for significance between two groups, a two-sided independent-samples t test was

used.

Metabolite enrichment and pathway analyses was performed using the Over Representation

and Quantitative Enrichment tools on the MetaboAnalyst 6.0 server

(https://www.metaboanalyst.ca/), integrated with the KEGG database

(https://www.genome.jp/kegg/). Input data included a list of compound names (KEGG and

HMDB IDs) for metabolites passing GC-MS quality control, categorized by their specific

class. Pathway-based analysis was performed with at least two entries in the metabolite set

library, which includes 80 metabolite sets based on KEGG human metabolic pathways (Dec

2023). Results were visualized as dot plots displaying key information on enrichment ratios,

p-values, and log2 values for the respective metabolite sets.

Supplementary Materials:

Fig. S1 to S4

SM1 Information on samples from volunteers (excel file)

SM2 Metabolite alternative names (excel file)

SM3 Metabolomics MS raw data (excel file)

SM4 Pathway Enrichment Analysis (excel file)

SM5 Fold Change and p-values with significantly increased metabolites in each condition for

each month (excel file)

SM6 Cytoadhesion counting raw data (excel file)

Fig. 1: Blood Plasma Metabolomic Profiling of *P. falciparum* Infections in The Gambia

(**A**) Region in The Gambia where plasma samples were taken from volunteers in neighboring

villages within a 5km radius (left). Overview of the change from high transmission to low

transmission months in The Gambia. Sample numbers are indicated by each month for

volunteers that had symptomatic or asymptomatic infections as well individuals with no

detected presence of *P. falciparum* (right). (**B**) Schematic showing the flow of the

experimental analysis for gas chromatography coupled with triple-quadrupole mass

spectrometry (GC–QqQ/MS) for widely targeted metabolites. (A, B) Figures created with

BioRender.com (**C**) Metabolite class enrichment for all 78 metabolites that passed quality

control. (**D**) Pathway enrichment analysis of the 78 metabolites used in analysis are shown

with p-value and enrichment ratio. (**E**) PCA plot for all 78 metabolites and all (n=199)

plasma samples for the three clinical conditions (symptomatic infections, asymptomatic

infections, and uninfected negative).

Fig. S1: Additional information on samples used

(**A**) Age distribution for all individuals from each of the conditions in each month. Grey

boxes represent the median age. (**B**) Sample distribution for all conditions in each month. The

median age in years is shown as well as the distribution by sex for males and females. **(C)**
Parasitemia determined by qPCR for each of the conditions with available data, analyzed
previously in [44]. Symptomatic October (n=7) and November (n=12), Asymptomatic October
(n=24), November (n=22), December (n=44), January (n=42), February (n=29), March (n=3),
April (n=4), and May (n=21). Statistical significance was determined by unpaired t-test
comparing symptomatic infections to asymptomatic infections in the same months (**
October p = 0.0085, November p = 0.0038). **(D)** Heatmap for all 78 metabolites and (n=199)
plasma samples with hierarchical clustering illustrating the changes in metabolite abundances
according to the median of each metabolite. Rows are metabolites, columns are samples.
Heatmap data are log2 normalized and centered around the average abundance computed
from all the samples for each metabolite. Red/green colors are ion signal higher/lower than
average. Samples and metabolites are clustered following the ward.D2 algorithm, with
Euclidean distance.

Fig. 2: Metabolites significantly enriched in symptomatic and asymptomatic samples
(**A**) Table showing metabolites that were uniquely enriched based on symptomatic infections
and seasonality (increased in asymptomatic infections during the high and low transmission
periods). **(B)** Heatmap from MS analysis showing metabolites that varied based on severity
and seasonality. All (n=199) samples, months, and conditions are included. Columns are
metabolites, rows are samples. Heatmap data are log2 normalized (within a range of [-2]-[2])
and centered around the average abundance computed from all the samples for each
metabolite. Red/green colors are ion signal higher/lower than average. Samples and
metabolites are clustered following the ward.D2 algorithm, with Euclidean distance. **(C)**
Scatter plots showing relative abundance levels, determine by MS metabolomic analysis, of
threonic acid and taurine in symptomatic infections, asymptomatic infections, and uninfected
negative individuals. High transmission (HT) and low transmission (LT) months are labeled.
For simplicity, November is shown for samples from individuals with symptomatic
infections, November, January and May are shown for samples from individuals with
asymptomatic infections, and January is shown for samples from uninfected negative
individuals. Mean values are represented in shaded boxes for all conditions and
corresponding months.

Fig. S2: Taurine levels rise in multiple individuals during the low transmission period.

(**A**) Spaghetti plot showing taurine levels in individuals (n=16) with persistent infections who

had samples taken for consecutive months during the low transmission period (December-

May). Values plotted are relative abundance determined from the MS analysis. (**B**) Scatter

plots showing relative abundance levels, determine by MS metabolomic analysis, of thirteen

metabolites that were significantly altered between uninfected negative individuals and

asymptomatic infections during the low transmission season in January. Mean values are

represented in shaded boxes for all metabolites in both conditions.

Fig. 3: Taurine markedly reduces iRBC cytoadhesion

(**A**) Plasma taurine concentrations for the top three highest and lowest values from

metabolomics MS analysis determined using a taurine assay kit. Statistical significance was

determined by paired t-test (* $p = 0.0115$). (**B**) Cytoadhesion binding assays showing 3D7

iRBCs binding to CD36. Parasites have been incubated for 30 hours in [0.04mM] methionine

or increasing concentrations of taurine. Additionally, parasites were incubated for one hour in

[0.16mM] taurine. Data is expressed as the percentage of iRBCs binding ±SD per mm^2 for

each condition compared to control, with the median represented at the center line. Results

shown are three biological replicates (n=3) from duplicate technical replicates. Statistical

significance was determined by paired t-test (* $p = 0.0305$ and 0.0152, **** $p < 0.0001$). (**C**)

Representative cytoadhesion assay images are shown for control and supplementation of

taurine physiological concentration [0.16mM]. (**D**) EPCR binding assay of Itvar19 FCR3

iRBCs in the presence of [0.16mM] taurine and [0.64mM] taurine. Data is expressed as the

percentage of iRBCs binding ±SD per mm^2 for each condition compared to control, with the

median represented at the center line. Results are from a total of three biological replicates

(n=3) done in duplicate. Statistical significance was determined by paired t-test (* $p = 0.0264$,

** $p = 0.0045$).

Fig. S3: Taurine reduces cytoadhesion without affecting parasite growth

(**A**) Five-day growth curve of 3D7 wild-type *P. falciparum* cultured under three conditions:

control, 0.16 mM taurine (physiological concentration), and 0.27 mM threonic acid

(physiological concentration). Error bars represent the standard deviation of three biological

replicates performed in triplicate with different donor blood (n = 3). (**B**) Replicate counts

(n=3) for cytoadhesion assays from (Fig. 3B). Data is expressed as the number of bound 3D7

iRBCs per mm^2 of coated receptor for each condition. (**C**) Cytoadhesion assays showing *P.*

falciparum 3D7 iRBCs binding to CD36 after 1-hour incubation with 0.27 mM threonic

acid. Data are presented as the percentage of iRBCs bound per mm² (±SD), relative to

control; median values are indicated by the center line. Results represent three biological

replicates (n□=□3), each performed in duplicate. Statistical significance was determined by

paired t-test (* p = 0.0305 and 0.0152, **** p < 0.0001). NS indicates not significant. **(D)**

Cytoadhesion assays examining parasite FCR3 Itvar19 iRBCs binding to CD36, with or

without pre-treatment with [0.16mM] taurine for one hour before the binding assay [parasites

were the same strain as those used from (Fig. 3D)]. Data is expressed as the percentage of

bound iRBCs ±SD per mm^2 for each condition compared to control, with the median

represented at the center line. Results are from a total of three biological replicates (n=3)

done in duplicate. Statistical significance was determined by paired t-test (*** p = 0.0003).

(**E)** Cytoadhesion assays for NF54 VAR2CSA iRBCS binding to CSA for control and

[0.64mM] taurine. Data is expressed as the percentage of bound iRBCs ±SD per mm^2 for

each condition compared to control, with the median represented at the center line. Results

are from a total of three biological replicates (n=3) done in duplicate. Statistical significance

was determined by paired t-test (* p = 0.0288). **(F)** Cytoadhesion assays assessing 3D7

iRBCs binding to CD36 under control conditions and with the addition of 10% human serum

with or without [0.64mM] taurine supplementation. Data is expressed as the percentage of

bound iRBCs ±SD per mm^2 compared to control, with the median represented at the center

line. Results are from a total of three biological replicates (n=3) done in duplicate. Statistical

significance was determined by paired t-test (* p= 0.035, ** p = 0.0066).

Fig. 4: Taurine-Mediated Reversal of iRBC Receptor Binding and Inhibition of

Cytoadhesion to C32 Cells

(**A**) Reversal of cytoadhesion of 3D7 iRBCs bound to CD36, incubated with or without

[0.64mM] taurine. Data is expressed as the fold change of bound iRBCs ±SD per mm^2

compared to control, with the median represented at the center line. Results are from a total

of three biological replicates (n=3) done in duplicate. Statistical significance was determined

by paired t-test (*** p = 0.0009). (**B**) Reversible cytoadhesion assays for FCR3 Itvar19

parasites bound to EPCR incubated with or without [0.64mM] taurine. Data is expressed as

the fold change of bound iRBCs ±SD per mm^2 compared to control, with the median

represented at the center line. Results are from a total of three biological replicates (n=3)

done in duplicate. Statistical significance was determined by paired t-test (** p = 0.0179).

(**C**) Cytoadhesion assays measuring *P. falciparum* FCR3 parasite strain binding to human

C32 melanoma cells expressing CD36 and ICAM-1 under control conditions and with

increasing taurine concentrations (0.16 mM, 0.64 mM, 0.93 mM). Data is expressed as
the fold change of bound iRBCs per mm² of confluent C32 cells (±SD), with medians
indicated by center lines. Results represent three biological replicates (n = 3), each in
duplicate. Statistical significance was determined by paired t-test (* p = 0.0231, **
p = 0.0076); NS = not significant. (**D**) Representative field of cytoadhesion assay from
(Fig. 4C).

Fig. S4: Taurine does not alter PfEMP1 surface expression

(**A**) FACS analysis for surface expression of PfEMP1 from parasites, incubated with or
without [0.64mM] taurine, from a total of two biological replicates (n=2) done in duplicate
using different antibodies against *var19* PfEMP1. Statistical significance was determined by
paired t-test (NS indicates not significant). (**B**) Cytoadhesion assays assessing 3D7 parasite
binding to CD36 under control conditions and with CD36 pre-incubated with [0.16mM]
taurine. Data is expressed as the percentage of bound iRBCs ±SD per mm^2 compared to
control, with the median represented at the center line. Results are from a total of three
biological replicates (n=3) done in duplicate. Statistical significance was determined by
paired t-test (NS indicates not significant). (**C**) Cytoadhesion assays measuring 3D7
iRBCs binding to human C32 melanoma cells expressing CD36 and ICAM-1 under control
conditions and [0.93mM] taurine. Replicate counts (n=2) for cytoadhesion assays. Data is
expressed as the number of bound iRBCs per mm^2 of confluent C32 cells for each condition
(±SD), with medians indicated by center lines. (**D**) Representative field of cytoadhesion assay
from (Fig. S4C). (**E**) Flow-cytometry gating strategy. Gating hierarchy for a representative
sample. From left to right: (a) In the FSC-A versus SSC-A plot, the cloud of RBC events is
identified. (b) Among the RBCs, in the FSC-A versus FSC-H plot, singlets (single cell
signals) are distinguished. (c) Among the singlets, the APC-A histogram is used to identify
iRBCs. (d) Finally, in the PE-A channel, cells expressing var19 on their surface are identified
based on PE-A labelling. (**F**) Flow cytometry analysis of FCR3 Var19 infected red blood
cells. Percentage of Var19 expression on the cell surface without or with [0.64mM] taurine
supplementation (negative controls included). Histogram legends for the expression data and
the percentage of Var19 expression on the surface with the geometric mean are shown.
Results are from three biological replicates (n=3).

References

1. W.H.O. World Malaria Report 2023. (2023).

2. Wahlgren, M., Goel, S. & Akhouri, R. R. Variant surface antigens of Plasmodium

falciparum and their roles in severe malaria. *Nat Rev Microbiol* (2017).

3. Miller, L. H., Baruch, D. I., Marsh, K. & Doumbo, O. K. The pathogenic basis of

malaria. *Nature* **415**, 673–679 (2002).

4. Scherf, A., Lopez-Rubio, J. J. & Riviere, L. Antigenic variation in Plasmodium

falciparum. *Annu Rev Microbiol* **62**, 445–470 (2008).

5. Deitsch, K. W. & Dzikowski, R. Variant Gene Expression and Antigenic Variation by

Malaria Parasites. *Annu Rev Microbiol* **71**, 625–641 (2017).

6. Baruch, D. I. *et al.* Cloning the P. falciparum gene encoding PfEMP1, a malarial variant

antigen and adherence receptor on the surface of parasitized human erythrocytes. *Cell* **82**,

77–87 (1995).

7. Smith, J. D. *et al.* Switches in expression of plasmodium falciparum var genes correlate

with changes in antigenic and cytoadherent phenotypes of infected erythrocytes. *Cell* **82**,

101–110 (1995).

8. Su, X. Z. *et al.* The large diverse gene family var encodes proteins involved in

cytoadherence and antigenic variation of plasmodium falciparum-infected erythrocytes.

Cell **82**, 89–100 (1995).

9. Gardner, M. J. *et al.* Genome sequence of the human malaria parasite Plasmodium

falciparum. *Nature* **419**, 498–511 (2002).

10. Smith, J. D., Rowe, J. A., Higgins, M. K. & Lavstsen, T. Malaria's deadly grip:

cytoadhesion of Plasmodium falciparum-infected erythrocytes: Plasmodium falciparum

cytoadhesion and var genes. *Cell Microbiol* **15**, 1976–1983 (2013).

11. Robinson, B. A., Welch, T. L. & Smith, J. D. Widespread functional specialization of

Plasmodium falciparum erythrocyte membrane protein 1 family members to bind CD36

analysed across a parasite genome. *Mol Microbiol* **47**, 1265–1278 (2003).

12. Hsieh, F.-L. *et al.* The structural basis for CD36 binding by the malaria parasite. *Nat*

Commun **7**, 12837 (2016).

13. Turner, L. *et al.* Severe malaria is associated with parasite binding to endothelial protein

C receptor. *Nature* **498**, 502–505 (2013).

14. Lau, C. K. Y. *et al.* Structural Conservation Despite Huge Sequence Diversity Allows

EPCR Binding by the PfEMP1 Family Implicated in Severe Childhood Malaria. *Cell*

Host & Microbe **17**, 118–129 (2015).

15. Urban, B. C. *et al.* Plasmodium falciparum-infected erythrocytes modulate the maturation

of dendritic cells. *Nature* **400**, 73–77 (1999).

16. Serghides, L., Smith, T. G., Patel, S. N. & Kain, K. C. CD36 and malaria: friends or foes?

Trends Parasitol **19**, 461–469 (2003).

17. Avril, M. *et al.* A restricted subset of var genes mediates adherence of Plasmodium

falciparum-infected erythrocytes to brain endothelial cells. *Proc Natl Acad Sci U S A* **109**,

E1782-1790 (2012).

18. Claessens, A. *et al.* A subset of group A-like var genes encodes the malaria parasite

ligands for binding to human brain endothelial cells. *Proc Natl Acad Sci U S A* **109**,

E1772-1781 (2012).

19. Lavstsen, T. *et al.* Plasmodium falciparum erythrocyte membrane protein 1 domain

cassettes 8 and 13 are associated with severe malaria in children. *Proc Natl Acad Sci U S*

A **109**, E1791-1800 (2012).

20. Tuikue Ndam, N. *et al.* Parasites Causing Cerebral Falciparum Malaria Bind Multiple

Endothelial Receptors and Express EPCR and ICAM-1-Binding PfEMP1. *The Journal of*

Infectious Diseases **215**, 1918–1925 (2017).

21. Lennartz, F. *et al.* Structure-Guided Identification of a Family of Dual Receptor-Binding

PfEMP1 that Is Associated with Cerebral Malaria. *Cell Host Microbe* **21**, 403–414

(2017).

22. Bozdech, Z. *et al.* The Transcriptome of the Intraerythrocytic Developmental Cycle of

Plasmodium falciparum. *PLoS Biol* **1**, e5 (2003).

23. Le Roch, K. G. *et al.* Discovery of gene function by expression profiling of the malaria

parasite life cycle. *Science* **301**, 1503–1508 (2003).

24. Lopez-Rubio, J. J., Mancio-Silva, L. & Scherf, A. Genome-wide analysis of

heterochromatin associates clonally variant gene regulation with perinuclear repressive

centers in malaria parasites. *Cell Host Microbe* **5**, 179–190 (2009).

25. Rovira-Graells, N. *et al.* Transcriptional variation in the malaria parasite Plasmodium

falciparum. *Genome Res* **22**, 925–938 (2012).

26. Babbitt, S. E. *et al.* Plasmodium falciparum responds to amino acid starvation by entering

into a hibernatory state. *Proc Natl Acad Sci U S A* **109**, E3278-3287 (2012).

27. Schneider, V. M. *et al.* The human malaria parasite Plasmodium falciparum can sense

environmental changes and respond by antigenic switching. *Proc Natl Acad Sci U S A*

120, e2302152120 (2023).

28. Marreiros, I. M. *et al.* A non-canonical sensing pathway mediates Plasmodium adaptation

to amino acid deficiency. *Commun Biol* **6**, 205 (2023).

29. Brancucci, N. M. B. *et al.* Lysophosphatidylcholine Regulates Sexual Stage

Differentiation in the Human Malaria Parasite Plasmodium falciparum. *Cell* **171**, 1532-

1544.e15 (2017).

30. Diffendall, G. *et al.* RNA polymerase III is involved in regulating Plasmodium

falciparum virulence. *Elife* **13**, RP95879 (2024).

31. Mancio-Silva, L. *et al.* Nutrient sensing modulates malaria parasite virulence. *Nature*

547, 213–216 (2017).

32. Babiker, H. A., Abdel-Muhsin, A. M., Ranford-Cartwright, L. C., Satti, G. & Walliker,

D. Characteristics of Plasmodium falciparum parasites that survive the lengthy dry season

in eastern Sudan where malaria transmission is markedly seasonal. *Am J Trop Med Hyg*

59, 582–590 (1998).

33. Andrade, C. M., Fleckenstein, H. & Thomson-Luque, R. Increased circulation time of

Plasmodium falciparum underlies persistent asymptomatic infection in the dry season.

Nat Med **26**, 1929–1940 (2020).

34. Portugal, S. *et al.* Treatment of Chronic Asymptomatic Plasmodium falciparum Infection

Does Not Increase the Risk of Clinical Malaria Upon Reinfection. *Clin Infect Dis* **64**,

645–653 (2017).

35. Thomson-Luque, R. *et al.* Plasmodium falciparum transcription in different clinical

presentations of malaria associates with circulation time of infected erythrocytes. *Nat*

Commun **12**, 4711 (2021).

36. Andrade, C. M. *et al.* Infection length and host environment influence on Plasmodium

falciparum dry season reservoir. *EMBO Mol Med* **16**, 2349–2375 (2024).

37. Fogang, B. *et al.* Asymptomatic Plasmodium falciparum carriage at the end of the dry

season is associated with subsequent infection and clinical malaria in Eastern Gambia.

Malar J **23**, 22 (2024).

38. Guery, M.-A. *et al.* Household clustering and seasonal genetic variation of Plasmodium

falciparum at the community-level in The Gambia. *eLife* **13**, RP103047 (2025).

39. Surowiec, I. *et al.* Metabolic Signature Profiling as a Diagnostic and Prognostic Tool in

Pediatric Plasmodium falciparum Malaria. *Open Forum Infectious Diseases* **2**, ofv062

(2015).

40. Beri, D. *et al.* Insights into physiological roles of unique metabolites released from

Plasmodium-infected RBCs and their potential as clinical biomarkers for malaria. *Sci Rep*

9, 2875 (2019).

41. Leopold, S. J. *et al.* Identifying the Components of Acidosis in Patients With Severe

Plasmodium falciparum Malaria Using Metabolomics. *The Journal of Infectious Diseases*

219, 1766–1776 (2019).

42. Lakshmanan, V. *et al.* Metabolomic analysis of patient plasma yields evidence of plant-

like α-linolenic acid metabolism in Plasmodium falciparum. *J Infect Dis* **206**, 238–248

(2012).

43. Abdrabou, W. *et al.* Metabolome modulation of the host adaptive immunity in human

malaria. *Nat Metab* **3**, 1001–1016 (2021).

44. Collins, K. A. *et al.* A Cohort Study on the Duration of Plasmodium falciparum

Infections During the Dry Season in The Gambia. *J Infect Dis* **226**, 128–137 (2022).

45. Huxtable, R. J. Physiological actions of taurine. *Physiol Rev* **72**, 101–163 (1992).

46. Debierre-Grockiego, F. *et al.* Fatty acids from Plasmodium falciparum down-regulate the

toxic activity of malaria glycosylphosphatidylinositols. *Infect Immun* **74**, 5487–5496

(2006).

47. Sasi, P. *et al.* Metabolic acidosis and other determinants of hemoglobin-oxygen

dissociation in severe childhood Plasmodium falciparum malaria. *Am J Trop Med Hyg*

77, 256–260 (2007).

48. Baliou, S. *et al.* Protective role of taurine against oxidative stress (Review). *Mol Med Rep*

24, 605 (2021).

49. Li, X. *et al.* Taurine potentiates artemisinin efficacy against malaria by modulating the

immune response in Plasmodium berghei-infected mice. *Parasit Vectors* **17**, 493 (2024).

50. Shi, X. *et al.* Vitamin C Inhibits Blood-Stage Plasmodium Parasites via Oxidative Stress.

Front Cell Dev Biol **9**, 639944 (2021).

51. Trautwein, E. & Hayes, K. Taurine concentrations in plasma and whole blood in humans:

estimation of error from intra- and interindividual variation and sampling technique. *The*

American Journal of Clinical Nutrition **52**, 758–764 (1990).

52. Wang, H., Hu, P. & Jiang, J. Pharmacokinetics and safety of calcium L-threonate in

healthy volunteers after single and multiple oral administrations. *Acta Pharmacol Sin* **32**,

1555–1560 (2011).

53. Rajapandi, T., Ackie, K. & Rajeev Hegde, K. Anti-plasmodial Activity of a Non-protein

Amino Acid Taurine. *BS* **5**, 34 (2019).

54. Ghandforoush-Sattari, M., Mashayekhi, S., Krishna, C. V., Thompson, J. P. & Routledge,

P. A. Pharmacokinetics of Oral Taurine in Healthy Volunteers. *Journal of Amino Acids*

2010, 1–5 (2010).

55. Garlick, P. J. Toxicity of Methionine in Humans. *The Journal of Nutrition* **136**, 1722S-

1725S (2006).

56. Srivastava, A. *et al.* Var2CSA Minimal CSA Binding Region Is Located within the N-

Terminal Region. *PLoS ONE* **6**, e20270 (2011).

57. Nunes-Silva, S. *et al.* Beninese children with cerebral malaria do not develop humoral

immunity against the IT4-VAR19-DC8 PfEMP1 variant linked to EPCR and brain

endothelial binding. *Malar J* **14**, 493 (2015).

58. Viebig, N. K. *et al.* A single member of the *Plasmodium falciparum var* multigene family

determines cytoadhesion to the placental receptor chondroitin sulphate A. *EMBO Reports*

6, 775–781 (2005).

59. Nacer, A. *et al.* Clag9 is not essential for PfEMP1 surface expression in non-cytoadherent

Plasmodium falciparum parasites with a chromosome 9 deletion. *PLoS One* **6**, e29039

(2011).

60. Schmidt, J. A. *et al.* Plasmodium Falciparum Malaria. *J. Clin. Invest.* **70**, 379–386

(1982).

61. Ho, M. *et al.* Clinical correlates of in vitro Plasmodium falciparum cytoadherence. *Infect*

Immun **59**, 873–878 (1991).

62. Ockenhouse, C. F. *et al.* Molecular basis of sequestration in severe and uncomplicated

Plasmodium falciparum malaria: differential adhesion of infected erythrocytes to CD36

and ICAM-1. *J Infect Dis* **164**, 163–169 (1991).

63. Beck, J. R. & Ho, C.-M. Transport mechanisms at the malaria parasite-host cell interface.

PLoS Pathog **17**, e1009394 (2021).

64. Jabre, I. *et al.* Histone lactylation: a new epigenetic mark in the malaria parasite

Plasmodium. Preprint at https://doi.org/10.1101/2024.08.19.608576 (2024).

65. Merrick, C. J. *et al.* Epigenetic dysregulation of virulence gene expression in severe

Plasmodium falciparum malaria. *J Infect Dis* **205**, 1593–1600 (2012).

66. Stacy, A. *et al.* Infection trains the host for microbiota-enhanced resistance to pathogens.

Cell **184**, 615-627.e17 (2021).

67. Li, X. *et al.* Taurine potentiates artemisinin efficacy against malaria by modulating the

immune response in Plasmodium berghei-infected mice. *Parasit Vectors* **17**, 493 (2024).

68. Vogt, A. M. *et al.* Release of Sequestered Malaria Parasites upon Injection of a

Glycosaminoglycan. *PLoS Pathog* **2**, e100 (2006).

69. Reyes, R. A. *et al.* Broadly inhibitory antibodies to severe malaria virulence proteins.

Nature **636**, 182–189 (2024).

70. Schaffer, S. & Kim, H. W. Effects and Mechanisms of Taurine as a Therapeutic Agent.

Biomolecules & Therapeutics **26**, 225–241 (2018).

71. Grajeda-Iglesias, C. *et al.* Oral administration of Akkermansia muciniphila elevates

systemic antiaging and anticancer metabolites. *Aging (Albany NY)* **13**, 6375–6405 (2021).

72. Durand, S. *et al.* The intracellular metabolome of starving cells. in *Methods in Cell*

Biology vol. 164 137–156 (Elsevier, 2021).

73. Vembar, S. S., Macpherson, C. R., Sismeiro, O., Coppée, J. Y. & Scherf, A. The pfalba1

rna-binding protein is an important regulator of translational timing in plasmodium

falciparum blood stages. *Genome Biol* **16**, (2015).

74. Fried, M. *et al.* Multilaboratory approach to preclinical evaluation of vaccine

immunogens for placental malaria. *Infect Immun* **81**, 487–495 (2013).

75. Bryant, J. M. *et al.* Crispr/cas9 genome editing reveals that the intron is not essential for

var2csa gene activation or silencing in plasmodium falciparum. *MBio* **8**, (2017).

76. Nacer, A. *et al.* clag9 Is Not Essential for PfEMP1 Surface Expression in Non-

Cytoadherent Plasmodium falciparum Parasites with a Chromosome 9 Deletion. *PLoS*

ONE **6**, e29039 (2011).

Acknowledgments: We thank Jessica Bryant for critical reading of this manuscript. We

thank Célia Peuziat and Sooraj Shivakumar for technical assistance with this project.

Funding: This work was supported by funding to AS from the Laboratoire d'Excellence

(LabEx) ParaFrap (ANR-11-LABX- 0024), ANR VAR2PTM (ANR-22-CE15-0023).

This work was also supported by funding to AC from MRC/LSHTM fellowship,

the Agence Nationale de la Recherche (ANR-18☐CE15☐0009☐01). This work was

additionally supported by funding to BG from the Fondation pour la Recherche

Médicale (Equipe FRM EQU202203014741).

Author contributions:

Conceptualization: GD, AS

Methodology: GD, PM, DD

Investigation: GD, PM, PC, FA, DD, SD

Visualization: GD

Funding acquisition: AS, AC, BG

Project administration: GD, AS

Supervision: AS, AC, BG

Writing – original draft: GD, AS

Writing – review & editing: GD, AS, AC, BG

Competing interests:

The funders had no role in the study design, data collection, and interpretation or decision to

submit the work for publication. The authors declare that they have no conflict of interest.

Data and materials availability: All data are available in the main text or the supplementary

materials.

999
1000

A

Human plasma collection

The Gambia

Longitude 13.2°N 13.6°N

5km

Latitude 15.0°W 14.5°W 14.0°W

	October	November	December	January	February	March	April	May
Symptomatic	10	28						
Asymptomatic	24	22	20	16	13	14	13	14
Negative			6	11	8			

High transmission | Low transmission

B

1. Volunteers from The Gambia → 2. *P. falciparum*-infected and uninfected human plasma → 3. GC-QqQ/MS → 4. Widely targeted metabolomics → 78 Metabolites used in analysis

x10⁸
0,9
0.1
Count vs Acquisition time (min)
6 12 18 24 30 36

C

Metabolite Class

78 Metabolites

- Amino acids 28.9%
- Carbohydrates 16.9%
- Bases 6%
- Nucleosides 2.4%
- Saturated FAs 6%
- TCA 6%
- Organic acids 12%
- PUFAs 2.4%
- Unclassified 4.8%
- Dicarboxylic acids 1.2%
- Prenols 1.2%
- MUFAs 2.4%
- SCFAs 6%
- Steroids 1.2%
- Phosphotidylcholine 1.2%
- Polyamines 1.2%

D

Overview of Enriched Metabolite Sets

Alanine, aspartate and glutamate metabolism
D-Amino acid metabolism
Galactose metabolism
Valine, leucine and isoleucine biosynthesis
Arginine biosynthesis
TCA cycle
and dicarboxylate metabolism
Butanoate metabolism
e and proline metabolism
and sucrose metabolism
curonate interconversion
te and CoA biosynthesis
Pyruvate metabolism
and threonine metabolism
nd tryptophan biosynthesis
se and mannose metabolism
Nitrogen metabolism
Glycolysis / Glucoenogenesis
Valine, leucine and isoleucine degradation
Phenylaline metabolism
Taurine and hypotaurine metabolism
Purine metabolism
Neomycin, kanamycin and gentamicin biosynthesis
Amino sugar and nucleotide sugar metabolism
Glutathione metabolism

P-value
0.15
0.10
0.05
0.00

Enrichment Ratio
5
10

-log10(p-value)
2.5 5.0 7.5 10.0

E

Individuals − PCA

Dim2 (10.9%)

Asymptomatic

Negative

Symptomatic

Conditions
- negative−AS−Dec
- negative−AS−Jan
- negative−AS−Feb
- positive−S−Oct
- positive−S−Nov
- positive−AS−Oct
- positive−AS−Nov
- positive−AS−Dec
- positive−AS−Jan
- positive−AS−Feb
- positive−AS−Mar
- positive−AS−Apr
- positive−AS−May

Dim1 (48.1%)
−10 −5 0 5 10 15

A Age Distribution

B Sample Distribution

Month	Condition	Median age (years)	% Male	% Female
October	Positive Symptomatic	19	50%	50%
	Positive Asymptomatic	9.5	37.5%	62.5%
November	Positive Symptomatic	20.5	50%	50%
	Positive Asymptomatic	18	45.5%	54.5%
December	Negative	30	33.3%	66.7%
	Positive Asymptomatic	12.5	40%	60%
January	Negative	11	27.3%	72.7%
	Positive Asymptomatic	16.5	37.5%	62.5%
February	Negative	9	25%	75%
	Positive Asymptomatic	13	46.2%	53.8%
March	Positive Asymptomatic	14	42.9%	57.1%
April	Positive Asymptomatic	18	35.7%	64.3%
May	Positive Asymptomatic	18.5	35.7%	62.3%

C Parasitemia

D

Color Key

log2FC median

-2 -1 0 1 2

Asymptomatic | **Symptomatic** | **Negative**

Asymptomatic: May, Apr, Mar, Feb, Jan, Dec, Nov, Oct
Symptomatic: Nov, Oct
Negative: Feb, Jan, Dec

Color Key

Row labels (top to bottom):

- ctose;Glucose;Mannose
- Inositol
- Citric acid;Isocitric acid
- Pantothenic acid
- Cholesterol
- Phosphoric acid
- Sebacic acid
- Arabinose
- Lysine
- Tyrosine
- Uric acid
- lactitol;Sorbitol;Mannitol
- Xanthine
- Uracil
- Ribose;Ribulose
- Succinic acid
- Glycerol-3-phosphate
- 4-hydroxymandelic acid
- Hypoxanthine
- Inosine
- Allantoin
- Palmitoleic acid
- Oleic acid;Elaidic acid
- Linoleic acid
- Serine
- 5-oxoproline
- 3-hydroxyisovaleric acid
- Homoserine
- 5-aminovaleric acid
- Glutamic acid
- Ornithine
- Methionine
- Threonine
- Aspartic acid
- Citrulline
- GABA
- 4-hydroxyproline
- Valine
- Phenylalanine
- Leucine
- Isoleucine
- Proline
- Arabitol
- Erythritol
- Ribitol
- Alpha-tocopherol
- Xylitol
- Xylose
- 2-aminoadipic acid
- 2-hydroxyglutaric acid
- Creatine
- Fumaric acid
- Lactic acid
- Malic acid
- Ketoisocaproic acid
- 2-hydroxybutyric acid
- Taurine
- Threonic acid
- Glyceric acid
- -phosphoethanolamine
- Glutamine
- Shikimic acid
- Tryptophan
- Asparagine
- Alanine
- Arachidonic acid
- Lauric acid
- Capric acid
- Myristic acid
- 3-hydroxybutyric acid
- Ketoisovaleric acid
- 2-oxoglutaric acid
- Guanosine
- Maltose;Trehalose
- Sucrose
- Fructose

A

Symptomatic	Asymptomatic		
Metabolites associated with Symptomatic Malaria (High Transmission Season) 2-oxoglutaric acid 3-Hydroxybutyric acid Palmitoleic acid	**Metabolites associated with Seasonality** (Comparison of Asymptomatic Malaria Cases)		
	(High Transmission Season)		(Low Transmission Season)
	Allantoin Palmitoleic acid	Glycerol-3-Phosphate 4-Hydroxymandelic acid	**Taurine** Glyceric acid Tryptophan Maltose;Trehalose Guanosine O-Phosphoethanolamine
	Metabolites increased in Chronic Asymtomatic Malaria (Low Transmission Season)		
	Taurine, Guanosine, Maltose;Trehalose, Threonic acid, Glyceric acid		
	Metabolites increased in Asymtomatic Malaria and not in uninfected individuals		
	Taurine 5-Aminovaleric acid Leucine	Ketoisovaleric acid Valine Isoleucine	O-Phosphoethanolamine Proline Allantoin

B

C

A

Longitudinal metabolite levels in the same individuals (n=16)

B

Valine

Taurine

Proline

Palmitoleic acid

O-phosphoethanolamine

Leucine

Ketoisovaleric acid

Isoleucine

Galactitol;Sorbitol;Mannitol

Arabinose

Allantoin

5-aminovaleric acid

2-oxoglutaric acid

Relative Abundance

Uninfected Individuals

Infected Asymptomatic Individuals

A

Plasma Taurine levels

B

CD36 Cytoadhesion

- Control
- [0.04mM] Methionine
- [0.02mM] Taurine
- [0.04mM] Taurine
- [0.08mM] Taurine
- [0.16mM] Taurine
- [0.16mM] Taurine 1-hour

C

CD36 Cytoadhesion

Control   [0.16mM] Taurine

0.1mm   0.1mm

D

EPCR Cytoadhesion

- Control
- [0.16mM] Taurine
- [0.64mM] Taurine

A **Taurine and Threonic Acid Growth Curve**

Legend:
● Control
■ [0.16mM] Taurine
▲ [0.27mM] Threonic acid

C **CD36 Cytoadhesion**

NS

Legend:
● Control
● [0.27mM] Threonic Acid 1-hour

B **3D7 : CD36 Cytoadhesion Replicate Counts**

Legend:
● Control
■ [0.04mM] Methionine
▲ [0.02mM] Taurine
▼ [0.04mM] Taurine
◆ [0.08mM] Taurine
● [0.16mM] Taurine
● [0.16mM] Taurine 1-hour

D **FCR3 *var19* : CD36 Cytoadhesion**

***

Legend:
● Control
■ Taurine [0.16mM]

E **NF54 *var2CSA* : CSA Cytoadhesion**

*

Legend:
● Control
▲ Taurine [0.64mM]

F **CD36 + Human Serum Binding**

*

**

Legend:
● Control
■ 10% Serum
▲ 10% serum + [0.64mM] Taurine

A

CD36 Reversible Binding

% Binding

- ● Control
- ▲ Taurine [0.64mM]

***

Condition

B

EPCR Reversible Binding

% Binding

- ● Control
- ▲ Taurine [0.64mM]

*

Condition

C

CD36 Human Endoethelial Cell Binding

Bound iRBC mm^2

ns

*

**

- ● Control
- ■ [0.16mM] Taurine
- ▲ [0.64mM] Taurine
- ◆ [0.93mM] Taurine

Condition

D

CD36 Human Endoethelial Cell Binding

Control   [0.93mM] Taurine

A

PfEMP1 Surface Expression

% Surface Expression

- Control
- [0.64mM] Taurine

ns

Condition

B

Taurine : CD36 Binding

% Binding

ns

- Control
- [0.16mM] Taurine during blocking

Condition

C

3D7 : CD36 : C32 Human Cell Binding

Bound iRBC mm^2

- Control
- [0.93mM] Taurine

Condition

D

3D7 : CD36 : C32 Human Cell Binding

Control   [0.93mM] Taurine

E

F

	Control	Taurine [0.64mM]

Replicate 1

	Percentage	Geometric Mean (PE-A)
Control	54.7 %	7124
Taurine [0.64mM]	58.8 %	8542

Percentage Geometric Mean (PE A)

Taurine [0.64mM]

Percentage Geometric Mean (PE A)

Taurine [0.64mM]

Percentage Geometric Mean (PE A)

Replicate 2

Taurine [0.64mM]

	Percentage	Geometric Mean (PE-A)
Control	37.5 %	7839
Taurine [0.64mM]	43.8 %	10394

Percentage Geometric Mean (PE A)

Taurine [0.64mM]

Percentage Geometric Mean (PE A)

Taurine [0.64mM]

Percentage Geometric Mean (PE A)

Taurine [0.64mM]

Replicate 3

	Percentage	Geometric Mean (PE-A)
Control	43.2 %	10629
Taurine [0.64mM]	44.5 %	10115