Title:

Selective disruption of lipid peroxide homeostasis in intratumoral regulatory T cells by targeting FSP1 enhances cancer immunity

Authors:
Jesse Garcia Castillo[1], Stephanie Silveria[1], Antoine Sauquet[1], Leo Schirokauer[1], Joseph Hendricks[2,3], Hei Sook Sul[3], James A. Olzmann[2,3], and Michel DuPage[1*]

Affiliations:
[1] Division of Immunology and Molecular Medicine, Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA
[2] Division of Molecular Therapeutics, Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA
[3] Department of Nutritional Sciences and Toxicology, University of California, Berkeley, Berkeley, CA 94720, USA

*Leading Corresponding Contact: dupage@berkeley.edu

Abstract:
A burgeoning approach to treat cancer is the pharmacological induction of ferroptotic cell death of tumor cells. However, the impact of disrupting anti-ferroptotic pathways in the broader tumor microenvironment (TME), such as in immune cells, is still undefined and may complicate treatments. Here, we show that Ferroptosis Suppressor Protein 1 (FSP1/*Aifm2*) is critically required for regulatory T cell (Treg) resistance to ferroptosis and their immunosuppressive function within the TME. Compared to other canonical ferroptosis regulators such as GPX4, GCH1, and NRF2, only FSP1 was induced upon T cell activation. Deletion of *Aifm2* in all T cells, or Tregs specifically, enhanced tumor control by selectively disrupting Treg immunosuppression within tumors without inciting autoimmune pathology in mice. As opposed to deletion of *Gpx4* in all T cells, T cell deletion of *Aifm2* did not impair antigen-specific CD8$^+$ T cell responses. These results reveal a unique opportunity for targeting a regulator of ferroptosis that can not only directly target cancer cells, but also simultaneously enhance anti-cancer immune responses without inciting autoimmunity.

Main Text:

Ferroptosis, an iron-dependent form of regulated cell death driven by unchecked lipid peroxide accumulation, has emerged as a promising approach for treating therapy-resistant cancers(*1–3*). Unlike apoptosis, ferroptosis can release danger-associated

molecular patterns (DAMPs) that activate immune responses(*4*, *5*). While glutathione
peroxidase 4 (GPX4) is the primary defense against ferroptosis, cancer cells often
express alternative protective ferroptotic regulators, such as ferroptosis suppressor
protein 1 (FSP1) and GTP cyclohydrolase 1 (GCH1), to resist GPX4 inhibition(*6–11*).

Regulatory T cells (Tregs), a suppressive subset of CD4$^+$ T cells, pose a major
barrier to effective cancer immunotherapy by inhibiting antitumor responses through
direct cell contact and the release of immunosuppressive molecules(*12–14*). Tregs are
uniquely adapted to the nutrient-deprived and hypoxic tumor microenvironment (TME)
by preferentially using oxidative phosphorylation (OXPHOS), importing lipids via CD36,
and utilizing lactate and fatty acids as alternative carbon sources for their
metabolism(*15–18*). Moreover, Tregs upregulate antioxidant pathways—including
glutathione synthesis and thioredoxin—to mitigate oxidative stress and preserve their
function in the TME(*19–21*). Although Tregs depend on GPX4 to prevent ferroptosis,
inhibiting GPX4 is not a viable strategy for cancer immunotherapy as its inhibition also
impairs effector CD8$^+$ and CD4$^+$ T cell survival(*22*, *23*).

In this study, we establish that FSP1/*Aifm2* is a critical effector protein for Treg
resistance to ferroptosis induction *in vitro* and Treg function *in vivo* in tumors. We found
that Tregs use FSP1 to prevent lipid peroxide accumulation despite having higher levels
of intracellular reactive oxygen species (ROS). *Aifm2*-deficiency was sufficient to
increase the sensitivity of Tregs to ferroptosis induction with GPX4 inhibition *in vitro*.
Aifm2-deficiency in Tregs decreased the expression of proteins essential for Treg
effector function, but did not impact effector T cell immune homeostasis or peripheral
tolerance in young or aged mice. Total ablation of *Aifm2* in all T cells did not disrupt
antitumor T cell responses but did diminish the immunosuppressive function of Tregs.
These findings are significant for the translation of pharmacological FSP1 inhibitors for
cancer as they show that FSP1 inhibition not only can directly sensitize cancer cells to
ferroptosis, but it may also enhance immune responses to cancer by selectively
blocking Treg function in the TME without generating autoimmune toxicity.
67
Results:
Tregs resist lipid peroxide accumulation and ferroptosis
To understand the regulation of ROS across different T cell populations, we
FACS-purified naive Tregs and conventional CD4$^+$ T cells (Tconv) for activation *in vitro*
and measuring their total ROS and lipid peroxide levels using fluorescent reagents
detected by flow cytometry **(Fig. 1A)**. In accordance with numerous studies, we
observed that Tregs had increased (~2-fold) intracellular ROS levels compared to Tconv
cells(*18*, *24*) **(Fig. 1B)**. Next, we directly measured lipid peroxidation by measuring the
amount of oxidized BODIPY C11 and observed that Tregs had decreased amounts of
lipid peroxides compared to Tconv cells **(Fig. 1C)**. While RSL3 increased oxidized C11-

BODIPY in both Tregs and Tconv cells, Tconv cells increased CD11-BODIPY^{OX}
significantly more than Tregs **(Fig. 1D)**. Importantly, only Ferrostatin-1 (Fer-1), a radical-
trapping antioxidant, but not the pan-caspase inhibitor Z-VAD-FMK or RIP1 kinase
inhibitor Nectrostatin-1 (Nec-1) treatments reversed CD11-BODIPY^{OX} elevation, as
expected **(Fig. 1D)**. Improved control of lipid oxidation ultimately made Tregs more
resistant to ferroptotic cell death with the inhibition of GPX4 across a series of
concentrations of RSL3 **(Fig. 1E)**. Decreased cell viability was confirmed to be due to
ferroptosis, as only treatment with Fer-1, but not other programmed cell death inhibitors,
was able to rescue cell viability **(Fig. 1F)**. Interestingly, the differences in sensitivity to
ferroptosis were not due to the expression of the essential Treg transcription factor
Foxp3, as *in vitro* induced Tregs (iTregs), were as sensitive to GPX4 inhibition as Tconv
cells **(Fig. 1E)**(*19*). To confirm these findings without activating or culturing T cells long
term *in vitro*, we isolated T cells from spleens of mice and treated them with RSL3 alone
or RSL3 and Fer-1 for 4 hours *ex vivo*. This revealed that while CD8$^+$ and CD4$^+$ Tconv
cells were highly sensitive to GPX4 inhibition, Tregs remained highly resistant to
ferroptosis *ex vivo* **(Fig. 1G)**. Therefore, despite having higher amounts of ROS, Tregs
reduce their accumulation of lipid peroxides, which likely increases their resistance to
ferroptosis.
96
FSP1 expression in Tregs confers resistance to ferroptosis
T cell activation increases levels of ROS, which can be tuned by the expression
of antioxidant genes to regulate cytosolic signaling pathways(*22, 25–29*). Analysis of
gene expression by RNA-sequencing of FACS-purified Tregs and Tconv cells that were
naive (CD44^{lo}CD62L^{hi}) versus four days after anti-CD3/anti-CD28 *in vitro* activation,
demonstrated that only *Aifm2* increased expression, but not other anti-ferroptotic
proteins, e.g. *Gpx4*, *Gch1*, or *Nfe2l2*, upon *in vitro* T cell activation **(Fig. 2A)**(*30*). In
addition, a comparison of freshly sorted *in vivo* activated (CD44^{hi}CD62L^{lo}) Tregs
compared to naive Tregs from mice showed a similar induction of *Aifm2* in activated
Tregs **(Fig. 2B)**(*30*). Moreover, direct comparison of *in vitro* activated Tregs versus
Tconv cells revealed a 2-3 fold greater induction of *Aifm2* and *Gch1* expression, and
reduced expression of *Nfe2l2* and *Ascl3*, in Tregs compared to Tconvs **(Fig. 2C)**. These
results demonstrate that *Aifm2*/FSP1 is induced during T cell activation, but that Tregs
induce higher expression of *Aifm2* and *Gch1* antioxidant genes with activation, which
may underlie their enhanced resistance to ferroptosis.
While GCH1 was shown to be crucial for T cell proliferative capacity, the role of
FSP1 in Treg biology is unexplored(*29*). To define the role of FSP1 in Tregs, we
crossed mice carrying LoxP-flanked *Aifm2^{fl}* alleles with *Foxp3-GFP-hCre* to specifically
delete *Aifm2* in Tregs (called Treg.*Aifm2$^{\Delta/\Delta}$* mice)(*31*). Treatment of *Aifm2*-deficient
Tregs with RSL3 to induce ferroptosis revealed that *Aifm2*-deficient Tregs now exhibited

significantly increased sensitivity to RSL3-mediated cell death compared to wildtype
Tregs (**Fig. 2D)**. Thus, FSP1 promotes Treg resistance to ferroptosis *in vitro*.
Finally, comparative gene expression analysis from human peripheral blood
mononuclear cells (PBMCs) confirmed that human Tregs that are activated (memory
Treg) also exhibit the highest expression of *Aifm2* compared both to naive Tregs as well
as all other activated T cell populations (**Fig. 2E-F**). These results indicate that
FSP1/*Aifm2* expression protects Tregs from ferroptosis and FSP1/*Aifm2* expression is
higher in both mouse and human activated Tregs compared to other T cell subsets.
125
Treg-specific FSP1 deletion does not impact peripheral tolerance
To directly test whether Treg-specific deletion of *Aifm2* disrupted Treg function *in*
vivo, we analyzed lymphoid and nonlymphoid tissues from wildtype versus
Treg.*Aifm2$^{\Delta/\Delta}$* mice at early (16 weeks) and late (28 weeks) stages of life. There were no
changes to the frequency of CD4$^+$ Tconv, CD8$^+$ T, or Tregs across lymphoid organs
(lymph node, spleen) at any age (**Fig. 3A and Fig. S1A-B**). However, in the organ
tissues examined (colon, skin, lung), while CD4$^+$ Tconv and CD8$^+$ T cell frequencies did
not change in Treg.*Aifm2$^{\Delta/\Delta}$* mice, we did note a modest reduction in Treg frequency in
the lung and colon of Treg.*Aifm2$^{\Delta/\Delta}$* mice **(Fig. 3A and Fig. S1C)**. Additionally, there
were no changes to the activation state of CD4$^+$ Tconv, CD8$^+$ T, or Tregs in all tissues
examined (**Fig. 3B and Fig. S1D**). These data suggest that Treg-specific *Aifm2*-
deficiency does not increase the prevalence of autoimmunity in mice.
Phenotyping of the Tregs from Treg.*Aifm2$^{\Delta/\Delta}$* mice revealed no significant
changes in the expression of canonical lineage defining Treg markers (i.e. CD25,
Foxp3, and NRP1) in either lymphoid or organ tissues **(Fig. 3C and Fig. S1E)**.
However, there was a significant decrease in CD73 expression on colonic and lung
Tregs **(Fig. 3D-E).** CD73, in conjunction with CD39, acts as an ectonucleoside that
breaks down ATP to generate adenosine, a critical immunosuppressive molecule
generated by Tregs. Interestingly, both CD73 and CD39 expression were significantly
decreased in the lymphoid tissues of aged mice **(Fig. S1E)**. Altogether, these results
show that *Aifm2*-deficiency in Tregs does not dramatically impact the frequencies or
phenotype of Tregs in lymphoid or non-lymphoid tissues. Thus, *Aifm2*-deficiency in
Tregs does not lead to a systemic breakdown in immunotolerance or organ-specific
autoimmunity.
150
FSP1-deficiency specifically in Tregs impairs immunosuppression in response to
cancer
Since the pharmacological inhibition of FSP1 in cancer cells is moving to clinical
trials, we next wanted to assess tumor Treg responses to the inhibition of FSP1. First,
we tested whether different subsets of T cells from tumors exhibited different levels of
resistance to ferroptosis induction, similar to experiments in Figure 1G. Here, we grew

MC38 colon carcinoma tumors subcutaneously for 14 days in wildtype mice, then
harvested tumors, made single cell suspensions, and treated all cells for four hours with
or without 1µM RSL3 to induce ferroptosis. Strikingly, Tregs again exhibited strong
resistance to the induction of ferroptosis with GPX4 inhibition as compared to Tconv
and CD8$^+$ T cells **(Fig. 4A)**. To test whether Treg-specific FSP1 activity promoted tumor
Treg resistance to ferroptosis and supported tumor growth, we compared MC38 tumor
growth in wildtype versus Treg.*Aifm2$^{Δ/Δ}$* mice **(Fig. 4B)**(*32*). Tumor growth was
significantly reduced in Treg.*Aifm2$^{Δ/Δ}$* mice **(Fig. 4C)**. While Treg.*Aifm2$^{Δ/Δ}$* mice did not
exhibit changes to Treg frequencies or tumor-specific CD8$^+$ T cells in tumors (**Fig. 4D**
and Fig. S2A), Treg.*Aifm2$^{Δ/Δ}$* mice did exhibit increased CD8 to Treg ratios in tumors,
indicative of a selective impact on Tregs compared to effector T cells in tumors with
FSP1-deficiency in Tregs (**Fig. 4D)**. In addition, we did not observe altered expression
of the exhaustion markers PD-1 or LAG-3 in CD8$^+$ T cells in Treg.*Aifm2$^{Δ/Δ}$* mice (**Fig.**
S2B). Importantly, to demonstrate that the reduction in tumor growth was due to the
accumulation of lipid peroxides in FSP1-deficient Tregs, we treated wildtype and
Treg.*Aifm2$^{Δ/Δ}$* tumor-bearing mice with the radical trapping antioxidant liproxstatin-1
(Lip-1), a ferroptosis inhibitor that is functional *in vivo in mice(33)* (**Fig. 4E**). Lip-1
treatment restored tumor growth in Treg.*Aifm2$^{Δ/Δ}$* mice, while Lip-1 treatment had no
effect on tumor growth in wildtype mice (**Fig. 4F**). These results indicate that FSP1
activity in Tregs is required to control their accumulation of lipid peroxides within the
TME, and disruption of FSP1 activity can release immunosuppression by Tregs and
promote better tumor control.
179
FSP1-deficiency in all T cells enhances T cell responses to infection and cancer
T cell deficiency for *Gpx4*- or *Gch-1* was recently shown to negatively impact T
cell responses in several disease settings(*23, 29, 34*). To test if *Aifm2*-deficiency also
impacts all T cell responses, which would occur with pharmacological inhibition for
cancer treatment, we generated *Aifm2*-deficient T cell mice (*CD4-Cre;Aifm2$^{fl/fl}$*, called
T.Aifm2$^{Δ/Δ}$) and infected these mice with *Listeria monocytogenes*, an intracellular
pathogen known for stimulating a robust CD8$^+$ T cell response (**Fig. S3A**)(*35, 36*).
Importantly, *T.Aifm2$^{Δ/Δ}$* mice had no deficiency in bacterial control, as there was no
difference in bacterial CFUs recovered from spleens or livers of wildtype versus
T.Aifm2$^{Δ/Δ}$ six days post infection (**Fig. S3B**). Upon *Listeria* infection, we observed a
decrease in the frequency of Tregs as a percentage of CD4$^+$ T conv cells, as well as of
total CD45$^+$ immune cells, in *T.Aifm2$^{Δ/Δ}$* (**Fig. S3C**). The frequency of total CD8$^+$ T cells,
Listeria-specific CD8$^+$ T cells, or CD4$^+$ Tconv cell populations, however, did not change
(**Fig. S3D-F**).
Next, to test the impact of deleting *Aifm2* in all T cells, as would happen with a
pharmacological FSP1 inhibitor, in the setting of cancer, we implanted MC38 tumors
into wildtype versus *T.Aifm2$^{Δ/Δ}$* mice and measured tumor growth (**Fig. 5A**). Tumor

growth was significantly decreased in mice harboring *Aifm2*-deficient T cells (**Fig. 5B**).
While there was no change in the frequency of Tregs in MC38 tumors, there was an
increase in the CD8:Treg ratios in tumors from *T.Aifm2$^{\Delta/\Delta}$* compared to wildtype mice,
as observed with Treg-specific *Aifm2*-deficiency (**Fig. 5C-D**)(*37, 38*). Assessment of
CD8$^+$ T cell exhaustion in tumors by staining for both PD-1 and Lag-3 proteins showed
no changes in total CD8$^+$ T cells **(Fig. 5E)**. However, tumor-specific CD8$^+$ T cells, while
unchanged in total frequencies, exhibited decreased frequencies of PD-1$^+$Lag3$^+$
MuLV/H2K^{b+} CD8$^+$ T cells (**Fig. 5F**)(*39*). These results starkly contrast with the
consequences of T cell deficiencies in other antioxidant genes, i.e. *Gpx4* or *Gch1*,
where deficient T cells completely lost their capacities to control infections or cancer(*23,*
29). Instead, deficiency for FSP1/*Aifm2* in all T cells completely preserves CD8$^+$ and
CD4$^+$ Tconv cell function while potently disrupting the function of Tregs.
209
Discussion:
In this study, we demonstrate that FSP1 (*Aifm2*) plays a critical role, and unique
role amongst T cells, in sustaining Treg function within tumors. Unlike other ferroptosis
resistance factors such as GPX4, which are essential for proper T cell function during
infections or cancer, FSP1 appears to be selectively required for Tregs without
compromising effector T cell responses. We show that FSP1 expression is induced
during T cell activation, distinguishing it from other ferroptosis regulators like GPX4,
GCH1, or NRF2. Our data indicate that FSP1 supports Treg survival and suppressive
function by limiting lipid peroxide accumulation despite high levels of ROS generation in
Tregs. This suggests that increased FSP1 expression in Tregs may have evolved to
prevent sublethal oxidative damage that would otherwise kill or impair Tregs.
These findings reveal a novel paradigm in which targeting FSP1 may specifically
modulate Treg-mediated immunosuppression in cancer without affecting effector T
cells. In contrast to GPX4-targeted approaches, which can dampen overall T cell
responses and carry a narrow therapeutic window due to GPX4's essential role in
multiple tissues, inhibition of FSP1 appears to selectively compromise Treg function(*23,*
34, 40, 41). In addition, Tregs are sensitive to specific antioxidant pathway activity as
the genetic deletion of KEAP1 leads to loss of Treg lineage identity that leads to
spontaneous autoimmunity due to hyperactivation of NRF2 and uncontrolled induction
of antioxidant pathways(*42*). Our tumor models showed that genetic ablation of FSP1 in
T cells resulted in better tumor control, potentially due to a reduction in Treg
suppressive capacity and a concomitant decrease in canonical exhaustion markers
(e.g., PD-1, Lag3) on tumor-specific CD8$^+$ T cells(*43*).
The implications of these findings are significant. By specifically targeting FSP1,
it may be possible to alleviate the immunosuppressive effects of Tregs within the TME,
thereby enhancing the efficacy of cancer immunotherapies. Moreover, FSP1 inhibition
exhibits great potential to synergize with immune checkpoint blockade (e.g., anti-PD-1

therapies) and cellular immunotherapies such as CAR T cells by preventing T cell
exhaustion and preserving cytotoxic function. PD-1 blockade has also been shown to
induce IFN-γ production by CD8⁺ T cells, which induced ferroptosis in cancer cells
directly by impacting the lipid composition through Ascl4 expression(*44*). Thus, this dual
approach, targeting tumor cells directly through ferroptosis induction while
simultaneously modulating Treg function, could offer a much broader therapeutic
strategy for treating cancer.
Our study also underscores the importance of metabolic regulation in Treg
biology. Tregs exhibit a unique metabolic profile characterized by a reliance on oxidative
phosphorylation and fatty acid utilization, which supports their survival in the nutrient-
deprived, hypoxic TME(*15–17*). Despite high ROS levels due to these forms of
metabolism, Tregs maintain low lipid peroxide accumulation, in part by maintaining
higher expression of FSP1. Disrupting this balance through FSP1 inhibition leads to
functional defects in Tregs without appearing to cause their complete elimination in vivo,
which is preferable for preserving immune homeostasis.
In conclusion, our results position FSP1 as a Treg-specific ferroptosis regulator
and a promising therapeutic target for enhancing cancer immunity. By selectively
modulating FSP1 activity, it may be possible to reduce Treg-mediated suppression in
tumors, preserve effector T cell responses, and ultimately improve the outcomes of
cancer immunotherapy. Future studies should focus on translating these findings into
pharmacological interventions, as well as exploring the potential of FSP1-targeting
strategies in combination with existing immunotherapies to achieve synergistic effects.
259

Materials and Methods:
Mice
CD4-Cre mice were obtained from Jackson laboratories (JAX:022071) and bred in
house. Foxp3-GFP-hCre mice were obtained from Jackson laboratories (JAX:023161)
and bred in house. *Aifm2^{fl}* mice were generated as described(*31*). For tumor studies,
syngeneic C57BL/6J mice were inoculated with 5.0×10^5 MC38 in PBS subcutaneously.
Tumor measurements were performed blindly across the entire experiment by a single
operator measuring three dimensions of the tumor with calipers three times per week.
For in vivo liproxstatin treatments (Selleck Chemicals), mice were injected
intraperitoneally with 10mg/kg of liproxstatin-1 dissolved in DMSO and diluted in
PEG300, Tween-80, and dH2O. All the experiments were conducted according to the
Institutional Animal Care and Use Committee guidelines of the University of California,
Berkeley.
273

274 **Cell lines**
275 MC38 cell lines were kindly provided by Dr. Jeff Bluestone's lab(*45*). All cell lines were
276 maintained in DMEM (GIBCO) supplemented with 10% FBS, sodium pyruvate (GIBCO),

277 10mM HEPES (GIBCO), and penicillin-streptomycin (GIBCO). Tumor cells were grown
278 at 37℃ with 5% CO_2.
279
280 **Listeria monocytogenes strains**
281 All strains of L. monocytogenes were derived from the wild-type 10403S strain. All
282 strains were cultured in filter-sterilized nutrient-rich Brain Heart Infusion (BHI) media
283 (BD Biosciences) containing 200 µg/mL streptomycin (Sigma-Aldrich).
284
Intravenous and Intratumoral Listeria infection
Overnight cultures were grown in BHI + 200 µg/mL streptomycin at 30℃. The following
day, bacteria were grown to logarithmic phase by diluting the overnight culture in fresh
BHI + 200 µg/mL streptomycin and culturing at 37℃ shaking. Log-phase bacteria were
washed and frozen in 9% glycerol/PBS. For intravenous infections, frozen stocks were
diluted in PBS to infect via the tail vein with $5x10^3$ CFU log-phase bacteria. The mice
were euthanized 6 days after intravenous injections and organs were collected for CFU
or flow analysis.
293
Primary Treg and T cell culture
Spleens and lymph nodes were collected from Foxp3-GFP-hCre;R26$^{LSL-RFP}$; *Aifm2$^{+/+}$*,
Foxp3-GFP-hCre;R26$^{LSL-RFP}$; *Aifm2$^{Δ/Δ}$*, CD4-Cre;*Aifm2$^{+/+}$* or CD4-Cre;*Aifm2$^{Δ/Δ}$*. Single
cell suspensions were generated and enriched for CD4+ T cells by negative selection
using EasySep magnetic bead kit (STEMCELL Technologies) and stained with anti-
mouse CD4 (RM4-5, BioLegend), anti-CD62L (MEL1-14, Biolegend), anti-CD8 (53-6.7,
Biolegend), anti-CD25 (PC61, BioLegend), anti-CD357 (GITR) (DTA-1, Biolegend).
Naïve Treg cells (CD4$^+$Foxp3GFP$^+$RFP$^+$CD62L$^+$ or CD4$^+$CD25$^+$GITR$^+$CD62L$^+$) and
naive CD4$^+$ T cells (CD4$^+$CD62L$^+$Foxp3GFP$^-$RFP$^-$ or CD4$^+$CD62L$^+$GITR$^-$CD25$^-$) were
sorted using an Aria Fusion sorter (BD Biosciences) with a 70µm nozzle. Cells were
activated with anti-CD3 and anti-CD28 coated beads (Dynabeads Mouse T-Activator
CD3/CD28, Invitrogen) at a ratio of 1:1 for Tconv or 1:3 ratio for Tregs (cell:bead). Cells
were kept at a concentration of 10^6 cells/ml in DMEM medium supplemented with 10%
FBS, non-essential amino acids, sodium pyruvate, L-glutamine, HEPES, and β-ME, and
cells cultured more than 24 hr were supplemented with 200–2,000 IU/ml recombinant
human IL-2 (NIH).
310
Tissue Collection and preparation for Flow cytometry
Flow cytometry was performed on an BD LSR Fortessa X20 (BD Biosciences), CyTEK
Aurora (CyTEK Biosciences) or LSRFortessa (BD Biosciences) and datasets were
analyzed using FlowJo software (Tree Star). Single cell suspensions were prepared in
ice-cold FACS buffer (PBS with 2mM EDTA and 1% BS) and subjected to red blood cell
lysis using ACK buffer (150mM NH$_4$Cl, 10mM KHCO$_3$, 0.1mM Na2EDTA, pH7.3). Dead

cells were stained with Live/Dead Fixable Blue or Aqua Dead Cell Stain kit (Molecular Probes) in PBS at 4℃. Cell surface antigens were stained at 4℃ using a mixture of fluorophore-conjugated antibodies. Surface marker stains for murine samples were carried out with anti-mouse CD3 (17A2, BioLegend), anti-mouse CD4 (RM4-5, BioLegend), anti-mouse CD8a (53-6.7, BioLegend), anti-mouse, CD44 (IM7, BioLeged), anti-mouse CD45 (30-F11, BioLegend), anti-H-2K^b MuLV p15E Tetramer-KSPWFTTL (MBL), anti-H-2K^b SIINFEKL Tetramer (NIH Tetramer Core) in PBS, 0.5% BSA. Cells were fixed using the eBioscience Foxp3/Transcription Factor staining buffer set (eBioscience), prior to intracellular staining. Intracellular staining was performed using anti-mouse Foxp3 (FJK-16S, eBioscience) at 4℃, according to manufacturer's instructions. Cells were resuspended in PBS and filtered through a 70-μm nylon mesh before data acquisition. Datasets were analyzed using FlowJo software (Tree Star).

Statistical Methods

p values were obtained from unpaired two-tailed Student's t tests for all statistical comparisons between two groups, and data were displayed as mean $\pm$ SEMs or mean $\pm$ SD. For multiple comparisons, one-way ANOVA or two-way ANOVA was used. For tumor growth curves, two-way ANOVA was used with Sidak's multiple comparisons test performed at each time point or by multiple regression analysis p values are denoted in figures by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

References:

1. S. J. Dixon, J. A. Olzmann, The cell biology of ferroptosis. *Nat Rev Mol Cell Biol* **25**, 424–442 (2024).

2. T. Nakamura, M. Conrad, Exploiting ferroptosis vulnerabilities in cancer. *Nature Cell Biology* **26**, 1407–1419 (2024).

3. K. Hadian, B. R. Stockwell, The therapeutic potential of targeting regulated non-apoptotic cell death. *Nature Reviews Drug Discovery* **22**, 723–742 (2023).

4. The release and activity of HMGB1 in ferroptosis. *Biochemical and Biophysical Research Communications* **510**, 278–283 (2019).

5. G. Kroemer, C. Galassi, L. Zitvogel, L. Galluzzi, Immunogenic cell stress and death. *Nature Immunology* **23**, 487–500 (2022).

6. K. Bersuker, J. M. Hendricks, Z. Li, L. Magtanong, B. Ford, P. H. Tang, M. A. Roberts, B. Tong, T. J. Maimone, R. Zoncu, M. C. Bassik, D. K. Nomura, S. J. Dixon, J. A. Olzmann, The CoQ oxidoreductase FSP1 acts parallel to GPX4 to inhibit ferroptosis. *Nature* **575**, 688–692 (2019).

7. S. Doll, F. P. Freitas, R. Shah, M. Aldrovandi, M. C. da Silva, I. Ingold, A. Goya Grocin, T. N. Xavier da Silva, E. Panzilius, C. H. Scheel, A. Mourão, K. Buday, M.

Sato, J. Wanninger, T. Vignane, V. Mohana, M. Rehberg, A. Flatley, A. Schepers, A. Kurz, D. White, M. Sauer, M. Sattler, E. W. Tate, W. Schmitz, A. Schulze, V. O'Donnell, B. Proneth, G. M. Popowicz, D. A. Pratt, J. P. F. Angeli, M. Conrad, FSP1 is a glutathione-independent ferroptosis suppressor. *Nature* **575**, 693–698 (2019).

8. V. A. N. Kraft, C. T. Bezjian, S. Pfeiffer, L. Ringelstetter, C. Müller, F. Zandkarimi, J. Merl-Pham, X. Bao, N. Anastasov, J. Kössl, S. Brandner, J. D. Daniels, P. Schmitt-Kopplin, S. M. Hauck, B. R. Stockwell, K. Hadian, J. A. Schick, GTP Cyclohydrolase 1/Tetrahydrobiopterin Counteract Ferroptosis through Lipid Remodeling. *ACS Central Science*, doi: 10.1021/acscentsci.9b01063 (2019).

9. E. Mishima, J. Ito, Z. Wu, T. Nakamura, A. Wahida, S. Doll, W. Tonnus, P. Nepachalovich, E. Eggenhofer, M. Aldrovandi, B. Henkelmann, K.-I. Yamada, J. Wanninger, O. Zilka, E. Sato, R. Feederle, D. Hass, A. Maida, A. S. D. Mourão, A. Linkermann, E. K. Geissler, K. Nakagawa, T. Abe, M. Fedorova, B. Proneth, D. A. Pratt, M. Conrad, A non-canonical vitamin K cycle is a potent ferroptosis suppressor. *Nature* **608**, 778–783 (2022).

10. D.-Y. Jin, X. Chen, Y. Liu, C. M. Williams, L. C. Pedersen, D. W. Stafford, J.-K. Tie, A genome-wide CRISPR-Cas9 knockout screen identifies FSP1 as the warfarin-resistant vitamin K reductase. *Nat Commun* **14**, 828 (2023).

11. M. Soula, R. A. Weber, O. Zilka, H. Alwaseem, K. La, F. Yen, H. Molina, J. Garcia-Bermudez, D. A. Pratt, K. Birsoy, Metabolic determinants of cancer cell sensitivity to canonical ferroptosis inducers. *Nat Chem Biol* **16**, 1351–1360 (2020).

12. Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell* **168**, 707–723 (2017).

13. J. B. Wing, A. Tanaka, S. Sakaguchi, Human FOXP3 Regulatory T Cell Heterogeneity and Function in Autoimmunity and Cancer. *Immunity* **50**, 302–316 (2019).

14. M. A. Moreno Ayala, Z. Li, M. DuPage, Treg programming and therapeutic reprogramming in cancer. *Immunology* **157**, 198–209 (2019).

15. A. Angelin, L. Gil-de-Gómez, S. Dahiya, J. Jiao, L. Guo, M. H. Levine, Z. Wang, W. J. Quinn 3rd, P. K. Kopinski, L. Wang, T. Akimova, Y. Liu, T. R. Bhatti, R. Han, B. L. Laskin, J. A. Baur, I. A. Blair, D. C. Wallace, W. W. Hancock, U. H. Beier, Foxp3 Reprograms T Cell Metabolism to Function in Low-Glucose, High-Lactate Environments. *Cell Metab* **25**, 1282–1293.e7 (2017).

16. R. Newton, B. Priyadharshini, L. A. Turka, Immunometabolism of regulatory T cells. *Nature Immunology* **17**, 618–625 (2016).

17. J. Miska, C. Lee-Chang, A. Rashidi, M. E. Muroski, A. L. Chang, A. Lopez-Rosas, P. Zhang, W. K. Panek, A. Cordero, Y. Han, A. U. Ahmed, N. S. Chandel, M. S.

Lesniak, HIF-1α Is a Metabolic Switch between Glycolytic-Driven Migration and Oxidative Phosphorylation-Driven Immunosuppression of Tregs in Glioblastoma. *Cell Rep* **27**, 226–237.e4 (2019).

18. T. Maj, W. Wang, J. Crespo, H. Zhang, W. Wang, S. Wei, L. Zhao, L. Vatan, I. Shao, W. Szeliga, C. Lyssiotis, J. R. Liu, I. Kryczek, W. Zou, Oxidative stress controls regulatory T cell apoptosis and suppressor activity and PD-L1-blockade resistance in tumor. *Nature Immunology* **18**, 1332–1341 (2017).

19. H. Kurniawan, D. G. Franchina, L. Guerra, L. Bonetti, L. Soriano-Baguet, M. Grusdat, L. Schlicker, O. Hunewald, C. Dostert, M. P. Merz, C. Binsfeld, G. S. Duncan, S. Farinelle, Y. Nonnenmacher, J. Haight, D. D. Gupta, A. Ewen, R. Taskesen, R. Halder, Y. Chen, C. Jäger, M. Ollert, P. Wilmes, V. Vasiliou, I. S. Harris, C. B. Knobbe-Thomsen, J. D. Turner, T. W. Mak, M. Lohoff, J. Meiser, K. Hiller, D. Brenner, Glutathione restricts serine metabolism to preserve regulatory T cell function. *Cell metabolism* **31**, 920 (2020).

20. Z. Yan, S. K. Garg, R. Banerjee, Regulatory T cells interfere with glutathione metabolism in dendritic cells and T cells. *J Biol Chem* **285**, 41525–41532 (2010).

21. Z. Yan, S. K. Garg, J. Kipnis, R. Banerjee, Extracellular redox modulation by regulatory T cells. *Nat Chem Biol* **5**, 721–723 (2009).

22. C. Xu, S. Sun, T. Johnson, R. Qi, S. Zhang, J. Zhang, K. Yang, The glutathione peroxidase Gpx4 prevents lipid peroxidation and ferroptosis to sustain Treg cell activation and suppression of antitumor immunity. *Cell Reports* **35** (2021).

23. M. Matsushita, S. Freigang, C. Schneider, M. Conrad, G. W. Bornkamm, M. Kopf, T cell lipid peroxidation induces ferroptosis and prevents immunity to infection. *J Exp Med* **212**, 555–568 (2015).

24. S. E. Weinberg, B. D. Singer, E. M. Steinert, C. A. Martinez, M. M. Mehta, I. Martínez-Reyes, P. Gao, K. A. Helmin, H. Abdala-Valencia, L. A. Sena, P. T. Schumacker, L. A. Turka, N. S. Chandel, Mitochondrial complex III is essential for suppressive function of regulatory T cells. *Nature* **565**, 495–499 (2019).

25. M. P. Murphy, R. M. Siegel, Mitochondrial ROS fire up T cell activation. *Immunity* **38**, 201–202 (2013).

26. S. H. Jackson, S. Devadas, J. Kwon, L. A. Pinto, M. S. Williams, T cells express a phagocyte-type NADPH oxidase that is activated after T cell receptor stimulation. *Nat Immunol* **5**, 818–827 (2004).

27. J. Kwon, K. E. Shatynski, H. Chen, S. Morand, X. de Deken, F. Miot, T. L. Leto, M. S. Williams, The nonphagocytic NADPH oxidase Duox1 mediates a positive feedback loop during T cell receptor signaling. *Sci Signal* **3**, ra59 (2010).

28. L. A. Sena, S. Li, A. Jairaman, M. Prakriya, T. Ezponda, D. A. Hildeman, C.-R.

Wang, P. T. Schumacker, J. D. Licht, H. Perlman, P. J. Bryce, N. S. Chandel, Mitochondria are required for antigen-specific T cell activation through reactive oxygen species signaling. *Immunity* **38**, 225–236 (2013).

29. S. J. F. Cronin, C. Seehus, A. Weidinger, S. Talbot, S. Reissig, M. Seifert, Y. Pierson, E. McNeill, M. S. Longhi, B. L. Turnes, T. Kreslavsky, M. Kogler, D. Hoffmann, M. Ticevic, D. da Luz Scheffer, L. Tortola, D. Cikes, A. Jais, M. Rangachari, S. Rao, M. Paolino, M. Novatchkova, M. Aichinger, L. Barrett, A. Latremoliere, G. Wirnsberger, G. Lametschwandtner, M. Busslinger, S. Zicha, A. Latini, S. C. Robson, A. Waisman, N. Andrews, M. Costigan, K. M. Channon, G. Weiss, A. V. Kozlov, M. Tebbe, K. Johnsson, C. J. Woolf, J. M. Penninger, The metabolite BH4 controls T cell proliferation in autoimmunity and cancer. *Nature* **563**, 564–568 (2018).

30. D. Wang, J. Quiros, K. Mahuron, C.-C. Pai, V. Ranzani, A. Young, S. Silveria, T. Harwin, A. Abnousian, M. Pagani, M. D. Rosenblum, F. Van Gool, L. Fong, J. A. Bluestone, M. DuPage, Targeting EZH2 Reprograms Intratumoral Regulatory T Cells to Enhance Cancer Immunity. *Cell Rep* **23**, 3262–3274 (2018).

31. H. P. Nguyen, D. Yi, F. Lin, J. A. Viscarra, C. Tabuchi, K. Ngo, G. Shin, A. Y.-F. Lee, Y. Wang, H. S. Sul, Aifm2, a NADH Oxidase, Supports Robust Glycolysis and Is Required for Cold- and Diet-Induced Thermogenesis. *Mol Cell* **77**, 600–617.e4 (2020).

32. M. A. Moreno Ayala, T. F. Campbell, C. Zhang, N. Dahan, A. Bockman, V. Prakash, L. Feng, T. Sher, M. DuPage, CXCR3 expression in regulatory T cells drives interactions with type I dendritic cells in tumors to restrict CD8+ T cell antitumor immunity. *Immunity* **56**, 1613–1630.e5 (2023).

33. J. Li, J. Liu, Z. Zhou, R. Wu, X. Chen, C. Yu, B. Stockwell, G. Kroemer, R. Kang, D. Tang, Tumor-specific GPX4 degradation enhances ferroptosis-initiated antitumor immune response in mouse models of pancreatic cancer. *Science Translational Medicine*, doi: 10.1126/scitranslmed.adg3049 (2023).

34. Y. Yao, Z. Chen, H. Zhang, C. Chen, M. Zeng, J. Yunis, Y. Wei, Y. Wan, N. Wang, M. Zhou, C. Qiu, Q. Zeng, H. S. Ong, H. Wang, F. V. Makota, Y. Yang, Z. Yang, N. Wang, J. Deng, C. Shen, Y. Xia, L. Yuan, Z. Lian, Y. Deng, C. Guo, A. Huang, P. Zhou, H. Shi, W. Zhang, H. Yi, D. Li, M. Xia, J. Fu, N. Wu, J. B. de Haan, N. Shen, W. Zhang, Z. Liu, D. Yu, Selenium–GPX4 axis protects follicular helper T cells from ferroptosis. *Nature Immunology* **22**, 1127–1139 (2021).

35. A. Chávez-Arroyo, D. A. Portnoy, Why is Listeria monocytogenes such a potent inducer of CD8+ T-cells? *Cell Microbiol* **22**, e13175 (2020).

36. A. Benson, S. Murray, P. Divakar, N. Burnaevskiy, R. Pifer, J. Forman, F. Yarovinsky, Microbial infection-induced expansion of effector T cells overcomes the suppressive effects of regulatory T cells via an IL-2 deprivation mechanism. *J Immunol* **188**, 800–810 (2012).

37. J. D. French, Z. J. Weber, D. L. Fretwell, S. Said, J. P. Klopper, B. R. Haugen,
Tumor-associated lymphocytes and increased FoxP3+ regulatory T cell frequency
correlate with more aggressive papillary thyroid cancer. *J Clin Endocrinol Metab* **95**,
2325–2333 (2010).

38. H. Ohtani, O. Yoshie, Morphometric analysis of the balance between CXCR3+ T
cells and FOXP3+ regulatory T cells in lymphocyte-rich and conventional gastric
cancers. *Virchows Arch* **456**, 615–623 (2010).

39. B. J. Hos, M. G. M. Camps, J. van den Bulk, E. Tondini, T. C. van den Ende, D.
Ruano, K. Franken, G. M. C. Janssen, A. Ru, D. V. Filippov, R. Arens, P. A. van
Veelen, N. Miranda, F. Ossendorp, Identification of a neo-epitope dominating
endogenous CD8 T cell responses to MC-38 colorectal cancer. *Oncoimmunology*
9, 1673125 (2019).

40. L. Chen, W. S. Hambright, R. Na, Q. Ran, Ablation of the Ferroptosis Inhibitor
Glutathione Peroxidase 4 in Neurons Results in Rapid Motor Neuron Degeneration
and Paralysis. *J Biol Chem* **290**, 28097–28106 (2015).

41. J. P. Friedmann Angeli, M. Schneider, B. Proneth, Y. Y. Tyurina, V. A. Tyurin, V. J.
Hammond, N. Herbach, M. Aichler, A. Walch, E. Eggenhofer, D. Basavarajappa, O.
Rådmark, S. Kobayashi, T. Seibt, H. Beck, F. Neff, I. Esposito, R. Wanke, H.
Förster, O. Yefremova, M. Heinrichmeyer, G. W. Bornkamm, E. K. Geissler, S. B.
Thomas, B. R. Stockwell, V. B. O'Donnell, V. E. Kagan, J. A. Schick, M. Conrad,
Inactivation of the ferroptosis regulator Gpx4 triggers acute renal failure in mice.
Nat Cell Biol **16**, 1180–1191 (2014).

42. P. Klemm, A. Rajendiran, A. Fragoulis, C. Wruck, A. Schippers, N. Wagner, T.
Bopp, K. Tenbrock, K. Ohl, Nrf2 expression driven by Foxp3 specific deletion of
Keap1 results in loss of immune tolerance in mice. *Eur J Immunol* **50**, 515–524
(2020).

43. S.-R. Woo, M. E. Turnis, M. V. Goldberg, J. Bankoti, M. Selby, C. J. Nirschl, M. L.
Bettini, D. M. Gravano, P. Vogel, C. L. Liu, S. Tangsombatvisit, J. F. Grosso, G.
Netto, M. P. Smeltzer, A. Chaux, P. J. Utz, C. J. Workman, D. M. Pardoll, A. J.
Korman, C. G. Drake, D. A. A. Vignali, Immune inhibitory molecules LAG-3 and PD-
1 synergistically regulate T-cell function to promote tumoral immune escape.
Cancer Res **72**, 917–927 (2012).

44. CD8+ T cells and fatty acids orchestrate tumor ferroptosis and immunity via
ACSL4. *Cancer Cell* **40**, 365–378.e6 (2022).

45. J. T. Sockolosky, E. Trotta, G. Parisi, L. Picton, L. L. Su, A. C. Le, A. Chhabra, S. L.
Silveria, B. M. George, I. C. King, M. R. Tiffany, K. Jude, L. V. Sibener, D. Baker, J.
A. Shizuru, A. Ribas, J. A. Bluestone, K. C. Garcia, Selective targeting of
engineered T cells using orthogonal IL-2 cytokine-receptor complexes. *Science*
359, 1037–1042 (2018).

Acknowledgments:
We thank Hector Nolla, Alma Valleros, Kartoosh Heydari and Anita Wong Lin of the UC
Berkeley Cancer Research Laboratory Flow Cytometry Facility. This research was
supported by 1DP2CA247830-01 from the National Institutes of Health (NCI),
R01GM112948 from the National Institutes of Health, and C23CR5612 from the UC
CRCC Cancer Research Coordinating Committee. M.D. is a Pew-Stewart Scholar and a
St. Baldrick's Scholar with generous support from Hope with Hazel. J.G.C is a HHMI
Gilliam Fellow. J.A.O is a Chan-Zuckerberg investigator.

Author Contributions:
Conceptualization and Methodology, M.D., J.A.O., J.G.C.; Investigation, J.G.C., S.S.,
A.S., L.S., J.H., J.A.O., and M.D.; Writing – Original Draft, J.G.C., M.D., J.A.O.; Writing
– Review & Editing, J.G.C., M.D., J.A.O., S.S.; Supervision, M.D., J.A.O., J.G.C.;
Funding Acquisition, M.D., J.A.O.

Resource Availability
Lead Contact
Further information and requests for resources and reagents should be directed to and
will be fulfilled by the lead contact, Michel DuPage (dupage@berkeley.edu).

Materials availability
Listeria monocytogenes strains used in the study will be made available from the lead
contact upon request.

Declaration of Interests:
J.A.O. is a member of the scientific advisory board for Vicinitas Therapeutics and has
patent applications related to ferroptosis.

Supplementary Materials
Materials and Methods
Figures S1 to S3

Figure Legends:
**Fig. 1 | Regulatory T cells (Tregs) naturally resist lipid peroxide accumulation and
ferroptosis.** (A) Schematic for *in vitro* T cell assays. (B) Representative flow plots (left)
and quantification (right) of bulk reactive oxygen species (ROS) by CellROX™ staining
in Tregs versus conventional CD4$^+$ T cells (Tconv) after activation and culture for 4 days
in vitro. Data pooled from 5 independent experiments. (C) Representative flow plots for
detection of oxidized C11-BODIPY in Tregs versus Tconv that were activated for 4 days
in vitro. (D) Quantification of oxidized C11-BODIPY in Tregs versus Tconv cells *in vitro*

+/- 1μM RSL3 in combination with 2μM Fer-1, 100μM Z-VAD-FMK, or 10μM Nec-1.

Data pooled from 3 experiments. (E) Cell viability of Tregs (nTregs), CD4$^+$ Tconv, and

induced Tregs (Foxp3$^+$ iTregs) treated with the indicated concentrations of RSL3 for 4

hours. Representative data from 2 independent experiments. (F) Cell viability of Tregs

versus CD4$^+$ Tconv treated *in vitro* +/- 1μM RSL3 in combination with 2μM Fer-1,

100μM Z-VAD-FMK, or 10μM Nec-1. Data pooled from 2 experiments. (G) Frequencies

of viable CD8$^+$ T (left), CD4$^+$ Tconv (middle), and Tregs (right) from spleen plotted as

the percent of CD45$^+$ cells +/- 1μM RSL3 and 2μM Fer-1. Data from n=7 mice. For all

plots, **P<0.05, **P<0.01, ***P<0.001 by two-way ANOVA (D-F) or one-way ANOVA

(G), or student t-test (B), mean ± s.d (B, D-F) or mean ± s.e.m (G).

Fig. 2 | FSP1 drives Treg resistance to ferroptosis. (A) Fold-change expression of

indicated anti-ferroptotic genes in naive (CD44^{lo}CD62L^{hi}) versus in vitro activated (+4

days with anti-CD3/CD28) Tregs. Data from n=5 mice. (B) Fold-change in expression of

indicated anti-ferroptotic genes from sorted naive (CD44^{lo}CD62L^{hi}) versus in vivo

activated (CD44^{hi}CD62L^{lo}) Tregs. (C) Fold change in expression of indicated

ferroptosis-related genes in 4 day in vitro activated Treg versus Tconv cells. Data from

n=5 mice. (D) Treatment of wildtype versus *Aifm2*-deficient Tregs with different

concentrations of RSL3 for 4 hours. (E) *Aifm2* expression from bulk RNA-Seq of human

PBMCs populations from Database of Immune Cell eQTLs, expression, epigenomics

(DICE). (F) Differential expression analysis of *Aifm2* expression across different PBMC

populations. For all plots, **P<0.05, **P<0.01, ***P<0.001 by one-way ANOVA (A-B),

student t-test (C), or two-way ANOVA (D), mean ± s.e.m (A-C) or mean ± s.d (D).

Fig. 3 | Deletion of *Aifm2* in Tregs does not disrupt peripheral tolerance in mice.

(A) Frequency of total CD8$^+$ T cells (left), CD4$^+$ Tconv cells (middle), and Tregs(right) as

a percentage of total CD45$^+$ immune cells in lymphoid tissues and organ tissues in 16

week mice. (B) Frequencies of activated (CD44^{hi}CD62^{lo}) CD8$^+$ T cells (left), CD4$^+$

Tconv cells (middle), and Tregs (right) in secondary lymphoid organs (SLOs) in 16

week mice. (C) gMFI of Foxp3, CD25, and NRP1 in Foxp3 Tregs across lymphoid

tissues and non-lymphoid tissues in 16 week mice. (D) Schematic of pathway for

conversion of ATP to Adenosine mediated by CD39 and CD73. (E) Representative

histograms and gMFI of CD73 and CD39 in wildtype versus *Aifm2*-deficient Tregs

across lymphoid tissues and nonlymphoid tissues in 16 week mice. Data from n=5 for

Treg.*Aifm2*$^{+/+}$ and n=6 for Treg.*Aifm2*$^{\Delta/\Delta}$ mice for 16 week mice. For all plots, **P<0.05,

**P<0.01, ***P<0.001 by two-way ANOVA (A, C, E) or student's t-test (B), mean ±

s.e.m.

Fig. 4 | Deletion of *Aifm2* in Tregs improves cancer immunity. (A) Schematic for

treating single cell suspensions from MC38 tumors +/- 1μM RSL3 (left). Frequencies of

viable CD8$^+$ (left), CD4$^+$ Tconv (middle), and Tregs (right) from tumor single cell
suspensions as a percentage of CD45$^+$ cells +/- 1µM RSL3 for 4 hours. Data from n=6
mice. (B) Experimental schematic for measuring tumor growth in wildtype versus mice
with *Aifm2*-deficient Tregs. (C) Growth of MC38 tumors in wildtype versus mice with
Aifm2-deficient Tregs. Data from n=5 for Treg.Aifm2$^{+/+}$ and n=6 from *Treg.Aifm2$^{Δ/Δ}$*. (D)
Frequency of Foxp3$^+$ Tregs as a percentage of total CD45$^+$ immune cells (left) and
CD8:Treg ratio (right) in MC38 tumors at D29. (E) Experimental schematic for
measuring tumor growth in wildtype versus mice with *Aifm2*-deficient Tregs with or
without 10mg/kg liproxstatin injected i.p daily. Data from n=3 for Treg.*Aifm2$^{+/+}$* + Vehicle,
n=4 for Treg.*Aifm2$^{+/+}$* + Lip-1, n=5 for Treg.*Aifm2$^{Δ/Δ}$*+ Vehicle, and n=5 for Treg.*Aifm2$^{Δ/Δ}$*
+ Lip-1. (F) Growth of MC38 tumors in wildtype versus mice with *Aifm2*-deficient Tregs
+/- Lip-1. For all plots, **P<0.05, **P<0.01, ***P<0.001 by student t-test (A, D) or two-
way ANOVA (C, F), mean ± s.e.m.
602
Fig. 5 | Total T cell deficiency for *Aifm2* preserves CD8$^+$ T cell function and
improves tumor immunity. (A) Experimental schematic for measuring tumor growth in
wildtype versus mice with *Aifm2*-deficient T cells. (B) Growth of MC38 tumors in
wildtype versus mice with *Aifm2*-deficient T cells. Data shown from n=5-6 mice/group
from one of two experiments. (C) Frequency of Foxp3$^+$ Tregs as a percentage total
CD45$^+$ immune cells in MC38 tumors at D29 from (B). (D) CD8:Treg ratio in MC38
tumors at D29 from (B). (E) Representative flow plot for PD-1 and LAG3 staining in total
CD8$^+$ T cells (left) and quantification of the frequency of PD-1$^+$Lag3$^+$ CD8$^+$ T cells in
MC38 tumors (right). (F) Frequency of tumor-specific (MuLV/H2K^{b+}) as a percentage of
CD8$^+$ T cells (left) and frequency of PD-1$^+$Lag3$^+$ of MuLV/H2K^{b+} CD8$^+$ T cells in MC38
tumors (right). For all plots, **P<0.05, **P<0.01, ***P<0.001 by two-way ANOVA (B) or
student t-test (C, D, F), mean ± s.e.m.

Garcia Castillo J. et al. Fig. 1

Fig. 1 | Regulatory T cells (Tregs) naturally resist lipid peroxide accumulation and ferroptosis. (A) Schematic for *in vitro* T cell assays. (B) Representative flow plots (left) and quantification (right) of bulk reactive oxygen species (ROS) by CellROX™ staining in Tregs versus conventional CD4+ T cells (Tconv) after activation and culture for 4 days *in vitro*. Data pooled from 5 independent experiments. (C) Representative flow plots for detection of oxidized C11-BODIPY in Tregs versus Tconv that were activated for 4 days *in vitro*. (D) Quantification of oxidized C11-BODIPY in Tregs versus Tconv cells *in vitro* +/- 1μM RSL3 in combination with 2μM Fer-1, 100μM Z-VAD-FMK, or 10μM Nec-1. Data pooled from 3 experiments. (E) Cell viability of Tregs (nTregs), CD4+ Tconv, and induced Tregs (Foxp3+ iTregs) treated with the indicated concentrations of RSL3 for 4 hours. Representative data from 2 independent experiments. (F) Cell viability of Tregs versus CD4+ Tconv treated *in vitro* +/- 1μM RSL3 in combination with 2μM Fer-1, 100μM Z-VAD-FMK, or 10μM Nec-1. Data pooled from 2 experiments. (G) Frequencies of viable CD8+ T (left), CD4+ Tconv (middle), and Tregs (right) from spleen plotted as the percent of CD45+ cells +/- 1μM RSL3 and 2μM Fer-1. Data from n=7 mice. For all plots, **P<0.05, **P<0.01, ***P<0.001 by two-way ANOVA (D-F) or one-way ANOVA (G), or student t-test (B), mean ± s.d (B, D-F) or mean ± s.e.m (G).

Garcia Castillo, J. et al. Fig. 2.

Fig. 2 | FSP1 drives Treg resistance to ferroptosis. (A) Fold-change expression of indicated anti-ferroptotic genes in naive (CD44^{lo}CD62L^{hi}) versus in vitro activated (+4 days with anti-CD3/CD28) Tregs. Data from n=5 mice. (B) Fold-change in expression of indicated anti-ferroptotic genes from sorted naive (CD44^{lo}CD62L^{hi}) versus in vivo activated (CD44^{hi}CD62L^{lo}) Tregs. (C) Fold change in expression of indicated ferroptosis-related genes in 4 day in vitro activated Treg versus Tconv cells. Data from n=5 mice. (D) Treatment of wildtype versus *Aifm2*-deficient Tregs with different concentrations of RSL3 for 4 hours. (E) *Aifm2* expression from bulk RNA-Seq of human PBMCs populations from Database of Immune Cell eQTLs, expression, epigenomics (DICE). (F) Differential expression analysis of *Aifm2* expression across different PBMC populations. For all plots, **P<0.05, **P<0.01, ***P<0.001 by one-way ANOVA (A-B), student t-test (C), or two-way ANOVA (D), mean ± s.e.m (A-C) or mean ± s.d (D).

Garcia Castillo, J. et al. Fig. 3.

Fig. 3 | Deletion of *Aifm2* in Tregs does not disrupt peripheral tolerance in mice.
(A) Frequency of total CD8$^+$ T cells (left), CD4$^+$ Tconv cells (middle), and Tregs(right) as a percentage of total CD45$^+$ immune cells in lymphoid tissues and organ tissues in 16 week mice. (B) Frequencies of activated (CD44^{hi}CD62^{lo}) CD8$^+$ T cells (left), CD4$^+$ Tconv cells (middle), and Tregs (right) in secondary lymphoid organs (SLOs) in 16 week mice. (C) gMFI of Foxp3, CD25, and NRP1 in Foxp3 Tregs across lymphoid tissues and non-lymphoid tissues in 16 week mice. (D) Schematic of pathway for conversion of ATP to Adenosine mediated by CD39 and CD73. (E) Representative histograms and gMFI of CD73 and CD39 in wildtype versus *Aifm2*-deficient Tregs across lymphoid tissues and nonlymphoid tissues in 16 week mice. Data from n=5 for Treg.*Aifm2*$^{+/+}$ and n=6 for Treg.*Aifm2*$^{\Delta/\Delta}$ mice for 16 week mice. For all plots, **P<0.05, **P<0.01, ***P<0.001 by two-way ANOVA (A, C, E) or student's t-test (B), mean ± s.e.m.

Garcia Castillo, J. et al. Fig. 4.

Fig. 4 | Deletion of *Aifm2* in Tregs improves cancer immunity. (A) Schematic for treating single cell suspensions from MC38 tumors +/- 1μM RSL3 (left). Frequencies of viable CD8$^+$ (left), CD4$^+$ Tconv (middle), and Tregs (right) from tumor single cell suspensions as a percentage of CD45$^+$ cells +/- 1μM RSL3 for 4 hours. Data from n=6 mice. (B) Experimental schematic for measuring tumor growth in wildtype versus mice with *Aifm2*-deficient Tregs. (C) Growth of MC38 tumors in wildtype versus mice with *Aifm2*-deficient Tregs. Data from n=5 for Treg.Aifm2$^{+/+}$ and n=6 from *Treg.Aifm2$^{\Delta/\Delta}$*. (D) Frequency of Foxp3$^+$ Tregs as a percentage of total CD45$^+$ immune cells (left) and CD8:Treg ratio (right) in MC38 tumors at D29. (E) Experimental schematic for measuring tumor growth in wildtype versus mice with *Aifm2*-deficient Tregs with or without 10mg/kg liproxstatin injected i.p daily. Data from n=3 for Treg.*Aifm2$^{+/+}$* + Vehicle, n=4 for Treg.*Aifm2$^{+/+}$* + Lip-1, n=5 for Treg.*Aifm2$^{\Delta/\Delta}$*+ Vehicle, and n=5 for Treg.*Aifm2$^{\Delta/\Delta}$* + Lip-1. (F) Growth of MC38 tumors in wildtype versus mice with *Aifm2*-deficient Tregs +/- Lip-1. For all plots, **P<0.05, **P<0.01, ***P<0.001 by student t-test (A, D) or two-way ANOVA (C, F), mean ± s.e.m.

Garcia Castillo, J. Fig. 5

Fig. 5 | Total T cell deficiency for *Aifm2* preserves CD8$^+$ T cell function and improves tumor immunity. (A) Experimental schematic for measuring tumor growth in wildtype versus mice with *Aifm2*-deficient T cells. (B) Growth of MC38 tumors in wildtype versus mice with *Aifm2*-deficient T cells. Data shown from n=5-6 mice/group from one of two experiments. (C) Frequency of Foxp3$^+$ Tregs as a percentage total CD45$^+$ immune cells in MC38 tumors at D29 from (B). (D) CD8:Treg ratio in MC38 tumors at D29 from (B). (E) Representative flow plot for PD-1 and LAG3 staining in total CD8$^+$ T cells (left) and quantification of the frequency of PD-1$^+$Lag3$^+$ CD8$^+$ T cells in MC38 tumors (right). (F) Frequency of tumor-specific (MuLV/H2K^{b+}) as a percentage of CD8$^+$ T cells (left) and frequency of PD-1$^+$Lag3$^+$ of MuLV/H2K^{b+} CD8$^+$ T cells in MC38 tumors (right). For all plots, **$P<0.05$, **$P<0.01$, ***$P<0.001$ by two-way ANOVA (B) or student t-test (C, D, F), mean ± s.e.m.