Inducible volatile chemical signalling drives antifungal activity of *Trichoderma hamatum* GD12

during confrontation with the pathogen *Sclerotinia sclerotiorum*

Gareth A. Thomas[1,2*], József Vuts[1], David M. Withall[1], John C. Caulfield[1], John Sidda[3], Murray R. Grant[3],

Christopher R. Thornton[2], Michael A. Birkett[1]

1. Protecting Crops and the Environment, Rothamsted Research, Harpenden, AL5 2JQ

2. Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, EX4 4QD

3. School of Life Sciences, University of Warwick, Coventry, CV4 7AL

*Corresponding author

Abstract

BACKGROUND: The use of beneficial soil fungi or their natural products offers a more sustainable

alternative to synthetic fungicides for pathogen management in crops. Volatile organic compounds

(VOCs) produced by such fungi act as semiochemicals that inhibit pathogens, with VOC production

influenced by physical interactions between competing fungi. This study explores the interaction

between the beneficial soil fungus *Trichoderma hamatum* GD12 strain (GD12), previously shown to

antagonize crop pathogens such as *Sclerotinia sclerotiorum*, to test the hypothesis that its antagonistic

effect is mediated by volatile chemical signalling. A GD12 mutant deficient in the chitinolytic enzyme

N-acetyl-β-glucosaminidase (*ΔThnag∷hph*), which shows reduced biocontrol activity, was also

examined**.**

RESULTS: In dual-culture confrontation assays, co-inoculation of GD12 and *S. sclerotiorum* led to

fungistatic interactions after 7 days, whereas *ΔThnag∷hph* showed no antagonism, indicating a loss

of antagonistic function. VOCs collected from individual and co-cultures were analysed by gas

chromatography – flame ionization detector (GC-FID) analysis and coupled GC-mass spectrometry

(GC-MS), revealing significant differences in VOC production between treatments, with VOC

production notably upregulated in the GD12 + *S. sclerotiorum* co-culture. Peak production of 6-pentyl-

2H-pyran-2-one occurred 17 days post-inoculation. This upregulation was absent in the *ΔThnag∷hph*

co-culture, suggesting VOCs may drive antagonism. Synthetic VOC assays revealed several compounds

inhibitory to *S. sclerotiorum*, including 1-octen-3-one, which also arrested the growth of key fungal

pathogens (*Botrytis cinerea*, *Pyrenopeziza brassicae*, and *Gaeumannomyces tritici*). Structural insights

into 1-octen-3-one's antifungal activity against *S. sclerotiorum* are also presented.

CONCLUSIONS: These findings support the hypothesis that the antagonistic properties of *T.*

hamatum GD12 against crop fungal pathogens can, in part, be attributed to VOC production. Further

research is needed to assess the potential of these semiochemicals as tools for pathogen

management in agriculture.

Keywords: *Trichoderma*, *Sclerotinia sclerotiorum*, volatile organic compounds, antagonism, 1-octen-

3-one

Introduction

Sclerotinia sclerotiorum (Lib.) de Bary (Family: Sclerotiniaceae) is a ubiquitous soil-borne fungal pathogen, affecting approximately 800 plant species worldwide, including economically important agricultural crops such as carrots, lettuce, sunflower, oilseed rape and potato (Boland *et al.*, 1994; Bolton *et al.*, 2006a). Management of *S. sclerotiorum* on agricultural crops relies mainly on the application of synthetic fungicides (Derbyshire and Denton-Giles, 2016), although the over-application of fungicides has increased selective pressure, leading to an increase in frequency of fungicide-resistant strains (Ma *et al.*, 2009). Alternative methods for controlling *S. sclerotiorum* include the use of crop rotations, which may also be ineffective due to the formation of vegetative sclerotia by *S. sclerotiorum,* which can remain viable in the soil for over eight years and are resistant to physical, chemical and biological degradation (Tribe, 1957; Adams, 1979; Bolton *et al.*, 2006a). Moreover, engineering crop resistance towards the pathogen has also proven challenging due to differing pathovars of the pathogen, and a lack of resistance in major crops, making breeding programmes a challenge (Bolton et al., 2006; Derbyshire et al., 2022). Therefore, more sustainable approaches for pathogen management on crops, that minimise reliance on synthetic fungicides, are needed.

Sustainable management strategies for the control of *S. sclerotiorum* include the exploitation of microbial biocontrol agents of the pathogen, including beneficial soil fungi. These decrease the negative potential of pathogens on crops through direct antagonism of the pathogen, competition for resources (e.g. nutrients), or through modification of plant defence responses (Ghorbanpour *et al.*, 2018). *Trichoderma* (Hypocreaceae) is a well-studied genus of beneficial soil fungi due to their ability to inhibit fungal pathogen development, induce plant defence responses against pathogens and promote plant growth (Druzhinina *et al.*, 2011; Woo *et al.*, 2023). GD12, a strain of *T. hamatum*, is effective at suppressing the growth of *S. sclerotiorum* in peat-based microcosms (Ryder *et al.*, 2012 Studholme *et al.*, 2013; Shaw *et al.*, 2016), with the suppressing capability of GD12 requiring the chitinase gene *N*-acetyl-β-glucosaminidase (Ryder et al., 2012). Genome sequencing of *Trichoderma hamatum* (Feng *et al.*, 2025) and specifically GD12 (Studholme et al., 2013) reveals the presence of silent gene clusters which could be activated in the presence of antagonistic microorganisms in soil, leading to the production of secondary metabolites which are not produced under standard laboratory conditions (Shaw et al., 2016). This antagonism can stimulate the induction of secondary metabolite biosynthetic gene clusters as evidenced by induction of genes encoding predicted polyketide synthases (PKSs) and Non-Ribosomal Peptide Synthetases (NRPSs) clusters during the interaction between *S. sclerotiorum* and *T. hamatum* in peat microcosms (Shaw *et al.*, 2016). However, the causal metabolites involved in these interactions are currently unknown.

Volatile organic compounds (VOCs) are a class of low molecular weight secondary metabolites produced by soil microorganisms, which contribute to their ability to compete against neighbouring organisms for resources in soil (Garbeva and Weisskopf, 2020; Weisskopf *et al.*, 2021). The ability of VOCs to travel between gas- and water-filled pockets in soil classifies them as long-distance messengers, compared to non-volatile secondary metabolites, which may be drivers of more local interactions (Kai *et al.*, 2016; Schulz-Bohm *et al.*, 2017; Westhoff *et al.*, 2017). Microbial VOCs are involved in a range of biological activities, including direct inhibition of pathogenic microorganisms (Fernando *et al.*, 2005), induced plant defence against pathogens (Ryu *et al.*, 2004) and plant growth promotion (Ryu *et al.*, 2003). Several studies indicate *Trichoderma* VOCs can specifically play an inhibitory role against a range of fungal pathogens (Amin *et al.*, 2010; Stoppacher *et al.*, 2010; Jeleń *et al.*, 2014a; Meena *et al.*, 2017; Wonglom *et al.*, 2020). These biological activities highlight the potential for microbial VOCs to be used as effective alternatives to pesticides and fertilisers (Thomas *et al.*, 2020, 2023).

Whilst beneficial soil microbes can produce VOCs when grown axenically under standard laboratory conditions, they exist in complex communities within the soil matrix. Genome sequencing of fungal species indicates that many secondary metabolite gene clusters are silent and telomeric under standard laboratory conditions and may require specific cultivation conditions to activate them, including stress inducing, or co-culturing with different species of microorganisms (Scherlach and Hertweck, 2021). Experimentally reproducing a more natural microbe interaction environment is feasible through inoculating different species of microorganisms within the same confined space. Such an approach may stimulate the antagonism that activate silents gene clusters, and hence facilitate the discovery of novel, bioactive compounds. It has been observed that ornesillic acid production was uniquely induced through co-culturing of *Streptomyces* and *Aspergillus* (Schroeckh *et al.*, 2009). This study subsequently led to an expansion in the discovery of novel secondary metabolite production through microbial co-culture (Knowles *et al.*, 2022). For example, in *T. harzianum,* co-culture with the endophyte *Talaromyces pinophilus* led to changes in secondary metabolite production relative to monoculture controls (Vinale *et al.*, 2017). The majority of these studies focus on changes in non-volatile compound production during physical interactions, however a growing body of evidence suggests co-culturing can also induce VOC production, with examples from fungal-fungal (Hynes *et al.*, 2007a; Evans *et al.*, 2008; El Ariebi *et al.*, 2016; Guo *et al.*, 2019a; O'Leary *et al.*, 2019), bacterial-bacterial (Tyc *et al.*, 2015, 2017) or fungal-bacterial (Albarracín Orio *et al.*, 2020) interactions.

Here, we aimed to determine the role of VOCs in the biocontrol capabilities of *T. hamatum* GD12 against *S. sclerotiorum*. We demonstrate; i) quantitative and qualitative changes in VOC production by *T. hamatum* occur during confrontation with *S. sclerotiorum*; ii) temporal changes in

VOC production during confrontation occur, with maximal induction day 17 post inoculation; iii) VOCs

produced by *T. hamatum* have antifungal activity against *S. sclerotiorum*; iv) identification of 1-octen-

3-one, which completely inhibits the growth of *S. sclerotiorum* as well as other agriculturally important

fungal pathogens; and iv) the structural features required for the antifungal activity of 1-octen-3-one

against *S. sclerotiorum*. This work highlights the power of using *Trichoderma*-pathogen co-culture to

reveal cryptic chemistries encoding bioactive VOCs for use as pathogen management tools in

agriculture.

Materials and methods

Dual-culture confrontation assays. Trichoderma hamatum GD12 isolated from a potato field (Great

Down, Devon, UK), (Thornton, pers. comm.), *ΔThnag::hph*::*hph* (Ryder et al., 2012) and *Sclerotinia*

sclerotiorum isolate 1 (isolated on oilseed rape petal, ADAS Rosemaund, Herefordshire, UK), (West,

pers. comm.), used in the study were maintained on Potato Dextrose Agar (PDA) (15 g Bacteriological

Agar No. 2, LabM, UK; 24 g Potato Dextrose Broth (PDB), Sigma, UK; 1000 mL distilled H_2O) slopes in

sterile, screw-capped plastic vials (ThermoScientific, UK). Circular plugs (5 mm diam.) were cut using

a sterilised cork-borer (Sigma, UK) from the leading edge of mycelia of 3-day old PDA plates of *T.*

hamatum (GD12, *ΔThnag::hph*) or *S. sclerotiorum* isolate 1. Each experiment comprised (a) a control,

containing uninoculated growth media (PDA), (b) the confrontation of a *T. hamatum* strain co-cultured

against itself (self-challenged), (c) the *S. sclerotiorum* co-cultured against itself (self-challenged), and

(d) the *T. hamatum* strain challenged against *S. sclerotiorum* (co-culture) (n=4). Plugs from individual

strains of *T. hamatum* were placed approximately 80 mm away from plugs of *S. sclerotiorum* on fresh

PDA plates (90 mm) and grown for 7 days under a 16 h/8 h fluorescent light/dark photoperiod at 24°C

until required for dynamic headspace collection experiments (detailed below).

Dynamic headspace collection (air entrainment). PDA plates containing 7-day-old fungal cultures (see

above) were enclosed individually in glass entrainment vessels (12 cm diam. x 6 cm height). Charcoal-

purified air (flowrate 600 mL/min) was pushed into each entrainment vessel and drawn (flowrate 500

mL/min), ensuring a positive pressure (100 mL/min) throughout the system. Air was drawn through a

glass tube containing Porapak Q (50 mg, 50/80 mesh, Supelco, Bellefonte, PA) held with two plugs of

silanized glass wool, for 20 h at ambient temperature (Pye volatile collection kits, Kings Walden, UK).

Before each collection, glass vessels were washed with Teepol detergent, rinsed with distilled water,

washed with acetone (ThermoFisher, UK), and then placed in a modified heating oven (180°C) for a

minimum of 2 h. Charcoal filters (10-14 mesh, 50 g) (Sigma, UK) were conditioned prior to each

experiment by attaching them to a supply of nitrogen in a modified heating oven (150°C) under a

constant stream of nitrogen. Volatile collections were performed under a 16 h/8 h light/dark photoperiod at 24°C. Porapak Q traps were cleaned by washing with freshly redistilled diethyl ether (2 mL) and heated to 132°C for a minimum of 2 h under a stream of nitrogen. Following collections, VOCs were eluted from the Porapak Q traps with freshly redistilled diethyl ether (750 µL) into 1.1 mL pointed vials (ThermoScientific, Germany), capped with an 8 mm Chromacol screw cap vial lid (ThermoScientific, Germany) with an 8 mm Silicone Red PTFE Septa (Kinesis, UK). The eluent was concentrated to 50 µL under a gentle stream of nitrogen and stored at -20°C prior to further analysis.

Time-course VOC collection experiment. Solid-phase microextraction (SPME) was selected as the method for VOC analysis for time-course experiments rather than dynamic headspace collection, as preliminary experiments showed repeated dynamic headspace collections of VOCs from fungal cultures led to drying out of growth media, impacting fungal growth. An SPME (100 µM Polydimethylsiloxane (PDMS) fibre, Supelco, UK) was introduced into the GC thermal desorption injector port to desorb for 10 min (temperature of injector=250°C). The SPME fibre was inserted through a clean septum and exposed to the headspace of the fungal culture within a clean glass entrainment vessel (12 cm diam. x 6 cm height) for 1 h. SPME samples were taken at 1, 2, 3, 4, 5, 6 ,7, 10, 17 and 24 days post inoculation (dpi) from cultures of (a) self-challenged *S. sclerotiorum* isolate 1, (b) self-challenged *T. hamatum* GD12, or (c) GD12 co-cultured with *S. sclerotiorum*. The first sampling timepoint (day 1 post inoculation) of the experiment was used as a baseline, with 80 mm of distance between the fungal mycelia. At day 2, the distance between the self-challenged GD12 treatments was 19-24 mm, and 7-15 mm for the GD12 co-cultured with *S. sclerotiorum* treatments. Self-challenged *S. sclerotiorum* treatments had already initiated contact by this stage of sampling. By day 3, contact between mycelia across all treatments had established.

Gas chromatography – flame ionization detector (GC-FID) analysis. Air entrainment samples were analysed on an Agilent 6890 GC equipped with a cool on-column injector, an FID and a HP-1 bonded-phase fused silica capillary column (50 m x 0.32 mm i. d. x 0.52 µm film thickness). The oven temperature was set at 30°C for 0.1 min, then increased at 5°C/min to 150°C for 0.1 min, then at 10°C/min to 230°C for a further 25 min. The carrier gas was hydrogen. VOCs adsorbed on SPME fibres were thermally desorbed by inserting the fibre directly into the OPTIC Programmable Temperature Vaporisor (PTV) unit (30 -> 250°C ballistically at a rate of 16°C/s).

Coupled GC-mass spectrometry (GC-MS). An Agilent Mass Selective Detector (MSD) 5973 coupled to an Agilent 6890N GC (fitted with a non-polar HP1 column 50 m length x 0.32 mm i. d. x 0.52 µM film thickness, J & W Scientific) was used for analysis. Sample injection was via cool-on column and MS ionization was by electron impact at 70 eV at 220°C. The GC oven temperature was maintained at 30°C

for 5 min and then programmed at 5°C/min to 250°C, run time 70 minutes. Tentative identifications

were confirmed by co-injections with authentic standards (Pickett et al., 1990).

Chemicals. 1-Pentanol (99%), 1-octen-3-ol (98%), 1-octene (98%), 2-undecanone (99%), 2-octanone

(98%), 3-octanone (99%), 2-pentylfuran (≥ 98%), 6-*n*-pentyl-2H-pyran-2-one (6-PAP) (≥ 96%), 1-octen-

3-one (96%) and 2-heptanone (98%) were all purchased from Sigma-Aldrich, UK. Diethyl ether (99.5%)

was purchased from Fisher and redistilled before use.

Synthetic compound assays. Synthetic standards of VOCs identified from air entrainments of *T.*

hamatum GD12 were applied to sterile qualitative filter paper (6 mm) (Whatman, UK) and placed onto

a Petri dish containing PDA. On a fresh plate of PDA, a plug (5 mm diam.) of *S. sclerotiorum* was

inoculated and the plate inverted over another plate containing the filter paper with the synthetic

VOC sample, and the two plates were then sealed with tape, ensuring no physical contact between

the VOC sample and the pathogen (Figure S1). Solutions of VOCs were prepared in freshly redistilled

diethyl ether to ensure that 20 µL application of solution gave a dose of 45.5 µM. This dose was

decided based on preliminary experiments, where 5 µL of each compound were applied neat to a

sterile filter paper, and the dose selected based on the least inhibitory compound (1-pentanol; 5 µL of

which equates to 45.5 µM). This dose is similar to that used in previous studies (Tyc *et al.*, 2015). For

compounds demonstrating significant ($p < 0.05$) antifungal activity relative to control treatments

when applied at 45.5 µM, doses onto filter discs were diluted to 22.75 µM, 11.125 µM, 4.55 µM, 2.275

µM, 0.91 µM and 0.455 µM in freshly redistilled diethyl ether from a stock solution, until no further

inhibition was observed. Mycelial measurements from *S. sclerotiorum* were taken using a 30 cm ruler

after 72 h. From the diameter of *S. sclerotiorum*, the area of colony was calculated as πr^2, where "r" is

equal to the radius of the colony. Control plates contained a sterile filter paper disc with 20 µL of

freshly redistilled diethyl ether alone.

Statistical analysis. For comparison of GC analysis from confrontation assays, peak area values were

individually measured (Agilent Chemstation) and $\log_{10}$-transformed. An adjustment of 0.001 was

applied to account for values recorded as zero. Statistical comparison of compounds present in both

mono- and co-culture treatments were analysed using an unpaired Student's t-test assuming equal

variances, one variate with grouped factor. To establish the antifungal activity of selected VOCs,

mycelial areas were statistically compared across treatments using one way analysis of variance

(ANOVA), followed by Tukey's Honest Significant Difference test at the $p < 0.05$ level, where multiple

comparisons were required. Genstat® (v21, ©VSN International, Hemel Hempstead, UK) was used for

all statistical analyses.

Results

Dual-culture confrontation assays demonstrated fungistatic interactions between *T. hamatum* GD12 when confronted with *S. sclerotiorum* after 7 days of growth (Figure 1). This was accompanied by the formation of yellow spores by *T. hamatum* GD12 in the interaction zone between the two fungi. Contrastingly, during confrontation with *T. hamatum ΔThnag::hph*, overgrowth of *S. sclerotiorum* was observed, depicted by a black arrow (Figure 1f), indicating a loss in the antagonistic capabilities of the *ΔThnag::hph* mutant.

Significant quantitative and qualitative changes in VOC production compared to self-challenged GD12 treatments were observed when *T. hamatum* GD12 was co-cultured with *S. sclerotiorum* (Figure 2; Table 1). Production of 6-pentyl-2H-pyran-2-one (6-PAP) dominated the headspace of GD12 co-cultured with *S. sclerotiorum*, with mean production of 6-PAP significantly greater in these co-cultures compared to self-challenged controls ($p < 0.001$) (Table 1). Of the 36 compounds detected, eight of which were confirmed by co-injection, 22 were unique to GD12-*S. sclerotiorum* co-cultures, suggesting that the VOCs were either biosynthesised *de novo* in the presence of *S. sclerotiorum* or produced below detectable limits of the GC in GD12 monocultures, including 2-pentylfuran (Figure 2; Table 1).

As co-culturing *T. hamatum* GD12 with *S. sclerotiorum* led to significant increases in VOC production, the temporal dynamics of *T. hamatum* GD12 VOC production was investigated. For three compounds detected by SPME, peak induction was greatest at day 17 post-co-culture, each subsequently decreasing by day 24 (Figure 3). Each compound showed similar trends in production across the different timepoints. In GD12 co-cultured with *S. sclerotiorum* treatments, compound 1 (KI = 1252) was not detected in the headspace until day 6. By day 10 there was an increase in production, which further increased at day 17, subsequently decreasing to day 24. 6-PAP shows a similar trend. Compound 2 (KI = 1994), detected by day 10, increasing until day 17, and then decreasing by day 24.

When co-cultured with *S. sclerotiorum*, quantitative and qualitative changes in VOC production relative to self-challenged *ΔThnag::hph* treatments were observed, although unlike with GD12, most VOCs were either down-regulated during co-cultivation, or not significantly different to self-challenged controls (Figure 4). This provides a negative control to directly link unique GD12 VOC production to biocontrol. Four of the 10 VOCs detected are up-regulated in self-challenged *ΔThnag::hph* treatments relative to *ΔThnag::hph* co-cultured with *S. sclerotiorum*, and five were not significantly different to controls (Table 2). Notably, 6-PAP, showing the greatest induction in GD12 co-cultured with *S. sclerotiorum* treatments, was present in *ΔThnag::hph* monocultures, but not in three out of four replicates in *ΔThnag::hph* co-cultured *S. sclerotiorum* treatments. Four VOCs (KI =

1552; KI = 2016; KI =2055; KI = 2264) were significantly greater in self-challenged *ΔThnag::hph* relative to co-culture treatments, and five were not significantly different. Taken together, these findings indicate a direct or indirect role for the *N*-acetyl-β-glucosaminidase enzyme in the induction of VOCs by *T. hamatum* GD12 during confrontation with *S. sclerotiorum*.

In preliminary experiments with *T. hamatum* VOCs and *S. sclerotiorum*, 6-PAP had no detectable inhibitory activity against *S. sclerotiorum* ($F_{2,6}$=2.17, p = 0.195) (Figure S2) and was therefore excluded from further bioassays. The mycelial area of *S. sclerotiorum* differed significantly depending on the specific VOC applied at the highest dose (45.5 µM), indicative of differences in their inhibitory activities ($F_{7,16}$= 171.36; p < 0.001; n= 3) (Figure 5). The mycelial area of *S. sclerotiorum* exposed to 1-pentanol was not significantly different to solvent control treatments (P > 0.05), and 2-heptanone treatments were not significantly different to 1-pentanol treatments (P > 0.05), so these compounds were not tested at reduced doses. 2-octanone and 2-undecanone demonstrated similar levels of inhibition, while 2-octanone was significantly more inhibitory than 3-octanone. Strikingly, 1-octen-3-one demonstrated effectively complete growth of inhibition of *S. sclerotiorum*. Importantly, this antifungal activity was effective against other economically fungal pathogens (*Botrytis cinerea*, *Pyrenopeziza brassicae* and *Gaeumannomyces tritici*) (Figure S3).

Of the tested VOCs, the five demonstrating the most significant inhibition were selected for further study at reduced doses, and all demonstrated significant inhibition when applied at reduced doses; 1-octen-3-one: ($F_{7,16}$= 246.03; p < 0.001); 2-octanone: ($F_{3,8}$= 42.29; p < 0.001); 3-octanone: ($F_{4,10}$= 31.74; p < 0.001); 2-pentylfuran: ($F_{5,12}$= 119.2; p < 0.001) and 2-undecanone: ($F_{7,16}$= 91.65; p < 0.001) (Figure 5). 2-octanone had a minimum inhibitory dose of 11.125 µM, 3-octanone, 2-pentylfuran had a minimum inhibitory dose of 4.55 µM and 2-undecanone, 2.275 µM (Figure 6). The compound showing the greatest inhibition, 1-octen-3-one, was still significantly inhibitory at a 100-fold dilution (0.445 µM).

To establish potential structural moieties required for antifungal activity of 1-octen-3-one against *S. sclerotiorum*, compounds with similar structural features to both 1-octene, 3-octanone and (*RS*)-1-octen-3-ol were tested (Figure S4), revealing significant differences in inhibitory activities ($F_{4,10}$= 114.44; p < 0.001) (Figure 7). 1-octene demonstrated no significant inhibition of *S. sclerotiorum* relative to solvent controls (p > 0.05), whereas both 3-octanone and (*RS*)-1-octen-3-ol showed significant inhibition of *S. sclerotiorum* relative to controls (p < 0.05). However, only 1-octen-3-one demonstrated 100% inhibition. When *S. sclerotiorum* was removed from the shared atmosphere with 1-octen-3-one, fungal growth of the pathogen was not restored 4 weeks after removing the pathogen

from the headspace (Figure 8). This is consistent with 1-octen-3-one having fungicidal activity at the
tested dose.

Discussion

In this study, we showed induction of VOC production by *T. hamatum* GD12 during co-culture
with the fungal pathogen *S. sclerotiorum*. This included VOCs not produced by *T. hamatum* when
grown axenically. This VOC induction was not observed during the interaction between *S. sclerotiorum*
and the *ΔThnag::hph* mutant, suggesting a role of the *N*-acteyl-β-glucosaminidase enzyme in the
direct or indirect facilitation of VOC induction. Whilst several VOCs were also produced in self-
challenged GD12 controls, the stimulation of VOC production in co-cultures indicates a *de novo*
biosynthesis in the presence of the pathogen, some of which we demonstrate possess an antifungal
role against *S. sclerotiorum*.

Many studies investigating VOC production from *Trichoderma* species utilise axenic fungal
growth, and VOCs are predominantly assigned to alcohols, ketones, alkanes, furans, mono- and-
sesquiterpenes (Jeleń *et al.*, 2014a). Several low molecular weight compounds reported here as being
produced by *T. hamatum* GD12 have been previously identified from other *Trichoderma* species,
including 3-octanone (Nemčovič *et al.*, 2008; Stoppacher *et al.*, 2010; Jeleń *et al.*, 2014a; Estrada-
Rivera *et al.*, 2019; Speckbacher *et al.*, 2020; Silva *et al.*, 2021) and 2-octanone (Jeleń *et al.*, 2014b;
Estrada-Rivera *et al.*, 2019; Speckbacher *et al.*, 2020). To our knowledge, this is the first report of 1-
octen-3-one being produced by *T. hamatum*, although it was identified from *T. virens* (Li *et al.*, 2018),
and a range of other fungi (Pennerman *et al.*, 2022). 6-PAP is a characteristic *Trichoderma* VOC
(Mendoza-Mendoza *et al.*, 2024), which produces a coconut aroma (Reithner *et al.*, 2007; Stoppacher
et al., 2010; Jeleń *et al.*, 2014a; Garnica-Vergara *et al.*, 2016; Estrada-Rivera *et al.*, 2019; Baazeem *et
al.*, 2021; Silva *et al.*, 2021). When comparing the *T. hamatum* GD12 VOCs identified via co-culture
with other studies, only 6-PAP has been previously reported (Jeleń *et al.*, 2014a; Baazeem *et al.*, 2021).
However, directly comparing volatile diversity across other studies should be undertaken with caution
as such studies employ different volatile sampling techniques, which can introduce biases for certain
compounds. For example, while many studies deploy SPME for headspace sampling, the diversity of
fungal volatiles recovered depends on the type of fibre used (Stoppacher et al., 2010; Jeleń et al.,
2014). Growth conditions of cultures will also likely vary across studies, which can influence volatile
production from *Trichoderma*, including age of cultures (Lee *et al.*, 2015), relative humidity and
temperature of growth conditions (Polizzi *et al.*, 2011), as well as media composition (Zhang *et al.*,
2014; González-Pérez *et al.*, 2018). Intraspecific differences in volatile production have been observed

for *T. hamatum*, as well as within other *Trichoderma* species. This likely relates to fungal evolution/adaptation to different geographical regions or ecological niches from which they were isolated (Jeleń et al., 2014; Lee et al., 2016), but also highlights the power of geographical/niche adaptation to drive the evolution of novel antifungals . Taken together, a range of factors can account for variation in VOC production by *Trichoderma*, both inter- and intra-specifically.

Co-cultivation with *S. sclerotiorum* revealed significant quantitative and qualitative changes in volatile production compared to the VOCs produced by *T. hamatum* GD12 in self-challenged controls. This induction was greatest 17 dpi, consistent with other studies (Hynes et al., 2007). 2-Pentylfuran, which was biosynthesised *de novo* in response to the interaction with *S. sclerotiorum*, has previously been reported from *T. hamatum* during axenic culture (Jeleń *et al.*, 2014a). Many of the upregulated compounds in our study are of the sesquiterpene-like class, which have previously been observed during physical fungal-fungal interactions (Hynes *et al.*, 2007a; Guo *et al.*, 2019a; Rajani *et al.*, 2021), including *T. hamatum* when challenged with the ectomycorrhizal fungus *Laccaria bicolor* (Guo *et al.*, 2019b) and *Trichoderma* in confrontation with *Sclerotium rolfsii* and *Macrophomina phaseolina* (Sridharan *et al.*, 2020). Sesquiterpenes are a well-known class of compounds involved in chemical signalling, and have been isolated across a range of *Trichoderma* species, including *T. hamatum* (Ma *et al.*, 2021), *T. brevicompactum* (Shi *et al.*, 2020), *T. virens* (Shi *et al.*, 2018, 2021; Hu *et al.*, 2019), *T. longibrachiatum* (Du *et al.*, 2020; Wang *et al.*, 2022), *T. asperellum* (Ding *et al.*, 2012), and *T. citrinoviride* (Liu *et al.*, 2020). Many of these sesquiterpene-like compounds possess antifungal activities against a range of phytopathogenic fungi, bacteria and marine phytoplankton. As well as their antimicrobial roles, microbial sesquiterpenes have a range of other biological activities including signalling, host growth promotion and defence (Avalos *et al.*, 2022). The upregulation of unknown sesquiterpenes during co-culture of *T. hamatum* and *S. sclerotiorum* could indicate a biological role for these compounds, and future work aims to isolate and identify these compounds to determine their role in the antagonistic response against *S. sclerotiorum*, and potential for integrating into biocontrol strategies.

6-PAP dominated the VOC profile of *T. hamatum* GD12 co-cultured with *S. sclerotiorum*, relative to self-challenged *T. hamatum* GD12 cultures, corroborating previous work which found significant increases in 6-PAP production when *T. harzianum* was co-inoculated with *R. solani* (Serrano-Carreón *et al.*, 2004; Flores *et al.*, 2019). Whilst significant increases in 6-PAP production by *T. hamatum* GD12 in co-culture were observed, no antifungal activity was observed when 6-PAP was applied in the inverted plate assay setup (Figure S2). However, several studies demonstrate an inhibitory role for 6-PAP against a range of fungal pathogens, including *Fusarium* species (Scarselletti and Faull, 1994; El-Hasan *et al.*, 2008; Jeleń *et al.*, 2014a; Rao *et al.*, 2022), *Botrytis cinerea* (Pezet *et*

al., 1999), *Cylindrocarpon destructans* (Jin *et al.*, 2020), and *Rhizoctonia solani* (Scarselletti and Faull,

1994), when the compound was in contact with the pathogens. These studies indicate that 6-PAP may

require direct contact for effective antifungal activity. As well as 6-PAP upregulation, 2-octanone

production was significantly upregulated in co-culture treatments, indicative of *T. hamatum - S.*

sclerotiorum antagonism. 2-octanone upregulation has also been observed during the interaction

between the fungal pathogen *Setophoma terrestris* and the beneficial soil bacteria *Bacillus subtilis*

(Albarracín Orio *et al.*, 2020), as well as the interaction between *T. atroviride* and *F. oxysporum*

(Speckbacher *et al.*, 2021), indicating a broader spectrum role for this compound in antagonistic fungal

interactions.

Many of the VOCs produced by *T. hamatum* GD12 during self-challenge and in co-cultures

with *S. sclerotiorum* show significant antifungal activity against *S. sclerotiorum*. The antifungal role of

2-octanone demonstrated here is in agreement with the inhibition of the soil fungal pathogen

Setaphoma terrestris, which could indicate broad-spectrum inhibitory activity against a range of fungal

pathogens (Albarracín Orio *et al.*, 2020). Similarly, 2-heptanone has previously shown significant

inhibition against *Curvularia lunata* (Xie *et al.*, 2020) and *Alternaria solani* (Zhang *et al.*, 2020). Here,

we report only moderate antifungal activity of 2-heptanone at the highest tested dose relative to

other compounds, which may relate to differences in doses tested across the studies, or specificity in

the antifungal activity of 2-heptanone against different pathogenic species. Alternatively, it may

reflect that anti-fungal role could be derived from 2-octanone via further modifications that many

occur in a more complex soil microbiome, as opposed to our two-component experimental system.

Specificity of antifungal activity has been observed for 2-undecanone, which has shown an inhibitory

role against *Verticillium dahlia*, *F. oxysporum*, *B. cinerea* and *Monilinia* spp., but not *Penicillium* spp.

(Calvo *et al.*, 2020) or *Rhizopus stolonifer* (Carter-House *et al.*, 2020). Having identified a range of

antifungal compounds, determining the modes of action of the antifungal activities against *S.*

sclerotiorum is an important next step. Furthermore, whilst the inhibitory properties of several *T.*

hamatum VOCs have been demonstrated against *S. sclerotiorum*, it is important to establish that

compounds at their inhibitory doses do not have phytotoxic effects. Interestingly, 2-pentylfuran,

which was upregulated in the presence of *S. sclerotiorum* and showed an antifungal role against the

pathogen, has also demonstrable plant growth-promoting capabilities (Zou *et al.*, 2010). Thus , 2-

pentylfuran could potentially be a promising candidate to replace synthetic chemical inputs due to its

ability to inhibit fungal pathogens without compromising plant growth.

To our knowledge, this is the first report demonstrating an antifungal role for 1-octen-3-one

against a pathogen. When structurally related compounds (1-octene, 3-octanone, (*R,S*)-1-octen-3-ol)

were tested for their antifungal activity at equivalent doses, 3-octanone and (*RS*)-1-octen-3-ol

demonstrated significant inhibition of *S. sclerotiorum*, whereas absolutely no growth of *S. sclerotiorum*

occurred when exposed to 1-octen-3-one. It is thus possible that fungicidal activity may be enhanced

via Michael-type acceptance by the α,β–unsaturated carbonyl structure within the latter compound

(Li *et al*., 2019). Several of the strobilurin class of fungicides (fungicides derived from *Strobilurus spp.*,

Nofiani *et al.*, 2018) also contain a conjugated ketone and alkene moiety. Findings here are contrary

to those reported by Xiong *et al*. (2017), who found significant inhibition of *F. tricinctum* and *F.*

oxysporum treated with 1-octen-3-ol, but no inhibition when fungi were treated with 1-octen-3-one

(Xiong *et al.*, 2017). However, VOCs were administered differently in each experiment, making cross-

comparison difficult. VOCs tested by Xiong et al. were supplemented into growth medium and in direct

contact with *Fusarium* species, whereas tested VOCs here were physically separated from *S.*

sclerotiorum, suggesting that the antifungal effect of 1-octen-3-one works at a distance. An important

consideration is that, in both studies, (*RS*)-1-octen-3-ol was tested as a racemic mixture, although

previous work has shown chirality can impact its antifungal activity (Yin *et al.*, 2019). Whilst 1-octen-

3-one shows an inhibitory role against *S. sclerotiorum*, future work should determine the role of the

compound on plant growth. For example, 1-octen-3-one exposure significantly inhibits *Arabidopsis*

growth and development (Lee *et al.*, 2019), therefore future work should focus on determining which

dose of 1-octen-3-one can inhibit *S. sclerotiorum* without compromising plant growth. Moreover, as

seen with antimicrobial drugs, a mixture of antifungals may have synergistic activities, yet overall

reduced potential phytotoxic effects, enabling the incorporation of 1-octen-3-one into biocontrol

strategies. As compounds identified here have *in vitro* antagonism against *S. sclerotiorum*, immediate

priorities will be to test these compounds against *S. sclerotiorum*, individually and in combinations,

using peat microcosms under glasshouse conditions. These data will inform future open field trials, to

examine their biological activities under more agriculturally relevant conditions.

Trichoderma spp. recognise plant pathogenic fungi when their lytic enzymes, including *N*-acetyl-*β*-D-

glucosaminidase, release small diffusible components from fungal cell walls (Druzhinina *et al.*, 2011).

These components can then bind G-Protein Coupled Receptors (GPCRs) on the cell surface of

Trichoderma, activating a downstream signalling cascade leading to the expression of secondary

metabolite biosynthesis genes, potentially including genes linked to volatile production. As

ΔThnag::hph cannot produce the chitinase enzymes required to break down fungal cell walls,

theoretically no breakdown products from *S. sclerotiorum* would bind to the GPCRs of *T. hamatum*,

preventing elicitation of the downstream signalling cascade and activation of secondary metabolite

biosynthesis genes. Significant reduction in 6-PAP production by the *ΔThnag::hph* mutant during co-

culture with *S. sclerotiorum* may also suggest a role for *N*-acetyl-*β*-D-glucosaminidase in 6-PAP

production, which several studies have previously demonstrated. For example, deletion of the *tga1*

gene, encoding the α-subunit of a heterochromatic G-protein 1 from *T. atroviride*, led to both a reduction in *N*-acetyl-*β*-D-glucosaminidase activity in mutant strains and an 8-fold reduction in 6-PAP production, relative to controls (Reithner *et al.*, 2005). Similarly, deletion of the gene encoding mitogen-activated protein kinase (*tmk1*) led to a 1.6-fold increase in production of 6-PAP and the enhancement of *nag1* expression relative to the parent strain (Reithner *et al.*, 2007). The reduction in 6-PAP production by *Δthnag::hph* may also explain why the mutant loses its antagonistic activity against *S. sclerotiorum* in peat microcosms (Ryder *et al.*, 2012).

In conclusion, this study suggests a role for volatile chemical signalling during the antagonistic response of *T. hamatum* GD12 against *S. sclerotiorum* and shows that certain *Trichoderma*-derived VOCs play an inhibitory role against the pathogen. Specifically, we identify1-octen-3-one as a potential novel antifungal VOC. Further glasshouse and field tests with antifungal compounds identified here are required to determine whether they inhibit pathogens at a larger scale under more agriculturally relevant conditions. Whilst several of these compounds have been identified here, or previously described, many *T. hamatum* compounds upregulated on confrontation with fungal plant pathogens remain to be identified and characterised.

Author contributions

Conceptualisation: MB, MG, CT. Data curation; GT, JV, JC, MB. Formal analysis; GT, JC, JV, MB, DW. Funding acquisition; MB, CT, MG. Investigation; MB, GT, MG, CT, JV, DW. Methodology; MB, GT, MG, CT, JS. Project administration; MB, MG, CT. Original draft; GT, MB, DW. Reviewing and editing; GT, MB, DW, JV, CT, MG, JS, JC.

Acknowledgements

Rothamsted Research receives strategic funding from the Biotechnology and Biological Sciences Research Council of the United Kingdom (BBSRC). We acknowledge support from the Growing Health Institute Strategic Programme (BB/X010953/1; BBS/E/RH/230003A). The work formed part of the Rothamsted Smart Crop Protection (SCP) strategic programme (BBS/OS/CP/000001) funded through BBSRC's Industrial Strategy Challenge Fund. GT's PhD studentship was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) Southwest Doctoral Training Programme award (project no. 1622285). The authors would also like to thank Jon West and Kevin King for providing *Sclerotinia*

sclerotiorum and *Botrytis cinerea*, Lisa Humbert for providing *Pyrenopeziza brassicae*, and Javier

Palma-Guerrero for providing *Gaeumannomyces tritici*.

References

Adams, P.B. (1979) Ecology of *Sclerotinia* Species . *Phytopathology* **69**: 896.

Albarracín Orio, A.G., Petras, D., Tobares, R.A., Aksenov, A.A., Wang, M., Juncosa, F., et al. (2020)
Fungal–bacterial interaction selects for quorum sensing mutants with increased production of
natural antifungal compounds. *Commun Biol*.

Amin, F., Razdan, V.K., Mohiddin, F.A., Bhat, K.A., and Sheikh, P.A. (2010) Effect of volatile
metabolites of *Trichoderma* species against seven fungal plant pathogens in-vitro. *Journal of*
Phytology **2**: 34–37.

Sridharan, A.P. Thankappan, S., G., K., and Uthandi, S. (2020) Comprehensive profiling of the VOCs of
Trichoderma longibrachiatum EF5 while interacting with *Sclerotium rolfsii* and *Macrophomina*
phaseolina. *Microbiol Res*.

El Ariebi, N., Hiscox, J., Scriven, S.A., Müller, C.T., and Boddy, L. (2016) Production and effects of
volatile organic compounds during interspecific interactions. *Fungal Ecol* **20**: 144–154.

Avalos, M., Garbeva, P., Vader, L., Van Wezel, G.P., Dickschat, J.S., and Ulanova, D. (2022)
Biosynthesis, evolution and ecology of microbial terpenoids. *Nat Prod Rep* **39**: 249–272.

Baazeem, A., Almanea, A., Manikandan, P., Alorabi, M., Vijayaraghavan, P., and Abdel-Hadi, A. (2021)
In Vitro Antibacterial, Antifungal, Nematocidal and Growth Promoting Activities of *Trichoderma*
hamatum FB10 and Its Secondary Metabolites. *Journal of Fungi 2021, Vol 7, Page 331* **7**: 331.

Boland, G.J., Hall, & R., and Hall, R. (1994) Index of plant hosts of *Sclerotinia sclerotiorum*. *CANADIAN*
JOURNAL OF PLANT PATHOLOGY.

Bolton, M.D., Thomma, B.P.H.J., and Nelson, B.D. (2006a) Sclerotinia sclerotiorum (Lib.) de Bary:
Biology and molecular traits of a cosmopolitan pathogen. *Mol Plant Pathol*.

Calvo, H., Mendiara, I., Arias, E., Gracia, A.P., Blanco, D., and Venturini, M.E. (2020) Antifungal
activity of the volatile organic compounds produced by *Bacillus velezensis* strains against
postharvest fungal pathogens. *Postharvest Biol Technol*.

Carter-House, D., Chung, J., McDonald, S., Mauck, K., and Stajich, J.E. (2020) Volatiles from *Serratia*
marcescens, *S. proteamaculans*, and *Bacillus subtilis* Inhibit Growth of *Rhizopus stolonifer* and
Other Fungi. *bioRxiv*.

Derbyshire, M.C. and Denton-Giles, M. (2016) The control of sclerotinia stem rot on oilseed rape
(*Brassica napus*): current practices and future opportunities. *Plant Pathol* **65**: 859–877.

Derbyshire, M.C., Newman, T.E., Khentry, Y., and Owolabi Taiwo, A. (2022) The evolutionary and
molecular features of the broad-host-range plant pathogen *Sclerotinia sclerotiorum*. *Mol Plant*
Pathol **23**:.

Ding, G., Chen, A.J., Lan, J., Zhang, H., Chen, X., Liu, X., and Zou, Z. (2012) Sesquiterpenes and
Cyclopeptides from the Endophytic Fungus *Trichoderma asperellum* Samuels, Lieckf. &
Nirenberg. *Chem Biodivers* **9**: 1205–1212.

Druzhinina, I.S., Seidl-Seiboth, V., Herrera-Estrella, A., Horwitz, B.A., Kenerley, C.M., Monte, E., et al.
(2011) *Trichoderma*: The genomics of opportunistic success. *Nat Rev Microbiol*.

Du, F.Y., Ju, G.L., Xiao, L., Zhou, Y.M., and Wu, X. (2020) Sesquiterpenes and Cyclodepsipeptides from
Marine-Derived Fungus *Trichoderma longibrachiatum* and Their Antagonistic Activities against
Soil-Borne Pathogens. *Marine Drugs 2020, Vol 18, Page 165* **18**: 165.

El-Hasan, A., Walker, F., and Buchenauer, H. (2008) *Trichoderma harzianum* and its metabolite 6-
pentyl-alpha-pyrone suppress fusaric acid produced by *Fusarium moniliforme*. *Journal of
Phytopathology* **156**: 79–87.

Estrada-Rivera, M., Rebolledo-Prudencio, O.G., Pérez-Robles, D.A., Rocha-Medina, M.D.C., González-
López, M.D.C., and Casas-Flores, S. (2019) *Trichoderma* histone deacetylase HDA-2 modulates
multiple responses in *Arabidopsis*. *Plant Physiol* **179**: 1343–1361.

Evans, J.A., Eyre, C.A., Rogers, H.J., Boddy, L., and Müller, C.T. (2008) Changes in volatile production
during interspecific interactions between four wood rotting fungi growing in artificial media.
Fungal Ecol.

Feng, Y., Shuai, X., Chen, J., Zhang, Q., Jia, L., Sun, L., et al. (2025) Unveiling the Genomic Features
and Biocontrol Potential of *Trichoderma hamatum* Against Root Rot Pathogens. *Journal of
Fungi* **11**: 126.

Fernando, W.G.D., Ramarathnam, R., Krishnamoorthy, A.S., and Savchuk, S.C. (2005) Identification
and use of potential bacterial organic antifungal volatiles in biocontrol. *Soil Biol Biochem* **37**:
955–964.

Flores, C., Nieto, M., Millán-Gómez, D. V., Caro, M., Galindo, E., and Serrano-Carreón, L. (2019)
Elicitation and biotransformation of 6-pentyl-α-pyrone in *Trichoderma atroviride* cultures.
Process Biochemistry **82**: 68–74.

Garbeva, P. and Weisskopf, L. (2020) Airborne medicine: bacterial volatiles and their influence on
plant health. *New Phytologist* **226**: 32–43.

Garnica-Vergara, A., Barrera-Ortiz, S., Muñoz-Parra, E., Raya-González, J., Méndez-Bravo, A., Macías-
Rodríguez, L., et al. (2016) The volatile 6-pentyl-2H-pyran-2-one from *Trichoderma atroviride*
regulates *Arabidopsis* thaliana root morphogenesis via auxin signaling and ETHYLENE
INSENSITIVE 2 functioning. *New Phytologist*.

Ghorbanpour, M., Omidvari, M., Abbaszadeh-Dahaji, P., Omidvar, R., and Kariman, K. (2018)
Mechanisms underlying the protective effects of beneficial fungi against plant diseases.
Biological Control **117**: 147–157.

González-Pérez, E., Ortega-Amaro, M.A., Salazar-Badillo, F.B., Bautista, E., Douterlungne, D., and
Jiménez-Bremont, J.F. (2018) The *Arabidopsis-Trichoderma* interaction reveals that the fungal
growth medium is an important factor in plant growth induction. *Scientific Reports 2018 8:1* **8**:
1–14.

Guo, Y., Ghirardo, A., Weber, B., Schnitzler, J.P., Philipp Benz, J., and Rosenkranz, M. (2019a) Trichoderma species differ in their volatile profiles and in antagonism toward ectomycorrhiza Laccaria bicolor. *Front Microbiol* **10**:.

Hu, Z., Tao, Y., Tao, X., Su, Q., Cai, J., Qin, C., et al. (2019) Sesquiterpenes with Phytopathogenic Fungi Inhibitory Activities from Fungus *Trichoderma virens* from *Litchi chinensis* Sonn. *J Agric Food Chem* **67**: 10646–10652.

Hynes, J., Müller, C.T., Jones, T.H., and Boddy, L. (2007a) Changes in volatile production during the course of fungal mycelial interactions between *Hypholoma fasciculare* and *Resinicium bicolor. J Chem Ecol*.

Jeleń, H., Błaszczyk, L., Chełkowski, J., Rogowicz, K., and Strakowska, J. (2014a) Formation of 6-n-pentyl-2H-pyran-2-one (6-PAP) and other volatiles by different Trichoderma species. *Mycol Prog*.

Jin, X., Guo, L., Jin, B., Zhu, S., Mei, X., Wu, J., et al. (2020) Inhibitory mechanism of 6-Pentyl-2H-pyran-2-one secreted by *Trichoderma atroviride* T2 against *Cylindrocarpon destructans*. *Pestic Biochem Physiol*.

Kai, M., Effmert, U., and Piechulla, B. (2016) Bacterial-plant-interactions: Approaches to unravel the biological function of bacterial volatiles in the rhizosphere. *Front Microbiol* **7**:.

Knowles, S.L., Raja, H.A., Roberts, C.D., and Oberlies, N.H. (2022) Fungal-fungal co-culture: a primer for generating chemical diversity. *Nat Prod Rep* **39**:.

Lee, S., Behringer, G., Hung, R., and Bennett, J. (2019) Effects of fungal volatile organic compounds on *Arabidopsis thaliana* growth and gene expression. *Fungal Ecol* **37**: 1–9.

Lee, S., Hung, R., Yap, M., and Bennett, J.W. (2015) Age matters: the effects of volatile organic compounds emitted by *Trichoderma atroviride* on plant growth. *Arch Microbiol* **197**: 723–727.

Lee, S., Yap, M., Behringer, G., Hung, R., and Bennett, J.W. (2016) Volatile organic compounds emitted by *Trichoderma* species mediate plant growth. *Fungal Biol Biotechnol* **3**: 7.

Li, N., Alfiky, A., Wang, W., Islam, M., Nourollahi, K., Liu, X., and Kang, S. (2018) Volatile Compound-Mediated Recognition and Inhibition Between *Trichoderma* Biocontrol Agents and Fusarium oxysporum. *Front Microbiol* **9**:.

Liu, X.H., Hou, X.L., Song, Y.P., Wang, B.G., and Ji, N.Y. (2020) Cyclonerane sesquiterpenes and an isocoumarin derivative from the marine-alga-endophytic fungus *Trichoderma citrinoviride* A-WH-20-3. *Fitoterapia* **141**: 104469.

Ma, H.X., Feng, X.J., Chen, Y., Chen, C.J., and Zhou, M.G. (2009) Occurrence and characterization of dimethachlon insensitivity in *Sclerotinia sclerotiorum* in Jiangsu Province of China. *Plant Dis* **93**: 36–42.

Ma, X.Y., Song, Y.P., Shi, Z.Z., and Ji, N.Y. (2021) Three sesquiterpenes from the marine-alga-epiphytic fungus *Trichoderma hamatum* Z36-7. *Phytochem Lett* **43**: 98–102.

Meena, M., Swapnil, P., Zehra, A., Dubey, M.K., and Upadhyay, R.S. (2017) Antagonistic assessment of *Trichoderma* spp. by producing volatile and non-volatile compounds against different fungal pathogens. *Archives of Phytopathology and Plant Protection* **50**: 629–648.

Mendoza-Mendoza, A., Esquivel-Naranjo, E.U., Soth, S., Whelan, H., Alizadeh, H., Echaide-Aquino, J.F., et al. (2024) Uncovering the multifaceted properties of 6-pentyl-alpha-pyrone for control of plant pathogens. *Front Plant Sci* **15**:.

Nemčovič, M., Jakubíková, L., Víden, I., and Farkaš, V. (2008) Induction of conidiation by endogenous volatile compounds in *Trichoderma* spp. *FEMS Microbiol Lett* **284**: 231–236.

Nofiani, R., de Mattos-Shipley, K., Lebe, K.E., Han, L.C., Iqbal, Z., Bailey, A.M., et al. (2018) Strobilurin biosynthesis in Basidiomycete fungi. *Nat Commun* **9**:.

O'Leary, J., Hiscox, J., Eastwood, D.C., Savoury, M., Langley, A., McDowell, S.W., et al. (2019) The whiff of decay: Linking volatile production and extracellular enzymes to outcomes of fungal interactions at different temperatures. *Fungal Ecol*.

Pennerman, K.K., Yin, G., and Bennett, J.W. (2022) Eight-carbon volatiles: prominent fungal and plant interaction compounds. *J Exp Bot* **73**: 487–497.

Pezet, R., Pont, V., and Tabacchi, R. (1999) Simple analysis of 6-pentyl-α-pyrone, a major antifungal metabolite of *Trichoderma* spp., useful for testing the antagonistic activity of these fungi. *Phytochemical Analysis* **10**:.

Polizzi, V., Adams, A., Picco, A.M., Adriaens, E., Lenoir, J., Van Peteghem, C., et al. (2011) Influence of environmental conditions on production of volatiles by *Trichoderma atroviride* in relation with the sick building syndrome. *Build Environ*.

Rajani, P., Rajasekaran, C., Vasanthakumari, M.M., Olsson, S.B., Ravikanth, G., and Uma Shaanker, R. (2021) Inhibition of plant pathogenic fungi by endophytic *Trichoderma* spp. through mycoparasitism and volatile organic compounds. *Microbiol Res*.

Rao, Y., Zeng, L., Jiang, H., Mei, L., and Wang, Y. (2022) *Trichoderma atroviride* LZ42 releases volatile organic compounds promoting plant growth and suppressing Fusarium wilt disease in tomato seedlings. *BMC Microbiology 2022 22:1* **22**: 1–12.

Reithner, B., Brunner, K., Schuhmacher, R., Peissl, I., Seidl, V., Krska, R., and Zeilinger, S. (2005) The G protein α subunit Tga1 of *Trichoderma atroviride* is involved in chitinase formation and differential production of antifungal metabolites. *Fungal Genetics and Biology* **42**: 749–760.

Reithner, B., Schuhmacher, R., Stoppacher, N., Pucher, M., Brunner, K., and Zeilinger, S. (2007) Signaling via the *Trichoderma atroviride* mitogen-activated protein kinase Tmk1 differentially affects mycoparasitism and plant protection. *Fungal Genetics and Biology*.

Ryder, L.S., Harris, B.D., Soanes, D.M., Kershaw, M.J., Talbot, N.J., and Thornton, C.R. (2012) Saprotrophic competitiveness and biocontrol fitness of a genetically modified strain of the plant-growth-promoting fungus *Trichoderma hamatum* GD12. *Microbiology*.

Ryu, C.M., Farag, M.A., Hu, C.H., Reddy, M.S., Kloepper, J.W., and Paré, P.W. (2004) Bacterial volatiles induce systemic resistance in *Arabidopsis*. *Plant Physiol*.

Ryu, C.M., Faragt, M.A., Hu, C.H., Reddy, M.S., Wei, H.X., Paré, P.W., and Kloepper, J.W. (2003) Bacterial volatiles promote growth in *Arabidopsis*. *Proc Natl Acad Sci U S A*.

Scarselletti, R. and Faull, J.L. (1994) In vitro activity of 6-pentyl-α-pyrone, a metabolite of *Trichoderma harzianum*, in the inhibition of *Rhizoctonia solani* and *Fusarium oxysporum* f. sp. lycopersici. *Mycol Res* **98**: 1207–1209.

Scherlach, K. and Hertweck, C. (2021) Mining and unearthing hidden biosynthetic potential. *Nat*
Commun.

Schroeckh, V., Scherlach, K., Nützmann, H.W., Shelest, E., Schmidt-Heck, W., Schuemann, J., et al.
(2009) Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in
Aspergillus nidulans. *Proc Natl Acad Sci U S A* **106**: 14558–14563.

Schulz-Bohm, K., Martín-Sánchez, L., and Garbeva, P. (2017) Microbial Volatiles: Small Molecules
with an Important Role in Intra- and Inter-Kingdom Interactions. *Front Microbiol* **8**: 1–10.

Serrano-Carreón, L., Flores, C., Rodríguez, B., and Galindo, E. (2004) *Rhizoctonia solani*, an elicitor of
6-pentyl-α-pyrone production by *Trichoderma harzianum* in a two liquid phases, extractive
fermentation system. *Biotechnol Lett*.

Shaw, S., Le Cocq, K., Paszkiewicz, K., Moore, K., Winsbury, R., De Torres Zabala, M., et al. (2016)
Transcriptional reprogramming underpins enhanced plant growth promotion by the biocontrol
fungus *Trichoderma hamatum* GD12 during antagonistic interactions with *Sclerotinia*
sclerotiorum in soil. *Mol Plant Pathol*.

Shi, X.-S., Song, Y.-P., Meng, L.-H., Yang, S.-Q., Wang, D.-J., Zhou, X.-W., et al. (2021) Isolation and
Characterization of Antibacterial Carotane Sesquiterpenes from *Artemisia argyi* Associated
Endophytic *Trichoderma virens* QA-8. *Antibiotics 2021, Vol 10, Page 213* **10**: 213.

Shi, Z.Z., Fang, S.T., Miao, F.P., Yin, X.L., and Ji, N.Y. (2018) Trichocarotins A–H and trichocadinin A,
nine sesquiterpenes from the marine-alga-epiphytic fungus *Trichoderma virens*. *Bioorg Chem*
81: 319–325.

Shi, Z.Z., Liu, X.H., Li, X.N., and Ji, N.Y. (2020) Antifungal and Antimicroalgal Trichothecene
Sesquiterpenes from the Marine Algicolous Fungus *Trichoderma brevicompactum* A-DL-9-2. *J*
Agric Food Chem **68**: 15440–15448.

Silva, L.R. da, Valadares-Inglis, M.C., Peixoto, G.H.S., Luccas, B.E.G. de, Muniz, P.H.P.C., Magalhães,
D.M., et al. (2021) Volatile organic compounds emitted by *Trichoderma azevedoi* promote the
growth of lettuce plants and delay the symptoms of white mould. *Biological Control* **152**:
104447.

Speckbacher, V., Ruzsanyi, V., Wigger, M., and Zeilinger, S. (2020) The *Trichoderma atroviride* Strains
P1 and IMI 206040 Differ in Their Light-Response and VOC Production. *Molecules 2020, Vol 25,*
Page 208 **25**: 208.

Speckbacher, V., Zeilinger, S., Zimmermann, S., Mayhew, C.A., Wiesenhofer, H., and Ruzsanyi, V.
(2021) Monitoring the volatile language of fungi using gas chromatography-ion mobility
spectrometry. *Anal Bioanal Chem*.

Stoppacher, N., Kluger, B., Zeilinger, S., Krska, R., and Schuhmacher, R. (2010) Identification and
profiling of volatile metabolites of the biocontrol fungus *Trichoderma atroviride* by HS-SPME-
GC-MS. *J Microbiol Methods*.

Studholme, D.J., Harris, B., Le Cocq, K., Winsbury, R., Perera, V., Ryder, L., et al. (2013) Investigating
the beneficial traits of *Trichoderma hamatum* GD12 for sustainable agriculture—insights from
genomics. *Front Plant Sci*.

Thomas, G., Rusman, Q., Morrison, W.R., Magalhães, D.M., Dowell, J.A., Ngumbi, E., et al. (2023)
Deciphering Plant-Insect-Microorganism Signals for Sustainable Crop Production. *Biomolecules*
13: 997.

Thomas, G., Withall, D., and Birkett, M. (2020) Harnessing microbial volatiles to replace pesticides
and fertilizers. *Microb Biotechnol* **13**: 1366–1376.

Tribe, H.T. (1957) On the parasitism of *Sclerotinia trifoliorum* by *Coniothyrium minitans*. *Transactions*
of the British Mycological Society **40**: 489–499.

Tyc, O., de Jager, V.C.L., van den Berg, M., Gerards, S., Janssens, T.K.S., Zaagman, N., et al. (2017)
Exploring bacterial interspecific interactions for discovery of novel antimicrobial compounds.
Microb Biotechnol **10**: 910–925.

Tyc, O., Zweers, H., de Boer, W., and Garbeva, P. (2015) Volatiles in inter-specific bacterial
interactions. *Front Microbiol* **6**:.

Vinale, F., Nicoletti, R., Borrelli, F., Mangoni, A., Parisi, O.A., Marra, R., et al. (2017) Co-Culture of
Plant Beneficial Microbes as Source of Bioactive Metabolites. *Sci Rep*.

Wang, Y., Li, X.M., Yang, S.Q., Zhang, F.Z., Wang, B.G., Li, H.L., and Meng, L.H. (2022) Sesquiterpene
and Sorbicillinoid Glycosides from the Endophytic Fungus *Trichoderma longibrachiatum* EN-586
Derived from the Marine Red Alga *Laurencia obtusa*. *Marine Drugs 2022, Vol 20, Page 177* **20**:
177.

Weisskopf, L., Schulz, S., and Garbeva, P. (2021) Microbial volatile organic compounds in intra-
kingdom and inter-kingdom interactions. *Nat Rev Microbiol* **19**: 391–404.

Westhoff, S., van Wezel, G.P., and Rozen, D.E. (2017) Distance-dependent danger responses in
bacteria. *Curr Opin Microbiol*.

Wonglom, P., Ito, S. ichi, and Sunpapao, A. (2020) Volatile organic compounds emitted from
endophytic fungus *Trichoderma asperellum* T1 mediate antifungal activity, defense response
and promote plant growth in lettuce (*Lactuca sativa*). *Fungal Ecol* **43**:.

Woo, S.L., Hermosa, R., Lorito, M., and Monte, E. (2023) *Trichoderma*: a multipurpose, plant-
beneficial microorganism for eco-sustainable agriculture. *Nat Rev Microbiol* **21**:.

Xie, S., Liu, J., Gu, S., Chen, X., Jiang, H., and Ding, T. (2020) Antifungal activity of volatile compounds
produced by endophytic *Bacillus subtilis* DZSY21 against Curvularia lunata. *Ann Microbiol*.

Xiong, C., Li, Q., Li, S., Chen, C., Chen, Z., and Huang, W. (2017) *In vitro* antimicrobial activities and
mechanism of 1-octen-3-ol against food-related bacteria and pathogenic fungi. *J Oleo Sci* **66**:
1041–1049.

Yin, G., Zhang, Y., Fu, M., Hua, S.S.T., Huang, Q., Pennerman, K.K., et al. (2019) Influence of R and S
enantiomers of 1-octen-3-ol on gene expression of *Penicillium chrysogenum*. *J Ind Microbiol*
Biotechnol **46**: 977–991.

Zhang, D., Yu, S., Yang, Y., Zhang, J., Zhao, D., Pan, Y., et al. (2020) Antifungal Effects of Volatiles
Produced by *Bacillus subtilis* Against *Alternaria solani* in Potato. *Front Microbiol*.

Zhang, F., Yang, X., Ran, W., and Shen, Q. (2014*) Fusarium oxysporum* induces the production of
proteins and volatile organic compounds by *Trichoderma harzianum* T-E5. *FEMS Microbiol Lett*
359: 116–123.

Zou, C., Li, Z., and Yu, D. (2010) *Bacillus megaterium* strain XTBG34 promotes plant growth by
producing 2-pentylfuran. *Journal of Microbiology* **48**: 460–466.

Figures.

Figure 1 | Dual-culture confrontation assays of (a) self-challenged *Trichoderma hamatum* GD12 strain;

(b) self-challenged *Sclerotinia sclerotiorum* (c) co-culture of *S. sclerotiorum* (top) and *T. hamatum*

GD12 (bottom); (d) self-challenged *T. hamatum* N-acetyl-β-glucosaminidase (*ΔThnag∷hph*) mutant;

(e) self-challenged *S. sclerotiorum*; (f) co-culture of *S sclerotiorum* (top) and *T. hamatum ΔThnag∷hph*

mutant (bottom).

Figure 2 | Representative gas chromatographic analysis of volatile organic compounds (VOCs)
collected by air entrainment from 7-day old cultures of (a) uninoculated growth media (control), (b)
self-challenged *S. sclerotiorum*, (c) self-challenged *T. hamatum* GD12, and (d) *T. hamatum* GD12 co-
inoculated with *S. sclerotiorum*. For an explanation of peak numbers, see Table 1.

Figure 3 | Production of VOCs in treatments of *Trichoderma hamatum GD12* and *S. sclerotiorurm* co-
culture treatments over the course of 24 days for three VOCs. Bars represent the peak area value of
each VOC (± SD) (n=3).

711   Figure 4 | Representative gas chromatographic (on an HP-1 column) analysis of VOCs from dynamic

712   headspace collections of 7-day old cultures of (a) uninoculated growth media (control), (b) self-

713   challenged *S. sclerotiorum*, (c) self-challenged *T. hamatum ΔThnag∷hph*, and (d) *T. hamatum*

714   *ΔThnag∷hph* co-inoculated with *S. sclerotiorum*.

Figure 5 | Antifungal activities of selected VOCs on the growth of *S. sclerotiorum*. *S. sclerotiorum* was incubated with selected VOCs at 45.5 μM doses and the inhibition rates were calculated relative to control plates (exposed to diethyl ether alone) after 3 days. Bars represent the mean mycelial area of *S. sclerotiorum* upon exposure to each VOC (± SD) (n=3). Different letters indicate significant differences between treatments according to Tukey's multiple comparisons test (p < 0.05). Y axis represents mycelial area (mm^2).

Figure 6 | Antifungal activities of selected VOCs on the growth of *S. sclerotiorum*, at reduced doses. Bars represent the mean mycelial area of *S. sclerotiorum* upon exposure to each VOC (± SD) (n=3). Different letters indicate significant differences between treatments according to Tukey's multiple comparisons test ($p < 0.05$). Y axis represents mycelial area (mm^2).

Figure 7 | Antifungal activities of selected VOCs on the growth of *S. sclerotiorum*, representing individual structural components of 1-octen-3-one, at 45.5 µM. Bars represent the mycelial area of *S. sclerotiorum* upon exposure to each VOC (± SD) (n=3). Different letters indicate significant differences between treatments according to Tukey's multiple comparisons test (p < 0.05).

Figure 8 | Antifungal activities of 6-PAP and 1-octen-3-one, on *Sclerotinia sclerotiorum* at 45.5 µM.

783 Table 1 | Composition of VOCs from dynamic headspace collections of 7-day old cultures of self-

784 challenged *T. hamatum* GD12, or GD12 co-inoculated with *S. sclerotiorum* (n=4) (mean peak area ±

785 SE). Data were analysed by Student's t-test (p < 0.05). KIs on a non-polar HP-1 GC column.

Peak no.	Compound	KI	GD12 vs S. sclerotiorum	GD12 vs GD12	p value
1	1-Pentanol*	754	7.29 (± 1.85)	2.42(±0.37)	0.016
2	2(5H)-Furanone	871	95.43 (± 24.50)	n.d.	NA
3	2-Heptanone*	876	26.08(±4.82)	26.47(±7.6)	0.813
4	1-Octen-3-one*	959	8.76(±3.92)	11.49(±9.23)	0.606
5	3-Octanone*	969	50.98(±17.87)	41.25(±22.83)	0.439
6	2-Octanone*	972	29.00(±6.52)	n.d.	NA
7	2-Pentylfuran*	981	142.00(± 48.95)	n.d.	NA
8	No i.d.	990	4.65(±0.44)	n.d.	NA
9	No i.d.	995	6.90(±1.81)	n.d.	NA
10	No i.d.	1049	30.67(±13.89)	n.d.	NA
11	No i.d.	1074	8.98(±1.59)	n.d.	NA
12	No i.d.	1137	25.29(±12.24)	20.34(±13.59)	0.931
13	2-n-Heptylfuran	1182	12.60(±3.31)	7.61(±4.01)	0.278
14	Cyclodecanone	1217	28.09(±9.8)	n.d.	NA
15	2-Butyl-cyclodecanone	1235	71.50(±29.64)	n.d.	NA
16	No i.d.	1252	53.66(±34.92)	n.d.	NA
17	2-Undecanone*	1276	9.00(±1.54)	n.d.	NA
18	6-Pentyl-2H-pyran-2-one*	1429	7596.41(±1617.76)	58.04(±20.08)	0.001
19	No i.d.	1457	27.35(±5.69)	n.d.	NA
20	No i.d.	1488	90.47(±24.10)	n.d.	NA
21	No i.d.	1497	14.21(±4.25)	n.d.	NA
22	No i.d.	1505	8.65(±1.61)	n.d.	NA
23	No i.d.	1518	8.12(±2.04)	n.d.	NA
24	No i.d.	1565	12.03(±5.38)	n.d.	NA
25	No i.d.	1608	6.67(±0.54)	n.d.	NA
26	No i.d.	1625	11.33(±2.90)	n.d.	NA
27	No i.d.	1632	14.44(±3.66)	n.d.	NA
28	No i.d.	1639	12.89(±3.51)	n.d.	NA
29	No i.d.	1710	334.61(±103.98)	5.19(±3.42)	0.034
30	No i.d.	1739	98.07(±42.20)	n.d.	NA
31	No i.d.	2018	68.41(±14.13)	7.67(±2.77)	0.002
32	No i.d.	2057	9.81(±2.14)	1.45(±0.67)	0.003
33	No i.d.	2198	2.40(±0.77)	0.65(±0.39)	0.105
34	No i.d.	2208	8.00(±1.56)	1.60(±0.69)	0.006
35	No i.d.	2240	11.00(±3.56)	1.55(±0.78)	0.043
36	No i.d.	2266	43.73(±8.74)	2.85(±0.72)	0.001

807 *n.d. not detected*

809 Table 2 | Composition of VOCs collected from dynamic headspace collections of 7-day old cultures of

810 self-challenged *T. hamatum ΔThnag::hph*, or *ΔThnag::hph* co-inoculated with *S. sclerotiorum* (n=4)

811 (mean peak area ± SE). Data were analysed by Student's t-test (p < 0.05). KIs on a non-polar HP-1 GC

812 column.

Peak no.	KI	Compound	Mean (± SEM)		p value
			nag vs *S. sclerotiorum*	nag vs nag	
1	1425	6-Pentyl-2H-pyran-2-one*	n.d.	52.06 (±17.40)	NA
2	1552	No i.d.	61.89(±21.1)	**793(±213.9)**	0.049
3	1624	No i.d.	189.32(±69.91)	526.03(±145.52)	0.278
4	1692	No i.d.	**1.9(±0.78)**	0	
5	1735	No i.d.	26.14(±9.06)	161.8(±53.61)	0.118
6	2016	No i.d.	51.09(±2.80)	**322.7(±66.90)**	0.001
7	2055	No i.d.	3.37(±1.20)	**80.3(±16.74)**	0.003
8	2208	No i.d.	74.87(±28.26)	336.83(±78.090	0.15
9	2243	No i.d.	76.49(±28.95)	367(±90.19)	0.145
10	2264	No i.d.	22.31(±2.91)	**271.00(±67.83)**	0.005