**Empathy for pain persists across live two-way video interactions and viewing of**

**prerecorded videos**

Jannik Heimann[1], Pauline Petereit[2], Anat Perry[3], Ulrike M. Krämer[1*]

[1] Institute of Medical Psychology, Center of Brain, Behavior and Metabolism, University of Lübeck,

Ratzeburger Allee 160, 23562 Lübeck, Germany

[2] Department of Neurology, University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

[3] Psychology Department, Hebrew University of Jerusalem, Mount Scopus, 91905 Jerusalem, Israel

* Corresponding author: University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany.

Email: ulrike.kraemer@uni-luebeck.de

Phone: 0049 451 3101 7421

15                                                          **Abstract**

The impact of digitally mediated social interaction on understanding others and sharing their emotions

has not been thoroughly investigated. We examined how live, video-mediated interaction, as opposed

to watching a prerecorded video, affects behavioral, neural, and physiological aspects of empathy for

pain.  Thirty-five  observers  watched  targets  undergoing  painful  electric  stimulation  in  an

electroencephalogram study. We hypothesized that reduced temporal presence, the immediacy or

delay with which information is transferred during social interactions, would result in diminished

behavioral and electrophysiological empathic responses. However, observer's behavioral empathic

responses were not diminished with reduced temporal presence. On a neural level, mid-frontal theta

was sensitive to the other's pain intensity, and we observed significant physiological coupling between

participants. Mu suppression, on the other hand, was not modulated by pain intensity. Importantly,

neural and physiological indices of empathy were independent of temporal presence. However,

exploratory analyses indicated a latency effect of temporal presence on pain-related theta activity with

an earlier theta increase in interactions with high temporal presence. The results suggest that the

temporal presence of individuals may not be necessary for empathy towards another's pain. Future

studies may investigate more naturalistic social interactions and include motivational aspects of

empathy. We discuss implications of these findings for debates on social presence and on second-

person neuroscience.

Key words: electroencephalography; empathic accuracy; neural oscillations; physiological coupling;

social cognition

Digitally mediated social interactions have become an integral part of everyday life, from social

media to video conferencing to online psychotherapy. However, the implications of mediated social

interactions for our understanding of others remain poorly understood. We recently proposed a

theoretical framework for studying the effects of social presence on social cognition, in particular on

empathy and mental state attribution (Krämer et al., 2025). In this framework, we defined social

presence as the degree to which others are physically and temporally present, that is the degree to

which verbal and nonverbal cues about others are available (physical presence) and the immediacy

with which these social cues are available (temporal presence). Extending a recent study of ours, in

which we investigated the influence of physical presence (Petereit et al., 2023), we here addressed

how temporal presence affects empathy for pain on behavioral, neural and physiological levels.

Temporal presence refers to the immediacy or delay with which information about the other is

transferred during social interactions, which in turn allows for the possibility of reciprocity. For example,

a direct conversation between two individuals demonstrates high temporal presence due to the

instantaneous exchange of information, while observing a prerecorded video reflects low temporal

presence because of the temporal delay between recording and viewing (Krämer et al., 2025).

Temporal alignment in social situations can be assumed to play an essential role in facilitating social

coordination and communication (Cañigueral et al., 2022; De Jaegher et al., 2010; Gallotti et al., 2017;

Schilbach et al., 2013). Only with high temporal presence, interacting individuals can directly react to

the other's behavior creating feedback-loops of reciprocal social exchange (Dumas, 2011; Hamilton &

Holler, 2023). Compared to live, bidirectional interactions, watching a video of another person reduces

the temporal presence of that person, which could also impair empathy due to the absence of reciprocal

social cue exchange. We understand empathy as an emotional response that involves affective sharing

of another's feelings and cognitive understanding of their mental and emotional states, while

maintaining a distinction between one's own and the other's states (Cuff et al., 2016; Zaki & Ochsner,

2012).

The effects of temporal presence in social interactions have been investigated especially in

regard to eye gaze as a social communication signal. The social significance and the signaling function

of direct gaze are impaired when temporal presence between interacting individuals is prevented

(Hamilton, 2016; Hamilton & Lind, 2016; Risko et al., 2016). When comparing direct, real-life interaction,

video-mediated interaction, and watching a prerecorded video, the skin conductance response to direct

gaze of others differed only between prerecorded videos and direct interaction, suggesting that the

direct gaze in the condition of preserved temporal presence was the critical factor for increased

autonomic arousal (Hietanen et al., 2020). Moreover, patterns of gaze behavior were altered when

comparing direct, live interactions and prerecorded videos, indicating differences in implicit mentalizing

between direct and mediated interaction (Morgan et al., 2022). Research on action observation in

infants, adults, and in macaque monkeys suggests stronger neural motor responses in live, direct action

observation compared to watching video-taped actions (Caggiano et al., 2011; Järveläinen et al., 2001;

Karimova et al., 2024; Shimada & Hiraki, 2006). A recent study comparing live and video-taped theater

performances reported more skin conductance responses when observing live performances,

suggesting increased physiological arousal with higher temporal presence (Gabriel et al., 2025). In a

further study, mimicry and the development of trust during interactions remained consistent across

direct, face-to-face interactions and live, video calls, whereas these features were found to be

diminished in prerecorded video interactions (Diana et al., 2023), which may indicate reduced

experience sharing with reduced temporal presence. These studies showed that temporal presence

affects important components of empathy and associated constructs, including implicit mentalizing,

mimicry, arousal in social situations and action sharing. Based on this, we hypothesized that temporal

presence also impacts more complex facets of empathy, which has not been shown before. Thus, the

current study investigated how reduced temporal presence affects our empathic abilities to share

experiences and understand others in pain on behavioral, neural and physiological levels (Cuff et al.,

2016; Decety & Jackson, 2004; Lieberman, 2007; Shamay-Tsoory, 2011; Zaki & Ochsner, 2012).

As in our previous study on physical presence and pain empathy (Petereit et al., 2023), we

used an established empathy-for-pain paradigm (Lamm et al., 2011; Rütgen et al., 2015; Singer et al.,

2004; Singer & Lamm, 2009).  In this paradigm, the behavioral readout, also called "empathic accuracy,"

is defined as the relationship between the ratings of the pain observer and the pain receiver (the target)

(Gauthier et al., 2008; Ickes et al., 1990; Jospe et al., 2020; Zaki & Ochsner, 2011). Empathic accuracy

is considered to reflect the cognitive dimension of empathy (Mackes et al., 2018; Rütgen et al., 2015),

whereas the observer's subjective experience of unpleasantness in response to the target's pain is

thought to reflect the affective component of pain empathy (Mu et al., 2008; Rütgen et al., 2015).

The neural correlates of empathy for pain are often described within the framework of shared

representations and simulation theories of empathy. These theories posit that observers represent the

targets' experience in a similar manner and in corresponding brain areas (Decety & Jackson, 2004;

Keysers & Gazzola, 2006; Lamm et al., 2011; Marsh, 2018; Preston & De Waal, 2002), which are

sometimes referred to as mirror neuron system (Rizzolatti & Craighero, 2004). Shared representations

are assumed to facilitate social understanding, and  this mechanism has been proposed as a neural

substrate for experience sharing and affective components of empathy (Bekkali et al., 2021; Gallese et

al., 2004; Keysers & Gazzola, 2006; Singer & Lamm, 2009), including empathy for pain (Hari & Kujala,

2009; Riecansky & Lamm, 2019). In the current study, reduced temporal presence disrupts the

exchange of social cues and contingent responding, or mutual adjustment. This disruption may weaken

engagement of neural circuits related to simulations and the subsequent empathic processes.

Accordingly, we focused on specific neural and physiological indices of shared representations and

pain empathy, namely the mu-rhythm, frontal theta and "physiological coupling". The so-called mu

rhythm (8-13 Hz) stems from the somatosensory cortex and has been repeatedly associated with pain

empathy, with mu power being suppressed in response to the pain of others  (Cheng et al., 2008; Peled-

Avron et al., 2018; Peng et al., 2021; Perry et al., 2010; Riecansky & Lamm, 2019; Whitmarsh et al.,

2011; Yang et al., 2009). Frontal theta activity (2-8 Hz; Misra et al., 2017; Mu et al., 2008; Peng et al.,

2021; Petereit et al., 2023) possibly stems from the anterior and mid-cingulate cortex (ACC/MCC, Misra

et al., 2017; Mitchell et al., 2008; Van Der Molen et al., 2017; Zhang et al., 2021) which is implicated in

both first-hand and vicarious pain as part of the "pain matrix" (Cheng et al., 2007; Fallon et al., 2020;

Hutchison et al., 1999; Jackson et al., 2005; Lamm et al., 2011; Xiao & Zhang, 2018), the core network

for nociceptive processing (Legrain et al., 2011). This is evidenced by its response to images of hands

in painful situations (Mu et al., 2008) and to facial expressions of varying pain intensities (Saarela et al.,

2006). Images of hands and faces in pain have both been shown to be associated with behavioral and

neural aspects of empathy for pain (Gallo et al., 2018; González-Roldan et al., 2011; Goubert et al.,

2005; Jauniaux et al., 2019; Murata et al., 2020; Saarela et al., 2006). Emotional facial expressions play

a major role in social signaling (Cañigueral et al., 2022; Güntekin & Başar, 2014). Reduced temporal

presence may hamper this reciprocal signaling by limiting the ability to express compassion or support.

In mediated interactions, the salience of another's suffering can diminish due to missing reciprocity and

reduced self-relevance of the other's experience (Conway et al., 2016; Schmitz & Johnson, 2007), as

meaningful reactions are constrained (Winczewski et al., 2016; Zaki, 2014). This decreased self-

relevance may lower engagement with the other's pain, leading us to expect reduced neural responses

under reduced temporal presence.

Another physiological measure of (pain) empathy is "physiological coupling", which refers to

the phenomenon of synchronized autonomous activity patterns, such as those observed in

electrodermal (EDA; skin conductance response, SCR) or cardiac activity (interbeat interval, IBI) across

interacting humans (Neumann & Westbury, 2011; Palumbo et al., 2017). Several studies have

demonstrated the relevance of physiological coupling in contexts of empathy (Jospe et al., 2020;

Levenson & Gottman, 1983; Pan et al., 2023; Qaiser et al., 2023), as well as in the context of pain

perception (Goldstein et al., 2017; Murata et al., 2020; Reddan et al., 2020), self-reported feelings of

social presence (Järvelä et al., 2016), and the effects of physical presence on empathy for pain (Petereit

et al., 2023). Investigating the relationship between physiological coupling and empathic accuracy has

provided mixed results (Jospe et al., 2020; Levenson & Ruef, 1992; Thorson, 2018; Zerwas et al.,

2021).

In our previous study, we investigated the effects of ***physical presence*** on empathy for pain

using the above-mentioned pain empathy paradigm (Petereit et al., 2023). Participants observed

another person undergoing painful electric stimulation, either in a direct, face-to-face or in a live, video-

mediated interaction. Physical presence did not affect empathic accuracy and shared unpleasantness,

and neither did it affect pain-related mu suppression. However, observer's theta power to the other's

pain and the physiological coupling between participants were diminished in mediated compared to

direct interactions. These results suggest that physical presence, i.e., the richness of social cues in an

interaction, is important for a nuanced physiological resonance with the other's pain experience.

Building on this work, we examined how **temporal presence** impacts pain empathy by

comparing live video calls with watching prerecorded videos of others in pain. Since reductions in

temporal presence disrupt the immediate exchange of information and reciprocity, we hypothesized

that temporal presence would affect empathy-related neurophysiological mechanisms in a similar way

to, or even more than, physical presence (Petereit et al., 2023). In detail, we hypothesized that reduced

temporal presence would result in diminished empathic responses. Specifically, at the behavioral level,

we expected reduced temporal presence to result in decreased empathic accuracy and less shared

unpleasantness of other's pain. At the physiological level, we hypothesized that heartbeat and skin

conductance coupling would decrease with reduced temporal presence. Finally, at the neural level, we

expected mu suppression and midfrontal theta activity to vary with levels of temporal presence.

157                                                      **Methods**

**Participants**

Thirty-five students from the University of Lübeck were recruited (28 females, 7 males, self-

identified; mean age: 24 years ($SD$ 5.7)). All participants served as target and observer. Before the

experiment, we made sure that the interacting participants were unfamiliar with each other. We used

the same power analysis as Petereit et al. (2023) since one can expect that the effect sizes are

comparable based on the similar experimental conditions. The power analysis showed that the sample

size of $N=30$ is sufficient to detect a small effect ($f^2=0.02$) of condition on empathic accuracy in a

generalized linear mixed model on single trials with a power of at least 0.98. The power calculation was

done in R using data simulation in the *simr* package (Green & MacLeod, 2016). Exclusion criteria

included current and past neurological, current psychiatric or cardiovascular disorders, current or

chronic pain disorders or current pain medication intake.

Due to the experimental design, the very first participant only served as target and the very last

participant only served as observer (see Experimental Procedure for further explanation). One

participant had to be excluded because of a non-reported sleeping disorder and could neither serve as

observer nor as target. One participant's behavioral data were lost due to technical issues. Some

participants had to be excluded because of bad data quality and were excluded for each data analysis

separately (EEG: 1, EDA: 1, own pain: 4 participants excluded). This led to the sample sizes of $N=31$

for the behavioral data, $N=31$ for the neural data in the observing conditions, $N=31$ for the SCR, $N=32$

for the ECG data, and $N=30$ for the neural data in the "own pain" condition. Participants received course

credit or €12 per hour. All participants provided written informed consent before taking part in the

experiment. The experiment was approved by the Ethics Committee of the University of Lübeck and

conducted according to the Declaration of Helsinki.

**Experimental design**

In prior social presence research, the terms "intimacy" and "immediacy" have been used for

physical and temporal presence, respectively (Cui et al., 2013; Petereit et al., 2023; Short et al., 1976).

We updated the terms to improve the precision and clarity of the two dimensions of social presence

(Krämer et al., 2025).

The experiment had three conditions: own pain, observed pain via live video call, and observed

pain via prerecorded video (Figure 1A). In the "own pain" condition, participants received electrical

stimulation and rated their pain intensity on a visual analog scale (VAS). In the observed pain conditions,

observers watched another participant (the target) receive the stimulation and rated both the target's

pain intensity and their own unpleasantness while watching, also using a VAS.

All participants completed all conditions and took on both roles (observer and target) across

two laboratory sessions (Figure 1B). This within-subjects design differs from our earlier work (Petereit

et al., 2023), in which participants served exclusively as either targets or observers.

The two primary conditions of interest were the counterbalanced live video call and prerecorded

video conditions. In both conditions, observers viewed the target's head, upper body, and the stimulated

hand. In the "video call" condition, both participants could see each other in real time and were aware

of the mutual visibility. In contrast, in the "prerecording" condition, observers viewed a recording of the

target's "own pain" condition, while the target could not see the observer. Cameras were positioned

directly above the screens to approximate face-to-face interaction as closely as possible. The videos

and the VAS were presented using PsychToolbox (PTB-3, Brainard, 1997; Kleiner et al., 2007).

To maintain comparability across conditions, the "own pain" condition was conducted via video

call with an experimenter acting as observer and was recorded for later presentation in the

"prerecording" condition.

Behavioral ratings, electrodermal activity (EDA), and electrocardiography (ECG) were recorded

from both participants across all conditions. Electroencephalography (EEG) was recorded from

observers during pain observation and from participants during their own pain experience, but not from

targets while being observed. To ensure visual consistency, targets nevertheless wore EEG caps in all

conditions. The experimental environment was standardized using gray standing walls surrounding the

participants.

212
**Figure 1.** A) Schematic overview of experimental conditions. In the actual experiment, joysticks and not tablets
were used for behavioral data acquisition and targets also wore EEG caps. B) Overview of the sequence of
interactions between participants. Note that within session 1 participant B underwent the "own pain" condition first
and then observed participant A. Participant B served as observer in session 1 and as target in session 2. The
video that was recorded in the "own pain" condition in session 1 was presented in the "prerecording" condition in
session 2. C) Overview of the trial structure with shock cue, shock and rating.

**Experimental procedure**

Each participant attended two laboratory sessions. In the first session, participants completed

the "own pain" condition and subsequently acted as observers in the "video call" and "prerecording"

conditions, which were presented in randomized order. In the second session, participants performed

the role of the target in the "video call" condition (Figure 1B).

After providing informed consent, participants read the instructions and were fitted with ECG,

EDA, and EEG electrodes. This was followed by the pain stimulus calibration (see next section) and

ten practice rating trials. Participants then completed the pain task in the "own pain" condition, followed

by the two observing conditions. After each condition, participants answered questions assessing the

perceived quality of the interaction. At the end of the experiment, they completed demographic and

personality questionnaires.

***Pain stimulus calibration***

Before the actual experiment, the pain stimulation was calibrated to the subjective pain

thresholds of the participants. The calibration started at 0 mA shock intensity and increased by steps of

0.5 mA. The maximum possible pain stimulation was at 10 mA. Only for the calibration, participants

rated each shock on a 9-point Likert scale from 0 ("not perceivable") to 8 ("worst possible pain"). The

in-between steps on the scale were labeled as 1 = "noticeable", 2 = "unpleasant", 3 = "slightly painful",

4 = "medium painful", 5 = "very painful", 6 = "extremely painful, 7 = "unbearably painful", 8 = "worst

possible pain". The shock intensity was increased until the participant rated the stimulus as 7 =

"unbearably painful". Then, the intensity decreased by steps of 0.5 mA until 0 mA or "not perceivable"

was rated. This procedure of increasing shock intensity was then repeated once. The final stimulus

intensity which was rated as 1 = "noticeable" was used as the lower limit and the intensity rated as 6 =

"extremely painful" was used as the upper limit in the actual experiment. Pain stimulus calibration results

did not differ between the "own pain" and "video call" conditions (lower limits: own pain $M$ = 0.9 mA,

video call $M$ = 0.9 mA, $t(33)$ = 1.22, $p$ = .231; upper limit: own pain $M$ = 6.5 mA, video call $M$ = 6.6 mA,

$t(33)$ = 0.38, $p$ = .703).

***Pain stimulation task***

The pain stimulation task was very similar to that described by  Rütgen et al. (2015). A Digitimer

DS 5 isolated bipolar constant current simulator (Sprenger et al., 2020) was used with a Digitimer bar

electrode (9 mm diameter, 30 mm apart, 10 volts) attached to the back of the left hand. Unlike in our

preceding study, we attached the electrode to the left and not the right hand to facilitate handling of the

right-hand joystick for the rating task. We used an abrasive gel underneath the electrode to improve the

electric conductivity of the skin. A schematic overview of a trial is given in Figure 1C. The pain

stimulation followed an auditory 500-ms cue, which was used for all different pain intensities alike.

Between the cue and the stimulation was a 2000-ms interval in all trials. The electric shock itself was

delivered for 500 ms in a series of 2-ms electric pulses, interspersed with 20-ms breaks. There was an

inter-trial interval with a pseudo-randomized duration of 6-9 seconds. The pain intensity ratings were

given in a total of 50% of the trials and were initialized with a vocal recording saying "Please rate". The

VAS ranged from 0 = "not painful at all" to 100 = "extremely painful". Targets rated the pain intensity for

themselves and observers rated the perceived targets' pain experience. Further, observers rated how

unpleasant it was for themselves to witness the others' pain experience on a VAS from 0 = "not

unpleasant at all" to 100 = "extremely unpleasant".

The range of the electric shocks were evenly split into twenty different levels of intensities in

the range obtained from the pain stimulus calibration for each participant. In total, there were 80 trials

with each intensity appearing four times in each of the three experimental conditions. This is the same

number of trials used previously in empathy for pain tasks (Petereit et al., 2023; Rütgen et al., 2015),

and rating 50% of the trials is sufficient according to prior power analysis. Shock intensity ordering was

fixed in all conditions and for all participants and was pseudo-randomized with less than four low- or

high-intensity shocks in a row. Targets were instructed to express their pain experience as they wished

and the observers were instructed to infer the others' pain as accurately as possible.

**Questionnaires**

Following the completion of each experimental condition, participants were prompted to

indicate how immediate they experienced the interaction, and how close towards and how attached

they felt to the other participant. These measures of perceived social presence were used to validate

the effectiveness of the social presence manipulation and were tested for differences between

conditions using a paired *t*-test. See Supplementary materials S1 for information about additional

questionnaires that were not analyzed in the current study.

**Physiological data acquisition**

EEG data were recorded with 64 sintered Ag/AgCl sensors in a cap with integrated eye and

ear lobe electrodes. Electrode placement was according to the international 10-20 system. The online

reference was placed on the right earlobe and an offline reference was on the left earlobe. Horizontal

and vertical EOG were recorded with 4 electrodes, including Fp2 as the upper vertical electrode, one

lower right vertical electrode and two horizontal electrodes next to the outer corners of the eyes. Data

were recorded with a Brain Products GmbH system (BrainAmp MR plus amplifiers,

BrainVisionRecorder software version 1.21.0102). The sampling rate was 500 Hz with an online high-

pass filter of 0.016 Hz, low-pass filter of 48 Hz and a notch filter at 50 Hz. Impedances were kept below

5 kΩ.

ECG data were recorded with bipolar Ag/AgCl recording electrodes and one reference

electrode, using a 50-Hz notch filter. One of the recording electrodes was placed on the left forearm,

the other one on the right lower calf of the participant (following Einthoven's lead II configuration,

Quigley et al., 2024). Skin conductance was measured with 2 electrodes placed on the thenar and

hypothenar of the right hand, using a 50-Hz notch filter. An electrode attached to the right forearm

served as ground for both ECG and skin conductance, which were recorded with the same amplifier

(BrainAmp ExG, BrainProducts GmbH). In the "video call" condition, data from the observer and target

were recorded synchronously by connecting all amplifiers to the same USB adapter feeding the data

into BrainVisionRecorder.

**Physiological data processing**

***EEG data***

The EEG preprocessing was done using EEGLAB (version 2023.0, Delorme & Makeig, 2004)

and ERPLAB (version 9.0, Lopez-Calderon & Luck, 2014), implemented in MATLAB R2022b (The

Mathworks, 2022). First, data were re-referenced to linked earlobes and bipolar horizontal and vertical

EOG channels were calculated. Noisy or bad data quality channels were marked to be ignored in further

analysis (observing conditions: $M$ = 3.1 channels, $SD$ = 2.7; "own pain" condition: $M$ = 7.1, $SD$ = 2.8).

Due to the first-hand experience of pain and consequent flinching, the data in the "own pain" condition

was more artifactual than in the observing conditions. However, the affected channels were positioned

in the peripheral areas of the skull, away from the central regions of the head that were of interest in

the current study. In the next step, bad data segments or big artifacts were identified visually and

removed. Before the independent components analysis (*runica* function in EEGLAB), data were

bandpass filtered (non-causal Butterworth impulse response function in ERPLAB; lower cut-off: 1 Hz,

upper cut-off: 40 Hz, 48 dB/oct roll-off). The ICA weights were back-projected to the original unfiltered

data without removed bad data segments (Stropahl et al., 2018). This procedure guarantees accurate

counting of rejected trials in the final step of threshold artifact rejection. Another bandpass filter was

applied (non-causal Butterworth impulse response function in ERPLAB; lower cut-off: 0.2 Hz, upper

cut-off: 40 Hz, 48 dB/oct roll-off) to the original data with the back-projected ICA weights. ICA

components clearly related to ocular artifacts (visual inspection) were removed (observing conditions:

*M* = 2.5 components, *SD* = 0.7; "own pain" condition: *M* = 2.7, *SD* = 0.8; Jung et al., 2000). Noisy

channels were interpolated (spherical interpolation). Afterwards, data were epoched into segments

time-locked to the shock onset with epochs ranging from -2500 to 5000 ms and a baseline correction

from -1000 to 0 ms. Finally, a semi-automatized voltage threshold-based artifact correction was

performed (-60/60 to -100/100 μV; rejected trials in observing conditions: *M* = 14%, *SD* = 3%, in "own

pain" condition: *M* = 15%, *SD* = 4%).

In the "prerecording" condition, EEG triggers had to be sent based on what was happening in

the video. Triggers in this case were sent using the logfile data from the corresponding previous

participant's "own pain" condition. However, since the video and the VAS for the observer had to be

presented in parallel, trigger timing deviated slightly more and more variable than expected based on

piloting. Because of this, trigger times in the "prerecording" condition were corrected based on the

trigger times in the logfiles of the corresponding "own pain" condition when the video had been recorded.

The correction of the trigger timings was performed based on the "experiment start" trigger as time point

zero in the "own pain" condition, in which the video was recorded. This "experiment start" trigger

corresponded to the start of the video in the "prerecording" condition and trigger timings were corrected

according to this time point zero.

Finally, we conducted a time-frequency analysis according to recommendations from Cohen

(2014). Data of all channels and in the frequency range of 1-40 Hz were Morlet wavelet convoluted

(cycles evenly spaced from 5 to 14 cycles for the range of 1 to 40 Hz) after Laplace transformation

(Cohen, 2014; Perrin et al., 1989). Power values were converted to decibels with a single-trial baseline

normalization of -700 to -200 ms before stimulus onset. The time window of -500 to 2500 ms was

extracted and downsampled to 100 Hz.

***ECG data***

ECG data preprocessing was done in MATLAB and EEGLAB. The data were bandpass filtered

(non-causal Butterworth impulse response function in ERPLAB; lower cut-off: 1 Hz, upper cut-off: 30

Hz, 48 dB/oct roll-off) and segmented into 10 s epochs (3 s before and 7 s after stimulus onset). The

R-peaks were detected using the MATLAB function *findpeaks*. Afterwards, the epochs were visually

screened for wrongly detected or missing R-peaks. Extrasystoles and other artifacts were removed.

The interbeat interval (IBI) in ms was calculated for every pair of heartbeats in each epoch. The IBI

values were assigned to every data point in the epoch with the original sampling rate. Every epoch was

baseline corrected to the mean baseline averaged across the two seconds before shock onset of all

trials within each participant per condition. Outliers, defined as values exceeding ±3 standard deviation

within each participant, were excluded before averaging, resulting in 18.2% excluded trials.

*Skin conductance data*

Skin conductance responses (SCR) were calculated using the Ledalab-Toolbox for MATLAB

(Benedek & Kaernbach, 2010a, 2010b, Version 3.4.8). Data were downsampled to 50 Hz and smoothed

with an adaptive Gaussian filter. Visually identified artifacts were spline-interpolated. To separate tonic

and phasic electrodermal activity, a continuous deconvolution analysis was conducted (Benedek &

Kaernbach, 2010b). For SCR amplitude analysis, SCRs greater than 0.01 µS were kept for further

analysis (Boucsein et al., 2012). An additional artifact correction was performed before averaging trials,

using a ±3 standard deviation exclusion criterion within each participant. Because participants operated

the joystick with the same hand where the EDA electrodes were attached, artifacts sometimes occurred

beyond the ITI during the first seconds of the next trial. Therefore, the correction was done for a time

window of -1 to 1 sec around the shock onset which was before the onset of the SCR. The artifact

correction resulted in 24.86% excluded trials. Individual SCR data were range-corrected by each

participant's maximum SCR (Hein et al., 2011; Lykken, 1972).

**Statistical analyses**

Before the details of each analysis are described, we will give an overview of the general

statistical analysis rationale:

Linear mixed models (LMM) were calculated for behavioral data, EEG power, and IBI data

using R (R Core Team, 2020). A generalized LMM (GLMM) with a gamma family and log link was

calculated for the right-skewed SCR data (skewness = 2.48). The fixed effects structure was dependent

on the hypotheses given the type of analysis. The details are given in the following subsections. The

model specifications for each final model are given in the Supplementary materials S2. The

experimental condition ("video call" vs. "prerecording") was dummy-coded by using the simple effect-

coding scheme (Daly et al., 2016). The random effects structure was identified by starting with the most

complex random effects structure (random intercept and random slope for condition*predictor-of-

interest interaction for each observer) (Barr et al., 2013) and successively removing the highest-order

random effects with the lowest variance until the model converged, as suggested by Singmann and

Kellen (2019). SCR data required a Gamma log link function for positive, right-skewed data (Lo &

Andrews, 2015). Accordingly, models were calculated with *lmer* and *glmer* functions in the *lme4*

package in R (Bates et al., 2015). LMM significances were tested with the *lmerTest* package

(Kuznetsova et al., 2017). The predictors target rating, shock intensity, physiological data and EEG

power were z-standardized to guarantee (G)LMM convergence. $R^2$ marginal and conditional were

calculated using the *MuMIn* package (Bartoń, 2010).

388            If the target's data (independent variable) significantly affected the observer's data (dependent

variable), it would suggest that the target's response influenced the observer's response. Therefore, it

would be considered empathic. A significant effect of condition ("video call" vs. "prerecording") would

imply that temporal presence influences the observer's response. A significant interaction between

target's data and temporal presence would mean that the observer's empathic response to the target's

pain response differed with the level of temporal presence. Depending on the direction of the effect,

this could indicate increased or diminished empathic responses in the "video call" or "prerecording"

condition. Thus, it would indicate increased or reduced empathy with lower or higher temporal presence.

In the case of non-significant results for the (G)LLMs, we calculated Bayes Factors (Schmalz et al.,

2023) to report evidence for or against the null hypothesis. Therefore, we used an approximation of the

Bayes Factor that is calculated by using the BIC from the obtained (G)LMMs (Faulkenberry, 2018;

Raftery, 1995). Since all mixed models test separate hypotheses, they do not require correction for

multiple comparisons. However, all post hoc tests and correlational analyses were Bonferroni-

corrected.

***Behavioral data***

404            The questions regarding the subjective quality of the interaction were compared between

conditions with a paired *t*-test. For the pain rating data, analyses of empathic accuracy (based on

perceived pain intensity) and of affective empathy (based on unpleasantness ratings) were conducted.

The distinction between a cognitive (empathic accuracy) and an affective (shared unpleasantness)

behavioral readout of empathy was based on classic definitions of empathy and prior studies showing

differences in neural and behavioral correlates of these constructs (Cuff et al., 2016; Rütgen et al.,

2015; Zaki & Ochsner, 2012). We related the observers' ratings of the targets' pain experience to the

targets' pain ratings as measure for empathic accuracy (Gauthier et al., 2008; Ickes, 1993; Ruben &

Hall, 2013; Sun et al., 2018). We related the observers' unpleasantness ratings to the targets' pain

ratings as measure for affective empathy (Rütgen et al., 2015; Shamay-Tsoory, 2011; Zaki & Ochsner,

2012). In the LMMs, we assessed whether observer's pain and unpleasantness ratings were predicted

by the interaction of the target's pain ratings and the temporal presence condition.

416           Although we considered the observers' ratings of the other's pain and their own unpleasantness

as indices of two different psychological constructs (empathic accuracy and affective empathy,

respectively), we checked if these ratings were correlated. We calculated Pearson correlation

coefficients for each dyad per condition separately. To average the resulting $p$-values of all correlations,

combined p-values were calculated using Stouffer's $Z$-score method for dependent $p$-values (Stouffer

et al., 1949). To obtain the distance between observer and target ratings, we calculated the difference

between these ratings for each dyad per condition separately and averaged the differences for each

condition.

***EEG data***

426           First, we explain the analyses for EEG data according to our hypotheses based on the literature

427    and based on the results of the whole time-frequency-electrode space analysis of Petereit et al. (2023).

428    Second, we explain the analysis of the whole time-frequency-electrode space, which closely matches

429    the analysis Petereit et al. (2023) presented. Finally, we explain the analysis of the own pain EEG data.

430           For all statistical analyses of the EEG data, the calculated LMMs contained a fixed effects

431    structure according to our hypotheses, including the interaction of temporal presence condition and the

432    shock intensity. The mean power in each time-frequency cluster was calculated for each trial to be

433    predicted by the model.

434           First, we analyzed the EEG data based on hypotheses derived from the literature for mu

suppression and from our previous study on physical presence (Petereit et al., 2023) for theta and beta

activity. For mu suppression, we analyzed the power within the frequency range 8-12 Hz at electrode

C3 over the somatosensory cortex in a time window from 500 to 1000 ms after shock onset (Genzer et

al., 2022; Hoenen et al., 2015; Perry et al., 2010). For theta activity, we looked at electrode Cz in a

time-window of 100-600 ms at frequencies of 3-9 Hz, and for (low) beta activity, we looked at electrode

CP1 in a time-window of 100-600 ms at frequencies of 11-20 Hz (Petereit et al., 2023).

Similar to our previous study, we conducted an exploratory whole-brain analysis by

investigating the whole time-frequency-electrode space (1-40 Hz, all electrodes). The data was

averaged across participants and permutation tests ($n$=1000) were calculated to obtain null hypothesis

distributions (Cohen, 2014). Three different permutation tests were calculated to investigate the effects

of shock intensity, condition and the interaction of both. For the effect of shock intensity, Spearman

correlation coefficients of shock intensity and EEG power were calculated. In the permutation test,

shock intensity labels were shuffled and randomly assigned. For the effect of condition, condition labels

were shuffled and randomly assigned. Then, condition differences were calculated. For the effect of the

interaction of shock intensity and condition, Spearman correlation coefficients of shock intensity and

EEG power were calculated, shuffled and condition labels were randomly assigned. Then, condition

differences were calculated. We used maximum value correction to correct for multiple comparisons.

To identify the largest clusters and the location of the maximum activity that survived the correction for

multiple comparisons, cluster sizes were calculated using the *bwconncomp* command in MATLAB. By

this, the frequency boundaries and time window boundaries as well as the electrodes of maximum

activity of the largest clusters were identified.

*IBI and SCR data*

To test for SCR coupling, observers' SCR was predicted in a GLMM using a fixed effects

structure of temporal presence condition in interaction with targets' SCR. The physiological coupling

was obtained by the relation of observers' SCR with targets' SCR. For the targets' SCR, we considered

the mean phasic activity in the time window of 2140-5140 ms after shock onset, which corresponded to

the 3-second time window of the maximum averaged response of targets to the shock onset. For

observers' SCR, four time windows (2.5-5.5, 3-6, 3.5-6.5, 4-7 s) were tested to account for a possible

time lag in observers' responses to the targets' pain expression. The 4-7 s time window was used for

analysis, because it yielded the highest correlation of target's SCR with observer's SCR averaged

across both conditions.

To test for IBI coupling, the same procedure as for SCR coupling was used, but with the IBIs

as inputs for the LMM. For the LMMs, the mean IBI in the time window of 2136-5136 ms after shock

onset was used for targets' IBI. This was the 3-second time window (Sperl et al., 2016) of maximum

averaged response of targets' IBI to the shock onset. For observers' IBI, five time windows (2.2-5.2,

2.5-5.5, 3-6, 3.5-6.5, 4-6.8 s) were tested to account for a possible time lag in observers' responses to

the targets' pain expression. The 2.5-5.5 s time window was used for analysis, because it yielded the

highest correlation of targets' IBI with observers' IBI averaged across both conditions.

**Exploratory analysis of the temporal presence effect on EEG latencies**

476            The exploratory whole-brain analysis yielded effects for mid-frontal theta and right posterior

alpha activity. During data analysis, we observed a latency difference between temporal presence

conditions in these frequency bands. To further explore this, we calculated the peak latency for every

participant in both observing conditions separately. Therefore, we automatically identified the maximum

activity (Kompus et al., 2015) in the time window of 100 to 1500 ms after shock onset for theta (1-6 Hz,

Cz) and alpha bands (9-13 Hz, PO8). The frequency bands resulted from the whole-brain analysis. The

latencies of the peaks were compared between conditions using a paired $t$-test.

**Relationships between different empathy measures**

485            We correlated different empathy measures to explore their relationship. Therefore, we

calculated Pearson's correlations between single-trial theta activation and pain or unpleasantness

ratings.  Secondly, we correlated empathic accuracy (correlation of observer and target ratings) and

physiological coupling (correlation between observer and target IBI or SCR). Correlation coefficients

were Fisher-$z$ transformed before averaging across subjects. For comparisons across conditions,

correlation coefficients were calculated per dyad and condition and then averaged. Combined p-values

were calculated using Stouffer's $Z$-score method for dependent $p$-values (Stouffer et al., 1949).

493                                                    **Results**

494    **Difference in perceived qualities of the interaction between temporal presence conditions**

495            After each condition, we asked the observers how they perceived the quality of the interaction,

specifically their feelings of immediacy, closeness of the interaction, and attachment to the other

participant. Observers rated the immediacy ($t(29)$ = 5.16, $p$ < .001, $p_{Bonferroni}$ < .001, $d$ = 0.94) and

closeness ($t(29)$ = 4.02, $p$ < .001, $p_{Bonferroni}$ = .001, $d$ = 0.72) higher after the video call compared to the

prerecorded video, but did not rate their feelings of attachment differently ($t(29) = 1.47$, $p = .153$,

$p_{Bonferroni} = .458$, $d = 0.28$). Results are shown in Figure 2A. These results provide support for the

construct validity of our paradigm.

**Temporal presence does not influence empathic accuracy**

To assess empathic accuracy, we calculated a LMM that contained targets' pain ratings,

condition ("video call" vs. "prerecording") and their interaction to predict observers' pain ratings. Model

parameters can be found in Table 1. Results show that observers' pain ratings were predicted by

targets' pain ratings indicating meaningful empathic accuracy of the observers. However, this effect

was not influenced by condition indicating that empathic accuracy did not differ between video calls and

prerecorded videos (Figure 2B). Correlating observer and target pain ratings yielded similar coefficients

for both conditions (video call: $r(31) = .65$, $p < .001$, $p_{Bonferroni} < .001$; prerecording: $r(31) = 0.64$, $p < .001$,

$p_{Bonferroni} < .001$). The mean distance between observer and target pain ratings was 1.99 ($SD = 15.8$) in

the "video call" and -1.57 ($SD = 16.5$) in the "prerecording" condition, indicating high accuracy in both

conditions considering the 0-100 VAS scale. Testing the model with and without the fixed effect of

condition, the obtained Bayes factor of $BF_{10} = 0.001$ indicates extreme evidence for the model without

the predictor "condition" given our data. The results demonstrate that the observers' empathic accuracy

was equally high across levels of temporal presence.

518
**Figure 2.** A) Observers' perceived qualities of the interaction. After both conditions, observers were asked to rate
the immediacy and closeness of the interaction with the target, and to indicate their attachment to the target.
Condition differences were found for perceived immediacy and closeness but not for attachment. B) LMMs on
single-trial behavioral data. Target ratings significantly predicted observers' pain and unpleasantness ratings, which
indicates empathic responses in observers. But this was similar in both conditions. Black solid and dotted lines
represent the "video call" and "prerecording" averages, respectively, while the green to yellow color coding reflects
the strength of random slopes per participant. * = $p<0.001$, n.s. = not significant.

**Temporal presence does not influence affective empathy**

To assess affective empathy, we calculated a LMM that contained targets' pain ratings, condition and their interaction to predict observers' unpleasantness ratings. Model parameters can be found in Table 1. Results show that observers' unpleasantness ratings were predicted by targets' pain ratings indicating affective empathy of the observers. Further, observers' unpleasantness ratings were generally slightly larger in the "video call" compared to the "prerecording" condition, indicated by a main effect of condition. However, the mean difference of only 1.66 ($SD = 21$) along the VAS, which ranged from 0-100, is negligible. Furthermore, the factors did not interact, showing that the effect of targets' pain ratings on observers' unpleasantness ratings did not vary with the temporal presence condition (Figure 2B). Correlating observers' unpleasantness and targets' pain ratings yielded similar coefficients for both conditions (video call: $r(31) = .58$, $p < .001$, $p_{Bonferroni} < .001$; prerecording: $r(31) = .57$, $p < .001$, $p_{Bonferroni} < .001$). Testing the model with and without the fixed effect of condition, the obtained Bayes factor of $BF_{10} = 0.028$ indicates very strong evidence for the model without the predictor condition given our data. The results indicate that the observers' affective empathy was equally high across levels of temporal presence.

Although we considered the observers' ratings of the other's pain and their own unpleasantness as indices of two different psychological constructs (empathic accuracy and affective empathy, respectively), it should be noted that the two ratings were correlated. When calculating the correlation of ratings across all participants and for each condition, the results confirmed that there is a significant relationship between observers' pain and unpleasantness ratings (video call: $r(31) = .90$, $p < .001$, $p_{Bonferroni} < .001$; prerecording: $r(31) = .90$, $p < .001$, $p_{Bonferroni} < .001$).

548
549 **Table 1.** Results of the single-trial (generalized) linear mixed models on observers' pain and
550 unpleasantness ratings, affective empathy, IBIs, SCRs, and EEG theta, mu, and beta power.

| Model | Random effects | SD | Fixed effects | b (SE) | z t (df) | p | R² marg R² cond |
|---|---|---|---|---|---|---|---|
| Observers' pain ratings ($n = 31$) | Intercept Condition Target rating Condition* target rating | 14.70 6.67 6.34 6.80 | Intercept Condition **Target rating** Condition*target rating | 40.26 (2.67) 1.44 (1.41) 15.37 (1.21) -0.05 (1.44) | 15.09 (30) 1.02 (28) 12.73 (25) -0.04 (28) | <.001 .316 **<.001** .971 | 0.287 0.629 |
| Observers' unpleasantnes s ratings ($n = 31$) | Intercept Condition Target rating Condition* target rating | 20.23 6.31 7.05 5.57 | Intercept **Condition** **Target rating** Condition*target rating | 30.34 (3.65) 3.61 (1.29) 11.20 (1.31) 0.57 (1.18) | 8.30 (30) 2.80 (30) 8.55 (26) 0.48 (30) | <.001 **.009** **<.001** .635 | 0.155 0.745 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Observers' theta power (*n* = 31) | Intercept Condition Intensity | 0.33 0.47 0.20 | Intercept Condition **Intensity** Condition*intensity | -2.65 (0.07) 0.14 (0.11) 0.12 (0.05) 0.02 (0.08) | -37.26 (30) 1.21 (30) 2.32 (30) 0.19 (4175) | <.001 .236 **.028** .848 | 0.003 0.034 |
| Observers' mu power (*n* = 31) | Intercept Condition | 0.96 0.73 | Intercept Condition Intensity Condition*intensity | -3.58 (0.18) 0.25 (0.18) 0.06 (0.06) -0.08 (0.12) | -19.64 (30) 1.47 (30) 1.08 (4197) -0.73 (4198) | <.001 .151 .280 .465 | 0.001 0.070 |
| Observers' beta power (*n* = 31) | Intercept | 0.33 | Intercept Condition Intensity Condition*intensity | -3.10 (0.07) -0.09 (0.09) -0.02 (0.04) -0.04 (0.08) | -44.17 (30) -1.20 (29) -0.59 (4201) -0.45 (4202) | <.001 .230 .556 .651 | 0.001 0.017 |
| Observers' SCR coupling with targets' SCR (*n* = 31) | Intercept Condition Target SCR Condition * target SCR | 0.57 0.85 0.16 0.24 | Intercept Condition **Target SCR** Condition*target SCR | -2.42 (0.09) -0.17 (0.14) 0.15 (0.03) 0.01 (0.06) | -26.39 -1.22 4.52 0.18 | <.001 .222 **<.001** .861 | 0.010 0.192 |
| Observers' IBI coupling with targets' IBI (*n* = 32) | Intercept Condition | 6.95 9.54 | Intercept Condition **Target IBI** Condition*target IBI | 7.38 (1.30) -0.79 (1.90) 1.71 (0.49) 0.77 (0.97) | 5.66 (30) -0.42 (31) 3.49 (4073) 0.80 (3244) | <.001 .680 **<.001** .426 | 0.003 0.081 |

551
552 Notes: IBI = interbeat interval, SCR = skin conductance response, Intensity = shock intensity, $R^2$ *marg* = $R^2$ marginal (explained
553 variance with fixed effects model), $R^2$ *cond* = $R^2$ conditional (explained variance with fixed and random effects model). Models on
554 behavioral data, EEG power, and IBI responses are linear mixed models, and models on SCRs are generalized linear mixed
555 models (note that the former provide Satterthwaite's degrees of freedom and that the latter do not provide degrees of freedom).
556

**Mu suppression, theta, and beta power are not sensitive to temporal presence**

Based on the whole-brain analysis of our last study on physical presence (Petereit et al., 2023)

and on the literature, we hypothesized that observers' mu, theta and beta activity should be modulated

both by the targets' pain level and by the level of temporal presence. We tested these hypotheses by

using LMMs on EEG power predicted by the interaction of condition and shock intensity in the frequency

bands of theta (3-9 Hz, electrode Cz, time window 100-600 ms), mu (8-12 Hz, electrode C3, time

window 500-1000 ms), and beta power (11-20 Hz, electrode CP1, time window 100-600 ms).

Visualizations are shown in the middle and right column of Figure 3A-C. LMM model parameters can

be found in Table 1.

We found an effect of the shock intensity on observers' frontal theta activity, indicating

increased theta activity in observers in response to more intense pain in others. Shock intensity did not

have an effect on observers' mu and beta power. Importantly, there were no significant effects of

condition or of the interaction of shock intensity and condition on mu, theta or beta activity. This

demonstrates that temporal presence did not influence neural activity in the hypothesized frequency

bands derived from the whole-brain analyses from our previous study. The Bayes factors for testing the

effect of condition in the LMMs for theta, mu and beta activity confirmed that there was no temporal

presence effect on EEG power in the given models (theta $BF_{10}$ = 0.0005, mu $BF_{10}$ = 0.0009, beta $BF_{10}$

= 0.0005). Together, these results provide strong evidence against an effect of temporal presence on

EEG power in the mu, theta, and beta frequency bands. See Supplementary materials S3 for analyses

of neural responses to own pain.

577
578
**Figure 3.** Time-power plots for the "own pain", "video call", and "prerecording" condition; for A) Theta (electrode
Cz), B) mu (C3), and C) beta (CP1) oscillations. Participants' own pain was related to theta, mu and beta activity.
In the observing conditions, only theta activity was related to the shock intensity (Supplementary materials S3).
This relationship did not differ significantly between the "video call" and the "prerecording" condition. Grey areas
indicate hypothesized time windows for a shock intensity and condition interaction. Observers' theta responses
occurred later than hypothesized. Shock intensities were dichotomized into low- and high-intensity for display
purposes only.
586

**Exploratory analyses of whole time-frequency-electrode space**

To analyze the neural effects of temporal presence and shock intensity beyond the

hypothesized time windows and frequency bands, we conducted a whole-time-frequency-electrode

space analysis (whole-brain analysis in the following) like we did in our previous study (Petereit et al.,

2023). For this, we used permutation tests to create null hypotheses distributions for the effects of shock

intensity, temporal presence condition, and their interaction. We compared our actual data with these

null hypotheses data to identify significant EEG responses. Results were maximum value corrected.

For the effect of shock intensity averaged over both conditions (Figure 4A), we found a significant cluster

at the frequencies 1-6 Hz in the time window of 270-1450 ms after shock onset with maximum power

at electrode Fpz. This indicates that observers' frontal (low) theta activity is related to the observed pain

of the target. A second significant cluster, showing a negative effect of shock intensity on power

between frequencies 9-13 Hz, was found for the time window of 870-1550 ms after shock onset with

the strongest deactivation at electrode PO8. This indicates that with increasing pain of the target, right

posterior alpha activity in observers was reduced. For the effects of condition and the interaction of

shock intensity and condition (Figure 4B-C), we found no activation that survived correction for multiple

comparisons. Together, these exploratory whole-brain analyses revealed robust effects of shock

intensity on frontal low-theta and posterior alpha activity, but no evidence for effects of temporal

presence.

605            We used the results of the whole-brain analysis to explore the relationship between behavioral

and neural empathic responses. On the single trial level, observer's frontal theta activity (1-6 Hz, Fpz,

270-1450 ms) was weakly, positively correlated to observer's pain ratings ("video call" condition: $r(30)$

= .14, $p < .001$, $p_{Bonferroni} < .001$; "prerecording" condition: $r(30) = .13$, $p < .001$, $p_{Bonferroni} < .001$) and

unpleasantness ratings ("video call" condition: $r(30) = .13$, $p < .001$, $p_{Bonferroni} < .001$; "prerecording"

condition: $r(30) = .13$, $p < .001$, $p_{Bonferroni} < .001$), indicating a relationship between neural and behavioral

empathic responses in empathy for pain.

613
614
**Figure 4.** A) – C) Exploratory analysis of the whole time-frequency-electrode space. Shock intensities were
dichotomized into low- and high-intensity for display purposes only. A) The effect of shock intensity. Two clusters
survived the maximum-value correction. The time-frequency-power plots display correlations of power and pain
intensities at electrodes of maximum amplitude for each cluster. Power time courses display the average power of
the significant frequency range for each cluster. Topographies show the average power across the significant
frequency and time range. B) The interaction of shock intensity and condition. Displayed is the effect of high-pain
minus low-pain intensities per condition averaged across all electrodes. C) The effect of experimental condition:
The heatmap shows the difference between conditions averaged across all electrodes and shock intensities. D)
Exploratory latency analysis of the theta peak. Participants' theta peak was delayed in the "prerecording" compared
to the "video call" condition.
625

**Exploratory analysis of the temporal presence effect on EEG activation latency**

Our theta analyses were based on hypotheses drawn from our previous study. However, as

can be seen in Figure 3A, the actual theta response of observers occurred later than expected (grey

areas indicate hypothesized time windows). To investigate this, we conducted exploratory analyses on

latency effects to determine whether temporal presence has an impact on peak latency in the theta and

alpha bands. The frequency band ranges were based on the exploratory whole-brain analysis, which

yielded effects for mid-frontal theta (1-6 Hz, Fpz) and right posterior alpha activity (9-13 Hz, PO8).

To test weather this latency effect was due to technical reasons and a delay of the video

presentation in the "prerecording" condition, we analyzed the latency of the auditory N1 to the sound

cue which indicated the shock onset. The auditory N1 peaked at 132 ms in the "video call" condition

and at 141 ms in the "prerecording" condition, showing that there were no general delays in the

"prerecording" condition (see Supplementary materials S4 for details).

As the theta and alpha activity delay was not caused by technical issues, we next tested

whether it was statistically robust. We therefore identified the latencies of the maximum theta amplitude

and of the minimum alpha amplitude on the individual subject level. We found a significant difference

in theta peak latencies between conditions (Figure 4D; electrode Cz; $t(30)$ = -2.78, $p$ = .009, $p_{Bonferroni}$ =

.019, $d$ = 0.54). The average theta peak was at 649 ms in the "video call" and at 901 ms in the

"prerecording" condition (difference: 252 ms). The analysis for the alpha decrease yielded no condition

difference (electrode cluster: PO8; peak in the "video call" condition at 984 ms, peak in the

"prerecording" condition at 1094 ms, $t(30)$ = -1.37, $p$ = .180, $p_{Bonferroni}$ = .360). These results indicate that

there is a temporal presence effect on frontal theta activity with an earlier theta peak in the "video call"

condition (higher temporal presence) compared to the "prerecording" condition (lower temporal

presence). Notably, the observed latency effect of temporal presence was not only visible in the time-

frequency data in the theta band, but also in the event-related potentials (see Supplementary materials

S4).

We noticed that a similar effect might be evident in our previous study about the effects of

physical presence on empathy for pain (Petereit et al., 2023). Therefore, we revisited the data from that

study and found a similar latency effect (electrode Cz, latency of the theta peak in the "direct" condition:

636 ms; "mediated" condition: 808 ms; difference: 172 ms; $t(28)$ = -1.92, $p$ = .033, $d$ = 0.30). This result

also shows delayed latencies in the condition with reduced social presence. The time course of the

theta power of the previous study can be seen in Figure 3D in Petereit et al. (2023). Thus, there seems

to be a theta latency difference between social presence conditions in both studies.

**Temporal presence does not modulate physiological coupling between observers and targets**

We used LMMs to test a possible effect of temporal presence on physiological coupling

between observers' and targets' SCR and IBI. Therefore, we calculated whether observers' data could

be predicted by targets' data and whether this effect was modulated by temporal presence. For both

SCR and IBI, target's data predicted the observer's data, showing significant physiological coupling in

terms of cardiac and electrodermal activity. However, no significant effect of condition or an interaction

of target data and condition was found on physiological coupling. Model parameters are shown in Table

1. Testing the SCR model with and without the fixed effect of condition, the obtained Bayes factors of

$BF_{10}$ = 0.0009 indicated extreme evidence for the model without the predictor condition given our data.

This indicates that the SCR coupling did not differ between the conditions with low and high temporal

presence. Testing the IBI model with and without the fixed effect of condition, the obtained Bayes factors

of $BF_{10}$ = 0.0003 indicates extreme evidence for the model without the predictor condition given our

data. This indicates that the IBI coupling did not differ between the conditions with low and high temporal

presence either. Together, these results provide strong evidence that temporal presence does not

modulate physiological coupling between observers and targets in either skin conductance responses

or interbeat intervals. Observers' SCR and IBI data in the observing conditions are shown in Figure 5A-

B and physiological coupling is shown in Figures 5C-D. Neither SCR nor IBI physiological coupling were

correlated to empathic accuracy (calculated across conditions; SCR: $r(28)$ = .25, $p$ = .178, $p_{Bonferroni}$ =

.355; IBI: $r(29)$ = .17, $p$ = .362, $p_{Bonferroni}$ = .725).

**Figure 5.** A) Observers' SCR by condition. B) Observers' IBI change by condition. Shock intensities were
dichotomized into low- and high-intensity for display purposes only. Grey areas indicate time window of maximum
correlation between target and observer data averaged across conditions. C) GLMM on skin-conductance coupling
by predicting observers' SCR with targets' SCR. D) LMM on heartbeat coupling by predicting observers' IBI with
targets' IBI. For both physiological read-outs, observers' empathic responses were coupled to targets' pain
responses. However, this relationship was not affected by temporal presence. Black solid and dotted lines
represent the "video call" and "prerecording" averages, respectively, while the green to yellow color coding reflects
the strength of random slopes per participant.

**Condition order, habituation effects, and sex/gender effects**

We did not find any condition order effects on observers' pain and unpleasantness ratings,

empathic accuracy, or affective empathy measures. Targets and observers showed habituation effects

in their pain and (in case of observers) unpleasantness ratings. However, empathic accuracy and

affective empathy did not change between first and second half of each condition. We did not find any

effects of observers' or targets' sex/gender on empathic accuracy or affective empathy. Details can be

found in Supplementary materials S5.

697                                                    **Discussion**

We examined pain empathy differences between a live, video-mediated interaction as opposed

to watching a prerecorded video of a similar pain scenario. In both conditions, observers watched

targets undergoing painful electric stimulation. Based on our theoretical framework, we expected

reduced temporal presence to impair empathic processing. However, we found that observers were just

as accurate in judging the other's perception of pain when they passively watched a prerecorded video

as when they were in a live video-mediated interaction. Moreover, observers experienced the other's

suffering as equally unpleasant in both conditions. At the neural level, we did not find mu suppression

over the somatosensory cortex to be sensitive to the other's pain in either condition. In contrast, mid-

frontal theta did track the other's pain intensity. Exploratory whole-time-frequency-space analyses

confirmed this result and additionally suggested that a decrease in right posterior alpha power was

involved in tracking the other's pain intensity. However, these neural effects were independent of

temporal presence. Further exploratory analyses revealed a possible latency effect of temporal

presence on pain-related theta activation with delayed responses in the condition with reduced temporal

presence. We observed significant skin-conductance and heartbeat coupling in the pain empathy

paradigm, but this was also independent of temporal presence. In conclusion, the pain paradigm elicited

meaningful empathic responses in observers at the behavioral, neural and physiological levels.

However, these effects were not sensitive to temporal presence, with the exception of a latency effect

of temporal presence on the theta response. The following section will first discuss the general findings

regarding empathy for pain, and then discuss the null findings concerning the effects of temporal

presence on empathy for pain.

**Behavioral, neural and physiological aspects of empathy for pain**

In agreement with other studies on pain empathy, the observers in our study were able to

accurately evaluate the other's pain experience (Ben Adiva et al., 2024; Gauthier et al., 2008; Goldstein

et al., 2017, 2018; Laursen et al., 2014; Leonard et al., 2013; Ruben & Hall, 2013; Shahri et al., 2024).

The observers in our study showed high empathic accuracy, with distances to the targets' ratings close

to zero and a medium-sized correlation between observer and target pain ratings. Also consistent with

other studies, participants showed affective empathy, reflected in increasing unpleasantness ratings in

reaction to the other's suffering (Fan & Han, 2008; Li et al., 2017; Mu et al., 2008; Peng et al., 2021;

Petereit et al., 2023; Rütgen et al., 2015). Further, our findings are consistent with previous studies

(Goldstein et al., 2017; Hein et al., 2011; Murata et al., 2020; Petereit et al., 2023; Reddan et al., 2020),

which demonstrated that empathy for pain is associated with physiological coupling between the pain

receiver and observer. Although the current study lacks a control condition without pain stimulation,

Figure 5A shows that observers indeed exhibit an SCR when observing others in pain, indicating an

empathic response. The relationship between physiological coupling and empathic accuracy seems to

be less clear, since there are mixed results (Jospe et al., 2020; Levenson & Ruef, 1992; Thorson, 2018;

Zerwas et al., 2021). Our results suggest no relationship between physiological coupling and empathic

accuracy. Further research is therefore needed.

Regarding neural indices of empathy, we found that mid-frontal theta activity was related to

others' pain, replicating previous findings in pain empathy research (Misra et al., 2017; Mu et al., 2008;

Peng et al., 2021; Petereit et al., 2023). Moreover, single-trial observers' theta activity correlated with

observers' pain and unpleasantness ratings, demonstrating a close relationship between neural and

behavioral indices of pain empathy. This correspondence was evident for both own pain

(Supplementary Materials S3) and vicarious pain, supporting the interpretation that mid-frontal theta

might reflect the perceived aversiveness or salience of others' suffering (Legrain et al., 2011; Seeley,

2019; Uddin, 2015).

Contrary to our hypothesis and much of the previous literature (Li et al., 2017; Peled-Avron et

al., 2018; Perry et al., 2010; Riecansky & Lamm, 2019; Whitmarsh et al., 2011), we did not observe mu

suppression in response to observing others' pain. As we did observe clear mu suppression in the "own

pain" condition, and as we applied a Laplace transformation (current source density, CSD) to improve

separation of occipital alpha and sensorimotor mu, we are confident that this absence of mu

suppression is not a methodological artefact. Notably, the null finding replicates our earlier study using

the same pain empathy paradigm (Petereit et al., 2023), suggesting that it is not incidental but may be

characteristic of this specific experimental context. One possible explanation is that mu suppression is

particularly sensitive to the observation of overt actions or somatosensory events (Fox et al., 2016;

Perry & Bentin, 2009; Whitmarsh et al., 2011), whereas the present paradigm primarily relied on facial

expressions of pain without visible limb movement or direct action observation. In such contexts,

affective and evaluative components of empathy may be more strongly engaged than sensorimotor

mirroring mechanisms (Saarela et al., 2006). Importantly, the absence of mu suppression occurred

alongside robust behavioral, physiological, and midfrontal theta responses to others' pain, indicating

that empathic processing was preserved despite the lack of a sensorimotor mirroring signature. Taken

together, these findings add to concerns about the reliability and specificity of mu suppression as a

general neural marker of empathy (Hobson & Bishop, 2016) and suggest that its involvement may

depend on task characteristics and the type of empathic cues available.

**Effect of temporal presence on empathy for pain**

764            We did not find effects of temporal presence on behavioral indices of pain empathy. Moreover,

we did not find temporal presence effects on neural mu, beta and theta activity and also not on

physiological markers of empathy for pain. However, there was a tendency of reduced theta power in

the condition without synchronous exchange of information compared to the synchronous condition

(see Figure 4A). Thus, with more power this effect might have reached significance. Furthermore, we

identified a temporal presence latency effect on the theta peak, which indicates delayed responses

without synchronous information exchange.

771            The results of the current study are similar to our previous study, which compared direct, face-

2-face and video call interaction ("physical presence"). In this study (Petereit et al., 2023), we did not

find effects of physical presence on behavioral indices of pain empathy, mu and beta power, and

heartbeat coupling. But in contrast to the current study on temporal presence, there were effects of

physical presence on theta power and EDA coupling in relation to pain empathy. This suggests that

physical presence, i.e., the number of available social cues, might have a stronger influence on the

responses to others' pain than the synchronous exchange of information between pain receiver and

observer. This demonstrates that the empathic physiological coupling between individuals is preserved

without bidirectional exchange of social cues when comparing live video calls and viewing prerecorded

videos.

781           Diana and colleagues (2023) found mimicry and trust to be lower in response to prerecorded

videos compared to direct, live interaction, which suggests that experience-sharing might be diminished

with reduced temporal presence. In contrast, we found that empathic responses to others' pain were

not affected by temporal presence. This suggests that our empathic abilities, like the understanding and

sharing of other's pain, are not impaired by lacking temporal presence. This is in line with Morgan and

colleagues (2022), who found that explicit mentalizing was unaffected by social presence when

comparing direct, live interactions and prerecorded videos in a theory of mind task.

788           As far as our measures of empathy are concerned, our results suggest that temporal presence,

and thus the possibility of reciprocal exchange (at least non-verbally in this case), does not modulate a

person's empathy for the other. Schilbach and colleagues (2013) stressed "the importance of

experiencing and interacting with others as our primary ways of knowing them" (p. 395). In this sense,

the emotional engagement and the social interaction with its complex interaction dynamics between

agents are considered key constituents of social cognition. Similarly, De Jaegher and colleagues (2010)

emphasized that co-regulation of coupling between interacting individuals is crucial for social

understanding. The relational dynamics and connectedness between individuals are supposed to affect

the cognitive processes involved in thinking about the other. Provided that our measures for empathy

are adequate, our results suggest that this is not necessarily the case. In our paradigm, the participants

were able to understand, share the other's pain, and physiologically resonate with the other just as

much without synchronous interaction and by merely observing the other's pain expression. Our data

suggest that the pain perceivers were as emotionally engaged (measured as affective empathy) in a

non-interactive condition as they were in an interactive condition. Thus, for the domain of pain empathy,

understanding and sharing the other's state might not be affected by the temporal presence between

pain receiver and observer. Empathy for pain likely relies on less complex social cues, which enables

understanding and sharing of experiences even without synchronous exchange of cues between

observer and target. This may contrast with more complex social interactions, such as sharing negative

autobiographical emotional events (Jospe et al., 2020), which may depend more heavily on feedback

loops of social exchange.

It can be said, however, that even the live, video mediated condition did not really offer

opportunities for social interaction. The participant's task was to estimate the other's pain intensity and

rate their own unpleasantness in response to the other's suffering, but they were not asked to interact

or respond to the other, or to display prosocial behavior. In contrast, due to the constraints of the EEG

recordings, they were asked to refrain from movements and talking. A paradigm that requires and allows

for more interaction might therefore show more pronounced effects of temporal presence on empathy.

For instance, more complex socio-cognitive phenomena, such as cooperation, prosocial motivation,

and trust, may be more sensitive to temporal presence. In comparison, empathy for pain may rely on

relatively direct, perceptual, affective cues. Nevertheless, our results are informative for second-person

neuroscience (Cañigueral et al., 2022; De Jaegher et al., 2010; Gallotti et al., 2017; Schilbach et al.,

2013), as they show that at least when it comes to empathic responses to others' pain, they remain

preserved and unaltered also with reduced temporal presence and reciprocal interaction between

individuals. Put more positively, our results are reassuring for social neuroscience research using

videos as stimuli, because the differences to direct interactions might not be as strong as one might

expect.

We cannot exclude though that temporal presence has an effect on empathy beyond those

measures of empathy we assessed here. Not being able to interact with the other person due to reduced

temporal presence may have reduced the self-relevance of the other person (Conway et al., 2016;

Schmitz & Johnson, 2007) and thereby the motivation for prosocial behavior or for showing compassion

(Ferguson et al., 2020; Hein et al., 2011; Weisz & Zaki, 2018; Zaki, 2014). As our experimental setup

did not allow the observers to express their compassion – other than through facial expressions – and

as we did not assess their motivation for prosocial behavior, this remains speculative. Future social

presence research should consider measuring motivational aspects of empathy.

Although the results of our hypothesis testing indicated no effect of temporal presence on

empathy for pain, we did identify a latency effect of temporal presence on frontal theta activity.

Participants' frontal theta activity in response to the others' pain was delayed in the condition with

reduced temporal presence with a difference of about 252 ms. This latency shift constitutes the primary

neural marker of reduced temporal presence in the present study, indicating an alteration in the

temporal dynamics of empathic processing rather than its overall magnitude. Possibly, differences in

visual attention and self-relevance could explain the latency effect. On a behavioral level, self-relevance

has been found to speed up visual search responses (Wade & Vickery, 2018) and to impact visual

attention (Svensson et al., 2023). On the neural level, self-relevant information increases neural

responses (Sui et al., 2013; Viskontas et al., 2009). Not being seen by the observed person may lead

to differences in attending visually to the others' pain expression compared to social encounters in

which both participants can see each other (Hietanen et al., 2020; Morgan et al., 2022). Thus,

differences in self-relevance may have affected attention processes in the current study, reflected in

delayed neural responses.

845            Interestingly, revisiting the data from our previous study about the effects of physical presence

on empathy for pain revealed a similar latency effect. The latency of the theta peak in the condition with

reduced social cues (live video call) was 172 ms later than in the condition with rich social cues (direct,

face-to-face). Together, these findings suggest that reduced social presence - across both temporal

and physical dimensions - consistently affects the timing of neural responses associated with empathy

for pain. Further studies are needed to specify the cognitive or motivational factors that are relevant

here, be it attention and self-relevance or other factors. The combination of preserved behavioral

responses and physiological coupling in empathy for pain with delayed frontal theta activity under

reduced temporal presence may indicate that empathic processing remains largely intact, while its

neural timing is altered. Rather than reflecting a reduction in empathy, the latency shift could point to

subtle changes or compensatory mechanisms in the processing dynamics supporting empathic

understanding when temporal presence is reduced.

**Limitations**

859            As with any null effect, the lack of condition differences in the current study could be due to the

860    sample size and too limited power. However, the prior power analysis suggested that a sample of thirty

861    participants is sufficient to detect also a small behavioral effect in a within-subject design. Moreover,

862    the Bayes factor analyses indicated quite strong evidence for the null hypotheses.

863            Our sample was quite homogeneous, consisting of young, predominantly female students. This

limits generalizability to male, older and non-student populations. In addition, younger people may be

more adept at digitally mediated interactions (Vossen & Valkenburg, 2016), which could offset any

impacts of temporal presence on empathy. Vice versa, effects of temporal presence on empathy might

be more pronounced in populations with less experience in digitally mediated social interactions.

Furthermore, the homogeneity of the sample might have contributed to the generally high levels

of empathic accuracy, as previous research has shown that people are more empathically accurate

with similar others (Han, 2018; Majdandžić et al., 2016) and with familiar others (Israelashvili & Perry,

2021). We only included dyads of strangers and guaranteed low levels of familiarity. However, as

participants were all similarly young students of the same University, it is likely that they perceived each

other as quite similar. The fact that participants were not more accurate in the live video calls compared

to the prerecorded video condition may be thus due to a ceiling effect. Future studies on social presence

might therefore consider including more heterogenous samples.

Although the "video call" condition met the criteria for an actual video call, the participants

remained in close proximity, and knew they were in adjacent rooms. Due to the within-subject design

of the study, it was not possible to physically separate the participants further. This limits the

generalizability of our findings, since real video calls usually also involve greater physical distance, and

the person's knowledge about this greater physical distance might lower the relevance of the other

person for the self. Future studies will have to show whether, in addition to technical transmission, the

actual physical distance plays a role for the participants' behavioral, physiological, and neural empathic

responses.

In terms of technical limitations, it should be noted that the hand, to which EDA electrodes were

attached, was also used for operating the joystick. Unfortunately, operating the joystick for the VAS

ratings introduced artifacts to the EDA data, which made it necessary to exclude trials. With more trials

and therefore more power, we might have been able to discover the probably nuanced effects of

temporal presence in the EDA data, as there was at least a trend for a condition difference (see Figure

5A).

891                                             **Conclusion**

Digitally-mediated social interactions have become increasingly common in daily life. Research

has just begun to explicitly compare direct and mediated interactions (Diana et al., 2023; Hietanen et

al., 2020; Järveläinen et al., 2001; Karimova et al., 2024; Prinsen & Alaerts, 2019; Shimada & Hiraki,

2006), especially with regard to social cognition (Jospe et al., 2020; Morgan et al., 2022; Petereit et al.,

2023; Pönkänen et al., 2011). In the current study on temporal presence and pain empathy, we found

no evidence that a lack of temporal presence between individuals significantly affects empathy for pain.

This indicates that major aspects of empathic responses to other's pain are unaltered in mediated

interactions. At the neural level, reduced temporal presence was primarily reflected in a latency shift of

frontal theta activity, suggesting an alteration in the temporal dynamics of empathic processing rather

than in its overall magnitude. Therefore, our findings are relevant for second-person neuroscience

(Schilbach et al., 2013), but also for online psychotherapy and telemedicine (Amichai-Hamburger et al.,

2014; Grondin et al., 2019). If empathic responses to others' pain are preserved when watching a

prerecorded video, then the cost of online therapy through live video interaction might be negligible, at

least in terms of understanding others' pain. Future studies will need to examine the effects of social

presence on empathy in more complex contexts and considering also motivational aspects of empathy.

**Funding**

This work was supported by the German Science Foundation (grant number KR3691/12-1).

**Data and Code availability**

The raw data, the final preprocessed data, which is prepared for statistical analysis, and the analysis

code can be found at https://osf.io/3jdf4/ (DOI 10.17605/OSF.IO/3JDF4).

**Acknowledgements**

We would like to thank Lea Henke, Kira Weckmann, Liina Elsner, and Nadine Horchler for their

assistance in acquiring the data. We also thank Andreas Sprenger and Marcus Heldmann for their

technical support.

**AI Usage Statement**

Artificial intelligence tools (ChatGPT and DeepL) were employed only to assist with language editing

and sentence-level rephrasing. All scientific concepts, analyses, and conclusions are entirely the

author's own or based on the previous scientific literature.

**References**

Amichai-Hamburger, Y., Klomek, A. B., Friedman, D., Zuckerman, O., & Shani-Sherman, T. (2014).

926           The future of online therapy. *Computers in Human Behavior*, *41*, 288–294.

927           https://doi.org/10.1016/j.chb.2014.09.016

928    Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory

929           hypothesis testing: Keep it maximal. *Journal of Memory and Language*, *68*(3), 255–278.

930           https://doi.org/10.1016/j.jml.2012.11.001

931    Bartoń, K. (2010). *MuMIn: Multi-Model Inference* (p. 1.47.5) [Dataset].

932           https://doi.org/10.32614/CRAN.package.MuMIn

933    Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using *lme4*.

934           *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

935    Bekkali, S., Youssef, G. J., Donaldson, P. H., Albein-Urios, N., Hyde, C., & Enticott, P. G. (2021). Is the

936           Putative Mirror Neuron System Associated with Empathy? A Systematic Review and Meta-

937           Analysis. *Neuropsychology Review*, *31*(1), 14–57. https://doi.org/10.1007/s11065-020-09452-

938           6

939    Ben Adiva, Y., Genzer, S., & Perry, A. (2024). Beyond physical sensations: Investigating empathy and

940           prosocial behavior in vicarious pain responders. *Social Cognitive and Affective Neuroscience*,

941           *19*(1), nsae039. https://doi.org/10.1093/scan/nsae039

942    Benedek, M., & Kaernbach, C. (2010a). A continuous measure of phasic electrodermal activity. *Journal*

943           *of Neuroscience Methods*, *190*(1–5), 80–91. https://doi.org/10.1016/j.jneumeth.2010.04.028

944    Benedek, M., & Kaernbach, C. (2010b). Decomposition of skin conductance data by means of

945           nonnegative deconvolution. *Psychophysiology*, *47*(4), 647–658. https://doi.org/10.1111/j.1469-

946           8986.2009.00972.x

947    Boucsein, W., Fowles, D. C., Grimnes, S., Ben-Shakhar, G., Roth, W. T., Dawson, M. E., & Filion, D.

948           L. (2012). Publication recommendations for electrodermal measurements. *Psychophysiology*,

949           *49*(8), 1017–1034. https://doi.org/10.1111/j.1469-8986.2012.01384.x

950    Caggiano, V., Fogassi, L., Rizzolatti, G., Pomper, J. K., Thier, P., Giese, M. A., & Casile, A. (2011).

951           View-Based Encoding of Actions in Mirror Neurons of Area F5 in Macaque Premotor Cortex.

952           *Current Biology, 21*(2), 144–148. https://doi.org/10.1016/j.cub.2010.12.022

Cañigueral, R., Krishnan-Barman, S., & Hamilton, A. F. D. C. (2022). Social signalling as a framework
for second-person neuroscience. *Psychonomic Bulletin & Review*, *29*(6), 2083–2095.
https://doi.org/10.3758/s13423-022-02103-2

Cheng, Y., Lin, C.-P., Liu, H.-L., Hsu, Y.-Y., Lim, K.-E., Hung, D., & Decety, J. (2007). Expertise
Modulates the Perception of Pain in Others. *Current Biology*, *17*(19), 1708–1713.
https://doi.org/10.1016/j.cub.2007.09.020

Cheng, Y., Yang, C.-Y., Lin, C.-P., Lee, P.-L., & Decety, J. (2008). The perception of pain in others
suppresses somatosensory oscillations: A magnetoencephalography study. *NeuroImage*,
*40*(4), 1833–1840. https://doi.org/10.1016/j.neuroimage.2008.01.064

Cohen, M. X. (2014). *Analyzing Neural Time Series Data: Theory and Practice*. The MIT Press.
https://doi.org/10.7551/mitpress/9609.001.0001

Conway, M. A., Pothos, E. M., & Turk, D. J. (2016). The self-relevance system? *Cognitive*
*Neuroscience*, *7*(1–4), 20–21. https://doi.org/10.1080/17588928.2015.1075484

Cuff, B. M. P., Brown, S. J., Taylor, L., & Howat, D. J. (2016). Empathy: A Review of the Concept.
*Emotion Review*, *8*(2), 144–153. https://doi.org/10.1177/1754073914558466

Cui, G., Lockee, B., & Meng, C. (2013). Building modern online social presence: A review of social
presence theory and its instructional design implications for future trends. *Education and*
*Information Technologies*, *18*(4), 661–685. https://doi.org/10.1007/s10639-012-9192-1

Daly, A., Dekker, T., & Hess, S. (2016). Dummy coding vs effects coding for categorical variables:
Clarifications and extensions. *Journal of Choice Modelling*, *21*, 36–41.
https://doi.org/10.1016/j.jocm.2016.09.005

De Jaegher, H., Di Paolo, E., & Gallagher, S. (2010). Can social interaction constitute social cognition?
*Trends in Cognitive Sciences*, *14*(10), 441–447. https://doi.org/10.1016/j.tics.2010.06.009

Decety, J., & Jackson, P. L. (2004). The Functional Architecture of Human Empathy. *Behavioral and*
*Cognitive Neuroscience Reviews*, *3*(2), 71–100. https://doi.org/10.1177/1534582304267187

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *Journal of Neuroscience Methods*,
*134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Diana, F., Juárez-Mora, O. E., Boekel, W., Hortensius, R., & Kret, M. E. (2023). How video calls affect
mimicry and trust during interactions. *Philosophical Transactions of the Royal Society B:*
*Biological Sciences*, *378*(1875), 20210484. https://doi.org/10.1098/rstb.2021.0484

Dumas, G. (2011). Towards a two-body neuroscience. *Communicative & Integrative Biology*, *4*(3), 349–
352. https://doi.org/10.4161/cib.4.3.15110

Fallon, N., Roberts, C., & Stancak, A. (2020). Shared and distinct functional networks for empathy and
pain processing: A systematic review and meta-analysis of fMRI studies. *Social Cognitive and*
*Affective Neuroscience*, *15*(7), 709–723. https://doi.org/10.1093/scan/nsaa090

Fan, Y., & Han, S. (2008). Temporal dynamic of neural mechanisms involved in empathy for pain: An
event-related     brain     potential     study.     *Neuropsychologia*,     *46*(1),     160–173.
https://doi.org/10.1016/j.neuropsychologia.2007.07.023

Faulkenberry, T. J. (2018). Computing Bayes factors to measure evidence from experiments: An
extension    of    the    BIC    approximation.    *Biometrical    Letters*,    *55*(1),    31–43.
https://doi.org/10.2478/bile-2018-0003

Ferguson, A. M., Cameron, C. D., & Inzlicht, M. (2020). Motivational effects on empathic choices.
*Journal     of     Experimental     Social     Psychology*,     *90*,     104010.
https://doi.org/10.1016/j.jesp.2020.104010

Fox, Bakermans-Kranenburg, M. J., Yoo, K. H., Bowman, L. C., Cannon, E. N., Vanderwert, R. E.,
Ferrari, P. F., & Van IJzendoorn, M. H. (2016). Assessing human mirror activity with EEG mu
rhythm:    A    meta-analysis.    *Psychological    Bulletin*,    *142*(3),    291–313.
https://doi.org/10.1037/bul0000031

Gabriel, D., Devos, J., Chapuis, S., Comte, A., & Billot, P.-E. (2025). Watching live performances
enhances subjective and physiological emotional responses compared to viewing the same
performance    on    screen.    *IBRO    Neuroscience    Reports*,    *19*,    381–390.
https://doi.org/10.1016/j.ibneur.2025.08.002

Gallese, V., Keysers, C., & Rizzolatti, G. (2004). A unifying view of the basis of social cognition. *Trends*
*in Cognitive Sciences*, *8*(9), 396–403. https://doi.org/10.1016/j.tics.2004.07.002

Gallo, S., Paracampo, R., Müller-Pinzler, L., Severo, M. C., Blömer, L., Fernandes-Henriques, C.,
Henschel, A., Lammes, B. K., Maskaljunas, T., Suttrup, J., Avenanti, A., Keysers, C., &

Gazzola, V. (2018). The causal role of the somatosensory cortex in prosocial behaviour. *eLife*,

1011                *7*, e32740. https://doi.org/10.7554/eLife.32740

1012    Gallotti, M., Fairhurst, M. T., & Frith, C. D. (2017). Alignment in social interactions. *Consciousness and*

1013                *Cognition*, *48*, 253–261. https://doi.org/10.1016/j.concog.2016.12.002

1014    Gauthier, N., Thibault, P., & Sullivan, M. J. L. (2008). Individual and Relational Correlates of Pain-

1015                related Empathic Accuracy in Spouses of Chronic Pain Patients. *The Clinical Journal of Pain*,

1016                *24*(8), 669–677. https://doi.org/10.1097/AJP.0b013e318173c28f

1017    Genzer, S., Ong, D. C., Zaki, J., & Perry, A. (2022). Mu rhythm suppression over sensorimotor regions

1018                is associated with greater empathic accuracy. *Social Cognitive and Affective Neuroscience*,

1019                *17*(9), 788–801. https://doi.org/10.1093/scan/nsac011

1020    Goldstein, P., Weissman-Fogel, I., Dumas, G., & Shamay-Tsoory, S. G. (2018). Brain-to-brain coupling

1021                during handholding is associated with pain reduction. *Proceedings of the National Academy of*

1022                *Sciences*, *115*(11). https://doi.org/10.1073/pnas.1703643115

1023    Goldstein, P., Weissman-Fogel, I., & Shamay-Tsoory, S. G. (2017). The role of touch in regulating inter-

1024                partner physiological coupling during empathy for pain. *Scientific Reports*, *7*(1), 3252.

1025                https://doi.org/10.1038/s41598-017-03627-7

1026    González-Roldan, A. M., Martínez-Jauand, M., Muñoz-García, M. A., Sitges, C., Cifre, I., & Montoya,

1027                P. (2011). Temporal dissociation in the brain processing of pain and anger faces with different

1028                intensities    of    emotional    expression.    *Pain*,    *152*(4),    853–859.

1029                https://doi.org/10.1016/j.pain.2010.12.037

1030    Goubert, L., Craig, K. D., Vervoort, T., Morley, S., Sullivan, M. J. L., Williams, D. C. A. C., Cano, A., &

1031                Crombez, G. (2005). Facing others in pain: The effects of empathy. *Pain*, *118*(3), 285–288.

1032                https://doi.org/10.1016/j.pain.2005.10.025

1033    Green, P., & MacLeod, C. J. (2016). SIMR: An R package for power analysis of generalized linear

1034                mixed models by simulation. *Methods in Ecology and Evolution*, *7*(4), 493–498.

1035    Grondin, F., Lomanowska, A. M., & Jackson, P. L. (2019). Empathy in computer-mediated interactions:

1036                A conceptual framework for research and clinical practice. *Clinical Psychology: Science and*

1037                *Practice*, *26*(4), 17–17. https://doi.org/10.1111/cpsp.12298

Güntekin, B., & Başar, E. (2014). A review of brain oscillations in perception of faces and emotional

pictures.                    *Neuropsychologia*,              *58*,                    33–51.

https://doi.org/10.1016/j.neuropsychologia.2014.03.014

Hamilton, A. F. D. C. (2016). Gazing at me: The importance of social meaning in understanding direct-

gaze cues. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1686),

20150080. https://doi.org/10.1098/rstb.2015.0080

Hamilton, A. F. D. C., & Holler, J. (2023). Face2face: Advancing the science of social interaction.

*Philosophical Transactions of the Royal Society B: Biological Sciences*, *378*(1875), 20210470.

https://doi.org/10.1098/rstb.2021.0470

Hamilton, A. F. D. C., & Lind, F. (2016). Audience effects: What can they tell us about social

neuroscience,   theory   of   mind   and   autism?   *Culture   and   Brain*,   *4*(2),   159–177.

https://doi.org/10.1007/s40167-016-0044-5

Han, S. (2018). Neurocognitive Basis of Racial Ingroup Bias in Empathy. *Trends in Cognitive Sciences*,

*22*(5), 400–421. https://doi.org/10.1016/j.tics.2018.02.013

Hari, R., & Kujala, M. V. (2009). Brain Basis of Human Social Interaction: From Concepts to Brain

Imaging. *Physiological Reviews*, *89*(2), 453–479. https://doi.org/10.1152/physrev.00041.2007

Hein, G., Lamm, C., Brodbeck, C., & Singer, T. (2011). Skin Conductance Response to the Pain of

Others   Predicts   Later   Costly   Helping.   *PLoS   ONE*,   *6*(8),   e22759.

https://doi.org/10.1371/journal.pone.0022759

Hietanen, J. O., Peltola, M. J., & Hietanen, J. K. (2020). Psychophysiological responses to eye contact

in   a   live   interaction   and   in   video   call.   *Psychophysiology*,   *57*(6).

https://doi.org/10.1111/psyp.13587

Hobson, H. M., & Bishop, D. V. M. (2016). Mu suppression – A good measure of the human mirror

neuron system? *Cortex*, *82*, 290–310. https://doi.org/10.1016/j.cortex.2016.03.019

Hoenen, M., Lübke, K. T., & Pause, B. M. (2015). Somatosensory mu activity reflects imagined pain

intensity of others: Mu activity reflects imagined pain intensity. *Psychophysiology*, *52*(12),

1551–1558. https://doi.org/10.1111/psyp.12522

Hutchison, W. D., Davis, K. D., Lozano, A. M., Tasker, R. R., & Dostrovsky, J. O. (1999). Pain-related

neurons   in   the   human   cingulate   cortex.   *Nature   Neuroscience*,   *2*(5),   403–405.

https://doi.org/10.1038/8065

Ickes, W. (1993). Empathic Accuracy. *Journal of Personality*, *61*(4), 587–610.
https://doi.org/10.1111/j.1467-6494.1993.tb00783.x

Ickes, W., Stinson, L., Bissonnette, V., & Garcia, S. (1990). *Naturalistic Social Cognition: Empathic*
*Accuracy in Mixed-Sex Dyads*.

Israelashvili, J., & Perry, A. (2021). Nuancing Perspective: Feedback Shapes the Understanding of
Another's Emotions. *Social Psychology*, *52*(4), 238–249. https://doi.org/10.1027/1864-
9335/a000452

Jackson, P. L., Meltzoff, A. N., & Decety, J. (2005). How do we perceive the pain of others? A window
into the neural processes involved in empathy. *NeuroImage*, *24*(3), 771–779.
https://doi.org/10.1016/j.neuroimage.2004.09.006

Järvelä, S., Kätsyri, J., Ravaja, N., Chanel, G., & Henttonen, P. (2016). Intragroup Emotions:
Physiological Linkage and Social Presence. *Frontiers in Psychology*, *7*.
https://doi.org/10.3389/fpsyg.2016.00105

Järveläinen, J., Schürmann, M., Avikainen, S., & Hari, R. (2001). Stronger reactivity of the human
primary motor cortex during observation of live rather than video motor acts: *Neuroreport*,
*12*(16), 3493–3495. https://doi.org/10.1097/00001756-200111160-00024

Jauniaux, J., Khatibi, A., Rainville, P., & Jackson, P. L. (2019). A meta-analysis of neuroimaging studies
on pain empathy: Investigating the role of visual information and observers' perspective. *Social*
*Cognitive and Affective Neuroscience*, *14*(8), 789–813. https://doi.org/10.1093/scan/nsz055

Jospe, K., Genzer, S., Klein Selle, N., Ong, D., Zaki, J., & Perry, A. (2020). The contribution of linguistic
and visual cues to physiological synchrony and empathic accuracy. *Cortex*, *132*, 296–308.
https://doi.org/10.1016/j.cortex.2020.09.001

Jung, T.-P., Makeig, S., Humphries, C., Lee, T.-W., McKeown, M. J., Iragui, V., & Sejnowski, T. J.
(2000). Removing electroencephalographic artifacts by blind source separation.
*Psychophysiology*, *37*(2), 163–178. https://doi.org/10.1111/1469-8986.3720163

Karimova, E. D., Ovakimian, A. S., & Katermin, N. S. (2024). Live vs video interaction: Sensorimotor
and visual cortical oscillations during action observation. *Cerebral Cortex*, *34*(4), bhae168.
https://doi.org/10.1093/cercor/bhae168

Keysers, C., & Gazzola, V. (2006). Towards a unifying neural theory of social cognition. In *Progress in*
*Brain   Research*   (Vol.   156,   pp.   379–401).   Elsevier.   https://doi.org/10.1016/S0079-
6123(06)56021-2

Kompus, K., Westerhausen, R., Craven, A. R., Kreegipuu, K., Põldver, N., Passow, S., Specht, K.,
Hugdahl, K., & Näätänen, R. (2015). Resting-state glutamatergic neurotransmission is related
to the peak latency of the auditory mismatch negativity (MMN) for duration deviants: An 1H-
MRS-EEG study. *Psychophysiology*, *52*(9), 1131–1139. https://doi.org/10.1111/psyp.12445

Krämer, U. M., Perry, A., & Jessen, S. (2025). *Social presence and its role in social interactions:*
*Insights,   challenges   and   a   call   for   action*.   Open   Science   Framework.
https://doi.org/10.31219/osf.io/f7rkg_v1

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). **lmerTest** Package: Tests in Linear
Mixed   Effects   Models.   *Journal   of   Statistical   Software*,   *82*(13).
https://doi.org/10.18637/jss.v082.i13

Lamm, C., Decety, J., & Singer, T. (2011). Meta-analytic evidence for common and distinct neural
networks associated with directly experienced pain and empathy for pain. *Neuroimage*, *54*(3),
2492–2502.

Laursen, H. R., Siebner, H. R., Haren, T., Madsen, K., Grønlund, R., Hulme, O., & Henningsson, S.
(2014). Variation in the oxytocin receptor gene is associated with behavioral and neural
correlates   of   empathic   accuracy.   *Frontiers   in   Behavioral   Neuroscience*,   *8*.
https://doi.org/10.3389/fnbeh.2014.00423

Legrain, V., Iannetti, G. D., Plaghki, L., & Mouraux, A. (2011). The pain matrix reloaded. *Progress in*
*Neurobiology*, *93*(1), 111–124. https://doi.org/10.1016/j.pneurobio.2010.10.005

Leonard, M. T., Issner, J. H., Cano, A., & Williams, A. M. (2013). Correlates of Spousal Empathic
Accuracy for Pain-related Thoughts and Feelings. *The Clinical Journal of Pain*, *29*(4), 324.
https://doi.org/10.1097/AJP.0b013e3182527bfd

Levenson, R., & Gottman, J. M. (1983). *Marital Interaction: Physiological Linkage and Affective*
*Exchange*.

Levenson, & Ruef, A. M. (1992). Empathy: A physiological substrate. *Journal of Personality and Social*
*Psychology*, *63*(2), 234.

Li, X., Meng, X., Li, H., Yang, J., & Yuan, J. (2017). The impact of mood on empathy for pain: Evidence

from an EEG study. *Psychophysiology*, *54*(9), 1311–1322. https://doi.org/10.1111/psyp.12882

Lieberman, M. D. (2007). Social Cognitive Neuroscience: A Review of Core Processes. *Annual Review*

*of Psychology*, *58*(1), 259–289. https://doi.org/10.1146/annurev.psych.58.110405.085654

Lo, S., & Andrews, S. (2015). To transform or not to transform: Using generalized linear mixed models

to analyse reaction time data. *Frontiers in Psychology*, *6*.

https://doi.org/10.3389/fpsyg.2015.01171

Lopez-Calderon, J., & Luck, S. J. (2014). ERPLAB: An open-source toolbox for the analysis of event-

related potentials. *Frontiers in Human Neuroscience*, *8*.

https://doi.org/10.3389/fnhum.2014.00213

Lykken, D. T. (1972). Range Correction Applied to Heart Rate and to GSR Data. *Psychophysiology*,

*9*(3), 373–379. https://doi.org/10.1111/j.1469-8986.1972.tb03222.x

Mackes, N. K., Golm, D., O'Daly, O. G., Sarkar, S., Sonuga-Barke, E. J. S., Fairchild, G., & Mehta, M.

A. (2018). Tracking emotions in the brain – Revisiting the Empathic Accuracy Task.

*NeuroImage*, *178*, 677–686. https://doi.org/10.1016/j.neuroimage.2018.05.080

Majdandžić, J., Amashaufer, S., Hummer, A., Windischberger, C., & Lamm, C. (2016). The selfless

mind: How prefrontal involvement in mentalizing with similar and dissimilar others shapes

empathy and prosocial behavior. *Cognition*, *157*, 24–38.

https://doi.org/10.1016/j.cognition.2016.08.003

Marsh, A. A. (2018). The neuroscience of empathy. *Current Opinion in Behavioral Sciences*, *19*, 110–

115. https://doi.org/10.1016/j.cobeha.2017.12.016

Misra, G., Wang, W., Archer, D. B., Roy, A., & Coombes, S. A. (2017). Automated classification of pain

perception using high-density electroencephalography data. *Journal of Neurophysiology*,

*117*(2), 786–795. https://doi.org/10.1152/jn.00650.2016

Mitchell, D. J., McNaughton, N., Flanagan, D., & Kirk, I. J. (2008). Frontal-midline theta from the

perspective of hippocampal "theta." *Progress in Neurobiology*, *86*(3), 156–185.

https://doi.org/10.1016/j.pneurobio.2008.09.005

Morgan, E. J., Carroll, D. J., Chow, C. K. C., & Freeth, M. (2022). The Effect of Social Presence on

Mentalizing Behavior. *Cognitive Science*, *46*(4), e13126. https://doi.org/10.1111/cogs.13126

Mu, Y., Fan, Y., Mao, L., & Han, S. (2008). Event-related theta and alpha oscillations mediate empathy

1155            for pain. *Brain Research*, *1234*, 128–136. https://doi.org/10.1016/j.brainres.2008.07.113

1156    Murata, A., Nishida, H., Watanabe, K., & Kameda, T. (2020). Convergence of physiological responses

1157            to pain during face-to-face interaction. *Scientific Reports*, *10*(1), 450.

1158            https://doi.org/10.1038/s41598-019-57375-x

1159    Neumann, D. L., & Westbury, H. R. (2011). *THE PSYCHOPHYSIOLOGICAL MEASUREMENT OF*

1160            *EMPATHY*.

1161    Palumbo, R. V., Marraccini, M. E., Weyandt, L. L., Wilder-Smith, O., McGee, H. A., Liu, S., & Goodwin,

1162            M. S. (2017). Interpersonal Autonomic Physiology: A Systematic Review of the Literature.

1163            *Personality    and    Social    Psychology    Review*,    *21*(2),    99–141.

1164            https://doi.org/10.1177/1088868316628405

1165    Pan, H., Chen, Z., Jospe, K., Gao, Q., Sheng, J., Gao, Z., & Perry, A. (2023). Mood congruency affects

1166            physiological synchrony but not empathic accuracy in a naturalistic empathy task. *Biological*

1167            *Psychology*, *184*, 108720. https://doi.org/10.1016/j.biopsycho.2023.108720

1168    Peled-Avron, L., Goldstein, P., Yellinek, S., Weissman-Fogel, I., & Shamay-Tsoory, S. G. (2018).

1169            Empathy during consoling touch is modulated by mu-rhythm: An EEG study. *Neuropsychologia,*

1170            *Special    Issue:    The    Neuroscience    of    Empathy*,    *116*,    68–74.

1171            https://doi.org/10.1016/j.neuropsychologia.2017.04.026

1172    Peng, W., Lou, W., Huang, X., Ye, Q., Tong, R. K.-Y., & Cui, F. (2021). Suffer together, bond together:

1173            Brain-to-brain synchronization and mutual affective empathy when sharing painful experiences.

1174            *NeuroImage*, *238*, 118249. https://doi.org/10.1016/j.neuroimage.2021.118249

1175    Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. (1989). Spherical splines for scalp potential and

1176            current density mapping. *Electroencephalography and Clinical Neurophysiology*, *72*(2), 184–

1177            187. https://doi.org/10.1016/0013-4694(89)90180-6

1178    Perry, A., & Bentin, S. (2009). Mirror activity in the human brain while observing hand movements: A

1179            comparison between EEG desynchronization in the μ-range and previous fMRI results. *Brain*

1180            *Research*, *1282*, 126–132. https://doi.org/10.1016/j.brainres.2009.05.059

1181    Perry, A., Bentin, S., Bartal, I. B.-A., Lamm, C., & Decety, J. (2010). "Feeling" the pain of those who are

1182            different from us: Modulation of EEG in the mu/alpha range. *Cognitive, Affective, & Behavioral*

1183            *Neuroscience*, *10*(4), 493–504. https://doi.org/10.3758/CABN.10.4.493

Petereit, P., Weiblen, R., Perry, A., & Krämer, U. M. (2023). Effects of social presence on behavioral,

neural, and physiological aspects of empathy for pain. *Cerebral Cortex*, 1–17.

https://doi.org/10.1093/cercor/bhad257

Pönkänen, L. M., Alhoniemi, A., Leppänen, J. M., & Hietanen, J. K. (2011). Does it make a difference if

I have an eye contact with you or with your picture? An ERP study. *Social Cognitive and*

*Affective Neuroscience*, *6*(4), 486–494. https://doi.org/10.1093/scan/nsq068

Preston, S. D., & De Waal, F. B. M. (2002). Empathy: Its ultimate and proximate bases. *Behavioral and*

*Brain Sciences*, *25*(1), 1–20. https://doi.org/10.1017/S0140525X02000018

Prinsen, J., & Alaerts, K. (2019). Eye contact enhances interpersonal motor resonance: Comparing

video stimuli to a live two-person action context. *Social Cognitive and Affective Neuroscience*,

*14*(9), 967–976. https://doi.org/10.1093/scan/nsz064

Qaiser, J., Leonhardt, N. D., Le, B. M., Gordon, A. M., Impett, E. A., & Stellar, J. E. (2023). Shared

Hearts and Minds: Physiological Synchrony During Empathy. *Affective Science*, *4*(4), 711–721.

https://doi.org/10.1007/s42761-023-00210-4

Quigley, K. S., Gianaros, P. J., Norman, G. J., Jennings, J. R., Berntson, G. G., & de Geus, E. J. C.

(2024). Publication guidelines for human heart rate and heart rate variability studies in

psychophysiology—Part 1: Physiological underpinnings and foundations of measurement.

*Psychophysiology*, *61*(9), e14604. https://doi.org/10.1111/psyp.14604

R Core Team. (2020). *R: A Language and Environment for Statistical Computing* (Version 4.0.3)

[Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/

Raftery, A. E. (1995). Bayesian Model Selection in Social Research. *Sociological Methodology*, *25*, 111.

https://doi.org/10.2307/271063

Reddan, M. C., Young, H., Falkner, J., López-Solà, M., & Wager, T. D. (2020). Touch and social support

influence interpersonal synchrony and pain. *Social Cognitive and Affective Neuroscience*,

*15*(10), 1064–1075. https://doi.org/10.1093/scan/nsaa048

Riecansky, I., & Lamm, C. (2019). The Role of Sensorimotor Processes in Pain Empathy. *Brain*

*Topography*, *32*(6), 965–976. https://doi.org/10.1007/s10548-019-00738-4

Risko, E. F., Richardson, D. C., & Kingstone, A. (2016). Breaking the Fourth Wall of Cognitive Science:

Real-World Social Attention and the Dual Function of Gaze. *Current Directions in Psychological*

*Science*, *25*(1), 70–74. https://doi.org/10.1177/0963721415617806

Rizzolatti, G., & Craighero, L. (2004). THE MIRROR-NEURON SYSTEM. *Annual Review of*

*Neuroscience*, *27*(1), 169–192. https://doi.org/10.1146/annurev.neuro.27.070203.144230

Ruben, M. A., & Hall, J. A. (2013). "I Know Your Pain": Proximal and Distal Predictors of Pain Detection

Accuracy. *Personality and Social Psychology Bulletin*, *39*(10), 1346–1358.

https://doi.org/10.1177/0146167213493188

Rütgen, M., Seidel, E.-M., Riecansky, I., & Lamm, C. (2015). Reduction of Empathy for Pain by Placebo

Analgesia Suggests Functional Equivalence of Empathy and First-Hand Emotion Experience.

*Journal of Neuroscience*, *35*(23), 8938–8947. https://doi.org/10.1523/JNEUROSCI.3936-

14.2015

Saarela, M. V., Hlushchuk, Y., Williams, A. C. D. C., Schurmann, M., Kalso, E., & Hari, R. (2006). The

Compassionate Brain: Humans Detect Intensity of Pain from Another's Face. *Cerebral Cortex*,

*17*(1), 230–237. https://doi.org/10.1093/cercor/bhj141

Schilbach, L., Timmermans, B., Reddy, V., Costall, A., Bente, G., Schlicht, T., & Vogeley, K. (2013).

Toward a second-person neuroscience. *Behavioral and Brain Sciences*, *36*(4), 393–414.

https://doi.org/10.1017/S0140525X12000660

Schmalz, X., Biurrun Manresa, J., & Zhang, L. (2023). What is a Bayes factor? *Psychological Methods*,

*28*(3), 705–718. https://doi.org/10.1037/met0000421

Schmitz, T. W., & Johnson, S. C. (2007). Relevance to self: A brief review and framework of neural

systems underlying appraisal. *Neuroscience & Biobehavioral Reviews*, *31*(4), 585–596.

https://doi.org/10.1016/j.neubiorev.2006.12.003

Seeley, W. W. (2019). The Salience Network: A Neural System for Perceiving and Responding to

Homeostatic Demands. *The Journal of Neuroscience*, *39*(50), 9878–9882.

https://doi.org/10.1523/JNEUROSCI.1138-17.2019

Shahri, F., Zabihzadeh, A., Taqipanahi, A., Haromi, M. E., Rasouli, M., Saeidi Nik, A., & Eddy, C. M.

(2024). I understand your pain but I do not feel it: Lower affective empathy in response to others'

social pain in narcissism. *Frontiers in Psychology*, *15*, 1350133.

https://doi.org/10.3389/fpsyg.2024.1350133

Shamay-Tsoory, S. G. (2011). The Neural Bases for Empathy. *The Neuroscientist*, *17*(1), 18–24.

https://doi.org/10.1177/1073858410379268

Shimada, S., & Hiraki, K. (2006). Infant's brain responses to live and televised action. *NeuroImage*,

*32*(2), 930–939. https://doi.org/10.1016/j.neuroimage.2006.03.044

Short, J., Williams, E., & Christie, B. (1976). *The social psychology of telecommunications*. John Wiley

& Sons.

Singer, T., & Lamm, C. (2009). The Social Neuroscience of Empathy. *Annals of the New York Academy*

*of Sciences*, *1156*(1), 81–96. https://doi.org/10.1111/j.1749-6632.2009.04418.x

Singer, T., Seymour, B., O'Doherty, J., Kaube, H., Dolan, R. J., & Frith, C. D. (2004). Empathy for Pain

Involves the Affective but not Sensory Components of Pain. *Science*, *303*(5661), 1157–1162.

https://doi.org/10.1126/science.1093535

Singmann, H., & Kellen, D. (2019). An Introduction to Mixed Models for Experimental Psychology. In D.

Spieler & E. Schumacher (Eds.), *New Methods in Cognitive Psychology* (1st ed., pp. 4–31).

Routledge. https://doi.org/10.4324/9780429318405-2

Sperl, M. F. J., Panitz, C., Hermann, C., & Mueller, E. M. (2016). A pragmatic comparison of noise burst

and electric shock unconditioned stimuli for fear conditioning research with many trials.

*Psychophysiology*, *53*(9), 1352–1365. https://doi.org/10.1111/psyp.12677

Sprenger, A., Spliethoff, P., Rother, M., Machner, B., & Helmchen, C. (2020). Effects of perceptible and

imperceptible galvanic vestibular stimulation on the postural control of patients with bilateral

vestibulopathy. *Journal of Neurology*, *267*(8), 2383–2397. https://doi.org/10.1007/s00415-020-

09852-x

Stouffer, S. A., Suchman, E. A., DeVinney, L. C., Star, S. A., & Williams Jr, R. M. (1949). *The american*

*soldier: Adjustment during army life. (Studies in social psychology in world war ii), vol. 1.*

Stropahl, M., Bauer, A.-K. R., Debener, S., & Bleichner, M. G. (2018). Source-Modeling Auditory

Processes of EEG Data Using EEGLAB and Brainstorm. *Frontiers in Neuroscience*, *12*, 309.

https://doi.org/10.3389/fnins.2018.00309

Sui, J., Rotshtein, P., & Humphreys, G. W. (2013). Coupling social attention to the self forms a network

for personal significance. *Proceedings of the National Academy of Sciences*, *110*(19), 7607–

7612. https://doi.org/10.1073/pnas.1221862110

Sun, B., Zhou, L., Xiao, W., Zhao, X., Zhang, W., & Li, W. (2018). Accurate empathy of romantic

partners increases pain ratings but promotes recovery. *Psychology Research and Behavior*

*Management*, *Volume 11*, 403–410. https://doi.org/10.2147/PRBM.S177457

Svensson, S., Golubickis, M., Johnson, S., Falbén, J. K., & Macrae, C. N. (2023). Self-relevance and

the activation of attentional networks. *Quarterly Journal of Experimental Psychology*, *76*(5),

1120–1130. https://doi.org/10.1177/17470218221112238

Thorson, K. R. (2018). *Understanding the Relationship Between Empathic Accuracy and Physiological*

*Synchrony: A Theoretical and Analytic Guide with Evidence From Three Contexts* [Dissertation,

New                        York                        University].

https://www.proquest.com/openview/cec8f3d43fdef74187d0c3cdcea8b0b7/

Uddin, L. Q. (2015). Salience processing and insular cortical function and dysfunction. *Nature Reviews*

*Neuroscience*, *16*(1), 55–61. https://doi.org/10.1038/nrn3857

Van Der Molen, M. J. W., Dekkers, L. M. S., Westenberg, P. M., Van Der Veen, F. M., & Van Der Molen,

M. W. (2017). Why don't you like me? Midfrontal theta power in response to unexpected peer

rejection            feedback.            *NeuroImage*,            *146*,            474–483.

https://doi.org/10.1016/j.neuroimage.2016.08.045

Viskontas, I. V., Quiroga, R. Q., & Fried, I. (2009). Human medial temporal lobe neurons respond

preferentially to personally relevant images. *Proceedings of the National Academy of Sciences*,

*106*(50), 21329–21334. https://doi.org/10.1073/pnas.0902319106

Vossen, H. G. M., & Valkenburg, P. M. (2016). Do social media foster or curtail adolescents' empathy?

A        longitudinal        study.        *Computers        in        Human        Behavior*,        *63*,        118–124.

https://doi.org/10.1016/j.chb.2016.05.040

Wade, G. L., & Vickery, T. J. (2018). Target self-relevance speeds visual search responses but does

not        improve        search        efficiency.        *Visual        Cognition*,        *26*(8),        563–582.

https://doi.org/10.1080/13506285.2018.1520377

Weisz, E., & Zaki, J. (2018). Motivated empathy: A social neuroscience perspective. *Current Opinion in*

*Psychology, Social Neuroscience*, *24*, 67–71. https://doi.org/10.1016/j.copsyc.2018.05.005

Whitmarsh, S., Nieuwenhuis, I. L. C., Barendregt, H. P., & Jensen, O. (2011). Sensorimotor Alpha

Activity is Modulated in Response to the Observation of Pain in Others. *Frontiers in Human*

*Neuroscience*, *5*. https://doi.org/10.3389/fnhum.2011.00091

Winczewski, L. A., Bowen, J. D., & Collins, N. L. (2016). Is Empathic Accuracy Enough to Facilitate

Responsive Behavior in Dyadic Interaction? Distinguishing Ability From Motivation.

*Psychological Science*, *27*(3), 394–404. https://doi.org/10.1177/0956797615624491

Xiao, X., & Zhang, Y.-Q. (2018). A new perspective on the anterior cingulate cortex and affective pain.

*Neuroscience        &        Biobehavioral        Reviews*,        *90*,        200–211.

https://doi.org/10.1016/j.neubiorev.2018.03.022

Yang, C.-Y., Decety, J., Lee, S., Chen, C., & Cheng, Y. (2009). Gender differences in the mu rhythm

during empathy for pain: An electroencephalographic study. *Brain Research*, *1251*, 176–184.

https://doi.org/10.1016/j.brainres.2008.11.062

Zaki, J. (2014). Empathy: A motivated account. *Psychological Bulletin*, *140*(6), 1608–1647.

https://doi.org/10.1037/a0037679

Zaki, J., & Ochsner, K. (2011). Reintegrating the Study of Accuracy Into Social Cognition Research.

*Psychological Inquiry*, *22*(3), 159–182. https://doi.org/10.1080/1047840X.2011.551743

Zaki, J., & Ochsner, K. N. (2012). The neuroscience of empathy: Progress, pitfalls and promise. *Nature*

*Neuroscience*, *15*(5), 675–680. https://doi.org/10.1038/nn.3085

Zerwas, F. K., Springstein, T., Karnilowicz, H. R., Lam, P., Butler, E. A., John, O. P., & Mauss, I. B.

(2021). "I feel you": Greater linkage between friends' physiological responses and emotional

experience is associated with greater empathic accuracy. *Biological Psychology*, *161*, 108079.

https://doi.org/10.1016/j.biopsycho.2021.108079

Zhang, X., Li, P., Otieno, S. C. S. A., Li, H., & Leppänen, P. H. T. (2021). Oxytocin reduces romantic

rejection-induced pain in online speed-dating as revealed by decreased frontal-midline theta

oscillations.        *Psychoneuroendocrinology*,        *133*,        105411.

https://doi.org/10.1016/j.psyneuen.2021.105411

1323