Title: Investigating Age-Related Decline in Sensorimotor Control Using Robotic Tasks
Abbreviated title: Robotic Assessment of Motor Aging

Laura Alvarez-Hidalgo[1], Ellie Edlmann[2], Gunnar Schmidtmann[3] & Ian S. Howard[1*]
[1] School of Computing, Engineering and Mathematics, University of Plymouth, Plymouth,
Devon, United Kingdom.
[2] Peninsula Medical School (Faculty of Health), University of Plymouth, Plymouth, Devon,
United Kingdom. ellie.edlmann@plymouth.ac.uk
[3] School of Health Professions (Faculty of Health), University of Plymouth, Plymouth,
Devon, United Kingdom. gunnar.schmidtmann@plymouth.ac.uk
10
11
12 *Correspondence should be addressed to: ISH: ian.howard@plymouth.ac.uk
13 Telephone: +44 1752 586324

Statement of Financial Interests or Conflicts of Interest: The authors declare that they
have no financial, personal, or professional interests that could be construed to have
influenced the paper.

Acknowledgements:
We thank the University of Plymouth Faculty of Health's Interdisciplinary Fellow Building
Programme for financial support for this project. Financial support for this project for LAH
was provided by the Engineering and Physical Sciences Research Council (EPSRC) and
the University of Plymouth. Financial support for ISH, EE, and GS was provided by the
University of Plymouth. During the preparation of this work, LAH and ISH utilized
ChatGPT-4o for proofreading.

Keywords: Sensorimotor Control, Aging and Motor Function, Robotic Assessment, Motor
Learning, Bimanual and Unimanual Coordination.

**Abstract (351/337)**
36
37 Background:

Aging is linked to widespread changes in sensorimotor control, leading to functional decline, mobility limitations, and an increased risk of falls. Certain aspects of the neurological examination and assessment of the motor system are subjective, underscoring the need for objective and quantitative tools to evaluate sensorimotor performance.

Methods:
To address this, we designed three experimental tasks using the vBOT planar robotic manipulandum to compare sensorimotor performance in healthy young adults (<35 years) and older adults (>60 years). Unlike previous assessments, our tasks uniquely integrate simultaneous bimanual control and altered dynamic conditions to comprehensively assess sensorimotor capabilities. The first task involved a bimanual de novo motor learning paradigm in which participants coordinated both hands to control a virtual 2D arm and performed 400 center-out and out-to-center trials. The second task assessed unimanual control using the right hand to perform center-out movements to eight targets. It introduced dynamic forces simulating object manipulation during the final 200 trials. The third task was similar but applied a viscous force field in the last 200 trials. It also included a scoring mechanism rewarding brisk movements, thereby encouraging participants to perform closer to their maximum capabilities. In this task, performance was also compared between the dominant and non-dominant arms. Group comparisons and within-subject differences were evaluated using appropriate statistical analyses.

Results:
Each task effectively identified age-related differences, but the viscous resistance task proved particularly sensitive, capturing significant age-related declines in force generation and movement duration. This scoring approach likely amplified performance differences, highlighting its suitability for detecting subtle effects of aging.

Older adults generally showed slower movement, longer task completion times, reduced peak forces, and prolonged reaction times. However, individual performance varied, with some older participants performing on par with younger individuals. In the third experiment, the dominant arm showed consistently better performance than the non-dominant arm, underscoring the sensitivity of the metrics to lateralized differences in motor control.

Conclusions:
These findings suggest that robot-based tasks have potential to quantify age-related differences in sensorimotor control and may support more precise clinical assessments.

### Significance Statement (87/120)

This study utilizes a robotic manipulandum, i.e., the vBOT, to assess the effect of aging. Through precisely controlled repeatable sensorimotor tasks, we quantify movement characteristics in both younger and older adults, providing clear insights into age-related motor decline. Our approach not only highlights the magnitude and variability of functional changes with age but also establishes rigorous and reliable quantitative benchmarks that could serve as critical tools for evaluating preventative therapies. In turn, this could guide the development of interventions aimed at preserving motor performance in older age.

### News & Noteworthy (75/75)

This study introduces a novel approach to assessing how aging affects movement dynamics using a robotic system. As populations age, understanding sensorimotor decline becomes increasingly crucial. Unlike conventional assessments, our method quantifies motor control metrics with precision, revealing previously undetectable, subtle age-related changes in movement efficiency, reaction times, and force generation. This framework establishes a rigorous benchmark for evaluating age-related decline, offering valuable clinical applications and informing the design of interventions to support older adults.

### Background (548/650)

Over the past century, life expectancy has shown a steady increase. In 1920, the global average at birth was just 36.4 years; by 2020, it had more than doubled to 72.5 years. Due to this significant rise, combined with other factors, the population over the age of 60 is steadily expanding. By 2030, this age group is estimated to grow from 1 billion in 2019 to 1.4 billion, with projections suggesting it will further increase to 2.1 billion by 2050 (1,2). This growth in the aging population introduces a range of challenges, particularly in the context of health and associated age-related changes in human motor control.

With increasing life expectancy and a growing population of individuals over 60 years of age, it is important to investigate quantifiable sensorimotor indicators that signal the onset of age-related decline in healthy older adults. Identifying and measuring these early markers can equip health professionals with precise, objective, and reliable tools to assess sensorimotor function, decline and the extent of movement improvements resulting from therapy.

Pioneering work using robotic manipulanda, such as the MIT-Manus (3) and Kinarm (4) devices, has significantly advanced the quantitative assessment of sensorimotor performance during upper-limb movements. These robotic systems precisely measure and control limb dynamics and kinematics, enabling detailed examination of motor control strategies, sensory integration, and adaptation. The MIT-Manus has been instrumental in characterizing impaired motor function post-stroke, revealing deficits in movement smoothness, accuracy, and force control, which correlate closely with clinical evaluations of motor impairment (5,6). Similarly, the Kinarm robotic system has facilitated investigations into sensorimotor integration and proprioception, allowing researchers to isolate and quantify specific deficits in motor coordination and sensory processing after

121 neurological injuries or disorders (7). Robotic manipulanda provide precise and objective
122 measurements, overcoming limitations of subjectivity and limited sensitivity inherent in
123 traditional assessments.
124
Robotic manipulanda are extensively utilized to assess motor deficits associated with
aging and various neurological conditions, providing objective, sensitive metrics beyond
traditional clinical assessments. The high precision and repeatability of robotic
measurements allow the early detection of subtle impairments in motor performance,
which are characteristic of aging. Studies employing the Kinarm, for example, have
identified age-related declines in proprioception, reaction time, and motor coordination,
highlighting the subtle sensorimotor changes that precede overt functional deficits in older
adults (8,9). Additionally, robotic assessments have been pivotal in evaluating the
progression and severity of conditions like Parkinson's disease and multiple sclerosis,
facilitating early diagnosis, monitoring disease progression, and assessing therapeutic
efficacy with greater accuracy and reliability (10,11).
136
In the current study, we developed a set of experiments to systematically investigate age-
related differences in sensorimotor control of the upper limb by employing a set of
targeted experimental tasks using the vBOT robotic manipulandum (12). Specifically, we
compared performance between healthy young adults (under 35 years) and older adults
(over 60 years) across three distinct robotic tasks designed to assess both bimanual and
unimanual sensorimotor capabilities. The tasks included coordinating both hands in a
novel bimanual control scenario, as well as unimanual movements in altered force
environments. Through these experiments, we aim to quantify age-related changes in key
movement metrics - including movement speed, reaction time, and force output - to
establish an objective and replicable framework for evaluating sensorimotor decline,
thereby supporting potential improvements in clinical assessments targeting motor
deficits associated with aging.
149

**Methods**

To develop assessments suitable for future participants in a clinical setting, it was
necessary to limit the duration of the tasks they would be required to perform. To ensure
participants were not fatigued by the assessment procedure, we restricted the overall
testing time to 30 minutes per assessment. Since the current study involves investigating
the appropriateness of the assessments we were developing, we ran two experimental
sessions per participant, although we envisage that running a single test session would
be the goal in the future if the assessments are used for clinical purposes rather than
research.
159
Participants: We recruited a total of 32 participants, consisting of 16 young adults and 16
older adults. Participants were screened to exclude individuals with a history of
neurological conditions, musculoskeletal disorders, or significant visual impairments that
could affect performance. Almost all participants took part in two separate experimental
sessions. Half of the young adults and half of the older adults were randomly assigned to
Experiments 1 and 2. Participants first completed Experiment 1, followed by Experiment

2. The remaining participants were assigned to Experiment 3. All but one of them completed it twice: once using the right arm and once using the left.

All participants were naïve to the objectives of the experiments and provided written informed consent prior to participation. Right-handedness was assessed using the Edinburgh Handedness Questionnaire (13), which confirmed that all participants were right-handed. All experiments followed the ethics protocol approved by the University of Plymouth's Faculty Research Ethics and Integrity Committee.

Experiments: We developed three different experiments. Experiment 1: Learning to Control a 2D Kinematic Arm, was a bimanual task that involved learning to control a two-dimensional (2D) virtual kinematic arm using both the left and right arms. Experiment 1 investigated 8 older adults (5 females and 3 males; mean age: 73.61 ± 6.24 years) and 8 younger adults (4 females and 4 males; mean age: 28.02 ± 2.92 years).

Experiment 2: Handle Mass Center-Out Movement, involved making unimanual right-handed point-to-point movements, initially in an unloaded null-field condition and later with a simulated mass at the handle. In Experiment 2, performance was investigated mainly reusing the same participants as in Experiment 1. It involved 8 older adults (5 females and 3 males; mean age: 73.61 ± 6.24 years) and 8 younger adults (3 females and 5 males; mean age: 28.69 ± 3.01 years).

Experiment 3: Handle Viscous Resistance Center-Out Movement, involved making unimanual point-to-point movements, initially in an unloaded null-field condition and later with simulated viscosity at the handle. In Experiment 3, almost all participants ran the task twice, once with each hand. The cohort used to investigate the effect of aging on right-hand performance included 8 older adults (4 females and 4 males; mean age: 69.09 ± 2.97 years) and 7 younger adults (4 females and 3 males; mean age: 22.42 ± 2.61 years). Left-hand performance was examined in all 7 younger adults and a smaller subset of 7 older participants, since one older participant was not available (leaving 4 females and 3 males; mean age: 68.35 ± 2.27 years).

**Apparatus:** Fig. 1 illustrates the experimental setup used in this study to analyze upper limb reaching movements. Participants interacted with a pair of vBOT robotic manipulanda (12). Each was equipped with a robotic handle that incorporated an activation switch, which participants needed to press during a trial to activate the controller and allow the experiment to proceed. During the experiment, participants were seated in a comfortable, cushioned chair and secured with two shoulder straps to reduce body movement. For the unimanual experiments, a participant's relevant forearm rested on an air sled, ensuring low-friction movement constrained to a 2D plane. For the bimanual experiments, both forearms were supported by air sleds. Throughout all experiments, participants could not directly observe their arms or hands; instead, veridical visual feedback was provided using a 2D virtual reality system, which displayed visual elements such as circles representing hand positions, target locations, and start points within the workspace.

Figure 1. Schematic Illustration of the vBOT Robotic Manipulandum. **A:** Plan view of the vBOT robotic manipulandum setup. **B:** Side view of the vBOT robotic manipulandum setup.

Handle positions were measured using optical motor encoders, sampled at 1000 Hz, while torque-controlled motors generated endpoint forces. A force transducer (Nano 25; ATI), mounted beneath the left handle, measured the forces applied by participants during unimanual trials. These force signals were low-pass filtered at 500 Hz using 4th-order analog Bessel filters prior to digitization. Force and position data were stored for offline MATLAB analysis.

## Experiment 1: Learning to Control a 2D Kinematic Arm

In this experiment, participants were asked to perform movements by controlling a simulated two-link (2D) planar arm, as used in a study to investigate *de novo* learning (14). The experimental setup and design are shown in Fig. 2.

**Figure 2**. **Experiment 1: Learning to Control a 2D Kinematic Arm. A:** Experimental setup. Eight pink target points were arranged in a circular pattern at specific angular positions. The pink center point represents the starting position during center-out trials, and the target during out-to-center trials. A red cursor indicates the endpoint of a virtual 2D robotic arm with its links displayed in yellow. **B:** Dual-arm control mechanism. The left hand controls the joint angle $\theta_1$, and the right hand controls the joint angle $\theta_2$. **C:** The robotic arm's reachable workspace is shown in black, illustrating the range of motion for the virtual arm. **D:** Trial design. Each participant completed a total of 400 trials during the null-field phase.

### Virtual 2D Arm

Participants interacted with a simulated two-link robotic arm displayed in yellow, whose endpoint was represented by a red cursor. As shown in Fig. 2A, eight possible pink target points (radius: 1.0 cm) were arranged in a circular pattern at a radius of 10.8 cm from a central pink point (radius: 1.0 cm) and located at workspace origin (0,0). Targets were positioned at 0°, 45°, 90°, 135°, 180°, 225°, 270°, and 315°. The center point also served as the starting position for center-out trials and as the target for out-center trials.

Participants controlled the virtual 2D arm by grasping two robotic handles, each linked to a specific joint angle of the virtual 2D arm. The left handle controlled joint angle $\theta_1$, and the right handle controlled joint angle $\theta_2$, as depicted in Fig. 2B. Vertical hand movements adjusted these joint angles: upward movements decreased the angles, while downward movements increased them. To constrain hand movements to the y-axis, participants operated within two vertical channels (15).

### Trials Procedure

The experiment involved two types of movements: center-out movements and out-to-center movements. In center-out movement trials, participants started at the central position and moved the red cursor to one of the eight peripheral targets. In contrast, in out-to-center movement trials, participants started at a peripheral target and moved the red cursor back to the central position. Each trial began with the participant pressing both activation switches on the back of the robotic handles.

At the start of each center-out trial, the potential peripheral targets were first displayed in pink, while the central position appeared in grey. Participants then positioned the arm's endpoint cursor at the starting location. Once the cursor remained stationary at this location for 500 ms, the vBOT emitted an auditory cue (a beep). Simultaneously, one of the eight target locations was selected pseudorandomly and turned yellow, signaling the direction in which the participant should perform the trial. The virtual arm also appeared at this point. As the cursor moved out of the start location, the central grey point disappeared. The trial ended when the red cursor reached the target and stopped moving, at which point the virtual arm disappeared as well.

After completion of a center-out movement trial, participants performed an out-to-center trial, which returned the cursor to the central position. The procedure was similar to the

274    center-out trials, except the movement began at a peripheral location, and the central
275    position turned yellow to indicate it as the target.
276
277    Experimental Block Design: As illustrated in Fig. 2D, the experiment consisted of 400
278    trials, divided into 25 blocks of 16 trials each. Each block included 8 center-out trials and
279    8 out-to-center trials.
280
281    Mechanical Channels: Participants' hand movements were constrained to two distinct
282    vertical channels, each corresponding to one hand. These channels were aligned along
283    the y-axis with a length of 30 cm (chanLen). Both channels were offset along the x-axis
284    by ±10 cm from the center of the workspace, with the midpoints of the sliders located at
285    (−10, 0) for the right hand and (+10, 0) for the left hand.
286
287    These linear channels were characterized by a stiffness value of 4,000 Nm⁻¹ ($k_{chan}$) and
288    a damping coefficient of 30 Nm⁻¹s ($b_{chan}$). The forces experienced by each hand within
289    the channels were determined by the following equation, which relates the force to
290    position and velocity:
291

292
$$\begin{bmatrix} F_{xh} \\ F_{yh} \end{bmatrix} = k_{chan} \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_h \\ y_h \end{bmatrix} + b_{chan} \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_h \\ \dot{y}_h \end{bmatrix} \tag{1}$$

293
294
295    Control of the Simulated Arm
296    The vertical position ($y$) of each handle was linearly scaled to map to the corresponding
297    joint angle ($\theta$). This mapping was defined as:
298

299
$$\theta = \theta_{min} + (\theta_{max} - \theta_{min}) \frac{\left( dir_{slider} \cdot y + \frac{\text{chanLen}}{2} \right)}{\text{chanLen}} \tag{2}$$

For the left handle, the vertical position ($y_{h1}$) was mapped to a joint angle ($\theta_1$), within the
range of $\theta_{min} = 0$ radians and $\theta_{max} = \pi$ radians. Similarly, the vertical position ($y_{h2}$) of the
right handle was mapped to a joint angle ($\theta_2$), within the range of $\theta_{min} - \pi$ radians and
$\theta_{max} = 0$ radians.

$$\theta_1 = \frac{(15 - y_{h1})}{30}\pi \tag{3}$$    306   $$\theta_2 = \frac{(15 - y_{h2})}{30}\pi - \pi \tag{4}$$

This allowed the y-position of the handles to be effectively converted into joint angles
within their respective ranges.
310
311    Simulating Forward Kinematics

To compute the endpoint and visualize the configuration of the 2D arm, its forward kinematics were used. The simulated arm link lengths $l_1$ and $l_2$ were both 16.2 cm long. By employing trigonometric analysis, the positions of the elbow and end effector in Cartesian coordinates can be described using matrix equations:

$$\begin{bmatrix} x_{elbow} \\ y_{elbow} \end{bmatrix} = \begin{bmatrix} l_1 \cos(\theta_1) \\ l_1 \sin(\theta_1) \end{bmatrix} + \begin{bmatrix} x_{shoulder} \\ y_{shoulder} \end{bmatrix} \tag{5}$$

$$\begin{bmatrix} x_{end} \\ y_{end} \end{bmatrix} = \begin{bmatrix} l_1 \cos(\theta_1) + l_2 \cos(\theta_1 + \theta_2) \\ l_1 \sin(\theta_1) + l_2 \sin(\theta_1 + \theta_2) \end{bmatrix} + \begin{bmatrix} x_{shoulder} \\ y_{shoulder} \end{bmatrix} \tag{6}$$

The shoulder position (-12, -2) was located slightly below the central start location and to its left just outside the circle on which the targets are located.

## Experiment 2: Handle Mass Center-Out Movement

This experiment investigated participants' ability to perform center-out movements under two conditions: in a null-field allowing free movement, and in a condition simulating the effect of moving a 5 kg mass located at the handle. The experimental design is summarized in Fig. 3.

**Figure 3. Experiment 2: Handle Mass Center-Out Movement. A:** Experimental setup. Eight yellow targets were arranged at specific angular locations. A grey center point represented the starting position, while a red cursor indicated the participant's hand

location. **B:** Experimental phases. Participants experienced either a null-field, allowing
free movement within the workspace, or the effect of a mass. The latter introduced inertial
resistance by simulating the sensation of moving a 5 kg point mass around the
workspace. **C:** Experimental conditions. The experiment consisted of 400 trials in total:
200 trials in the null-field phase, followed by 200 trials in the point mass phase.

Experimental Setup
Participants were instructed to perform center-out movements toward one of eight
possible yellow targets positioned in a circular pattern around a grey center point. As
depicted in Fig. 3A, the starting position was represented by a central grey circle (radius:
1.25 cm) at the center of the workspace (0,0), and a red cursor (radius: 0.5 cm) displayed
the hand's position in the workspace. Additionally, eight yellow target circles (radius: 0.5
cm) were arranged in a circular pattern at a radius of 12 cm from the center, located at
angular positions of 0º, 45º, 90º, 135º, 180º, 225º, 270º, and 315º. Only one target
appeared on the screen at a time.

Participants controlled the red cursor by holding the left vBOT robotic handle. At the start
of each trial, participants viewed an empty screen and grasped the left vBOT robotic
handle. The vBOT guided their hand to the designated center position. After remaining
stationary at this location for 300 ms, an auditory beep signaled the beginning of the trial.
At the same time, the hand position cursor, as well as a single yellow target, appeared on
the screen. At this point, participants were required to move their hand from the center
position toward the displayed target. The trial concluded when the participant successfully
reached the target.

Experimental Block Design
Movements were undertaken under two conditions. During this center-out movement,
participants encountered either a null-field or a 5 kg mass (see Fig. 3B). Each condition
consisted of 400 trials, with the first 200 conducted in a null-field phase and the remaining
200 in an endpoint mass phase. Approximately midway through the experimental
condition, participants were given a 10-second rest. However, participants could take a
break at any time simply by releasing the activation switch. The timeline of trials is shown
in Fig. 3C.

Simulating Mass
To ensure a stable simulation, moving the mass was modeled as being connected to the
hand cursor via a stiff spring. Additionally, viscous damping was incorporated in the
simulation, to enhance stability. At the start of the simulation, the mass' position is
initialized to match the current location of the hand cursor. The mass' velocity,
acceleration, and the forces acting on it are all set to zero.

373
$$\begin{bmatrix} x_m \\ y_m \end{bmatrix} = \begin{bmatrix} x_{cur} \\ y_{cur} \end{bmatrix} \tag{7}$$

$$\begin{bmatrix} v_{xm} \\ v_{ym} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \tag{8}$$

$$\begin{bmatrix} a_{xm} \\ a_{ym} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \tag{9}$$

$$\begin{bmatrix} F_{xm} \\ F_{ym} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \tag{10}$$

381 Within the main simulation update loop, which operates at a nominal frequency of 1 kHz,
382 the motion of the mass is updated in real time. At each time step, the position of the mass
383 is updated using Euler integration based on its velocity:

$$\begin{bmatrix} x_m \\ y_m \end{bmatrix} = \begin{bmatrix} x_m \\ y_m \end{bmatrix} + \begin{bmatrix} v_{xm} \\ v_{ym} \end{bmatrix} Dt \tag{11}$$

385 Here, Dt represents the timestep since the last update, typically 1 ms. The mass'
386 acceleration is then calculated using the following equation:

$$\begin{bmatrix} a_m \\ a_m \end{bmatrix} = \frac{1}{m} \begin{bmatrix} F_{xm} \\ F_{ym} \end{bmatrix} \tag{12}$$

388 where m = 5 kg is the mass and $\begin{bmatrix} F_{xm} \\ F_{ym} \end{bmatrix}$ is the force acting on it. Using Euler integration,

389 the velocity of the mass is updated based on its acceleration:

$$\begin{bmatrix} v_{xm} \\ v_{ym} \end{bmatrix} = \begin{bmatrix} v_{xm} \\ v_{ym} \end{bmatrix} + \begin{bmatrix} a_{xm} \\ a_{ym} \end{bmatrix} Dt \tag{13}$$

391 The force acting on the mass is determined by the extension of the spring, which connects
392 the mass to the endpoint of the arm. The spring has a stiffness constant of $k_{mass}$ = 2000
393 N/m:

$$\begin{bmatrix} F_{xm} \\ F_{ym} \end{bmatrix} = -k_{mass} \left( \begin{bmatrix} x_m \\ y_m \end{bmatrix} - \begin{bmatrix} x_{end} \\ y_{end} \end{bmatrix} \right) \tag{14}$$

395 The endpoint velocity, which is directly available from the handle velocity, is used to
396 calculate an additional viscous damping force for further stability enhancement. The
397 viscous damping force is given by:

$$\begin{bmatrix} F_{xdamp} \\ F_{ydamp} \end{bmatrix} = -b_{end} \begin{bmatrix} \dot{x}_{end} \\ \dot{y}_{end} \end{bmatrix} \tag{15}$$

399   Here, the damping viscosity coefficient ($b_{end}$) is 10 Nm$^{-1}$s. Finally, the spring force (Eq.
400   14) and damping force (Eq. 15) are combined to compute the total force experienced at
401   the handle.
402
403

Experiment 3: Handle Viscous Resistance Center-Out Movement
This experiment was similar to Experiment 2. However, after the initial 200 trials, during
the second phase of the experiment, instead of introducing a 5 kg simulated point mass
at the handle, a viscous force field that resisted hand movement was introduced. This
resistance was proportional to the velocity of movement, producing a sensation similar to
moving underwater. The experimental design is illustrated in Fig. 4.
410
411   Experiment 3 was first carried out with their right hand. To compare dominant and non-
412   dominant arm movement performance, some participants also repeated the experiment
413   with their left hands.
414

415   Viscous Force Field
416   To implement viscous resistance in the direction of movement, we used the force
417   equation:

418
$$F = -\eta v \tag{16}$$

419
420   Where the viscous-field constant $\eta$ has a value of 0.3 N.s/m.

421
422

**Figure 4. Experiment 3: Handle Viscous Resistance Center-Out Movement. A:** Experimental setup. Eight yellow targets were arranged at specific angular locations. A grey center point represented the starting position, while a red cursor displayed the participant's hand location. **B:** Experimental phases. Participants experienced either a null-field, allowing free movement within the workspace, or a viscous-field, which introduced resistance. **C:** Right-Hand Experimental Condition. Participants performed the experiment with their right arm for all 400 trials: 200 trials in a null-field phase, followed by 200 trials in a viscous-field phase. **D:** Left-Hand Experimental Condition. Participants repeated the same experimental procedure performed with their right hand, but this time using their left hand.

Experimental Block Design

Each condition consisted of 400 trials, with the first 200 conducted in a null-field phase and the remaining 200 in a viscous force-field phase. Approximately midway through the experimental condition, participants were given a 10-second rest period. Participants could also take a break at any time simply by releasing the activation switch. The timeline of trials to assess right-hand performance is shown in Fig. 4C. In this experiment, participants received a score reflecting the number of times they achieved a brisk movement, thereby encouraging them to move faster than they otherwise might have done.

Dominant verses Non-Dominant Arm Usage

To assess how dominant verses non-dominant arm use affects performance on movement characteristics and performance, each participant performed the experiment twice: once using their right arm and once using their left arm (refer to Fig. 4D). This also provided an effective means to examine the robustness of the movement analyses, as it permitted intra-participant comparisons that were unaffected by the different levels of performance that occur across participants.

**Data Analysis**

The experimental data was processed offline using MATLAB R2024a. For the de novo arm experiment (Experiment 1), we examined the trajectory of the cursor, which corresponded to the endpoint of the virtual arm. For the null-mass experiment (Experiment 2) and the null-viscous experiment (Experiment 3), we also examined the trajectory of the cursor, which corresponded to the location of the robot handle.

Each trial began with an auditory cue (a beep). Movement onset was defined as the moment the hand exceeded a distance of 1.25 cm from the starting position and its speed surpassed 5 cm/s. Movement termination was defined as the moment the hand entered within 1.25 cm of the target and its speed dropped below 5 cm/s.

### Hand Trajectory Plots

For the de novo arm experiment (Experiment 1) we plot the mean ± standard error (SE) of movement trajectories over the first and final blocks of trials to provide an intuitive visualization of participants' movement behavior.

### Performance Metrics

We quantified kinematic accuracy with six metrics:

Absolute Maximum Perpendicular Error (AMPE): For each trial, we computed the absolute maximum perpendicular deviation of the movement trajectory from the ideal straight line connecting the start and target cursor locations in extrinsic space. By taking the absolute value, this metric avoids positive and negative deviations from canceling each other out across trials and provides a clear overall measure of how accurately participants followed a straight path from the starting point to the target.

**Movement Duration:** The time taken to complete a movement was calculated as the difference between the movement onset and termination times.

**Movement Length:** The total path length was determined by summing the distances between consecutive (x, y) cursor positions along the trajectory, from movement start to movement end.

**Reaction Time:** The time elapsed between the cue to start moving and actual movement onset.

**Maximum Speed:** The highest speed attained during the movement.

**Maximum Force:** The peak force exerted in the direction of movement.

### Statistical Analysis

Statistical analyses were performed using JASP software (16). When appropriate, Mauchly's test was applied to assess the assumption of sphericity. If this assumption was violated, Greenhouse-Geisser corrections were used to adjust the degrees of freedom. The significance level ($\alpha$) was set at 0.05 for all tests. Omega squared ($\omega^2$) was reported as a measure of effect size, indicating the proportion of variance in the dependent variable explained by the independent variables.

Repeated measures ANOVAs were used to examine differences in key metrics across critical phases within each experimental condition. When a significant main effect was observed, where appropriate we proceeded with post hoc comparisons using the Holm-Bonferroni correction. Bonferroni-corrected p-values are indicated as pBonf.

To compare metrics across the conditions, of age (younger verses older participants and the use of the right verses left hand), separate ANOVAs were conducted on the initial

508  exposure metrics (defined as the mean of the first 25 exposure blocks) and the final
509  exposure metrics (mean of the last 25 exposure blocks).
510
511  **Results**
512  Experiment 1: Learning to Control a 2D Kinematic Arm.
513
514  The first experiment was a novel bimanual kinematic arm task in which participants
515  controlled the endpoint of a simulated two-dimensional (2D) arm to complete a center-out
516  task. Their left- and right-hand positions, restricted to vertical movements, determined the
517  arm's joint angles. The experiment consisted of a total of 400 trials.
518
Fig. 5 shows the mean and standard error (SE) of both outwards and inwards endpoint
movement trajectories of the virtual arm at two critical stages, for both younger and older
participants. Panel A illustrates the first 16 trials of the younger participants, where
movements are highly erratic, reflecting exploratory behavior as they attempt to gain
control of the arm's endpoint. By the end of the experiment, as shown in Panel B,
trajectories become much straighter, and deviations from the mean direction are markedly
reduced. Similar improvements are observed in older participants. Panel C displays the
first 16 trials for the older group, also showing highly erratic movements. However, by the
end of the 400 trials, their performance improves significantly as well. Qualitatively,
younger participants appear to perform slightly better than older participants, in terms of
the straightness of their movements. We also note that for both sets of participants, inward
movements were straighter than outward movements, no doubt arising from the fact that
it was easier to move back to the central location than outwards to one of many peripheral
targets.
533

534
535

**Figure 5.** Plots showing the mean ± standard error (SE) of virtual arm endpoint trajectories over 16 trials for outward and inward movements: A: Under-35 years, initial trials (1 - 16); B: Under-35 years, final trials (385–400); C: Over-60 years, initial trials (1–16); D: Over-60 years, final trials (385–400).

Behavioral trends related to age were evident in the kinematic arm task data. Younger adults, in general, took less time to complete each trial and made straighter movements in extrinsic space. These quantitative indicators of motor performance across age groups may serve as a means for evaluating the effectiveness of future rehabilitation strategies.

Fig. 6 displays movement metrics used to quantify the performance of the endpoint movement of the virtual arm. Each metric represents the average of 16 consecutive trials and is plotted against block pairs. Across all metrics, there is a general tendency for performance to improve with trial number. Panel A shows that movement duration starts off high for both younger and older participants but decreases considerably by the end of the 400 trials. Younger participants exhibit consistently shorter movement durations than older participants. Panel B presents the absolute maximum perpendicular error (AMPE) across block pairs, with a general downward trend. Younger participants tend to show slightly lower AMPE values compared to older participants. Panel C depicts reaction time - defined as the interval between the cued start of a trial and movement initiation. Younger participants consistently demonstrate shorter reaction times. Panel D shows the maximum movement speed as a function of block pair, revealing a slight overall increase with practice. Younger participants move faster than older participants on average. Panel E displays trajectory path length, which is consistently shorter in younger participants across all trial blocks.

**Comparisons of results within age group and conditions**

Repeated-measures ANOVAs were used to examine whether there were any significant variations in the measured metrics across the initial and final two blocks (32 trials) for both younger and older participants.

Both younger and older participants showed significant changes over time in Duration, AMPE, Reaction Time, and Path Length, with large effect sizes across all four metrics. Reaction Time showed particularly large effects, especially in older adults ($\omega^2 = 0.664$). No significant changes were observed in Peak Speed for either group. Detailed results are summarized in Table 1.

**Table 1.** Repeated-measures ANOVA results for each age group across the entire experiment: Separate repeated-measures ANOVAs were conducted for each performance metric to assess overall changes across the full duration of the task within each age group. Reported values include F-statistics, p-values, and effect sizes ($\omega^2$). Significant p-values are shown in bold.

| AGE GROUP | METRIC | F | P | $\omega^2$ |
|---|---|---|---|---|
| Younger | Duration | 35.808 | **< 0.001** | 0.643 |

| | Metric | F | P | ω² |
|---|---|---|---|---|
| | AMPE | 20.611 | **0.003** | 0.460 |
| | Reaction Time | 22.820 | **0.002** | 0.530 |
| | Peak Speed | 0.014 | 0.909 | 0.000 |
| | Path Length | 25.384 | **0.001** | 0.546 |
| Older | Duration | 44.317 | **< 0.001** | 0.741 |
| | AMPE | 39.080 | **< 0.001** | 0.546 |
| | Reaction Time | 107.030 | **< 0.001** | 0.664 |
| | Peak Speed | 0.121 | 0.738 | 0.000 |
| | Path Length | 19.747 | **0.003** | 0.482 |

**Summary within age group and conditions**

The repeated-measures ANOVAs revealed some significant improvements in task performance across training for both younger and older participants. Specifically, Duration, Absolute Mean Position Error (AMPE), Reaction Time, and Path Length all significantly decreased from the initial to the final blocks of training, indicating enhanced movement efficiency and response preparation over time. In the younger group, all four metrics showed marked improvements with large effect sizes, such as $\omega^2 = 0.643$ for Duration. Similarly, the older group demonstrated significant gains, with particularly strong effects observed for Reaction Time ($F = 107.303$, $p < 0.001$, $\omega^2 = 0.664$). Overall, the results demonstrate clear learning effects within both age groups, with improvements across multiple performance metrics except for Peak Speed.

**Comparisons of Results Across Age Group Conditions**

To investigate age-related differences in performance, we compared average values for each metric between younger and older participants using an ANOVA, analysing the first half of trials (1–200) and the second half (201–400) separately.

During Phase 1, no significant age-related differences were observed, though Duration showed a trend toward significance with a moderate effect size ($\omega^2 = 0.131$). In Phase 2, significant differences were found in Duration ($p = 0.008$, $\omega^2 = 0.343$) and Reaction Time ($p = 0.041$, $\omega^2 = 0.202$), indicating that older participants performed slower and responded more slowly in the later stage of the task. No differences were found in AMPE, Peak Speed, or Path Length. Detailed results are summarized in Table 2.

**Table 2.** ANOVA results comparing younger and older participants within each experimental phase: Separate one-way ANOVAs were conducted for each performance metric during Phase 1 (P1, trials 1–200) and Phase 2 (P2, trials 201–400). Reported values include F-statistics, p-values, and effect sizes ($\omega^2$). Significant p-values are shown in bold.

| PHASE OF EXPERIMENT | METRIC | F | P | ω² |
|---|---|---|---|---|
| P1 | Duration | 3.419 | 0.086 | 0.131 |

| | | | | |
|---|---|---|---|---|
| | AMPE | 0.186 | 0.673 | 0.000 |
| | Reaction Time | 1.838 | 0.197 | 0.050 |
| | Peak Speed | 0.679 | 0.424 | 0.000 |
| | Path Length | 0.997 | 0.335 | 0.000 |
| P2 | Duration | 9.367 | **0.008** | 0.343 |
| | AMPE | 0.296 | 0.595 | 0.000 |
| | Reaction Time | 5.038 | **0.041** | 0.202 |
| | Peak Speed | 0.565 | 0.465 | 0.000 |
| | Path Length | 0.980 | 0.339 | 0.000 |

**Summary cross age**

In the first half, there were no significant differences between age groups for any of the measures. In the second half of trials, only movement Duration and Reaction Time are significantly shorter in the younger participants (p = 0.008 and p = 0.041, respectively). No other metrics showed a statistically significant difference between the two age groups.

**Figure 6.** Movement metrics quantifying the trajectory at the endpoint of the virtual arm, plotted against block pairs. Younger participants are plotted in blue, and older participants are plotted in green. **A:** Duration of movement. **B:** Absolute maximum perpendicular error (AMPE). **C:** Movement reaction time. **D:** Maximum trajectory speed. **E:** Trajectory path length.

## Experiment 2: Handle Mass Center-Out Movement

The second experiment was a right-hand center-out task involving two field conditions: a
null field and a simulated endpoint mass. Participants made 12 cm movements from a
central starting position to one of eight targets in each of two 200-trial phases.

The conditions were designed to investigate performance during unloaded and loaded
movements. Behavioral trends related to age were evident: younger adults typically made
faster, shorter-duration movements, generated greater peak forces in the direction of
movement, and exhibited quicker reaction times.

Fig. 7 shows the performance of younger and older participants across several metrics
quantifying hand trajectories during the null-mass center-out task. Compared to the older
participants, younger participants take less time to complete the movement, have a faster
reaction time, move with a higher peak speed, generate more force in the direction of
movement, and have shorter path lengths. However, AMPE is quite similar across the
groups.

**Comparisons of results within age group and conditions**

Repeated-measures ANOVAs were used to examine the significance of differences in the
measured metrics between pairs of blocks (16 trials each) located at the initial null
exposure, final null exposure, initial mass dynamic exposure, and final mass dynamic
exposure. Post hoc comparisons were conducted to further assess significant differences
between these phases.

Significant phase effects were observed in several performance metrics for both age
groups. Bonferroni-corrected post hoc tests showed that the introduction of the mass
dynamic ($p_{Bonf}$ 2) led to consistent performance changes in AMPE, Peak Speed, Peak Y-
Force, and Path Length for younger participants, and in Duration, AMPE, Reaction Time,
and Path Length for older participants. Adaptation across the mass dynamic phase ($p_{Bonf}$
3) was evident primarily in AMPE and Path Length in both groups. Only older participants
exhibited significant changes in Reaction Time. Path Length showed the largest effect
sizes in both groups, indicating strong sensitivity to dynamic perturbation. Detailed results
are summarized in Table 3.

**Table 3.** Repeated-measures ANOVA results for each age group across task phases:
Each metric was analyzed using repeated-measures ANOVAs to assess changes across
three task transitions: (1) from the initial to final null blocks, (2) from the final null blocks
to the introduction of the mass dynamic, and (3) from the start to the end of the mass
dynamic blocks. The table presents F-statistics, p-values, effect sizes ($\omega^2$), and
Bonferroni-corrected post hoc p-values for each comparison. Significant p-values are
shown in bold.

| AGE GROUP | METRIC | F | P | $\omega^2$ | $P_{BONF}$ (1) | $P_{BONF}$ (2) | $P_{BONF}$ (3) |
|-----------|--------|---|---|------------|----------------|----------------|----------------|

| Younger | Duration | 21.168 | **< 0.001** | 0.539 | 0.244 | **< 0.001** | 0.077 |
| | AMPE | 18.210 | **< 0.001** | 0.493 | 0.009 | **0.017** | **0.003** |
| | Reaction Time | 5.227 | **0.007** | 0.165 | 0.158 | 0.118 | 0.914 |
| | Peak Speed | 6.705 | **0.002** | 0.161 | 1.000 | **< 0.001** | 0.094 |
| | Peak Y-Force | 26.530 | **< 0.001** | 0.608 | 1.000 | **0.011** | 1.000 |
| | Path Length | 40.052 | **< 0.001** | 0.649 | 0.305 | **< 0.001** | **0.001** |
| Older | Duration | 17.697 | **< 0.001** | 0.398 | **0.030** | 0.001 | **0.006** |
| | AMPE | 9.258 | **< 0.001** | 0.384 | 0.155 | **0.009** | **0.017** |
| | Reaction Time | 6.113 | **0.004** | 0.129 | 0.947 | **0.026** | **0.003** |
| | Peak Speed | 5.860 | **0.005** | 0.058 | 0.087 | 0.124 | 1.000 |
| | Peak Y-Force | 20.825 | **< 0.001** | 0.519 | 1.000 | **0.013** | 0.663 |
| | Path Length | 55.636 | **< 0.001** | 0.776 | 0.310 | **< 0.001** | **0.001** |

**Summary within age group and conditions**

For younger participants, AMPE decreased significantly by the end of the null blocks, increased with the introduction of the mass dynamic, and decreased again by its end. Duration and Peak Speed increased significantly with the mass dynamic but did not change significantly otherwise. Peak Y-Force and Path Length also increased significantly with the mass dynamic; Path Length then decreased significantly by the end of this phase. Reaction Time showed no significant changes.

For older participants, Duration decreased significantly during the null phase, increased with the mass dynamic, and decreased again by its end. AMPE and Reaction Time both increased significantly with the mass dynamic and decreased significantly by its end. Peak Y-Force and Path Length increased significantly with the mass dynamic, and Path Length also decreased significantly by the end of this phase. Peak Speed showed no significant changes.

Compared to younger participants, older participants showed significant changes in Reaction Time and Duration across all phases, indicating a broader sensitivity to the mass dynamic. In contrast, younger participants showed no significant changes in Reaction Time and more selective effects, with AMPE showing the clearest pattern of adaptation.

**Cross-age statistical analyses**

To investigate age-related differences in performance, we compared average values for each metric between younger and older participants using an ANOVA, analysing the first half of trials (1–200) and the second half (201–400) separately.

In Phase 1, significant age-related differences were observed in Duration, Peak Speed, and Path Length (all $p < 0.05$), with moderate to large effect sizes. Differences in Reaction Time and Peak Y-Force approached significance. In Phase 2, only Reaction Time

686 reached the conventional threshold for significance (p = 0.050), with a moderate effect
687 size ($\omega^2$ = 0.184). Detailed results are summarized in Table 4.
688

689 **Table 4.** ANOVA results comparing younger and older participants within each
690 experimental phase: Separate one-way ANOVAs were conducted for each metric within
691 Phase 1 (P1, trials 1–200) and Phase 2 (P2, trials 201–400). Reported values include F-
692 statistics, p-values, and effect sizes ($\omega^2$). Significant p-values are shown in bold.
693

| PHASE OF EXPERIMENT | METRIC | F | P | $\omega^2$ |
|---|---|---|---|---|
| P1 | Duration | 5.779 | **0.031** | 0.230 |
| | AMPE | 0.643 | 0.436 | 0.000 |
| | Reaction Time | 3.767 | 0.073 | 0. 147 |
| | Peak Speed | 5.640 | **0.032** | 0.225 |
| | Peak Y-Force | 4.185 | 0.060 | 0.166 |
| | Path Length | 7.950 | **0.014** | 0.303 |
| P2 | Duration | 3.721 | 0.074 | 0.145 |
| | AMPE | 0.021 | 0.886 | 0.000 |
| | Reaction Time | 4.601 | 0.050 | 0.184 |
| | Peak Speed | 3.141 | 0.098 | 0.118 |
| | Peak Y-Force | 2.929 | 0.109 | 0.108 |
| | Path Length | 2.833 | 0.114 | 0.103 |

**Summary cross-age**

697 In the null phase movement duration is significantly shorter in the younger participants (*p*
698 = 0.031), who also exhibit higher peak speed (p=0.032) and longer path lengths (*p* =
699 0.014). No other metrics showed a statistically significant difference between the two age
700 groups.
701

702
703 **Figure 7. Movement metrics quantifying the hand trajectory during the null-mass**
704 **center-out task, plotted against blocks**. Younger participants are shown in blue, and
705 older participants in green. **A:** Movement duration. **B:** Absolute maximum perpendicular
706 error (AMPE). **C:** Reaction time. **D:** Maximum trajectory speed. **E:** Maximum force in the
707 direction of movement. **F:** Trajectory path length.

708
709
710
711
712 Experiment 3: Handle Viscous Resistance Center-Out Movement - age related
713 performance

**Figure 8.** Movement metrics quantifying the hand trajectory during the right-hand null-viscous center-out task, plotted against blocks. Younger participants are shown in blue, and older participants in green. A: Movement duration. B: Absolute maximum perpendicular error (AMPE). C: Reaction time. D: Maximum trajectory speed. E: Maximum force in the direction of movement. F: Trajectory path length.

Participants again performed a right-hand center-out task involving 12 cm movements from a central start position to one of eight targets, under two conditions: a null field and a viscous field. The task comprised two phases of 200 trials each to assess performance in unloaded and loaded movements.

Younger adults generally made faster, shorter-duration movements, produced greater peak force in the movement direction, and had quicker reaction times. The dominant arm consistently showed superior performance, highlighting the metrics' sensitivity to individual differences in motor ability.

Fig. 8 illustrates the performance of younger and older participants across multiple metrics that characterize hand trajectories during the null-viscosity center-out task. Relative to the older group, younger participants complete the movements more quickly, exhibit shorter reaction times, reach higher peak speeds, apply greater force in the movement direction, and follow shorter paths. In contrast, AMPE remains comparable between the two groups

Repeated-measures ANOVAs were used to examine the significance of differences in the measured metrics between pairs of blocks (16 trials each) located at the initial null exposure, final null exposure, initial viscous dynamic exposure, and final viscous dynamic exposure. Post hoc comparisons were conducted to further assess significant differences between these phases.

Significant phase effects were observed across most metrics in both age groups. For younger participants, Duration, AMPE, Peak Y-Force, and Path Length showed consistent changes across all task phases. Reaction Time and Peak Speed also changed significantly over time, although Reaction Time was stable during the mass introduction phase. For older participants, strong effects were observed in Duration, AMPE, Peak Y-Force, and Path Length, with adaptation particularly evident between the null and mass dynamic phases. Reaction Time did not differ significantly across phases in this group. Notably, Peak Y-Force showed extremely large effect sizes in both groups ($\omega^2 > 0.85$), indicating a strong sensitivity to the dynamic manipulation. Detailed results are summarized in Table 5.

**Table 5.** Repeated-measures ANOVA results for each age group with post hoc comparisons: ANOVAs were conducted for each performance metric within each age group across three key phases of the task. Bonferroni-corrected post hoc comparisons are reported for: (1) initial vs final null blocks, (2) final null blocks vs introduction of the mass dynamic, and (3) start vs end of the mass dynamic blocks. Significant p-values are shown in bold.

| AGE GROUP | METRIC | F | P | $\omega^2$ | $P_{BONF}$ (1) | $P_{BONF}$ (2) | $P_{BONF}$ (3) |
|---|---|---|---|---|---|---|---|
| Younger | Duration | 48.947 | **< 0.001** | 0.463 | **< 0.001** | **< 0.001** | **< 0.001** |
| | AMPE | 50.677 | **< 0.001** | 0.510 | **< 0.001** | **0.001** | **0.002** |
| | Reaction Time | 18.853 | **< 0.001** | 0.201 | **0.005** | 0.987 | **0.034** |
| | Peak Speed | 18.361 | **< 0.001** | 0.096 | 0.098 | 0.002 | **< 0.001** |
| | Peak Y-Force | 250.107 | **< 0.001** | 0.855 | 0.059 | < 0.001 | **< 0.001** |
| | Path Length | 37.046 | **< 0.001** | 0.555 | **< 0.001** | < 0.001 | **< 0.001** |
| Older | Duration | 53.759 | **< 0.001** | 0.684 | **< 0.001** | 0.141 | **< 0.001** |
| | AMPE | 31.797 | **< 0.001** | 0.550 | **< 0.001** | 0.141 | **0.027** |
| | Reaction Time | 4.953 | **0.005** | 0.063 | 0.253 | 0.868 | 0.236 |
| | Peak Speed | 11.089 | **< 0.001** | 0.116 | 1.000 | **< 0.001** | **< 0.001** |
| | Peak Y-Force | 388.621 | **< 0.001** | 0.916 | 1.000 | **< 0.001** | **< 0.001** |
| | Path Length | 17.334 | **< 0.001** | 0.401 | **0.009** | 1.000 | **0.007** |

**Summary within age and conditions**

For younger participants, Duration, AMPE, Reaction Time, and Path Length decreased significantly during the null phase. The introduction of the viscous dynamic led to significant increases in Duration, AMPE, Peak Y-Force, and Path Length, while Peak Speed decreased. By the end of the viscous phase, Duration, AMPE, Reaction Time, and Path Length had decreased again, while Peak Speed and Peak Y-Force increased.

769 For older participants, Duration and AMPE decreased significantly during the null phase.
770 Duration increased with the viscous dynamic and then decreased again, while AMPE only
771 decreased by the end. Peak Speed decreased with the dynamic and increased later, and
772 Peak Y-Force increased throughout. Path Length decreased over both phases. Reaction
773 Time showed no significant changes.
774
775 Younger participants showed more consistent adaptation across all measures, with
776 significant changes at each phase, while older participants exhibited fewer significant
777 changes, particularly in AMPE and Reaction Time, and showed less sensitivity to the
778 introduction of the viscous dynamic. This suggests younger adults adapted more readily
779 to both the removal and addition of dynamic forces.
780
**Cross age statistical analyses**

To investigate age-related differences in performance, we compared average values for
each metric between younger and older participants using an ANOVA, analysing the first
half of trials (1–200) and the second half (201–400) separately.

In Phase 1, significant age-related differences were found for Duration, Peak Speed,
Peak Y-Force, and Path Length, with moderate to large effect sizes. In Phase 2, strong
group differences were observed in Duration, Peak Speed, and Peak Y-Force, with large
effect sizes ($\omega^2 > 0.47$). No significant differences were observed for AMPE, Reaction
Time, or Path Length during this phase. Detailed results are summarized in Table 6.

**Table 6.** ANOVA results comparing younger and older participants within each
experimental phase: Separate one-way ANOVAs were conducted for each performance
metric in Phase 1 (P1, trials 1–200) and Phase 2 (P2, trials 201–400). Reported values
include F-statistics, p-values, and effect sizes ($\omega^2$). Significant p-values are shown in bold.

| PHASE OF EXPERIMENT | METRIC | F | P | $\omega^2$ |
|---|---|---|---|---|
| P1 | Duration | 6.411 | **0.025** | 0.265 |
| | AMPE | 0.940 | 0.350 | 0.000 |
| | Reaction Time | 3.224 | 0.096 | 0.129 |
| | Peak Speed | 16.316 | **0.001** | 0.505 |
| | Peak Y-Force | 9.588 | **0.009** | 0.364 |
| | Path Length | 4.703 | **0.049** | 0.198 |
| P2 | Duration | 14.343 | **0.002** | 0.471 |
| | AMPE | 0.0004 | 0.984 | 0.000 |
| | Reaction Time | 0.844 | 0.375 | 0.000 |
| | Peak Speed | 17.613 | **0.001** | 0.526 |
| | Peak Y-Force | 18.594 | **< 0.001** | 0.540 |
| | *Path Length* | 0.688 | 0.422 | 0.000 |

794
795 **Summary cross age**

796
These qualitative results partially align with statistical analyses across the two age groups
in the two phases, P1 and P2, of the experiments. Movement duration is significantly
shorter in the younger participants (*p* = 0.025 and *p* = 0.002), who also exhibit higher
peak movement speed, (p = 0.001 and p = 0.001) and greater force in the direction of
motion, (*p* = 0.009 and p < 0.001). Path length is also significantly different in P1 (p =
0.049) No other metrics showed a statistically significant difference between the two age
groups.
804
805
806 Experiment 3: Handle Viscous Resistance Center-Out Movement - Comparison Between
807 Hands.

808
809
**Figure 9.** Movement metrics quantifying the hand trajectories during the null-viscous
center-out task, plotted across blocks. The results from the experiments with participants
using their right hand are shown in blue, and the corresponding results for the left-hand
experiment are shown in green. A: Movement duration. B: Absolute maximum
perpendicular error (MPE). C: Reaction time. D: Maximum trajectory speed. E: Maximum
force in the direction of movement. F: Trajectory path length.
816
The experiment was then repeated with the left hand and a cross-hand comparison made.
Fig. 9 shows the metrics across experimental trials, plotted as the mean ± SE. The most
marked difference is that right-hand AMPE is notably lower than left-hand AMPE,

although there were also other differences. Left-hand peak speed was higher than right-
hand peak speed, and left-hand path length was shorter. Statistical analyses confirm
these observations.

To investigate differences in performance between the right and left hands, we compared
average values for each metric between metrics using a repeated measures ANOVA,
analysing the first half of trials (1 – 200) and the second half (201 – 400) separately.

In Phase 1, significant age-related differences were observed in AMPE, Reaction Time,
and Path Length, with AMPE showing the largest effect ($\omega^2 = 0.133$). In Phase 2,
AMPE, Reaction Time, and Path Length also reached significance, though with smaller
effect sizes. No significant differences were found in Peak Speed or Peak Y-Force
during either phase. Detailed results are summarized in Table 7.

**Table 7.** ANOVA results comparing younger and older participants within each
experimental phase (third dataset): One-way ANOVAs were conducted separately for
Phase 1 (P1, trials 1–200) and Phase 2 (P2, trials 201–400). The table reports F-values,
p-values, and effect sizes ($\omega^2$). Significant p-values are shown in bold.

| PHASE OF EXPERIMENT | METRIC | F | P | $\omega^2$ |
|---|---|---|---|---|
| P1 | Duration | 3.021 | 0.106 | 0.056 |
| | AMPE | 10.167 | **0.007** | 0.133 |
| | Reaction Time | 14.011 | **0.002** | 0.047 |
| | Peak Speed | 0.059 | 0.811 | 0.000 |
| | Peak Y-Force | 0.143 | 0.712 | 0.000 |
| | Path Length | 4.853 | **0.046** | 0.036 |
| P2 | Duration | 0.004 | 0.949 | 0.000 |
| | AMPE | 9.138 | **0.010** | 0.151 |
| | Reaction Time | 5.104 | **0.042** | 0.012 |
| | Peak Speed | 2.115 | 0.170 | 0.005 |
| | Peak Y-Force | 3.514 | 0.083 | 0.010 |
| | Path Length | 7.618 | **0.016** | 0.069 |

**Summary cross hands**

837
838 Right-hand AMPE values were significantly lower than left-hand values ($p = 0.007$ and
839 p=0.01) and path lengths were significantly shorter ($p = 0.046$ and p = 0.016). No other
840 metrics showed a statistically significant difference between the two hands. Reaction
841 times were significantly shorter with the left hand (p = 0.002 and p = 0.042),

842 **Discussion (1582)**
843

## The Effects of Aging on Sensorimotor Performance

In this study, we examine quantitative measures to test motor performance relevant to everyday tasks. We designed three experimental tasks using the vBOT robotic manipulandum (12), to accurately compare performance between healthy young adults (<35) and older adults (>60) in a controlled and repeatable fashion.

Across all three experiments, younger participants consistently demonstrated robust learning and adaptation in motor performance. They showed significant reductions in movement duration, Absolute Mean Position Error (AMPE), reaction time, and path length over the course of training, particularly in kinematic and viscous dynamic tasks. Younger individuals also adapted more consistently and quickly to the introduction of novel dynamics, such as added mass or viscous resistance, showing clear changes across all measured metrics. In comparison to older participants, younger adults had significantly shorter durations, higher peak speeds, greater directional force generation, and more extensive trajectory adjustments during the initial phases of exposure. These findings highlight a greater sensitivity to task dynamics and a higher capacity for motor adaptation among the younger cohort.

Analysis of right verses left-hand performance in Experiment 3 revealed significant asymmetries. AMPE and path length were both significantly lower for the right hand, indicating more accurate and efficient movements. In contrast, reaction times were significantly shorter with the left hand, possibly reflecting differences in motor planning or strategy rather than execution. No significant differences were found in peak speed or peak force between the hands. These hand-specific performance differences underscore the sensitivity and relevance of our methodology in capturing fine-grained variations in motor behavior. The ability to detect these subtle asymmetries demonstrates the utility of our approach for quantifying motor performance with high resolution and for identifying individual differences in motor control strategies.

The three experimental tasks differed significantly in their capacity to reveal age-related sensorimotor differences. Experiment 1, involving novel bimanual control of a virtual arm, highlighted initial difficulties among older adults in acquiring new motor skills, though performance improved significantly with practice. Experiment 2 assessed the management of inertial loads, clearly demonstrating older adults' reductions in peak forces and prolonged movement durations. However, Experiment 3, featuring viscous resistance and a scoring system designed to reward brisk movements, was particularly effective at differentiating younger from older adults. By explicitly encouraging participants to move quickly, the scoring mechanism appeared to push individuals closer to their limits, thereby likely exaggerating performance differences. Consequently, this task identified pronounced age-related deficits in force production, reaction time, and movement duration, suggesting its superior effectiveness for clinical assessment of subtle motor impairments associated with aging.

Our findings that movements take longer and slow down with age are consistent with previous research (17). Individual variability was found to be high, and some older adults performed similarly to younger adults. Variability may result from differences in lifestyle,

890 regular physical activity, cognitive engagement, or other unexplored genetic and
891 environmental factors (18). As individuals age, sensorimotor performance often declines
892 (18,19), impacting everyday activities such as drinking a glass of water or walking down
893 the street (20). This decline is largely due to age-related alterations in the brain and body
894 that affect motor control (21).
895
896 The decline in sensorimotor performance with aging is thought to arise from multifaceted
897 neural mechanisms, primarily involving structural and functional changes within the
898 central nervous system.
899
900 One of the most significant factors contributing to sensorimotor decline is a reduction in
901 neuroplasticity - the brain's ability to adapt and reorganize in response to new
902 experiences or environmental changes (22–24). Additionally, brain volume tends to
903 decrease over time (25,26), which can impair the regulation of motor regions, further
904 contributing to motor dysfunction (20).
905
One of the prominent neural changes is the progressive loss of gray matter volume,
particularly affecting regions such as the motor cortex, basal ganglia, and cerebellum,
which are critical for motor planning, execution, and coordination (20). Moreover, aging
leads to reduced integrity of white matter tracts, including corticospinal and sensorimotor
pathways, resulting in impaired signal conduction, reduced cortical excitability, and
increased variability in motor responses (27). A reduction in neurotransmitter availability
- particularly dopamine - further impairs motor control, reduces precision, and limits the
adaptability of motor responses, thereby contributing significantly to sensorimotor deficits
in older adults (28).
915
In addition to neural deterioration, biomechanical factors substantially exacerbate age-
related sensorimotor decline. Aging is accompanied by progressive muscle atrophy
(sarcopenia), characterized by reductions in muscle mass, strength, and contractile
speed, which significantly impair motor function and reaction times (18). There is also a
decline in proprioceptive acuity due to decreased density and sensitivity of peripheral
sensory receptors (including muscle spindles and mechanoreceptors), impairing the
ability to accurately sense joint position and movement—a capability crucial for precise
motor control (29). Furthermore, increased joint stiffness and reduced flexibility, resulting
from degenerative changes in connective tissues, limit movement range and fluidity,
compounding sensorimotor impairment in older populations (30).
926
When motor function becomes further impaired by neurological diseases such as
Parkinson's disease, stroke, or other movement disorders, abnormal movements such as
tremors (31), bradykinesia (32), rigidity (33), or dyskinesia (34) become prevalent. These
pathological conditions exaggerate the sensorimotor deficits already seen in healthy
aging, introducing additional complexity for both assessment and rehabilitation (7,35).
Such changes might explain the observed decline in sensorimotor performance.
933
Current Clinical Paradigms

To assess human motor performance, several clinical paradigms have been developed.
The clinical assessment "Timed Up & Go" (TUG) was designed as a brief evaluation of
basic mobility skills in frail and older adults (36). During the test, participants are instructed
to stand up, walk three meters to a chair, and then sit down again. This assessment has
proven to be an effective tool for evaluating mobility, walking, and balance (37–39).
Moreover, it has been demonstrated to be a reliable method for identifying adults at
increased risk of falls (40).
942
For the upper body, there has been continuous research aimed at developing tests that
evaluate multiple aspects of motor function. Researchers have specifically focused on
creating clinical assessments for upper-limb motor abilities. To evaluate upper-limb motor
function, researchers introduced the Wolf Motor Function Test (WMFT) (41), in which
participants perform tasks that simulate daily activities, such as lifting or picking up
objects. This test is widely used in hospitals, particularly for neurorehabilitation purposes
(35,42,43). Other well-known assessments include the Action Research Arm Test
(ARAT), which evaluates post-stroke patients by measuring grasp, grip, pinch, and other
motor tasks (44). Another commonly used assessment is the Jebsen–Taylor Hand
Function Test (JTHFT), also designed for post-stroke patients. This test is sensitive to
changes over time, making it an effective tool for monitoring recovery (45).
954
955 Some tests assess both upper- and lower-limb movement. One example is the Fugl–
956 Meyer Assessment (FMA), which is typically used to evaluate individuals with stroke or
957 neurological impairments. It assesses balance and coordination in both the upper and
958 lower extremities and is widely used in research and clinical settings (46).
959
960 Robotic Manipulanda Assessment
961 The experimental results presented here demonstrate that the vBOT system, in
962 conjunction with the experimental procedures, can detect subtle changes in performance
963 caused by age.
964
965 Previous research employing robotic manipulanda to assess sensorimotor ability has
966 predominantly used standardized upper-limb reaching tasks designed to objectively
967 evaluate specific aspects of motor and sensory function. Common movement paradigms
968 include:
969
970 Center-out reaching tasks: Participants typically initiate movements from a central starting
971 position to peripheral targets arranged radially around the starting point. These tasks
972 assess motor accuracy, reaction time, movement speed, trajectory smoothness, and
973 force control (10,47).
974
975 Visuomotor adaptation paradigms: Participants perform reaching movements under
976 altered visual feedback (e.g., rotated cursor) or novel force environments (e.g., viscous,
977 inertial, or perturbation fields). Such tasks evaluate sensory integration, adaptation,
978 learning rates, and the ability to generalize learned sensorimotor mappings (4,48,49).
979

980 Bimanual coordination tasks: Tasks requiring simultaneous or coordinated movements of
981 both arms have been used to investigate interlimb coordination, sensorimotor integration,
982 and neural control mechanisms underlying complex motor behavior (50,51).
983
984 Proprioceptive assessment paradigms: Robotic manipulanda have been used to
985 passively move a participant's limb to specific positions, after which participants are asked
986 to reproduce or discriminate limb positions. These paradigms quantify proprioceptive
987 acuity, position sense, and kinesthetic deficits in both neurological disorders and aging
988 populations (7,8).
989
990 Force generation and control tasks: Participants are asked to exert specified forces or
991 match force profiles, either statically or dynamically. These tasks measure maximum
992 voluntary force, accuracy of force modulation, and motor deficits related to muscle
993 weakness or impaired neuromuscular control (11).
994
Collectively, these paradigms enable detailed, quantitative assessments of various
components of sensorimotor function, providing insights into motor control deficits,
adaptation processes, and sensory impairments associated with aging, neurological
disease, or injury. However, in cases of severe movement disruption—where patients
may have difficulty initiating movement or are unable to perform certain tasks—additional
or alternative metrics may be necessary to accurately assess functional capacity and
impairment.
1002
Compared to widely-used clinical assessments such as WMFT or ARAT, the robotic
approach provides finer resolution and greater sensitivity, especially in detecting subtle
deficits or tracking small changes over short intervals Clinically, the proposed robotic
tasks could be implemented in outpatient rehabilitation settings, allowing practitioners to
routinely quantify subtle sensorimotor deficits, monitor progression, and evaluate
treatment efficacy.
1009
Emerging methods using robotic manipulanda, as utilized in this study, offer highly
sensitive, quantitative assessments capable of detecting subtle impairments and
characterizing abnormal movement patterns (47). Future developments in robotic
technologies combined with machine learning algorithms could further enhance
diagnostic accuracy, enable precise tracking of disease progression, and allow
personalized rehabilitation interventions tailored to the individual's specific sensorimotor
impairments (11).
1017
Conclusions (201)
This study demonstrates that robotic manipulanda, specifically the vBOT (12) can be used
to sensitively and quantitatively assess age-related differences in sensorimotor control.
Across three experimental tasks, older adults consistently exhibited deficits in movement
timing, force generation, and reaction speed, although some individual variability was
observed. The use of both unimanual and bimanual paradigms enabled detailed
characterization of performance, including lateralized differences between dominant and
non-dominant limbs.

Our results reinforce the growing consensus that robotic systems can complement traditional clinical assessments by providing objective, high-resolution data on motor function. Such tools offer considerable promise for early detection of motor decline, tracking disease progression, and the development of rehabilitation interventions.

We note that our proof-of-concept study involved participants who were physically fit, particularly among the older participant group. This represents a sample skewed towards higher-functioning individuals. As a result, the observed differences between younger and older adults were probably smaller than those that would be seen in a clinical setting with less able patients. Obtaining more representative participant samples will be a priority in future research. Further studies should also include more diverse participant groups and explore how robotic assessments can be integrated into clinical workflows for both age-related and neurological conditions.

## Supplementary Material
The supplementary material presents the experimental data in an alternative format.

## Disclosures
The authors declare that they have no financial, personal, or professional conflicts of interest that may have influenced the content of this paper.

## Author Contributions
LAH, ISH, GS, and EE conceived the study. LAH and ISH implemented the study, and LAH conducted data collection. Data analysis was carried out by LAH and ISH. The initial draft of the manuscript was written by LAH. All authors - LAH, ISH, GS, and EE - reviewed and edited the manuscript.

## References

1.  Grinin L, Grinin A, Korotayev A. Global Aging and our Futures. World Futur. 2023 Jul 4;79(5):536–56.

2.  WHO Global. Ageing and health. 2024 [cited 2024 Dec 17]. Ageing and health. Available from: https://www.who.int/news-room/fact-sheets/detail/ageing-and-health

3.  Hogan N, Krebs HI, Charnnarong J, Srikrishna P, Sharon A. MIT-MANUS: a workstation for manual therapy and training. I. In: [1992] Proceedings IEEE International Workshop on Robot and Human Communication [Internet]. 1992 [cited 2024 Mar 25]. p. 161–5. Available from: https://ieeexplore.ieee.org/abstract/document/253895

4.  Scott SH. Apparatus for measuring and perturbing shoulder and elbow joint positions and torques during reaching. J Neurosci Methods. 1999 Jul 15;89(2):119–27.

5. Krebs HI, Volpe BT, Aisen ML, Hogan N. Increasing productivity and quality of care: robot-aided neuro-rehabilitation. J Rehabil Res Dev. 2000;37(6):639–52.

6. Krebs HI, Hogan N. Therapeutic Robotics: A Technology Push. Proc IEEE Inst Electr Electron Eng. 2006 Sep 1;94(9):1727–38.

7. Dukelow SP, Herter TM, Moore KD, Demers MJ, Glasgow JI, Bagg SD, et al. Quantitative assessment of limb position sense following stroke. Neurorehabil Neural Repair. 2010 Feb;24(2):178–87.

8. Herter TM, Scott SH, Dukelow SP. Systematic changes in position sense accompany normal aging across adulthood. J Neuroengineering Rehabil. 2014 Mar 25;11:43.

9. Kitchen NM, Miall RC. Proprioceptive deficits in inactive older adults are not reflected in fast targeted reaching movements. Exp Brain Res. 2019 Feb 1;237(2):531–45.

10. Bosecker C, Dipietro L, Volpe B, Krebs HI. Kinematic robot-based evaluation scales and clinical counterparts to measure upper limb motor performance in patients with chronic stroke. Neurorehabil Neural Repair. 2010 Jan;24(1):62–9.

11. Coderre AM, Zeid AA, Dukelow SP, Demmer MJ, Moore KD, Demers MJ, et al. Assessment of upper-limb sensorimotor function of subacute stroke patients using visually guided reaching. Neurorehabil Neural Repair. 2010;24(6):528–41.

12. Howard IS, Ingram JN, Wolpert DM. A modular planar robotic manipulandum with end-point torque control. J Neurosci Meth. 2009;181(2):199–211.

13. Oldfield RC. The assessment and analysis of handedness: The Edinburgh inventory. Neuropsychologia. 1971 Mar 1;9(1):97–113.

14. Howard IS, Alvarez-Hidalgo L. Haptic Force Feedback Enhances de Novo Learning of Arm Kinematics and Dynamics [Internet]. bioRxiv; 2025 [cited 2025 May 10]. p. 2025.04.29.651298. Available from: https://www.biorxiv.org/content/10.1101/2025.04.29.651298v1

15. Scheidt RA, Reinkensmeyer DJ, Conditt MA, Rymer WZ, Mussa-Ivaldi FA. Persistence of motor adaptation during constrained, multi-joint, arm movements. J Neurophysiol. 2000 Aug;84(2):853–62.

16. JASP Team. JASP (Version 0.19.0)[Computer software] [Internet]. 2024. Available from: https://jasp-stats.org/

17. Moulton RH, Rudie K, Dukelow SP, Scott SH. Quantitatively assessing aging effects in rapid motor behaviours: a cross-sectional study. J NeuroEngineering Rehabil. 2022 Jul 26;19(1):82.

18. Hunter SK, Pereira HM, Keenan KG. The aging neuromuscular system and motor performance. J Appl Physiol. 2016 Oct;121(4):982–95.

19. Yoshimura N, Tsuda H, Aquino D, Takagi A, Ogata Y, Koike Y, et al. Age-Related
Decline of Sensorimotor Integration Influences Resting-State Functional Brain
Connectivity. Brain Sci. 2020 Dec;10(12):966.

20. Seidler RD, Bernard JA, Burutolu TB, Fling BW, Gordon MT, Gwin JT, et al. Motor
control and aging: Links to age-related brain structural, functional, and biochemical
effects. Neurosci Biobehav Rev. 2010 Apr 1;34(5):721–33.

21. Clark BC, Taylor JL. Age-related changes in motor cortical properties and voluntary
activation of skeletal muscle. Curr Aging Sci. 2011 Dec;4(3):192–9.

22. Burke SN, Barnes CA. Neural plasticity in the ageing brain. Nat Rev Neurosci. 2006
Jan;7(1):30–40.

23. Kourosh-Arami M, Hosseini N, Komaki A. Brain is modulated by neuronal plasticity
during postnatal development. J Physiol Sci. 2021 Nov 17;71(1):34.

24. Marzola P, Melzer T, Pavesi E, Gil-Mohapel J, Brocardo PS. Exploring the Role of
Neuroplasticity in Development, Aging, and Neurodegeneration. Brain Sci. 2023
Dec;13(12):1610.

25. Raz N, Gunning-Dixon F, Head D, Dupuis J, Acker J. Neuroanatomical correlates of
cognitive aging: Evidence from structural magnetic resonance imaging.
Neuropsychology. 1998 Feb 1;12:95–114.

26. Fjell AM, Walhovd KB. Structural brain changes in aging: courses, causes and
cognitive consequences. Rev Neurosci. 2010;21(3):187–221.

27. Cassady K, Ruitenberg MFL, Reuter-Lorenz PA, Tommerdahl M, Seidler RD. Neural
Dedifferentiation across the Lifespan in the Motor and Somatosensory Systems.
Cereb Cortex N Y N 1991. 2020 May 18;30(6):3704–16.

28. Kuehn E, Dinse J, Jakobsen E, Long X, Schäfer A, Bazin PL, et al. Body Topography
Parcellates Human Sensory and Motor Cortex. Cereb Cortex N Y N 1991. 2017 Jul
1;27(7):3790–805.

29. Goble DJ, Coxon JP, Van Impe A, Geurts M, Doumas M, Wenderoth N, et al. Brain
activity during ankle proprioceptive stimulation predicts balance performance in young
and older adults. J Neurosci Off J Soc Neurosci. 2011 Nov 9;31(45):16344–52.

30. Kulmala JP, Korhonen MT, Kuitunen S, Suominen H, Heinonen A, Mikkola A, et al.
Which muscles compromise human locomotor performance with age? J R Soc
Interface. 2014 Nov 6;11(100):20140858.

31. Deuschl G, Bain P, Brin M. Consensus statement of the Movement Disorder Society
on Tremor. Ad Hoc Scientific Committee. Mov Disord Off J Mov Disord Soc. 1998;13
Suppl 3:2–23.

32. Berardelli A, Rothwell JC, Thompson PD, Hallett M. Pathophysiology of bradykinesia
in Parkinson's disease. Brain J Neurol. 2001 Nov;124(Pt 11):2131–46.

33. Fearon C, Doherty L, Lynch T. How Do I Examine Rigidity and Spasticity? Mov Disord
Clin Pract. 2015;2(2):204–204.

34. Jankovic J. Motor fluctuations and dyskinesias in Parkinson's disease: Clinical
manifestations. Mov Disord. 2005;20(S11):S11–6.

35. Edwards DF, Lang CE, Wagner JM, Birkenmeier R, Dromerick AW. An Evaluation of
the Wolf Motor Function Test in Motor Trials Early After Stroke. Arch Phys Med
Rehabil. 2012 Apr 1;93(4):660–8.

36. Podsiadlo D, Richardson S. The Timed "Up & Go": A Test of Basic Functional Mobility
for Frail Elderly Persons. J Am Geriatr Soc. 1991 Feb;39(2):142–8.

37. Shumway-Cook A, Brauer S, Woollacott M. Predicting the Probability for Falls in
Community-Dwelling Older Adults Using the Timed Up & Go Test. Phys Ther. 2000
Sep 1;80(9):896–903.

38. Ng SS, Hui-Chan CW. The Timed Up & Go Test: Its Reliability and Association With
Lower-Limb Impairments and Locomotor Capacities in People With Chronic Stroke.
Arch Phys Med Rehabil. 2005 Aug 1;86(8):1641–7.

39. Kristensen MT, Foss NB, Kehlet H. Timed "Up & Go" Test as a Predictor of Falls
Within 6 Months After Hip Fracture Surgery. Phys Ther. 2007 Jan 1;87(1):24–30.

40. Nocera JR, Stegemöller EL, Malaty IA, Okun MS, Marsiske M, Hass CJ, et al. Using
the Timed Up & Go test in a clinical setting to predict falling in Parkinson's disease.
Arch Phys Med Rehabil. 2013 Jul;94(7):1300–5.

41. Wolf SL, Catlin PA, Ellis M, Archer AL, Morgan B, Piacentino A. Assessing Wolf motor
function test as outcome measure for research in patients after stroke. Stroke. 2001
Jul;32(7):1635–9.

42. Fritz SL, Blanton S, Uswatte G, Taub E, Wolf SL. Minimal Detectable Change Scores
for the Wolf Motor Function Test. Neurorehabil Neural Repair. 2009 Sep 1;23(7):662–
7.

43. Turtle B, Porter-Armstrong A, Stinson M. The reliability of the graded Wolf Motor
Function Test for stroke. Br J Occup Ther. 2020 Sep;83(9):585–94.

44. Lyle RC. A performance test for assessment of upper limb function in physical
rehabilitation treatment and research. Int J Rehabil Res Int Z Rehabil Rev Int Rech
Readaptation. 1981;4(4):483–92.

45. Jebsen RH, Taylor N, Trieschmann RB, Trotter MJ, Howard LA. An objective and
standardized test of hand function. Arch Phys Med Rehabil. 1969 Jun;50(6):311–9.

46. Sullivan KJ, Tilson JK, Cen SY, Rose DK, Hershberg J, Correa A, et al. Fugl-Meyer
assessment of sensorimotor function after stroke: standardized training procedure for
clinical practice and clinical trials. Stroke. 2011 Feb;42(2):427–32.

47. Krebs HI, Hogan N, Aisen ML, Volpe BT. Robot-aided neurorehabilitation. IEEE Trans
Rehabil Eng Publ IEEE Eng Med Biol Soc. 1998 Mar;6(1):75–87.

48. Shadmehr R, Mussa-Ivaldi FA. Adaptive representation of dynamics during learning
of a motor task. J Neurosci. 1994 May 1;14(5):3208–24.

49. Cressman EK, Henriques DYP. Sensory recalibration of hand position following
visuomotor adaptation. J Neurophysiol. 2009 Dec;102(6):3505–18.

50. Diedrichsen J. Optimal task-dependent changes of bimanual feedback control and
adaptation. Curr Biol CB. 2007 Oct 9;17(19):1675–9.

51. Howard IS, Ingram JN, Wolpert DM. Composition and decomposition in bimanual
dynamic learning. J Neurosci. 2008 Oct 15;28(42):10531–40.

1182