1 **SCLC-TumorMiner: A Directly Accessible Genomics Resource for Precision Oncology – Big**
2 **Data for Small Cells**
3

4 Fathi Elloumi[1,*], Anjali Dhall[1,*], Daiki Taniyama[1], Augustin Luna[1,2], Yasuhiro Arakawa[1,3], Sudhir
5 Varma[1,4], Jeffrey Wang[1], Anisha Tehim[1,5], Mark Raffeld [6], Kenneth Aldape[6], Christophe Redon[1],
6 Roshan Shresta[1], William Reinhold[1], Mirit Aladjem[1], Jaydira Del Rivero[1], Nitin Roper[1], Anish
7 Thomas[1] and Yves Pommier[1,#]
8

[1] Laboratory of Molecular Pharmacology and Developmental Therapeutics Branch, Center for
Cancer Research, NCI, NIH
[2] NLM, NIH
[3] Department of Clinical Pharmacology and Therapeutics, The Jikei University School of
Medicine, Tokyo, Japan
[4] HiThru Analytics LLC
[5] Cornell University, NY
[6] Laboratory of Pathology, Center for Cancer Research, NCI, NIH
17

18 [*] Equal contributions
19

20 To whom correspondence should be addressed: pommier@nih.gov; fathi.Elloumi@nih.gov
21
22

SUMMARY

RNAseq and DNAseq are fast, cost-effective and quantitative methods to dissect cancer cells.
However, for each sample, they generate thousands of data points, and comparing patient samples
multiplies the complexity. To handle these difficulties, we developed TumorMiner, a web-based
tool for clinicians and basic researchers. Here we present our analyses and website for Small Cell
Lung Cancer (SCLC). SCLC_TumorMiner includes 235 samples from untreated and relapse
patients across the NCI, the University of Rochester, Tongji University and the University of
Cologne (https://discover.nci.nih.gov/SclcTumorMinerCDB/). SCLC_TumorMiner allows the
molecular classification of tumors based on the canonical NAPY classification and NMF, genomic
network analyses exemplified by the Myc and Notch pathways, and the identification of risk-
factors and predictive gene expression biomarkers for cell surface and intracellular targets such as
DLL3, TROP2, SEZ6, CEA, TRPM5, SLFN11 and proapoptotic and multidrug resistance genes.
The architecture of TumorMiner allows its extension to proteomic databases, and to other cancers
and institutions worldwide to elucidate cancer pathways, achieve precision medicine and serve as
a medical assistant software.
38

INTRODUCTION

Small cell lung cancers are rapidly proliferative tumors often diagnosed with metastases (Extensive-stage; ES-SCLC). SCLC accounts for approximately 13%–15% of all lung cancer cases and remains one of the most lethal malignancies. Survival rate is less than one year for most patients, and 95% of patients die within 5 years despite initial response to cisplatin-etoposide combination therapy. Upon relapse, tumors evolve and become resistant to chemotherapy reflecting profound changes in their biology and genomic network.

In addition to mutations of the tumor suppressors TP53 and RB1 in 80% of SCLC, selective activation of transcription factors characterizes the disease. SCLCs have been divided into two groups. 70-80% are neuroendocrine (NE) driven by two pioneer neurogenic transcription factors, ASCL1 (Acheate-scute homolog 1) and NEUROD1 (Neurogenic differentiation factor 1). The remaining non-NE SCLCs are driven by two other transcription factors POU2F3 (POU class 2 homeobox 3) and YAP1 (Yes-associated protein 1). Consequently, SCLCs have been classified in 4 groups: SCLC-A, SCLC-N, SCLC-P and SCLC-Y [1]. Additionally, an Inflamed/Immune subset has been identified as SCLC-I [2]. Multiple genomic studies have been conducted in the past 10 years, confirming the NAPY and NAPI classifications in untreated patients. The common histopathological classification of SCLC is presently based on immunohistochemistry for NE markers: synaptophysin (encoded by the *SYP* gene), chromogranin (encoded by *CHGA*) and neural cell adhesion molecule (encoded by *NCAM1*). Immunohistochemistry of ASCL1, NEUROD1, POU2F3 and YAP1 [3] is not routinely performed.

Immune checkpoint (PD-L1) inhibitors have recently been added to chemotherapy as standard of care for 1st line treatment of ES-SCLC. Atezolizumab (Tecentrip®) was approved in 2019 in combination with carboplatin and etoposide [4], and durvalumab (Imfinzi®) in 2020 in combination with etoposide and either carboplatin or cisplatin [5]. However, immune checkpoint inhibitors only provide a modest overall survival benefit of 2-3 months [5,6].

The most significant therapeutic recent breakthrough is Tarlatamab®, a bispecific T-cell engager (BiTE®) targeting the cell surface glycoprotein Notch ligand DLL3 (Delta-like ligand 3), a transcriptional target of ASCL1 [7]. Tarlatamab was granted FDA approval in May 2024 for ES-SCLC non-responsive to platinum-based therapy or at relapse [8]. However, Tarlatamab is only active in 50% of patients who need to be hospitalized to control treatment-related immune hyper-reactions, and it is prescribed without mandatory evaluation of DLL3 expression.

The substantial benefit provided by Tarlatamab and immune checkpoint inhibitors is fueling the development of novel therapeutic strategies targeting not only DLL3 [9], but also other proteins expressed at the surface of SCLC cells [10]. Two main approaches are actively pursed beyond T-cell engagers, Chimeric antigen receptor T-cell (CAR-T) therapies such as AMG119 [11] and antibody drug conjugates (ADCs). The efficacy of ADCs is not only dependent on the expression of their cell surface target(s) but also on the type of cytotoxic payload delivered to tumor cells. The three main classes of payloads are tubulin inhibitors, as in Rova-T [12], topoisomerase I (TOP1) inhibitors, as in Enhertu (Trastuzumab deruxtecan) and Trodelvy (Sacizutumab govitecan), and DNA binders. Despite their potency, tubulin inhibitors have two main limitations: they are substrates for drug efflux transporters, and they are neurotoxic. The advantage of TOP1 inhibitors is their selectivity for replicating cells and the availability of a testable biomarker, Schlafen 11 (SLFN11) [13], which has a wide range of expression in SCLC cells [7] and tumors [14] and acts as a lethal factor by irreversible blocking stressed replication forks [15,16].

RNA sequencing (RNA-seq) is a well-established, sensitive and routinely used method. However, even when only applied to protein-coding genes, it generates large datasets, which are

difficult to analyze instantaneously in a clinical setting. To address this issue, we developed the
SCLC_TumorMiner database integrating 235 SCLC patient samples (154 untreated and 81
relapse) from our NCI clinic and the published datasets of the University of Cologne (UoC) [17],
Tongji University (TU) [18], and the University of Rochester (RU) [19] (Table 1, Table S1). The
interface of TumorMiner is an extension of on our widely used SCLC_CellMinerCDB website
including the genomics and drug responses of 118 SCLC cell lines from multiple open sources
(https://discover.nci.nih.gov/rsconnect/SclcCellMinerCDB) [7]. Thus SCLC_TumorMiner and
SCLC_CellMinerCDB can be used to compare tumor samples with cell lines.
Here we demonstrate that SCLC_TumorMiner allows instantaneous: 1/ genome-wide analyses
of the transcriptome and mutations of patient samples; 2/ Pathway analyses based on the
hypothesis that genes functioning in a given genetic network have cross-correlated expression; 3/
Patient sample classification using both NAPY and/or NMF (non-negative matrix factorization);
and 4/ Identification of prognosis and predictive biomarkers for targeted therapies.
98

99 **RESULTS**
100
Introduction to SCLC_TumorMiner
The SCLC_TumorMiner website is built on the architecture of our cancer cell line CellMiner
cross-database   websites   (https://discover.nci.nih.gov/).   Thus,   users   familiar   with
SCLC_CellMinerCDB (https://discover.nci.nih.gov/SclcCellMinerCDB/) will have no difficulty
navigating back and forth between the tumor samples and cell line databases [7].
Figure 1 (https://discover.nci.nih.gov/SCLC_TumorMinerCDB/) introduces the website
with a snapshot of the default page display. The upper part provides tabs for each of the website
tools and applications. They are numbered 1-10 for convenience on the figure. The "2 variable
Analyses" tab (numbered "1" in Figure 1A) allows the display of different data types or parameters
on each of the axes: "xsq" (RNA-seq), "sig" (Signatures), "mut" (gene mutations), "mda"
(Miscellaneous data), "hll" (Hallmark gene sets) and "act" (Drug Response). These parameters
can be displayed on each axis independently of each other. For instance, plotting ASCL1
expression vs. NE (Neuro Endocrine) signature shows the expected high correlation (Pearson
correlation (r)=0.79, p-value= e-51) (Figure S1A).
Univariate analysis data can be downloaded as Tab-delimited format. Figure S1B compares
the expression distribution of NEUROD1, ASCL1, POU2F3 & YAP1 in the samples from
untreated and relapse patients. It demonstrates a loss of the bimodal distributions for ASCL1 and
POU2F3 in the relapse samples while YAP1 expression is also significantly reduced in the relapse
samples (Figures S1A-B and 1). As result the assignment of samples to each of the NAPY groups
cannot be applied to more than 50% of the relapse samples while, as reported previously [1], it
discriminates the samples from untreated patients (Figure 1B)
The "Multivariable Analyses" tab (labeled "2" in Figure 1A) shows as default the Heatmap
"Display" of ASCL1 expression (xsq: RNA-seq) vs. the expression of key genes driving the
biology of SCLC: INSM1 & CHGA used as immunohistochemistry NE markers, MKI67
(encoding Ki67 used as proliferation marker), as well as the SCLC pioneer transcription factors
NEUROD1, POU2F3 & YAP1, and the oncogene MYC (Figure 2A). Each row displays the
selected genes and the columns the 235 individual patient samples as a Heatmap). Users can
explore different inputs and outputs by entering chosen gene names in the "Response Identifier"
or "Predictor Identifiers" windows (left panels of Figure 2A). Examples of "Multivariate

Analyses" for the NAPY genes are shown in Figure 2A-E. They will be discussed in the next
"Network and Pathway Analyses" section.
In the "Multivariate Analyses" tab, the default visualization setting is "Display". Users can
also choose the "Linear Regression" tab (found in bottom left corner of the website under the label
"Algorithm"; Figure 2A). This feature computes the correlations between each of the "Predictive
Identifiers" and the "Response Identifier". Figure S2A-C shows the results for ASCL1 for the
genes entered in Figure 2A. The "Plot" option demonstrates the high predictive value of the
combined expression of INSM1, CHGA, MKI67, NEUROD1, POU2F3, YAP1 and MYC as
predictors of ASCL1 expression (Figure S2A) (Pearson correlation r = 0.89; p-value = 6.8e-83;
numbers displayed on top of the graph). The contribution of each of the 7 predictor genes are
available under the "Technical Details" tab (Figure S2B), demonstrating the prominent
contribution of INSM1, CHGA as positive predictors and NEUROD1, POU2F3 and MYC as
negative predictors. Users can also choose "Lasso" (Least Absolute Shrinkage and Selection
Operator), a machine-learning and statistical method that identifies genomic biomarkers predicting
the expression of the "Response Identifier". For YAP1 expression, the top gene identified is the
Neuron Restrictive Silencing Factor REST, a key downstream effector of the Notch pathway [7].
The "Expression Distribution" tab (labeled "3" in Figure 1A) displays the expression of
any chosen gene across the 4 SCLC_TumorMinerCDB datasets (the two untreated NCI & RU
datasets and the two relapse TU & UoC datasets). Schlafen 11 (SLFN11) expression is displayed
as default on the website (Figure S3A). SLFN11 expression tends to be suppressed in relapse
samples, which can explain the resistance of relapse patients to DNA-targeted chemotherapies
(such as etoposide, topotecan, cisplatin and lurbinectedin). Users can display and download data
for any gene from each of the 4 datasets by inserting the corresponding gene HUGO name in the
"Gene symbol" box of the website.
The "Mutation Variants" tab (labeled "4" in Figure 1A) displays mutations for up to 5
genes at a time. Colors reflect the types of mutations, and the variants can be viewed and
downloaded using the "View variants" tab (Figure S3B). As expected TP53 is the most frequently
mutated gene (81% of the 114 samples with Exome-sequencing data). The second most frequently
mutated gene is TTN, which encodes Titin. This is because titin is the largest human protein, and
TTN the largest target for exome sequencing, thereby reflecting the high mutation burden of
SCLC. Accordingly, TTN gene mutation is correlated with Tumor Mutation Burden (TMB)
(Pearson correlation = 0.37, p-value = 6e-05 in the "2-variable analyses" tool of
SCLC_TumorMiner; not shown in the figures). As expected, the next most mutated gene is RB1
with an overall mutation frequency of 55% (Figure S3B). Although inactivation of SLFN11 leads
to resistance to chemotherapy [15,20,21], its mutation frequency is low (only 4% of all 114 sequenced
samples; Figure S3B) (13% vs. 2% in the untreated).
The "Prognosis Biomarkers", "Predictive Biomarkers" and "MyPatient" tabs (numbered 5,
6 & 7 in Figure 1A) will be discussed in the remaining sections of the manuscript with examples
displayed in Figures 5, 7 and S6.
169
Genomic networks and canonical pathways of SCLC patient samples
We previously reported [7] that for the SCLC cell lines, the expression of genes functioning together
in a given pathways are cross-correlated (see Fig. 5 in [7]). The main pathways for SCLC comprise
the neuroendocrine (NE) pathways driven prominently by ASCL1 and less frequently by
NEUROD1, and the non-NE pathways driven by POU2F3 and YAP1, the MYC genes and genes
of the Notch pathway [1,7].

As in the SCLC cell lines (https://discover.nci.nih.gov/SclcCellMinerCDB/) [7] and
consistent with prior studies of clinical samples, the group of NE-SCLC is prominent across our
four clinical datasets based on the expression of the two genes used as immunochemistry NE
markers, INSM1 and CHGA (Figures 2A). As expected, ASCL1 expression is highly correlated
with the NE markers, while NEUROD1
Next, we used the "Multivariable Analysis" tool to investigate the networks/pathways of
each of the 4 NAPY genes (Figure 1B). The approach is an extension of our previous genomic
network analyses in the 118 SCLC cell lines for ASCL1 and YAP1 (Figure 5 in ref. [7]). Here we
also analyzed NEUROD1 and POU2F3 (Figures 2C-E) in the tumor samples and compared the
235 tumor samples to the 118-cell line collection ((Figure S2C). Figure 2A shows that multiple
samples express both ASCL1 and NEUROD1 and that NEUROD1-expressing samples do not
closely match INSM1 or CHGA. Figure 2A also shows a significant overlap between YAP1,
POU2F3 and MYC expression.
Analysis of the ASCL1 coexpression network [7] (Figure 2B) shows that most of the samples
expressing ASCL1 also express INSM1 (Pearson correlation r = 0.72; p-value = 3.6e-38 in the "2
variable analysis"). As expected, NKX2-1 and PROX1, the transcriptional coactivators of ASCL1
[7,22] (Figure 5 in [7]) show high coexpression (Pearson correlations r = 0.67 & 0.55; p-values = 3.2e-
32 and 9.3e-20, respectively using the "2 variable analysis" tool). The downstream transcriptional
target of ASCL1, DLL3 (encoding the Delta Like Canonical Notch Ligand 3), a validated SCLC
therapeutic target, is highly expressed in approximately two-third of the samples, and its
expression is highly correlated with ASCL1 (Pearson correlation r = 0.82; p-value = 5.7e-58 in the
"2 Variable Analyses"; highest correlate with ASCL1 expression in "Compare Patterns" of the "2
Variable Analyses"). *SEZ6* (encoding Seizure 6-like), another ASCL1 target gene and an emerging
cell surface target for Antibody Drug Conjugates is also highly correlated with both ASCL1 and
DLL3 (Figure 2B). Of note, SEZ6 expression is highly correlated with SEZ6L (SEZ6 homolog
like) in spite of different chromosome locations (17q11.2 and 22q12.1, respectively), suggesting
that both genes function in the ASCL1 pathways. The "Compare Analyses" tool of
SCLC_TumorMinerCDB identifies additional genes associated the ASCL1 genetic network, such
as GRP (Gastrin Releasing Peptide) (4th most highly correlated gene after DLL3 and SEC11C
(SEC11 Homolog C, Signal Peptidase Complex Subunit)). Taken together, the 9 "Predictor genes"
listed in Figure 2B are highly predictive of ASCL1 expression (r = 0.87; p-value = 6.5e-73; Figures
S2C) and comprise a significant part of the ASCL1 network/pathways.
Analysis of the NEUROD1 coexpression network shows that NEUROD1 expression only
accounts for a fraction of the NE samples as many samples have high INSM1 expression without
expressing high levels of NEUROD1 (Figure 2C, compare the 1st and 2nd rows). NKX2-2 (NK2
transcription factor related, locus 2 (Drosophila)), which is involved in the morphogenesis of the
central nervous system and a known regulator of NEUROD1 in pancreatic islet cells together with
the Neurogenin genes (NEUROG1 and NEUROG3) [23] are co-expressed with NEUROD1 (Figure
2C). The coexpression of PAX6 (Paired Box Homeotic Gene-6), which encode a transcription
factor important for the development of the eyes, nose, pancreas and the central nervous system,
is logical as it has been shown to control NEUROD1 expression [24]. Using the "2 variable Analysis"
tool and the "Compare Patterns" tools of SCLC_TumorMinerCDB (see Figure 1A; red dotted
box), we found coexpression of two genes encoding cell surface proteins and potential targets for
immune therapies, *GFRA1* (Glial Cell Line-Derived Neurotrophic Factor Receptor Alpha) and
SEMA6A (encoding one of the semaphorins) [25] (Figure 2C). The SCLC cell lines [7] also express
significant levels of SEMA6A with high correlation with NEUROD1 (p = 0.57 with p-value =

2.7e-11). Whether NEUROD1 directly drives the expression of GFRA1 and SEMA6A remains to
be determined. Taken together, the 8 NEUROD1 "Predictor genes" are highly predictive of
NEUROD1 expression (r = 0.85; p-value = 2.6e-65; Figures S2A-C) and could be used to establish
a prominent NEUROD1 subtype in tumor samples. They also identify a significant part of the
NEUROD1 network/pathway.
Analysis of the POU2F3 network shows the expected inverse expression between POU2F3
and INSM1 (Figure 2D, compare the 1st and 2nd rows) as POU2F3 SCLC cells are derived from
the Tuft cells of the bronchial epithelium and are non-NE [26-28]. Figure 2D shows that the tumor
samples expressing POU2F3 also coexpress 4 transcription factors known to drive POU2F3
expression [26,29]: ASCL2, COLCA2 (also known as OCAT-2 or POU2AF3), C11orf53 (also known
as OCAT-1 or POU2AF2) and SOX9. Consistent with the Tuft lineage of the POU2F3-driven
SCLC (SCLC-P), POU2F3 is also highly correlated with the Tuft-ionocyte progenitor cell-specific
genes FOXI1 and AVIL [27,28], and with TRPM5 (Transient Receptor Potential Cation Channel
Subfamily M Member 5), which encodes a cell surface ion channel protein involved in immune
defense, and was recently shown to be highly expressed in bronchial Tuft cells [30]. This suggests
that TRPM5 could be considered as cell surface target for SCLC-P tumors. Figure 2D also shows
that the POU2F3 network/pathway is coordinated with the NOTCH pathways as POU2F3
expression is highly correlated with REST and its transcriptional activator NOTCH1 (Pearson
correlations r = 0.67 and 0.46; p-values = 9e-22 and 2.7e-9, respectively in the "2 variable analysis"
for the untreated samples). MYC expression is also highly correlated with POU2F3 (Pearson
correlation r = 0.74; p-value = 5.6e-28 in the "2 variable analysis" in the untreated samples), which
is consistent with independent reports [18,28,31]. Taking together the expression of the 11 genes
displayed in Figure 2D could be used as predictor for POU2F3 expression and for classifying a
given tumor sample as SCLC-P (Pearson correlation = 0.94; p-value = 3.3e-112) (Figure S2C).
Analysis of the YAP1 network confirms that samples with high YAP1 expression lack
INSM1 expression. Consistent with the SCLC cell lines analyses (Figure 5 in [7]), TAZ (encoded
by *WWTR1*), the transcriptional co-activator partner of YAP1 is coexpressed with YAP1 (Pearson
correlation r = 0.81; p-values = 1.24e-56). Similarly, TEAD2 and TEAD3, the downstream
transcription factors of YAP/TAZ are coexpressed with YAP1. Figure 2E also shows that the
downstream transcriptional targets of YAP1 (see Figure 5C-D in [7]) including SMAD6, VGLL4,
PTPN14, and the cell adhesion genes CCN1, CTNNA1 and CTNNA2 are coexpressed with YAP1.
Also consistent with the SCLC cell line data (see Figure 5C-D in [7]) the NOTCH pathway (REST,
NOTCH1 and NOTCH2) is co-expressed in the SCLC-Y tumor samples. Together the expression
of the 12 genes displayed in Figure 2E is a statistically significant predictor for YAP1 expression
and classification of a given tumor sample as SCLC-Y (Pearson correlation = 0.89; p-value = 1.9e-
81) (Figure S2D).
Analysis of the Notch network using REST, the transcriptional target and effector of the
Notch pathway (Figure 2F) shows, as expected, coexpression of REST with the 4 Notch receptors,
NOTCH1-4 and with YAP/TAZ, consistent activation of the Notch pathway by YAP1-mediated
transcription [32]. By contrast, the Notch ligands (JAG1/2, DLL3, DLL1, DLL4 and DLK1) are
significantly suppressed in the Notch receptor-positive samples and coexpressed with DLL3 and
ASCL1 (Figure 2F), consistent with the conclusion that the YAP/Notch/REST network controls
the NE fate of SCLC cells [32] and vice-versa as DLL3 suppresses Notch signaling in SCLC [33].
The Myc pathway (including MYC, MYCL and MYCN) is activated in 52% of the
untreated patient samples where MYCL dominates (36.4% of the samples) and is linked to ASCL1
expression (SCLC-A) (Figures 2B & 2G). By contrast MYC is highly expressed in a smaller

fraction of the untreated patients (15.6%) and primarily in the SCLC-P samples (Figures 2D) [34].
Consistent with the fact that POU2F3 disappears in the relapse samples and that ASCL1 expression
is also attenuated in the samples of relapse patients, the expression of both MYCL and MYC is
reduced in the relapse samples; with a greater reduction for MYCL at relapse (Figures 2G).
To test the robustness and conservation of the 5 networks displayed in Figure 2 (ASCL1,
NEUROD1, POU2F3, YAP1 and Notch) we compared their predictive value in TumorMiner and
CellMinerCDB. Figure S2C shows that the 5 genomic networks are highly conserved in tumor
samples and cell lines.
276
Classification of SCLC tumor samples
Although all the samples from the untreated patients can be assigned to the NAPY classification,
less than half (47%) of the relapse samples can be classified using RNA-seq expression levels > 5
(log2(FPKM+1)) (Figure 1B). To overcome this limitation, we first performed cluster analyses
using 12 genes by adding to the 4 NAPY genes (highlighted in red in Figure 3) the two NE genes
(INSM1 and CHGA), MYC and REST and 4 innate immune genes of the major histocompatibility
complex (MHC): HLA-A, HLA-B, HLA-C and B2M (Beta-2-Microglobulin) that are part of the
recently defined inflamed signature of SCLCs (SCLC-I) [2]. This 12 gene algorithm successfully
classified the untreated patients in 2 groups (Figure 3A; see annotations under the heat map). The
largest is the SCLC-NE group (≈ 70% of the patient samples) with high expression of ASCL1,
INSM1, CHGA and/or NEUROD1, and the 2nd the POU2F3 group (SCLC-P). As discussed above
the SCLC-P also includes samples with high MYC, REST and to a lesser extent YAP1 expression.
The inflamed group characterized by high MHC expression (HLA-A/B/C and B2M) is more
diffuse (≈ 50% of the patient samples) and tends to be excluded from the SCLC-NE group and to
encompass the SCLC-P group.
For relapse patient samples, the 12 gene signature did not provide clear clusters (Figure
3B), and the ASCL1, NEUROD1, INSM1 and CHGA genes failed to form defined clusters. The
only visible separation is for the MHC genes with ≈ 50% of the samples MHC positive (SCLC-I)
(Figure 3B; see annotation under the heatmap).
To overcome the difficulty of classifying relapse patient samples, we performed NMF
(Non-Negative Matrix Factorization) analyses [2] using the top-5000 protein coding genes most
variably expressed in the samples from untreated and relapse patients (Figure 3C, Table S2 and
S3). Among those genes, 65% are common for the untreated and relapse samples (Figure S5A).
Cophenetic analysis showed that 4 clusters gave a satisfactory division of the samples (Figure
S5B-C). The top 30 selectively expressed genes for each of the 4 clusters for the samples of
untreated patients (NMF1u, NMF2u, NMF3u and NMF4u) are displayed in Figure 3D. Similarly,
the top 30 selectively expressed genes for the clusters of relapse patient samples (NMF1r, NMF2r,
NMF3r and NMF4r) are displayed in Figure 3E. Overall, only a few genes are common in the
NMF clusters of untreated and relapse patient samples: 5 out of 30 between NMF1u and NMF1r,
3 out of 30 in NMF2u and NMF2r, and 2 out of 30 in NMF3u and NMF3r (see diagonal lines
between panels D and E in Figure 3) See (Table S4 and S5).
Pathway analyses for each of the clusters show that NMF1u consists of genes involved in
protein, RNA metabolism and cell cycle division (Figure S5A), and includes the NE genes ASCL1
(Figure 3H) and GRP (coding for the gastrin releasing peptide; see above) (Figure 3D). The NE
enrichment of NMF1u is easily visible through the mapping ASCL1, NEUROD1, INSM1 and
CHGA in the NMFu clusters (Figure 3F). NMF2u also shows a strong NE enrichment (Figure 3F),
which is confirmed by Reactome pathway analysis (Figure S5A). NMF3u coincide with POU2F3

expression (Figures 3F and 3H), and with immune and interleukin pathways (Figure S5A) as well
as with the MYC pathway (Figures 3D; MYC and MYB annotated in red). NMF4u partially
extends to NMF3u and shows positive enrichment for immune, cytokine and interferon pathways
(Figures 3D and S5A). It is also enriched for YAP1 (Figure 3H).
Pathway analyses for the 4 relapse NMF clusters show that NMF1r includes genes and
pathways related to NE differentiation and cell cycle progression (Figures 3E and S5B) as well as
ASCL1 (Figure 3H). NMF2r and NMF1r overlap with genes involved in cell metabolism as well
as NE pathways (including NEUROD1) (Figures 3E, 3H and S5B). NMF3r includes genes
involved in metabolism and complement. It also includes YAP1 (Figure 3H). NMF4r consists
specifically of immune genes (Figures 3E and S5B) and some YAP1-expressing samples (Figures
3G and 3H).
Figure 4A summarizes the Reactome pathway analyses across the samples. It demonstrates
the similarities between the untreated and relapse NMF1 and NMF2 clusters with enrichment for
NE and cell cycle pathways. It also demonstrates the prominent enrichment for immune pathways
both in the untreated and relapse NMF 3 and 4 clusters. The complete pathway enrichment results
for untreated and relapse SCLC are provided in Table S6 and Table S7, respectively.
330
NMF clusters coincide with molecular signatures
Having performed NMF classification, we derived the NMF signatures (4 untreated and 4 relapse;
see Methods) and integrated them into SCLC_TumorMiner. This feature enables direct
explorations and correlation analyses with other parameters included on the website. Figure 4
explores the correlations between the NMF signatures and the transcriptional signatures included
in SCLC_TumorMiner: NE, RepStress, CES, Epithelial, Stroma, Immune, APM, Ychrom. The
signatures were computed as the mean expression of the genes included in each signature (see
Method section). Figure 4B tabulates the correlations between each NMF and the 8 signatures
(Pearson correlation coefficients obtained by comparing NMFs and individual signatures). The
colors reflect the direction and intensity of the correlations (red for highly positive and blue for
highly negative). Consistent with the top gene signatures (see Figure 3D-E) and the Reactome
pathway analyses (see Figure S5A-B), the NMF1u, NMF2u, NMF1r and NMF2r signatures are
highly correlated with the NE signature as well as with the RepStress and CES (Chromosome
Centromere and Kinetochore score) signatures. They are also significantly devoid of the Immune,
APM (Antigen Presentation Machinery) and Stroma signatures (Figure 4B). Conversely, the
NMF3u, NMF3r, NMF4u and NMF4r are enriched for the Immune, APM, Stroma and epithelial
signatures and devoid of the NE, RepStress and CES signatures.
Figure 4C shows examples of the significant correlations NMF2r and NMF4r with 4 of the
8 signatures as snapshots of "2 Variable Analyses". Together these results demonstrate the
functional significance of the NMF classification for SCLC patient biopsies.
351
Prognosis biomarkers and risk factors
The SCLC_TumorMiner website is designed to assess prognosis biomarkers (Figures 5 and S6).
Users can test risk factors based on gene expression, mutations, clinical metadata, gene and NMF
signatures. The default display on the website is for SLFN11 (Schlafen 11) (Figure S6A) whose
expression is known to predict response to chemotherapeutic agents targeting DNA replication
[15,20]. SLFN11's wide range of expression in SCLC samples (Figure 6E and S8) makes it a
quantitative biomarker. Taking the 4 datasets together, high SLFN11 expression is associated with
significantly better survival (Figures S6E), which agrees with a recent publication [35] demonstrating

preferential response of SLFN11-expressing cells to the DNA-targeted chemotherapies used to
treat SCLC.
The prognostic value of each of the eight NMF gene signatures can be explored in the
"Prognosis Biomarkers" tool of SCLC_TumorMiner. NMF4r (inflammatory-immune) is a good
prognosis biomarker for the entire set of 205 patients (Figure 5A-C). The good prognosis value of
the APM, immune and Stroma signatures (Figures 5D and S6B) also indicates that tumors with
high immune and stroma infiltration respond better to treatments.
Notably, none of the NAPY genes show significant predictive value, even in the untreated
patients despite their differential expression (see Figure 1). Similarly, MYC, MYCL, REST and
NOTCH do not show significant predictive value. On the other hand, high expression of some
genes is associated with high risk in untreated patients (Figures 5E). For instance, consistent with
a recent study reporting that high expression of Serine and Arginine Rich Splicing Factor 1
(SRSF1) gene is linked with poor survival in SCLC patients [36], we also observe that high
expression of SRSF1 is a risk factor for the untreated patients (HR 2 groups = 1.7; p-value = 0.02),
while its significance in the relapse patients (NCI + UR datasets) is less (HR 2 groups = 1.3; p-
value = 0.19).
In addition to univariate analyses, user can perform multivariable analyses by selecting
combination of multiple genes or signatures. Figures 5E-F shows the example of combining
MKI67, which encodes the proliferation biomarker Ki67, an established poor prognosis biomarker,
with SEZ6 and SRSF1 in the 150 samples of untreated patients. High expression of the 3 genes
together shows significantly poor prognosis (HR 2 groups = 1.7; p-value = 0.01). Additional
predictive biomarkers can be searched using the Lasso analysis option (highlighted in yellow at
the bottom left in Figure 5A).
383
Predictive biomarkers for cell surface targets
ADCs are rapidly changing cancer treatments by allowing tumor-targeted delivery of
chemotherapeutic agents, which otherwise exhibit dose-limiting toxicity. Thirteen ADC have been
approved in the past 6 years while the FDA only approved one small molecule, Lurbinectedin for
metastatic SCLC. The most recently FDA-approved ADC (January 2025) is Datroway®
(Datopotamab Deruxtecan) delivering the TOP1 inhibitor Deruxtecan to cancer cells expressing
TROP2 (TACSTD2) [37]. In parallel, CAR-T cells and bi- and tri-specific T-cell engagers are in
active development. Among the T-cell engagers, Imdelltra® (Tarlatmab-dlle) is approved for the
treatment of ES-SCLC [8] without DLL3 biomarker(s) for patient selection. Although the DLL3-
targeted ADC with a tubulin inhibitor payload, Rovalpituzumab teserine (ROVA-T) showed
clinical activity [12], it was discontinued due to its dose-limiting toxicity [38].
Figures 6A-B displays the expression of clinical relevant cell surface targets (Table S8). In
the untreated patient samples, DLL3 shows a broad range of expression with high expression in
multiple NE samples (Figure 6A and 6C and S7A) belonging to the NMF1u & NMF2u groups
(Figure 6C). This result is consistent with those of the SCLC cell lines [7]. Notably, we find reduced
DLL3 expression at relapse (Compare Figures 6C and D; see also Figure S7A), consistent with the
reduced ASCL1 expression at relapse (see Figure 1A). Together these results highlight the
potential of measuring DLL3 expression as biomarker for DLL3-targeted therapies and suggest
that DLL3-targeted therapies should be moved to front-line treatment.
Because TROP2 ADCs, Datroway® and Trodelvy® , are FDA-approved for triple-negative
breast cancer (TNBC), and Trodelvy® has been granted Breakthrough Therapy Designation for
ES-SCLC patients [39], we looked at TACTSD2 (TROP2) expression across our 235 samples

(Figures 6A-D and S7A). Like DLL3, TROP2 expression is low in relapse tumors (Figure 6 and
S7A). Only the samples of untreated patients with low NE score and high APM score highly
express TROP2 (Figure 6A). Accordingly, TROP2 expression is highest in the NMF3u (POU2F3-
endriched) & NMF4u (Immune-enriched) clusters (Figures 6 and S7A). Further analyses in the
SCLC cell lines of CCLE also show that none of the 50 SCLC cell lines express TROP2 above
log2(FPKM+1) >5 while 24% of the cell lines expressed DLL3 above this threshold. These results
suggest potential utility of Trodelvy® or Datroway® in newly diagnosed patients with non-NE
(POU2F3), NMF4 and APM (Inflammatory/immune) signatures.
Figures 6 and S7A display the expression of 6 additional cell surface targets with high
expression [log2(FPKM+1) >5] in at least 10 samples. DLK1 (Delta Like Non-Canonical Notch
Ligand 1) is highly expressed many of the untreated patient samples with NE signature (Figures
6A and S7A). It is also expressed in relapse tumors although less than in the untreated group
(Figure 6A-D). NMF analyses show DLK1 expression in the NMF1u and NMF2u and in
approximately 50% of the samples in the NMF4u group (Inflammatory and Immune group)
(Figure 6C). These results are encouraging as a DLK1-ADC is being developed and tested in NE
tumors [40].
Two of the carcinogenic embryonic antigens (CEACAM5 and 6), which are targeted with
several ADCs in clinical trials for colon, gastric cancers and NSCLC, are highly expressed in both
the untreated and relapse patient samples with NMF1u and NMF2r signatures (Figures 6A-D and
S8). These results suggest the potential of CEACAM5/6-targeted ADCs in SCLC patients. It also
suggests the potential value of measuring CEA in the blood of SCLC patients as a biomarker [41,42].
Claudin 18 (CLDN18) is also the target of ADCs in clinical trials for pancreatic cancer.
Figure 6A shows that it is highly expressed in a few untreated patients but not in relapse patients.
As discussed above, SEZ6, a downstream target of ASCL1 [43] shows high expression across
untreated and relapse samples matching the NE signature (Figures 6A-D and S7A). Notably, the
overlap between SEZ6 and DLL3 is only partial. We also found that the B7-H3 (encoded by
CD276, and belonging to the immunoglobulin superfamily), a common ADC cell surface target,
is expressed in a broad range of SCLC samples (Figures 6A-D and S7A). These results validate
the potential of SEZ6- and B7-H3-targeted ADCs in SCLC [44].
435

Predictive biomarkers for chemotherapy and ADC payloads
Because ADCs act by delivering chemotherapeutic payloads, we explored the intracellular targets
and determinants of response to these payloads. Figures 6E-J and S7B-C summarize the expression
of intracellular drug targets (Figure E-F), drug efflux transporters (Figure G-H) and proapoptotic
genes (Figure I-J) known to drive response to chemotherapeutic drugs.
Schlafen 11 (SLFN11) shows a broad expression range (over 8-fold) both in the untreated
and relapse patient samples, in agreement with SLFN11's wide distribution across the SCLC cell
lines irrespective of the NAPY subgroups [7] (https://discover.nci.nih.gov/SclcCellMinerCDB/).
Consistent with SLFN11being an immune- and interferon-response gene [15], the NMF4
inflammatory/immune subgroups show significantly high expression of SLFN11 both in the
untreated and relapse patient samples (Figure 6E and F). Comparing SLFN11 between the
untreated vs. relapse samples shows significant suppression of SLFN11 expression in the relapse
patient samples (two-tailed p < 0.0001) (Figure S7B and S8). The broad range of SLFN11
expression suggests that further studies are warranted to confirm the relevance of SLFN11 to
predict response of patients treated with topotecan and TOP1 inhibitor payload ADCs [3,14,45].

Since TOP1 is the primary cellular targets of topotecan and TOP1-ADC, and TOP2 the
target of etoposide, we looked at their expression (Figure 6E-F). Both are highly expressed in the
samples from untreated and relapse patients with a slightly lower expression in the
inflammatory/immune subgroups (NMFu and NMFr). Thus, TOP1 and TOP2 levels are irrelevant
as predictive biomarkers.
Lack of O-6-Methylguanine-DNA Methyl-Transferase (MGMT) is well-established as
predicting the activity of DNA alkylating agents, and foremost of temozolomide [46], which is
occasionally used in SCLC [47]. Figure 6 (right panels E & F) shows low expression MGMT in some
samples, suggesting the potential of measuring MGMT expression for temozolomide treatments.
Drug efflux transporters of the ABC family are a major cause of resistance to tubulin
inhibitors that are commonly used as ADC payloads. Yet, measuring ABC transporters is not done
in clinical routine. Figure 6G-H displays the expression of the main ABC transporter genes.
ABCB1, which encodes P-glycoprotein (PgP) has notably low expression both in the untreated
and relapse samples, while ABCC2 and ABCC3 show significant overexpression in the NMF3r
cluster (Figure 6H). Together, these results suggest that measuring ABC transporters should be
considered for ADCs with anti-tubulin payloads.
Because broad resistance of cancers to chemotherapy is dependent on their reduced
apoptotic threshold, we explored the apoptotic genes. Figures 6I-J and S7C shows global
suppression of proapoptotic gene expression in relapse patient, which may account, in part for the
overall resistance of relapse patients to chemotherapy.
471
Analysis of individual patients using the "MyPatient" tab of SCLC_TumorMiner
One of the long-term goals of SCLC_TumorMiner is to provide an interface for classifying tumors
and providing prognosis and rational therapies to a given patient (Figure 7A).
As discussed above (see Figure 1B), untreated patient samples can be assigned to at least
one of the NAPY groups based on high expression cut-off of log2(FPKM+1) > 5. In our dataset,
as expected, most samples are prominently NE (74% ASCL1, 20% NEUROD1), with 14% of the
samples dual ASCL1 & NEUROD1. The POU2F3 group represents 10% of the samples, which is
consistent with prior publications [1]. The YAP1 group is the smallest with 8% of the samples. We
found only 2 samples (1%) with both ASCL1 & YAP1 (see Figures 1B). Thus, we conclude that
the NAPY classification can effectively classify samples from untreated patients.
The situation is different for the relapse samples (see Figures 1B and S9A). Less than half
of the samples (46%) can be assigned to the NAPY groups: 31% to the ASCL1 group, 14% to the
NEUROD1 group (i.e. a total of 45% are NE), only 1 sample to the POU2F3 group, and no sample
to the YAP1 group. Thus, the NAPY classification is deficient for relapse patient samples.
To overcome this limitation, based on our data, we developed open-access NMF classifiers
(listed in Table S5). We first selected the genes with a 2-fold selectivity compared to the genes in
the other 3 NMF groups. This led to a list of genes in each NMF group:128 for NMF1r, 49 for
NMF2r, 299 for NMF3r and 293 for NMF4r. The same was done for the untreated samples
providing 54 genes for NMF1u, 137 for NMF2u, 85 for NMF3u and 386 for NMF4u (Table S4).
Figure S9B shows the mean average scores for each NMFr across the 81 samples from relapse
patients. A threshold of 3.8 was chosen to differentiate the samples assigned to each NMF class.
This approach enables the computation of mean average NMF scores for individual samples. We
implemented this approach to score automatically individual samples in the "MyPatient" tab of
SCLC_TumorMiner (Figure 7B). Upon typing a patient identifier (code), SCLC_TumorMiner
automatically displays key parameters for classifying the patient sample.

497    The "My Patient" tab also displays predictive biomarkers for ADCs, BiTEs or CAR-T
498 therapies (Figure 7C). This is done by the TumorMiner software, which queries a dataset listing
499 46 cell surface targets for ADCs currently FDA-approved and in clinical development (Table S8).
500 The website displays the cell surface targets expressed above a threshold of 5 (log2(FPKM+1)).
501 For example, in Figure 7C, the displayed patient tumor sample (ID: S00035T) overexpresses
502 SEZ6, NCAM1, DLK1, FN1 and SLFN11, implying that the patient could be candidate for ADCs
503 comprising SEZ6, NCAM1, DLK1 or FN1 antibodies and a TOP1 payload based on high SLFN11
504 expression.

506    These results highlight the translational potential of the SCLC_TumorMiner platform
507 for individualized patient care. While traditional NAPY classification remains useful for untreated
508 tumors, its limited applicability to relapsed samples underscores the importance of alternative
509 classifiers. By implementing NMF-based transcriptomic classifiers and integrating biomarker-
510 based predictions for targeted therapies, the "MyPatient" module enables real-time, personalized
511 tumor profiling. This facilitates not only tumor subtype identification but also actionable
512 therapeutic insights, bridging molecular characterization with clinical decision-making in SCLC.
513 As the platform evolves with more comprehensive datasets, it holds promise for guiding precision
514 medicine strategies in both newly diagnosed and treatment-refractory patients.

516 **DISCUSSION**

517 Mining cancer patient clinical and genomics data is critical for understanding the state of the
518 disease, finding biomarkers, predicting disease outcome or drug response, and ultimately shifting
519 patient care to a logic-based and flexible paradigm. During the last decade, new tools have emerged
520 such as cBioPortal (http://cbioportal.org), Xena (https://xena.ucsc.edu)), GEPIA2
521 (http://gepia.cancer-pku.cn/), UALCAN (https://ualcan.path.uab.edu/) or CARIS
522 (https://www.carislifesciences.com/) offering different functions for data visualization and
523 analyses mining The Cancer Genome Atlas (TCGA) or public datasets. Here we present "SCLC-
524 TumorMiner" a new web portal for mining patient clinical, genomics and treatment response data
525 focusing on SCLC, which was not included in TCGA. SCLC_TumorMiner hosts samples from
526 both relapse and untreated patients. Its architecture and displays are constructed on the architecture
527 of SCLC_CellMinerCDB (https://discover.nci.nih.gov/SclcCellMinerCDB), which enables users
528 to compare tumor samples with a broad collection of SCLC cell lines derived primarily from
529 untreated patients [7,48].
530    SCLC_TumorMiner (https://discover.nci.nih.gov/SclcTumorMinerCDB/) contains a
531 battery of unique tools (see Figure 1A) allowing instantaneous display of omics data for individual
532 genes and individual patient samples (*2-Variable Analyses*, *Prognosis Biomarkers* and *Predictive*
533 *Biomarkers*) or for multiple genes to explore molecular networks (*Multivariable Analyses* and
534 *Compare Patterns* tools). It integrates 8 pathway *Signatures* relevant to SCLC biology (NE,
535 RepStress, CES, APM, Immune, EMT, Stroma, Y-Chrom), 50 Hallmark signatures and 46 cell
536 surface targets relevant to the current development of Antibody Drug Conjugates (ADCs),
537 Bispecific T-cell engagers (BiTEs) and CAR-T cell therapies. Data can be downloaded from the
538 SCLC_TumorMiner website for further analyses, as demonstrated in this report by the different
539 plots comparing gene expression in untreated and treated patients and in samples from different
540 NMF groups (see Figures 2G, 3H, 6C-J, S1B-C and S7).
541    A biologically relevant feature of TumorMiner is its ability to explore the logic of
542 molecular networks/pathways based on the premise that individual components of molecular

pathways are quantitatively linked and significantly correlated [7]. Figures 2 and S2 demonstrate
that each of the NAPY genes has its own network (genomic identity) across tumor samples and
that these networks are highly conserved across patient samples and cell line models (Figure S2C),
implying their robustness. A notable finding is the interconnection between the Notch, the
YAP/TAZ and the ASCL1-NE pathways whereby expression of the YAP/TAZ pathway is tightly
linked to activation of the Notch pathway; and vice-versa that activation of the NE (ASCL1-SEZ6)
pathway is associated with expression of the Notch ligands (DLL3, DLK1, DLL1, DLL4 and
JAG2) and negatively correlated with the Notch receptors and their transcription effector REST
(see Figure 2F). These results suggest a feedback loop involving the non-NE pathways including
YAP/TAZ and the Notch receptors, and the NE pathway including ASCL1 and the Notch ligands
[32,33]. Whether and how YAP/TAZ functions upstream of the Notch transcription factor REST
remains to be established.
Cancer cell plasticity and heterogeneity are two outstanding characteristics of SCLC
[10,19,28,49-52]. This is exemplified by the fact that at relapse, patients become resistant to
chemotherapy after initially responding often dramatically. Our analyses reveal several key
differences between the samples of untreated and relapse patients.
One such difference is the loss of the bimodal expression of NEUROD1, ASCL1, POU2F3
and YAP1 in the samples from relapse patients (see Figure 1). This could have, at least two
explanations, which are not mutually exclusive: 1/ differentiated cancer cells expressing ASCL1
(NE cells) or POU2F3 (Tuft-ionocyte-like cells) are eliminated by the 1st line chemotherapy (see
Figures 1A and S1B), leaving behind less differentiated resistant cancer cells; 2/ chemotherapy is
less effective toward the less differentiated and more plastic tumor cells with basal cell
characteristics [28], which remain in the tumor and become prominent at relapse. Both explanations
are consistent with the conclusion that SCLC tumors are endowed with intratumoral heterogeneity
(ITH) and plasticity [10,19,28,49-51].
Another notable difference is the significant downregulation of MYCL in relapse tumors
(see Figures 2G and S1C). This reduction is likely consequential to the fact that ASCL1 is a
transcriptional driver of MYCL [53] (as visualized in Figures 2C) [53] and that ASCL1 expression is
significantly suppressed in the relapse tumor samples (see Figures 1A and S1B).
Prognosis biomarkers (i.e. Risk factors) are also different between relapse and untreated
patient samples. While MKI67, SEZ6 and SRSF1 [which encodes a splicing factor associated with
poor prognosis [36]], are high risk factors for untreated patients, they are not significant in relapse
patients (see Figures 5 and S6C).
With respect to the chemoresistance of the relapse tumors, it is likely multifactorial. Indeed,
while the expression of SLFN11 and of proapoptotic genes (BAD, BAX, BID, NOXA and PUMA)
is high in chemotherapy-naïve patient samples, it is significantly reduced in relapse tumors (see
Figures 6 and S7), suggesting that the initial response of patients to various chemotherapies is
driven, in part by SLFN11 and pro-apoptotic genes. Conversely, resistance of the relapse patients
may be related to the downregulation of the cell death and SLFN11 pathways. Whether these
pathways are epigenetically suppressed in relapse tumors and could be reactivated by HDAC and
EZH2 inhibitors remains to be tested [45,54,55].
Cell surface targets for bispecific antibodies T-cell engagers (BiTEs), antibody drug
conjugates (ADCs), and CAR-T are in full development. Regarding the FDA-approved therapies
targeting DLL3 and TROP2, we find that expression of DLL3, the target of Tarlatamab
(Imdelltra®) is generally low at relapse, consistent with DLL3 being a direct transcriptional target
of ASCL1, which is significantly suppressed at relapse [7,53]. This suggests moving Tarlatamab to

1st line therapy. Also, since IHC analyses apparently fail to predict Tarlatamab responses [56], adding
RNAseq to IHC could identify patients who respond (approximately 40%) to Tarlatamab [8,56] and
the other DLL3-targeted therapies in development including other BiTEs, CAR-T cells or ADCs
[9,44]; thereby avoiding unnecessary toxicities to non-responders and limiting treatment cost.
Surprisingly, expression of TROP2 (encoded by *TACSTD2*) is generally low both in the untreated
and relapse patient samples. This questions the potential of Trodelvy® (Sacituzumab govitecan) or
Datroway® (Datopotamab deruxtecan) in SCLC patients irrespective of their disease stage.

Based on our observations, SEZ6 appears promising for two of the ADCs in clinical
development: ABBV-011 and ABBV-706. Yet, like DLL3, SEZ6 is a downstream transcriptional
target of ASCL1 [43,44] (see Figure 2) and is therefore likely to be targeted more effectively in
untreated patients. Our systematic screening for other cell surface targets reveals that CEA
(carcinoembryonic antigen) encoded by *CEACAM5* and *CEACAM6* is highly expressed in multiple
samples from relapse patients, suggesting the potential of CEA-ADCs in development [57-59]. An
advantage of targeting CEA is that, in addition to measuring CEACAAM5/6 transcripts and by
IHC in tumor samples, clinical trials could include CEA blood sampling [60,61]. Of note, we find that
TRPM5, which encodes a sodium channel transporter, is selectively expressed in the POU2F3-
expressing tumors and that GFRA1 (Glial Cell Line-Derived Neurotrophic Factor Receptor Alpha)
and SEMA6A (encoding one of the semaphorins) [25] are specifically expressed in NEUROD1
expressing SCLC (Figure 2), suggesting they might be considered as novel cell surface targets for
SCLC.

As we consider moving toward precision medicine, multivariable analyses will be needed.
For instance, for a given cell surface target, clinicians may have to choose between an ADC with
a TOP1 or a tubulin payload or a T-cell engager (BiTE). One could propose guiding payload choice
with at least two biomarkers: SLFN11 and ABC drug efflux transporters. If the tumor expresses
high SLFN11 and drug efflux transporters, TOP1 inhibitors will likely be superior to a tubulin
inhibitor ADC, and conversely a tubulin inhibitor ADC payload would be logical for tumors with
low SLFN11 and ABC transporters expression. For immune checkpoint inhibitors and BiTEs, one
could consider testing the predictive value of high APM and Immune signatures, which would
reflect the presence of immune cells in the tumor.

Ultimately, we envision that the "*MyPatient*" tab of SCLC_TumorMiner could be
developed for clinical practice as bulk RNA-seq and DNA-seq are readily performed on sample
biopsies. Sequencing results (FASTQ files) and clinical parameters could be loaded into
SCLC_TumorMiner. The "*My_Patient*" tool would provide a medical interface displaying a
patient's genomic parameters, dictating which therapy would be most appropriate and predicting
risk-factors. Based on our analyses, RNA-seq values can readily be set to separate samples. RNA-
seq expression value (log2(FPKM+1) > 5 and <3 discriminate between high and low gene
expression, respectively. Tumor_Miner allows the display and comparison of the molecular data
of the given patient in the context the historical cohort of SCLC patients. Additionally, the
Tumor_Miner architecture and interface could be readily expanded by integrating protein
expression (proteomics) and epigenetic (DNA methylome) parameters as we have already done in
our CellMiner websites (https://discover.nci.nih.gov/).

LIMITATIONS OF THE STUDY

Adding more samples with annotated data would further increase the statistical power of
SCLC_TumorMiner. However, our experience has been that material transfer agreements for
sharing FASTQ files and clinical information between institutions and individual researchers
remain a limiting factor. SCLC was not included in TCGA because surgical tumors biopsies are
not commonly done in SCLC patients. Obtaining high quality samples in sufficient quantity should
be a priority to obtain the necessary omics and pathological data required to precisely analyze
individual patients in a disease which remains currently lethal within months. Different genomic
platforms were used in our SCLC_TumorMiner database. This creates batch effects, which are
difficult to correct and can lead to erroneous conclusions unless one is aware of it. One solution is
to harmonize the technical platforms and avoid changing them without the absolute need to do so.
Although navigating websites remains a challenge, the rapid advance of Artificial Intelligence (AI)
is likely to humanize the website interface.
645

MATERIAL AND METHODS

Data content and Infrastructure
Patient clinical and genomics data were collected from our DTB clinic at the Center for Cancer
Research, NCI-NIH, from the University of Rochester (RU), the University of Cologne (UoC) and
Tongji university (TU). Our genomics patient data currently includes a total of 235 samples from
212 patients with RNA-seq and/or whole exome seq (WES) samples. The NIH samples were
obtained from patients enrolled in IRB-approved protocols and had consented to the analysis of
their tumors.
Clinical data from NIH and collaborators were gathered and manually curated by a team
of DTB physicians. They integrate patient demographics, survival, treatment with response and
sample information including sample type, biopsy site, tumor stage and pathology indicators (see
Table S2). NCI patient samples were correlated with treatment response, and for each patient we
selected for the first drug(s) or treatment(s) post-sampling as well as its response class (CR:
complete response, PR: partial response, PD: progressive disease or SD: stable disease). Due to
the current limited number of treatments, we decided that the response will be of value 1 (good
responder: PR, CR) or 0 (bad responder: SD, PD).
All genomics patient data were processed with our on-site data-handling pipelines using
the raw data. We generated 235 RNA-seq and 114 WES normalized data. Treatment response were
available for 21 patients. Table S9 shows details by disease, data sets and experiments.
We also computed for each sample gene expression Hallmark enrichment and genomics signatures
scores such as the NE, RepStress, EMT, APM, the level of stroma and immune infiltrate
contamination or a new Y chromosome signature scores. Table S10 presents the list of signatures
as well as the phenotypic data quarriable on TumorMiner.
SCLC-TumorMiner has similar design as the CellMinerCDB tool and websites
(https://discover.nci.nih.gov/) dedicated to mining pharmaco-genomics data across broad ranges
of cancer cell line sets and databases. SCLC-TumorMiner is implemented as a shiny application
and   hosted   using   Posit   Connect   on   CBIIT   servers   at
https://discover.nci.nih.gov/SclcTumorMinerCDB/.

674
675

RNA sequencing
The NIH/NCI and Rochester university (RU) samples were made from FFPE samples using the Illumina access protocol whereas the Universities of Cologne (UoC) and Tongji university (TU) were from Frozen using PolyA protocol. All the SCLC samples were processed with other diseases samples such as Adrenal cortical, Pancreatic NET or GINET samples by running the NCI CCBR RNAseq pipeline (https://github.com/skchronicles/RNA-seek.git). STAR (2.7.6a) was run to map reads to hg38 reference genome (release 36. Then RSEM was used to have gene expression values in log2(FPKM+1). We applied the "RemoveBatchEffect" function from the package Limma to remove the impact of the library preparation protocols (polyA, total RNA or access) and keeping in consideration the diseases differences.

Mutations
Somatic mutations were based on paired tumor-normal samples and ran with the CCBR exome sequencing pipeline (https://github.com/mtandon09/CCBR_GATK4_Exome_Seq_Pipeline). Briefly, BWA MEM (version 0.7.17) was run to map reads to the hg38 reference genome. Then Mutect2 in GATK 4.2 was used to call the variants. Variants with 6 or more total filtered reads were selected for display. We used Annovar (hg38 version) to annotate the mutations, including their predicted effect on amino-acid sequence and whether they are predicted to be deleterious, as well as their prevalence in the ExAC normal population. A mutation was called deleterious if its predicted effect on the RNA sequence was frameshift, nonsense or splicesense, or a missense mutation with a SIFT score <0.05 or PolyPhen2 HDIV score>=0.85. We selected all deleterious variants for the mutation gene level score.
        The fraction of alternate alleles (the mutation ratio) was used as the probability that any cell has that mutation. Furthermore, we assumed that the probability that any mutation is present in a gene is independent of the probability of any other mutation on that gene. Given those assumptions, we calculate the overall mutation probability for that gene in any particular cell as:
        $1-(1-p1)*(1-p2)*...*(1-pn)$
where p1, p2, ..., pn are the mutation ratios for each individual mutation on that gene. 1-pj is the probability that jth mutation is not present for a particular cell. Thus, the product is the probability that none of the mutations are present and 1 minus the product is the probability that at least one mutation is present. The expression gives a number that is between 0 and 1, which we multiply by 100 to get the mutation score for that gene.

Tumor Mutation Burden (TMB)
The TMB was computed using the R package MAFtools [62] based on Mutect2 somatic variants that have variant allele frequency (VAF) >= 5%

Genomic data for survival
If a patient has multiple samples, we select the primary tumor sample (if available) for survival analysis or we choose one based on biopsy site. The list of selected samples is downloadable from the website.

Unsupervised Clustering Approach
We implemented unsupervised clustering based on Non-Negative Matrix Factorization (NMF) algorithm using log-transformed normalized RNA-seq data to classify SCLC patients into molecular subtypes. NMF uses dimensionality reduction approach in which it decomposes the

721 gene expression matrix X (n × m), into the product of two lower-dimension non-negative matrices
722 (See Equation 1), where n represents number of patients and m is number of genes.

$$X \approx WH \qquad (1)$$

with X = n × m matrix, W = n × k matrix, and H = m × k, where k is a latent factor; optimal number
of factors (k) was determined based on matrix stability using cophenetic correlation coefficient.
We selected only protein-coding genes for both untreated and relapse SCLC samples, then we
selected the top 5000 highly variable genes for both datasets using a non-parametric approach,
Median Absolute Deviation (MAD) using "mad" function in R (version 4.2.3). After which, we
performed consensus NMF clustering evaluating values of k ranging from 2 to 8, based on
cophenetic correlation coefficient. k = 4 provided most stable and robust clusters for both untreated
and relapse SCLC datasets (See Figure S4). Clustering was performed using "NMF" library in R
(version 4.2.3).

Pairwise Differential Expression Analysis

After defining the NMF subgroups, differential gene analyses were performed using the "Limma"
package in R (version 4.2.3). To determine the up or down-regulated genes across the NMF-
subgroups, we performed pairwise comparisons for untreated and relapse SCLC separately. This
included all possible combinations of comparisons: NMF1 vs NMF2, NMF1 vs NMF3, NMF3 vs
NMF4, NMF2 vs NMF3, NMF2 vs NMF4, NMF3 vs NMF4. In addition, each subgroup was
compared with the aggregated sum of other subgroups, for instance: NMF1 vs Others
(NMF2+NMF3+NMF4), NMF2 vs Others (NMF1+NMF3+NMF4), NMF3 vs Others
(NMF1+NMF2+NMF4) and NMF4 vs Others (NMF1+NMF2+NMF3).
Subsequently, we utilized these pairwise comparisons to identify subgroup-specific genes
for each NMF cluster for both untreated and relapse samples. Significantly upregulated genes in
NMF1 in comparison to NMF2, NMF3 and NMF4 with $\log_2FC >1$ and p-value < 0.05 are listed
as NMF1-specific genes. Similarly, this criterion was implemented for the selection of specific
genes for the NMF2, NMF3 and NMF4 subgroups. The subgroup-specific genes identified using
this approach were used to construct NMF-derived gene expression signatures.

Gene Set Enrichment Analysis (GSEA)

To investigate the functional relevance of genes we performed pre-ranked Gene Set Enrichment
Analysis (GSEA). From previous analysis, we obtained pairwise comparisons for each NMF
groups and used t-statistic parameter, which provided magnitude and direction of differential
expression to rank the genes. Ranked genes were then analyzed using the fGSEA R package
(version 4.2.3) with REACTOME gene sets [https://www.gsea-
msigdb.org/gsea/msigdb/human/collections.jsp#C5]. fGSEA is a fast implementation of the
GSEA algorithm providing normalized enrichment scores (NES) and adjusting for multiple testing
using permutation-based p-values and false discovery rate (FDR).

Statistical Methods

Correlations, scatter plots, heatmaps, survival analysis, regression model predictions, and pathway
enrichment scores were generated using the R-Project for Statistical Computing based mostly on
the R packages: ggplot2, plotly, heatmaply, survival, glmnet, GSVA and GSEABase. These
packages are downloadable from the comprehensive R archive network (CRAN) or Bioconductor
websites.

Table 1. SCLC samples and Datasets
768

		Untreated		Relapse		
		UoC	**TU**	**NCI**	**RU**	**Total**
Number of samples		56	98	50	31	235
Genomics	RNA Seq	56	98	50	31	235
	WES	0	98	16	0	114
Prior Treatment	Yes	0	0	50	31	81
	No	52	98	0	0	150
	Unknown	4	0	0	0	4
Biopsy Site	Lung	55	98	7	10	170
	Liver	0	0	21	1	22
	Lymph Node	0	0	10	11	21
	Pleura	1	0	1	4	6
	Adrenal Gland	0	0	4	0	4
	Mediastin	0	0	3	0	3
	Others	0	0	4	5	9
Tumor Stage	Stage 1	25	30	0	0	55
	Stage 2	7	26	0	0	33
	Stage 3	17	41	2	2	62
	Stage 4	7	1	48	14	70
	Unknown	0	0	0	15	15

779 **Figures with legends**

780
781
Figure 1. Introduction to SCLC_TumorMinerCDB (https://discover.nci.nih.gov/SclcTumorMinerCDB/). **A**.
Snapshot of the introductory "2 variable Analyses" display (1). Both the horizontal and vertical axes offer a choice of
"Data Type" shown in the upper inset. Users must type their choice in the "Identifier" box (here ASCL1 for the
horizontal axis and POU2F3 for the vertical axis). Other options include choosing the "Range" for the axes and the
"Select Subsets" to "include" or "exclude". The + sign and folder indicate that further selections can be made such as
choosing the "Dataset" to display. To display more than one dataset, users need to press the Command key while
clicking the individual datasets (lower inset). Other features of SCLC_TumorMinerCDB include: (2) "Multivariable
Analyses" with examples shown in panel B and Figure 2; (3) "Expression Distribution" across the 4 Datasets with
example shown in Figure S1; (4) "Mutation variants"; (5) "Prognosis Biomarkers" with example shown in Figures 5
and S4; (6) "Predictive Biomarkers" with example shown in Figure 6; (7) "My Patient" to display the genomic
characteristics of a selected patient with example shown in Figures 7 and S6. Additional features include the ability
to "Download Data" for further analyses (8) as shown in Figures 7, S1, S4 and S5, "Search IDs" (9), as well as a
detailed "Help" section (10). **B**. NAPY distribution of samples from untreated and relapse patients. Samples were
assigned to each NAPY group based on gene expression log2(FPKM+1) > 5.0 (see horizontal and vertical dotted lines
in panel A).
797

Figure 2. Multivariable Analyses showing co-regulation of the neuroendocrine (INSM1, CHGA), NAPY (NEUROD1, ASCL1, POU2F3 and YAP1), NOTCH (NOTCH1, NOTCH2, REST) and MYC (MYC, MYCL and MYCN) pathways. **A**. Default display for the "Multivariable Analyses" tab. Color scale goes from red to blue to reflect expression levels for each gene (rows) and patients (columns). Users can choose different data types (upper inset of panel A) and then type the "Response Identifier" to display as the first row on the heatmap. "Predictor Identifier (s)" need to be typed with a space between each. "Tissues of origin" can be selected. The default algorithm (bottom left) is "Display". Users can also perform Linear and Lasso analyses like in CellMinerCDB (https://discover.nci.nih.gov/cellminercdb/). **B-E**. Multivariable analyses for the NAPY genes used as "Response Identifiers". INSM1 is included in each analysis as a NE biomarker. **F**. Multivariable analysis for the Notch pathway. **G**. Differential expression of the 3 MYC genes in samples from untreated and relapse patients. Linear regression analyses for each of "Response Identifiers" (in red, 1st rows in panels B-F) show that the "Predictor Identifiers" are highly significant (Pearson correlations > 0.85 and p-values > 2.5e-65; see Figure S2C).

813
814
Figure 3. Classification of the untreated (primary) and relapse 245 SCLC samples comparing untreated (UoC and TU
datasets) and relapse samples (NCI and RU). **A-B**. Hierarchical clustering of the untreated and relapse samples based
on the expression of the four genes of the NAPY classification (NEUROD1, ASCL1, POU2F3 and YAP1 in red) and
of genes reflecting commonly used biomarkers for clinical neuroendocrine (NE) classification (INSM1 and CHGA)
as well as MYC, REST and 4 exemplary innate immune genes (HLA-A, B, C and B2M). **C**. Flow chart summarizing
the NMF approach for selecting 5,000 protein-coding transcripts based on MAD (Median Absolute Deviation) for the
untreated and relapse samples. **D-E**. Display of the NMF clusters for the untreated and relapse samples (D and E,
respectively). The percentages of samples within each NMF are indicated in parenthesis. NMFs for untreated and
relapse are abbreviated NMFu and NMFr, respectively. The rows show the top 30 genes for each NMF with log fold-
change > 1 and p-value < 0.05. Only 8 genes (13%) are common across untreated and relapse samples for the NMF1
and NMF2 clusters, as indicated by the oblique lines. **F-G**. NMF distribution of the canonical NAPY genes (red) and
reference NE genes as well as MYC and REST. **H**. Box plots comparing the differential expression of the NAPY
genes, MYC and REST across the 4 untreated (top) and relapse NMFs (bottom). Statistical comparisons were
performed by Student t-test. **** indicates p < 0.001; ** p < 0.01; * p < 0.05. Shaded area represent low expression
(log2(FPKM+1) <3).

Figure 4. Genomic signatures associated with each of the NMF's from the untreated and relapse patient biopsy samples. **A**. Cluster image map of the 8 NMFs (4 untreated and 4 relapse) and top Reactome pathways. **B**. Correlation between each of the NMF's and the eight genomic signatures of SCLC_TumorMinerCDB for the samples from untreated (left) and relapse (right) patients. Correlations were obtained using the "2 variable Analyses" of the SCLC_TumorMinerCDB website ("Plot Data" or "View Data" tabs for each of the NMF on the horizontal axis). For the untreated samples the corresponding NMF's were computed after limiting the analysis using "Select Subsets" for the untreated Datasets: TU + UoC. Conversely for the relapse samples, analyses were done using the corresponding relapse samples: NCI + UR. **C**. Snapshots of "2 Variable Analyses" using the 4 datasets (inset) and displaying the correlations between the NMF2r signature and the RepStress (upper left) and the Immune signatures (upper right). The bottom 2 panels show the correlations between NMF4r and the NE (bottom left) and APM (bottom right) signatures.

Blue = Low risk   Red = High risk
Number are p-values for Hazard Ration (HR) 2 groups

Figure 5. Exploration of prognosis biomarkers. Panels A and F are screen snapshot of the SCLC_TumorMinerCDB website. **A**. High NMF4r expression is a good prognosis biomarker for all 4 Data sets. **B-E**. Tabulated results for the prognosis value of each of the NMFs (panels B-C), for 3 signatures (panel D) and genes (panel E) for each of the patient group. Combined expression of SEZ6, SRSF1 and MKI67 is a bad prognosis biomarker for the combined untreated Data sets (TU and UoC). Numbers are p-values for Hazard Ratio (HR) and colors highlight significant value (p < 0.05) with low-risk parameter in blue and high-risk in red. **F**. Gene expression of more than one gene can be tested. Here expression of SEZ6, the splicing factor SRSF1 and the proliferation marker Ki67 are poor prognosis biomarkers in untreated patients (TU and UoC). The option "Run Lasso" highlighted in yellow at the bottom left of panel A enables the unbiased search of novel prognosis biomarkers among selected Gene Sets lists that are displayed upon clicking "Run Lasso". Users are invited to run this feature on the website.

Figure 6. Predictive biomarkers with focus on ADCs for untreated patients (left panels A, C, E, G & I) versus patients after relapse (right panels B, D, F, H & J). **A-B**. Heatmaps showing the differential expression of 8 selected ADC targets. Samples were clustered based on the ADC target surface receptor genes listed between panels A and B. DLL3 and TROP2 (TACSTD2) are both FDA-approved targets. DLK1, CEACAM5, CEACAM6, CLN18, SIZ6 and B7-H3 are displayed because of their selective expression in subsets of patients. The top parts of the figure show for each patient the biopsy site, clinical stage, sample type and dataset. The bottom part shows for each patient whether the sample is enriched for the neuroendocrine (NE) and antigen-presenting-machinery (APM) signatures. **C-D**. Relationship between the listed ADC antibody targets and the NMF subgroups (NMFu and NMFr stand for the NMF's of samples from untreated and relapse patients, respectively. The blue-shaded areas include samples with low gene expression (log2(FPKM+1) <3) and the pink-shaded areas samples with high expression (log2(FPKM+1) >5). **E-F**. Expression of intracellular drug targets/biomarkers in the different NMF subgroups. **G-H**. Expression of drug efflux transporters in the different NMF subgroups. **I-J**. Expression of major pro-apoptotic genes in the different NMF subgroups. Statistical comparisons were performed by Student t-test. **** indicates p < 0.001; ** p < 0.01; * p < 0.05.

Figure 7. MyPatient module of SCLC_TumorMinerCDB. **A**. Proposed use of the "MyPatient" module. Genomic analyses are performed immediately after biopsy and uploaded into TumorMiner to provide results guiding the patient treatment as genomic classification, prognostic and predictive biomarkers are automatically displayed using "MyPatient" tab. **B**. Snapshot of the "MyPatient" tab in SCLC_TumorMinerCDB. Users are prompted to enter a "patient identifier" or select a patient from the available table. **C**. Snapshot of the "MyPatient" tab in SCLC_TumorMiner following the selection of a given patient. Here is an example for patient "S00035".

Supplemental Figures

Figure S1 (related to Fig 1). **A**. Example of a "2 Variable Analysis" snapshot of the websites displaying the
correlation between ASCL1 expression and the NE signature. **B**. Distribution and differential expression of the NAPY
genes across the untreated (n = 154) and relapse (n = 81) patient samples. Colored areas define samples with significant
expression ((log2(FPKM+1) > 5.0 in pink) and low expression (log2(FPKM+1) <3 in blue). Statistical comparisons
were performed by Student t-test. **** indicates p < 0.001; ** p < 0.01; * p < 0.05. Data were downloaded from the
SCLC_TumorMinerCDB website using the download tabs. **C**. Differential expression of the two MYC genes (MYCL,
MYC) and of the two innate immune response genes HLA-A and B2M across the untreated (n = 154) and relapse (n
= 81) patient samples. Colored areas define samples with significant expression ((log2(FPKM+1) > 5.0 in pink) and
low expression (log2(FPKM+1) <3 in blue). Statistical comparisons were performed by Student t-test. **** indicates
p < 0.001; ** p < 0.01; * p < 0.05. Data were downloaded from the SCLC_TumorMinerCDB website using the
download tabs.

Figure S2 (related to Figs 1-2). **A**. Snapshot of the plot corresponding to the predictive value of the 8 variables ("Predictor Identifiers") listed in Figure 2A under "Multivariable Analyses". **B**. Snapshot of the "Technical Details" corresponding to the plot shown in panel A and the Heatmap shown in Figure 2A. **C**. Linear regression analyses for the Heatmaps shown in panels B-E of Figure 2. Pearson correlation coefficients and p-values were obtained from the "Plot" tab (as shown in panel A for Figure 2A). **D**. Expression of the Notch pathway canonical genes REST, NOTCH1-4 predict the expression of YAP1 (Pearson correlation (r) = 0.83; p-value = 2.4e-60).

Figure S3 (related to Figs 1). **A**. Snapshot of the "Expression distribution" tab from the SCLC_TumorMinerCDB website in each of the 4 datasets of SCLC_TumorMinerCDB. Gene expression (genexp) is expressed as log2(FPKM+1). Statistical comparisons are performed by Student t-test. **** indicates $p < 0.001$; ** $p < 0.01$; * $p < 0.05$. The "Download Plot Data" tab at the bottom left enables users to export the data. **B**. Snapshot of the "Mutation Variants" tab in the SCLC_TumorMinerCDB website. Number to the right indicate the percentage of samples with deleterious mutations.

Figure S4 (related to Fig. 3). **A**. Venn-diagram showing the overlap of Untreated (top 5000) and Relapse (top 5000) most variable genes, selected using Median Absolute deviation (MAD) approach. **B** and **C** NMF rank summary for untreated and relapse SCLC samples, respectively. The cophenetic correlation coefficient is plotted for range of clusters (k=2:8), depicting k=4 as optimal number of clusters for both cohorts. Consensus matrix heatmaps at k=4 show well defined and stable clusters. Right: pie charts showing the distribution of samples in each NMF cluster.

A

B

Figure S5 (matching Fig. 3). **A**. PrerankGSEA Reactome pathway analysis for the 4 NMF groups including the samples of untreated patients. **B**. PrerankGSEA Reactome pathway analysis for the 4 NMF groups including the samples of relapse patients.

921　**Figure S6 (related to Fig. 5)**. **A**. Good prognostic value of SLFN11 expression. **B**. Good prognostic value (risk factor)
922　of the APM signature. **C**. Poor prognosis value (risk factor) on SEZ6 expression.

Figure S7 (related to Fig. 6). Distribution of predictive biomarkers across the 235 samples comparing samples from untreated (n = 154) and relapse (n = 81) patients. Colored areas defined samples with significant expression ((log2(FPKM+1) > 5.0 in pink) and low expression (log2(FPKM+1) <3 in blue). **A**. Surface biomarkers for ADCs. **B**. Broad range for SLFN11 expression across patient samples and downregulation at relapse. **C**. Broad expression range for proapoptotic genes across untreated patient samples and significant downregulation at relapse. Statistical comparisons were performed by Student t-test. **** indicates p < 0.001; ** p < 0.01; * p < 0.05.

931
932
933   **Figure S8 (related to Fig. 6)**. Expression of CEACAM5 and SLFN11 across the 235 samples of SCLC_TumorMiner.
934   The figure is a snapshot of the website. The red dotted lines separate the samples with high gene expression
935   (log2(FPKM+1) > 5.
936

938
939 **Figure S9 (related to Fig. 7)**. Mean average NMF expression for each and all 81 samples from patients at relapse. A
940 mean average threshold value log2(FPKM+1) > 3.8 is sufficient to assign each sample to one of the 4 NMF's. The
941 gray area defines non-significant values.
942

REFERENCES

1. Rudin, C.M., Poirier, J.T., Byers, L.A., Dive, C., Dowlati, A., George, J., Heymach, J.V., Johnson, J.E., Lehman, J.M., MacPherson, D., et al. (2019). Molecular subtypes of small cell lung cancer: a synthesis of human and mouse model data. Nat Rev Cancer. 10.1038/s41568-019-0133-9.

2. Gay, C.M., Stewart, C.A., Park, E.M., Diao, L., Groves, S.M., Heeke, S., Nabet, B.Y., Fujimoto, J., Solis, L.M., Lu, W., et al. (2021). Patterns of transcription factor programs and immune pathway activation define four major subtypes of SCLC with distinct therapeutic vulnerabilities. Cancer Cell *39*, 346-360 e347. 10.1016/j.ccell.2020.12.014.

3. Qu, S., Fetsch, P., Thomas, A., Pommier, Y., Schrump, D.S., Miettinen, M.M., and Chen, H. (2022). Molecular Subtypes of Primary SCLC Tumors and Their Associations With Neuroendocrine and Therapeutic Markers. Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer *17*, 141-153. 10.1016/j.jtho.2021.08.763.

4. Pacheco, J., and Bunn, P.A. (2019). Advancements in Small-cell Lung Cancer: The Changing Landscape Following IMpower-133. Clin Lung Cancer *20*, 148-160 e142. 10.1016/j.cllc.2018.12.019.

5. Paz-Ares, L., Dvorkin, M., Chen, Y., Reinmuth, N., Hotta, K., Trukhin, D., Statsenko, G., Hochmair, M.J., Ozguroglu, M., Ji, J.H., et al. (2019). Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. Lancet *394*, 1929-1939. 10.1016/S0140-6736(19)32222-6.

6. Nabet, B.Y., Hamidi, H., Lee, M.C., Banchereau, R., Morris, S., Adler, L., Gayevskiy, V., Elhossiny, A.M., Srivastava, M.K., Patil, N.S., et al. (2024). Immune heterogeneity in small-cell lung cancer and vulnerability to immune checkpoint blockade. Cancer Cell. https://doi.org/10.1016/j.ccell.2024.01.010.

7. Tlemsani, C., Pongor, L., Elloumi, F., Girard, L., Huffman, K.E., Roper, N., Varma, S., Luna, A., Rajapakse, V.N., Sebastian, R., et al. (2020). SCLC-CellMiner: A Resource for Small Cell Lung Cancer Cell Line Genomics and Pharmacology Based on Genomic Signatures. Cell Rep *33*, 108296. 10.1016/j.celrep.2020.108296.

8. Ahn, M.-J., Cho, B.C., Felip, E., Korantzis, I., Ohashi, K., Majem, M., Juan-Vidal, O., Handzhiev, S., Izumi, H., Lee, J.-S., et al. (2023). Tarlatamab for Patients with Previously Treated Small-Cell Lung Cancer. New England Journal of Medicine *389*, 2063-2075. 10.1056/NEJMoa2307980.

9. Su, P.L., Chakravarthy, K., Furuya, N., Brownstein, J., Yu, J., Long, M., Carbone, D., Li, Z., and He, K. (2024). DLL3-guided therapies in small-cell lung cancer: from antibody-drug conjugate to precision immunotherapy and radioimmunotherapy. Mol Cancer *23*, 97. 10.1186/s12943-024-02012-z.

10. Simpson, K.L., Rothwell, D.G., Blackhall, F., and Dive, C. (2025). Challenges of small cell lung cancer heterogeneity and phenotypic plasticity. Nat Rev Cancer *25*, 447-462. 10.1038/s41568-025-00803-0.

11. Zhou, D., Byers, L.A., Sable, B., Smit, M.-A.D., Sadraei, N.H., Dutta, S., and Upreti, V.V. (2024). Clinical Pharmacology Profile of AMG 119, the First Chimeric Antigen Receptor T (CAR-T) Cell Therapy Targeting Delta-Like Ligand 3 (DLL3), in Patients with Relapsed/Refractory Small Cell Lung Cancer (SCLC). The Journal of Clinical Pharmacology *64*, 362-370. https://doi.org/10.1002/jcph.2346.

12. Rudin, C.M., Pietanza, M.C., Bauer, T.M., Ready, N., Morgensztern, D., Glisson, B.S.,
Byers, L.A., Johnson, M.L., Burris, H.A., 3rd, Robert, F., et al. (2017). Rovalpituzumab
tesirine, a DLL3-targeted antibody-drug conjugate, in recurrent small-cell lung cancer: a
first-in-human, first-in-class, open-label, phase 1 study. Lancet Oncol *18*, 42-51.
10.1016/S1470-2045(16)30565-4.
13. Thomas, A., and Pommier, Y. (2019). Targeting Topoisomerase I in the Era of Precision
Medicine. Clin Cancer Res *25*, 6581-6589. 10.1158/1078-0432.CCR-19-1089.
14. Zhang, B., Stewart, C.A., Wang, Q., Cardnell, R.J., Rocha, P., Fujimoto, J., Solis Soto,
L.M., Wang, R., Novegil, V., Ansell, P., et al. (2022). Dynamic expression of Schlafen 11
(SLFN11) in circulating tumour cells as a liquid biomarker in small cell lung cancer. Br J
Cancer. 10.1038/s41416-022-01811-9.
15. Jo, U., and Pommier, Y. (2022). Structural, molecular, and functional insights into Schlafen
proteins. Experimental & Molecular Medicine. 10.1038/s12276-022-00794-0.
16. Murai, J., Tang, S.-W., Leo, E., Baechler, S.A., Redon, C.E., Zhang, H., Al Abo, M.,
Rajapakse, V.N., Nakamura, E., Jenkins, L.M.M., et al. (2018). SLFN11 Blocks Stressed
Replication Forks Independently of ATR. Molecular Cell *69*, 371-384.e376.
10.1016/j.molcel.2018.01.012.
17. George, J., Lim, J.S., Jang, S.J., Cun, Y., Ozretic, L., Kong, G., Leenders, F., Lu, X.,
Fernandez-Cuesta, L., Bosco, G., et al. (2015). Comprehensive genomic profiles of small
cell lung cancer. Nature *524*, 47-53. 10.1038/nature14664.
18. Liu, Q., Zhang, J., Guo, C., Wang, M., Wang, C., Yan, Y., Sun, L., Wang, D., Zhang, L.,
Yu, H., et al. (2024). Proteogenomic characterization of small cell lung cancer identifies
biological insights and subtype-specific therapeutic strategies. Cell *187*, 184-203 e128.
10.1016/j.cell.2023.12.004.
19. Lissa, D., Takahashi, N., Desai, P., Manukyan, I., Schultz, C.W., Rajapakse, V., Velez,
M.J., Mulford, D., Roper, N., Nichols, S., et al. (2022). Heterogeneity of neuroendocrine
transcriptional states in metastatic small cell lung cancers and patient-derived models. Nat
Commun *13*, 2023. 10.1038/s41467-022-29517-9.
20. Murai, J., Thomas, A., Miettinen, M., and Pommier, Y. (2019). Schlafen 11 (SLFN11), a
restriction factor for replicative stress induced by DNA-targeting anti-cancer therapies.
Pharmacol Ther *201*, 94-102. 10.1016/j.pharmthera.2019.05.009.
21. Zoppoli, G., Regairaz, M., Leo, E., Reinhold, W.C., Varma, S., Ballestrero, A., Doroshow,
J.H., and Pommier, Y. (2012). Putative DNA/RNA helicase Schlafen-11 (SLFN11)
sensitizes cancer cells to DNA-damaging agents. Proc Natl Acad Sci U S A *109*, 15030-
15035. 10.1073/pnas.1205943109.
22. Pozo, K., Kollipara, R.K., Kelenis, D.P., Rodarte, K.E., Ullrich, M.S., Zhang, X., Minna,
J.D., and Johnson, J.E. (2021). ASCL1, NKX2-1, and PROX1 co-regulate subtype-specific
genes   in   small-cell   lung   cancer.   iScience   *24*,   102953.
https://doi.org/10.1016/j.isci.2021.102953.
23. Anderson, K.R., Torres, C.A., Solomon, K., Becker, T.C., Newgard, C.B., Wright, C.V.,
Hagman, J., and Sussel, L. (2009). Cooperative transcriptional regulation of the essential
pancreatic islet gene NeuroD1 (beta2) by Nkx2.2 and neurogenin 3. J Biol Chem *284*,
31236-31248. 10.1074/jbc.M109.048694.
24. Gosmain, Y., Marthinet, E., Cheyssac, C., Guerardel, A., Mamin, A., Katz, L.S., Bouzakri,
K., and Philippe, J. (2010). Pax6 controls the expression of critical genes involved in

1035   pancreatic alpha cell differentiation and function. J Biol Chem *285*, 33381-33393.
1036   10.1074/jbc.M110.147215.

1037   25.   Mastrantonio, R., You, H., and Tamagnone, L. (2021). Semaphorins as emerging clinical
1038   biomarkers and therapeutic targets in cancer. Theranostics *11*, 3262-3277.
1039   10.7150/thno.54023.

1040   26.   Huang, Y.H., Klingbeil, O., He, X.Y., Wu, X.S., Arun, G., Lu, B., Somerville, T.D.D.,
1041   Milazzo, J.P., Wilkinson, J.E., Demerdash, O.E., et al. (2018). POU2F3 is a master
1042   regulator of a tuft cell-like variant of small cell lung cancer. Genes Dev *32*, 915-928.
1043   10.1101/gad.314815.118.

1044   27.   Montoro, D.T., Haber, A.L., Biton, M., Vinarsky, V., Lin, B., Birket, S.E., Yuan, F., Chen,
1045   S., Leung, H.M., Villoria, J., et al. (2018). A revised airway epithelial hierarchy includes
1046   CFTR-expressing ionocytes. Nature *560*, 319-324. 10.1038/s41586-018-0393-7.

1047   28.   Ireland, A.S., Hawgood, S.B., Xie, D.A., Barbier, M.W., Lucas-Randolph, S., Tyson, D.R.,
1048   Zuo, L.Y., Witt, B.L., Govindan, R., Dowlati, A., et al. (2024). Basal cell of origin resolves
1049   neuroendocrine–tuft lineage plasticity in cancer. bioRxiv, 2024.2011.2013.623500.
1050   10.1101/2024.11.13.623500.

1051   29.   Wu, X.S., He, X.Y., Ipsaro, J.J., Huang, Y.H., Preall, J.B., Ng, D., Shue, Y.T., Sage, J.,
1052   Egeblad, M., Joshua-Tor, L., and Vakoc, C.R. (2022). OCA-T1 and OCA-T2 are
1053   coactivators of POU2F3 in the tuft cell lineage. Nature *607*, 169-175. 10.1038/s41586-022-
1054   04842-7.

1055   30.   Abdel Wadood, N., Hollenhorst, M.I., Elhawy, M.I., Zhao, N., Englisch, C., Evers, S.B.,
1056   Sabachvili, M., Maxeiner, S., Wyatt, A., Herr, C., et al. (2025). Tracheal tuft cells release
1057   ATP and link innate to adaptive immunity in pneumonia. Nature Communications *16*, 584.
1058   10.1038/s41467-025-55936-5.

1059   31.   Baine, M.K., Febres-Aldana, C.A., Chang, J.C., Jungbluth, A.A., Sethi, S., Antonescu,
1060   C.R., Travis, W.D., Hsieh, M.S., Roh, M.S., Homer, R.J., et al. (2022). POU2F3 in SCLC:
1061   Clinicopathologic and Genomic Analysis With a Focus on Its Diagnostic Utility in
1062   Neuroendocrine-Low SCLC. Journal of thoracic oncology : official publication of the
1063   International Association for the Study of Lung Cancer *17*, 1109-1121.
1064   10.1016/j.jtho.2022.06.004.

1065   32.   Shue, Y.T., Drainas, A.P., Li, N.Y., Pearsall, S.M., Morgan, D., Sinnott-Armstrong, N.,
1066   Hipkins, S.Q., Coles, G.L., Lim, J.S., Oro, A.E., et al. (2022). A conserved
1067   YAP/Notch/REST network controls the neuroendocrine cell fate in the lungs. Nat Commun
1068   *13*, 2690. 10.1038/s41467-022-30416-2.

1069   33.   Kim, J.W., Ko, J.H., and Sage, J. (2022). DLL3 regulates Notch signaling in small cell
1070   lung cancer. iScience *25*, 105603. 10.1016/j.isci.2022.105603.

1071   34.   Ireland, A.S., Micinski, A.M., Kastner, D.W., Guo, B., Wait, S.J., Spainhower, K.B.,
1072   Conley, C.C., Chen, O.S., Guthrie, M.R., Soltero, D., et al. (2020). MYC Drives Temporal
1073   Evolution of Small Cell Lung Cancer Subtypes by Reprogramming Neuroendocrine Fate.
1074   Cancer Cell *38*, 60-78 e12. 10.1016/j.ccell.2020.05.001.

1075   35.   Masuda, K., Yoshida, T., Motoi, N., Shinno, Y., Matsumoto, Y., Okuma, Y., Goto, Y.,
1076   Horinouchi, H., Yamamoto, N., Watanabe, S.-i., et al. (2025). Schlafen 11 Expression in
1077   Patients With Small Cell Lung Cancer and Its Association With Clinical Outcomes.
1078   Thoracic Cancer *16*, e15529. https://doi.org/10.1111/1759-7714.15529.

1079   36.   Jiang, L., Huang, J., Higgs, B.W., Hu, Z., Xiao, Z., Yao, X., Conley, S., Zhong, H., Liu,
1080   Z., Brohawn, P., et al. (2016). Genomic Landscape Survey Identifies SRSF1 as a Key

1081   Oncodriver in Small Cell Lung Cancer. PLoS Genet *12*, e1005895.
1082   10.1371/journal.pgen.1005895.

1083 37. Meric-Bernstam, F., Yuca, E., Evans, K.W., Zhao, M., Maejima, T., Karibe, T., Raso,
1084   M.G., Tang, X., Zheng, X., Rizvi, Y.Q., et al. (2025). Antitumor Activity and Biomarker
1085   Analysis for TROP2 Antibody–Drug Conjugate Datopotamab Deruxtecan in Patient-
1086   Derived Breast Cancer Xenograft Models. Clinical Cancer Research *31*, 573-587.
1087   10.1158/1078-0432.Ccr-24-1948.

1088 38. Blackhall, F., Jao, K., Greillier, L., Cho, B.C., Penkov, K., Reguart, N., Majem, M.,
1089   Nackaerts, K., Syrigos, K., Hansen, K., et al. (2021). Efficacy and Safety of
1090   Rovalpituzumab Tesirine Compared With Topotecan as Second-Line Therapy in DLL3-
1091   High SCLC: Results From the Phase 3 TAHOE Study. Journal of thoracic oncology :
1092   official publication of the International Association for the Study of Lung Cancer *16*, 1547-
1093   1558. 10.1016/j.jtho.2021.02.009.

1094 39. Dowlati, A., Chiang, A.C., Cervantes, A., Babu, S., Hamilton, E., Wong, S.F., Tazbirkova,
1095   A., Sullivan, I.G., van Marcke, C., Italiano, A., et al. (2025). Phase 2 Open-Label Study of
1096   Sacituzumab Govitecan as Second-Line Therapy in Patients With Extensive-Stage SCLC:
1097   Results From TROPiCS-03. Journal of thoracic oncology : official publication of the
1098   International Association for the Study of Lung Cancer. 10.1016/j.jtho.2024.12.028.

1099 40. Sun, N.Y., Kumar, S., Kim, Y.S., Varghese, D., Mendoza, A., Nguyen, R., Okada, R.,
1100   Reilly, K., Widemann, B., Pommier, Y., et al. (2024). Identification of DLK1, a Notch
1101   ligand, as an immunotherapeutic target and regulator of tumor cell plasticity and
1102   chemoresistance in adrenocortical carcinoma. bioRxiv. 10.1101/2024.10.09.617077.

1103 41. Han, K., Chen, Y., Sun, X., Wen, L., Wu, Y., Chen, S., Wei, L., Yu, J., Zeng, T., Jiang, L.,
1104   and Tan, L. (2025). Combining serum CDK1 with tumor markers for the diagnosis of small
1105   cell lung cancer. Clin Transl Oncol *27*, 2005-2013. 10.1007/s12094-024-03722-y.

1106 42. Muley, T., Herth, F.J., Heussel, C.P., Kriegsmann, M., Thomas, M., Meister, M.,
1107   Schneider, M.A., Wehnl, B., Mang, A., and Holdenrieder, S. (2024). Prognostic value of
1108   tumor markers ProGRP, NSE and CYFRA 21-1 in patients with small cell lung cancer and
1109   chemotherapy-induced remission. Tumour Biol *46*, S219-S232. 10.3233/TUB-230016.

1110 43. Kudoh, S., Tenjin, Y., Kameyama, H., Ichimura, T., Yamada, T., Matsuo, A., Kudo, N.,
1111   Sato, Y., and Ito, T. (2020). Significance of achaete-scute complex homologue 1 (ASCL1)
1112   in pulmonary neuroendocrine carcinomas; RNA sequence analyses using small cell lung
1113   cancer cells and Ascl1-induced pulmonary neuroendocrine carcinoma cells. Histochem
1114   Cell Biol *153*, 443-456. 10.1007/s00418-020-01863-z.

1115 44. Ajay, A., Wang, H., Rezvani, A., Savari, O., Grubb, B.J., McColl, K.S., Yoon, S., Joseph,
1116   P.L., Kopp, S.R., Kresak, A.M., et al. (2025). Assessment of targets of antibody drug
1117   conjugates in SCLC. NPJ Precis Oncol *9*, 1. 10.1038/s41698-024-00784-7.

1118 45. Sun, F.D., and Das, M.S. (2024). Targeting the EZH2-SLFN11 pathway-a lesson in
1119   developing molecularly-informed treatments for recurrent small cell lung cancer.
1120   Translational Cancer Research *13*, 6608-6612. 10.21037/tcr-24-1755.

1121 46. Thomas, A., Tanaka, M., Trepel, J., Reinhold, W.C., Rajapakse, V.N., and Pommier, Y.
1122   (2017). Temozolomide in the Era of Precision Medicine. Cancer Res *77*, 823-826.
1123   10.1158/0008-5472.CAN-16-2983.

1124 47. Schultz, C.W., Zhang, Y., Elmeskini, R., Zimmermann, A., Fu, H., Murai, Y., Wangsa, D.,
1125   Kumar, S., Takahashi, N., Atkinson, D., et al. (2023). ATR inhibition augments the

1126   efficacy of lurbinectedin in small-cell lung cancer. EMBO Molecular Medicine *15*, e17313.
1127   https://doi.org/10.15252/emmm.202217313.

1128   48.   Polley, E., Kunkel, M., Evans, D., Silvers, T., Delosh, R., Laudeman, J., Ogle, C., Reinhart,
1129   R., Selby, M., Connelly, J., et al. (2016). Small Cell Lung Cancer Screen of Oncology
1130   Drugs, Investigational Agents, and Gene and microRNA Expression. J Natl Cancer Inst
1131   *108*. 10.1093/jnci/djw122.

1132   49.   Chan, J.M., Quintanal-Villalonga, Á., Gao, V.R., Xie, Y., Allaj, V., Chaudhary, O.,
1133   Masilionis, I., Egger, J., Chow, A., Walle, T., et al. (2021). Signatures of plasticity,
1134   metastasis, and immunosuppression in an atlas of human small cell lung cancer. Cancer
1135   Cell *39*, 1479-1496.e1418. https://doi.org/10.1016/j.ccell.2021.09.008.

1136   50.   Stewart, C.A., Gay, C.M., Xi, Y., Sivajothi, S., Sivakamasundari, V., Fujimoto, J.,
1137   Bolisetty, M., Hartsfield, P.M., Balasubramaniyan, V., Chalishazar, M.D., et al. (2020).
1138   Single-cell analyses reveal increased intratumoral heterogeneity after the onset of therapy
1139   resistance in small-cell lung cancer. Nature Cancer. 10.1038/s43018-019-0020-z.

1140   51.   Zhang, W., Girard, L., Zhang, Y.A., Haruki, T., Papari-Zareei, M., Stastny, V., Ghayee,
1141   H.K., Pacak, K., Oliver, T.G., Minna, J.D., and Gazdar, A.F. (2018). Small cell lung cancer
1142   tumors and preclinical models display heterogeneity of neuroendocrine phenotypes. Transl
1143   Lung Cancer Res *7*, 32-49. 10.21037/tlcr.2018.02.02.

1144   52.   Qi, J., Zhang, J., Liu, N., Zhao, L., and Xu, B. (2022). Prognostic Implications of Molecular
1145   Subtypes in Primary Small Cell Lung Cancer and Their Correlation With Cancer
1146   Immunity. Frontiers in oncology *12*, 779276. 10.3389/fonc.2022.779276.

1147   53.   Borromeo, M.D., Savage, T.K., Kollipara, R.K., He, M., Augustyn, A., Osborne, J.K.,
1148   Girard, L., Minna, J.D., Gazdar, A.F., Cobb, M.H., and Johnson, J.E. (2016). ASCL1 and
1149   NEUROD1 Reveal Heterogeneity in Pulmonary Neuroendocrine Tumors and Regulate
1150   Distinct Genetic Programs. Cell Rep *16*, 1259-1272. 10.1016/j.celrep.2016.06.081.

1151   54.   Choudhury, N.J., Lai, W.V., Makhnin, A., Heller, G., Eng, J., Li, B., Preeshagul, I., Santini,
1152   F.C., Offin, M., Ng, K., et al. (2024). A Phase I/II Study of Valemetostat (DS-3201b), an
1153   EZH1/2 Inhibitor, in Combination with Irinotecan in Patients with Recurrent Small-Cell
1154   Lung Cancer. Clin Cancer Res *30*, 3697-3703. 10.1158/1078-0432.CCR-23-3383.

1155   55.   Tang, S.W., Thomas, A., Murai, J., Trepel, J.B., Bates, S.E., Rajapakse, V.N., and
1156   Pommier, Y. (2018). Overcoming Resistance to DNA-Targeted Agents by Epigenetic
1157   Activation of Schlafen 11 (SLFN11) Expression with Class I Histone Deacetylase
1158   Inhibitors. Clin Cancer Res *24*, 1944-1953. 10.1158/1078-0432.CCR-17-0443.

1159   56.   Rudin, C.M., Reck, M., Johnson, M.L., Blackhall, F., Hann, C.L., Yang, J.C., Bailis, J.M.,
1160   Bebb, G., Goldrick, A., Umejiego, J., and Paz-Ares, L. (2023). Emerging therapies
1161   targeting the delta-like ligand 3 (DLL3) in small cell lung cancer. J Hematol Oncol *16*, 66.
1162   10.1186/s13045-023-01464-y.

1163   57.   Paul, S., Konig, M.F., Pardoll, D.M., Bettegowda, C., Papadopoulos, N., Wright, K.M.,
1164   Gabelli, S.B., Ho, M., van Elsas, A., and Zhou, S. (2024). Cancer therapy with antibodies.
1165   Nat Rev Cancer *24*, 399-426. 10.1038/s41568-024-00690-x.

1166   58.   Shen, L., Sun, X., Chen, Z., Guo, Y., Shen, Z., Song, Y., Xin, W., Ding, H., Ma, X., Xu,
1167   W., et al. (2024). ADCdb: the database of antibody-drug conjugates. Nucleic Acids Res
1168   *52*, D1097-D1109. 10.1093/nar/gkad831.

1169   59.   DeLucia, D.C., Cardillo, T.M., Ang, L., Labrecque, M.P., Zhang, A., Hopkins, J.E., De
1170   Sarkar, N., Coleman, I., da Costa, R.M.G., Corey, E., et al. (2021). Regulation of
1171   CEACAM5 and Therapeutic Efficacy of an Anti-CEACAM5-SN38 Antibody-drug

Conjugate in Neuroendocrine Prostate Cancer. Clin Cancer Res *27*, 759-774.
10.1158/1078-0432.CCR-20-3396.

60.   Du, Y., Wen, Y., and Huang, J. (2024). Analysis of variation of serum CEA, SCC,
CYFRA21-1 in patients with lung cancer and their diagnostic value with EBUS-TBNA. J
Med Biochem *43*, 363-371. 10.5937/jomb0-37083.

61.   Yu, M., and Wang, P. (2024). A comparative study of ProGRP and CEA as serological
markers in small cell lung cancer treatment. Discov Oncol *15*, 485. 10.1007/s12672-024-
01323-3.

62.   Mayakonda, A., Lin, D.C., Assenov, Y., Plass, C., and Koeffler, H.P. (2018). Maftools:
efficient and comprehensive analysis of somatic variants in cancer. Genome Res *28*, 1747-
1756. 10.1101/gr.239244.118.

1183