Introducing furanocoumarin biosynthetic genes in tomato results in coumarins accumulation and impacted growth

Alexandre Bouillé[1], Cloé Villard[1], Gianni Galati[1], Marwa Roumani[1], Aude Fauvet[1], Jérémy Grosjean[1], Lennart Hoengenaert[2,3], Wout Boerjan[2,3], John Ralph[4,5], Frédérique Hilliou[6], Christophe Robin[1], Alain Hehn[1], Romain Larbat[7*]

Corresponding author*
Phone contact: +33 (0)2 41 20 44 03

1- Université de Lorraine, INRAE, LAE, 54000 Nancy, FR
2- Department of Plant Biotechnology and Bioinformatics, Ghent University, Ghent, BE
3- Center for Plant Systems Biology, VIB, Ghent, BE
4- Department of Biochemistry, University of Wisconsin, Madison, WI 53706, USA
5- Department of Energy Great Lakes Bioenergy Research Center, The Wisconsin Energy Institute, University of Wisconsin, Madison, WI 53726, USA
6- INRAe, Institut Sophia Agrobiotech, Université Côte D'Azur, CNRS, Sophia Antipolis, France
7- Institut Agro, University of Angers, INRAE, IRHS, SFR QUASAV, F-49000 Angers, FR

#scopoletin #coumarins #furanocoumarins #tomato plant #growth #psoralen #specialized metabolism

Graphical abstract

Abstract

Over the past three decades, efforts to decipher plant metabolism have shed light on key enzymes driving specialized metabolite biosynthesis. Although only few pathways have been completely investigated to date, their characterization paves the way for exploring the potential effects of specialized metabolites on plant physiology. Among them is the linear furanocoumarin pathway, which was recently completed to produce up to psoralen. In this study, we report the first metabolic engineering of the linear furanocoumarin pathway to enable artificial psoralen production in tomato, through the integration of four genes coding for the enzymes: Umbelliferone Synthase, Demethylsuberosin Synthase, Marmesin Synthase and Psoralen Synthase. Interestingly, coumarins were produced instead of furanocoumarins. Using morphophysiological, metabolomic, and transcriptomic analyses, we suggest how coumarins, particularly scopoletin, can impact growth and affect plant physiology, even at low concentrations.

As coumarins have increasingly attracted interest for agricultural applications due to

their minimal environmental impact, this work both expands and challenges their

potential by highlighting the physiological costs and benefits they may impose on tomato.

1. Introduction

The around 400 million years of evolution since the terrestrialization of plants have driven

the emergence of numerous mechanisms for interacting with an ever-evolving

environment (Morris *et al.,* 2018). The production of specialized metabolites is a key

example of such adaptation as many of these compound play vital roles in enabling

plants to cope with biotic and abiotic stresses (Maeda & Fernie, 2021; Weng *et al.,* 2021).

Investment of resources and energy in processes that promote plant defense,

particularly the biosynthesis of specialized metabolites, may come at the expense of

processes that support plant growth (He *et al.,* 2022). Plants must therefore balance the

trade-off between investing in the costly biosynthesis of specialized metabolites and

allocating resources for growth (Neilson *et al.,* 2013).

Land plants produce a wide range of specialized metabolites (Chassagne *et al.,* 2019).

Among them, linear furanocoumarins are defensive metabolites restricted to certain

lineages (Sarker & Nahar, 2017). Its pathway has been largely elucidated over the past

two decades, particularly in Apiaceae and Moraceae (**Fig. 1**) (Bouillé et al., 2025).

Furanocoumarins are derived from the ubiquitous coumarin umbelliferone, which itself

is derived from *p*-coumaroyl-CoA (Vialart *et al.,* 2012; Roselli *et al.,* 2017). The

biosynthesis of linear furanocoumarins then starts with the C6-prenylation of

umbelliferone to produce demethylsuberosin by a prenyltransferase (Karamat *et al.,*

2014; Munakata *et al.,* 2016). Demethylsuberosin is then cyclized by marmesin synthase,

a cytochrome P450 (P450), into marmesin (Villard *et al.,* 2021; K. Wang *et al.,* 2024),

which is subsequently converted to psoralen via C7-decarboxylation through *syn*-

elimination of acetone by psoralen synthase, another P450 (Jian *et al.,* 2020; Larbat *et al.,*

2007, 2009). The psoralen backbone can then be hydroxylated at C5 or C8 by additional

P450s to yield bergaptol or xanthotoxol, respectively (**Fig. 1**) (Krieger *et al.,* 2018; Ji *et al.,*

2024). These structures can be further decorated by other enzymes, such as *O*-

methyltransferases, *O*-prenyltransferases and/or other P450s, to generate more complex

alkylated structures (Hehmann *et al.,* 2004; Zhao *et al.,* 2016; Han *et al.,* 2023; Bouillé *et*

al., 2025). Along with some of its derivatives, psoralen exhibits a broad spectrum of

defensive effects, mainly driven by its ability to form covalent bonds with nucleic acids

(*i.e.,* via photocycloaddition) and to alter herbivore defences (*e.g.,* through cytochrome

P450 inhibition) (Neal & Wu, 1994; Kitamura *et al.,* 2005). For such reasons, psoralen is

considered the first toxic metabolite in the linear furanocoumarin pathway.

In this study, we aimed to investigate the potential physiological and growth-related costs

associated with the production of a novel specialized metabolite in plants using

metabolic engineering. Considering the importance of psoralen in plant-environment

interactions and the unknown costs its biosynthesis may impose on plants, we chosen

to investigate the recently completed psoralen pathway. In that regard, using the

GoldenBraid multigenic assembly system (Sarrion-Perdigones *et al.,* 2013), we

integrated four linear furanocoumarin biosynthetic genes into the tomato genome

(*Solanum lycopersicum*, Solanaceae), a plant that does not naturally produce these

compounds, to enable the conversion of the ubiquitous *p*-coumaroyl-CoA into psoralen

(**Fig. 1**). Metabolic analyses of the transgenic tomato line (referred here to as "Psoralen

Pathway" or PP line) confirmed the detection of small quantities of psoralen, the

expected product, but also highlighted larger modifications in the accumulation of other

metabolite families, among them coumarins and particularly scopoletin. This

modulation of the metabolic composition among PP line was accompanied by major

transcriptomic changes and a significant modification in the biomass distribution. Our

results shed light on the physiological costs associated with the implementation of the

psoralen pathway in tomato and offer novel perspectives on the use of coumarins in

agriculture.

Figure 1: Simplified diagram of the psoralen pathway (PP) among linear furanocoumarin in some restricted angiosperms taxa. The pathway from *p*-coumaroyl-CoA to psoralen is framed, with the enzymes engineered in the tomato PP line encircled in purple and the specific genes inserted in the tomato genome in bold, while the carbon number on the furanocoumarin backbone is depicted in blue. Abbreviations: 2-OxoGlutarate dependent Dioxygenase (2OGD), COumarin SYnthase (COSY), PrenylTransferase (PT), CYtochrome P450s (CYP), US (*UMBELLIFERONE SYNTHASE*), DS (*DEMETHYLSUBEROSIN SYNTHASE*), MS (*MARMESIN SYNTHASE*), PS (*PSORALEN SYNTHASE*). Species: Apiaceae (*Angelica dahurica, Angelica sinensis, Pastinaca sativa, Petrolinum crispum, Peucedanum praeruptorum*), Rutaceae (*Ruta graveolens*), Brassicaceae (*Arabidopsis thaliana*), Moraceae (*Ficus carica*). From Larbat *et al.,* 2007, 2009; Vialart *et al.,* 2012; Karamat *et al.,* 2014; Munakata *et al.,* 2016; Roselli *et al.,* 2017; Vanholme *et al.,* 2019; Jian *et al.,* 2020; Villard *et al.,* 2021; Wang *et al.,* 2024.

2. Material and methods

2.1. Biological material and Standards

The tomato plants (*Solanum lycopersicum*) cultivar *West Virginia* 106 (Wva106) was used as the wild-type control and for plant transformation. Analytical standards were sourced from Sigma-Aldrich (Saint-Louis, MO, USA; for *p*-coumarate, scopoletin, and esculin), Extrasynthese (Lyon, France; for umbelliferone, psoralen, and esculetin), TransMIT (Giessen, Germany; for marmesin), Phytolab (Nantes, France; for scopolin), and R. Munakata from Kyoto University (demethylsuberosin).

2.2. Creation of the PP transgene

The Psoralen Pathway (PP) T-DNA was created by combining the genes controlling the 4-steps metabolic pathway from *p*-coumaroyl-CoA to psoralen (**Fig. 1**). The four genes were *Ps*Diox (ncbi: KY081678; Roselli *et al.,* 2017) coding for Umbelliferone Synthase (US), *Ps*PT1 (ncbi: KM017083; Munakata *et al.,* 2016) coding for Demethylsuberosin Synthase (DS), *Fc*CYP76F112 (ncbi: MW348922 ; Villard *et al.,* 2021) coding for Marmesin Synthase (MS) and *Ps*CYP71AJ3 (ncbi: C0SJS2 ; Larbat *et al.,* 2009) coding for Psoralen Synthase (PS). The selection marker used in the creation of the PP transgene was the *NEOMYCIN PHOSPHOTRANSFERASE* (*Ec*NPTII) gene conferring resistance to kanamycin, which originates from the plCSL11024 vector. The coding sequence of the five genes of interest were domesticated by removing internal restriction sites for *Bsa*I and *Bsm*BI, and by adding flanking extensions compatible with the GoldenBraid 2.0 multigene cloning system (GB2.0; Sarrion-Perdigones *et al.,* 2013). Each domesticated gene was cloned into the pUPD vector using the *Bsm*BI restriction enzyme, and assembled with the *Ca*MV P35S promoter (pUPD-35S, Addgene plasmid #68163) and the tNOS terminator (pUPD-tNOS, Addgene plasmid #68188) to form individual transcription unit (TU). TUs were assembled into the plasmid pDBGB1α1 or pDBGB1α2 by restriction/ligation using the restriction enzymes *Bsa*I or *Bsm*BI, respectively, and the T4 DNA ligase, as described in Sarrion-Perdigones *et al.,* 2011. The resulting TUs were further assembled two-by-two into pDGB1 α- or Ω-level plasmids, through restriction/ligation, to build the final PP T-DNA into pDGB1Ω1. Integrity of this final vector pDGB1Ω1_PP was validated by PCR analysis (see primers section 2.3), restriction analysis (**Fig. S1a**), and sequencing of the coding genes (Eco-Seq, Macrogen). PCR amplification was performed using the SapphireAmp® polymerase (Takara Bio) and following the reaction cycle: Initial denaturation (98°C for 5min), 35 cycles composed of a denaturation step (98°C for 10s),

an annealing step (55°C for 15s) and an elongation step (72°C for 10s), followed by a final

elongation step (72°C for 1min). Restriction of 1µg of plasmid DNA was performed using

*Bsa*I, *Bsm*BI, *Pvu*II, and *Eco*RI enzymes with the FastDigest system (Thermo Fisher

Scientific), following the manufacturer's recommendations.

2.3.   Plant transformation and regeneration of the PP line

The pDGB1Ω1_PP vector was co-transformed into *Agrobacterium tumefaciens* EHA105

with the replication helper vector pSOUP. The transformed cells were selected on solid

LB medium containing spectinomycin (100mg/L), tetracycline (5mg/L), and rifampicin

(20mg/L) (Hellens *et al.,* 2000; Sarrion-Perdigones *et al.,* 2013). A pre-culture was

performed overnight at 28°C, 200rpm, in 10mL of LB medium with appropriate

antibiotics. A 50mL culture was then started with 5mL of the pre-culture to reach an

OD_{600nm} of 0.4-0.6. The culture was centrifuged for 10 min at 7000 ×g, then resuspended

in 40mL of sterile water and diluted to 30mL to achieve an OD_{600nm} of 0.08. Axenic

cotyledon fragments approximately 4mm in size were harvested from 7-day-old tomato

seedlings grown under controlled conditions (24°C, 16h/8h light/dark cycles; $150µmol.m^{-2}.s^{-1}$,

Fitoclima 1200, Aralab). The cotyledon fragments were cultured for 24h at 24°C in

darkness before being co-incubated with the Agrobacteria solution for 30min at room

temperature (gentle agitation every 5min). The fragments were then dried on axenic paper

and placed adaxial side up on co-cultivation medium for 48h at 24°C in darkness.

Partially transformed tissues were selected on kanamycin-enriched callus medium

(150mg/L) in Petri dishes (15 fragments per dish) at 24°C (16h/8h light/dark cycles;

$150µmol.m^{-2}.s^{-1}$) and subcultured weekly. After 3 months of subcultures, differentiated

seedlings from calli were cut and transferred to rooting medium before being acclimated

in soil in phytotrons (24°C; 16h/8h light/dark cycles; $400 µmol.m^{-2}.s^{-1}$; 70% humidity).

Several heterozygous descendants from the regenerated tomato line were obtained by

self-pollination. The insertion of the 5 transgenes into the genome was assessed by PCR

(**Fig. S1b**) using genomic DNA extracted with the kit following the supplier's

recommendations   and   the   primers   described   as   follow: *Ps*Diox_F:

CTAGTCGAAATGGCTCCATCTCC;   *Ps*Diox_R: CGAAGATCCATTCCAAGCTCACATTTG;

*Ps*PT1_F: CTAGTCGAAATGGCTCAAACAATTATGC;   *Ps*PT1_R:

GAAGATCCATTCCAAGCTCACCTC;   *Fc*CYP76F112_F:

CTATTCTAGTCGAAATGGATATTTTCACCTC;   *Fc*CYP76F112_R:

CGAAGATCCATTCCAAGCCTAATGATG;   *Ps*CYP71AJ3_F:

CTATTCTAGTCGAAATGAAGATGCTTGAG;   *Ps*CYP71AJ3_R:

GATCGAAGATCCATTCCAAGCTCATC;   NPTII_F:

CAATTTACTATTCTAGTCGAAATGGTTGAACAAG;   NPTII_R:

GAAGATCCATTCCAAGCTCAGAAGAAC. The expression of the 5 genes in each individual

was confirmed by RT-PCR for each tomato line (**Fig. S1c**). Total RNAs was extracted using

the E.Z.N.A.® HP kit (Omega Bio-Tek) instructions, treated with DNase (AMPD1-1KT,

Sigma) to remove genomic DNA contaminants, and then reverse-transcribed into cDNA

using the High-Capacity RNA-to-cDNA Kit (Applied Biosystems™), according to the

supplier's recommendation.

2.4.   Plant cultivation and measurement of morpho-physiological traits

Four sets of plant cultivation and harvest were performed over a period of four months.

Three sets of 6-8 plants per genotype were used for transcriptomic and metabolomic

analyses (23 WT, 19 PP), and one additional set (23 WT, 19 PP) for morphophysiological

measurements, for a total of 46 WT and 38 PP plants. All plants were staggered under

identical conditions. The first three were used for the metabolomics and transcriptomics

analyses. The fourth series was dedicated to measuring morphometric traits, including

carbon and nitrogen allocation, and lignin composition/content in the stems. At harvest,

stem and petiole tissues were separated from the leaves and roots. For metabolomics

analyses, petioles and internodes were combined.

Tomato plants were grown in 1.5L pots with 0.5L of potting soil (Stender D400, NPK

14/10/18, pH 6.2, sieved to 5 mm). The climatic conditions in the growth chamber were

set at 24°C with 16h/8h light/dark cycles, LED illumination at $400\mu mol.m^{-2}.s^{-1}$, and 70%

humidity. Plants were watered to maintain 70% soil moisture three times a week with

osmosed water. Weekly, tomatoes were fertilized with horticultural fertilizer (50 mL at 2

$g.L^{-1}$) (9:16:36 of N:P:K, MultiTec™). Tomato plants were harvested at 28 days after sowing,

at the vegetative stage. Roots were gently washed with a brush and untangled for length

measurements. For metabolomic and transcriptomic analyses, tissues were rapidly

snap-frozen in liquid nitrogen before being finely ground and stored at -80°C. Leaf areas

without petioles were measured using a planimeter (Li-3100 C Area Meter, Li-Cor,

Lincoln, Nebraska, USA). Tissues used for carbon, nitrogen, nitrate and lignin content
measurement were dried in an oven at 70°C for 1 week and then finely ground using an
orbital mill (Pulverisette 6, Fritsch, France) before being stored at room temperature in a
dry-preserving package. Resulting powder was used for carbon, nitrogen and lignin
measurements.

2.5. Total nitrogen and carbon analyses
Total C and N contents in dry tissues were determined using 3mg of dry matter, according
to the Dumas method, with an elemental auto-analyzer (Unicube, Elementar,
Langenselbold, Germany).

2.6. Cell wall characterization
Lignin and its components were quantified according to Lu *et al.,* 2021. Briefly, cell wall
residues (CWR) were extracted using successive boiling in solvent baths. Lignin content
was quantified in percentage of CWR according to the Cysteine-Assisted Sulfuric Acid
(CASA) method. The lignin composition of CWR was measured using the thioacidolysis
method as described by Rolando *et al.,* 1992, and the released fragments were analyzed
using GC-MS at the VIB Metabolomics Core as described in (Hoengenaert *et al.,* 2022).
The composition of the lignin was analyzed as described in (Hoengenaert *et al.,* 2022).
Briefly, 20 mg of stems CWR were cellulase-digested and resuspended in DMSO. Scans
of HSQC NMR spectra (Heteronuclear Single-Quantum Coherence) ^{1}H–^{13}C short-range
correlation experiments were acquired for 6h, followed by a second 54h acquisition for
improve sensitivity.

2.7. Metabolites extraction
Frozen powders from leaves, stems and roots (100 mg FW) were extracted twice using
maceration (16h in the dark at 22°C with slow vertical agitation) followed by sonication
(37 kHz, 10min at room temperature), each time with 800 µL of methanol/deionized water
(80/20). The macerates were then centrifuged (10min, 16100 g, 22°C). The supernatants
(1.6 mL) were collected, dried (5-7h under vacuum, Concentrator Plus, Eppendorf) and
then resuspended in 100µL of methanol/deionized water (80/20). The extracts were
filtered (0.2 µm, Minisart® RC4 membrane, Sartorius, REF17821) and stored at -20°C.
Extracts were added with taxifolin (285µM) as an internal quantification standard.

2.8. Metabolomics analysis

Chromatographic analyses coupled with High Resolution Mass Spectrometry were

performed on a Thermo Scientific Vanquish UHPLC system equipped with a binary pump,

an autosampler, a temperature-controlled column (40°C) and a Diode Array Detector.

Extracted samples were separated on a XB-C18 Kinetex (150 × 2.1 mm, 2.6 μm)

(Phenomenex Inc., Torrance, CA, USA) using a gradient of mobile phase composed of

0.1% formic acid in water (A) and 0.1% formic acid in methanol (B) at a flow rate of 200

μL·min^{-1}. The separation of the compounds was performed using a 30 min mobile phase

gradient (A: B; v/v). The elution method is as follows: (90:10) at 0 min, (90:10) up to 2 min,

(70:30) up to 10 min, (5:95) up to 20 min remaining 5 min, (90:10) up to 26 min. The

molecules are detected according to their absorbance at different wavelengths between

190 and 800 nm.

HRMS[1] detection was performed on an Orbitrap IDX^{TM} (ThermoFisher Scientific, Bremen,

Germany) mass spectrometer in positive and negative electrospray ionization (ESI)

modes. The capillary voltages were set at 3.5 kV and 2.5 kV for positive and negative

modes, respectively. The source gases were set (in arbitrary unit min^{-1}) to 40 (sheath gas),

8 (auxiliary gas) and 1 (sweep gas) and the vaporizer temperature and ion transfer tube

was respectively 320°C and 275°C. Full scan MS[1] spectra were acquired from 120 to 1200

m/z at a resolution of 60,000. MS^2 analysis was performed on the pooled-samples vial

using the data dependent acquisition (DDA) mode and fragments were analyzed by

Orbitrap at a resolution of 15000. For this analysis, the AcquireX data acquisition

workflow was applied. The raw data were uploaded on the Compound discoverer 3.3

software (ThermoFisher Scientific, Bremen, Germany) to run the untargeted

metabolomics workflow (peak detection, chromatogram alignment and peak grouping in

features). In addition, the full MS^1–MS^2 dataset was exported as a mascot generic format

(mgf) file and analyzed through the Sirius 5 software (Dührkop *et al.,* 2019). Briefly, the

procedure implies the automatic annotation of molecular formula using SIRIUS and

ZODIAC (Dührkop *et al.,* 2019; Ludwig *et al.,* 2019). Then, molecular structure prediction

was proposed through CSI: FingerID (Hoffmann *et al.,* 2021). Finally, feature

was assigned to chemical families using CANOPUS (Djoumbou Feunang *et al.,* 2016;

Dührkop *et al.,* 2019; H. W. Kim *et al.,* 2021). Further annotation was performed using

Compound Discoverer 3.3 by predicting elemental composition, searching in local and

public mass and formula databases such as ChemSpider, HMDB, and LipidMaps, and

querying spectral databases including mzCloud, MoNA, and GNPS.

Formal identification of coumarins and furanocoumarins, was afforded via comparisons

with commercial standards according at least to their retention time (RT), mass (MS[1]) and

mass fragmentation (MS[2]) when available. A standard curve was established for the

scopoletin quantification using as internal standard the taxifolin (285µM).

2.9. Construction and analysis of RNAseq libraries

Total RNA from leaves and roots was extracted from 3 plants per genotype (WT and PP)

as previously described (Roumani *et al.,* 2022). RNA concentrations were determined

from their absorbance at 260 nm. Their purity was evaluated by the 260/280 nm

absorbance ratio, their quality was checked by electrophoresis on a 1% agarose gel and

their integrity by determining and considering an RNA Integrity Number (RIN) >7 (Agilent

BioAnalyzer 2100). Transcript data was obtained by RNA-Seq Illumina pair-end

sequencing (2× 150pb) using the NovaSeq 6000 methods for 30 million read pairs. Raw

data were cleaned using Trimmomatic (Bolger *et al.,* 2014). The quality of the sequences

was assessed using FastQC. The transcripts were mapped on the tomato genome

(ITAG2.4) using the BWA-MEM software (Li & Durbin, 2009). Count tables were generated

using the FeatureCounts software (Version 1.6.4) and Samtools merge workflows (Liao *et*

al., 2014). The differential gene expression was performed with the glmfit option within

the edgeR package for a FDR <0.05 (Robinson *et al.,* 2010) and using AskoR (Alves-

Carvalho *et al.,* 2021). We performed, within AskoR, a clustering using kmeans

classification algorithms (Coseq R package) (Godichon-Baggioni *et al.,* 2019). Functional

annotations, such as gene ontologies and interproscan features, were extracted from

ITAG2.4_protein_functional.gff3 file. Differentially expressed genes (DEGs) between WT

and PP lines DEGs were used to perform GO enrichment analysis (GOEA) restricted to

the Biological Processes on ShinyGO (version 0.80) using the "*Solanum lycopersicum*

genes SL3.0" and a FDR cutoff <0.05 (Ge *et al.,* 2020).

The results of the RNAseq differential analysis were confirmed by following the

expression level of 2 genes (*Sl*COSY and *Sl*PRCA) in addition to the 5 engineered genes in

leaves and roots of the PP and control tomato plants. The list of targeted genes and

primers used is given in **Table S2h**. The expression level was followed by RT-qPCR. For

that purpose, a total of 200 ng of DNA-free RNA samples was reverse transcribed in a final volume of 20 μL using the RNA to cDNA kit (Life technologies, Tokyo, Japan). cDNAs were then diluted 10 times. Gene expression was measured by qPCR on a CFX Duet Realtime PCR System (Bio-Rad). Each PCR reaction was made in 20 μL containing primer pairs at 0.3 μM, 40 ng of cDNAs in 1× Mesa Blue Premix (Eurogentec) and followed the amplification protocol consisting in a denaturing step at 95 °C for 30 s, 40 cycles at 95 °C for 5 s and 60 °C for 30 s. The gene expression of each sample was calculated with three analytical replicates and normalized to the two internal control genes *ACT* and *GAPDH*.

2.10. Statistical Analysis

Measured and calculated morphometric and physiological variables, as well as relative and absolute quantified metabolites were statistically analyzed by comparing WT and PP lines using a non-parametric Kruskal-Wallis test.

3. Results

3.1. The PP line regeneration

To insert the Psoralen Pathway (PP) in tomato plants, we first leveraged the GoldenBraid technology to assemble the selected four genes encoding the enzymes required for the conversion of *p*-coumaroyl-CoA into psoralen. The T-DNA was build-up with 5 genes organized as transcription units (TUs) (Larbat *et al.,* 2009; Munakata *et al.,* 2016; Roselli *et al.,* 2017; Villard *et al.,* 2021), each independently under the control of the constitutive *CaMV*35S promoter and the tNOS terminator (**Fig. 1**; **Fig. S1**). Then, we integrate the PP T-DNA into the genome of the *Wva*106 tomato cultivar using an *agrobacterium*-mediated stable transfection. Plant selection was allowed by a fifth similarly assembled TU including the kanamycin resistance. By assessing genomic integration via PCR on gDNA, we successfully regenerated two independently transfected lines (**Fig. S1b**). Among the two lines, qPCR revealed that only one line (the PP line) showed a constitutively high expression levels of each transgene in the investigated leaves and roots, compared to the WT (**Table S2g,h; Fig. S1c**). Our study focused on this particular line, which exhibited a phenotype similar to the WT (**Fig. S1e**).

3.2. The PP line accumulates coumarins

To investigate the impact of Psoralen Pathway integration on tomato plant metabolic composition, we compared the metabolomes of PP and WT plants in leaves, stems, and roots using untargeted LC-HRMS analysis of hydro-methanolic extracts. To strengthen the analysis, given that only one PP line was obtained, we performed three independent cultivation rounds under identical growing conditions, resulting in 23 WT and 19 PP plants. The metabolome comparison between PP and WT in all the three-round cultivation of plants and across the three organs, highlighted an overall tendency to a higher accumulation of dozens to hundred metabolic features in the PP line. By considering only significant metabolic features that differentially accumulated (Benjamini-Hochberg p<0.05), 82 metabolic features across leaves, stems and roots were retained (**Table S1**). One metabolic feature was less abundant in the PP line whereas all others were enriched in comparison to the WT. Putative annotation of these features allowed to categorize 20 of them as fatty acids and 19 as phenylpropanoids (*i.e.,* 10 coumarins, 7 phenolamides, the furanocoumarin: psoralen, and one miscellaneous compound) (**Fig. 2a**). The abundance of only eight of the metabolic features were significantly affected in at least two organs, and among them scopoletin, the only metabolite that was found to be differential in leaves, stems and roots (**Fig. 2b**).

Figure 2: Untargeted comparative metabolomic profiling of the Psoralen Pathway (PP) and the Wild-Type (WT) tomato lines. (**a**) Number of metabolic features significantly over-accumulated in the PP line compared to the WT. Metabolic features are categorized according to the NP-Classifier ontology (Kim *et al.,* 2021). Shikimate and phenylpropanoids are highlighted in blue from dark to light: coumarins*, phenolamides*, others, furanocoumarins. Asterisks*

351 correspond to metabolic subclasses significantly enriched (**Table S1b**). (**b**) Venn diagram
352 describing the distribution of the PP line's over-accumulated features across leaves (green),
353 stems (brown) and roots (grey) compared to the control line. Number of replicates for leaves,
354 stems, roots: WT (n=23), PP (n=19). The data are based on 3 independent cultivation rounds of
355 tomato plants grown in the same phytotronic conditions.
356

We then performed targeted analyses focusing on coumarins and metabolites of the

psoralen pathway. Psoralen and its precursors (*i.e.,* demethylsuberosin and marmesin)

were detected only in trace amounts across the analyzed organs of the PP line, and were

absent from the WT (**Fig. 3a**). The PP line exhibited significant increases in *p*-coumaric

acid and umbelliferone levels in the leaves (50- and 31-fold, respectively in comparison

to WT), with no significant changes in other plant parts. Scopoletin concentrations were

significantly increased in all organs of the PP line (23- to 1799-fold), and especially in

stems where it reached concentrations close to 5 μg/g of DW (**Fig. 3b**). Esculetin and its

glycoside form esculin accumulated significantly higher in the PP line than in the WT (**Fig.**

3a). On the opposite, scopolin accumulation was not different between the WT and the

PP line.

369
Figure 3: Comparative targeted profiling of coumarins and furanocoumarins in the Psoralen
Pathway (PP) and the Wild-Type (WT) tomato lines. (**a**) Relative accumulation of reported
coumarins and furanocoumarins in the PP line (blue) versus the WT line (grey), expressed as
mean ± standard deviation. Percentage of replicates containing the metabolite in the PP line is
illustrated using black pie charts. Only significantly differentially accumulated metabolites,
assessed by the Kruskal-Wallis test (p<0.05*), are depicted according to their increasing fold
change in the PP line compared to the WT line. 'Up' refers to compounds detected only in the PP
line. (**b**) Absolute quantification of scopoletin in µg/g of dry weight. Data are expressed as
percentages (a) or absolute quantity for each metabolite relative to the most abundant peak area,
normalized to the internal standard taxifolin, and reported per mg of dry weight; Number of
replicates for leaves, stems, roots: WT (n=23), PP (n=19). The data are based on 3 independent
cultivation rounds of tomato plants grown in the same phytotronic conditions.
382

3.3. Scopoletin is released by a lignin-degradative method

Scopoletin has recently been described to be able to incorporate into the lignin polymer
of *Arabidopsis thaliana* (Brassicaceae) and *Populus sp* (Salicaceae) transgenic lines
producing this metabolite in the stems (Hoengenaert *et al.,* 2022; N. Wang *et al.,* 2025).
Accordingly, the lignin composition was analyzed in the PP line and compared to that in
the WT. Lignin content (**Fig. 4a**) and composition (H, G, S) (**Fig. 4b-c**) in the stems of the
PP line were not significantly affected compared to WT. However, analysis of the

thioacidolysis released monomers indicated a slight but significant increase in the release of scopoletin in the PP line (2.8×) compared to the WT line (**Fig. 4c**), suggesting the presence of scopoletin in the cell wall as well as in the lignin.

Figure 4: Impact of the Psoralen Pathway (PP) line on lignin content (a) and composition for G, S (b), and H monomers, as well as for scopoletin (c), in stems compared to the Wild-Type (WT) line. The average percentage of monomers and scopoletin are calculated based on the content of H, G, and S units and scopoletin in the lignin. The error bars indicate the standard deviation. The lignin content was measured as percentage of Cell Wall Residue (CWR) using the Cysteine-Assisted Sulfuric Acid (CASA) method (Lu *et al.,* 2021). Data were compared using the Kruskal-Wallis's test (p-value<0.05) (p-value for scopoletin % = 0.00106); WT (n=8) and PP (n=9).

3.4. Specific genes regulation in the PP line

To further understand the metabolic changes observed in the PP line, we then investigated its transcriptome compared to the WT. Results highlighted 1014 and 53 differentially expressed genes (DEGs) in the leaves and roots, respectively, of the PP line (**Fig. 5a; Table S2**). In leaves, the 1014 DEGs comprised 659 upregulated and 355 downregulated genes in the PP line compared to WT. In roots, 41 genes were upregulated and 12 downregulated. Seventeen DEGs were common to leaves and roots. Gene Ontology Enrichment Analysis (GOEA) was performed on the up- and downregulated DEGs identified. In leaves, GOEA analysis highlighted an overall upregulation of genes involved in processes related to response to stress and wounding (**Fig. 5b**), whereas the main trend was a downregulation of genes associated to chlorophyll biosynthesis (**Fig. 5c**). In roots, the relatively short list of DEGs did not lead to any significant enrichment using GOEA.

Figure 5: Transcriptomic regulation in the Psoralen Pathway (PP) line compared to the Wild-Type (WT) line. (**a**) Differentially Expressed Genes (DEGs) in leaves and/or roots (FDR<0.05). Gene Ontology Enrichment Analysis (GOEA) on biological processes for (**b**) upregulated and (**c**) downregulated genes considering only leaves (False Discovery Rate: FDR<0.05). The position of the circles represents the fold change by related to the ratio PP/WT. The size of the circles represents the number of Differentially Expressed Genes (DEGs) in each Gene Ontology (GO) category. The color of the circles indicates the statistical significance. The data consist of 3 replicates of PP and WT per tissue.

Using a differential gene analysis based on 3 replicates in leaves and roots comparing the PP and WT lines, we highlighted only strongly DEGs in the PP line (**Fig. 5**). Finer regulatory patterns were uncovered through a K-means clustering approach, which revealed 11 expression clusters based on gene expression profiles in leaves and roots, comparing the PP and WT lines (**Fig. S2**). Among them, five clusters displayed expression patterns that distinguished the PP line from the WT. The smaller and most specific expression cluster was cluster 8 regrouping 309 genes with a higher expression in both the leaves and roots of the PP line in comparison to WT. In addition, cluster 2 (1449 genes) contained genes specifically over-expressed in the roots of the PP line, and cluster 7 (1113 genes), over-expressed genes in the leaves. Conversely, the clusters 1 (743 genes) and 5 (1466 genes) grouped genes downregulated in leaves and roots of the PP line compared to the WT. GOEA of cluster 8 indicates the induced expression of genes potentially linked to the PP

line's response to stress, likely signaled *via* jasmonate and in response to lipids (**Fig. 6a**).

As an extension of cluster 8, the same trend was observed as for the genes from cluster

7 (upregulated in leaves of the PP line) (**Fig. 6b**). The cluster 2 (upregulated genes in roots

of the PP line), were enriched in processes related to hormonal regulations (**Fig. 6c**).

Among the processes related to downregulated genes (**Fig. 6d,e**), DNA replication and

cell cycle appeared affected in the roots of the PP line (cluster 5), whereas lipid and sterol

biosynthesis were, in leaves (cluster 1).

Figure 6: Gene Ontology Enrichment Analysis (GOEA) on biological processes in the Psoralen

Pathway (PP) line compared to Wild-Type (WT) line for clusters 8 (a), 7 (b), 2 (c), 5 (d) and 1 (e).

(*False Discovery Rate:* FDR<0.05). The position of the circles represents the fold change by

related to the ratio PP/WT. The size of the circles represents the number of Differentially

Expressed Genes (DEGs) in each GO category. The color of the circles indicates the statistical

significance. Gene clusters were grouped using the k-means method. (**a**) Cluster 8 includes 309

upregulated genes and most of those were specifically differentially expressed in the leaves and

roots of the PP line compared to the WT line (10/17 genes). (**b**) Cluster 7 contains 1113

upregulated genes in leaves. (**c**) Cluster 2 contains 1449 upregulated genes in roots. (**d**) Cluster

1 and (**e**) cluster 5 contain 743 and 1465 downregulated genes in roots and leaves, respectively.
The data consist of 3 replicates of PP and WT per tissue.

3.5. The PP line showed reduced biomass and elevated nitrogen content
To identify potential impacts on the growth and development of the PP line compared to

the WT line, we performed a comparative resource allocation and morphology analysis

(**Table S3**). After 28 days of growth, PP and WT plants exhibited comparable morphology

and size (**Fig. S1e**), with similar leaf numbers and stem lengths between days 13 and 28,

indicating no significant developmental changes (**Table 1**). However, dry biomass in the

PP line was significantly reduced by 21.3% compared to the WT. This decrease was

uniformly distributed across plant organs with the exception of stem which had a similar

amount of dry biomass compared to the WT. As a results, the resource allocation pattern

in the PP line was significantly different from the WT, with a decrease in the proportion of

matter in the leaves (-10%) and an increase in the stem (+20%) (**Table 1**). The decrease in

the biomass accumulation in leaves was in accordance with an increase in the Specific

Leaf Area (SLA) in the PP line.

	Variable	WT average ± SE	PP average ± SE	p-value
	Foliar blades (g)	2.46±0.36	1.74±0.24	**0.0004***
	Petioles (g)	0.49±0.07	0.4±0.05	**0.0222***
Dry weights	Stems (g)	1.11±0.28	1.05±0.17	0.4142
	Roots (g)	0.49±0.05	0.39±0.06	**0.0025***
	Stems/Roots ratio	8.20±1.07	8.15±0.70	0.9349
	Foliar blades	54.06	48.6	**0.0042***
Tissue allocation (% by DW)	Petioles	10.76	11.17	0.2529
	Stems	24.39	29.32	**0.0178***
	Roots	10.76	10.89	0.9349
	Stems (cm)	49.95±7.42	51.33±6.02	0.6532
Development	Roots (cm)	44.21±15.1	44.22±9.38	0.7439
	Leaf number	8	8	1
	D13/D28 plant height	0.1±0.02	0.1±0.02	0.8064

Key growth analysis parameters	Stems/Roots ratio	1.24±0.43	1.2±0.28	1
	SLA (cm2.g⁻¹)	236.48±33.67	329.73±35.81	**0.0002***
	SRL (cm.g⁻¹)	90.95±39.82	115.21±31.85	0.0724
	SSL (cm.g⁻¹)	45.06±14.62	47.84±6.17	**0.0274***

Table 1: Mean alterations in growth and development parameters between WT and the PP line. The significant differences between the two genotypes are supported by a non-parametric Kruskal-Wallis test. Significant differences are indicated in bold with an asterisk. Measurements were taken at a vegetative growth stage (28 days after sowing; D28) and compared for development with measurements at an early growth stage (13 days after sowing; D13). Dry biomass allocation patterns are defined by the percentage of mean dry biomass in each compartment. Abbreviations: SLA (Specific Leaf Area); SRL (Specific Root Length); SSL (Specific Shoot Length). WT (n=23), PP (n=19).

To better understand the cause of the loss in biomass in the PP line, we analyzed carbon (%C) and nitrogen (%N) accumulation across the different tissues (**Table 2**). Results showed that nitrogen content was significantly higher in the PP line compared to the WT, particularly in the shoots (+53%). Focusing on ion uptakes in leaves and roots (**Table S4**), we identified a significant increase of nitrate ions in roots (×3.3). Conversely, carbon content was slightly lower in the PP line, most notably in the stems (-13%). As a result, the carbon-to-nitrogen ratio (C/N) was significantly decreased in the PP line, especially in the shoots.

	Carbone (%)			Nitrogen (%)			C/N ratio		
	WT mean (mean ± STDEV)	PP mean	p-value	WT mean (mean ± STDEV)	PP mean	p-value	WT mean (mean ± STDEV)	PP mean	p-value
Foliar blades	39.01±0.41	38.66±1.02	0.1616	1.72±0.3	2.63±0.33	**0.0005***	22.47±3.96	14.82±1.59	**0.0002***
Petioles	34.57±0.62	33.89±0.66	**0.0484***	1.34±0.22	1.96±0.3	**0.0005***	25.46±3.91	17.57±2.49	**0.0002***
Stems	31.95±2.9	27.78±5.75	0.0542	1.01±0.18	1.22±0.21	**0.0301***	31.96±4.74	22.85±3.68	**0.0004***
Roots	40.79±0.7	40.52±0.63	0.885	2.02±0.12	2.27±0.17	**0.006***	20.33±0.95	17.89±1.29	**0.0004***

Table 2: Mean alterations in Carbon (C), Nitrogen (N) contents and C/N ratio between the Wild-Type (WT) and the Psoralen Pathway (PP) lines. The variables significance between the two genotypes is supported by a non-parametric Kruskal-Wallis test. Significant differences are indicated in bold with an asterisk. Measurements were taken at a vegetative growth stage (28 days after sowing). WT (n=10), PP (n=9).

4. Discussion

The biosynthesis of specialized metabolites necessarily triggered a physiological cost on plants, either by diverting resources for their production or through the pleiotropic effects of the specialized metabolites themselves (Neilson *et al.,* 2013). This study aimed to better understand the impacts of constitutive expression of four genes from the linear furanocoumarin biosynthesis pathway on tomato plant physiology and growth.

4.1. Furanocoumarins might be toxic for tomato plants

Genetic modification of tomato with our construct aimed at producing psoralen surprisingly resulted in a low transformation success rate (0.33%), especially when compared to longer pathways already inserted with success in tomato using the same GoldenBraid system with TUs also under the *CaMV*35S promoter (H. Kim *et al.,* 2025; Sonawane *et al.,* 2017). Furthermore, psoralen was found only as trace amounts despite that all PP transgenes were constitutively expressed (**Fig. S1c**; **Table S2g**). In parallel, transcriptomic data showed significantly over-expressed genes falling in the GO categories "response to stress", "response to chemicals" and other direct or indirect stress-related GO classifications in the PP line (*e.g.,* photoinhibition, stress phytohormones, etc) (**Fig. 5b,c**; **Fig. 6a-c**). These findings suggest a psoralen pathway-dependent phytotoxicity in tomato plants, potentially linked to the production of furanocoumarins (**Fig. 3a**). During the seven last decades, incremental progress has highlighted a large spectrum of furanocoumarin's toxicity on a vast range of organisms, including plants, occurring through a diversity of processes (*e.g.,* photocycloaddition, lipid peroxidation, cytochrome P450 inhibition) (Baskin *et al.,* 1967; Bennet & Bonner, 1953; Diawara *et al.,* 1993; Frank *et al.,* 1998; Greenberg *et al.,* 2001; Hale *et al.,* 2004; Hassan *et al.,* 2016; Kitamura *et al.,* 2005; Meepagala *et al.,* 2021; Razavi, 2010; X.-B. Wang *et al.,* 2008). We therefore hypothesize that psoralen, or core intermediates and/or derivatives of its biosynthetic pathway, may be toxic, or even lethal, to tomato above a certain threshold. This toxicity may have impaired the regeneration of additional transgenic tomato lines. However, although our transcriptomic data clearly indicated a chemically related phytotoxicity in the PP line, no clear signal allowed us to pinpoint if this toxicity was related to the production of psoralen and/or any intermediates, or even to the accumulation of any of the other accumulated compounds, in particular

coumarins. However, the fact that the PP line plants appeared healthy (**Fig. S1e**) suggests that they were able to circumvent the induced phytotoxicity.

4.2. The low accumulation of furanocoumarins may reflect metabolic competition favouring the coumarin biosynthesis pathway

Metabolic engineering approaches often fall short of expectations due to our limited understanding of the self-management of specialized metabolite biosynthetic pathways in plants (Lynch *et al.,* 2021). In the PP line, psoralen as well as its precursors from umbelliferone could be detected in almost all the plant parts but at a level below the quantification limit (**Fig. 3a**), proving that our metabolic engineering approach is working but not sufficiently well to allow psoralen accumulation. Given that all genes encoding the enzymes essential for psoralen accumulation were appropriately expressed and the corresponding enzymes proven functional (**Fig. S1e**; **Table S2g**), we suggest a metabolic competition for its precursor, *p*-coumaroyl-CoA, which may be steered instead toward coumarins, which were the more enriched in the PP line. Indeed, simple coumarins like umbelliferone, esculetin, and scopoletin derive from *p*-coumaroyl-, caffeoyl-, and feruloyl-CoA, respectively, via 2OGD-mediated *ortho*-hydroxylation followed by a cyclisation that occurs spontaneously and/or is catalyzed by a COSY (C. Y. Kim *et al.,* 2023; Vanholme *et al.,* 2019) (**Fig. 1**). The *ortho*-hydroxylation of *p*-coumaroyl-CoA constitutes the first intended metabolic step for psoralen production in the PP line. To allow umbelliferone accumulation for psoralen production, we chosen the C2'H 2OGD from *Pastinaca sativa* (Apiaceae) (*Ps*Diox) which was characterized *in vitro* as being specific for *p*-coumaroyl-CoA (Roselli *et al.,* 2017). The marked and significant accumulation of coumarins in the PP line including the formally identified esculin and scopoletin (**Fig. 3**, **Fig. S3**), suggests that, *in vivo*, *Ps*Diox could share the same catalytic properties as its orthologs from *Ruta graveolens* (Rutaceae), *A. thaliana* and *Ipomea batatas* (Convolvulaceae) that also accept caffeoyl-CoA and feruloyl-CoA as substrates (Matsumoto *et al.,* 2012; Vialart *et al.,* 2012; Yang *et al.,* 2015). This hypothesis is supported by the accumulation of scopoletin and esculetin in plants overexpressing F6'H (Beesley *et al.,* 2023; N. Wang *et al.,* 2025).

Non-exclusive hypotheses for the limited accumulation of psoralen may also involve active regulation of competing metabolic pathways that bypass the psoralen branch (**Fig. 3**). Indeed, comparative metabolomics analysis highlighted the accumulation of phenolamides (*i.e.,* fused hydroxycinnamates and amines), a natural defence compound family in tomato plants (Roumani et al., 2022), in addition to coumarins (**Fig. 2**, **Table S1**). In parallel, comparative transcriptomics pointed to the overexpression in the PP line of genes of the core phenylpropanoid pathway putatively involved in the coumarin and phenolamide biosynthesis pathways (*e.g.*, BAHD acyltransferases, Tyramine N-feruloyl transferase) (**Table S2c**). Such metabolic competition for a common substrate such as *p*-coumaroyl-CoA could also be further established by other factors like metabolic channeling as now established in several species (Dahmani *et al.,* 2023; Lallemand *et al.,* 2013; Laursen *et al.,* 2016). Heterologous enzymes from different plant species may be less competitive than endogenous ones in establishing metabolic channeling of intermediates and co-substrates, ultimately at the expense of psoralen accumulation.

4.3. Coumarin accumulation might underly the PP line's altered physiology and growth

The metabolomics analyses highlighted 82 metabolic features almost exclusively over-accumulated in the transgenic line. Annotation outlined that primarily 4 metabolic families (fatty acids, phenolamides, coumarins and furanocoumarins) were involved. Among them, coumarin accumulation stood out from the other families for at least one of the following reasons: (i) the accumulation of all individual coumarins followed a presence/absence pattern in the PP/WT comparison (**Fig. 3a**; **Table S1**), (ii) the accumulation of coumarin metabolites was detected in the three tissues (leaf, stem, roots) in accordance to the constitutive expression of the PP transgenes (**Fig. 3a**) and (iii) based on the targeted analyses the accumulation of coumarins reached several µg/g of DW (*i.e.,* scopoletin) whereas the furanocoumarin accumulation did not reach the quantification limit (**Fig. 3**). These observations made us consider that coumarins accumulation may be the main stimulus of the transcriptomic, physiological and developmental characteristics of the PP line. This would be consistent with previous studies in which coumarins (*i.e.,* umbelliferone, scopoletin, coumarin, and daphnoretin)

were applied to *A. thaliana* and *Lactuca sativa* (Asteraceae), resulting in allelopathic effects such as membrane peroxidation, Reactive Oxygen Species (ROS) production, mitotic disruption, and photoinhibition (Araniti *et al.,* 2017; Graña *et al.,* 2017; Z. Yan *et al.,* 2016). Our transcriptomics analysis highlighted the upregulation, in the PP line in comparison to WT, of a majority of genes associated with responses to stress, chemicals, organic substances and wounding (**Fig. 5b**; **Fig. 6a-c**). These categories comprise genes involved in ROS detoxification (*e.g.,* glutathione-S-transferases, cytochrome P450s, peroxidases, detoxification proteins, chaperones, and heat-shock proteins) (**Table S2**). On the contrary, the main categories of downregulated genes in the leaves of the PP line compared to WT were associated to chlorophyll biosynthesis (**Fig. 5c**; **Fig. 6d**). These transcriptomics patterns suggest an active regulation of the plant oxidative status, potentially mediated by an increase in the ROS detoxication processes and a decrease in the ROS production linked to photosynthesis. This suggestion is supported by the healthy appearance of the PP line despite the activation of all these stress signaling pathways (**Fig. S1e**). As now well established, photoinhibition generally acts as a protective mechanism to limit photoinduced ROS accumulation, as reviewed by Didaran *et al.,* 2024. In parallel, the lipid over-accumulation observed in the leaves of the PP line (**Table 1**), together with the over-expression of genes associated with lipid metabolism (*e.g.,* phospholipases, aldehyde dehydrogenases, lipid biosynthesis) and membrane repair (*e.g.,* lipid desaturases, acyltransferases) (**Fig. 6a**; **Table S2**), further supports the occurrence of membrane damage. This lipid stress may result from coumarin-induced destabilization of chloroplast integrity and photochemistry, as previously observed in *A. thaliana* (Araniti *et al.,* 2017).

To further understand the impact of coumarins on tomato physiology in the PP line, we investigated its effects on growth, development and resource allocation compared to the WT line. First, we observed no developmental alterations in the PP line, in contrast to the pronounced growth impairments reported in various plant species metabolically engineered to over-accumulate scopoletin (Beesley *et al.,* 2023; Hoengenaert *et al.,* 2022; N. Wang *et al.,* 2025). However, F6'H knock-down lines of *Manihot esculenta* (Euphorbiaceae) with reduced scopoletin accumulation have revealed key roles for this compound in cassava development (*e.g.*, auxiliary budding, leaf appearance, apical

dominance) (Mukami *et al.,* 2024). All these elements suggests that the physiological effects of coumarins on development may depend on their constitutive accumulation levels and may be species-specific. This also underscores how poorly understood the physiological regulation of coumarins remains, despite their strong potential to act in a phytohormone-like manner, as hypothesized for an auxin-like effect of scopoletin by Graña *et al.,* 2017.

Morphological analysis of the PP line revealed a significant biomass reduction, particularly of the leaves (**Table 1**). A similar biomass loss was previously reported in *A. thaliana* supplemented with coumarin or scopoletin (Araniti *et al.,* 2017; Graña *et al.,* 2017) and was partly attributed to photoinhibition observed in the leaves, similarly to what we found in the PP line according to transcriptomic data (**Fig. 6d**). In addition, we also observed a downregulation of a few genes putatively involved in sugar storage (*i.e.,* starch synthase) and numerous genes putatively involved in sugar metabolism and cell wall remobilization (**Table S2c,d**). Their differential expression may help explain the biomass loss observed, which cannot be exclusively attributed to carbon content reduction, as this decrease is substantially lower than the 21% drop in biomass (**Tables 2 and 3**). Considering that most of the biomass components were measured (*e.g.,* carbon, nitrogen, ions) (**Table 2; Table S4**), we suggest a concomitant reduction in oxygen-rich compounds such as sugars. In parallel, significant increase of specific leave surface (SLA; +39%) underlies a biomass loss that is distributed within thinner leaves (**Table 1**). Taken together, these elements suggest that coumarins may trigger a 'slimming effect', leading to biomass reduction, primarily in tomato leaves, as a potential response to coumarin-induced phytotoxicity.

Thioacidolysis revealed that scopoletin was incorporated into the cell wall (**Fig. 4c**), albeit at a level approximately 10 to 100 times lower than found in *A. thaliana* and *Glycine max* (Fabaceae) (Beesley *et al.,* 2023; Hoengenaert *et al.,* 2022). Although only trace amounts of scopoletin accumulated in the cell wall, which prevented validation of its integration into the lignin polymer by NMR, it is reasonable to consider that a small fraction of scopoletin could be incorporated into the lignin polymer. Although the stem cell wall appears to be a key storage compartment for coumarins (Beesley *et al.,* 2023), glycosylated coumarins are stored in vacuoles, *e.g.,* in *A. thaliana* roots (Robe *et al.,*

2021). Both vacuolar storage and cell wall transfer are well established toxicity mitigation

processes (Weng *et al.,* 2021). Our metabolic and phenotypic data suggest that even a

modest accumulation of scopoletin (~1-5 µg/g DW), has to be mitigated in the PP line.

Finally, we found that the significant increase in nitrogen content in the upper parts of the

PP line coincided with enhanced nitrate uptake by the roots. Catecholic coumarins such

as esculetin and fraxetin are well-known to facilitate iron uptake from the soil in plants

(H. H. Tsai & Schmidt, 2017). Our data suggest that coumarins accumulation, primarily

esculetin, is concomitant with nitrate uptake, highlighting a potential role of coumarins

in this process. A similar contribution of coumarins already reported in *Triticum durum*

and *Zea mays* (both Poaceae) (Abenavoli *et al.,* 2001; Lupini *et al.,* 2018).

5. Conclusion

The production of psoralen in tomato proved to be more challenging than the mere

expression of the biosynthetic genes as it potentially led to pronounced phytotoxicity. Our

results suggests that the accumulation of coumarins at concentrations of 1-5 µg/g of DW

in tomato plants could induces several levels of regulation: (i) xenobiotic stress leading

to oxidative stress, which the plant can adapt; (ii) a photoinhibition and an increase in

metabolism inducing a biomass slimming possibly in response to oxidative stress; and

(iii) a modification of the nitrogen and nitrate levels. Our results indicate that, although

coumarins hold substantial agronomic potential (non-toxic to pollinators, facilitating iron

uptake, shaping root microbiota, etc) (Cosme et al., 2021; H.-H. Tsai et al., 2018; Zhou et

al., 2023), their role in plant defense entails a growth cost even at low content. A

counterpart that should be carefully considered and ideally mitigated, following a more

precise delineation of the regulatory pathways of coumarins underlying the identified

physiological impacts, before pursuing potential viable applications.

6. Acknowledgments

We would like to thank Alexandre Olry, Claude Gallois and David Marcolet members of

the Experimental phytotronic platform of Lorraine PEPLor (Université de Lorraine, France)

for their technical assistance, Cédric Paris from the Metabolomic and Structural Analytic

Platform facility (PASM, Université de Lorraine, France) as well as Geert Goeminne and

Sandrien Desmet from the Metabolomic Core (VIB, Gent, Belgium). We thank Sarah Liu

for her assistance on NMR sample preparation. We thank Florent Magot (BBV, Université

de Tours) and Florent Ducrocq for assistance in the critical analysis of the metabolomic

data as well as Eugène Maurey for the critical statistical analysis of morphophysiological

data.

7. Competing interests

None declared

8. Author contributions

GG, CV and AB, designed and assembled the PP genetic construction. AB, MR, CV, AF

and GG participated to transgenic lines regeneration. AB, AF and MR cultivated and

harvested the tomato plants. AB, FH and RL generated and analyzed the transcriptomic

data. AB, JG and RL generated and analyzed the metabolomic data. AB, LH, JR and WB

analyzed the lignin composition. AB, CR and RL generated and analyzed the

morphophysiological data. AB, RL and AH conceptualized the project and wrote the

article. WB, AH and RL supervised the project. All co-authors reviewed the article.

9. Funding sources

AB was financially supported by a PhD grant provided by the French Government and

Grand Est Region. AB benefited from the mobility programs DREAM provided by 'Lorraine

Université d'Excellence', funded by the ANR 'Investissements d'avenir' (grant no. 15-004)

and from the DESSE facility funded by the INRAE. WB was funded by the interuniversity

Bijzonder Onderzoeksfonds (iBOF) NextBioRef. JR was supported by the Great Lakes

Bioenergy Research Center, U.S. Department of Energy, Office of Science, Biological and

Environmental Research Program, under Award Number DE-SC0018409.

10. Bibliography

Abenavoli, M. R., De Santis, C., Sidari, M., Sorgonà, A., Badiani, M., & Cacco, G.
(2001). Influence of coumarin on the net nitrate uptake in durum wheat. New Phytologist,
150(3), 619-627. https://doi.org/10.1046/j.1469-8137.2001.00119.x
Alves-Carvalho, S., Gazengel, K., Bretaudeau, A., Robin, S., Daval, S., & Legeai, F.
(2021). AskoR, A R Package for Easy RNASeq Data Analysis. Proceedings of The 1st

International Electronic Conference on Entomology, 10646. https://doi.org/10.3390/IECE-10646

Araniti, F., Scognamiglio, M., Chambery, A., Russo, R., Esposito, A., D'Abrosca, B., Fiorentino, A., Lupini, A., Sunseri, F., & Abenavoli, M. R. (2017). Highlighting the effects of coumarin on adult plants of *Arabidopsis thaliana* (L.) Heynh. By an integrated -omic approach. Journal of Plant Physiology, 213, 30-41. https://doi.org/10.1016/j.jplph.2017.02.013

Baskin, J. M., Ludlow, C. J., Harris, T. M., & Wolf, F. T. (1967). Psoralen, an inhibitor in the seeds of *Psoralea subacaulis* (Leguminosae). Phytochemistry, 6(9), 1209-1213. https://doi.org/10.1016/S0031-9422(00)86083-5

Beesley, A., Beyer, S. F., Wanders, V., Levecque, S., Bredenbruch, S., Habash, S. S., Schleker, A. S. S., Gätgens, J., Oldiges, M., Schultheiss, H., Conrath, U., & Langenbach, C. J. G. (2023). Engineered coumarin accumulation reduces mycotoxin-induced oxidative stress and disease susceptibility. Plant Biotechnology Journal, 21(12), 2490-2506. https://doi.org/10.1111/pbi.14144

Bennet, & Bonner. (1953). Isolation of Plant Growth Inhibitors from *Thamnosma montana*. 6. American Journal of Botany, 29-33. https://doi.org/10.2307/2438489

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic : A flexible trimmer for Illumina sequence data. Bioinformatics, 30(15), 2114-2120. https://doi.org/10.1093/bioinformatics/btu170

Bouillé, A., Larbat, R., Kumari, R., Olry, A., Charles, C., Nelson, D. R., Thornton, J., Villard, C., & Hehn, A. (2025). Lineage-specific patterns in the Moraceae family allow identification of convergent P450 enzymes involved in furanocoumarin biosynthesis. New Phytologist. https://doi.org/10.1111/nph.20381

Chassagne, F., Cabanac, G., Hubert, G., David, B., & Marti, G. (2019). The landscape of natural product diversity and their pharmacological relevance from a focus on the Dictionary of Natural Products®. Phytochemistry Reviews, 18(3), 601-622. https://doi.org/10.1007/s11101-019-09606-2

Cosme, M., Fernández, I., Declerck, S., van der Heijden, M. G. A., & Pieterse, C. M. J. (2021). A coumarin exudation pathway mitigates arbuscular mycorrhizal incompatibility in *Arabidopsis thaliana*. Plant Molecular Biology, 106(4-5), 319-334. https://doi.org/10.1007/s11103-021-01143-x

Dahmani, I., Qin, K., Zhang, Y., & Fernie, A. R. (2023). The formation and function of plant metabolons. The Plant Journal, 114(5), 1080-1092. https://doi.org/10.1111/tpj.16179

Diawara, M. M., Trumble, J. T., White, K. K., Carson, W. G., & Martinez, L. A. (1993). Toxicity of linear furanocoumarins to *Spodoptera exigua*: Evidence for antagonistic interactions. Journal of Chemical Ecology, 19(11), 2473-2484. https://doi.org/10.1007/BF00980684

Didaran, F., Kordrostami, M., Ghasemi-Soloklui, A. A., Pashkovskiy, P., Kreslavski, V., Kuznetsov, V., & Allakhverdiev, S. I. (2024). The mechanisms of photoinhibition and repair in plants under high light conditions and interplay with abiotic stressors. Journal of Photochemistry and Photobiology B: Biology, 259, 113004. https://doi.org/10.1016/j.jphotobiol.2024.113004

Djoumbou Feunang, Y., Eisner, R., Knox, C., Chepelev, L., Hastings, J., Owen, G., Fahy, E., Steinbeck, C., Subramanian, S., Bolton, E., Greiner, R., & Wishart, D. S. (2016). ClassyFire : Automated chemical classification with a comprehensive, computable

taxonomy. Journal of Cheminformatics, 8(1), 61. https://doi.org/10.1186/s13321-016-0174-y

Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A. A., Melnik, A. V., Meusel, M., Dorrestein, P. C., Rousu, J., & Böcker, S. (2019). SIRIUS 4 : A rapid tool for turning tandem mass spectra into metabolite structure information. Nature Methods, 16(4), 299-302. https://doi.org/10.1038/s41592-019-0344-8

Frank, S., Caffieri, S., Raffaelli, A., Vedaldi, D., & Dall'Acqua, F. (1998). Characterization of psoralen-oleic acid cycloadducts and their possible involvement in membrane photodamage. Journal of Photochemistry and Photobiology B: Biology, 44(1), 39-44. https://doi.org/10.1016/S1011-1344(98)00103-1

Ge, S. X., Jung, D., & Yao, R. (2020). ShinyGO : A graphical gene-set enrichment tool for animals and plants. Bioinformatics, 36(8), 2628-2629. https://doi.org/10.1093/bioinformatics/btz931

Godichon-Baggioni, A., Maugis-Rabusseau, C., & Rau, A. (2019). Clustering transformed compositional data using K -means, with applications in gene expression and bicycle sharing system data. Journal of Applied Statistics, 46(1), 47-65. https://doi.org/10.1080/02664763.2018.1454894

Graña, E., Costas-Gil, A., Longueira, S., Celeiro, M., Teijeira, M., Reigosa, M. J., & Sánchez-Moreiras, A. M. (2017a). Auxin-like effects of the natural coumarin scopoletin on A*rabidopsis* cell structure and morphology. Journal of Plant Physiology, 218, 45-55. https://doi.org/10.1016/j.jplph.2017.07.007

Graña, E., Costas-Gil, A., Longueira, S., Celeiro, M., Teijeira, M., Reigosa, M. J., & Sánchez-Moreiras, A. M. (2017b). Auxin-like effects of the natural coumarin scopoletin on Arabidopsis cell structure and morphology. Journal of Plant Physiology, 218, 45-55. https://doi.org/10.1016/j.jplph.2017.07.007

Greenberg, R. B., Alberti, M., Hearst, J. E., Chua, M. A., & Saffran, W. A. (2001). Recombinational and Mutagenic Repair of Psoralen Interstrand Cross-links in *Saccharomyces cerevisiae*. Journal of Biological Chemistry, 276(34), 31551-31560. https://doi.org/10.1074/jbc.M103588200

Hale, A. L., Meepagala, K. M., Oliva, A., Aliotta, G., & Duke, S. O. (2004). Phytotoxins from the Leaves of *Ruta graveolens*. Journal of Agricultural and Food Chemistry, 52(11), 3345-3349. https://doi.org/10.1021/jf0497298

Han, L., Zhang, L., He, Y., Liao, L., Li, J., Xu, S., Zhao, Y., Bian, X., & Xia, Y. (2023). Three carbon-/oxygen-prenyltransferases responsible for furanocoumarin synthesis in *Angelica dahurica*. Industrial Crops and Products, 200, 116814. https://doi.org/10.1016/j.indcrop.2023.116814

Hassan, Mohd. Z., Osman, H., Ali, M. A., & Ahsan, M. J. (2016). Therapeutic potential of coumarins as antiviral agents. European Journal of Medicinal Chemistry, 123, 236-255. https://doi.org/10.1016/j.ejmech.2016.07.056

He, Z., Webster, S., & He, S. Y. (2022). Growth–defense trade-offs in plants. Current Biology, 32(12), R634-R639. https://doi.org/10.1016/j.cub.2022.04.070

Hehmann, M., Lukačin, R., Ekiert, H., & Matern, U. (2004). Furanocoumarin biosynthesis in *Ammi majus L.* : Cloning of bergaptol O-methyltransferase. European Journal of Biochemistry, 271(5), 932-940. https://doi.org/10.1111/j.1432-1033.2004.03995.x

Hellens, R. P., Edwards, E. A., Leyland, N. R., Bean, S., & Mullineaux, P. M. (2000). pGreen : A versatile and flexible binary Ti vector for *Agrobacterium*-mediated plant

transformation. Plant Molecular Biology, 42(6), 819-832.
https://doi.org/10.1023/A:1006496308160

Hoengenaert, L., Wouters, M., Kim, H., De Meester, B., Morreel, K., Vandersyppe,
S., Pollier, J., Desmet, S., Goeminne, G., Ralph, J., Boerjan, W., & Vanholme, R. (2022).
Overexpression of the scopoletin biosynthetic pathway enhances lignocellulosic
biomass processing. Science Advances, 8(28), eabo5738.
https://doi.org/10.1126/sciadv.abo5738

Hoffmann, M. A., Nothias, L.-F., Ludwig, M., Fleischauer, M., Gentry, E. C.,
Witting, M., Dorrestein, P. C., Dührkop, K., & Böcker, S. (2021). Assigning confidence to
structural annotations from mass spectra with COSMIC. Bioinformatics.
https://doi.org/10.1101/2021.03.18.435634

Ji, J., Han, X., Zang, L., Li, Y., Lin, L., Hu, D., Sun, S., Ren, Y., Maker, G., Lu, Z., &
Wang, L. (2024). Integrative multi-omics data elucidating the biosynthesis and regulatory
mechanisms of furanocoumarins in *Angelica dahurica*.
https://doi.org/10.1101/2024.07.23.604792

Jian, X., Zhao, Y., Wang, Z., Li, S., Li, L., Luo, J., & Kong, L. (2020). Two CYP71AJ
enzymes function as psoralen synthase and angelicin synthase in the biosynthesis of
furanocoumarins in *Peucedanum praeruptorum Dunn*. Plant Molecular Biology, 104(3),
327-337. https://doi.org/10.1007/s11103-020-01045-4

Karamat, F., Olry, A., Munakata, R., Koeduka, T., Sugiyama, A., Paris, C., Hehn, A.,
Bourgaud, F., & Yazaki, K. (2014). A coumarin-specific prenyltransferase catalyzes the
crucial biosynthetic reaction for furanocoumarin formation in parsley. The Plant Journal,
77(4), 627-638. https://doi.org/10.1111/tpj.12409

Kim, C. Y., Mitchell, A. J., Kastner, D. W., Albright, C. E., Gutierrez, M. A.,
Glinkerman, C. M., Kulik, H. J., & Weng, J.-K. (2023). Emergence of a proton exchange-
based isomerization and lactonization mechanism in the plant coumarin synthase
COSY. Nature Communications, 14(1), 597. https://doi.org/10.1038/s41467-023-36299-
1

Kim, H., Liu, L., Han, L., Park, K., Kim, H. J., Nguyen, T., Nazarenus, T. J., Cahoon,
R. E., Haslam, R. P., Ciftci, O., Napier, J. A., & Cahoon, E. B. (2025). Oilseed-based
metabolic engineering of astaxanthin and related ketocarotenoids using a plant-derived
pathway : Lab-to-field-to-application. Plant Biotechnology Journal, pbi.70148.
https://doi.org/10.1111/pbi.70148

Kim, H. W., Wang, M., Leber, C. A., Nothias, L.-F., Reher, R., Kang, K. B., Van Der
Hooft, J. J. J., Dorrestein, P. C., Gerwick, W. H., & Cottrell, G. W. (2021). NPClassifier : A
Deep Neural Network-Based Structural Classification Tool for Natural Products. Journal
of Natural Products, 84(11), 2795-2807. https://doi.org/10.1021/acs.jnatprod.1c00399

Kitamura, N., Kohtani, S., & Nakagaki, R. (2005). Molecular aspects of
furocoumarin reactions : Photophysics, photochemistry, photobiology, and structural
analysis. Journal of Photochemistry and Photobiology C: Photochemistry Reviews,
6(2-3), 168-185. https://doi.org/10.1016/j.jphotochemrev.2005.08.002

Krieger, C., Roselli, S., Kellner-Thielmann, S., Galati, G., Schneider, B., Grosjean,
J., Olry, A., Ritchie, D., Matern, U., Bourgaud, F., & Hehn, A. (2018). The CYP71AZ P450
Subfamily : A Driving Factor for the Diversification of Coumarin Biosynthesis in
Apiaceous Plants. Frontiers in Plant Science, 9, 820.
https://doi.org/10.3389/fpls.2018.00820

Lallemand, B., Erhardt, M., Heitz, T., & Legrand, M. (2013). Sporopollenin
Biosynthetic Enzymes Interact and Constitute a Metabolon Localized to the
Endoplasmic Reticulum of Tapetum Cells. Plant Physiology, 162(2), 616-625.
https://doi.org/10.1104/pp.112.213124

Larbat, R., Hehn, A., Hans, J., Schneider, S., Jugdé, H., Schneider, B., Matern, U.,
& Bourgaud, F. (2009). Isolation and Functional Characterization of CYP71AJ4 Encoding
for the First P450 Monooxygenase of Angular Furanocoumarin Biosynthesis. Journal of
Biological Chemistry, 284(8), 4776-4785. https://doi.org/10.1074/jbc.M807351200

Larbat, R., Kellner, S., Specker, S., Hehn, A., Gontier, E., Hans, J., Bourgaud, F., &
Matern, U. (2007). Molecular Cloning and Functional Characterization of Psoralen
Synthase, the First Committed Monooxygenase of Furanocoumarin Biosynthesis.
Journal   of   Biological   Chemistry,   282(1),   542-554.
https://doi.org/10.1074/jbc.M604762200

Laursen, T., Borch, J., Knudsen, C., Bavishi, K., Torta, F., Martens, H. J., Silvestro,
D., Hatzakis, N. S., Wenk, M. R., Dafforn, T. R., Olsen, C. E., Motawia, M. S., Hamberger,
B., Møller, B. L., & Bassard, J.-E. (2016). Characterization of a dynamic metabolon
producing the defense compound dhurrin in sorghum. Science, 354(6314), 890-893.
https://doi.org/10.1126/science.aag2347

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–
Wheeler   transform.   Bioinformatics,   25(14),   1754-1760.
https://doi.org/10.1093/bioinformatics/btp324

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts : An efficient general
purpose program for assigning sequence reads to genomic features. Bioinformatics,
30(7), 923-930. https://doi.org/10.1093/bioinformatics/btt656

Lu, F., Wang, C., Chen, M., Yue, F., & Ralph, J. (2021). A facile spectroscopic
method for measuring lignin content in lignocellulosic biomass. Green Chemistry,
23(14), 5106-5112. https://doi.org/10.1039/D1GC01507A

Ludwig, M., Nothias, L.-F., Dührkop, K., Koester, I., Fleischauer, M., Hoffmann, M.
A., Petras, D., Vargas, F., Morsy, M., Aluwihare, L., Dorrestein, P. C., & Böcker, S. (2019).
ZODIAC : Database-independent molecular formula annotation using Gibbs sampling
reveals unknown small molecules. Bioinformatics. https://doi.org/10.1101/842740

Lupini, A., Araniti, F., Mauceri, A., Princi, M. P., Sorgonà, A., Sunseri, F., Varanini,
Z., & Abenavoli, M. R. (2018). Coumarin enhances nitrate uptake in maize roots through
modulation of plasma membrane H+ - ATP ase activity. Plant Biology, 20(2), 390-398.
https://doi.org/10.1111/plb.12674

Lynch, J. H., Huang, X.-Q., & Dudareva, N. (2021). Silent constraints : The hidden
challenges faced in plant metabolic engineering. Current Opinion in Biotechnology, 69,
112-117. https://doi.org/10.1016/j.copbio.2020.12.014

Maeda, H. A., & Fernie, A. R. (2021). Evolutionary History of Plant Metabolism.
Annual Review of Plant Biology, 72(1), 185-216. https://doi.org/10.1146/annurev-
arplant-080620-031054

Matsumoto, S., Mizutani, M., Sakata, K., & Shimizu, B.-I. (2012). Molecular cloning
and functional analysis of the ortho-hydroxylases of p-coumaroyl coenzyme A/feruloyl
coenzyme A involved in formation of umbelliferone and scopoletin in sweet potato,
Ipomoea   batatas   (L.)   Lam.   Phytochemistry,   74,   49-57.
https://doi.org/10.1016/j.phytochem.2011.11.009

Meepagala, K. M., Bracken, A. K., Fronczek, F. R., Johnson, R. D., Wedge, D. E., &
Duke, S. O. (2021). Furanocoumarin with Phytotoxic Activity from the Leaves of *Amyris*
elemifera   (Rutaceae).   ACS   Omega,   6(1),   401-407.
https://doi.org/10.1021/acsomega.0c04778

Morris, J. L., Puttick, M. N., Clark, J. W., Edwards, D., Kenrick, P., Pressel, S.,
Wellman, C. H., Yang, Z., Schneider, H., & Donoghue, P. C. J. (2018). The timescale of
early land plant evolution. Proceedings of the National Academy of Sciences, 115(10).
https://doi.org/10.1073/pnas.1719588115

Mukami, A., Juma, B. S., Mweu, C., Oduor, R., & Mbinda, W. (2024). CRISPR-Cas9-
induced targeted mutagenesis of feruloyl CoA 6'-hydroxylase gene reduces postharvest
physiological deterioration in cassava roots. Postharvest Biology and Technology, 208,
112649. https://doi.org/10.1016/j.postharvbio.2023.112649

Munakata, R., Olry, A., Karamat, F., Courdavault, V., Sugiyama, A., Date, Y.,
Krieger, C., Silie, P., Foureau, E., Papon, N., Grosjean, J., Yazaki, K., Bourgaud, F., & Hehn,
A. (2016). Molecular evolution of parsnip (*Pastinaca sativa*) membrane-bound
prenyltransferases for linear and/or angular furanocoumarin biosynthesis. New
Phytologist, 211(1), 332-344. https://doi.org/10.1111/nph.13899

Neal, J. J., & Wu, D. (1994). Inhibition of Insect Cytochromes P450 by
Furanocoumarins.   Pesticide   Biochemistry   and   Physiology,   50(1),   43-50.
https://doi.org/10.1006/pest.1994.1056

Neilson, E. H., Goodger, J. Q. D., Woodrow, I. E., & Møller, B. L. (2013). Plant
chemical   defense : At   what   cost?   Trends   in   Plant   Science,   18(5),   250-258.
https://doi.org/10.1016/j.tplants.2013.01.001

Razavi, S. M. (2010). Plant Coumarins as Allelopathic Agents. International
Journal of Biological Chemistry, 5(1), 86-90. https://doi.org/10.3923/ijbc.2011.86.90

Robe, K., Conejero, G., Gao, F., Lefebvre-Legendre, L., Sylvestre-Gonon, E.,
Rofidal, V., Hem, S., Rouhier, N., Barberon, M., Hecker, A., Gaymard, F., Izquierdo, E., &
Dubos, C. (2021). Coumarin accumulation and trafficking in *Arabidopsis thaliana*: A
complex   and   dynamic   process.   New   Phytologist,   229(4),   2062-2079.
https://doi.org/10.1111/nph.17090

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR : A Bioconductor
package   for   differential   expression   analysis   of   digital   gene   expression   data.
Bioinformatics, 26(1), 139-140. https://doi.org/10.1093/bioinformatics/btp616

Rolando, C., Monties, B., & Lapierre, C. (1992). Thioacidolysis. In S. Y. Lin & C. W.
Dence (Éds.), Methods in Lignin Chemistry (p. 334-349). Springer Berlin Heidelberg.
https://doi.org/10.1007/978-3-642-74065-7_23

Roselli, S., Olry, A., Vautrin, S., Coriton, O., Ritchie, D., Galati, G., Navrot, N.,
Krieger, C., Vialart, G., Bergès, H., Bourgaud, F., & Hehn, A. (2017). A bacterial artificial
chromosome (BAC) genomic approach reveals partial clustering of the furanocoumarin
pathway   genes   in   parsnip.   The   Plant   Journal,   89(6),   1119-1132.
https://doi.org/10.1111/tpj.13450

Roumani, M., Le Bot, J., Boisbrun, M., Magot, F., Péré, A., Robin, C., Hilliou, F., &
Larbat, R. (2022). Transcriptomics and metabolomics analyses reveal high induction of
the phenolamide pathway in tomato plants attacked by the leafminer *Tuta absoluta*.
Metabolites, 12(6), 484. https://doi.org/10.3390/metabo12060484

Sarker, S. D., & Nahar, L. (2017). Progress in the chemistry of naturally occurring
coumarins. In A. D. Kinghorn, H. Falk, S. Gibbons, & J. Kobayashi (Éds.), Progress in the

Chemistry of Organic Natural Products 106 (Vol. 106, p. 241-304). Springer International Publishing. https://doi.org/10.1007/978-3-319-59542-9_3

Sarrion-Perdigones, A., Falconi, E. E., Zandalinas, S. I., Juárez, P., Fernández-del-Carmen, A., Granell, A., & Orzaez, D. (2011). GoldenBraid : An iterative cloning system for standardized assembly of reusable genetic modules. PLoS ONE, 6(7), e21622. https://doi.org/10.1371/journal.pone.0021622

Sarrion-Perdigones, A., Vazquez-Vilar, M., Palaci, J., Castelijns, B., Forment, J., Ziarsolo, P., Blanca, J., Granell, A., & Orzaez, D. (2013). GoldenBraid 2.0 : A comprehensive DNA assembly framework for plant synthetic biology. PLANT PHYSIOLOGY, 162(3), 1618-1631. https://doi.org/10.1104/pp.113.217661

Sonawane, P. D., Pollier, J., Panda, S., Szymanski, J., Massalha, H., Yona, M., Unger, T., Malitsky, S., Arendt, P., Pauwels, L., Almekias-Siegl, E., Rogachev, I., Meir, S., Cárdenas, P. D., Masri, A., Petrikov, M., Schaller, H., Schaffer, A. A., Kamble, A., ... Aharoni, A. (2017). Plant cholesterol biosynthetic pathway overlaps with phytosterol metabolism. Nature Plants, 3(1), 16205. https://doi.org/10.1038/nplants.2016.205

Tsai, H. H., & Schmidt, W. (2017). Mobilization of Iron by Plant-Borne Coumarins. Trends in Plant Science, 22(6), 538-548. https://doi.org/10.1016/j.tplants.2017.03.008 Vanholme, R., Sundin, L., Seetso, K. C., Kim, H., Liu, X., Li, J., De Meester, B., Hoengenaert, L., Goeminne, G., Morreel, K., Haustraete, J., Tsai, H.-H., Schmidt, W., Vanholme, B., Ralph, J., & Boerjan, W. (2019). COSY catalyses trans–cis isomerization and lactonization in the biosynthesis of coumarins. Nature Plants, 5(10), 1066-1075. https://doi.org/10.1038/s41477-019-0510-0

Tsai, H.-H., Rodríguez-Celma, J., Lan, P., Wu, Y.-C., Vélez-Bermúdez, I. C., & Schmidt, W. (2018). Scopoletin 8-Hydroxylase-mediated fraxetin production Is crucial for iron mobilization. Plant Physiology, 177(1), 194-207. https://doi.org/10.1104/pp.18.00178

Vialart, G., Hehn, A., Olry, A., Ito, K., Krieger, C., Larbat, R., Paris, C., Shimizu, B., Sugimoto, Y., Mizutani, M., & Bourgaud, F. (2012). A 2-oxoglutarate-dependent dioxygenase from *Ruta graveolens L.* exhibits *p*-coumaroyl CoA 2'-hydroxylase activity (C2'H) : A missing step in the synthesis of umbelliferone in plants: C2'H involved in umbelliferone synthesis. The Plant Journal, 70(3), 460-470. https://doi.org/10.1111/j.1365-313X.2011.04879.x

Villard, C., Munakata, R., Kitajima, S., Velzen, R., Schranz, M. E., Larbat, R., & Hehn, A. (2021). A new P450 involved in the furanocoumarin pathway underlies a recent case of convergent evolution. New Phytologist, 231(5), 1923-1939. https://doi.org/10.1111/nph.17458

Wang, K., Zeng, H., Dai, Y., Wang, Z., Tang, H., Li, J., Lu, X., Jiang, N., Xie, G., Zhu, Y., Zhao, Y., & Qin, M. (2024). Three types of enzymes complete the furanocoumarins core skeleton biosynthesis in *Angelica sinensis*. Phytochemistry, 222, 114102. https://doi.org/10.1016/j.phytochem.2024.114102

Wang, N., Takada, M., Sakamoto, S., Vanholme, R., Goeminne, G., Kim, H., Nagano, S., Takata, N., Kamimura, N., Uesugi, M., Izumi-Nakagawa, A., Masai, E., Mitsuda, N., Boerjan, W., Ralph, J., & Kajita, S. (2025). Effects of feruloyl-CoA 6'-hydroxylase 1 overexpression on lignin and cell wall characteristics in transgenic hybrid aspen. Frontiers in Plant Science, 16, 1543168. https://doi.org/10.3389/fpls.2025.1543168

997  Wang, X.-B., Li, G.-H., Li, L., Zheng, L.-J., Huang, R., & Zhang, K.-Q. (2008).
998 Nematicidal coumarins from *Heracleum candicans Wall*. Natural Product Research,
999 22(8), 666-671. https://doi.org/10.1080/14786410701766463

1000  Weng, J.-K., Lynch, J. H., Matos, J. O., & Dudareva, N. (2021). Adaptive
1001 mechanisms of plant specialized metabolism connecting chemistry to function. Nature
1002 Chemical Biology, 17(10), 1037-1045. https://doi.org/10.1038/s41589-021-00822-6

1003  Yan, Z., Wang, D., Cui, H., Zhang, D., Sun, Y., Jin, H., Li, X., Yang, X., Guo, H., He,
1004 X., Pan, L., Ren, X., Guo, K., & Qin, B. (2016). Phytotoxicity mechanisms of two coumarin
1005 allelochemicals from *Stellera chamaejasme* in lettuce seedlings. Acta Physiologiae
1006 Plantarum, 38(10), 248. https://doi.org/10.1007/s11738-016-2270-z

1007  Yang, S.-M., Shim, G. Y., Kim, B.-G., & Ahn, J.-H. (2015). Biological synthesis of
1008 coumarins in *Escherichia coli*. Microbial Cell Factories, 14(1), 65.
1009 https://doi.org/10.1186/s12934-015-0248-y

1010  Zhao, Y., Wang, N., Zeng, Z., Xu, S., Huang, C., Wang, W., Liu, T., Luo, J., & Kong,
1011 L. (2016). Cloning, functional characterization, and catalytic mechanism of a bergaptol
1012 O-methyltransferase from *Peucedanum praeruptorum Dunn*. Frontiers in Plant Science,
1013 7. https://doi.org/10.3389/fpls.2016.00722

1014  Zhou, H., Jian, Y., Shao, Q., Guo, F., Zhang, M., Wan, F., Yang, L., Liu, Y., Yang, L., Li,
1015 Y., Yang, P., Li, Z., Li, S., & Ding, W. (2023). Development of sustainable insecticide
1016 candidates for protecting pollinators: Insight into the bioactivities, selective mechanism
1017 of action and QSAR of natural coumarin derivatives against aphids. Journal of Agricultural
1018 and Food Chemistry, 71(47), 18359-18374. https://doi.org/10.1021/acs.jafc.3c03493