Lipid-mediated GPR32 signaling reprograms macrophage metabolism to

impair anti-tuberculous immunity

Joaquina Barros[1,2,3], Mariano Maio[1,2,3], Martín Manuel Ledesma[2], Leandro

Ferrini[1,3], María Emilia Boix[1], Natacha Faivre[3,4], Sarah Monard[3,4], José Luis

Marín Franco[2], Arnaud Métais[4], Florencia Sabbione[2], Federico Fuentes[2],

Tomas Martín Grosso[1], Agustina Juliana Errea[5], Xavier Aragone[6], Milagro

Sanchez Cunto[6], Domingo Palmero[6], Mario Matteo[6], Matías Ostrowski[1], Rafael

J Argüello[7], Ranjan Kumar Nanda[8], Emilie Layre[3,4], Olivier Neyrolles[3,4], Christel

Vérollet[3,4], Geanncarlo Lugo Villarino[3,4†], and Luciana Balboa[1,2,3,*†]

[1] Instituto de Investigaciones Biomédicas en Retrovirus y SIDA (INBIRS).

Facultad de Medicina. Universidad de Buenos Aires- Consejo Nacional de

Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina

[2] Instituto de Medicina Experimental (IMEX)-CONICET, Academia Nacional de

Medicina, Buenos Aires, Argentina

[3] International Associated Laboratory (LIA) CNRS IM-TB/HIV (1167), Buenos

Aires, Argentina / International Research Project Toulouse, France

[4] Institut de Pharmacologie et de Biologie Structurale (IPBS), Université de

Toulouse, CNRS, UPS, Toulouse, France

[5] Instituto de Estudios Inmunológicos y Fisiopatológicos - CONICET -

Universidad Nacional de La Plata, La Plata, Argentina

[6] Instituto Prof. Dr. Raúl Vaccarezza; Hospital de Infecciosas Dr. F.J. Muñiz,

Buenos Aires, Argentina

[7] Aix-Marseille University, CNRS, INSERM, CIML, Centre d'Immunologie de

Marseille-Luminy, Turing Centre for Living Systems, Marseille, France

[8] Translational Health Group, International Center of Genetic Engineering and

Biotechnology, New Delhi, India

Corresponding Author: Luciana Balboa, Paraguay 2155, Piso 11 CABA

(C1121ABG), Argentina, + 54(011) 4508-3689, luciana_balboa@hotmail.com

The authors have declared that there is no conflict of interest.

[†] These authors contributed equally to this work

Abstract

Mycobacterium tuberculosis, the causative agent of tuberculosis (TB), has evolved strategies to evade innate immunity and establish persistent infection. However, the mechanisms by which M. tuberculosis reprograms human macrophage metabolism remain incompletely defined. Tuberculous pleural effusion (TB-PE), a common extrapulmonary manifestation that frequently coexists with pulmonary TB, offers a unique, clinically relevant immunometabolic window into the TB microenvironment. Here, using patient-derived TB-PE samples, we demonstrate that this microenvironment induces a metabolic state in human macrophages that compromises their antimicrobial function. Lipidomic analysis identified an enrichment of the specialized pro-resolving mediator Resolvin D5 (RvD5), which signals through GPR32 to suppress macrophage microbicidal activity. The acellular fraction of TPE was sufficient to induce RvD5 secretion by monocytes, correlating with increased expression of RvD5 biosynthetic enzymes in pleural monocytes from TB patients. Mechanistically, RvD5-GPR32 signaling inhibited glycolysis without promoting oxidative phosphorylation, reducing HIF-1α activity and impairing intracellular M. tuberculosis control. HIF-1α stabilization restored antimicrobial function. These findings uncover the RvD5-GPR32-HIF-1α axis as a mechanism of metabolic immune suppression and a potential target for host-directed TB therapy.

Introduction

The success of *M. tuberculosis* (Mtb) as a pathogen lies, in part, in its efficient adaptation to the intracellular environment of human macrophages, which involves regulating the host's metabolic pathways to facilitate immune evasion. Macrophages are essential immune cells that can serve as a niche for intracellular mycobacterial persistence but can also limit infection by activating their microbicidal properties. In fact, innate immunity appears sufficient to clear Mtb infections in certain patients, conferring resistance[1]. This activation of macrophages towards the microbicidal program (called the M1 activation program) depends on a profound metabolic change that promotes glycolysis to the detriment of mitochondrial oxidative phosphorylation[2,3].

Tuberculous pleural effusion (TB-PE) is characterized by an accumulation of fluid in the pleural cavity of infected patients, marked by elevated levels of proteins, lipids, and specific leukocyte populations. Recent studies have begun to characterize the metabolites present in these samples, revealing differential abundance of amino acids, citric acid cycle intermediates, and free fatty acids compared with malignant pleural effusions[4]. TB-PE is typically a serous fluid and contains very few viable mycobacteria, indicating a paucibacillary infection. The development of TB-PE is likely due to a low-burden mycobacterial infection in the pleural space, which typically arises from initial parenchymal lesions. This infection triggers an immune response that not only increases the production of pleural fluid but also impairs its drainage[5].

TB-PE is the second most prevalent form of extrapulmonary TB, after TB lymphadenitis, accounting for 15-25% of TB cases worldwide. In immunocompromised populations, it may represent as much as 50% of all cases[6]. Compared to samples obtained from blood or bronchoalveolar lavage (BAL) of TB patients, TB-PE provides a more accurate reflection of the microenvironment encountered during infection. It captures the specific cellular composition and environmental factors at the site of infection, which is anatomically related to the lungs. In fact, there is a significant association between pleural TB and pulmonary TB as coexisting parenchymal disease has been identified in 48%[7] and 86%[8] of computed tomography scans conducted on patients with pleural TB. Furthermore, TB-PE samples better reflect the physiological concentrations of metabolites and compounds because they are

not diluted with saline, unlike BAL. These distinctive characteristics make TB-
PE a valuable tool for TB research.

While it has been suggested that type-1 immunity is dominantly present in TB-
PE[9], infection control within the pleural cavity is considered insufficient, as
patients frequently experience recurrent episodes of TB if left untreated[10]. This
raises the possibility of active mechanisms that impede the development of a
long-term protective immune response. In this regard, we demonstrated that the
acellular fraction of TB-PE induces the differentiation of monocytes into
immunosuppressive macrophages[11–13], converting them into lipid-laden or
foamy macrophages, thereby affecting their effector functions against bacilli[14].
We have also demonstrated that microbicidal M1 macrophages exposed to a
lipid fraction isolated from TB-PE, enriched in metabolites derived from
polyunsaturated fatty acids (PUFA), are unable to carry out the metabolic
reprogramming associated with their pro-inflammatory profile, showing reduced
glycolytic activity, as well as increased mitochondrial respiration and reduced
microbicidal activity[15]; however, the mechanism of action as well as the specific
metabolites involved in this effect remain unclear, limiting the potential to
develop targeted therapies aimed at enhancing anti-TB immune response.

In response to infection, both inflammatory and pro-resolving processes are
deployed in TB-PE. Among the mechanisms aimed at limiting the inflammatory
process and activating tissue repair and healing are those catalyzed by a family
of lipids, recently termed "specialized resolution lipid mediators" (SPMs)[16].
SPMs derive from PUFA such as linoleic acid (C18: Δ2, n-6), arachidonic acid
(C20: Δ4, n-6), eicosapentaenoic acid (EPA, 20:5, n-3), and docosahexaenoic
acid (DHA, 22:6, n-3), which are produced through enzymatic reactions
catalyzed by lipoxygenases (LOX) and/or cyclooxygenases (COX). SPMs are
composed of DHA-derived D-series protectins and resolvins, EPA-derived E-
series resolvins, and arachidonic acid-derived lipoxins.

By binding to specific receptors, resolvins modulate various inflammatory
processes, including neutrophil migration, macrophage phagocytosis, and the
expression of pro-inflammatory mediators, thereby mitigating inflammatory
pathologies[17]. The G protein-coupled receptor 32 (GPR32) has been shown to
bind to resolvins RvD1, RvD3, and RvD5[18–20]. Although direct evidence
supporting the RvD5-GPR32 axis in lung infections remains limited,

investigations into related resolvins, such as RvD1[21], yield promising results in
models of lung injury, indicating reduced neutrophil infiltration and diminished
alveolar damage. These findings strongly implicate the RvD1-GPR32 axis as a
critical regulatory pathway. This mechanism is especially pertinent in the
context of bacterial pneumonia and viral infections, like influenza, where a
carefully calibrated immune response is essential. Despite the known
importance of resolvins, no study to date has investigated the role of RvD5 in
TB. In this study, we performed a lipidomic analysis of TB-PE and found high
levels of the lipid mediator RvD5. We characterized its function by exposing M1
macrophages to this lipid mediator and found that RvD5 is sufficient to impair
their microbicidal activity and glycolytic metabolism. We functionally validate
that the direct binding mediates this impairment between RvD5 and its receptor
GPR32, expressed in macrophages, whose activation inhibits HIF1α. Restoring
HIF1α with a chemical activator was sufficient to rescue the functionality of M1
macrophages exposed to RvD5.
140
141

Material and methods

Human Subjects

Buffy coats from healthy donors were prepared at the Centro Regional de Hemoterapia Garrahan (Buenos Aires, Argentina) according to the institutional guidelines (Resolution No. CEIANM-664/07). Informed consent was obtained from each donor before blood collection.

Pleural effusions (PE) were obtained by therapeutic thoracentesis by physicians at the Hospital F. J Muñiz (Buenos Aires, Argentina). The diagnosis of TB pleurisy was based on a positive Ziehl–Neelsen stain or Löwenstein–Jensen culture from PE and/or histopathology of pleural biopsy, and was further confirmed by an Mtb-induced IFNγ response and an adenosine deaminase-positive test[10]. None of the patients had multidrug-resistant TB. For this study, we used acellular fractions of PE stored at -80°C for lipid analysis, and fresh paired pleural fluids and blood mononuclear cells (PFMC and PBMC) for assessment of metabolic profiles and GPR32 expression. PE samples were divided by etiology. We utilized five pools of the acellular fraction from TB patients (10-12 samples per pool), one pool of malignant PE (PE-C) with six patients (including mesothelioma, lung carcinoma, or metastatic PE), one pool of parapneumonic PE (PE-PNI) with eight patients, and three pools of heart failure-related transudates (HF-PE, five samples per pool). The research was carried out in accordance with the Declaration of Helsinki (2013) of the World Medical Association. It was approved by the Institutional Review Board of Hospital F. J. Muñiz (approval number PRIISA.BA – 5334, renewed in December 2024) via the Buenos Aires Computerized Health Research Registry Platform. Written informed consent was obtained before sample collection.

Bacterial strain

The Mtb laboratory strain, H37Rv, was grown at 37°C in Middlebrook 7H9 medium supplemented with 10% albumin-dextrose-catalase (both from Becton, Dickinson, New Jersey, USA) and 0.05% Tween-80 (Sigma-Aldrich, St. Louis, MO, USA). The red fluorescent protein (RFP)-expressing Mtb strain was gently provided by Dr. Fabiana Bigi (INTA, Castelar, Argentina).

<u>Preparation of PE pools</u>

The PE were collected in heparin tubes and pools of the acellular fractions were

prepared, as previously described[15].

<u>Preparation of human monocyte-derived Macrophages</u>

Monocytes were isolated from PBMC, as previously described[15], and

differentiated in the presence of human recombinant Granulocyte-Macrophage

Colony-Stimulating Factor (GM-CSF, Peprotech, New York, USA) at 50 ng/ml.

Cells were allowed to differentiate for five to seven days into M0 macrophages.

<u>Macrophages treatments</u>

Macrophages were polarized toward M1 profile with IFNγ (10 ng/ml) plus LPS

(10 ng/ml), exposed (or not) to either RvD5 (33 nM), vehicle (ethanol), or 20%

v/v of PE during 24 h to generate M1 and treated-M1 cells, respectively. When

indicated, macrophages were pre-incubated with GPR32-blocking antibodies

(10 µg/mL; GTX71225, GeneTex) or their rabbit IgG isotype control (GeneTex)

for 30 minutes before exposure to RvD5. Alternatively, M1 macrophages were

generated only with IFNγ and either stimulated with Mtb γ-irradiated H37Rv

strain (NR-49098, BEI Resources, USA) (equivalent to a multiplicity of infection

"MOI" of 1:1) or infected with Mtb, and exposed (or not) to RvD5 (33 nM) for 24

h. Also, when specified, cells were exposed (or not) to DMOG (200 µM) during

LPS or Mtb infection for 24 h.

In some experiments, M1 macrophages were treated with lipids extracted from

PE. Briefly, lipids were isolated from PE, dried under a gentle nitrogen stream,

and reconstituted in a small volume of methanol. This volume was normalized

to the original volume of each pleural fluid sample to maintain the relative lipid

abundances for subsequent cellular assays.

<u>Determination of lactate</u>

Lactate production in the culture medium was measured using the Lactate Kit

from Wiener (Argentina). The absorbance was read using a Biochrom Asys

UVM 340 Microplate Reader and software.

<u>Lipid preparation and lipidomic analysis of PUFAs from TB-PE</u>

Lipid extraction from TB-PE was based on the Bligh and Dyer protocol, as

previously described[15]. PUFAs were then extracted on a solid phase as follows.

4 µL of the internal standards mixture (deuterium-labeled compounds) was

added to the lipid extracts, which were resuspended in 300 µL of cold methanol.

After centrifugation at 2000 g for 15 min at 4°C, the supernatants were

transferred into 2 mL 96-well deep plates and diluted to 2 mL with H_2O.

Samples were then submitted to solid-phase extraction (SPE) using an OASIS

HLB 96-well plate (30 mg/well, Waters), pretreated with MeOH (1 mL), and

equilibrated with 10% MeOH (1 mL). After the sample application, the extraction

plate was washed with 10% MeOH (1 mL). After drying under aspiration, lipid

mediators were eluted with 1 mL of MeOH. Before LC-MS/MS analysis,

samples were evaporated under nitrogen gas and reconstituted in 10 µL of

MeOH.

The samples were analyzed at the MetaToul lipidomic platform (I2MC, INSERM

1048, Toulouse, France) as previously described[22]. Briefly, lipid mediators were

separated on a ZorBAX SB-C18 column (2.1 mm, 50 mm, 1.8 µm) (Agilent

Technologies) using an Agilent 1290 Infinity HPLC system (Agilent

Technologies) coupled to an ESI-triple quadruple G6460 mass spectrometer

(Agilent Technologies). Data was acquired in Multiple Reaction Monitoring

(MRM) mode with optimized conditions (ion optics and collision energy). Peak

detection, integration, and quantitative analysis were performed using Mass

Hunter Quantitative Analysis software (Agilent Technologies) based on

calibration lines constructed from commercially available eicosanoid standards

(Cayman Chemicals).

<u>Determination of intracellular RvD5 abundances</u>

Monocytes from healthy donors were stimulated with TB-PE (20% v/v) for 30

minutes. Cells were washed with ice-cold PBS (without Mg^{2+} and Ca^{2+}) and

scraped into ice-cold methanol/5 mM EGTA (2:1 v/v), snap-frozen in liquid

nitrogen, and stored at -80°C until SPE and LC-MS analysis, as described

above.

<u>SCENITH assays</u>

SCENITH experiments were performed using the SCENITH kit, which contains
all reagents and anti-puromycin antibodies (www.scenith.com). Briefly, PFMCs,
PBMCs, and macrophages were treated for 40 minutes at 37°C in the presence
of the indicated inhibitors of various metabolic pathways and puromycin. After
incubation, puromycin was stained with fluorescently labeled anti-Puromycin
monoclonal antibodies (clone R4743L-E8) conjugated to Alexa Fluor 647 or
Alexa Fluor 488, and analyzed by flow cytometry. The impact of the various
metabolic inhibitors was quantified, as previously described[23,24].

<u>Measurement of cell respiration with Seahorse flux analyzer</u>
Bioenergetics in human macrophages were determined using a Seahorse
XFe24 analyzer. ATP production rates and relative contribution from glycolysis
and oxidative phosphorylation (OXPHOS) were measured by the Seahorse XF
Real-Time ATP Rate Assay kit. Macrophages (250,000 cells/well) were cultured
in triplicate per condition. The assay was performed in XF Assay Modified
DMEM. Three consecutive measurements were performed under basal
conditions and after sequential addition of oligomycin (OM) and rotenone
(ROT/AA) (Agilent, USA). Extracellular acidification (ECAR) and oxygen
consumption rates (OCR) were measured. Mitochondrial ATP production rate
was determined by the decrease in the OCR after oligomycin addition.
Conversely, the complete inhibition of mitochondrial respiration with rotenone
plus antimycin A allows accounting for mitochondrial-associated acidification
and, when combined with proton efflux rate (PER) data, calculation of the
glycolytic ATP production rate. OCR and ECAR values were normalized by first
performing brightfield imaging before the assay, then using ImageJ software to
calculate the cellular area per condition, and finally importing these cellular area
values into Agilent's Wave software to divide the metabolic data by cell quantity,
thereby accounting for variations in cell numbers between experiments and
conditions.
Additionally, the Mito Stress Test was performed to measure mitochondrial
oxidative phosphorylation rate. Basal respiration was calculated as the last
measurement before the addition of OM minus the non-mitochondrial
respiration, which is the minimum rate measurement after the addition of
ROT/AA. Estimated ATP production is the last measurement before OM

addition minus the minimum rate after OM. Maximal respiration rate (Max) was

defined as the OCR after addition of OM and FCCP at 2µM (for 2.5 minutes)

minus the non-mitochondrial OCR. Spare respiratory capacity (SRC) was

defined as the difference between maximum and basal respiration. Three

consecutive measurements were performed under basal conditions and after

the sequential addition of the following electron transport chain inhibitors.

<u>RNA-Single Cell Sequencing Analysis</u>

Previously published single cell transcriptomic data available in Genome

Sequence Archive in the BIG Data Center, Beijing Institute of Genomics,

Chinese Academy of Sciences, under accession numbers HRA000910 and

HRA00036[25], were used to evaluate the expression of enzymes involved in

SMPs' biosynthesis (ALOX5, ALOX12, ALOX 15 and ALOX15B), including five

samples of PBMC and six samples of PFMC from TB patients. The Seurat

package (version 4.3.0.1) in RStudio was used to read the pre-processed files.

The filtering process involved discarding cells with fewer than 200 or more than

5000 total genes, fewer than 200 or more than 20,000 total counts, and

mitochondrial read levels above 10%. Expression levels were then normalized

using the log transformation (scale factor = 10,000), based on the 2,000 most

variable genes. The samples were integrated using the reciprocal PCA (RPCA)

algorithm. The different cell clusters were identified using the shared nearest

neighbor (SNN) modularity optimization algorithm, and dimensionality reduction

was performed using the UMAP (Uniform Manifold Approximation and

Projection) algorithm. Individual cells were annotated using the SingleR

package version 2.2.0 and the HumanPrimaryCellAtlasData database (label.fine

version).

<u>Intracellular expression of HIF1α</u>

Macrophages were permeabilized with triton 0.3% in FACS buffer and

incubated with PE-anti-HIF1α (clone 546-16, Biolegend) for 30 minutes. After

that, cells were analyzed by flow cytometry using the FACSCalibur cytometer

(BD Biosciences, San Jose, CA, USA). The macrophage population was gated

according to its Forward and Size Scatter properties. The median fluorescence

intensity (MFI) was analyzed using FCS Express V3 software (*De Novo*
Software, Los Angeles, CA, USA).

GPR32 surface expression

To monitor GPR32 expression in CD14$^+$ cells from pleural effusion and blood
paired samples of TB vs. non-TB patients, cells were incubated with a primary
anti-GPR32 antibody (GTX71225; GeneTex) for 40 minutes at 4°C. Next, the
cells were incubated with a secondary antibody conjugated to Alexa Fluor 488
(anti-Rabbit IgG, 111-546-144) to label anti-GPR32. After an anti-CD14-
PerCP/Cy7 (clone HCD14, Biolegend) was employed, the cell viability marker
Zombie Violet (Biolegend) was used.

Fluorescence microscopy

Fluorescence microscopy images of fixed IFNγ-activated (M1) macrophages
exposed (or not) to RvD5 that were first infected with Mtb-RFP (MOI 1-3:1) for
24 h and then fixed. Cell nuclei were labeled with DAPI, and the autophagy
marker LC3 protein (anti-LCE3B; dilution 1:300; #NB100-2220) was identified
by indirect immunofluorescence using Alexa Fluor 488 conjugated antibodies,
whereas the late endosome marker, LAMP2, was identified by indirect
immunofluorescence using a specific conjugated antibody (anti-LAMP2 Alexa
Fluor 633; dilution 1:200; #A21126). Samples were imaged on a FluoView
FV1000 confocal microscope (Olympus, Tokyo, Japan) equipped with a Plapon
60X/NA1.42 objective and on the Zeiss LSM 800 confocal microscope (Zeiss,
Oberkochen, Germany).

IL-1β determination

The amount of human IL-1β was measured by ELISA, according to the
manufacturer's instructions (ELISA MAX Deluxe Kits from Biolegend). The
detection limit was 15.6 pg/ml.

Infection of human macrophages with Mtb

Infections were performed in the biosafety level 3 laboratories at the *Unidad*
Operativa Centro de Contención Biológica (UOCCB), ANLIS-MALBRAN (in
2022) and at *Instituto de Investigaciones Biomédicas en Retrovirus y SIDA*

(INBIRS)-CONICET-UBA (in 2023-24), according to the biosafety institutional
guidelines (both institutes located in Buenos Aires). Macrophages were infected
with Mtb H37Rv strain at a MOI of 0.5-3:1 during 1 h at 37°C. Next, extracellular
bacteria were removed gently by washing with pre-warmed PBS, and cells were
cultured in RPMI-1640 medium supplemented with 10% FBS and gentamicin
(50 µg/ml). The extracellular bacteria were then removed gently by washing with
pre-warmed PBS, and cells were cultured in RPMI-1640 medium supplemented
with 10% FBS and gentamicin (50 µg/ml) for 4, 24 and 72h. At different time
points, macrophages were lysed in 0.1% SDS and neutralized with 20% BSA in
Middlebrook 7H9 broth. Serial dilutions of the lysates were plated in triplicate,
onto 7H11-Oleic Albumin Dextrose Catalase (OADC, Becton Dickinson) agar
medium for CFU scoring at 21 days later.
356

<u>Quantification and Statistical Analysis</u>
Data are presented as scatter plots where each symbol represents one
independent donor or biological sample, with means and standard error of the
mean (SEM) shown unless otherwise indicated. Each experiment corresponds
to a separate donor. Statistical tests were selected based on experimental
design and data distribution. Comparisons between two unpaired groups used
the Mann–Whitney test or unpaired t-test; comparisons between two paired
groups used the Wilcoxon signed-rank test or paired t-test. Comparisons among
three or more groups used one-way or two-way ANOVA followed by post hoc
tests: Dunnett's test for comparisons against a single control; Tukey's test for
comparisons among all groups; Holm–Sidak test for specified comparisons in
multi-factor designs; or the Benjamini–Krieger–Yekutieli two-stage linear step-
up procedure to control the false discovery rate. Categorical data were analyzed
with Fisher's exact test, correlation analyses used Spearman's rank correlation
coefficient, and specialized designs employed nested t-tests for hierarchical
data or mixed-effects models for repeated measures. Specific tests for each
figure are detailed in the corresponding legends. A p-value < 0.05 was
considered significant. Analyses were performed using GraphPad Prism
(version 8.0.1) or R.

Results

Enrichment in specialized resolution lipid mediators correlates with impaired macrophage glycolysis in tuberculous pleural effusions.

We previously found that PUFA metabolites in TB-PE inhibit the glycolytic activity of M1 macrophages[15]. To explore whether this inhibition was specific to pleural TB lipids, we compared the ability of the PUFA extracted from the acellular fraction of TB-PE and non-TB patients to inhibit glycolysis in M1 macrophages (activated by IFNγ and LPS), by assessing lactate release from these cells. The lipid fractions isolated from TB-PE showed greater inhibition of lactate release by M1 macrophages than PE from non-tuberculous patients with other pathologies, such as heart failure, pneumonia, and cancer (**Figure 1A**). To further investigate the candidate components responsible for this inhibition, a targeted lipidomic analysis was conducted on lipids isolated from TB and non-TB patients (n=13 and 5, respectively). Among the 30 screened PUFA metabolites, four exhibited significantly higher levels in TB-PE compared to non-TB PE (**Figure 1B-D**). Notably, these molecules were specialized pro-resolving lipid mediators (SPMs) derived from omega-3 fatty acids, i.e., 7 Maresin R1 (7MaR1), RvD5, Protectin Dx (PDx), and 18-HEPE (**Figure 1B-D**). Furthermore, the abundance of these SPMs was significantly correlated with the decline in glycolysis induced by each PE in monocyte-derived M1 macrophages from healthy donors (**Figure 1E**). Our data indicate that SPMs are highly abundant in TB-PE, and their levels correlate with TB-PE's capacity to suppress M1 macrophage glycolytic activity.

401
Figure 1. Enrichment in specialized resolution lipid mediators correlates
with impaired macrophage glycolysis in tuberculous pleural effusions. (A)
M1 macrophages were exposed or not to the polyunsaturated fatty acid (PUFA)
extracted from the acellular fraction of tuberculous (TB; five pools and eight
individual samples) or non-tuberculous (non-TB; five pools) effusions. The
percentage of release of lactate in the supernatant compared to untreated M1
cells is shown. Data are presented as individual values with the mean ± SEM.
Statistical significance was determined by a Mann-Whitney test. In the figure,
sample pools are distinguished by color and symbol. Effusions from tuberculous
(TB) patients are shown as yellow circles (●), with pooled samples containing
black dots. The three cardiac failure pools are black circles (●), the single
pneumonia pool is dark gray (●) and the single malignancy pool is light gray
represent the single malignancy pool (●). **(B)** Heat map illustrating the relative
abundance (Z-score normalized) of 30 PUFA lipid species detected through
lipidomic analysis in pleural effusions classified as TB and non-TB. Lipids are
grouped according to their precursor PUFA. The specialized pro-resolving
mediator (SPM) family is highlighted (red box). Non-TB effusions comprise
heart failure (HF), pneumonia (PNI), and cancer (C). **(C)** The volcano plot
depicts fold changes in lipid levels between TB (n=13) and non-TB (n=5) pleural
effusion samples, along with their statistical significance (p-value), for all
detected metabolites. **(D)** Abundances of selected SPMs (RvD5, 7-MaR1, 18-
HEPE, and PDx). Samples and pool symbols are consistent with panel A. A
two-way ANOVA with Benjamini, Krieger, and Yekutieli correction for multiple
comparisons was performed. **(E)** Correlations between lactate release by M1
macrophages exposed to TB or non-TB pleural effusions and the abundance of
specific lipids. Samples and pool symbols are consistent with panel A.
Spearman's correlation coefficient (r) and significance levels are reported. LA:

linoleic acid, AA: arachidonic acid, DHA: docosahexaenoic acid, EPA:
eicosapentaenoic acid.

Local Synthesis of RvD5 by Monocytes and Expression of its Receptor
GPR32 on Macrophages in Tuberculous Pleuritis.

Our next objective was to investigate the potential origins of SMPs in the TB-
PE. To address this, we analyzed the expression of key enzymes involved in
SPM biosynthesis, namely arachidonic acid 5-lipoxygenase (5-LO, ALOX5),
along with arachidonic acid 12- and 15-lipoxygenating paralogues (ALOX12,
ALOX15, and ALOX15B)[26], in peripheral blood mononuclear cells (PBMC) and
pleural fluid mononuclear cells (PFMC) from TB patients. This analysis utilized
publicly available single-cell RNA-Seq datasets (accession numbers
HRA000910 and HRA00036). According to the literature[27], lymphocytes (mainly
T cells) dominate leukocyte infiltrates, while monocytic populations are
underrepresented in TB-PE compared to blood (**Figure 2A)**. While monocytes
from both compartments showed minimal expression of ALOX12, ALOX15, and
ALOX15B, that of ALOX5 was significantly upregulated in pleural CD14$^+$ cells
compared to circulating monocytes (**Figure 2B, S1A**). In CD14$^+$ monocytes
specifically, ALOX5 expression increased ~4.4-fold in PFMC versus PBMC
(predicted means: ~0.33 to ~1.48), with no significant changes observed in
other cell types (**Figure 2C, S1B**). This selective upregulation suggests that
pleural CD14$^+$ cells are the primary mononuclear cell population responsible for
local specialized pro-resolving mediators in the tuberculous pleural
environment.

We also examined RvD5 levels in monocytes from healthy donors exposed to
the acellular fraction of TB-PE and compared them with those of unexposed
monocytes. RvD5 was significantly detected in extracts from monocytes treated
with the acellular fraction of TB-PE (**Figure 2D**). Together, these findings
indicate that monocytes produce RvD5 in response to environmental factors in
the pleural cavity of patients with pleural TB.

Building on these results, we next explore the potential involvement of the
RvD5-GPR32 axis in pleural TB pathogenesis. Previous studies have shown

that GPR32 is highly expressed on macrophages and its activation induces
macrophage polarization from a pro-inflammatory to a pro-resolutive
phenotype[28]. We first confirmed the presence of the RvD5 receptor, GPR32, on
cells within the pleural cavity. Expression was significantly higher in pleural
CD14+ cells than in their circulating counterparts, a pattern consistent in both
TB and non-TB patients. Furthermore, GPR32 expression in pleural CD14+ cells
was clearly increased in TB patients compared with non-TB patients **(Figure
2E-F)**. This demonstrates the receptor's expression in cells locally exposed to
RvD5.

**Figure 2. CD14+ cells synthesize RvD5 and its receptor GPR32 in
tuberculous pleural effusions. (A)** Single cell RNA-seq data visualized
through a 2D UMAP plot: visualization of mononuclear cells from blood and
pleural fluid of TB patients, with cells color-coded by major cell types (PBMC,
n=5; PFMC, n=5). **(B)** Expression of enzymes involved in specialized pro-
resolving mediator (SPM) biosynthesis. Violin plots depict the expression levels
of key lipoxygenase genes (ALOX5, ALOX12, ALOX15, ALOX15B) across B
cell, monocyte (CD14+), NK cell, and CD4 and CD8 T cell lineages. Data are
shown for peripheral blood mononuclear cells (PBMCs; upper panels, n=5) and
pleural fluid mononuclear cells (PFMCs; lower panels, n=6) from patients with
TB. **(C)** Box-and-whisker plots represent the mean-to-variability ratio (μ/σ) of
normalized ALOX5 expression in monocyte cells (CD14+) from TB patients. The
ratio was calculated separately for PBMC (n=5 patients) and PFMC (n=4

patients). The total number of cells in each group is indicated on the x-axis. Each box summarizes the distribution of μ/σ values across patients within each group, with boxes showing the interquartile range (IQR), whiskers extending to 1.5×IQR, and center lines indicating median values. Statistical comparison was performed using a nested t-test. **(D)** Quantification of RvD5 in monocytes treated in vitro with either PBS 1x or TB-PE (n=6). The proportion of RvD5-producing monocytes out of a total of 6 independent donors is shown above each bar. Statistical analysis was performed using Fisher's exact test. **(E)** Flow cytometry analysis was performed to compare GPR32 expression between CD14+ cells from peripheral blood and pleural effusions. Representative dot plots showing GPR32 expression on CD14$^+$ cells from paired blood and TB pleural mononuclear cells. Representative flow cytometry dot plots show GPR32 expression on paired peripheral blood and pleural fluid CD14$^+$ mononuclear cells from a TB patient. **(F)** GPR32 levels, expressed as median fluorescence intensity normalized to fluorescence-minus-one controls (MFI/FMO), on CD14$^+$ cells from paired blood and pleural fluid samples. Data includes TB patients (n=4) and non-TB patients (n=3). Individual data points are shown as scatter plots; horizontal bars represent the mean ± SEM. Statistical comparisons were performed using a paired t-test (paired blood vs. pleural expression) and an unpaired t-test (TB vs. non-TB groups). PBMC: peripheral blood mononuclear cells; PFMC: pleural fluid mononuclear cells.

The RvD5-GPR32 axis mediates the inhibition of macrophage glycolysis by tuberculous pleural effusions

To determine whether RvD5 was responsible for the inhibition of macrophage glycolysis triggered by TB-PE, we conducted functional assays using commercially available RvD5. As previously shown[29], compared to non-activated macrophages (M0), M1 macrophages typically exhibited high glycolytic activity by increasing lactate release (**Figure 3A**). When physiological doses of RvD5 (17 and 33nM), usually found in TB-PE, were added to M1 macrophages, it efficiently inhibited lactate production similarly to TB-PE (**Figure 3A and S3A**). Notably, RvD5 treatment did not affect cell viability (**Figure S3B**). Given the pivotal role of HIF1α in the TB-PE-induced impairment of M1 macrophage functions[15], we explored whether RvD5 influenced the upregulation of HIF1α in M1 cells. Our findings revealed that RvD5 reduced HIF1α expression in M1 macrophages (**Figure 3B**). Similarly, RvD5 suppressed lactate release when M1 macrophages (activated by IFNγ) were challenged with irradiated *M. tuberculosis* (**Figure 3C**).

Given the high expression of GPR32 observed in CD14$^+$ pleural cells, we investigated whether the inhibitory effects of exogenous RvD5 are mediated

through this receptor. First, in line with the literature[30], we confirmed GPR32
expression in M1 macrophages (**Figure 3D**). Next, we observed that blocking
GPR32 fully restored lactate release in M1 macrophages exposed to either
RvD5 or TB-PE (**Figure 3E**), indicating that GPR32 signaling by TB-PE lipids
was sufficient to alter the metabolism of M1 macrophages. Moreover, blockade
of GPR32 also restored HIF1α levels in both RvD5 and PE-treated M1
macrophages (**Figure 3F**).

These findings collectively demonstrate that RvD5 inhibits glycolysis in M1
macrophages through a GPR32-dependent mechanism, suppressing HIF1α
activity.

**Figure 3. The RvD5-GPR32 axis mediates the inhibition of macrophage
glycolysis induced by TB pleural effusions. (A)** Lactate release from
unstimulated (M0) or LPS/IFNγ-activated (M1) macrophages treated for 24h
with RvD5 (33 nM), 20% v/v TB pleural effusion (TB-PE), or vehicle (ethanol)
(n=8). Data were analyzed by one-way ANOVA with Dunnett's multiple
comparisons test. **(B)** HIF1α expression in M0 or M1 macrophages exposed to
RvD5 (n=6) or vehicle (n=4). Data was analyzed using a mixed-effects model
with Sidak´s multiple comparisons test. **(C)** Lactate release from unstimulated
(M0) or IFNγ-activated (M1) macrophages exposed to irradiated *M. tuberculosis*
(iMtb) and treated with RvD5 or vehicle for 24h (n=7). Data was analyzed by
one-way ANOVA with Dunnett's multiple comparisons test. **(D)** Representative
flow cytometry histogram showing GPR32 surface expression on LPS/IFNγ-
stimulated (M1) human macrophages, with an isotype control. **(E)** Lactate
release from LPS/IFNγ-stimulated M1 macrophages pre-treated with a
neutralizing anti-GPR32 antibody (αGPR32) or isotype control before addition of

RvD5 or 20% v/v TB-PE (n=6). **(F)** HIF1α expression in LPS/IFNγ-stimulated
M1 macrophages pre-treated with αGPR32 or isotype control before addition of
RvD5 or 20% v/v TB-PE (n=5). The horizontal dotted line indicates the mean
value from untreated M0 macrophages. Data in (E) and (F) were analyzed by
two-way ANOVA. For the factor of antibody treatment, Holm-Sidak's multiple
comparisons test was applied (αGPR32 vs. isotype). For the stimulation
condition factor, Tukey's multiple comparisons test was applied (M1 vs. PE-M1
vs. RvD5-M1). All quantitative data are presented as scatter plots showing
individual replicates with mean ± SEM.
566
567

Activation of the RvD5-GPR32 axis reduces glycolysis without affecting
mitochondrial bioenergetics in M1 macrophages
To investigate if the TB-associated microenvironment affects the metabolic
profile of pleural monocyte/macrophages, we compared the *ex vivo* metabolic
profile of CD14[+] cells from TB-PE with those from pleural effusions of other
etiologies (nonTB-PE). This was achieved by monitoring changes in protein
translation upon treatment with oligomycin and/or 2-deoxy-D-glucose using the
SCENITH assay[24]. Our findings indicated a reduced glycolytic capacity in cells
residing in the TB pleural cavity compared to blood cells (**Figure 4A**).
Interestingly, pleural macrophages from TB patients undergoing antibiotic
treatment exhibited a significant increase in glycolysis compared to those from
untreated patients (**Figure 4A**). Consistent with our previous findings showing
that TB patients display elevated glycolysis in systemic monocytes[23]. These
data indicate a similar increased glycolytic activity in the blood of patients with
pleural TB, which tends to normalize with antibiotic treatment (**Figure 4A**).
583

Additionally, we evaluated the impact of RvD5 on the relative contributions of
metabolic pathways to the cellular energy metabolism of M1 macrophages
using SCENITH. According to the expected glycolytic profile of M1 activation
program[31], M1 macrophages exhibited a lower reliance on OXPHOS alongside
an increase in the glycolytic capacity compared to unstimulated cells (M0)
(**Figure 4B**). However, this metabolic shift in M1 macrophages was impaired
upon addition of exogenous RvD5 **(Figure 4B and Figure S2 and S3)**. Of note,
while TB-PE has been shown to promote lipid body accumulation in
macrophages[11,14], RvD5 did not induce lipid body formation, at least at the
tested doses **(Figure S4)**. These findings indicate that RvD5 reduces the

glycolytic activity of M1 macrophages independently of lipid body accumulation,
unlike TB-PE.

To examine the effect of RvD5 on the OXPHOS metabolic state in
macrophages, we performed the Seahorse XF Cell Mito Stress Test in M0, M1,
and RvD5-treated M1 cells. To characterize the overall energy phenotypes of
these macrophages, basal oxygen consumption rate (OCR) was plotted against
extracellular acidification rate (ECAR). The resulting metabolic phenogram
revealed that RvD5 shifts the energy metabolism of M1 macrophages towards a
less glycolytic profile without impacting aerobic metabolism (**Figure S5A**). This
shift is further reinforced by a small but significant decrease in basal ECAR
levels observed in RvD5-treated M1 cells compared to untreated M1 cells
(**Figure 4C**), aligning with the metabolic characterization from SCENITH and
lactate measurements described earlier (**Figure 3A and 4B**).

Analysis of respiratory parameters indicates substantial reductions in ATP-
linked OCR and maximal respiratory capacity in M1 macrophages compared to
M0 cells. While RvD5 treatment did not reverse these M1-induced deficits, it
caused a further, specific reduction in maximal respiratory capacity (**Figure 4D**).

To further evaluate intracellular ATP production, we measured the basal ECAR
and OCR to calculate ATP production rates from glycolysis (GlycoATP) and
mitochondrial OXPHOS (MitoATP). M1 macrophages exhibited higher ATP
production rates from glycolysis compared to RvD5-treated M1 macrophages
(**Figure S5A, C-D**), with no differences in ATP production from mitochondrial
respiration (**Figure S5A-B and D**). Consistent with SCENITH findings, the
relative contribution of MitoATP to total ATP production increased, while the
contribution of GlycoATP decreased in RvD5-treated M1 macrophages
compared to untreated M1 cells (**Figure 4E**). Furthermore, in line with these
results, M1 macrophages showed lower mitochondrial mass than control cells,
but this parameter remained unchanged upon RvD5 treatment **(Figure S6A-B)**.
Similarly, transmission electron microscopy (TEM) revealed no differences in
mitochondrial morphology between RvD5-treated and untreated M1
macrophages (**Figure S6C**). These findings suggest that RvD5 reduces

glycolysis without increasing aerobic metabolism, thereby promoting a less
energetically active metabolic state.

Finally, we assessed whether specific inhibition of GPR32 could restore M1
metabolism in RvD5-treated macrophages. Our results revealed that blocking
GPR32 led to a notable shift in RvD5-M1 cells, driving them towards heightened
glycolysis, as evidenced by an increase in basal ECAR (**Figure 4F and Figure
S7A and C**). Notably, key respiratory parameters, including basal OCR and
maximal respiratory capacity, were significantly diminished in RvD5-M1
macrophages following GPR32 inhibition (**Figure 4G**), suggesting a
compensatory reduction in OXPHOS to accommodate the increased glycolysis.
These findings were further corroborated by ATP production analysis, which
yielded a significant increase in GlycoATP levels and a corresponding decrease
in MitoATP levels upon GPR32 inhibition (**Figure S7B-D**). Additionally, the
relative contribution of GlycoATP to total ATP production increased, while that
of MitoATP decreased, compared with cells treated with the isotype control
(**Figure 4H**). This metabolic reprogramming highlights a pronounced shift
towards glycolytic reliance at the expense of the OXPHOS activity.

Collectively, these results underscore the critical role of the RvD5-GPR32 axis
in suppressing glycolytic activity in M1 macrophages, maintaining a
hypometabolic state without increasing OXPHOS flux.

**Figure 4. Activation of the RvD5-GPR32 Axis Reduces Glycolysis Without
Affecting Mitochondrial Bioenergetics in M1 macrophages. (A)** Basal
glycolysis measured via SCENITH in CD14⁺ cells from paired blood
mononuclear cells (PBMC) and pleural fluid mononuclear cells (PFMC) of
untreated TB patients (n=5), treated TB patients (n=3), and non-TB patients
(n=2). Statistical analysis: paired t-test for PFMC vs. PBMC within each group;
Mann–Whitney test for untreated vs. treated TB patients. **(B)** Mitochondrial
dependence and glycolytic capacity analyzed via SCENITH in M0, M1, and
RvD5-treated M1 macrophages (n=4). Data was analyzed using one-way
ANOVA with Dunnett's multiple comparisons test. **(C)** Extracellular acidification
rate (ECAR) measured in M0 and M1 macrophages treated or not with RvD5
(n=6). Data was analyzed using one-way ANOVA with Sidak's multiple
comparisons test. **(D)** Oxygen consumption rate (OCR) traces and respiration
parameters for M0 and M1 macrophages treated or not with RvD5. Respiratory
parameters include basal respiration, ATP-linked respiration, proton leak,
maximal respiration, spare respiratory capacity (SRC), and non-mitochondrial
respiration. Representative trace from one of six independent experiments is
shown; bar graphs show means ± SEM (n=6). Data was analyzed using two-
way ANOVA with the Benjamini, Krieger and Yekutieli two-stage linear step-up
procedure. **(E)** Percent contributions of ATP from mitochondrial oxidative
phosphorylation (MitoATP) and glycolysis (GlycoATP) in M0 and M1
macrophages treated or not with RvD5 (n=5). Data was analyzed using one-
way ANOVA with Sidak's multiple comparisons test. **(F)** ECAR in M1
macrophages treated with RvD5, with or without neutralizing anti-GPR32
antibody (αGPR32) (n=3). Statistical analysis: paired t-test. **(G)** OCR traces and
respiration parameters for RvD5-treated M1 macrophages in the presence or
absence of αGPR32 (n=3). Data were analyzed using two-way ANOVA with

680   Benjamini, Krieger, and Yekutieli procedure. **(H)** Percent contributions of
681   MitoATP and GlycoATP in RvD5-treated M1 macrophages in the presence or
682   absence of αGPR32 (n=3). Statistical analysis: paired t-test. All quantitative
683   data are presented as scatter plots showing individual donors; means ± SEM
684   are indicated.
685

RvD5 boosts a pro-resolving phenotype in M1 macrophages at the
expense of its mycobacterial control potential

Since RvD5 is an SPM, we assessed the acquisition of pro-resolutive features

by M1 macrophages exposed to RvD5. M1 macrophages displayed low

efferocytic activity, which was significantly enhanced by RvD5 treatment, as

evidenced by an increased engulfment of apoptotic neutrophils (**Figure S8A**).

This improvement was accompanied by an upregulation of MerTK, a major

efferocytic receptor[32], following RvD5 treatment (**Figure S8B**), along with an

increased release of the anti-inflammatory cytokine IL-10 (**Figure S8C**). In

contrast, the expression levels of other efferocytosis receptors, such as CD14[33]

and the scavenger receptor CD36[34], remained unchanged between M1

macrophages and RvD5-treated M1 cells (**Figure S8D**).

Then, we examined the capacity of M1 macrophages to engulf viable Mtb.

There was no significant change in the initial bacterial uptake between RvD5-

treated M1 macrophages and untreated cells (**Figure 5A**). However, RvD5

treatment lowers IL-1β secretion by M1 macrophages in response to Mtb

infection and impaired strongly their microbicidal activity (**Figure 5B-C**).

Additionally, RvD5 prevented the induction of autophagy in Mtb-infected

macrophages (**Figure 5D**), a key mechanism through which macrophages

control bacterial growth[35]. Consistent with the autophagy-dependent

mycobacterial degradation in enhancing antigen presentation to T cells[36,37], we

observed reduced CD4 T cell activation in antigen-presenting cell (APC)-

dependent responses against irradiated Mtb (iMtb) when RvD5 was added to

the cocultures (**Figure S19A-D**). Of note, this effect was absent in APC-

independent T cell activation (**Figure S9E-G**), providing additional evidence that

RvD5 compromises macrophage function.

Altogether, our results indicate that the omega-3-derived lipid mediator RvD5
reprograms the proinflammatory profile of M1 macrophages toward a more pro-
resolving, immunomodulatory phenotype.
717

718
Figure 5. RvD5 Impairs Mycobacterial control by M1 Macrophages. (A)
Initial uptake of *Mtb*. Unstimulated (M0) or IFNγ-activated (M1) macrophages,
treated or not with RvD5, were infected with *Mtb*-RFP. The percentage of RFP⁺
cells was quantified 4 h post-infection. Results are expressed relative to uptake
by unstimulated cells (n=4). Statistical analysis: Wilcoxon matched-pairs signed-
rank test comparing M1 macrophages with vs. without RvD5. **(B)** IL-1β levels.
Supernatants collected 24 h post-infection were analyzed by ELISA (n=9).
Statistical analysis: Wilcoxon matched-pairs signed-rank test comparing M1
macrophages with vs. without RvD5. **(C)** Mycobacterial growth. Colony-forming
units (CFU) were quantified 72 h post-infection in unstimulated or M1
macrophages with and without RvD5 (n=6). Data was analyzed by one-way
ANOVA followed by Holm-Sidak's multiple comparisons test. All quantitative
data are presented as scatter plots of individual donors, with means ± SEM
shown. **(D)** Representative confocal micrographs show M1 macrophages
infected with *Mtb*-RFP for 24 h in the presence or absence of RvD5. Nuclei are
stained with DAPI (blue), and the autophagy marker LC3 is stained with an
Alexa Fluor 488-conjugated antibody (green). The percentage of infected cells
(RFP⁺) exhibiting LC3 puncta per field was quantified. Statistical analysis:
Nested t-test from two independent experiments, analyzing 10 images per
condition.
739

HIF1α stabilization restores M1 macrophage features inhibited by the
GPR32-RvD5 axis

To investigate whether the inhibition of HIF1α mediates the inhibition of the
microbicidal activity of M1 macrophages induced by the -RvD5-GPR32, we
stabilized HIF1α expression by using Dimethyloxalylglycine (DMOG).
Remarkably, HIF1α stabilization in RvD5-treated M1 macrophages restored
their impaired lactate and IL-1β release (**Figure 6A-B**), as well as their
microbicidal activity, evidenced by reduced bacillary loads and a higher
localization of Mtb in compartments marked by the late endosomal marker,
LAMP2 (**Figure 6C-D**). Interestingly, RvD5-induced pro-resolution features,
such as enhanced efferocytic capacity, upregulated MerTK expression, and
increased IL-10 production, were diminished following DMOG treatment (**Figure
S10**). These findings highlight that the RvD5-GPR32-HIF1α axis suppresses
macrophage microbicidal functions and increases susceptibility to Mtb.
Significantly, these effects can be reversed through pharmacological
stabilization of HIF1α.

756
**Figure 6. Chemical HIF1α Stabilization Restores RvD5-mediated M1
Macrophage metabolism and function. (A)** Lactate release from unstimulated
(M0) or LPS/IFNγ-activated (M1) macrophages treated with RvD5 in the
presence or absence of the HIF-1α stabilizer DMOG for 24 h (n=11). **(B–D)**
IFNγ-activated macrophages (M1), treated or not with RvD5 and/or DMOG,
were infected with *Mtb*. **(B)** IL-1β levels measured by ELISA in supernatants
collected 24 h post-infection (n=8). **(C)** Mycobacterial growth assessed by
colony-forming units (CFU) quantified 72 h post-infection (n=6). For panels A–C,
data were analyzed by one-way ANOVA followed by Sidak's multiple
comparisons test. All quantitative data are presented as scatter plots of

individual donors, with means ± SEM. **(D)** Colocalization of *Mtb*-RFP with LAMP2. Representative confocal micrographs show macrophages infected with *Mtb*-RFP for 24 h. Nuclei are stained with DAPI (blue), and the late endosome marker LAMP2 is stained with an Alexa Fluor 633-conjugated antibody (cyan). The lower panels show magnified views of the boxed areas (scale bar = 10 µm). The right panel shows Manders colocalization coefficients for the fraction of *Mtb* signal overlapping with LAMP2 staining. Data was from a representative experiment (10 images per condition) and was analyzed by one-way ANOVA followed by Tukey's multiple comparisons test.

Discussion

Pleural TB presents unique challenges for healthcare providers, particularly concerning its treatment. Effective host defense against TB involves infiltration of peripheral blood cells into the pleural space, leading to accumulation of immune cells and soluble factors, including lipid mediators, in tuberculous pleural fluid. SPMs play a critical role in limiting inflammation and preventing collateral damage to healthy tissues. However, it is essential to carefully regulate the potential adverse effects of SPMs on the microbicidal functions of inflammatory macrophages to achieve a balanced approach that supports adequate pathogen clearance while minimizing excessive inflammation. While previous studies have highlighted the beneficial roles of SPMs in enhancing the microbicidal activities of macrophages and neutrophils[21,38–40], our study is the first to establish the involvement of the RvD5-GPR32-HIF1α axis in TB. We demonstrate that RvD5, present in pleural TB, activates GPR32, thereby inhibiting HIF1α -dependent microbicidal activity in macrophages.

Our study centers on two pivotal findings. First, we identified the RvD5-GPR32 axis as a key mediator of macrophage glycolysis suppression in pleural effusions. We demonstrated that RvD5 is not only present in pleural effusions but can also be produced by monocytes exposed to the acellular fraction of these effusions. Notably, CD14[+] cells in the pleural cavity constitute the primary mononuclear cell population responsible for local production of SPMs, such as resolvins. The accumulation of RvD5 within the pleural cavity correlates with reduced glycolytic capacity in CD14[+] cells compared to their circulating counterparts. This suggests that upon entering the pleural cavity, phagocytes undergo a metabolic shift characterized by attenuated glycolytic activity, influenced by the local microenvironment. In line with this, in vitro exposure of macrophages to RvD5 led to a similar reduction in glycolysis.

We show that pleural macrophages from TB patients undergoing antibiotic treatment exhibit a significant increase in glycolytic activity compared to those from untreated patients. This observation indicates the presence of active mechanisms within the pleural cavity that suppress long-term protective

immune responses, likely modulated by SPMs. Such findings offer a potential

explanation for the frequent relapses observed in untreated pleural TB patients,

suggesting that while glycolysis in macrophages is locally suppressed during

active disease, systemically, monocytes in TB patients maintain a dysregulated

glycolytic profile, as we recently reported[23]. This dichotomy supports the notion

of contrasting metabolic environments at local and systemic levels. Even though

using drugs that alter metabolism in all cells might have deleterious implications

for patients, identifying strategies to do so selectively in macrophages has

important clinical implications. We propose that upon entering the pleural cavity,

monocytes upregulate ALOX5 transcript levels, thereby promoting local

biosynthesis of SPMs. These SPMs, in turn, may modulate the metabolism of

inflammatory macrophages, shifting them toward reduced glycolysis. This

immunosuppressive metabolic state, however, appears reversible with TB

treatment, underscoring the dynamic interplay between local immune

modulation and systemic immune dysregulation in pleural TB.

Several studies have identified associations between ALOX5 polymorphisms

and increased susceptibility to TB[41,42] or the development of multisystem TB[43].

These studies primarily focus on the role of ALOX5 in synthesizing pro-

inflammatory lipid mediators, such as leukotrienes. In contrast, our findings

identify RvD5 as a novel and underappreciated player in this context.

Supporting our observations, ALOX5-deficient mice have been shown to exhibit

increased resistance to Mtb infection[44]. Interestingly, ALOX5 expression has

been implicated in both pro-inflammatory and immunosuppressive outcomes

across various pathological contexts. For example, in rheumatoid arthritis,

increased ALOX5 expression has been associated with harmful inflammation by

driving pyroptosis in CD4[+] T cells[45]. Conversely, in gliomas, ALOX5 promotes

an immunosuppressive phenotype by facilitating M2 polarization in

macrophages[46]. These diverse outcomes are likely the eclectic array of

products ALOX5 can generate, depending on substrate availability. Indeed,

ALOX5 can metabolize arachidonic acid, EPA, or DHA to produce oxylipins with

distinct biological functions, including 5-HETE, lipoxins, and resolvins[47]. This

substrate-driven variability highlights the complexity of ALOX5's involvement in

TB and other diseases. It suggests that its effects may change depending on

microenvironmental factors, such as lipid availability, cellular metabolic states,

and the balance between pro-inflammatory and pro-resolving mediators.

Understanding these dynamics is critical for fully elucidating the dual roles of

ALOX5 and for considering therapeutic strategies that target its metabolic

pathways in TB and beyond, and inhibition of ALOX5 might have undesired

effects due to its pleiotropic function. On the positive side, GPR32 belongs to

the most intensively studied drug target family, mainly because of its substantial

involvement in human pathophysiology and its pharmacological tractability[48].

Controlling excessive inflammation in the pleural space is crucial to prevent

tissue damage and maintain host-pathogen balance. While RvD5 has been

shown to impair the microbicidal functions of macrophages, other SPMs

prevalent in pleural TB, such as PDx and MaR1, do not exhibit these

detrimental effects[21]. This underscores an opportunity to selectively enhance

SPMs that preserve microbicidal activity while minimizing tissue damage,

thereby avoiding lipid mediators such as RvD5. For instance, in tuberculous

meningitis, elevated concentrations of SPMs in cerebrospinal fluid have been

associated with improved patient survival[49]. These results support the idea of

targeted interventions to harness the benefits of SPMs while mitigating the

risks.

The literature extensively explores the balance between "proinflammatory" and

"protective" lipid mediators, often manipulated through dietary supplementation

with omega-3 fatty acids (n-3 PUFAs, precursors of protective lipids) versus

omega-6 fatty acids (n-6 PUFAs, mostly precursors of proinflammatory lipids).

However, many studies lack a detailed examination of the mechanisms of

action for individual lipids. The outcomes of such supplementation studies have

been inconsistent, ranging from detrimental effects, such as the impairment of

murine macrophage responses to Mtb with omega-3 supplementation[50,51], to

beneficial outcomes where n-3 PUFAs enhance immune responses against

Mtb[50,52]. These conflicting findings underscore the need for further research to

delineate the specific roles and mechanisms of individual lipid mediators in the

context of TB.

Although some reports propose that certain SPMs, such as RvD1[53] or MaR1[54],
can enhance cellular respiration in neutrophils and macrophages, our findings
did not demonstrate that RvD5 increases absolute mitochondrial respiration in
macrophages stimulated with LPS and IFNγ. This discrepancy may be due to
the strong metabolic imprinting induced by LPS and IFNγ, which results in
decreased OXPHOS that RvD5 cannot counteract. Alternatively, the effect of
RvD5 may be limited to reducing glycolysis without activating compensatory
energetic mechanisms. These insights further support the complex, context-
dependent nature of SPM-mediated metabolic regulation, warranting more in-
depth studies to fully understand their roles and therapeutic potential in TB.

Our second key finding reveals that activation of GPR32 by RvD5 inhibits HIF1α
-dependent microbicidal activity in macrophages, shedding light on a critical
pathway that modulates immune responses in TB. Previously, we demonstrated
that lipids extracted from the acellular fraction of TB pleural effusions suppress
HIF1α expression, leading to reduced glycolytic and microbicidal phenotypes in
macrophages[15]. The present study goes beyond our previous results, as we
now show that RvD5 is explicitly a central lipid mediator driving this effect
through GPR32 binding. While GPR32 is the best-characterized receptor for
RvD5, emerging evidence suggests that RvD5 may also interact with GPR101,
which is also expressed in macrophages[55]. As GPR101 is constitutively active,
any role for RvD5 would likely involve enhancing or fine-tuning its basal
signaling rather than classical activation. The potential contribution of this
alternative receptor to the observed metabolic effects was not explored in this
study and remains an open avenue for future research.

Our results demonstrate that RvD5 impairs Mtb clearance, at least partially, by
limiting the induction of autophagy in infected macrophages. Importantly, this
suppression of autophagy could be reversed by stabilizing HIF1α, consistent
with previous findings that HIF1α promotes autophagy in cancer cells[56].
Additionally, it has been shown that Mtb can inhibit IFNγ-induced autophagy via
ALOX5-dependent induction of the Mir155-Mir31-WNT-SHH signaling pathway,
which is further supported by findings indicating that pharmacological
perturbation of ALOX5 leads to improved control of mycobacterial intracellular

loads[57]. However, the role of RvD5 specifically was not investigated in that
study. Interestingly, unlike RvD5, other resolvins such as RvD1 have been
reported to promote autophagy in various contexts[58,59]. These findings
emphasize the functional specificity of different SPMs and caution against
generalizing biological functions across the resolvin family.

In conclusion, our study provides valuable insights into the pathophysiology of
pleural TB by uncovering RvD5 as a critical lipid mediator that modulates
macrophage metabolism and function through GPR32. This work underlines the
importance of understanding the specific mechanisms by which lipid mediators
influence immune cell metabolism and TB severity. These results pave the way
for the development of targeted therapeutic strategies that aim to balance
immune regulation, preserve microbicidal activity, and mitigate the detrimental
effects of excessive inflammation in TB.

Acknowledgements

We thank the staff of the Regional Center of Hemotherapy of the Garrahan Hospital (Buenos Aires). We thank MetaToul (Toulouse metabolomics & fluxomics facilities, https://mth-metatoul.com/), which is part of the French National Infrastructure for Metabolomics and Fluxomics, MetaboHUB-ANR-11-INBS-0010 (doi.org/10.15454/VRJK-KY76). We also thank S. Fried and A. Lucas from the We-Met platform at I2MC, in Toulouse, France. We also thank Life Science Editors for editing services (www.lifescienceeditors.com).

This work was supported by the Argentinean National Agency of Promotion of Science and Technology (PICT-2019-01044 and PICT-2020-00501 to LB); the Argentinean National Council of Scientific and Technical Investigations (CONICET, PIP 11220200100299CO to LB); the ICGEB CRP/ARG23-02 Research Grant to LB; Fondation Bettencourt Schueller (EXPORE-TB grant to ON); MSDAVENIR (the FIGHT-TB grant to ON); and the French ANR JCJC-Epic-SCENITH ANR-20-CE14-0028 and CoPoC Inserm-transfert MAT-PI-17493-A-04 to RA.

Author contributions

Conceptualization: CV, GLV, and LB. Methodology: JB and LB. Investigation: JB, MMaio, LF, EB, NF, SM, JLMF, AM, FS, FF, TMG, MMatteo, EL, CV, GLV, and LB. Resources: AJE, XA, MSC, DP, MO, RA, RKN, ON, CV, GLV, and LB. Formal Analysis: JB, MML, EL, and LB. Writing: JB, MML, EL, ON, CV, GLV, and LB. Visualization: JB, FF, and LB. Supervision: EL, CV, GLV, and LB. Project Administration: LB. Funding acquisition: RA, ON, and LB. Corresponding author: LB is responsible for ownership and responsibility that are inherent to aspects of this study.

Declaration of interests

The authors declare no competing interests.

References

1. Koeken, V.A.C.M., Verrall, A.J., Netea, M.G., Hill, P.C., and van Crevel, R. (2019). Trained innate immunity and resistance to Mycobacterium tuberculosis infection. Clinical Microbiology and Infection *25*, 1468–1472. https://doi.org/10.1016/J.CMI.2019.02.015.

2. Biswas, S.K., and Mantovani, A. (2012). Orchestration of metabolism by macrophages. Preprint, https://doi.org/10.1016/j.cmet.2011.11.013 https://doi.org/10.1016/j.cmet.2011.11.013.

3. O'Neill, L.A.J., Kishton, R.J., and Rathmell, J. (2016). A guide to immunometabolism for immunologists. Nat Rev Immunol *16*, 553–565. https://doi.org/10.1038/nri.2016.70.

4. Che, N., Ma, Y., Ruan, H., Xu, L., Wang, X., Yang, X., and Liu, X. (2018). Integrated semi-targeted metabolomics analysis reveals distinct metabolic dysregulation in pleural effusion caused by tuberculosis and malignancy. Clinica Chimica Acta *477*, 81–88. https://doi.org/10.1016/J.CCA.2017.12.003.

5. Seibert, A.F., Haynes, J., Middleton, R., and Bass, J.B. (1991). Tuberculous pleural effusion; Twenty-year experience. Preprint, https://doi.org/10.1378/chest.99.4.883 https://doi.org/10.1378/chest.99.4.883.

6. Sharma, S.K., Mohan, A., and Kohli, M. (2021). Extrapulmonary tuberculosis. Expert Rev Respir Med *15*, 931–948. https://doi.org/10.1080/17476348.2021.1927718.

7. Macías, A., Sánchez-Montalvá, A., Salvador, F., Villar, A., Tórtola, T., Saborit, N., and Molina, I. (2019). Epidemiology and diagnosis of pleural tuberculosis in a low incidence country with high rate of immigrant population: A retrospective study. International Journal of Infectious Diseases *78*, 34–38. https://doi.org/10.1016/j.ijid.2018.10.005.

8. Kim, H.J., Lee, H.J., Kwon, S.Y., Yoon, H. Il, Chung, H.S., Lee, C.T., Han, S.K., Shim, Y.S., and Yim, J.J. (2006). The prevalence of pulmonary parenchymal tuberculosis in patients with tuberculous pleuritis. Chest *129*, 1253–1258. https://doi.org/10.1378/chest.129.5.1253.

9. Sharma, S.K., Mitra, D.K., Balamurugan, A., Pandey, R.M., and Mehra, N.K. (2002). Cytokine polarization in miliary and pleural tuberculosis. J Clin Immunol *22*, 345–352. https://doi.org/10.1023/A:1020604331886/METRICS.

10. Light, R.W. (2010). Update on tuberculous pleural effusion. Preprint, https://doi.org/10.1111/j.1440-1843.2010.01723.x https://doi.org/10.1111/j.1440-1843.2010.01723.x.

11. Genoula, M., Franco, J.L.M., Dupont, M., Kviatcovsky, D., Milillo, A., Schierloh, P., Moraña, E.J., Poggi, S., Palmero, D., Mata-Espinosa, D., et al. (2018). Formation of foamy macrophages by tuberculous pleural effusions is triggered by the interleukin-10/signal transducer and activator of transcription 3 axis through ACAT upregulation. Front Immunol. https://doi.org/10.3389/fimmu.2018.00459.

12. Lastrucci, C., Bénard, A., Balboa, L., Pingris, K., Souriant, S., Poincloux, R., Al Saati, T., Rasolofo, V., González-Montaner, P., Inwentarz, S., et al. (2015). Tuberculosis is associated with expansion of a motile, permissive and immunomodulatory CD16+ monocyte population via the IL-10/STAT3 axis. Cell Res. https://doi.org/10.1038/cr.2015.123.

13. Souriant, S., Balboa, L., Dupont, M., Pingris, K., Kviatcovsky, D., Cougoule, C., Lastrucci, C., Bah, A., Gasser, R., Poincloux, R., et al. (2019). Tuberculosis Exacerbates HIV-1 Infection through IL-10/STAT3-Dependent Tunneling Nanotube Formation in Macrophages. Cell Rep. https://doi.org/10.1016/j.celrep.2019.02.091.

14. Genoula, M., Marín Franco, J.L., Maio, M., Dolotowicz, B., Ferreyra, M., Milillo, M.A., Mascarau, R., Moraña, E.J., Palmero, D., Matteo, M., et al. (2020). Fatty acid

oxidation of alternatively activated macrophages prevents foam cell formation, but Mycobacterium tuberculosis counteracts this process via HIF-1α activation. PLoS Pathog *16*, e1008929. https://doi.org/10.1371/journal.ppat.1008929.

15. Marín Franco, J.L., Genoula, M., Corral, D., Duette, G., Ferreyra, M., Maio, M., Dolotowicz, M.B., Aparicio-Trejo, O.E., Patiño-Martínez, E., Charton, A., et al. (2020). Host-Derived Lipids from Tuberculous Pleurisy Impair Macrophage Microbicidal-Associated Metabolic Activity. Cell Rep. https://doi.org/10.1016/j.celrep.2020.108547.

16. Serhan, C.N. (2014). Pro-resolving lipid mediators are leads for resolution physiology. Nature *510*, 92–101. https://doi.org/10.1038/NATURE13479.

17. Centanni, D., Henricks, P.A.J., and Engels, F. (2023). The therapeutic potential of resolvins in pulmonary diseases. Eur J Pharmacol *958*, 176047. https://doi.org/10.1016/J.EJPHAR.2023.176047.

18. Dalli, J., Winkler, J.W., Colas, R.A., Arnardottir, H., Cheng, C.Y.C., Chiang, N., Petasis, N.A., and Serhan, C.N. (2013). Resolvin D3 and Aspirin-Triggered Resolvin D3 Are Potent Immunoresolvents. Chem Biol *20*, 188. https://doi.org/10.1016/J.CHEMBIOL.2012.11.010.

19. Krishnamoorthy, S., Recchiuti, A., Chiang, N., Yacoubian, S., Lee, C.H., Yang, R., Petasis, N.A., and Serhan, C.N. (2010). Resolvin D1 binds human phagocytes with evidence for proresolving receptors. Proc Natl Acad Sci U S A *107*, 1660–1665. https://doi.org/10.1073/PNAS.0907342107/SUPPL_FILE/PNAS.200907342SI.PDF.

20. Chiang, N., Fredman, G., Bäckhed, F., Oh, S.F., Vickery, T., Schmidt, B.A., and Serhan, C.N. (2012). Infection regulates pro-resolving mediators that lower antibiotic requirements. Nature 2012 484:7395 *484*, 524–528. https://doi.org/10.1038/nature11042.

21. Ruiz, A., Sarabia, C., Torres, M., and Juárez, E. (2019). Resolvin D1 (RvD1) and maresin 1 (Mar1) contribute to human macrophage control of M. tuberculosis infection while resolving inflammation. Int Immunopharmacol *74*. https://doi.org/10.1016/J.INTIMP.2019.105694.

22. Le Faouder, P., Baillif, V., Spreadbury, I., Motta, J.P., Rousset, P., Chêne, G., Guigné, C., Tercé, F., Vanner, S., Vergnolle, N., et al. (2013). LC-MS/MS method for rapid and concomitant quantification of pro-inflammatory and pro-resolving polyunsaturated fatty acid metabolites. J Chromatogr B Analyt Technol Biomed Life Sci. https://doi.org/10.1016/j.jchromb.2013.06.014.

23. Maio, M., Barros, J., Joly, M., Vahlas, Z., Franco, J.L.M., Genoula, M., Monard, S., Vecchione, M.B., Fuentes, F., Polo, V.G., et al. (2024). Elevated glycolytic metabolism of monocytes limits the generation of HIF-1α-driven migratory dendritic cells in tuberculosis. Elife *12*. https://doi.org/10.7554/ELIFE.89319.3.

24. Argüello, R.J., Combes, A.J., Char, R., Gigan, J.P., Baaziz, A.I., Bousiquot, E., Camosseto, V., Samad, B., Tsui, J., Yan, P., et al. (2020). SCENITH: A Flow Cytometry-Based Method to Functionally Profile Energy Metabolism with Single-Cell Resolution. Cell Metab. https://doi.org/10.1016/j.cmet.2020.11.007.

25. Cai, Y., Wang, Y., Shi, C., Dai, Y., Li, F., Xu, Y., Zhang, P., Kong, F., Deng, G., Wen, Z., et al. (2022). Single-cell immune profiling reveals functional diversity of T cells in tuberculous pleural effusion. Journal of Experimental Medicine *219*. https://doi.org/10.1084/JEM.20211777/212978.

26. Schebb, N.H., Kühn, H., Kahnt, A.S., Rund, K.M., O'Donnell, V.B., Flamand, N., Peters-Golden, M., Jakobsson, P.J., Weylandt, K.H., Rohwer, N., et al. (2022). Formation, Signaling and Occurrence of Specialized Pro-Resolving Lipid Mediators— What is the Evidence so far? Front Pharmacol *13*. https://doi.org/10.3389/FPHAR.2022.838782/FULL.

27.   Lee, J., Lim, J.K., Yoo, S.S., Lee, S.Y., Cha, S.I., Park, J.Y., and Kim, C.H. (2016). Different characteristics of tuberculous pleural effusion according to pleural fluid cellular predominance and loculation. J Thorac Dis 8, 1935–1942. https://doi.org/10.21037/JTD.2016.06.54.

28.   Schmid, M., Gemperle, C., Rimann, N., and Hersberger, M. (2016). Resolvin D1 Polarizes Primary Human Macrophages toward a Proresolution Phenotype through GPR32. The Journal of Immunology 196, 3429–3437. https://doi.org/10.4049/JIMMUNOL.1501701.

29.   Palsson-Mcdermott, E.M., Curtis, A.M., Goel, G., Lauterbach, M.A.R., Sheedy, F.J., Gleeson, L.E., Van Den Bosch, M.W.M., Quinn, S.R., Domingo-Fernandez, R., Johnson, D.G.W., et al. (2015). Pyruvate kinase M2 regulates hif-1α activity and il-1β induction and is a critical determinant of the warburg effect in LPS-activated macrophages. Cell Metab. https://doi.org/10.1016/j.cmet.2014.12.005.

30.   Park, J., Langmead, C.J., and Riddy, D.M. (2020). New Advances in Targeting the Resolution of Inflammation: Implications for Specialized Pro-Resolving Mediator GPCR Drug Discovery. ACS Pharmacol Transl Sci 3, 88. https://doi.org/10.1021/ACSPTSCI.9B00075.

31.   Viola, A., Munari, F., Sánchez-Rodríguez, R., Scolaro, T., and Castegna, A. (2019). The metabolic signature of macrophage responses. Front Immunol 10, 466337. https://doi.org/10.3389/FIMMU.2019.01462/BIBTEX.

32.   Scott, R.S., McMahon, E.J., Pop, S.M., Reap, E.A., Caricchio, R., Cohen, P.L., Shelton Earp, H., and Matsushima, G.K. (2001). Phagocytosis and clearance of apoptotic cells is mediated by MER. Nature 411, 207–211. https://doi.org/10.1038/35075603.

33.   Devitt, A., Moffatt, O.D., Raykundalia, C., Capra, J.D., Simmons, D.L., and Gregory, C.D. (1998). Human CD14 mediates recognition and phagocytosis of apoptotic cells. Nature 1998 392:6675 392, 505–509. https://doi.org/10.1038/33169.

34.   Savill, J., Hogg, N., Ren, Y., and Haslett, C. (1992). Thrombospondin cooperates with CD36 and the vitronectin receptor in macrophage recognition of neutrophils undergoing apoptosis. J Clin Invest 90, 1513–1522. https://doi.org/10.1172/JCI116019.

35.   Golovkine, G.R., Roberts, A.W., Morrison, H.M., Rivera-Lugo, R., McCall, R.M., Nilsson, H., Garelis, N.E., Repasy, T., Cronce, M., Budzik, J., et al. (2023). Autophagy restricts Mycobacterium tuberculosis during acute infection in mice. Nature Microbiology 2023 8:5 8, 819–832. https://doi.org/10.1038/s41564-023-01354-6.

36.   Dang, A.T., Teles, R.M.B., Liu, P.T., Choi, A., Legaspi, A., Sarno, E.N., Ochoa, M.T., Parvatiyar, K., Cheng, G., Gilliet, M., et al. (2019). Autophagy links antimicrobial activity with antigen presentation in Langerhans cells. JCI Insight 4. https://doi.org/10.1172/JCI.INSIGHT.126955.

37.   Jagannath, C., Lindsey, D.R., Dhandayuthapani, S., Xu, Y., Hunter, Jr., R.L., and Eissa, N.T. (2009). Autophagy enhances the efficacy of BCG vaccine by increasing peptide presentation in mouse dendritic cells. Nat Med 15, 267–277.

38.   Croasdell, A., Lacy, S.H., Thatcher, T.H., Sime, P.J., and Phipps, R.P. (2016). Resolvin D1 Dampens Pulmonary Inflammation and Promotes Clearance of Nontypeable Haemophilus influenzae. J Immunol 196, 2742–2752. https://doi.org/10.4049/JIMMUNOL.1502331.

39.   Wu, B., Capilato, J., Pham, M.P., Walker, J., Spur, B., Rodriguez, A., Perez, L.J., and Yin, K. (2016). Lipoxin A4 augments host defense in sepsis and reduces Pseudomonas aeruginosa virulence through quorum sensing inhibition. FASEB J 30, 2400–2410. https://doi.org/10.1096/FJ.201500029R.

40. Wu, B., Walker, J., Spur, B., Rodriguez, A., and Yin, K. (2015). Effects of Lipoxin A4 on antimicrobial actions of neutrophils in sepsis. Prostaglandins Leukot Essent Fatty Acids *94*, 55–64. https://doi.org/10.1016/J.PLEFA.2014.11.005.

41. Shen, C., Wu, X.R., Wang, B. Bin, Sun, L., Jiao, W.W., Wang, J., Feng, W.X., Xiao, J., Miao, Q., Liu, F., et al. (2013). ALOX5 is associated with tuberculosis in a subset of the pediatric population of North China. Genet Test Mol Biomarkers *17*, 284–288. https://doi.org/10.1089/GTMB.2012.0426.

42. Herb, F., Thye, T., Niemann, S., Browne, E.N.L., Chinbuah, M.A., Gyapong, J., Osei, I., Owusu-Dabo, E., Werz, O., Rüsch-Gerdes, S., et al. (2008). ALOX5 variants associated with susceptibility to human pulmonary tuberculosis. Hum Mol Genet *17*, 1052–1060. https://doi.org/10.1093/HMG/DDM378.

43. Zhang, J., Wang, M. gui, and He, J.Q. (2023). Association between a single nucleotide polymorphism of the ALOX5 gene and susceptibility to multisystem tuberculosis in a Chinese Han population. Microb Pathog *183*. https://doi.org/10.1016/J.MICPATH.2023.106289.

44. Bafica, A., Scanga, C.A., Serhan, C., Machado, F., White, S., Sher, A., and Aliberti, J. (2005). Host control of Mycobacterium tuberculosis is regulated by 5-lipoxygenase-dependent lipoxin production. Journal of Clinical Investigation. https://doi.org/10.1172/JCI23949.

45. Cai, H., Zhang, J., Xu, H., Sun, W., Wu, W., Dong, C., Zhou, P., Xue, C., Nan, Y., Ni, Y., et al. (2024). ALOX5 drives the pyroptosis of CD4+ T cells and tissue inflammation in rheumatoid arthritis. Sci Signal *17*. https://doi.org/10.1126/SCISIGNAL.ADH1178/SUPPL_FILE/SCISIGNAL.ADH1178_MDAR_REPRODUCIBILITY_CHECKLIST.PDF.

46. Chen, T., Liu, J., Wang, C., Wang, Z., Zhou, J., Lin, J., Mao, J., Pan, T., Wang, J., Xu, H., et al. (2024). ALOX5 contributes to glioma progression by promoting 5-HETE-mediated immunosuppressive M2 polarization and PD-L1 expression of glioma-associated microglia/macrophages. J Immunother Cancer *12*, e009492. https://doi.org/10.1136/JITC-2024-009492.

47. Kahnt, A.S., Häfner, A.K., and Steinhilber, D. (2024). The role of human 5-Lipoxygenase (5-LO) in carcinogenesis - a question of canonical and non-canonical functions. Oncogene 2024 43:18 *43*, 1319–1327. https://doi.org/10.1038/s41388-024-03016-1.

48. Hauser, A.S., Attwood, M.M., Rask-Andersen, M., Schiöth, H.B., and Gloriam, D.E. (2017). Trends in GPCR drug discovery: new agents, targets and indications. Nat Rev Drug Discov *16*, 829. https://doi.org/10.1038/NRD.2017.178.

49. Colas, R.A., Nhat, L.T.H., Thuong, N.T.T., Gómez, E.A., Ly, L., Thanh, H.H., Mai, N.T.H., Phu, N.H., Thwaites, G.E., and Dalli, J. (2019). Proresolving mediator profiles in cerebrospinal fluid are linked with disease severity and outcome in adults with tuberculous meningitis. FASEB J *33*, 13028–13039. https://doi.org/10.1096/FJ.201901590R.

50. Jordao, L., Lengeling, A., Bordat, Y., Boudou, F., Gicquel, B., Neyrolles, O., Becker, P.D., Guzman, C.A., Griffiths, G., and Anes, E. (2008). Effects of omega-3 and -6 fatty acids on Mycobacterium tuberculosis in macrophages and in mice. Microbes Infect *10*, 1379–1386. https://doi.org/10.1016/J.MICINF.2008.08.004.

51. Bonilla, D.L., Ly, L.H., Fan, Y.Y., Chapkin, R.S., and Mcmurray, D.N. (2010). Incorporation of a Dietary Omega 3 Fatty Acid Impairs Murine Macrophage Responses to Mycobacterium tuberculosis. PLoS One *5*, e10878. https://doi.org/10.1371/JOURNAL.PONE.0010878.

52. Nienaber, A., Baumgartner, J., Dolman, R.C., Ozturk, M., Zandberg, L., Hayford, F.E.A., Brombacher, F., Blaauw, R., Parihar, S.P., Smuts, C.M., et al. (2020). Omega-3 Fatty Acid and Iron Supplementation Alone, but Not in Combination, Lower Inflammation and Anemia of Infection in Mycobacterium tuberculosis-Infected Mice. Nutrients *12*, 1–22. https://doi.org/10.3390/NU12092897.

53. Li, L., Cheng, S.Q., Sun, Y.Q., Yu, J.B., Huang, X.X., Dong, Y.F., Ji, J., Zhang, X.Y., Hu, G., and Sun, X.L. (2023). Resolvin D1 reprograms energy metabolism to promote microglia to phagocytize neutrophils after ischemic stroke. Cell Rep *42*. https://doi.org/10.1016/J.CELREP.2023.112617.

54. Calderin, E.P., McNally, L., Hill, B.G., and Hellmann, J. (2021). Specialized pro-resolving lipid mediators stimulate mitochondrial metabolism in macrophages. The Journal of Immunology *206*, 97.09-97.09. https://doi.org/10.4049/JIMMUNOL.206.SUPP.97.09.

55. Chiang, N., and Serha, C.N. (2020). Specialized pro-resolving mediator network: an update on production and actions. Essays Biochem *64*, 443–462. https://doi.org/10.1042/EBC20200018.

56. Wang, P., Long, M., Zhang, S., Cheng, Z., Zhao, X., He, F., Liu, H., and Ming, L. (2017). Hypoxia inducible factor-1α regulates autophagy via the p27-E2F1 signaling pathway. Mol Med Rep *16*, 2107. https://doi.org/10.3892/MMR.2017.6794.

57. Holla, S., Kurowska-Stolarska, M., Bayry, J., and Balaji, K.N. (2013). Selective inhibition of IFNG-induced autophagy by Mir155- and Mir31-responsive WNT5A and SHH signaling. Autophagy *10*, 311. https://doi.org/10.4161/AUTO.27225.

58. Prieto, P., Rosales-Mendoza, C.E., Terron, V., Toledano, V., Cuadrado, A., López-Collazo, E., Bannenberg, G., Martín-Sanz, P., Fernández-Velasco, M., and Boscá, L. (2015). Activation of autophagy in macrophages by pro-resolving lipid mediators. Autophagy *11*, 1729–1744. https://doi.org/10.1080/15548627.2015.1078958.

59. Xiong, W., Wang, H., Zhang, H., Xing, Y., Gao, W., Chen, L., Chen, L., and Dai, Z. (2023). Resolvin D1 attenuates depressive-like behavior in LPS-challenged mice by promoting microglial autophagy. Inflammopharmacology *31*, 2061–2075. https://doi.org/10.1007/S10787-023-01234-9.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6