Altered Stress and Fear Responses in the VPA Rat Model of Autism: Behavioral Dissociation Across Tactile, Nociceptive, and Social Contexts

Debora Hashiguchi[1*], Ana Luiza Dias[2], Rodrigo Neves Romcy-Pereira[1*]

[1]Laboratory of Neural Circuit Plasticity and [2]Laboratory of Computational Neuroscience. Brain Institute, Federal University of Rio Grande do Norte, UFRN, Natal, RN, 59076-550, Brazil.

Corresponding author

E-mail: hashiguchi.debora@gmail.com, rnrpereira@neuro.ufrn.br

Author's contribution

Author DH: Conceptualization, data curation, formal analysis, investigation, methodology, project administration, supervision; validation, visualization, writing – original draft preparation, and writing – review and editing.

Author ALD: Formal analysis, and writing – review and editing.

Author RNRP: Conceptualization, formal analysis, funding acquisition, methodology, resources, writing – review and editing.

Abstract

Sensitivity to environmental stimuli is a fundamental aspect of human behavior, and its dysregulation is associated with stress-related and anxiety disorders. In autism spectrum disorder (ASD), altered sensory processing may contribute to increased vulnerability to such conditions. To better understand this relationship, we evaluated autonomic and behavioral responses to tactile, nociceptive, and social stressors in juvenile Wistar rats prenatally exposed to valproic acid (VPA), a widely used animal model of ASD. VPA-treated and saline-treated control (CTL) rats underwent a battery of stress-related behavioral tests. Defecation, freezing and vocalization behaviors were analyzed in response to handling, electro-tactile stimulation, nociceptive foot shock (fear conditioning) and social stress stimuli (emotional contagion). VPA-treated rats maintained increased defecation during handling and a higher prevalence of defecation during electro-tactile stimulation, without corresponding changes in freezing. During fear conditioning, these animals showed delayed onset and heightened freezing responses. Furthermore, unlike CTL rats, VPA-treated rats lacked correlations among freezing, defecation, and vocalization. In the emotional contagion paradigm, observation of shock in the partner increased the prevalence of animals freezing and reduced the prevalence of animals vocalizing in both VPA and CTL groups (oV+ and oC+) compared to their respective controls (oV- and oC-). However, these effects were more sustained in oV+ rats, which also exhibited prolonged freezing behavior and an earlier reduction in vocalization rate. These findings indicate that VPA-treated rats display heightened stress reactivity, habituation deficit and disrupted coordination of fear responses, supporting the VPA model as a relevant tool for investigating the neurobiological basis of stress vulnerability and social dysfunction in ASD.

Keywords: Autism, VPA animal model, stress, tactile sensitivity, nociceptive sensitivity, emotional sensitivity

Introduction

The ability to respond to environmental changes is a fundamental characteristic of human existence. Excessive worry, fear, and heightened sensitivity to threats characterize stress and anxiety disorders. Evidence indicates that individuals with ASD are especially susceptible to experiencing anxiety (1–7). Studies suggest that innocuous stimuli are more frequently appraised as stressful by affected individuals compared to normotypical individuals, which may result in behaviors similar to the characteristics inherent to ASD, such as difficulties in coping with changes, abnormal responsiveness to sensory stimuli and social demands, thus increasing the vulnerability of these individuals to stress disorders (1,8–12).

In humans and animals, the primary arousal system in the human body is the autonomic nervous system (ANS). The ANS responds rapidly to stress by activating and inhibiting its sympathetic and parasympathetic branches. These responses manifest in a variety of physiological changes, including increased heart rate, blood pressure, respiratory rate, pupil dilation, and colonic motility (13–17). These responses collectively prepare the body to confront or avoid the stressor, commonly known as the fight-or-flight response. Behavioral responses to these physiological changes include increased alertness and focus, enhanced sensory perception, and suppression of nonessential functions. Consequently, studies use changes in cardiac activity, sweating, skin temperature, and colonic motility as indicators to assess ANS functionality (18–22). Increased fecal pellet release, driven by increased colonic motility, and freezing behavior, which involves the integration of central nervous system, ANS, and sensorimotor functions, have been observed in stressed rats (23–26). However, the behavioral response

elicited varies depending on the level of disturbance in ANS homeostasis. Furthermore, stress-induced emission of ultrasonic vocalizations (USVs) within the frequency range of 20–35 kHz, also known as 22 kHz calls, has been widely correlated with the level of stress experienced by the animal (27,28). Calls lasting less than 100 ms are typically associated with acute distress, while longer calls, which can extend up to 3000 ms, are frequently observed under sustained stressful conditions (29–33). All of these behaviors are recognized as stress signals and considered vital for social communication, particularly in signaling danger within a group.

The valproic acid (VPA) rat model is extensively employed as a preclinical model of autism spectrum disorder (ASD) due to its recapitulation of key neural and behavioral alterations characteristic of the condition (34–37). Despite its widespread application, the model's validity for investigating stress susceptibility remains underexplored. This study aimed to assess the feasibility of employing the VPA rat model for translational research on stress responses to tactile, nociceptive, and adverse social stimuli. To achieve this, four experimental paradigms were employed: (1) a manipulation test and (2) an electro-tactile sensitivity test to evaluate stress responses to tactile stimuli; (3) a fear conditioning paradigm to assess stress responses to nociceptive stimuli; and (4) an emotional contagion paradigm to investigate stress responses to negative social stimuli.

Material and methods

Animals

Experimental and control animals were obtained by breeding male and female Wistar rats from the Brain Institute Animal Facility (BISIC) at the Federal University of Rio Grande do Norte (UFRN). Breeding pairs were >60 days old and weighed between 250–300 g at the time of mating. On gestation day 12.5, 14 pregnant females were injected with either sodium valproate (n = 8; 250 mg/mL saline solution, 500mg/Kg, i.p., VPA; Sigma-Aldrich

P-4543) or saline (n = 6; 0.15M NaCl) solutions. A group of pregnant females (n = 9) was

left undisturbed for the generation of naive animals to be used in the emotional contagion

paradigm (as demonstrators). In total, we used 23 litters generated by mating 17 females

with 14 males. Six females were mated twice and contributed with two litters (first litter

control and second litter VPA, n = 1; first litter control and second litter naive, n = 1; first

litter naive and second litter VPA, n = 3; or first litter VPA and second litter naive, n = 1).

After birth, pups were left in their homecage with the dam until weaning at P21.

Newborn rats used in the tactile sensitivity tests were transferred at P21 to new clean

cages (3-6 littermates/cage) without sexual segregation. A total of 58 animals were

included in these experiments: 28 VPA-treated rats (15 females and 13 males) and 30

CTL rats (15 females and 15 males). Behavioral tests were conducted between P33 and

P37. Newborn animals used in the nociceptive and negative social experiments were

transferred at P21 to new clean cages in pairs. In each cage, a VPA-treated or CTL rat

was housed with a naive rat of the same age and sex. A total of 122 animals were

included in these experiments: 31 VPA-treated rats (15 females and 16 males), 30 CTL

rats (15 females and 15 males) and 61 naive rats (30 females and 31 males). The same

cohorts of VPA and CTL animals were used for the nociceptive and negative social

experiments. Behavioral tests were performed on P38 and P42.

All VPA-treated rats showed a characteristic tail kink due to the VPA embryological effect.

All experimental procedures used in this study were approved by the Ethics Committee on

the Use of Animals (CEUA-UFRN, protocol nº 017/2020). Throughout the experiments, all

animals were kept in polypropylene boxes (55 x 46 x 30 cm) in a conditioned environment,

with temperature set to 24°C, 70% humidity, a 12h/12h light-dark cycle (lights-on: 7am;

lights-off: 7pm), with food and water freely available.

Behavioral tests

Touch sensitivity test

To investigate touch sensitivity, we used gentle handling as a stimulus. Over a period of four days, rats aged P33 to P36 were transferred from the vivarium to an experimental room within their housing boxes. Once in the experimental room, each rat was individually removed from its cage and handled by the experimenter. Handling involved four sessions/day of one minute each, during which the animals received gentle hand strokes on their fur while held in the experimenter's hand. Between sessions, animals were transferred back to their home cage where they stayed for five minutes before the next session. Handling sessions were carried out at the end of the lights-on phase, between 4pm and 7pm. Once the handling was completed, the animals were returned to the vivarium.

Electro-tactile sensitivity test

To investigate electro-tactile sensitivity of the glabrous skin of the paws, we used footshocks as stimuli. The experimental apparatus consisted of a shock chamber measuring 32 cm in length, 25 cm in width, and 30 cm in height, featuring a metallic grid floor for the delivery of the electrical foot-shock stimuli. The chamber had a translucent roof with small openings on the sides, two metallic walls, and two translucent walls, through which video recordings were obtained. Video monitoring captured the entire area of the apparatus, through two high-definition webcams (Logitech C920). Foot-shock stimuli were applied through the grid by a programmable digital stimulus generator (STG 4002 Multi Channel Systems) controlled by software (MC stimulus 4002). Each stimulus consisted of a rectangular pulse of 1 sec duration and varying amplitude (50-550 µA; 50 µA steps). A digital ultrasonic microphone (Ultramic 250K, Dodotronics) was also positioned through the opening at the roof of the chamber to record vocalizations.

Microphone and video camera recordings were synchronized to the foot-shocks by an Arduino device programmed to activate a light flash and a sound buzz simultaneous to the electrical stimulus.

All animals were tested one day after the last handling session, which was used as habituation to the experimenter and to the experimental room. The experiment took place at the end of the lights-on phase, between 4pm and 7pm and was conducted under dim light. For the experiment, the rats were placed in the middle of the apparatus facing one of the metal walls. After a period of 90 sec of free exploration, single electrical stimuli of 1 sec were applied with increasing intensity in 50 µA steps with random inter-stimulus intervals between 30 and 90 sec. The test was considered completed when the animal displayed jump or flee, for two consecutive stimuli.

Nociceptive stress response

To investigate the nociceptive stress response, we used a contextual fear conditioning paradigm with repeated unpredictable footshocks. The apparatus consisted of an operant conditioning chamber (32L x 25W x 30H cm each) with a metal grid floor, a translucent roof with small openings on the sides, surrounded by two metallic and two translucent walls. Electrical foot shock stimuli were applied by a programmable digital stimulus generator (STG 4002 MultiChannel Systems) controlled by software (MC stimulus 4002). Video monitoring captured the entire area of the apparatus, through a high-definition webcam (Logitech C920) positioned above the apparatus. A digital ultrasonic microphone (Ultramicrophone 250K, Dodotronic; 16 bits ADC; 250kHz sampling rate) was positioned through the opening at the roof of the chamber to record vocalizations at approximately 30 cm from the chamber floor. Microphone and video camera recordings were synchronized to the foot-shocks by an Arduino device programmed to activate a light flash and a sound buzz simultaneous to the electrical stimulus. Videos recordings in .mp4

format and audio recordings in .wav format were stored on a hard disk for posterior

analysis.

The experiment took place at the end of the lights-on phase, between 2pm and 5pm

and was carried out under dim light. At P38, rats were placed in the middle of the

apparatus facing one of the metal walls. Following a 4-min period of free exploration that

served as baseline (block B0), five electrical shock stimuli of 1-sec duration and random

inter-stimulus interval 0-180 sec were applied (blocks B1-B5). The experiment finished 4

minutes after the last shock was applied. We used a shock intensity (750uA) that

corresponded to two times the nociceptive threshold defined during the electro-tactile

sensitivity test. At the end of the experiment, each rat was transferred to a new clean

(neutral) cage for one hour before returning to its home cage.

Social stress response

In order to investigate the animal's response to a cagemate stress demonstration,

we used the emotional contagion paradigm, in which an experimental animal (observer,

OBS) observing a conspecific (demonstrator, DEM) undergo a series of aversive foot

shocks. The apparatus used consisted of two compartments: a conditioning chamber

(shock chamber) adjacent to a neutral chamber (32L x 25W x 30H cm) was separated by a

translucent plexiglass wall with two rows (4 and 6 cm above the floor) of 8 open roles each

(1cm diameter). An ultrasonic microphone was positioned in the shock chamber through a

roof aperture in the chamber (30 cm from the floor) to record vocalizations. Control

software and equipment synchronization were the same as previously described. Audio

(.wav files) and video recordings (.mp4 files) were save and stored in hard disk for

posterior analysis of freezing and vocal behaviors. Habituation and testing were conducted

during the lights-off phase, between 7:00 p.m. and 3:00 a.m., under dim light.

Over the following two days of fear conditioning, OBS animals were habituated to

the neutral chamber for 10 minutes each day (P39-P40). They were placed in the center of

the neutral box, facing the shock chamber, and allowed to freely explore the apparatus

and then, returned to their home cages. At P41, both OBS and DEM were brought to the

experimental apparatus. The DEM animal was placed in the center of the shock chamber,

facing one of the side walls, while the OBS was placed in the center of the neutral

chamber, facing one of the side walls. Following a 4-min period of free exploration (B0),

which served as a baseline, five electrical shock stimuli of 1-sec duration, 750 µA intensity,

delivered at random inter-stimulus interval 0-360 sec, were applied to the DEM animal

(blocks B1-B5). In order to maximize the OBS-to-DEM attention, all shock stimuli were

delivered to the DEM only when the OBS animal faced the DEM. Following the last

stimulus, both animals remained for an additional 4 min in the apparatus. Throughout the

experiment, the OBS animal was able to make visual contact with the DEM through a clear

plexiglass division. All experiments were conducted under dim light conditions.

In order to assign the identity of the call emitter while using one microphone in the

DEM chamber, we designed a pilot experiment (Fig S1). Vocalizations above -76 dB were

considered as originating from animals in the same chamber where the microphone was

placed (DEM chamber). Calls with power below -85 dB were considered predominantly

originating from animals in the adjacent chamber (OBS chamber). In summary, high

intensity vocalizations were considered DEM calls, whereas low intensity vocalizations

were considered OBS calls.

Figure S1: USV power. (A) Maternal separation (MS) experimental design at P08 and

graphic with USVs mean power by animal. Dotted line represents the lower power of USV

emitted in the microphone chamber (– 85.84 kHz). Dashed line represents the higher

power of USV emitted in the opposite chamber (– 84.03 kHz). (B) Maternal separation

(MS) experimental design at P14 and graphic with USVs mean power by animal. Dotted

line represents the lower power of USV emitted in the microphone chamber (– 90.08 kHz).

Dashed line represents the higher power of USV emitted in the opposite chamber (– 83.60 kHz). (C) Fear conditioning (FC) experimental design at P35-38 and graphic with USVs mean power by animal. Dotted line represents the lower power of USV emitted in the microphone chamber (– 92.12 kHz). Dashed line represents the higher power of USV emitted in the opposite chamber (– 76.54 kHz). [#] between-groups significant difference.

Behavioral analysis

Quantification of fecal pellets was recorded at the end of each handling session and immediately after all other experiments. To maintain objective quantification of variables, video and audio files (.mp4 and .wav) were randomly coded and analyzed by a skilled experimenter.

For the determination of the tactile threshold, the following behaviors were considered: grid investigation and paw retraction. The tactile threshold was defined as the lowest current at which the animal exhibited any of the previous behaviors. For the determination of the nociceptive threshold, the following behaviors were considered: jumping and fleeing. The nociceptive threshold was defined as the lowest current at which the animal displayed any of these behaviors. A freezing episode was identified as the behavioral state where the animal remained completely still without any intentional movement for a duration exceeding 3 seconds. It was quantified as the relative freezing time throughout the experiment. The analyzes of nociceptive and social negative experiments compared measurements during the baseline (B0) with either their combination during all shock blocks (B1-5) or individually with their values per block (B1 to B5). Following B0 (baseline, 4 min before first shock), each subsequent block (B1-B4) extended from 2s prior to a shock to 2s before the next shock. The last block (B5) extended from 2s before the fifth shock to 4 min after this shock. All video recordings were blindly analyzed by behavioral logging using BORIS software v.7.13.8.

The vocal behavior during the experiments was analyzed by quantifying the number of USVs produced and their acoustic properties. The analysis consisted of four steps: (1) Call detection, (2) Curation, (3) Segmentation and (4) Feature extraction. They were performed using software routines developed in our laboratory (Bessa, Rafael dos Santos de & Romcy-Pereira, 2023) and a custom-modified version of DeepSqueak software v2.6.2. Automatic USV detection was performed by computing a time-dependent entropy score of the power distribution across frequencies in the audio spectrogram with time resolution = 2 ms; frequency resolution = 65 Hz; frequency range, 20-100 kHz using the Chronux software package (Bokil et al., 2010). USV curation and segmentation consisted in a software-assisted refinement of USV time and frequency boundary boxes, which were performed by a trained investigator. The software performed a pre-identification of the USVs, which could be confirmed or rejected by the investigator. The investigator could include USVs that were not identified by the software or exclude regions of noise incorrectly identified as USVs. The criteria used to consider a putative USV were as follows: minimum frequency of 20 kHz, maximum frequency of 125 kHz, minimum duration of 5 ms, and a maximum interval between consecutive signals of 30 ms. For the purposes of this study, statistical analysis focused on the number of calls produced, call duration, and fundamental frequency of long 22 kHz calls (< 35-kHz; > 200 ms).

Statistics

We applied parametric tests when data distributions conformed to normality criteria, otherwise non-parametric tests were used. For proportions and percentages, data were z-transformed before applying a parametric test. Fisher's exact test or Chi-square were employed to compare data on contingency tables from two independent samples. Mann-Whitney and Student-t tests were employed to compare two independent distributed samples. Paired Student-t test, Wilcoxon signed-rank test employed to compare single

group paired samples. Friedman followed by post hoc Dunn's tests and repeated

measures (RM) one-way ANOVA with subsequent Holm-Sidak post-hoc test was

employed in single group repeated measures. Kruskal-Wallis was employed to compare

multiple groups independent data. Two-way RM ANOVA followed by post-hoc Holm-Sidak

test was employed to compare to between-groups with repeated measures. Two-way RM

ANOVA with subsequent Dunnets post-hoc test was employed to compare to within-group

repeated measures. Within-group analysis was used on data spanning multiple

experimental time blocks. Spearman correlation was applied to evaluate linear relationship

between behavioral variables. Simple linear regression was applied to evaluate

correlations. Correction for multiple tests was applied to original p-values when necessary,

by computing a new set of adjusted p-values through the two-stage False Discovery Rate

(FDR) proposed by Benjamini, Krieger and Yekutieli (38). Statistical significance level was

set at 0.05. Statistical analyses were conducted using GraphPad Prism software (v9.0.0).

285   Results

286 Increased sensitivity and delayed habituation to repeated
287 handling in VPA-treated rats

Over four handling sessions (Fig 1A), a similar proportion of VPA-treated and CTL

animals displayed defecation behavior, defined as the release of at least one fecal pellet

(Fig 1B). Analysis across experimental days showed a higher prevalence of VPA-treated

animals releasing fecal pellets compared to controls on the second day (Fig 1C). Within-

group analysis showed a significant decrease in the relative number of CTL animals

defecating after the first session indicating an early habituation to the handling stress.

Conversely, such a decrease occurred in the VPA group by the third day onward. We also

quantified the number of fecal pellets produced by responsive animals, in order to assess

the magnitude of their behavioral responses. Our results showed that there was no overall

significant difference in the number of fecal pellets released by CTL and VPA animals

during the four experimental days (Fig 1D). However, 33% of VPA-treated rats released

more than 10 fecal pellets compared to 6% of CTL rats, indicating higher sensitivity to

handling stress among VPA-treated animals. More fecal pellets were released by VPA

animals on the second day compared to CTL animals (Fig 1E). Temporal analysis across

handling sessions showed that while CTL rats habituated (reduced release of fecal pellets)

from second day on, VPA-treated animals showed delayed habituation starting from day 3

on. We also assessed differences between sexes by comparing female and male subjects

within each group. No significant differences were observed either in the prevalence or in

the magnitude of the defecation response (Data not shown).

Fig 1. Stress responses elicited by handling. (A) Schematic illustration of the

experiment. Four handling sessions of 1 min with 5 min intervals between handlings, once

a day for four consecutive days. (B) Percentage of animals releasing fecal pellets over all

experimental days. [#]$p > 0.050$. (C) Percentage of animals releasing fecal pellets on each

experimental day. [#]$p = 0.030$; CTL *$p = 0.001$; VPA *$p = 0.008$. (D) Number of fecal

pellets released over all experiment days. Dotted line for reference at 10 fecal pellets. [#]$p >$

0.050. (E) Number of fecal pellets on each experimental day. [#]$p = 0.030$; CTL *$p \leq 0.004$;

VPA *$p = 0.008$. Panels B and C, Fisher's exact test [$N_{CTL} = 30$; 15 female, 15 male. N_{VPA}

$= 28$; 15 female, 13 male]. Panel D, Mann-Whitney test. Panel E, Mann-Whitney and

Friedman tests. [$N_{CTL} = 19$; 10 female, 9 male. $N_{VPA} = 21$; 10 female, 11 male] [#], Between-

group significant differences. *, Within-group significant difference compared to D1.

Increased susceptibility of VPA-treated rats to electro-tactile stimulation

During the electro-tactile sensitivity test (Fig 2A), the VPA group showed a significantly prevalence of animals releasing fecal pellets as compared to the CTL group (Fig 2B). However, the number of pellets released per responsive animal did not differ between VPA and CTL groups (Fig 2C). Analysis of freezing behavior demonstrated no significant group differences in either the prevalence of animals displaying freezing (Fig 2D) or the relative duration of freezing among responsive animals (Fig 2E).

Fig 2: Stress responses elicited by electro-tactile stimuli. (A) Experimental design: the rats were placed in the middle of the apparatus facing one of the metal walls. After a period of 90 sec of free exploration, single electrical stimuli of 1 sec were applied with increasing intensity on 50 µA steps, random inter-stimuli intervals between 30 and 90 sec. The test ended with the presentation of nociceptive variables by two consecutive stimuli. (B) Percentage of animals that released fecal pellets. [#]$p < 0.001$. (C) Number of fecal pellets released. [#]$p > 0.050$. (D) Percentage of animals displaying freezing behavior. [#]$p > 0.050$. (E) Percentage of freezing behavior. [#]$p > 0.050$. Panels B and D, Fisher's exact test [N_{CTL} = 30; 15 female, 15 male. N_{VPA} = 28; 15 female, 13 male]. Panel C, Mann-Whitney [N_{CTL} = 11; 7 female, 4 male. N_{VPA} = 24; 14 female, 10 male]. Panel E, [N_{CTL} = 8; 5 female, 3 male. N_{VPA} = 12, 6 female, 6 male]. [#], Between-group significant differences.

Additionally, assessments of tactile and nociceptive thresholds indicated no significant differences between VPA-treated and CTL animals (Fig S2B). In order to assess sex differences, we compared the behavior of females and males in each group. Our results showed no sex differences in stress susceptibility, magnitude of defecation response or freezing behavior in the VPA or CTL group. Similarly, no sex differences were

345 detected in the nociceptive threshold. However, female CTL rats exhibited higher tactile

346 thresholds compared to their male counterparts, whereas no sex differences were

347 observed among VPA-treated animals (Fig S2C).

349 Delayed onset and increased magnitude of freezing in VPA-

350 treated rats during fear conditioning

VPA and CTL groups had similar prevalence of animals displaying defecation

behavior during fear conditioning (Fig 3A and 3B). No significant difference was detected

in the number of fecal pellets produced by responsive rats in either group (Fig 3C).

Freezing behavior was also similarly prevalent in CTL and VPA-treated animals (Fig 3D).

Within-group analyses revealed that, in both the VPA and CTL groups, the prevalence of

freezing behavior significantly increased from the second block (B2) onward compared to

baseline (B0; Fig 3E). Comparing the groups, responsive animals differ in their overall

freezing behavior (Fig 3F). Initially, VPA-treated rats exhibited a delayed onset of freezing

compared to CTL. While CTL animals showed a significant increase in freezing behavior

from B1 onward relative to baseline (B0), VPA-treated rats only displayed a comparable

increase starting at B2. Moreover, VPA-treated animals exhibited a more pronounced

freezing response, peaking at B4, where a statistically significant difference was observed

compared to CTL animals.

Figure 3: Nociceptive stress response (fear conditioning). (A) Fear conditioning

experimental design: Following a 4-min period of free exploration, five electrical shock

stimuli were applied. After the last stimulus the animal remained in the chamber for 4-min

of free exploration. (B) Percentage of animals releasing fecal pellets. (C) Number of fecal

pellets released. (D) Percentage of animals that displayed freezing behavior over all

experimental blocks. (E) Percentage of animals that displayed freezing behavior on each

experimental block. Dotted lines at 40 and 80% for reference. *p < 0.001. (F) Percentage

of time in freezing behavior in each experimental block. B4, #p = 0.046; CTL, *p = 0.010;

VPA, *p < 0.001. (G) Percentage of animals that emit USV. #p > 0.050. (H) Percentage of

animals that emitted USV on each experimental block. CTL *p ≤ 0.014, VPA *p ≤ 0.026.

(I) Number of calls per experimental block emitted by vocalizing animals. CTL *p ≤ 0.003,

VPA *p ≤ 0.014. (J) Principal frequency (F0) of calls emitted in blocks B_{1-5}. (K) Median call

duration in blocks B_{1-5}. (L) Correlation analysis between relative freezing behavior and

number of long 22-kHz calls emitted by CTL and VPA animals. Each point represents an

individual. The trend line indicates the direction and strength of the correlation between

behaviors. *p = 0.001. (M) Correlation analysis between relative freezing behavior and

number of fecal pellets released by CTL and VPA animals. Each point represents an

individual. The trend line indicates the direction and strength of the correlation between

behaviors. Panels B, D-E, G-H, Fisher's exact test. B [N_{CTL} = 31, 15 female, 16 male; N_{VPA}

= 30, 14 female, 16 male]; D-E [N_{CTL} = 27; 13 female, 14 male. N_{VPA} = 29; 14 female, 15

male]; G-H [N_{CTL} = 28; 13 female, 15 male. N_{VPA} = 29; 14 female, 15 male]. Panel C Mann-

Whitney test [N_{CTL} = 30; 15 female, 15 male. N_{VPA} = 29; 14 female, 15 male]. Panel F,

Student-t test and RM one-way ANOVA [N_{CTL} = 27; 13 female, 14 male. N_{VPA} = 29; 14

female, 15 male]. Panel I, Mann-Whitney and Friedman tests [N_{CTL} = 22; 11 female, 11

male. N_{VPA} = 21; 9 female, 12 male]. Panels J-K, Mann-Whitney test [N_{CTL} = 12; 7 female,

5 male. N_{VPA} = 9; 5 female, 4 male]. Panels L, M, Simple linear regression. L [N_{CTL} = 12; 7

female, 5 male. N_{VPA} = 9; 5 female, 4 male]. M [N_{CTL} = 25; 13 female, 12 male. N_{VPA} = 28;

14 female, 14 male]. #, Between-group significant difference. *, Within-group significant

difference compared to B0. @, correlation significant difference.

The prevalence of vocalizing animals did not differ significantly between groups,

either when analyzed globally or across individual experimental blocks (Fig 3G and 3H).

Both VPA and CTL groups exhibited an increased prevalence of ultrasonic vocalization (USV) emitters from block B4 onward, compared to baseline (B0). Although not statistically significant, a slightly higher proportion of CTL animals vocalized across all blocks relative to the VPA group. Among animals classified as responsive (i.e., those that emitted at least one USV), no between-group differences were found in the total number of 22-kHz calls (Fig 3I). However, a significant within-group increase in call number was observed in both groups at B5. There were no significant differences between groups in the mean duration of the 22-kHz calls or mean fundamental frequency (F_0; Fig 3J and 3K). Notably, however, vocalizations from VPA-treated animals displayed a bimodal distribution of mean call durations, in contrast to the unimodal pattern observed in CTL. Additionally, a subset of VPA-treated rats (22%; 2/9) produced calls with F_0 values exceeding the third quartile of the CTL distribution (i.e., >30 kHz).

Given that freezing, 22-kHz ultrasonic vocalizations, and defecation are behavioral indicators of fear, stress, and anxiety, we examined the correlations among these variables. Of the animals tested, 12 out of 27 CTL and 9 out of 29 VPA-treated rats exhibited both freezing behavior and 22-kHz vocalizations, while 26 out of 27 CTL and 28 out of 29 VPA-treated rats displayed freezing behavior and produced fecal pellets. These subsets were included in the respective correlation analyses. In the CTL group, a significant positive correlation was observed between freezing duration and the number of 22-kHz calls, along with a small but significant negative correlation between freezing and the number of fecal pellets released (Fig 3L and 3M). In contrast, no significant correlations between these behavioral measures were found in the VPA group. Finally, analyses of sex differences showed no significant effects on either the prevalence and magnitude of defecation, freezing and vocal response (data not shown).

Memory retrieval was assessed 24 hours after fear conditioning training (Fig S3). Both groups showed a similar proportion of animals exhibiting freezing behavior, with

423   freezing levels significantly elevated relative to baseline (B0). Among responsive animals,

424   freezing levels during the retrieval session did not differ significantly between VPA-treated

425   and CTL rats. To evaluate potential sex differences in freezing behavior, females and

426   males within each treatment group were compared. No significant differences were

427   observed between sexes in either the control or VPA groups (data not shown).

428 Prenatal exposure to VPA promotes delayed fear response
429 during emotional contagion

During the emotional contagion test, OBS animals—both VPA-treated and CTL—were

exposed to a DEM animal (naive) that either received a series of five-foot shocks or no

shocks (Fig 4A). During the experiment, we assessed their behavioral responses by

measuring freezing and 22-kHz vocalizations. Our results showed that witnessing DEM

animals receiving successive foot shocks engaged more animals of both groups (oC+ and

oV+) in freezing behavior. This effect was not observed in OBS witnessing DEM animals

that did not received shock (oC- and oV-). Between-group comparisons showed that, at B2

there was a significantly larger engagement in freezing of oC+ animals compared to oC-,

as well as of oV+ animals compared to oV-. Notably, in the oV+ group, this elevated

freezing response persisted through B3 (Fig 4B). In addition, within-group analysis

(compared to B0 baseline), a significantly higher prevalence of freezing behavior was

observed in the oC+ group at B2 and B4 and in the oV+ group at B2-B5. We also

observed an increase in the relative number of animals expressing freezing behavior at

B4-B5 in oC- and oV- groups, but they were not statistically significant.

Figure 4: Behavioral analysis during negative social stress (emotional contagion).

(A) Experimental design: Following a 4-min period of free exploration, five electrical shock

stimuli were applied to the demonstrator. After the last stimulus the animals remained in

the chamber for four minutes of free exploration. (B) Percentage of observer animals

displaying freezing behavior. (C) Percentage of observer animals displaying freezing

behavior on each experimental block. [#], $p = 0.036$; *, $p \leq 0.036$. (D) Percentage of time

displaying freezing behavior on each experimental block. Dotted line at 20% for reference.

[#], $p \leq 0.029$; *, $p \leq 0.015$. (E) Percentage of animals that vocalized long 22-kHz calls over

all the experiment. (F) Percentage of animals that vocalized (at all frequencies) on each

experimental block. *$p = 0.007$. (G) Number of calls on each experimental block.

Horizontal (dotted and dashed) lines at 20 and 100 calls for reference. Vertical line

separate each block. *, $p \leq 0.0306$. (H) Call rate on each experimental block. * $p < 0.03$.

(I) Principal frequency of calls over blocks B_{1-5}. (J) Call duration over blocks B_{1-5}. Dashed

line represents 0.1 seconds for reference. (K) Correlation analysis between relative

freezing behavior and number of long 22-kHz calls emitted by OBS animals. Each point

represents an individual. The trend line indicates the direction and strength of the

correlation between behaviors. Panels B,C, E and F, Fisher's exact test. Panels D and H,

two-way ANOVA and RM two-way ANOVA. Panel G, Kruskal-Wallis and Friedman tests.

$N_{oC-} = 15$; 7 female, 8 male. $N_{oV-} = 15$; 7 female, 8 male. $N_{oC+} = 16$; 8 female, 8 male. N_{oV+}

$= 15$; 8 female and 7 male) for panels B and C. $N_{oC-} = 13$; 5 female, 8 male. $N_{oV-} = 15$; 7

female, 8 male. $N_{oC+} = 16$; 8 female, 8 male. $N_{oV+} = 15$; 8 female, 7 male for panel D. N_{oC-}

$= 15$; 7 female, 8 male. $N_{oV-} = 13$; 6 female, 7 male. $N_{oC+} = 15$; 8 female, 7 male. $N_{oV+} =$

15; 8 female, 7 male for panels E-J. $N_{oC-} = 13$; 5 female, 8 male. $N_{oV-} = 13$; 6 female, 7

male. $N_{oC+} = 15$; 8 female, 7 male. $N_{oV+} = 15$; 8 female, 7 male for panel K.[#], Between

groups: Shock (+) vs. nonShock (-) significant differences. *, Within-group significant

difference compared to B0. *, Within-group significant difference compared to Tac. [&],

female-male within-group difference.

When comparing the freezing behavior of responsive animals (i.e., those exhibiting any

freezing) at B2, both oC+ and oV+ rats displayed significantly higher levels of freezing

compared to their respective controls (oC- and oV-). Notably, the difference between oV+

and oV- rats persisted through B3 (Fig 4C). Within-group analysis revealed that both oC+

and oV+ animals exhibited increased freezing in response to witnessing repeated shocks

administered to their DEM partner, relative to baseline (B0). This increase reached

statistical significance starting at B2. Notably, while the heightened freezing in oC+ rats

was restricted to B2, oV+ rats maintained an elevated freezing response from B2 through

B4.

The prevalence of animals engaging in vocalizations during the experiment (i.e. those

animals that emitted at least one 22-kHz or 50-kHz call) was similar between groups,

despite a trend to less engagement of animals in oC+ and oV+ groups as more stimuli

were applied to DEM (Fig 4D and 4E). Within-group analysis indicated that the oC+, oV+,

and oC– groups exhibited a reduction in the proportion of animals vocalizing in response

to witnessing their DEM partner receive a shock, relative to B0. This reduction reached

statistical significance beginning at B3. Notably, in oC+ rats, the decrease in vocalization

was restricted to B3, whereas in oV+ rats, it persisted through B5. Interestingly, the oC-

group also showed a significant reduction in vocalization at B5 (Fig. 4E). Changes in the

number of vocalizations produced by OBS animals were also observed compared to the

baseline, with no differences between groups (Fig 4F). Among responsive animals, a

reduction in vocal behavior was observed from block B3 onward in the oC+ and oV+, as

well as in oV- groups. In contrast, the oC⁻ animals exhibited a significant decrease as early

as block B1, relative to B0. Analysis of vocalization rate (USVs per minute) revealed that

oV+ animals exhibited a consistent reduction during sessions B2, B3, and B5. In contrast,

both oC+ and oC- groups showed a decline in vocal behavior only from session B4

onward. The oV- group maintained a stable vocalization rate across all sessions. Notably,

both oC+ and oV+ animals displayed a peak in vocalization rate during session B1,

followed by a progressive decline in subsequent blocks. (Fig 4G).

When basic bioacoustic properties (duration and f_0) of the ultrasonic calls were

analyzed, we noticed that calls produced by oC+ and oV+ animals had longer durations at

blocks B1-B5 when compared to calls produced at B0 (Fig 4H). However, no differences

were observed in f_0 either across blocks or between groups (Fig 4I). Correlation analysis

between freezing and vocal production released during the emotional contagion showed

no significant differences (Fig 4J). No sex-related differences were observed in the

susceptibility and magnitude of the freezing and vocal responses (data not shown).

Considering that the behavioral responses displayed by OBS animals should depend

on the behavior of DEM rats, we also quantified freezing and vocalizations produced by

these animals. Exposure to shock significantly increased prevalence of displaying freezing

compared to non-shocked controls (dV− and dC−), regardless of pairing with VPA-treated

or CTL counterparts. (Fig S4A-B). No significant differences were observed between the

dV+ and dC+ groups or between the dV− and dC− groups. Among responsive animals,

both dC+ and dV+ groups exhibited a significant increase in freezing behavior following

the delivery of shock stimuli, as expected (Fig S4C). Compared to B0, dC+ and dV+

increased freezing from B1 to a plateau between B2-B5. dV- and dC- animals showed no

changes in freezing behavior relative to B0 levels. Importantly, no differences were

observed in freezing between either dC+ vs. dV+ or dC- vs. dV- (p > 0.05).

Among dV+ and dC+ animals, there was a high prevalence of vocalizing animals

compared to dV- and dC- rats (Fig S4D). The vocal engagement across experimental

blocks revealed (1) no group differences between dV+ vs. dC+ or between dV- vs. dC-; (2)

significant differences between dV+ vs. dV- (with onset at B2) and between dC+ vs. dC-

(with onset at B1) and (3) an increase relative to B0 levels in dV+ and in dC+ (from B2

onwards; Fig S4E). dV- and dC- animals showed no changes in vocal engagement relative

to B0 levels.

DEM rats exhibited robust engagement in fear-related ultrasonic vocalizations (USVs)

following foot shock exposure during the emotional contagion experiment. No significant

differences in the number of USVs were observed between dV+ and dC+ groups;

however, both groups demonstrated a marked increase in USV emissions from day B4

onward relative to their respective controls (dC- and dV-; Fig S4F).Besides, significant

vocal production increased was observed in dV+ (at B2-B5), dC+ (at B2, B4-B5), as well

as dV- (at B1) animals, compared to B0. No significant within-group changes were

observed in calls emitted by dC- rats. Bioacoustic properties such as call duration and f_0

did not change between dV+ and dC+ (Fig S4G and S4H). Owing to the low number of

vocalizations observed in dV- and dC- animals (median of 3 per block), their acoustic

properties were not analyzed.

Discussion

Animal models remain essential for elucidating the mechanisms underlying emotional

dysregulation in neurodevelopmental disorders and for developing therapeutic strategies.

In the present study, we systematically examined the stress responsiveness across

multiple behavioral paradigms and the emotional contagion response in a rat model of

autism induced by the prenatal exposure to VPA.

Unfamiliar handling in rodents is known to elicit stress-related behaviors such as

avoidance and increased defecation, which typically attenuate with repeated exposure to

the experimenter (39–41). In the present study, the greater number of VPA-treated ani-

mals releasing fecal pellets, as well as the greater number of fecal pellets released during

the first experimental days, compared do CTL, suggest an increased susceptibility to

stress responses to innocuous stimuli. Furthermore, although both VPA-treated and CTL

animals showed a reduction in fecal pellet production after four days of handling, suggest-

ing successful habituation responses in both groups, the delayed reduction observed in

VPA rats indicates a slower adaptation to nonthreatening stimuli, consistent with reports of

atypical sensory processing, stress management, and impaired desensitization in individu-

als with ASD. Evidence from human studies consistently demonstrate altered responses

to tactile stimuli in ASD, characterized by both hyper- and hypo-behavioral reactivity, but

with hyper-reactivity being the most consistently observed phenotype (42–49). These be-

havioral manifestations suggest autonomic nervous system (ANS) dysregulation. This al-

teration is accompanied by elevated baseline electrodermal activity and heart rate, corrob-

orating the dysregulation of the ANS (13–15). A recent study examining ANS responses to

affective touch (e.g., gentle stroking of hairy skin) in children reported smaller autonomic

reactivity in those with ASD (13). Similarly, the increased defecation response of VPA rats

during early handling sessions may reflect analogous autonomic dysregulation and an ele-

vated susceptibility for stress responses to innocuous stimuli.

Although a greater proportion of VPA-treated animals exhibited defecation in response

to electro-tactile stimulation compared to controls, suggesting increased susceptibility,

neither the freezing behavior nor the number of fecal pellets differed significantly between

groups, indicating comparable magnitudes of stress responses. These findings are

consistent with our observations of tactile and nociceptive thresholds and align with

previous studies demonstrating preserved nociceptive thresholds to electrical stimulation

in VPA-exposed rats (50). Notably, prior investigations into tactile sensitivity in VPA-

treated animals have consistently reported increased responsiveness to mechanical

allodynia and diminished sensitivity to thermal nociceptive stimuli (35,36,51–55).

During the fear conditioning paradigm, VPA and CTL animals were similarly

susceptible to stress-induced defecation, freezing, and 22-kHz vocalization behaviors,

following a similar time-course in both groups. Although studies examining the dynamics of

the freezing response in VPA-treated animals remain limited, our findings are consistent

with a previous report showing increased shock-induced freezing in VPA rats relative to

CTL controls (56). In contrast, other studies assessing overall freezing across the entire

experimental session have reported preserved freezing behavior in VPA-treated animals

(36,50,57). These discrepancies may arise from variations in experimental design, strain-

specific factors, or developmental stage differences. Further analysis of our data indicated

a delayed onset of freezing behavior in the VPA group, potentially reflecting impairments in

the integration or coordination of stress-related emotional responses, including sensory

processing, decision-making, motor planning and execution, or disruptions in emotional

and motivational regulation. This delayed response, alongside preserved vocalization,

reinforces the hypothesis of altered integration or coordination of emotional processing

under stress. Moreover, the observed negative correlation between freezing behavior and

fecal pellet release—along with the progressive increase in freezing observed in CTL

animals—suggests that defecation predominates at lower stress intensities, whereas

freezing becomes more prominent as stress levels escalate. The lack of this correlation in

VPA-treated rats provides additional evidence for a disruption in the integration of

physiological responses associated with emotional stress.

During fear retrieval, behavioral responses were limited, particularly for vocalizations

and defecation, restricting the analysis to freezing behavior. No group differences were

observed in freezing magnitude or incidence. These findings are consistent with prior

reports indicating preserved retrieval responses in VPA rats and further suggest that acute

associative memory of the conditioned stimulus is intact in this model.

Nociceptive responses in individuals with autism spectrum disorder (ASD) vary across

developmental stages. Human studies have shown that children with ASD often exhibit

hypersensitivity to pressure and cold pain, while adolescents tend to display thermal pain

thresholds within the normative range. In adults, thermal pain thresholds are preserved,

but pain ratings are elevated and more variable (58–65). These developmental discrepancies in pain sensitivity further underscore the complexity and heterogeneity of sensory and stress-related phenotypes in ASD. Our findings—demonstrating preserved nociceptive sensitivity but altered stress habituation in adolescent rats—parallel this developmental trajectory and suggest translational relevance for the VPA model in this domain.

In the emotional contagion paradigm, both CTL and VPA observer rats exhibited freezing and 22-kHz vocalizations in response to a distressed conspecific, consistent with prior findings in CTL animals (66,67). However, VPA observers displayed prolonged freezing and delayed habituation, suggestive of increased sensitivity to social distress or impairments in social buffering mechanisms. Notably, demonstrator animals showed comparable freezing behavior across groups (Fig. S3B–C) and only minor differences in vocalizations (Fig. S3D–F), indicating that the sensory cues available to observers were largely equivalent. Additionally, the acoustic properties of demonstrator calls were similar across conditions, supporting the conclusion that CTL and VPA observers were exposed to comparable emotional signals (Fig. S3G–H). Together, these results indicate that VPA-treated animals exhibit altered habituation and heightened responsiveness to social distress, reflecting a dysregulated emotional contagion response. This phenotype underscores the utility of the VPA model for investigating stress reactivity and social emotion regulation deficits relevant to ASD.

While emotional contagion has been well-characterized in various rodent strains, to our knowledge, no previous studies have examined this phenomenon in a rodent model of autism. Our findings thus provide novel evidence that, despite their established social deficits, juvenile VPA animals still perceive social stressors as salient, though more challenging to process. In addition, the preserved sensitivity to social cues but fail to downregulate stress responses following social exposure of VPA-treated animals results

are consistent with human data, where both children with ASD and TD peers show

increased cortisol levels in response to social interaction. However, while TD children

demonstrate cortisol habituation by the end of the exposure, this attenuation is absent in

children with ASD. Additionally, among children exhibiting the highest cortisol reactivity—

including both ASD and TD groups—those with ASD engaged in lower levels of social

interaction. Complementary findings from the same research group further report that

elevated cortisol responses to social challenges are positively associated with increased

daily stress and sensory processing difficulties (68–71). This pattern underscores the

translational value of the VPA model for investigating stress regulation abnormalities in

ASD. By capturing both behavioral and physiological features of impaired social stress

adaptation, the model offers an important platform for elucidating the underlying

neurobiological mechanisms and for identifying potential targets for therapeutic

intervention.

Notably, across all experimental paradigms, VPA-treated animals consistently exhibited

a trend toward increased involuntary responses (defecation and freezing) and slightly

reduced voluntary vocalizations relative to control animals, even in the absence of

statistically significant group differences.

Sex differences in stress responsiveness were not observed across any of the para-

digms employed. This aligns with previous research demonstrating that juvenile rats ex-

posed to VPA do not exhibit significant sex-based differences in social play behavior (72).

These findings are consistent with a human study in which VPA-exposed ASD children

have a much reduced autism male bias (1.2-2.4:1) compared to no VPA-exposed ASD

children (3.3:1) (73,74). One possible explanation for the attenuated sex differences is that

VPA acts as a neurodevelopmental risk factor whose effects may override or bypass sex-

dependent protective mechanisms. Alternatively, it has been proposed that females with

ASD may engage in compensatory social behaviors that mask symptomatology, contrib-

uting to diagnostic underrepresentation and an artificially reduced prevalence (75–78). The

absence of sex effects in our study may also reflect limitations in the sensitivity of the em-

ployed behavioral assays to detect subtle sex-dependent phenotypic variations or may

result from the homogeneity of the VPA-induced phenotype across sexes. To better eluci-

date sex-specific mechanisms, future studies should incorporate hormonal profiling or cir-

cuit-level analyses and more nuanced behavioral assessments designed to detect subtle,

sex-specific traits.

Finally, we would like to identify some limitations of the present study that include the

lack of assessment of hearing as they would be particularly important to control for individ-

uality in hearing-dependent fear responses, as well as the lack of measurements of

corticosterone levels, which would have been valuable to control for eventual stress-

related HPA dysfunctions VPA-treated animals. Considering these limitations will provide

future experiments with important directions for improvement.

Conclusion

VPA-treated rats exhibited altered stress and fear-related behaviors across multiple

paradigms. Compared to controls, they showed heightened and prolonged physiological

responses to handling and electro-tactile stimulation, as well as delayed and exaggerated

freezing during fear conditioning. The typical correlations among fear-related behaviors

observed in control animals were disrupted in the VPA group, indicating a decoupling of

behavioral stress markers. During emotional contagion, VPA-treated observers

demonstrated more sustained freezing and earlier reductions in vocalization when

witnessing a conspecific in distress, suggesting altered emotional processing. These

findings support a profile of increased emotional reactivity and impaired integration of fear-

related responses in VPA-treated animals. Collectively, these findings suggest that the

VPA model provides a valid platform for investigating the neurobiological mechanisms of

stress susceptibility and social dysfunction in ASD.

Acknowledgments

680 We are grateful to Renan Araújo de Lima, Ana Luiza Campos Vila Real and Juliana Alves Brandão

681 Medeiros de Sousa for their technical assistance with experimental apparatus design, behavioral

682 experiments and histological procedures.

References

1. Hernández-González O, Fresno-Rodríguez A, Spencer-Contreras RE, Tárraga-Mínguez R,
González-Fernández D, Sepúlveda-Opazo F. Research Mapping of Trauma Experiences in
Autism Spectrum Disorders: A Bibliometric Analysis. Healthcare. 2023 Apr 28;11(9):1267.

2. Keith JM, Jamieson JP, Bennetto L. The importance of adolescent self-report in autism
spectrum disorder: integration of questionnaire and autonomic measures. J Abnorm Child
Psychol. 2019 Apr;47(4):741–54.

3. Van Steensel FJA, Bögels SM, Perrin S. Anxiety Disorders in Children and Adolescents with
Autistic Spectrum Disorders: A Meta-Analysis. Clin Child Fam Psychol Rev. 2011
Sep;14(3):302–17.

4. Lecavalier L, Wood JJ, Halladay AK, Jones NE, Aman MG, Cook EH, et al. Measuring
Anxiety as a Treatment Endpoint in Youth with Autism Spectrum Disorder. J Autism Dev
Disord. 2014 May;44(5):1128–43.

5. Li Q, Zhao W, Kendrick KM. Affective touch in the context of development, oxytocin
signaling, and autism. Front Psychol. 2022 Nov 23;13:967791.

6. Ng-Cordell E, Wardell V, Stewardson C, Kerns CM. Anxiety and Trauma-Related Disorders in
Children on the Autism Spectrum. Curr Psychiatry Rep. 2022 Mar;24(3):171–80.

7. Kalvin CB, Marsh CL, Ibrahim K, Gladstone TR, Woodward D, Grantz H, et al. Discrepancies
between parent and child ratings of anxiety in children with autism spectrum disorder. Autism
Research. 2020 Jan;13(1):93–103.

8. Groden J, Diller A, Bausman M, Velicer W, Norman G, Cautela J. The Development of a Stress
Survey Schedule for Persons with Autism and Other Developmental Disabilities. 2001;31.

9. Kerns CM, Newschaffer CJ, Berkowitz SJ. Traumatic Childhood Events and Autism Spectrum
Disorder. J Autism Dev Disord. 2015 Nov;45(11):3475–86.

10. Haruvi-Lamdan N, Horesh D, Golan O. PTSD and autism spectrum disorder: Co-morbidity,
gaps in research, and potential shared mechanisms. Psychological Trauma: Theory, Research,
Practice, and Policy. 2018 May;10(3):290–9.

11.   Rumball F, Happé F, Grey N. Experience of Trauma and PTSD Symptoms in Autistic Adults:
Risk of PTSD Development Following DSM □5 and Non□DSM □5 Traumatic Life Events.
Autism Research. 2020 Dec;13(12):2122–32.

12.   Li ST, Chien WC, Chung CH, Tzeng NS. Increased risk of acute stress disorder and post-
traumatic stress disorder in children and adolescents with autism spectrum disorder: a nation-
wide cohort study in Taiwan. Front Psychiatry. 2024 Jan 31;15:1329836.

13.   Bufo MR, Guidotti M, Mofid Y, Malvy J, Bonnet-Brilhault F, Aguillon-Hernandez N, et al.
Atypical Response to Affective Touch in Children with Autism: Multi-Parametric Exploration
of the Autonomic System. JCM. 2022 Dec 1;11(23):7146.

14.   Kushki A, Drumm E, Pla Mobarak M, Tanel N, Dupuis A, Chau T, et al. Investigating the
Autonomic Nervous System Response to Anxiety in Children with Autism Spectrum Disorders.
Whitehouse A, editor. PLoS ONE. 2013 Apr 5;8(4):e59730.

15.   Kushki A, Brian J, Dupuis A, Anagnostou E. Functional autonomic nervous system profile in
children with autism spectrum disorder. Mol Autism. 2014;5(1):39.

16.   Mulkey SB, Du Plessis AJ. Autonomic nervous system development and its impact on
neuropsychiatric outcome. Pediatr Res. 2019 Jan;85(2):120–6.

17.   Million M, Wang L, Martinez V, Taché Y. Differential Fos expression in the paraventricular
nucleus of the hypothalamus, sacral parasympathetic nucleus and colonic motor response to
water avoidance stress in Fischer and Lewis rats. Brain Research. 2000 Sep;877(2):345–53.

18.   Posada-Quintero HF, Chon KH. Innovations in Electrodermal Activity Data Collection and
Signal Processing: A Systematic Review. Sensors. 2020 Jan 15;20(2):479.

19.   Gullett N, Zajkowska Z, Walsh A, Harper R, Mondelli V. Heart rate variability (HRV) as a way
to understand associations between the autonomic nervous system (ANS) and affective states: A
critical review of the literature. International Journal of Psychophysiology. 2023 Oct;192:35–
42.

20.   Greenlee JL, Lorang E, Olson RH, Rodriquez G, Yoon DM, Hartley S. Comparative analysis of
electrodermal activity metrics and their association with child behavior in autism spectrum
disorder. Developmental Psychobiology. 2024 Feb;66(2):e22461.

21.   Agorastos A, Mansueto AC, Hager T, Pappi E, Gardikioti A, Stiedl O. Heart Rate Variability as
a Translational Dynamic Biomarker of Altered Autonomic Function in Health and Psychiatric
Disease. Biomedicines. 2023 May 30;11(6):1591.

22.   Ali MK, Chen JDZ. Roles of Heart Rate Variability in Assessing Autonomic Nervous System
in Functional Gastrointestinal Disorders: A Systematic Review. Diagnostics. 2023 Jan
12;13(2):293.

23.   Maillot C, Million M, Wei JY, Gauthier A, Taché Y. Peripheral corticotropin-releasing factor
and stress-stimulated colonic motor activity involve type 1 receptor in rats. Gastroenterology.
2000 Dec;119(6):1569–79.

24.   Reed DE, Zhang Y, Beyak MJ, Lourenssen S, Blennerhassett MG, Paterson WG, et al. Stress
increases descending inhibition in mouse and human colon. Neurogastroenterology Motil. 2016
Apr;28(4):569–80.

25. Roelofs K. Freeze for action: neurobiological mechanisms in animal and human freezing. Phil
Trans R Soc B. 2017 Apr 19;372(1718):20160206.

26. Storsberg S, Stryjek R, Modlińska K, Gottswinter K, D'Hanis W, Kröber A, et al. Predator odor
induced defensive behavior in wild and laboratory rats: A comparative study. Physiology &
Behavior. 2018 Oct;194:341–7.

27. Kim EJ, Kim ES, Covey E, Kim JJ. Social Transmission of Fear in Rats: The Role of 22-kHz
Ultrasonic Distress Vocalization. Chapouthier G, editor. PLoS ONE. 2010 Dec 1;5(12):e15077.

28. Wöhr M, Schwarting RKW. Affective communication in rodents: ultrasonic vocalizations as a
tool for research on emotion and motivation. Cell Tissue Res. 2013 Oct;354(1):81–97.

29. Yee N, Schwarting RKW, Fuchs E, Wöhr M. Juvenile stress potentiates aversive 22-kHz
ultrasonic vocalizations and freezing during auditory fear conditioning in adult male rats. Stress.
2012 Sep;15(5):533–44.

30. Brudzynski SM, Bihari F, Ociepa D, Fu XW. Analysis of 22 kHz ultrasonic vocalization in
laboratory rats: Long and short calls. Physiology & Behavior. 1993 Aug;54(2):215–21.

31. Brudzynski SM. Biological Functions of Rat Ultrasonic Vocalizations, Arousal Mechanisms,
and Call Initiation. Brain Sciences. 2021 May 9;11(5):605.

32. Tonoue T, Ashida Y, Makino H, Hata H. Inhibition of shock-elicited ultrasonic vocalization by
opioid peptides in the rat: A psychotropic effect. Psychoneuroendocrinology. 1986;11(2):177–
84.

33. Blanchard RJ, Blanchard DC, Agullana R, Weiss SM. Twenty-two kHz alarm cries to
presentation of a predator, by laboratory rats living in visible burrow systems. Physiology &
Behavior. 1991 Nov;50(5):967–72.

34. Nicolini C, Fahnestock M. The valproic acid-induced rodent model of autism. Experimental
Neurology. 2018 Jan;299:217–27.

35. Schneider T, Przewłocki R. Behavioral Alterations in Rats Prenatally Exposed to Valproic
Acid: Animal Model of Autism. Neuropsychopharmacol. 2005 Jan;30(1):80–9.

36. Markram K, Rinaldi T, Mendola DL, Sandi C, Markram H. Abnormal Fear Conditioning and
Amygdala Processing in an Animal Model of Autism. Neuropsychopharmacol. 2008
Mar;33(4):901–12.

37. Bringas ME, Carvajal-Flores FN, López-Ramírez TA, Atzori M, Flores G. Rearrangement of
the dendritic morphology in limbic regions and altered exploratory behavior in a rat model of
autism spectrum disorder. Neuroscience. 2013 Jun;241:170–87.

38. Benjamini Y, Krieger AM, Yekutieli D. Adaptive linear step-up procedures that control the
false discovery rate. Biometrika. 2006 Sep 1;93(3):491–507.

39. Bigelow LJ, Pope EK, MacDonald DS, Rock JE, Bernard PB. Getting a handle on rat
familiarization: The impact of handling protocols on classic tests of stress in *Rattus norvegicus*.
Lab Anim. 2023 Jun;57(3):259–69.

40. Costa R, Tamascia ML, Nogueira MD, Casarini DE, Marcondes FK. Handling of Adolescent
Rats Improves Learning and Memory and Decreases Anxiety. Journal of the American
Association for Laboratory Animal Science. 2012;51(5).

41. Hurst JL, West RS. Taming anxiety in laboratory mice. Nat Methods. 2010 Oct;7(10):825–6.

42. Ramappa S, Anderson A, Jung J, Chu R, Cummings KK, Patterson G, et al. An Observed
Assessment of Sensory Responsivity in Autism Spectrum Disorders: Associations with
Diagnosis, Age, and Parent Report. J Autism Dev Disord. 2023 Oct;53(10):3860–72.

43. Williams ZJ. Examining the latent structure and correlates of sensory reactivity in autism: a
multi-site integrative data analysis by the autism sensory research consortium. 2023;

44. Baranek GT, David FJ, Poe MD, Stone WL, Watson LR. Sensory Experiences Questionnaire:
discriminating sensory features in young children with autism, developmental delays, and
typical development. Child Psychology Psychiatry. 2006 Jun;47(6):591–601.

45. Narzisi A, Fabbri-Destro M, Crifaci G, Scatigna S, Maugeri F, Berloffa S, et al. Sensory
Profiles in School-Aged Children with Autism Spectrum Disorder: A Descriptive Study Using
the Sensory Processing Measure-2 (SPM-2). JCM. 2022 Mar 17;11(6):1668.

46. Mikkelsen M. Autism spectrum disorder in the scope of tactile processing. Developmental
Cognitive Neuroscience. 2018;

47. Kern JK, Trivedi MH, Garver CR, Grannemann BD, Andrews AA, Savla JS, et al. The pattern
of sensory processing abnormalities in autism. Autism. 2006 Sep;10(5):480–94.

48. Cascio CJ. Self-reported Pleasantness Ratings and Examiner-Coded Defensiveness in Response
to Touch in Children with ASD: Effects of Stimulus Material and Bodily Location. J Autism
Dev Disord. 2016;

49. Espenhahn S, Godfrey KJ, Kaur S, McMorris C, Murias K, Tommerdahl M, et al. Atypical
Tactile Perception in Early Childhood Autism. J Autism Dev Disord. 2023 Jul;53(7):2891–904.

50. Banerjee A, Engineer CT, Sauls BL, Morales AA, Kilgard MP, Ploski JE. Abnormal emotional
learning in a rat model of autism exposed to valproic acid in utero. Front Behav Neurosci
[Internet]. 2014 Nov 12 [cited 2024 Aug 19];8. Available from:
http://journal.frontiersin.org/article/10.3389/fnbeh.2014.00387/abstract

51. Dendrinos G, Hemelt M, Keller A. Prenatal VPA Exposure and Changes in Sensory Processing
by the Superior Colliculus. Front Integr Neurosci [Internet]. 2011 [cited 2024 Aug 19];5.
Available from: http://journal.frontiersin.org/article/10.3389/fnint.2011.00068/abstract

52. Edalatmanesh MA, Nikfarjam H, Vafaee F, Moghadas M. Increased hippocampal cell density
and enhanced spatial memory in the valproic acid rat model of autism. Brain Research. 2013
Aug;1526:15–25.

53. Juybari KB, Sepehri G, Meymandi MS, Vakili Shahrbabaki SS, Moslemizadeh A, Saeedi N, et
al. Sex dependent alterations of resveratrol on social behaviors and nociceptive reactivity in
VPA-induced autistic-like model in rats. Neurotoxicology and Teratology. 2020
Sep;81:106905.

54. Mohammadi S, Asadi-Shekaari M, Basiri M, Parvan M, Shabani M, Nozari M. Improvement of
autistic-like behaviors in adult rats prenatally exposed to valproic acid through early
suppression of NMDA receptor function. Psychopharmacology. 2020 Jan;237(1):199–208.

55. Luhach K, Kulkarni GT, Singh VP, Sharma B. Attenuation of neurobehavioural abnormalities
by papaverine in prenatal valproic acid rat model of ASD. European Journal of Pharmacology.
2021 Jan;890:173663.

56. Lin HC, Gean PW, Wang CC, Chan YH, Chen PS. The Amygdala Excitatory/Inhibitory
Balance in a Valproate-Induced Rat Autism Model. Chang AYW, editor. PLoS ONE. 2013 Jan
29;8(1):e55248.

57. Sui L, Chen M. Prenatal exposure to valproic acid enhances synaptic plasticity in the medial
prefrontal cortex and fear memories. Brain Research Bulletin. 2012 Apr;87(6):556–63.

58. Riquelme I, Hatem SM, Montoya P. Abnormal Pressure Pain, Touch Sensitivity,
Proprioception, and Manual Dexterity in Children with Autism Spectrum Disorders. Neural
Plasticity. 2016;2016:1–9.

59. Riquelme I, Hatem SM, Montoya P. Reduction of Pain Sensitivity after Somatosensory Therapy
in Children with Autism Spectrum Disorders. J Abnorm Child Psychol. 2018 Nov;46(8):1731–
40.

60. Riquelme I, Hatem SM, Sabater-Gárriz Á, Montoya P. A multidimensional investigation of the
relationship between skin-mediated somatosensory signals, emotion regulation and behavior
problems in autistic children. Front Neurosci. 2023 Aug 10;17:1227173.

61. Duerden EG, Taylor MJ, Lee M, McGrath PA, Davis KD, Roberts SW. Decreased Sensitivity
to Thermal Stimuli in Adolescents With Autism Spectrum Disorder: Relation to
Symptomatology and Cognitive Ability. The Journal of Pain. 2015 May;16(5):463–71.

62. Yasuda Y, Hashimoto R, Nakae A, Kang H, Ohi K, Yamamori H, et al. Sensory cognitive
abnormalities of pain in autism spectrum disorder: a case–control study. Ann Gen Psychiatry.
2016 Dec;15(1):8.

63. Fründt O, Grashorn W, Schöttle D, Peiker I, David N, Engel AK, et al. Quantitative Sensory
Testing in adults with Autism Spectrum Disorders. J Autism Dev Disord. 2017
Apr;47(4):1183–92.

64. Vaughan S, McGlone F, Poole H, Moore DJ. A Quantitative Sensory Testing Approach to Pain
in Autism Spectrum Disorders. J Autism Dev Disord. 2020 May;50(5):1607–20.

65. Failla MD, Gerdes MB, Williams ZJ, Moore DJ, Cascio CJ. Increased pain sensitivity and pain-
related anxiety in individuals with autism. PR9. 2020 Nov;5(6):e861.

66. Atsak P, Orre M, Bakker P, Cerliani L, Roozendaal B, Gazzola V, et al. Experience Modulates
Vicarious Freezing in Rats: A Model for Empathy. Ferrari PF, editor. PLoS ONE. 2011 Jul
13;6(7):e21855.

67. Bruchey AK, Jones CE, Monfils MH. Fear conditioning by-proxy: Social transmission of fear
during memory retrieval. Behavioural Brain Research. 2010 Dec;214(1):80–4.

68. Corbett BA, Schupp CW, Lanni KE. Comparing biobehavioral profiles across two social stress
paradigms in children with and without autism spectrum disorders. Mol Autism. 2012;3(1):13.

69. Corbett BA, Swain DM, Newsom C, Wang L, Song Y, Edgerton D. Biobehavioral profiles of
arousal and social motivation in autism spectrum disorders. Child Psychology Psychiatry. 2014
Aug;55(8):924–34.

70. Corbett BA, Muscatello RA, Blain SD. Impact of Sensory Sensitivity on Physiological Stress
Response and Novel Peer Interaction in Children with and without Autism Spectrum Disorder.
Front Neurosci [Internet]. 2016 Jun 23 [cited 2024 Aug 19];10. Available from:
http://journal.frontiersin.org/Article/10.3389/fnins.2016.00278/abstract

71. Corbett BA, Schupp CW, Simon D, Ryan N, Mendoza S. Elevated cortisol during play is
associated with age and social engagement in children with autism. Mol Autism. 2010;1(1):13.

72. Melancia F, Schiavi S, Servadio M, Cartocci V, Campolongo P, Palmery M, et al. Sex□specific
autistic endophenotypes induced by prenatal exposure to valproic acid involve anandamide
signalling. British J Pharmacology. 2018 Sep;175(18):3699–712.

73. Christensen J, Gr TK. Prenatal Valproate Exposure and Risk of Autism Spectrum Disorders and
Childhood Autism.

74. Loomes R, Hull L, Mandy WPL. What Is the Male-to-Female Ratio in Autism Spectrum
Disorder? A Systematic Review and Meta-Analysis. Journal of the American Academy of Child
& Adolescent Psychiatry. 2017 Jun;56(6):466–74.

75. Dworzynski K, Ronald A, Bolton P, Happé F. How Different Are Girls and Boys Above and
Below the Diagnostic Threshold for Autism Spectrum Disorders? Journal of the American
Academy of Child & Adolescent Psychiatry. 2012 Aug;51(8):788–97.

76. Mandy W, Chilvers R, Chowdhury U, Salter G, Seigal A, Skuse D. Sex Differences in Autism
Spectrum Disorder: Evidence from a Large Sample of Children and Adolescents. J Autism Dev
Disord. 2012 Jul;42(7):1304–13.

77. Young H, Oreve MJ, Speranza M. Clinical characteristics and problems diagnosing autism
spectrum disorder in girls. Archives de Pédiatrie. 2018 Aug;25(6):399–403.

78. Lockwood Estrin G, Milner V, Spain D, Happé F, Colvert E. Barriers to Autism Spectrum
Disorder Diagnosis for Young Women and Girls: a Systematic Review. Rev J Autism Dev
Disord. 2021 Dec;8(4):454–70.

Supporting informations

S1_Fig: USV power. (A) Maternal separation (MS) at P08 schematic illustration and

graphic with USVs mean power by animal. [#]p < 0.0001. (B) Maternal separation (MS) at

P14 schematic illustration and graphic with USVs mean power by animal. [#]p < 0.0001. (C)

Fear conditioning (FC) at P35-38 schematic illustration and graphic with USVs mean

power by animal. Dotted line represents the lower power of USV emitted in the

microphone chamber. Dashed line represents the higher power of USV emitted in the

opposite chamber. Panels A, B and C, Mann-Whitney test. N opposite = 41. N microphone

= 35 for panel A. N opposite = 55. N microphone = 69 for panel B. N opposite = 153. N

microphone = 153 for panel C. [#], between-group significant differences.

S2_Fig: Tactile and nociceptive thresholds. (A) Tactile and nociceptive thresholds. [#]$p >$

0.050; *$p < 0.001$. (B) Female and male tactile thresholds. [&]$p = 0.007$. Panel A, paired

Student-t test [N_{CTL} = 30; 15 female, 15 male. N_{VPA} = 28; 15 female, 13 male]. Panel B,

Mann-Whitney [N_{CTL} = 30; 15 female, 15 male. N_{VPA} = 28; 15 female, 13 male]. [&],

between-group female/male significant differences.

S3_Fig: Memory retrieval. (A) Experimental design: the rats were placed in the middle of

the apparatus facing one of the metal walls. After five minutes, animals return for its home

cage. (B) Percentage of animals that displayed freezing behavior over the experiment. (C)

Percentage of animals that present freezing behavior throughout the experiment. *$p =$

0.0003. Panel B, Fisher's exact test. Panel C, Student-t and paired Student-t tests. N_{CTL} =

27; 13 female, 14 male. N_{VPA} = 28; 15 female, 13 male. *, within-group significant

differences.

S4_Fig: Emotional contagion DEM. (A) Emotional contagion experimental design:

Following a 4-min period of free exploration, five electrical shock stimuli were applied to

the demonstrator. After the last stimulus the animals remained in the chamber for four

minutes of free exploration. (B) Percentage of demonstrator animals that present freezing

behavior on each experimental block. CTL [#]$p \leq 0.011$; VPA [#]$p \leq 0.021$; dC+ *$p < 0.001$;

dV+ *$p \leq 0.012$. (C) Demonstrator's freezing behavior percentage in each experimental

block. CTL [#] $p \leq 0.008$; VPA [#]$p < 0.001$; dC+ *$p < 0.001$; dV+ *$p \leq 0.022$. (D) Percentage

of animals that emitted USVs throughout the experiment (B0-5). CTL [#]$p < 0.001$; VPA [#] $p \leq$

0.011. (E) Percentage of animals that emitted USVs on each experimental block. CTL [#]$p \le$

0.002; VPA [#]$p \le 0.006$; dC+ *$p \le 0.042$; dV+ *$p \le 0.027$. (F) Number of USVs on each

experimental block. CTL [#] $p \le 0.017$; VPA [#] $p \le 0.024$; dV- *$p = 0.017$; dC+ *$p \le 0.001$; dV+

*$p \le 0.042$. (G) USVs duration emitted in blocks B_{1-5}. (H) USVs principal frequency emitted

in blocks B_{1-5}. [#], Between groups: Shock (+) vs. nonShock (-) significant differences. *,

Within-group significant difference compared to B0.

A

B **% animals defecating**

CTL   VPA

#

37   86

C

N. fecal pellets

Groups

D **% animals freezing**

26   43

E

% freezing

Groups