Identification of transporters essential for survival of *Leishmania*

promastigotes in the digestive tract of sand flies

Authors

Jovana Sadlova[1,¶], Ulrich Dobramysl[2,3,¶], Barbora Vojtková[1], Tomáš Bečvář[1], Çağla Alagöz[4,5], Sandro

Möri[4], Richard J. Wheeler[2,3], Petr Volf[1], Eva Gluenz[4,6*], Andreia Albuquerque-Wendt[1,4,6-8*]

Affiliations

[1]Department of Parasitology, Faculty of Science, Charles University, Prag, Czech Republic

[2]Medawar Building for Pathogen Research, Nuffield Department of Medicine, University of Oxford,

Oxford, United Kingdom

[3]Institute of Immunology & Infection Research, University of Edinburgh, Ashworth Laboratories,

Edinburgh, United Kingdom

[4]Institute of Cell Biology, University of Bern, Bern, Switzerland

[5]Graduate School for Cellular and Biomedical Sciences (GCB), University of Bern, Bern, Switzerland

[6]School of Infection and Immunity, University of Glasgow, Sir Graeme Davies Building, Glasgow,

United Kingdom

[7]Parasite Chemotherapy Unit, Swiss Tropical and Public Health Institute, Allschwil, Switzerland

[8]Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade Nova

de Lisboa, Lisbon, Portugal

[*]Corresponding authors

andreia.albuquerquewendt@swisstph.ch (AAW)

eva.gluenz@unibe.ch (EG)

[¶]these authors contributed equally

Author contributions

Conceptualisation, AAW, EG

Methodology, UD, RJW, AAW, EG, JS, PV

Software, UD, RJW

Formal analysis, AAW, JS, UD, EG

Investigation, AAW, JS, UD, BV, TB, CA, SM

Resources, EG, PV

Data curation, AAW, EG, UD

Writing original draft, AAW

Writing review and editing, all authors

Visualisation, AAW

Supervision, AAW, RJW, EG, JS, PV

Project administration, EG, PV

Funding Acquisition, AAW, EG, RJW, PV

Competing Interest Statement:

The authors declare no competing interest. The funders had no role in study design, data collection

and analysis, decision to publish, or preparation of the manuscript.

Keywords: Transportome, *Leishmania*, Sand flies, bar-seq, phenotypic screen, V-ATPase

Manuscript contents:

Main Text

Figures 1 to 4

Supplementary Figures 1 to 7

Supplementary File 1

Supplementary Tables 1 to 8

Abstract

Leishmania amastigotes ingested by female phlebotomine sand flies are exposed to a harsh and

dynamic environment that differs markedly from the intracellular niche in the mammalian host in

temperature, pH and nutrient availability. Membrane transporter proteins, channels and pumps play a

crucial role in maintaining cellular physiology under changing environments. A systematic loss-of-

function screen of the *L. mexicana* transporter deletion mutants in macrophage and mouse infections

previously identified transporter genes important for the amastigote stage. To test which transporters

are important for the promastigote stage in the insect vector, we measured the fitness of gene deletion

mutants in *Lu. longipalpis* sand flies. Pooled libraries of different complexities, consisting of 71 to 317

barcoded parasite lines allowed for an estimation of the bottleneck size in experimental infections,

providing a foundation for similar experimental bar-seq studies. The fitness of each mutant parasite

line was measured by tracking population composition over a course of 9 days in the sand flies and

compared with the growth fitness of promastigotes over 7 days in laboratory cultures. There was a

high correlation of fitness scores *in vitro* and *in vivo*, but 34 mutants showed a loss of fitness only *in*

vivo, including deletion mutants of vacuolar H+ ATPase (V-ATPase) subunits. V-ATPase deletion

mutants expressed low levels of the metacyclic-specific transcript *sherp in vitro* and failed to generate

metacyclic promastigotes in sand flies, indicating that V-ATPase function is required for parasite

differentiation and progression through the *Leishmania* life cycle.

Author Summary

Leishmania parasites cause leishmaniases - a group of neglected tropical diseases that affect millions

of people worldwide. These parasites must survive in two radically different environments: inside a

mammalian host and within the gut of a blood-feeding sand fly. To thrive in the sand fly, *Leishmania*

undergo extensive physiological changes and depend on transporter proteins to move nutrients and

other molecules across their cell membranes. In this study, we focused on identifying which of these

transporters are critical for the parasite's survival inside the sand fly. We used a library of genetically

engineered *Leishmania* promastigotes - the parasite form adapted to the insect vector - to assess the

importance of more than 300 different transporter genes. We discovered that 34 of these transporters

are essential for successful colonization of the sand fly. Among them, one key protein complex - the

vacuolar H$^+$ ATPase (V-ATPase) pump – was found to be crucial for parasite survival in the insect

vector. Our findings deepen our understanding of how *Leishmania* adapts to life within the sand fly

and highlight potential molecular targets for disrupting its transmission.

Introduction

Protozoan parasites of the genus *Leishmania* (Kinetoplastida: Trypanosomatidae) are the causative

agents of leishmaniases (1), a group of neglected tropical diseases. Over 20 *Leishmania* species share

a digenetic life cycle, alternating between an insect vector and a mammalian host (1). Females of more

than 90 species of phlebotomine sand flies - *Phlebotomus* in the Eastern Hemisphere and *Lutzomyia* in

the Western Hemisphere - serve as their primary vectors (2).

When a female sand fly takes a blood meal from an infected mammalian host, it ingests immotile

amastigote forms, which are encased in a chitin-rich peritrophic matrix (PM) (2). During this stage,

parasites face dramatic environmental changes, including a temperature drop (from ~37 °C to ~26 °C),

pH shift (from acidic to neutral/alkaline), and altered nutrient availability (from blood components to

sugar meals and microbiome metabolites). These signals trigger rapid differentiation from amastigote

into promastigotes, often within 6 hours post feeding (3,4).

Inside the sand fly gut, *Leishmania* promastigotes differentiate into a series of different morphotypes.

(4)The initial amastigote to procyclic promastigote differentiation is marked by key metabolic

changes, including a ~10-fold increase in uptake of carbon sources (e.g., glucose and non-essential

amino acids), and elevated secretion of glycolytic end-products (5–8). These weakly motile, short-

flagellated procyclic forms undergo binary fission for at least 48-96 hours before slowing down their

replication and differentiating into highly motile elongated nectomonad promastigote forms (9). After

72-96 hours post blood feeding, the toxic products of blood digestion cause a reduction in the number

of procyclic forms (10) and enzyme-mediated PM disintegration ensures that nectomonads escape the

PM-encased blood meal into the midgut lumen (9,11). There, they bind to the midgut epithelium via

parasite- and vector-derived surface molecules such as lipophosphoglycans (LPGs) or mucin-like O-

glycoconjugates, which prevent their expulsion during defecation (12–15). Parasites of the *Leishmania*

subgenus then differentiate into replicative leptomonad promastigotes and migrate to the anterior

midgut (3). During this phase, flagellar motility is crucial for *Leishmania* parasites to migrate through

the thoracic midgut toward the stomodeal valve, the structure at the junction between the sand fly

foregut and midgut – a process required for transmission. For example, Beneke et al. (2019) used

genetically engineered *L. mexicana* promastigotes with impaired flagellar motility to infect *Lu.*

longipalpis and observed that these species require directional motility to successfully colonise the fly

(16). Furthermore, Cuvillier et al. (2003) showed that overexpression of constitutively active variant

of the ADP-ribosylation factor-like protein 3A (ARL-3A) in *L. amazonensis* promastigotes resulted in

cells with a short, non-motile flagellum, which failed to colonise *Lu. longipalpis* sand flies (17). When

reaching the stomodeal valve, parasites undergo differentiation into two main morphologically distinct

forms: replicative, short-flagellated, non-motile haptomonads and non-replicative, long-flagellated

highly motile metacyclic promastigotes (18,19). Notably, recent single cell transcriptomic evidence

from *L. major* promastigotes isolated from *Ph. duboscqi*, suggest that metacyclic promastigotes may

be further divided into two transcriptionally distinct sub-forms; replicative early metacyclics and non-

replicative late metacyclics (4). Moreover, the same study provides convincing evidence that in

addition to metacyclics, which are traditionally viewed as the primary infective stage, haptomonads

also play a significant role in transmission (4).

In mature sand fly infections, parasites secrete promastigote secretory gel (PSG) - a viscous matrix

rich in filamentous proteophosphoglycans (fPPGs) - which fills the thoracic midgut (20–24). In

addition, they secrete chitinase (25), which damages the insect´s alimentary canal (26). Along with

fPPGs, this disruption alters sand fly feeding behavior and promotes regurgitation during subsequent

blood meals (12,27,28). Collectively, these changes enhance parasite transmission by increasing the

number of parasites egested into the skin of the mammalian host (28).

Although our understanding of the *Leishmania* life cycle within sand flies still lags behind that of

other vector-borne diseases, several molecular determinants of vector colonisation have been

identified (3,12,16,24,29–42). Recent advances in reverse genetics and high-throughput barcode

sequencing (bar-seq) in *Leishmania* sp. (43) have significantly accelerated discovery of parasite genes

essential for promastigote fitness and vector colonisation (16,32,44). For instance, proteins required

for flagellar assembly (IFT88, LmxM.27.1130) and motility (e.g. the central pair protein PF16,

LmxM.20.1400 or the inner dynein arm protein IC140, LmxM.27.1630) (16) were found to play a

crucial role in persistence in sand flies and migration to the stomodeal valve (17). In a separate screen

targeting the parasite's kinome, ATM (LmxM.02.0120) and PI4K (LmxM.33.3590), were identified as

atypical (aPK) and phosphatidylinositol 3' kinase-related (PIKK) protein kinases, respectively,

conditionally essential for survival only in sand flies, suggesting unique pathways are involved in

vector-stage survival (32). Another kinase, MPK9 (LmxM.19.0180), was identified as essential for

sand fly colonisation (32). In an independent study, MPK9 was shown to influence flagellar length

(45), reinforcing the importance of flagellum integrity for the parasite survival inside the vector (45).

Moreover, single-cell RNA sequencing is beginning to reveal molecular markers of parasite

development in insect stages (4,46). Despite these advances, the role of transporter proteins in sand fly

colonisation remains largely underexplored. Exceptions include two nucleotide sugar transporters

involved in lipophosphoglycan (LPG) biosynthesis. One, LPG2 (LmxM.33.3120), encodes a GDP-

mannose transporter responsible for incorporating the initial and repeating mannose units into the

mannose-rich LPG structure and was shown to be essential for development of *L. donovani* and *L.*

major in several sand fly species, including *Ph. argentipes*, *Ph. papatasi*, *Ph. duboscqi* and *Ph.*

perniciosus (31,33,35,47). Another, LPG5A/B gene array (LmxM.24.0360-65 3120), encodes a UDP-

galactose transporter, results in reduced colonisation of *L. major* in *Ph. duboscqi* (35). This transporter

incorporates galactose units into the same repetitive LPG backbone, highlighting the importance of

glycan modifications in vector attachment and colonisation. This is further supported by ablation of

additional (non-transporter) genes involved in LPG biosynthesis (15,16,31,47). Interestingly, LPG

mediated binding between sand flies and *Leishmania* is known to be crucial in parasite-vector systems

where the vector is specific, i.e. each glycoconjugate presented by each different *Leishmania* sp.

enables extraordinary specificity to a single vector species (48,49). In contrast, permissive vectors

such as those within the genus *Lutzomyia*, have been reported to present additional binding

mechanisms involving O-glycosylated proteins (50).

Recently, we systematically assessed the fitness contribution of 312 predicted *L. mexicana* membrane

transporters, channels and pumps during the exponential phase of promastigote growth *in vitro*, in

amastigotes in human induced pluripotent stem derived macrophages (iMACs) and over 6 weeks in

mice (*in vivo*) (44). Using bar-seq, we showed that deletion of at least 40 transporters compromised

amastigote survival *in vivo* (44). The vacuolar H^+ ATPase (V-ATPase) emerged as a crucial proton

pump for the survival of parasites *in vivo*, and *in vitro* under conditions of low external pH (44). Here,

we extend that work by conducting an independent comprehensive systematic loss-of-function screen

targeting 316 single putative transporter-encoding genes and 17 gene arrays. We assessed mutant

fitness over a one-week time course *in vitro* and in a sand fly model of infection *in vivo*. This screen

revealed some mutants that show gain-of-fitness phenotypes and many with loss-of-fitness

phenotypes. While there was a positive correlation between mutant fitness *in vitro* and in the flies,

these results indicate a vital function for ion pumps, sugar nucleotide transporters, and transporters of

some other classes, notably several mitochondrial carrier proteins, for survival and fitness within their

sand fly vector. Moreover, the V-ATPase is required for effective completion of the developmental

cycle from promastigotes to metacyclics *in vivo*.

Results and Discussion

An expanded gene deletion screen of the L. mexicana *transportome reveals that most transporters are*

dispensable for promastigote survival in vitro

(44)To interrogate the fitness phenotypes of mutant promastigotes over an extended time course in

laboratory cultures and in experimental infections of *Lutzomyia longipalpis* sand flies, we generated

new CRISPR-Cas9 mutant libraries to allow for the inclusion of additional genes and gene arrays, and

to assemble pooled libraries of different complexities for the sand fly infections. In addition to the

previously defined "transportome" (44) four newly identified genes were targeted: two from the

Acetate Uptake Transporter (AceTr) family, one with similarity to the Selenoprotein P Receptor

(SelP-Receptor) family, and one Multidrug/Oligosaccharidyl-lipid/Polysaccharide (MOP) Flippase, as

classified by the Transporter Classification Database (TCDB) (51), thus expanding the number of

proteins in the *L. mexicana* "transportome" to 316 (Supplementary Table 1) (16,43,44,52). This

approach successfully generated 304 viable mutant populations, each resistant to both Blasticidin and

Puromycin selection drugs (Supplementary Tables 2,3), which were broadly organised by TCDB

families into four sub-libraries (Supplementary Table 4). Diagnostic PCR confirmed the successful

deletion of all copies of the targeted genes for 154 lines (null mutants); the remaining 132 lines where

the targeted gene was still detectable were classified as 'refractory to deletion' (Supplementary Table

3). Of the confirmed single-gene deletions, 46 had not been successfully generated in our previous

screen (44). This new mutant library also identified three additional single-member superfamilies

whose transporters appear dispensable *in vitro*, namely the mitochondrial EF hand Ca^{2+} uniporter

regulator (MICU; *LmxM.07.0110*), the Proton-dependent Oligopeptide Transporter (POT;

LmxM.32.0710) and the Selenoprotein P Receptor (SelP-Receptor; *LmxM.28.2380*) in addition to the

eight superfamilies that were previously shown to be dispensable (44).

Deletion of genes arranged in tandem arrays

A total of 17 gene arrays were targeted, including nine not previously targeted, from the following

families: AceTr (1 array), Amino Acid/Auxin Permease (AAAP, 4 arrays), Cyclin M Mg^{2+} Exporter

(CNNM, 1 array), Equilibrative Nucleoside Transporter (ENT, 1 array), Major Facilitator (MFS, 2

arrays), Mitochondrial Carrier (MC, 4 arrays), P-type ATPase (P-ATPase, 1 array), Voltage-gated Ion

Channel (VIC, 2 arrays), Zinc (Zn^{2+})-Iron (Fe^{2+}) Permease (ZIP, 1 array) (Supplementary Table 1).

Upon analysis of the 13 drug resistant mutants that survived the selection, only one AAAP array

mutant (*LmxM.34.5350* and *LmxM.34.5360*) was found to be null (Supplementary Table 1 and 3). For

the *LmxM.18.1290* and *LmxM.18.1300* array where a null mutant was previously achieved (44), only

double puromycin, blasticidin resistant mutant populations retaining at least one copy of the targeted

gene were recovered (Supplementary Table 1 and 3, Supplementary Figure 3). As discussed

previously (44), technical challenges such as high sequence similarity and underestimated gene copy

numbers can hinder the isolation of null mutants from tandem arrays. For instance, at first, we failed in

the isolation of a null mutant for the glucose transporter array which harbours three genes (LmGT1-
LmGT3), despite its successful deletion in *L. mexicana* using a strategy of gene knockout by
homologous recombination (53,54). In this case, we repeated transfections using both puromycin and
blasticidin selection, but only double drug-resistant populations emerged that had retained at least
some LmGT genes. It was only after a third round of transfection and subsequent selection of clonal
cell lines that we successfully isolated three clones lacking the entire array (Supplementary Figure 6A-
C). The doubling times of two null mutant clones were significantly increased (8.73 and 8.24 h)
compared to that of the parental cells (5.14 h) (Supplementary Figure 6D-E). This suggests that
although the GT-array null mutants are viable, mutants that somehow retained one or several of the
genes from the targeted array may have a significant growth advantage in mixed populations. These
data show that while it is possible to achieve null array mutants, the technical challenges in identifying
and isolating array mutants precludes phenotype screens at scale with this bar-seq method.

Less than 30% of the Leishmania *transportome is essential for promastigote survival*

Consolidating data across the two independently generated libraries indicates that 225 (~71%)
transporter-encoding genes are dispensable for promastigote survival in standard *in vitro* laboratory
cultures (Supplementary Table 3; Supplementary Figure 1, 2) (44). The successful deletion of these
genes is positive proof that they are not essential for cell proliferation under the tested conditions,
although they may still contribute to fitness. Conclusive statements cannot be made however about the
importance of genes where a deletion attempt failed. For the 91 genes refractory to deletion in two
independent screens (Supplementary Table 3), further attempts at gene deletion may yet prove
successful. Data released from the LeishGEM genome wide gene deletion screen
(https://browse.leishgem.org/) (55) already reports several transporter gene deletion mutants that were
not retrieved in the current screen. Examples include transporter-encoding genes located on
supernumerary chromosomes like the putative acidocalcisome inositol 1,4,5-triphosphate
receptor/Ca^{2+} release channel (LmxM.16.0280), one of the four amino acid transporters of the AAT1
locus, AAT1.4 (LmxM.30.0350), a glycosomal ABC transporter, GAT1 (LmxM.30.0540) and a major

facilitator (LmxM.30.0720); on diploid chromosomes a porphyrin transporter (LmxM.17.1430), the

amino acid transporter, AAT23.2 (LmxM.27.0680) and a mitochondrial carrier (LmxM.29.2240)

(Supplementary Table 3). Conditional gene knockout strategies would be required for conclusive

functional validation and stronger support for a claim of "essentiality" (56).

Prolonged culturing of transporter deletion mutants identifies novel growth fitness phenotypes in vitro

We next asked whether gene disruption had any effect on the relative growth rates of the surviving

promastigotes in standard laboratory cultures, over a one-week time course. To assess their relative

fitness, all 304 viable isolated barcoded transporter mutants and 13 array mutants, were combined into

a single masterpool. To this pool we added five barcoded parental control lines (SBL1-5), eight non-

transporter null mutants with previously characterised phenotypes [three independently barcoded

Δ*LPG1* (*LmxM.25.0010*, normal growth, important for sand fly colonisation), two independently

barcoded Δ*PF16* (*LmxM.20.1400*, normal growth, essential for sand fly colonisation), three

independently barcoded Δ*IFT88* (*LmxM.27.1130*, very slow growth, essential for sand fly

colonisation) (17), and one non-transporter null mutant Δ*LmxM.15.0240* (nonspecific lipid-transfer

protein) (44) (Supplementary Table 4). This masterpool was split into three separate flasks and grown

in standard M199 culture medium for seven days. Cultures were diluted into fresh medium twice

during this period (Fig 1). DNA was sampled at baseline (0 hours), and after 24, 48, and 144 hours

(Fig 1) and each mutant's relative representation over time was assessed by measuring DNA barcode

abundance at the sampled time points and calculating the proportion of each barcode at a given time

point relative to its representation in the starting population (Supplementary Table 5). This showed

that the parental control cells and a majority of the mutants maintained a flat trajectory, indicating that

they proliferated in the population at similar rates. There were a small number of mutants with an

upwards trajectory, indicating their proportion within the pool increased over time, and a larger set

where barcode proportions sharply decreased over time (Fig 1C, Supplementary Table 5,

Supplementary Figure 4,5). To quantify this, fitness scores were calculated by comparing the change

in barcode proportions for a given mutant to that of the mode of the cell line change distribution

(Supplementary Table 5). After 144 hours, seven mutants displayed enhanced fitness in the

promastigote *in vitro* pool (fitness score above 2 and $p < 0.05$, Fig 1D): two mutants lacking a

mitochondrial carrier (Δ*LmxM.15.0120* and Δ*LmxM.34.3330*), two ABC transporters (Δ*LmxM.06.0100*

and Δ*LmxM.11.1290*), two amino acid permeases (Δ*LmxM.30.1820* and Δ*LmxM.27.0680*) and one

hypothetical protein of the MSF family (Δ*LmxM.34.2810b*). In contrast, 107 transporter mutants

exhibited significantly reduced fitness (score below 0.5 and $p < 0.05$, Fig 1D), with two barcodes

dropping below the detection limit at 48 h and eleven at 144 h (zero read counts in all three replicates,

Supplementary Table 5). Amongst the mutants disappearing rapidly from the population was a

confirmed deletion of ABCB3, which acts in heme and cytosolic iron/sulfur clusters biogenesis and is

required for *L. major* virulence (57). This severe loss of fitness in culture may explain why previous

attempts to generate null mutants for this transporter were unsuccessful (44,57). Still detectable at the

lowest levels were confirmed null mutants for a predicted sodium/hydrogen exchanger of the CPA1

family (Δ*LmxM.14.0980* score 0.014, p=0.0001) and a putative calcium-transporting P-ATPase

(Δ*LmxM.32.1010*, score 0.009, p=0.004). Also significantly depleted were mutants for predicted

ADP/ATP carrier proteins: Δ*LmxM.07.0530* (MCP15, refractory to deletion, MCP nomenclature taken

from (58)) and the tandem array *LmxM.19.0200_LmxM.19.0210* (MCP5, refractory to deletion). The

deletion of the fourth predicted ADP/ATP carrier predicted in the *L. mexicana* genome

(LmxM.14.0990, MCP16) resulted in a less severe but also significant loss of fitness (score 0.17 and *p*

= 0.0016), indicating that these mitochondrial carrier proteins perform vital non-redundant functions

in promastigotes.

In this study, the culture conditions were designed to provide ample nutrients, a buffered environment

and constant temperature. However, over the 144 hours, cells experienced changes in population

density and two culture dilutions. After 48 h of growth and a density of $>3 \times 10^7$ parasites/ml, the

population doubling time increased from 5.36 h to 6.87 h, suggesting entry into stationary phase just

before dilution (Supplementary Table 6). Consequently, these mutants experienced a variety of

stresses, including microenvironmental pH shifts, nutrient depletion, and waste metabolite

accumulation. Thus, although the design of the knockout libraries has the inherent limitation that only

viable mutants can be assessed, these changeable conditions allowed for the detection of fitness

phenotypes and indicate that many of the null mutants would be outcompeted over time in mixed

populations. Future work comparing transporter requirements across different media and culture

conditions may reveal additional phenotypes. While M199 medium is widely used for *Leishmania*

cultivation, it was originally developed for fibroblasts (59) and is unlikely to reproduce the exact

conditions encountered within the sand fly gut.

Loss and gain of fitness in transporter deletion mutants in vivo

We next asked which mutants would be able to tolerate this more varied and harsher environment in

the alimentary tract of phlebotomine sand flies. To evaluate the relative fitness of all viable

transporter mutants *in vivo*, we created four sub-pools (named P1-P4) which were added separately to

blood feeds presented to female *Lutzomyia longipalpis* sand flies. Each pool contained five barcoded

parental control lines (SBL1-5), three non-transporter null controls with established *in vitro* and *in*

vivo promastigote phenotypes (ΔLPG1, ΔPF16, ΔIFT88), and on average 86 barcoded mutants per

pool. We reasoned that this smaller pool size would reduce the chance of mutants being lost at random

considering the small volume ingested by the sand flies. Assuming that in experimental conditions

each *Lu. longipalpis* female feed 0.8-1 µl of blood (59), each fly would be expected to ingest 8´000-

10'000 if the blood-cell suspension was prepared at 1×10^7 parasites/ml, guaranteeing an average of

93-116 parasites per mutant line, from a pool of 86 barcoded mutant lines – a level within the range

used in comparable bar-seq studies (16,32). We also infected sand flies with the masterpool to allow

for a direct test of the effect of bigger pool sizes on the outcome of the bar-seq screen. DNA was

collected from the parasite-blood mixture at 0 hours (pre-infection) and from infected sand flies after 2

days (48 h) and 9 days (216 h) post blood meal (PBM) (Fig 2A). The barcode proportions for each

mutant at each time point were quantified by sequencing.

Bar-seq performance by pool size and infection bottlenecks

To confirm the validity of our pool sizes and evaluate how many cell lines go extinct due to

population bottlenecks in the process of establishing an infection, we turned to a more comprehensive

mathematical analysis. First, we developed an evenness score representing how uniformly the mutant

lines were represented in the pools before the start of the assay. As a guide, higher values are better

and a value below 0.5 indicates a pool in which a few mutant lines dominate, with other cell lines

having low barcode read counts (see Materials and Methods). This evenness parameter was on average

0.79 for pool P1, 1.24 for pool P2, 1.02 for pool P3, 0.87 for pool P4, and 1.06 for the masterpool,

with a standard deviation across the pool replicates of less than 0.03 in all cases (Fig 3A). Next, we

checked how many cell line barcodes became undetectable in each pool's replicates at the 2-day and 9-

day time points as an estimate of how many barcodes became extinct. Using a Pólya urn model we

then estimated that between 16 and 265 cells (with a median of 50 cells) establish an infection within a

single sand fly (Fig 3B; see Materials and Methods for the mathematical derivation). This indicates

that for pools P1-P4 the number of flies chosen in the experiment was sufficient, using the criterion

that 90% of hypothetical equal-fitness cell lines survive the experimental population bottleneck on

average. However, the number of flies chosen for the masterpool was not sufficient to reach this

threshold, due to the high number of cell lines it contains despite good evenness (Fig 3C), therefore

was not included in subsequent analysis. It is important to meet the 90% survival threshold, otherwise

the interpretation of low-proportion barcodes and extinction as cell line fitness becomes difficult to

disentangle from random extinction events.

In each cohort, the parental control lines remained stable over the 9-day infection (Supplementary

Figure 5C-F), with many of the mutant barcodes following the same trajectory as the parentals,

indicating no fitness change. The trajectories of the control mutants Δ*IFT88*, Δ*LPG1*, and Δ*PF16*

indicated depletion of these mutants, as expected (16), albeit with some variation between the different

pools (Supplementary Figure 5C-F). To quantify these changes, fitness scores were calculated to

identify mutants that became significantly depleted or enriched in the flies (Supplementary Table 5).

The fitness scores of the mutants 9 days PBM in flies showed a positive correlation (Pearson r =

0.6055) with the fitness scores of promastigotes measured after 144 h in culture (Fig 2B). Across all

sub-pools, 80 barcoded transporter mutants displayed significantly reduced fitness (score < 0.5, $p <$

0.05), while nine mutants showed increased fitness (score > 2, $p < 0.05$) in the sand fly (Fig 2C-F).

There was a small number of mutants that showed a loss-of-fitness phenotype only in the fly (Fig 2B),

including a UDP-Gal nucleotide sugar transporter (LmxM.22.1010 HUT1L), one amino acid

transporter (LmxM.32.1420), two ABC transporters (LmxM.32.3260 ABCI4 and LmxM.32.1800

ABCD3, the ortholog of the *T. brucei* glycosomal transporter GAT2(61), one uncharacterised MSF

transporter (LmxM.01.0440), a putative calcium motive p-type ATPase (LmxM.34.2080) and a

subunit of the V-ATPAse (discussed below). While the substrates of most of these transporters remain

to be defined, they may contribute to the parasite's metabolic adaptation to changing environments in

the fly, or to the secretion of glycoconjugates (e.g. the HUT1L mutant). The importance of

glycoconjugate secretion *in vivo* is supported by the phenotype of the control mutant ∆*LPG1*, which

became strongly depleted in the flies. Loss of the Golgi GDP-Man transporter LPG2, which lacks a

broader range of glycoconjugates including LPG (33,35,47,62), also resulted in mutants with low

fitness scores in flies (16), as well as in *in vitro* cultures, although these did not pass the statistical

significance test.

The gain-of-fitness mutants included three ABC transporters; ABCG3 (LmxM.06.0100), ABCH1

(LmxM.11.0040) and ABCA6 (LmxM.11.1290), one MFS protein (LmxM.34.2810), one UDP-

galactose transporter (LmxM.24.0365), aquaglyceroporin 1 (AQP1, LmxM.30.0020), two folate-

biopterin (FBT) transporters; FT1 (LmxM.10.0400) and LmxM.19.0920 and one voltage-gated

calcium channels (VGCC) of the VIC family (LmxM.17.1440). The latter encodes one of two L-type

VGCCs in *Leishmania*, previously shown to be sensitive to VGCC inhibitors (63). Interestingly,

∆*LmxM.17.1440* promastigotes also exhibited significantly increased fitness *in vitro* after 24h and 144

h (this study and (63)) while their abundance decreased in macrophages at 120 h (score = 0.5, $p < 0.05$

(63)). In contrast, the second VGCC (LmxM.33.0480) showed consistently reduced fitness both *in*

vitro and *in vivo*, suggesting that these VGCCs have distinct roles in calcium homeostasis across life

cycle stages. Similarly, stage-specific phenotypes were observed for the two predicted magnesium

transporters of *L. mexicana*. Mutants where MGT2 (LmxM.25.1090) was targeted, but only double

drug-resistant populations, refractory to gene deletion were isolated, resulted in decreased fitness in

sand flies (score 0.01, p=0.008), while the MGT1 mutant (∆*LmxM.15.1310*, null) was enriched in the

flies 9 days PBM. An MGT1 null mutant was previously found to have significantly reduced fitness in

mice (44). Prior work suggests AQP1 may function in volume regulation, osmotaxis and antimony

[Sb(III)] uptake (64), although its broader biological role remains unclear. While direct measurements

sand fly gut or macrophage phagolysosome osmolality are unavailable, estimates suggest values range

from 85-448 mOsm/kg in a hematophagous insect gut (64) and 275-295 mOsm/kg in mammalian

blood (66). The phagolysosome is likely mildly hyperosmotic due to ion influx, digestion and

acidification. How these osmotic shifts across host environments are buffered in the absence of AQP1

remains to be investigated. Similarly, the phenotypes of the FBT family mutants requires further

study. Folate transporter FT1 (LmxM.10.0400) is known to be highly expressed in actively dividing

promastigotes (66,67). *Leishmania*, like many other eukaryotic cells, are folate (vitamin B9)

auxotrophs, requiring external folate uptake. Host folate availability can vary widely depending on

nutritional status and microbiota, implying that *Leishmania*'s 13 FBTs (Supplementary Figure 1B)

may be specialised for stage-specific roles. The mutant for the FBT family biopterin transporter BT1

(Δ*LmxM.34.5150*) was significantly depleted from flies 9 days PBM (score 0.003, p=0.02) and

completely lost from the promastigote *in vitro* culture.

Whether the higher fitness scores reflected more rapid proliferation or better survival or persistence in

the fly following excretion of the digested bloodmeal cannot be deduced from these data. Similarly, a

loss of barcodes from the population could indicate a higher death rate or slower proliferation. In the

in vitro assay, exponential growth of the population was precisely measured, showing a rate of 20.3

doublings over the 144 h time course (Supplementary Table 6). In the fly, the promastigotes normally

progress through a series of replicative and non-replicative developmental stages. How many exact

doublings a wild type *L. mexicana* promastigote is expected to undergo during 9 days in *Lu.*

longipalpis is not precisely determined, but it is likely to be fewer than in the constant environment of

a culture flask (12). Previous reports, suggest that promastigotes may take ~ 29 h (Supplementary

Table 6) to double in the digestive tract of a female sand fly (12). However, this is a very rough

estimate, since factors like parasite death or loss from the fly during defecation of bloodmeal

remnants, are likely the dominant reasons for severe dropouts in the sand fly. Thus, small differences

in growth rate would be hard to detect in the fly, but easily detectable in culture, especially in the high-

glucose M199 medium that maximises the growth rate of *Leishmania* promastigotes. Likely as a result

of that, we observed a greater number of mutants with small but significant changes in fitness *in vitro*

(107/304) compared to *in vivo* in sand flies (80/304) (Fig 2B, Supplementary Table 5). We can

speculate that this reflects differences between the assays, where the *in vitro* populations underwent a

larger number of doublings, combined with the technical advantage of sampling larger amounts of

DNA from the *in vitro* cultured cells, allowing for the reliable detection of small differences in

replication rate, which may not be detectable in the fly assay.

Future experiments, assessing fitness of different mutants and morphotypes within defined sections of

the sand fly gut, and in flies may provide further insights into the functions of specific transporter

proteins, as well as exposing sand flies to multiple blood meals, which can alter promastigote

differentiation and vector competence (69).

The V-ATPase of L. mexicana *is critical for survival and parasite differentiation in the sand fly gut*

(69)We recently showed that (44)loss of V-ATPase subunits had little effect on the growth of

promastigotes in standard culture medium at neutral pH but it proved detrimental to promastigotes in

acidified culture medium, and caused significant loss of fitness in macrophages and mice (44). Here,

we found that loss of V-ATPase subunits also had a profoundly detrimental effect on promastigotes *in*

vivo. Consistent with the finding that V-ATPase subunits are non-essential for promastigotes in

culture, the barcode trajectories of V-ATPase mutants showed only a moderate decline after 48 hours

of promastigote *in vitro* growth (Fig 4A) when the cells had reached a density of $>1 \times 10^7$ cells ml^{-1}

(Fig 1B); at that density, dilution into fresh medium was required to maintain the population in log

phase. Despite the decline in abundance, the barcodes of V-ATPase mutants were still represented

within the pool after 144 h of continued exponential growth. In the sand flies, the decline was more

pronounced within just 48 h PBM (Fig 4B), when the infected bloodmeal was still surrounded by

peritrophic matrix, and declined further at 9 days PBM, resulting in lower fitness scores than observed

in vitro (Supplementary Table 5).

To investigate this phenotype further, we introduced an ectopic copy of the V-ATPase subunit E (V_1E,

Fig 4C) into the V_1E null mutant cell line ($\Delta LmxM.36.3100$, KO) to generate an addback control (AB)

and compared the growth profiles of the three cell lines *in vitro* and *in vivo* (Fig 4D). The growth rates

of the KO, AB and parental cell lines were measured over 6 days, with one dilution at the end of log

phase on day 3. All three cell lines grew at a comparable rate for the first three days; after dilution the

null mutant cell line continued exponential growth but with a slightly longer doubling time. The

growth of the AB cell line was indistinguishable from that of the parental control (Fig 4D). We next

infected female *Lu. longipalpis* flies with the KO, AB or the parental *L. mex* Cas9 T7 cell line to

measure parasite abundance, location, and developmental stages over time (Fig 4 E-G). At 48 h PBM,

100% of flies infected with parental and AB lines showed heavy infections (> 1000 promastigotes/gut,

Fig 4E) and parasites were located within the endoperitrophic space surrounded by the peritrophic

matrix (Fig 4F). In contrast, the KO persisted in only 40% of flies, and infections were mostly

moderate (100-1000 promastigotes/gut). By day 9 PBM, most flies infected with parental and AB

lines developed heavy infections with colonization of the stomodeal valve in 97% of cases. In contrast,

the KO only established moderate or light infections, which were confined to the abdominal or

proximal thoracic midgut, and did not reach the cardia (Fig 4E,F).

Quantitative analysis of promastigotes at 9 days PBM showed significant differences in the

distribution of parasite morphotypes between KO, AB and parental cell lines. Morphoptype-specific

transcriptional profiles have recently been described for *L. major* isolated from *Phlebotomus duboscqi*

guts (4). A definitive set of molecular markers is currently lacking for most *L. mexicana* morphotypes.

Morphology and location were therefore used to distinguish different promastigote developmental

stages in the sand fly. Elongated cells morphometrically classified as "nectomonads" predominated in

the KO, whereas leptomonads were less abundant, and metacyclic promastigotes were completely

absent when compared to AB and parental lines (Fig 4G, Supplementary Table 7). Additionally, the

KO exhibited a significantly longer body and flagella (16 µm, p=0.000; 16.48 µm, p=0.005) compared

to the parental cells (10.67 µm; 14.91 µm) (Supplementary Table 7). Expression of an episomal copy

of the deleted gene in the KO, significantly restored body length (11.57 µm, p=0.000), but not

flagellum length (16.11 µm, p=0.747). To test whether metacyclic promastigotes formed in *in vitro*,

promastigote cultures of V-ATPase V_1E KO, AB and parental cell lines were left to grow to stationary

phase (Supplementary Figure 7). RNA was extracted every day for 5 days to measure the expression

of the metacyclic marker *sherp* (70) by quantitative real-time PCR. The results showed an increase in

the level of *sherp* transcript on day 5 in the parental and AB lines but not in the KO cell line (Fig 4H).

This suggests that V-ATPAse KOs experience a delay, or block, in metacyclogenesis, consistent with

the absence of metacyclics observed in sand flies.

In laboratory cultures, promastigotes in stationary-phase culture may encounter metabolic stress as

cultures grow dense, including waste accumulation, nutrient depletion, and cell crowding, which may

cause the longer doubling times of the V-ATPase mutants (Fig 4A,D). In the sand fly, these challenges

are intensified by competition with the host and its microbiota for limited resources and change of the

external pH in the sand fly gut, which shifts from ~6 in unfed or sugar-fed insects to ~8.15 following a

blood meal (71). Blood ingestion also triggers diuresis, a process in which hematophagous insects,

including sand flies, expel excess water and concentrate ingested blood, facilitating more blood

uptake. This process is mediated by water absorption in the midgut and urine production by the

malpighian tubules (71,72). Our results suggest that the V-ATPase may contribute to parasite fitness

under these different stresses, beyond adaptation to a low pH environment. The vacuolar proton

ATPase (V-ATPase) is a conserved multi-subunit protein complex that acts as a rotary proton pump,

using the energy from ATP hydrolysis to translocate protons across membranes. Their structure and

biochemical activity of V-ATPase has been extensively characterized in many systems, and the

primary function is the acidification of intracellular organelles. Their specific physiological functions

are however varied and context-dependent, including adaptation to pH stress, regulatory functions in

the endocytic pathway and autophagy (73,74). In kinetoplastids, V-ATPases contribute to the

acidification of acidocalcisomes (75,76) and the *T. brucei* V-ATPase was shown to regulate

endocytosis in bloodstream forms (69,77). The *L. mexicana* V-ATPase is concentrated near the

flagellar pocket (44), compatible with a role in endocytosis and endo-/lysosomal trafficking.

Moreover, autophagy (78–80) has been shown to be important for the differentiation to amastigotes

and metacyclic promastigotes *in vitro* (81). A function in autophagy could explain why

metacyclogenesis was delayed or prevented in V-ATPase mutants, both *in vitro* and in the insect

vector. Taken together, the mutant phenotypes *in vitro*, in the insect vector and in a mammalian host

(44) identify the V-ATPase as being key to the parasite's ability to tolerate and respond to changing

environments at every stage in its life cycle.

Materials and Methods

Leishmania parasites

Promastigote forms of the *L. mexicana* cell line *L. mex* Cas9 T7 (52) and all generated mutants in this

study were either grown in T25 cm^2 flasks at 27 °C or flat bottom well plates at 27 °C + 5 % CO_2 in

filter-sterilised M199 medium (Life Technologies) supplemented with 2.2 g/L $NaHCO_3$, 0.005%

hemin, 40 mM 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES) pH 7.4 and 10 % FCS

[referred in the text as "standard M199"]. 50 μg/ml Nourseothricin Sulphate and 32 μg/ml

Hygromycin B were added to the medium for the maintenance of the *sp*Cas9 and T7 RNA polymerase

transgenes (52).

Phlebotomine sand flies

A laboratory colony of *Lutzomyia longipalpis* (originating from Jacobina, Brazil) was maintained in

the insectary of the Charles University (Prague, Czechia) under standard conditions (at 26 °C fed on

50% sucrose solution with a 14 h light/10 h dark photoperiod) as described previously (75). The use of

laboratory mice for sand fly breeding has been approved by the Ministry of Education, Youth and

Sports number MSMT-25062/2023-6. Mice were kept in the animal facility of Charles University in

Prague in accordance with institutional guidelines and Czech legislation (Act No. 246/1992 and

359/2012 coll. on protection of animals against cruelty in present statutes at large), which complies

with all relevant European Union and international guidelines for experimental animals.

Revised transportome of Leishmania mexicana

This study included the genes that were previously reported in the 'TransLeish' mutant screen (44),

plus four newly identified putative membrane transporter genes: LmxM.03.0370 and LmxM.03.0390,

encode for proteins of the The Acetate Uptake Transporter (AceTr) Transporter Classification Data

Base (TCDB) family (51), LmxM.28.2380, belonging to the Selenoprotein P Receptor (SelP-

Receptor) family and LmxM.28.2410, encoding a protein of the Multidrug/Oligosaccharidyl-

lipid/Polysaccharide (MOP) Flippase TCDB family, thus updating the current size of the *L. mexicana*

transportome to a total of 316 putative members.

CRISPR-Cas9 gene knockouts

Gene deletions were done using the CRISPR-Cas9 barcoding method previously described (52).

Diagnostic PCRs for the validation of gene deletions was done as reported in (44) using ORF_Fw and

ORF_Rv primers (Supplementary Table 2). In addition to targeting each gene individually, a total of

17 tandem arrays were targeted and 8 non-transporter null mutant control cell lines were produced

[three independently barcoded Δ*LPG1*, two independently barcoded Δ*PF16*, and three independently

barcoded Δ*IFT88* (Supplementary Table 1). Fitness screens were done with populations, without drug

pressure, for which gene deletions were assessed by diagnostic PCR, without further subcloning.

Exceptionally, for the deletion of the glucose transporter array (LmGT1-GT3), new primers were

designed that captured the dissimilar UTR regions flanking the GT array. Drug resistant mutants were

cloned by limiting dilution and ORF verification primers for validation of resulting null mutant clones

were also redesigned, so that all three copies could be recognised (Supplementary Figure 6).

Generation of V-ATPase subunit E add-back cell line

The LmxM.36.3100 gene was cloned into the pT-add plasmid via restriction digestion cloning

(Supplementary File 1). Restriction sites *Spe*I and *EcoR*I were inserted into the 5' and 3' end of the

gene, respectively, with the following primers:

SubE_forward_SpeI 5' TCAAGACTAGTATGAGCGAGGCACGCCAAAT 3'

SubE_reverse_EcoRI 5' AATACAGAATTCTTACAGTGGCGCCTCGGTGT 3'

The ΔLmxM.36.3100 cell line was transfected with 5 µg of the newly generated pTadd-

LmxM.36.3100 plasmid as described elsewhere (79). After approximately 15 hours following

transfection, drug resistant cells were selected by addition of phleomycin at a final concentration of 25

µg/ml and cells were kept in presence of drug for the following 3 passages.

Pooling of cells for bar-seq experiments

The barcoded mutant and parental cell lines were combined in mixed pools, adding similar numbers of

each individual cell line. For the *in vitro* screen, a total of 290 individually targeted transporter

mutants, 13 array transporter mutants, 5 barcoded parental lines (SBL1-5; barcodes introduced into the

SSU locus (16), and 9 non-transporter knock-out mutants, of which 8 acted as controls; 3 Δ*IFT88*

(LmxM.27.1130), 3 Δ*LPG1* (LmxM.25.0010) and 2 Δ*PF16* (LmxM.20.1400) different barcoded

mutants, were combined into a pool of 1×10^5 cells/ml (Supplementary Table 4), which was split into

three aliquots for replicate measurements of *in vitro* growth.

For sand fly infections, four separate experiments were conducted using distinct pools (Supplementary

Table 4, Pool membership). Pool 1 contained 75 individually targeted transporter mutant and one

array mutant, Pool 2 contained 71 individual transporter mutants and three array mutants, Pool 3

contained, 77 individual transporter mutants and 5 array mutants and Pool 4 contained 76 individual

transporter mutants and 4 array mutants. Each pool also contained five barcoded parental lines (SBL1-

5) and three non-transporter knock-out control mutants (Δ*IFT88, ΔLPG1, ΔPF16*). Each of the four

pools was split into three equal aliquots (replicates) in preparation for the infection of female

Lutzomyia longipalpis sand flies.

Growth curves in vitro

For the *in vitro* growth assay, the mixed pool was split into 3 equal aliquots (replicates), which were left to grow in separate flasks for 48 h, diluted to 1×10^6 cells/ml, grown for an additional 24 h, diluted to 1×10^5 cells/ml and grown for an additional 72 h (total 144 hours) at 27 °C + 5 % CO_2 in standard M199. For the growth curves of individual cell lines, log phase promastigotes (between 2 and 4×10^6 parasites/ml) of V-ATPase Subunit E (LmxM.36.3100) knockout, add-back (AB) and *L. mex* Cas9 T7 (C9T7) cell lines were seeded in standard M199 at a density of 1×10^5 cells/ml. Parasites were left to grow at 27 °C for 3 consecutive days and on day 3, were diluted back to 1×10^5 cells/ml density and left to grow at 27 °C. Growth was assessed by counting the cells every 24 hours with a CASY cell counter (Cambridge Bioscience) using a 60 µm capillary and measurement range set between 2 and 15 µm. For each condition measurements from three replicate flasks were recorded.

Sand fly infections

For infections with pooled barcoded mutant populations, each pool was seeded at a density of 2×10^6 parasites/ml and grown for 24 h at 26 °C in standard M199 with 250 µg/ml of Amikacin (Amikin). On the day of infection, a total of either 3×10^7 (Pools 1-3), or 1.8×10^7 (Pool 4) logarithmic growing parasites were washed three times with sterile 0.9% NaCl saline solution (Braun) and then resuspended in 200 µl of saline, which were then mixed with 2.8 ml of ram's defibrinated blood (LabMedia), previously heat inactivated at 56 °C for 35 minutes. For each separate pool, three groups of 120-180 female sand flies, 4-5 days old, were allowed to feed on the parasite-blood mixture, through a skin membrane from a 1-day old chick, as previously described (62). Fully engorged females were separated and maintained at 26 °C with free access to 40% sucrose solution. Infected sand flies were dissected at days 2 (48 h) and 9 (216 h) post blood-meal (PBM) (Supplementary Table 8). At day 2 PBM, a total of 1 to 5 female sand fly guts were checked to qualitatively assess the progress, localisation and intensity of infection by light microscopy. Parasite abundance was graded into three qualitative categories: negative, light (<100 parasites/gut), moderate (100-1000

parasites/gut) and heavy (>1000 parasites/gut), as described elsewhere (59). For infections with

individual promastigote cell lines; female sand flies (5–9 days old) were infected by feeding through a

chick-skin membrane parasites from log-phase cultures (3–4 day cultures), washed twice in sterile

saline solution and resuspended in heat-inactivated ram blood at concentration of 1 x 10^6

promastigotes/ml. Engorged sand flies were maintained as described above. Female flies were

dissected at days 2 and 9 PBM and the abundance and localisation of *Leishmania* promastigotes in the

sand fly digestive tract was examined as described above. Experiments were performed in duplicates.

Sampling of DNA for sequencing

For the promastigote *in vitro* cultures, gDNA was extracted at 0 h, 24 h, 48 h and 144 h of growth

from approximately $1x10^7$ cells from each replicate culture, using the Wizard SV Genomic DNA

Purification System (Promega) according to the manufacturer's instructions, eluting in 40 μl of bi-

distilled water (Ambion). For the *in vivo* experiments, gDNA was extracted from the pool after mixing

in standard M199 and from the parasite-blood mixture used for the infection (time point 0). For

extraction of genomic DNA from whole infected sand flies, a total of 17 to 82 specimens were

collected from each batch at 2- and 9-days post blood meal (PBM) (Supplementary Table 5). For both

cells and tissues, the High Pure PCR Template Preparation Kit (Roche) was used and all samples were

eluted in 100 μl of VWR Life Sciences PCR grade water as previously described (16).

Bar-seq library preparation and sequencing

The preparation of bar-seq amplicon libraries was done as previously reported (44), with minor

changes. For the initial bar-seq amplicon PCR, 100 ng of gDNA isolated from promastigote cultures,

250 ng of gDNA isolated from *Leishmania*-blood meal mix and 500 ng gDNA isolated from whole

Leishmania-infected female sand flies, were used. To account for the excess of host gDNA present in

blood and sand fly derived samples, the number of cycles for the same PCR was increased from 31 for

Leishmania culture derived samples to 35 for blood and sand fly samples. Raw sequencing files (fastq)

for all samples generated during this study were deposited in the European Nucleotide Archive (ENA)

study accession PRJEB90861 (ERP173867).

Quantification of barcoded cell line fitness

For each sample, we grouped and counted raw sequencing reads containing barcode sequences,

normalised this to the sample total reads and calculated the change in the proportion of reads of each

cell line in each sample compared to the zero time point, as previously reported (45). The majority of

cell lines in a pool do not exhibit a fitness change, therefore we identified the mode (peak) of the

distribution of changes in proportion as the reference to compare to in each replicate. For each time

point and cell line we assigned fitness scores by dividing the median of the cell line's change in

proportion over all replicates with the median over all modes. A fitness score above one indicates that

proportion of barcodes from a particular cell line have increased faster relative to the bulk of the

pooled cell lines, corresponding to faster growth and/or better survival than the bulk of the pooled cell

lines from the start of the assay up to that time point. A fitness score below one indicates the inverse.

P-values were calculated using a paired t-test of the log-transformed cell line changes in proportions

against the corresponding reference values, testing the null hypothesis that the cell line change in

proportion from all replicates of a particular cell line in a given time point cannot be distinguished

from the change of the bulk of the pooled cell lines in all replicates of the same time point. Cell lines

were labelled as having a strong fitness phenotype in a given time point if their p-value was below

0.05 and their fitness score was either below 0.5 (deleterious phenotype) or above 2 (beneficial

phenotype).

Population bottleneck assessment

To determine the appropriate number of sand flies for infection to avoid mutant lines within a pool

going extinct by chance due to population bottlenecks, we first needed to determine the number of

parasites that establish an infection in a sand fly. To do so, we first assessed how evenly barcodes

were represented within each replicate of each pool by fitting a symmetrical Dirichlet distribution to

the barcode proportions using maximum likelihood estimation. The single resulting dimensionless

parameter determines the pool evenness: a small value indicates that very few cell lines dominate the

pool while a very large value corresponds to all barcodes being equally represented. Next, we treat the

experiment as

a Pólya urn model, we identify parasites in a hypothetical pool sample (using the same evenness as

measured from our actual pools) as coloured balls in an urn, with the colour representing the M

different barcodes from mutant lines in the pool. We then model the establishment of infection and

detection of

the barcode by drawing N balls from this urn (corresponding to detecting the barcode from that

parasite

cell) and count the number of different colours present C (corresponding to the number of different

mutant lines detected). The experimentally observed fraction of surviving mutant lines is S=C/N. The

fraction of surviving mutant lines can also be modelled as a function of the population bottleneck N,

the number of cell lines M and the pool evenness α is given by

$$S_\alpha(M, N) = \sum_{i=1}^{M} (1 - [1 - p_i]^N),$$

668 with the abundance of individual mutant lines p_i distributed according to a symmetric Dirichlet

669 distribution with the concentration parameter α. Lengthy but straightforward calculation yields that

670 the expectation value of the survived fraction is then given as

$$E[S_\alpha(M, N)] = 1 - \exp\left(\frac{\Gamma(M\alpha)\Gamma(N + [M-1]\alpha)}{\Gamma([M-1]\alpha)\Gamma(N + M\alpha)}\right).$$

Knowing the experimentally determined survived fraction S, the number of cell lines in the pool M

and the pool evenness α, we can now determine the population bottleneck size N by numerically

inverting the above relationship. Therefore, we can estimate the number of cells that, on average,

successfully establish an infection as N/K where K is the number of sand flies used in an experiment,

because each sand fly can be treated as independent from all others (Fig 3B).

This calculation uses the measured pool evenness from time point zero and treats our hypothetical

pool cell lines to have equal fitness and hence equal chance of survival. However, this is an

approximation as the actual fraction of surviving mutants also depends on the intrinsic fitness of the

individual mutant lines, which will tend to increase extinction events as deletion mutations tend to be

detrimental. As a result, this procedure underestimates of the population bottleneck size N and

therefore overestimates the number of sand flies required.

In vivo *parasite morphometry*

Midgut smears of infected sand flies were fixed with methanol, stained with Giemsa, examined by

light microscopy (Olympus BX51) with an oil immersion objective and photographed (Olympus

DP72). Body length, flagellar length and body width of 200 randomly selected promastigotes were

measured on day 9 PBM using Fiji (65). Four morphological forms were distinguished, based on

criteria previously described (10,28). Briefly; elongated nectomonads (EN), body length ⏃14 μm;

leptomonads (LE) body length <14 m and flagellar length ⏃2 times body length and metacyclic

promastigotes (MP), flagellar length >2 times body length and body length < 14 μm. Haptomonads

were not distinguished in this study as they are often remaining attached to the gut and can be

underrepresented on gut smears. Differences in proportion of morphological forms and measurements

were compared by a Tukey's HSD (honestly significant difference) test, using the software SPSS

version 27.

Quantification of sherp *expression by qPCR*

Cultures of V-ATPase V_1E KO, AB and parental cell lines were seeded at a density of 1×10^5 cells/ml

on day 0 and grown at 27 °C / 5 % CO_2 in standard M199, with five replicate flasks for each line.

Starting from day 1 of growth, total RNA was extracted from 1×10^8 cells every day for 5 consecutive

days, using PureLink RNA Mini Kit (Ambion by Life Technologies, Invitrogen) according to the

manufacturer's protocol, with the exception of the homogenization step. To clear the genomic DNA in

the sample, 10 μg of purified RNA was subjected to DNase I treatment (New England Biolabs),

according to the manufacturer's protocol. Afterwards, RNA cleanup was performed with the Monarch

RNA cleanup kit (New England Biolabs), according to the manufacturer's protocol. To ensure a

complete cleanup of genomic DNA, the DNase I treatment and subsequent RNA cleanup were

performed twice. cDNA was synthesized from 200 ng RNA samples using the iScript cDNA synthesis

kit (Biorad) according to the manufacturer's instructions. The qPCR was done with the SsoAdvanced

Universal SYBR Green Supermix (Biorad), according to the manufacturer's protocol. 1 ng cDNA

sample was used for each qPCR reaction. The *sherp* primer sequences were taken from (70): Forward:

5'- CTGCAGCCTTGTTGCTCA-3' ; Reverse: 5'- ACCAGGAGACAAGGGACCA-3'. The 18S

rRNA was used as a control, with primer sequences taken from (83): Forward: 5'-

CTGCAGCCTTGTTGCTCA-3' ; Reverse: 5'- ACCAGGAGACAAGGGACCA-3'. All cDNAs were

run in triplicates with each primer pair, in a qPCR-compatible 96 well plate, using Biorad C1000

touch thermal cycler. For the calculation of relative *sherp* transcript abundance, the quantification

cycles (Cqs) of each reaction with *sherp* or 18S rRNA was averaged. Then, the average Cq of 18S

rRNA was subtracted from *sherp* Cqs for the corresponding sample (ΔCq), for normalization. Relative

abundance was calculated by the formula $2^{-\Delta Cq}$; this value was multiplied by 1000 to generate the

plots.

Acknowledgements

721 We like to thank Pamela Nicholson and Daniela Steiner (Next Generation Sequencing Platform,

722 University of Bern) for help with Illumina sequencing, Vít Dvořák for maintaining the colony of

723 *Lutzomyia longipalpis*, Kristýna Srstková for technical support during sand fly experiments and all

724 past and current members of the Gluenz and Volf labs for helpful discussions.

Funding statement

AAW was the recipient of a Marie Skłodowska-Curie Individual Fellowship (trans-LEISHion-EU

FP7, No. 798736) and is supported by a Marie Skłodowska-Curie Global Fellowship (LeishBlock-

Horizon, No. 101148623). RJW is supported by a Wellcome Trust Henry Dale Fellowship

(211075/Z/18/Z). This work was supported by a UKRI Medical Research Council grant

(MR/V000446/1; This UK funded award was part of the EDCTP2 programme supported by the

European Union), the Wellcome Trust (221944/A/20/Z, 200807/Z/16/Z, 104627/Z/14/Z) and the

Wellcome Centre for Integrative Parasitology (WCIP) core Wellcome Centre Award (104111/Z/14/Z)

and a project grant from the Swiss National Science Foundation (310030_220011). This work was

partially supported by national funds through FCT – Fundação para a Ciência e a Tecnologia, I.P.,

under the R&D unit Global Health and Tropical Medicine (UID/04413/2025) and the Associated

Laboratory in Translation and Innovation Towards Global Health REAL (LA/P/0117/2020). The

funders had no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

Data availability statement

742   All data supporting the findings of this study are available within the article and its supplementary

743   materials, which have been deposited to Figshare (https://figshare.com/) under Doi:

744   10.6084/m9.figshare.29481254.

References

References

1.   Pace D. Leishmaniasis. J Infect. 2014 Nov;69(S1):S10–8.

2.   Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L. Phlebotomine sandflies and
the spreading of leishmaniases and other diseases of public health concern. Med Vet Entomol. 2013
Jun;27(2):123–47.

3.   Dostálová A, Volf P. *Leishmania* development in sand flies: parasite-vector interactions
overview. Parasit Vectors. 2012 Dec;5(1):276.

4.   Catta-Preta CMC, Ghosh K, Sacks DL, Ferreira TR. Single-cell atlas of *Leishmania*
development in sandflies reveals the heterogeneity of transmitted parasites and their role in infection.
Proc Natl Acad Sci. 2024 Dec 24;121(52):e2406776121.

5.   Hart DT, Coombs GH. *Leishmania mexicana*: Energy metabolism of amastigotes and
promastigotes. Exp Parasitol. 1982 Dec;54(3):397–409.

6.   McConville MJ, Saunders EC, Kloehn J, Dagley MJ. *Leishmania* carbon metabolism in the
macrophage phagolysosome- feast or famine? F1000Research. 2015 Oct 1;4:938.

7.   Rosenzweig D, Smith D, Opperdoes F, Stern S, Olafson RW, Zilberstein D. Retooling
Leishmania metabolism: from sand fly gut to human macrophage. FASEB J. 2008 Feb;22(2):590–602.

8.   Saunders EC, Ng WW, Kloehn J, Chambers JM, Ng M, McConville MJ. Induction of a
Stringent Metabolic Response in Intracellular Stages of *Leishmania mexicana* Leads to Increased
Dependence on Mitochondrial Metabolism. Wilson ME, editor. PLoS Pathog. 2014 Jan
23;10(1):e1003888.

9.   Gossage SM, Rogers ME, Bates PA. Two separate growth phases during the development of
Leishmania in sand flies: implications for understanding the life cycle. Int J Parasitol. 2003
Sep;33(10):1027–34.

10.   Pruzinova K, Sadlova J, Myskova J, Lestinova T, Janda J, Volf P. *Leishmania* mortality in
sand fly blood meal is not species-specific and does not result from direct effect of proteinases. Parasit
Vectors. 2018 Dec;11(1):37.

11.   Sadlova J, Homola M, Myskova J, Jancarova M, Volf P. Refractoriness of *Sergentomyia*
schwetzi to *Leishmania* spp. is mediated by the peritrophic matrix. Traub-Csekö YM, editor. PLoS
Negl Trop Dis. 2018 Apr 4;12(4):e0006382.

12.   Rogers ME, Chance ML, Bates PA. The role of promastigote secretory gel in the origin and
transmission of the infective stage of *Leishmania mexicana* by the sandfly *Lutzomyia longipalpis*.
Parasitology. 2002 May;124(5):495–507.

13.   Sádlová J, Volf P. Peritrophic matrix of *Phlebotomus duboscqi* and its kinetics during
Leishmania major development. Cell Tissue Res. 2009 Aug;337(2):313–25.

14.   Myšková J, Dostálová A, Pěničková L, Halada P, Bates PA, Volf P. Characterization of a
midgut mucin-like glycoconjugate of Lutzomyia longipalpis with a potential role in *Leishmania*
attachment. Parasit Vectors. 2016 Dec;9(1):413.

15.   Jecna L, Dostalova A, Wilson R, Seblova V, Chang KP, Bates PA, et al. The role of surface
glycoconjugates in *Leishmania* midgut attachment examined by competitive binding assays and
experimental development in sand flies. Parasitology. 2013 Jul;140(8):1026–32.

16.   Beneke T, Demay F, Hookway E, Ashman N, Jeffery H, Smith J, et al. Genetic dissection of a
Leishmania flagellar proteome demonstrates requirement for directional motility in sand fly infections.
PLoS Pathog. 2019 Jun;15(6).

17.   Cuvillier A, Miranda JC, Ambit A, Barral A, Merlin G. Abortive infection of *Lutzomyia*
longipalpis insect vectors by aflagellated LdARL-3A-Q70L overexpressing *Leishmania amazonensis*
parasites. Cell Microbiol. 2003 Oct;5(10):717–28.

18.   Bates PA. Transmission of *Leishmania* metacyclic promastigotes by phlebotomine sand flies.
Int J Parasitol. 2007 Aug;37(10):1097–106.

19.   Yanase R, Moreira-Leite F, Rea E, Wilburn L, Sádlová J, Vojtkova B, et al. Formation and
three-dimensional architecture of *Leishmania* adhesion in the sand fly vector. eLife [Internet]. 2023
May 10 [cited 2025 Jul 7];12. Available from: https://elifesciences.org/articles/84552

20.   Rogers M, Kropf P, Choi BS, Dillon R, Podinovskaia M, Bates P, et al.
Proteophosophoglycans Regurgitated by Leishmania-Infected Sand Flies Target the L-Arginine
Metabolism of Host Macrophages to Promote Parasite Survival. Ribeiro JMC, editor. PLoS Pathog.
2009 Aug 21;5(8):e1000555.

21.   Ilg T, Stierhof YD, Wiese M, McConville MJ, Overath P. Characterization of phosphoglycan-
containing secretory products of *Leishmania*. Parasitology. 1994 Mar;108(S1):S63–71.

22.   Stierhof YD, Bates PA, Jacobson RL, Rogers ME, Schlein Y, Handman E, et al. Filamentous
proteophosphoglycan secreted by *Leishmania* promastigotes forms gel-like three-dimensional
networks that obstruct the digestive tract of infected sandfly vectors. Eur J Cell Biol. 1999
Oct;78(10):675–89.

23.   Volf P, Hajmova M, Sadlova J, Votypka J. Blocked stomodeal valve of the insect vector:
similar mechanism of transmission in two trypanosomatid models. Int J Parasitol. 2004
Oct;34(11):1221–7.

24.   Yanase R, Pruzinova K, Owino BO, Rea E, Moreira-Leite F, Taniguchi A, et al. Discovery of
essential kinetoplastid-insect adhesion proteins and their function in *Leishmania*-sand fly interactions.
Nat Commun. 2024 Aug 13;15(1):6960.

25.   Rogers ME, Hajmová M, Joshi MB, Sadlova J, Dwyer DM, Volf P, et al. *Leishmania*
chitinase facilitates colonization of sand fly vectors and enhances transmission to mice. Cell
Microbiol. 2008 Jun;10(6):1363–72.

26.   Schlein Y, Jacobson RL, Messer G. Leishmania infections damage the feeding mechanism of
the sandfly vector and implement parasite transmission by bite. Proc Natl Acad Sci. 1992 Oct
15;89(20):9944–8.

27.   Ilg T, Stierhof YD, Craik D, Simpson R, Handman E, Bacic A. Purification and Structural
Characterization of a Filamentous, Mucin-like Proteophosphoglycan Secreted by *Leishmania*
Parasites. J Biol Chem. 1996 Aug;271(35):21583–96.

28.   Rogers ME, Ilg T, Nikolaev AV, Ferguson MAJ, Bates PA. Transmission of cutaneous
leishmaniasis by sand flies is enhanced by regurgitation of fPPG. Nature. 2004 Jul;430(6998):463–7.

29.   Rogers ME. The Role of Leishmania Proteophosphoglycans in Sand Fly Transmission and
Infection of the Mammalian Host. Front Microbiol [Internet]. 2012 [cited 2025 Apr 10];3. Available
from: http://journal.frontiersin.org/article/10.3389/fmicb.2012.00223/abstract

30.   Kamhawi S, Ramalho-Ortigao M, Van M. Pham, Kumar S, Lawyer PG, Turco SJ, et al. A
Role for Insect Galectins in Parasite Survival. Cell. 2004 Oct;119(3):329–41.

31.   Sacks DL, Modi G, Rowton E, Späth G, Epstein L, Turco SJ, et al. The role of
phosphoglycans in *Leishmania* –sand fly interactions. Proc Natl Acad Sci. 2000 Jan 4;97(1):406–11.

32.   Baker N, Catta-Preta CMC, Neish R, Sadlova J, Powell B, Alves-Ferreira EVC, et al.
Systematic functional analysis of *Leishmania* protein kinases identifies regulators of differentiation or
survival. Nat Commun. 2021 Feb 23;12(1):1244.

33.   Späth GF, Lye LF, Segawa H, Sacks DL, Turco SJ, Beverley SM. Persistence without
pathology in phosphoglycan-deficient *Leishmania major*. Science. 2003 Aug;301(5637):1241–3.

34.   Sádlová J, Price HP, Smith BA, Votýpka J, Volf P, Smith DF. The stage-regulated HASPB
and SHERP proteins are essential for differentiation of the protozoan parasite *Leishmania major* in its
sand fly vector, *Phlebotomus papatasi*: *Leishmania* metacyclogenesis in the sand fly. Cell Microbiol.
2010 Dec;12(12):1765–79.

35.   Secundino N, Kimblin N, Peters NC, Lawyer P, Capul AA, Beverley SM, et al.
Proteophosphoglycan confers resistance of *Leishmania major* to midgut digestive enzymes induced by
blood feeding in vector sand flies: Proteophosphoglycan protects *L. major*. Cell Microbiol. 2010 Jan
20;12(7):906–18.

36.   Gazanion E, Seblova V, Votypka J, Vergnes B, Garcia D, Volf P, et al. *Leishmania infantum*
nicotinamidase is required for late-stage development in its natural sand fly vector, *Phlebotomus*
perniciosus. Int J Parasitol. 2012 Apr;42(4):323–7.

37.   Jecna L, Dostalova A, Wilson R, Seblova V, Chang KP, Bates PA, et al. The role of surface
glycoconjugates in *Leishmania* midgut attachment examined by competitive binding assays and
experimental development in sand flies. Parasitology. 2013 Jul;140(8):1026–32.

38.   Doehl JSP, Sádlová J, Aslan H, Pružinová K, Metangmo S, Votýpka J, et al. *Leishmania*
HASP and SHERP Genes Are Required for In Vivo Differentiation, Parasite Transmission and
Virulence Attenuation in the Host. Beverley SM, editor. PLOS Pathog. 2017 Jan 17;13(1):e1006130.

39.   Sunter JD, Yanase R, Wang Z, Catta-Preta CMC, Moreira-Leite F, Myskova J, et al.
Leishmania flagellum attachment zone is critical for flagellar pocket shape, development in the sand
fly, and pathogenicity in the host. Proc Natl Acad Sci. 2019 Mar 26;116(13):6351–60.

40.   Halliday C, Yanase R, Catta-Preta CMC, Moreira-Leite F, Myskova J, Pruzinova K, et al.
Role for the flagellum attachment zone in *Leishmania* anterior cell tip morphogenesis. Hill KL, editor.
PLOS Pathog. 2020 Oct 22;16(10):e1008494.

41.   Corrales RM, Vaselek S, Neish R, Berry L, Brunet CD, Crobu L, et al. The kinesin of the
flagellum attachment zone in *Leishmania* is required for cell morphogenesis, cell division and
virulence in the mammalian host. Hill KL, editor. PLOS Pathog. 2021 Jun 18;17(6):e1009666.

42.   Alcoforado Diniz J, Chaves MM, Vaselek S, Miserani Magalhães RD, Ricci-Azevedo R, De
Carvalho RVH, et al. Protein methyltransferase 7 deficiency in *Leishmania major* increases neutrophil
associated pathology in murine model. De Oliveira CI, editor. PLoS Negl Trop Dis. 2021 Mar
2;15(3):e0009230.

43.   Beneke T, Gluenz E. Bar-seq strategies for the LeishGEdit toolbox. Mol Biochem Parasitol.
2020 Sep;239:111295.

44.   Albuquerque-Wendt A, McCoy C, Neish R, Dobramysl U, Alagoz C, Beneke T, et al.
TransLeish: Identification of membrane transporters essential for survival of intracellular *Leishmania*
parasites in a systematic gene deletion screen. Nat Commun. 2025 Jan 2;16(1):299.

45.   Bengs F, Scholz A, Kuhn D, Wiese M. LmxMPK9, a mitogen☐activated protein kinase
homologue affects flagellar length in *Leishmania mexicana*. Mol Microbiol. 2005 Mar;55(5):1606–15.

46.   Louradour I, Ferreira TR, Duge E, Karunaweera N, Paun A, Sacks D. Stress conditions
promote *Leishmania* hybridization in vitro marked by expression of the ancestral gamete fusogen
HAP2 as revealed by single-cell RNA-seq. eLife. 2022 Jan 7;11:e73488.

47.   Svárovská A, Ant TH, Seblová V, Jecná L, Beverley SM, Volf P. *Leishmania major*
Glycosylation Mutants Require Phosphoglycans (lpg2−) but Not Lipophosphoglycan (lpg1−) for
Survival in Permissive Sand Fly Vectors. Traub-Cseko YM, editor. PLoS Negl Trop Dis. 2010 Jan
12;4(1):e580.

48.   Kamhawi S. Phlebotomine sand flies and *Leishmania* parasites: friends or foes? Vol. 22,
Trends in Parasitology. 2006. p. 439–45.

49.   Volf P, Myskova J. Sand flies and *Leishmania*: specific versus permissive vectors. Trends
Parasitol. 2007 Mar;23(3):91–2.

50.   Myskova J, Svobodova M, Beverley SM, Volf P. A lipophosphoglycan-independent
development of *Leishmania* in permissive sand flies. Microbes Infect. 2007 Mar;9(3):317–24.

51.   Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, et al. The
Transporter Classification Database (TCDB): 2021 update. Nucleic Acids Res. 2021 Jan
8;49(D1):D461–7.

52.   Beneke T, Madden R, Makin L, Valli J, Sunter J, Gluenz E. A CRISPR Cas9 high-throughput
genome editing toolkit for kinetoplastids. R Soc Open Sci. 2017 May;4(5):1–16.

53.   Burchmore RJS, Rodriguez-Contreras D, Mcbride K, Barrett MP, Modi G, Sacks D, et al.
Genetic characterization of glucose transporter function in *Leishmania mexicana* [Internet]. Available
from: www.pnas.orgcgidoi10.1073pnas.0630165100

54.   Rodriguez-Contreras D, Feng X, Keeney KM, Bouwer HGA, Landfear SM. Phenotypic
characterization of a glucose transporter null mutant in *Leishmania mexicana*. Mol Biochem Parasitol.
2007 May;153(1):9–18.

55.   Aellig S, Billington K, Damasceno JD, Davidson L, Dobramysl U, Etzensperger R, et al.
LeishGEM: genome-wide deletion mutant fitness and protein localisations in *Leishmania*. Trends
Parasitol. 2024 Aug;40(8):675–8.

56.   Jones NG, Catta-Preta CMC, Lima APCA, Mottram JC. Genetically Validated Drug Targets
in *Leishmania*: Current Knowledge and Future Prospects. Vol. 4, ACS Infectious Diseases. American
Chemical Society; 2018. p. 467–77.

57.   Martínez-García M, Campos-Salinas J, Cabello-Donayre M, Pineda-Molina E, Gálvez FJ,
Orrego LM, et al. LmABCB3, an atypical mitochondrial ABC transporter essential for *Leishmania*
major virulence, acts in heme and cytosolic iron/sulfur clusters biogenesis. Parasit Vectors. 2016 Jan
5;9:7.

58.   Colasante C, Diaz PP, Clayton C, Voncken F. Mitochondrial carrier family inventory of
Trypanosoma brucei brucei: Identification, expression and subcellular localisation. Mol Biochem
Parasitol. 2009 Oct;167(2):104–17.

59.   Morgan JF, Morton HJ, Parker RC. Nutrition of Animal Cells in Tissue Culture. I. Initial
Studies on a Synthetic Medium.,. Exp Biol Med. 1950 Jan 1;73(1):1–8.

60.   Volfová V, Jančářová M, Volf P. Sand fly blood meal volumes and their relation to female
body weight under experimental conditions. Parasit Vectors. 2024 Aug 23;17(1):360.

61.   Yernaux C, Fransen M, Brees C, Lorenzen S, Michels PAM. Trypanosoma brucei glycosomal
ABC transporters: Identification and membrane targeting. Mol Membr Biol. 2006;23(2):157–72.

62.   Descoteaux A, Mengeling BJ, Beverley SM, Turco SJ. *Leishmania donovani* has distinct
mannosylphosphoryltransferases for the initiation and elongation phases of lipophosphoglycan
repeating unit biosynthesis. Mol Biochem Parasitol. 1998 Jul 1;94(1):27–40.

63.   Benaim G, García-Marchán Y, Reyes C, Uzcanga G, Figarella K. Identification of a
sphingosine-sensitive Ca^{2+} channel in the plasma membrane of *Leishmania mexicana*. Biochem
Biophys Res Commun. 2013 Jan;430(3):1091–6.

64.   Plourde M, Ubeda JM, Mandal G, Monte-Neto RLD, Mukhopadhyay R, Ouellette M.
Generation of an aquaglyceroporin AQP1 null mutant in *Leishmania major*. Mol Biochem Parasitol.
2015 Aug;201(2):108–11.

65.   Beyenbach KW. The plasticity of extracellular fluid homeostasis in insects. J Exp Biol. 2016
Sep 1;219(17):2596–607.

66.   Najem O, Shah MM, Zubair M, De Jesus O. Serum Osmolality. In: StatPearls [Internet].
Treasure Island (FL): StatPearls Publishing; 2025 [cited 2025 May 21]. Available from:
http://www.ncbi.nlm.nih.gov/books/NBK567764/

67.   Richard D, Kündig C, Ouellette M. A new type of high affinity folic acid transporter in the
protozoan parasite *Leishmania* and deletion of its gene in methotrexate-resistant cells. J Biol Chem.
2002 Aug 16;277(33):29460–7.

68.   Ouameur AA, Girard I, Légaré D, Ouellette M. Functional analysis and complex gene
rearrangements of the folate/biopterin transporter (FBT) gene family in the protozoan parasite
Leishmania. Mol Biochem Parasitol. 2008 Dec;162(2):155–64.

69.   Baker N, Hamilton G, Wilkes JM, Hutchinson S, Barrett MP, Horn D. Vacuolar ATPase
depletion affects mitochondrial ATPase function, kinetoplast dependency, and drug sensitivity in
trypanosomes. Proc Natl Acad Sci U S A. 2015 Jul;112(29):9112–7.

70.   Giraud E, Martin O, Yakob L, Rogers M. Quantifying *Leishmania* Metacyclic Promastigotes
from Individual Sandfly Bites Reveals the Efficiency of Vector Transmission. Commun Biol. 2019
Feb 28;2(1):84.

71.   Santos VC, Nunes CA, Pereira MH, Gontijo NF. Mechanisms of pH control in the midgut of
Lutzomyia longipalpis: roles for ingested molecules and hormones. J Exp Biol. 2011 May
1;214(9):1411–8.

72.   Te Brugge V, Ianowski JP, Orchard I. Biological activity of diuretic factors on the anterior
midgut of the blood-feeding bug, *Rhodnius prolixus*. Gen Comp Endocrinol. 2009 May;162(1):105–
12.

73.   Vasanthakumar T, Rubinstein JL. Structure and Roles of V-type ATPases. Trends Biochem
Sci. 2020 Apr;45(4):295–307.

74.   Collins MP, Forgac M. Regulation and function of V-ATPases in physiology and disease.
Biochim Biophys Acta Biomembr. 2020 Dec 1;1862(12):183341.

75.   Vercesi AE, Moreno SN, Docampo R. Ca^{2+}/H^+ exchange in acidic vacuoles of *Trypanosoma*
brucei. Biochem J. 1994 Nov 15;304 (Pt 1)(Pt 1):227–33.

76.   Docampo R. Advances in the cellular biology, biochemistry, and molecular biology of
acidocalcisomes. Microbiol Mol Biol Rev MMBR. 2024 Mar 27;88(1):e0004223.

77.   Xu ZS, Li FJ, Hide G, Lun ZR, Lai DH. Vacuolar ATPase depletion contributes to
dysregulation of endocytosis in bloodstream forms of *Trypanosoma brucei*. Parasit Vectors. 2020 Apr
25;13(1):214.

78.   He C. Balancing nutrient and energy demand and supply via autophagy. Curr Biol CB. 2022
Jun 20;32(12):R684–96.

79.   Finbow ME, Harrison MA. The vacuolar H$^+$-ATPase: a universal proton pump of eukaryotes.
Biochem J. 1997 Jun 15;324 (Pt 3)(Pt 3):697–712.

80.   Maxson ME, Grinstein S. The vacuolar-type H$^+$-ATPase at a glance – more than a proton
pump. J Cell Sci. 2014 Dec 1;127(23):4987–93.

81.   Williams RA, Tetley L, Mottram JC, Coombs GH. Cysteine peptidases CPA and CPB are
vital for autophagy and differentiation in *Leishmania mexicana*. Mol Microbiol. 2006 Aug;61(3):655–
74.

82.   Volf P, Volfova V. Establishment and maintenance of sand fly colonies. J Vector Ecol. 2011
Mar;36(SUPPL.1).

83.   Badirzadeh A, Taheri T, Taslimi Y, Abdossamadi Z, Heidari-Kharaji M, Gholami E, et al.
Arginase activity in pathogenic and non-pathogenic species of *Leishmania* parasites. Jardim A, editor.
PLoS Negl Trop Dis. 2017 Jul 14;11(7):e0005774.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Figure Legends

Fig 1. Fitness of promastigote mutants *in vitro*.

(A) Overview of the experiment timeline for *in vitro* growth of promastigotes in culture. DNA sampling time-points are indicated by yellow arrows. Dark pink arrows denote dilutions, to (a) 1 x 10^6 parasites/ml and (b) 1 x 10^5 parasites/ml. (B) Growth profile of the masterpool of promastigote mutants over time. Data points are the average of three measurements; yellow dots indicate where gDNA was sampled. The dotted line indicates dilution of the cultures. (C) Trajectories of the average of normalised reads of the promastigote masterpool, relative to time-point "0 hours" (T_0). Red dotted line highlight relative barcode abundance of 1. Controls are shown in colour: dark blue, SBL1-5 parental cell lines; Cyan, ΔLPG1; Magenta, ΔIFT88; Yellow, ΔPF16. Grey, all other barcoded cell lines. (D) Volcano plot showing fitness scores against p-values of mutants from the promastigote masterpool after 144 hours of growth. Dashed lines demarcate fitness score thresholds of < 0.5 and > 2, and a significance threshold of $p < 0.05$. Barcodes meeting both threshold criteria are coloured black, non-significant values are grey. Controls are coloured as in D.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Fig 2. Fitness of promastigote mutants *in vivo*.

(A) Overview of the experiment timeline for sand fly infections. DNA sampling time-points are indicated by yellow arrows. (B) Fitness scores from the promastigote masterpool after 144 hours *in vitro* growth plotted against the fitness scores from all mutants 216 hours after infection of sand flies. Black dashed lines mark fitness score thresholds of < 0.5 and > 2. (C-F) Volcano plots showing fitness scores against p-values after 216 h (9 days PBM) in sand flies, separated by sub-pools (P1-P4). Dashed lines demarcate fitness score thresholds of < 0.5 and > 2, and a significance threshold of $p < 0.05$. Barcodes meeting both threshold criteria are colored black, non-significant are grey. Controls are shown in colour: Dark blue dots, SBL1-5 parental cell lines; Cyan, ΔLPG1; Magenta, ΔIFT88; Yellow, ΔPF16. Black numbers denote the abbreviated GeneIDs (LmxM.xx.xxxx) of selected mutants with the highest fitness changes.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Fig 3. Pool bottlenecks and required number of flies.

(A) Pool evenness. A small value indicates that the pool is dominated by a few barcoded cell lines with high cell counts. Conversely, a large value means that all cell lines have comparable cell counts. Normal values for pools of cell lines fall into the range 0.7-1.3. (B) Inferred per-fly population bottleneck sizes. Each marker shows the average number of cells that successfully established a detectable infection within a single sand fly for a given replicate, time point and pool according to our mathematical model. The dashed line shows the median population bottleneck size (50 cells per fly). (C) Number of flies used in each pool replicate vs. number of flies required to meet a 90% survival threshold. The solid black line shows the threshold number of flies needed as a function of the pool evenness such that 90% of cell lines survive on average if all cell lines share the same fitness. If this criterion is not met (grey shaded area), it becomes difficult to interpret low-proportion barcodes and extinction as indicators for fitness rather than random events. All pools except for the masterpool lie above the threshold, hence the masterpool was excluded from further analysis.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Fig 4. V-ATPase is required for differentiation and colonisation in the sand fly vector.

(A) Trajectories of the average of normalised reads for V-ATPase deletion mutants from the promastigote masterpool (grey lines), relative to T_0. Dark blue lines, SBL1-5 parental controls. (B) Barcode trajectories for V-ATPase deletion mutants in sand flies, normalised to the start of the experiment. Colour code as in A. (C) Schematic of V-ATPase pump with each subunit labelled. Subunit E, for which the null mutant was individually characterised, is highlighted in magenta. (D) Growth of parental (PAR, dark blue), V-ATPase V_1E null mutant (KO, magenta) and V-ATPase V_1E add-back (AB, grey) mutant promastigotes *in vitro*. After 3 days of continuous growth, cultures were diluted back to 1 x 10^5 parasites/ml and monitored for an additional 3 days. (E) Parasite abundance in the digestive tract of dissected sand flies infected with PAR, AB or KO parasite lines, assessed at 2 days PBM (left) and 9 days PBM (right). (F) Location of PAR, AB or KO parasite lines at 2 days PBM (left) and 9 days PBM (right). NI, non-infected; ES, endoperitrophic space; AMG, abdominal midgut; TMG, thoracic midgut; CAR, cardia; SV, stomodeal valve. (G) Percentage of promastigote morphotypes of PAR, AB or KO parasite lines observed in infected gut smears after 9 days PBM. (H) *Sherp* transcript quantification by qPCR for PAR, AB and KO parasite lines grown *in vitro*. Each point shows the abundance of s*herp* normalised to 18S rRNA, calculated from three replicate measurements. (I) Representative micrographs of promastigote morphotypes found in each cell line *in vivo* at 9 days PBM. Scale bar = 5 µm.

A V-ATPase subunit trajectories *in vitro*

B V-ATPase subunit trajectories *in vivo*

C V-ATPase / Subunit E

D Growth profile *in vitro*

E Parasite abundance *in vivo*

F Parasite location *in vivo*

G Promastigote morphotypes *in vivo*

H SHERP relative abundance

I Nectomonad / Leptomonad / Metacyclic