Identification of transporters essential for survival of *Leishmania* promastigotes

in the digestive tract of sand flies

Authors

Jovana Sadlova[1], Barbora Vojtková[1], Tomáš Bečvář[1], Ulrich Dobramysl[2,3], Sandro Möri[4], Çağla

Alagöz[4,5], Richard J. Wheeler[2,3], Petr Volf[1], Eva Gluenz[4,6*], Andreia Albuquerque-Wendt[1,4,6-8*]

Affiliations

[1]Department of Parasitology, Faculty of Science, Charles University, Prag, Czech Republic

[2]Medawar Building for Pathogen Research, Nuffield Department of Medicine, University of Oxford,

Oxford, United Kingdom

[3]Institute of Immunology & Infection Research, University of Edinburgh, Ashworth Laboratories,

Edinburgh, United Kingdom

[4]Institute of Cell Biology, University of Bern, Bern, Switzerland

[5]Graduate School for Cellular and Biomedical Sciences (GCB), University of Bern, Bern, Switzerland

[6]School of Infection and Immunity, University of Glasgow, Sir Graeme Davies Building, Glasgow,

United Kingdom

[7]Parasite Chemotherapy Unit, Swiss Tropical and Public Health Institute, Allschwil, Switzerland

[8]Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade Nova de

Lisboa, Lisbon, Portugal

*Corresponding authors

andreia.albuquerquewendt@swisstph.ch (AAW)

eva.gluenz@unibe.ch (EG)

Author contributions

Conceptualisation, AAW, EG

Methodology, UD, RJW, AAW, EG, JS, PV

Software, UD, RJW

Formal analysis, AAW, JS, UD, EG

Investigation, AAW, JS, UD, BV, TB, CA, SM

Resources, EG, PV

Data curation, AAW, EG, UD

Writing original draft, AAW

Writing review and editing, all authors

Visualisation, AAW

Supervision, AAW, RJW, EG, JS, PV

Project administration, EG, PV

Funding Acquisition, AAW, EG, RJW, PV

Competing Interest Statement:

49 The authors declare no competing interest. The funders had no role in study design, data collection and

50 analysis, decision to publish, or preparation of the manuscript.

52 **Keyword**s: Transportome, *Leishmania*, Sand flies, bar-seq, phenotypic screen, V-ATPase

Manuscript contents:

Main Text

Figures 1 to 4

Supplementary Figures 1 to 6

Supplementary File 1

Supplementary Tables 1 to 8

Abstract

Leishmania amastigotes ingested by female phlebotomine sand flies are exposed to a harsh and dynamic environment, markedly different from that of their mammalian host. Within the sand fly's alimentary tract, these parasite forms encounter shifts in temperature, pH and nutrient availability, which trigger significant morphological and physiological adaptations. Membrane transporter proteins, channels and pumps play a crucial role in facilitating the movement of solutes across eukaryotic membranes. Previously, a systematic loss-of-function screen of the *L. mexicana* "transportome" identified forty transporter deletion mutants that caused significant loss of fitness in macrophage and mouse infections. Here, using an independently generated library of over 300 barcoded gene deletion mutants, we monitored their growth fitness for seven days *in vitro* and tested which transporters are required for *Leishmania* promastigotes to successfully colonise *Lutzomyia longipalpis* sand flies for nine days. Overall, fitness scores correlated between promastigotes from long-term *in vitro* culture and *in vivo* sand fly infections. More importantly, for 34 mutants, a significant loss of fitness was observed exclusively *in vivo*. Moreover, deletion of the vacuolar H$^+$ ATPase (V-ATPase) proved detrimental for parasite persistence and promastigote differentiation in the sand fly, uncovering a key role for the V-ATPase at different stages throughout the *Leishmania* life cycle.

Author Summary

Leishmania parasites cause leishmaniases - a group of neglected tropical diseases that affect millions of people worldwide. These parasites must survive in two radically different environments: inside a mammalian host and within the gut of a blood-feeding sand fly. To thrive in the sand fly, *Leishmania* undergo extensive physiological changes and depend on transporter proteins to move nutrients and other molecules across their cell membranes. In this study, we focused on identifying which of these transporters are critical for the parasite's survival inside the sand fly. We used a genetically engineered library of *Leishmania* promastigotes - the parasite form adapted to the insect vector - to assess the importance of more than 300 different transporter genes. We discovered that 34 of these transporters are essential for successful colonization of the sand fly. Among them, one key protein complex - the

87 vacuolar H^+ ATPase (V-ATPase) pump – was found to be crucial for parasite survival in the insect

88 vector. Our findings deepen our understanding of how *Leishmania* adapts to life within the sand fly and

89 highlight potential molecular targets for disrupting its transmission.

Introduction

Leishmania (Kinetoplastida: Trypanosomatidae) are unicellular eukaryotic protozoa and the causative

agents of leishmaniases (1), a group of neglected tropical diseases. Over 20 *Leishmania* species share a

digenetic life cycle, alternating between an insect vector and a mammalian host (1). Females of more

than 90 species of phlebotomine sand flies - *Phlebotomus* in the Eastern Hemisphere and *Lutzomyia* in

the Western Hemisphere - serve as their primary vectors (2).

When a female sand fly takes a blood meal from an infected mammalian host, it ingests immotile

amastigote forms, which are encased in a chitin-rich peritrophic matrix (PM) (2). During this stage,

parasites face dramatic environmental changes, including a temperature drop (from ~37 °C to ~26 °C),

pH shift (from acidic to neutral/alkaline), and altered nutrient availability (from blood components to

sugar meals and microbiome metabolites). These signals trigger rapid differentiation from amastigote

into promastigotes, often within 6 hours post-feeding (3,4).

Inside the sand fly gut, *Leishmania* promastigotes undergo several differentiation stages. While stage-

specific transcriptional profiles have recently been described for *L. major* isolated from *Phlebotomus*

duboscqi guts(4), a definitive set of molecular markers for each *L. mexicana* promastigote morphotype

is currently lacking. Morphology, location and physiology are therefore still widely used to distinguish

different promastigote developmental stages in the sand fly. The initial amastigote to procyclic

promastigote differentiation is marked by key metabolic changes, including a ~10-fold increase in

uptake of carbon sources (e.g., glucose and non-essential amino acids), and elevated secretion of

glycolytic end-products (5–8).

These weakly motile, short-flagellated procyclic forms undergo binary fission for at least 48-96 hours

before slowing down their replication and differentiating into highly motile elongated nectomonad

promastigote forms (9).

After 72-96 hours post blood feeding, the toxic products of blood digestion cause a reduction in the number of procyclic forms (10) and enzyme-mediated PM disintegration ensures that nectomonads escape the PM-encased blood meal into the midgut lumen (9,11). There, they bind to the midgut epithelium via parasite- and vector-derived surface molecules such as lipophosphoglycans (LPGs) or mucin-like O-glycoconjugates, which prevent their expulsion during defecation (12–15).

Parasites of the *Leishmania* subgenus then differentiate into replicative leptomonad promastigotes and migrate to the anterior midgut (3). During this phase, flagellar motility is crucial for *Leishmania* parasites to migrate through the thoracic midgut toward the stomodeal valve, the structure at the junction between the sand fly foregut and midgut – a process required for transmission. For example, Beneke et al. (2019) used genetically engineered *L. mexicana* promastigotes with impaired flagellar motility to infect *Lu. longipalpis* and observed that these species require directional motility to successfully colonise the fly (16). Furthermore, Cuvillier et al. (2003) showed that overexpression of constitutively active variant of the ADP-ribosylation factor-like protein 3A (ARL-3A) in *L. amazonensis* promastigotes resulted in cells with a short, non-motile flagellum, which failed to colonise *Lu. longipalpis* sand flies (17).

When reaching the stomodeal valve, parasites undergo terminal differentiation into two morphologically distinct forms: replicative, short-flagellated, non-motile haptomonads and non-replicative, long-flagellated highly motile metacyclic promastigotes (18,19). Notably, recent single cell transcriptomic evidence from *L. major* promastigotes isolated from *Ph. duboscqi*, suggest that metacyclic promastigotes may be further divided into two transcriptionally distinct sub-forms; replicative early metacyclics and non-replicative late metacyclics (4). Moreover, the same study provides convincing evidence that in addition to metacyclics, which are traditionally viewed as the primary infective stage, haptomonads also play a significant role in transmission (4).

In mature sand fly infections, parasites secrete promastigote secretory gel (PSG) - a viscous matrix rich in filamentous proteophosphoglycans (fPPGs) - which fills the thoracic midgut (20–24). In addition, they secrete chitinase (25), which damages the insect´s alimentary canal (26). Along with fPPGs, this disruption alters sand fly feeding behavior and promotes regurgitation during subsequent blood meals

(12,27,28). Collectively, these changes enhance parasite transmission by increasing the number of

parasites egested into the skin of the mammalian host (28).

Although our understanding of the *Leishmania* life cycle within sand flies still lags behind that of other

vector-borne diseases, several molecular determinants of vector colonisation have been identified

(3,12,16,24,29–42). Recent advances in reverse genetics and high-throughput barcode sequencing (bar-

seq) in *Leishmania* sp. (43) have significantly accelerated discovery of parasite genes essential for

promastigote fitness and vector colonisation (16,32,44). For instance, proteins required for flagellar

assembly (IFT88, LmxM.27.1130) and motility (e.g. the central pair protein PF16, LmxM.20.1400 or

the inner dynein arm protein IC140, LmxM.27.1630) (16) were found to play a crucial role in persistence

in sand flies and migration to the stomodeal valve (17). In a separate screen targeting the parasite's

kinome, ATM (LmxM.02.0120) and PI4K (LmxM.33.3590), were identified as atypical (aPK) and

phosphatidylinositol 3' kinase-related (PIKK) protein kinases, respectively, conditionally essential for

survival only in sand flies, suggesting unique pathways are involved in vector-stage survival (32).

Another kinase, MPK9 (LmxM.19.0180), was identified as essential for sand fly colonisation (32). In

an independent study, MPK9 was shown to influence flagellar length (45), reinforcing the importance

of flagellum integrity for the parasite survival inside the vector (45). Moreover, single-cell RNA

sequencing is beginning to reveal molecular markers of parasite development in insect stages (4,46).

Despite these advances, the role of transporter proteins in sand fly colonisation remains largely

underexplored. Exceptions include two nucleotide sugar transporters involved in lipophosphoglycan

(LPG) biosynthesis. One, LPG2 (LmxM.33.3120), encodes a GDP-mannose transporter responsible for

incorporating the initial and repeating mannose units into the mannose-rich LPG structure and was

shown to be essential for development of *L. donovani* and *L. major* in several sand fly species, including

Ph. argentipes, *Ph. papatasi*, *Ph. duboscqi* and *Ph. perniciosus* (31,33,35,47). Another, LPG5A/B gene

array (LmxM.24.0360-65 3120), encodes a UDP-galactose transporter, results in reduced colonisation

of *L. major* in *Ph. duboscqi* (35). This transporter incorporates galactose units into the same repetitive

LPG backbone, highlighting the importance of glycan modifications in vector attachment and

colonisation. This is further supported by ablation of additional (non-transporter) genes involved in LPG

biosynthesis (15,16,31,47). Interestingly, LPG mediated binding between sand flies and *Leishmania* is

known to be crucial in parasite-vector systems where the vector is specific, i.e. each glycoconjugate presented by each different *Leishmania* sp. enables extraordinary specificity to a single vector species (48,49). In contrast, permissive vectors such as those within the genus *Lutzomyia*, have been reported to present additional binding mechanisms involving O-glycosylated proteins (50).

Recently, we systematically assessed the fitness contribution of 312 predicted *L. mexicana* membrane transporters, channels and pumps in promastigotes in culture (*in vitro*) during exponential phase of growth, in amastigotes in human induced pluripotent stem derived macrophages (iMACs) and over 6 weeks in mice (*in vivo*) (44). Using bar-seq, we showed that deletion of at least 40 transporters compromised amastigote survival *in vivo* (44). The vacuolar H^+ ATPase (V-ATPase) emerged as a crucial proton pump for the survival of parasites *in vivo*, and *in vitro* under conditions of low external pH (44). Here, we extend that work by conducting an independent comprehensive systematic loss-of-function screen targeting 316 single putative transporter-encoding genes and 17 gene arrays. We assessed mutant fitness over a one-week time course *in vitro* and in a sand fly model of infection *in vivo*. This screen revealed some mutants that show gain-of-fitness phenotypes and many with loss-of-fitness phenotypes. While there was a positive correlation between mutant fitness *in vitro* and in the flies, these results indicate a vital function for ion pumps, sugar nucleotide transporters, and transporters of some other classes, notably several mitochondrial carrier proteins, for survival and fitness within their sand fly vector. Moreover, the V-ATPase is required for effective completion of the developmental cycle from promastigotes to metacyclics *in vivo*.

Results and Discussion

An expanded gene deletion screen of the L. mexicana *transportome reveals that most transporters are dispensable for promastigote survival* in vitro

We previously reported the identification *L. mexicana* transportome, comprising of 312 putative membrane transporters, channels and pumps, and their functional evaluation in the mammalian host parasitic stage in a gene deletion fitness screen (44). Here we expanded on this by studying the fitness phenotypes of promastigote forms under prolonged *in vitro* culture and in sand fly infection assays *in*

vivo. For this screen, we generated arrayed CRISPR/Cas9 mutant libraries targeting the previously

identified "transportome" (44) plus four newly identified genes: two from the Acetate Uptake

Transporter (AceTr) family, one with similarity to the Selenoprotein P Receptor (SelP-Receptor) family,

and one Multidrug/Oligosaccharidyl-lipid/Polysaccharide (MOP) Flippase, as classified by the

Transporter Classification Database (TCDB) (51), thus expanding the number of proteins in the *L.*

mexicana "transportome" to 316 (Supplementary Table 1) (16,43,44,52). This approach successfully

generated 304 viable mutant populations, each resistant to both Blasticidin and Puromycin selection

drugs (Supplementary Tables 2,3), broadly organised by TCDB families into four sub-libraries

(Supplementary Table 4). Diagnostic PCR confirmed the successful deletion of all copies of the targeted

genes for 154 lines (null mutants); the remaining 132 lines where the targeted gene was still detectable

were classified as 'refractory to deletion' (Supplementary Table 3). Of the confirmed single-gene

deletions, 46 had not been successfully generated in our previous screen (44). This new mutant library

also identified three additional single-member superfamilies whose transporters appear dispensable *in*

vitro, namely the mitochondrial EF hand Ca^{2+} uniporter regulator (MICU; *LmxM.07.0110*), the Proton-

dependent Oligopeptide Transporter (POT; *LmxM.32.0710*) and the Selenoprotein P Receptor (SelP-

Receptor; *LmxM.28.2380*) in addition to the eight superfamilies that were previously shown to be

dispensable (44).

Deletion of genes arranged in tandem arrays

This screen also expanded the analysis of transporter genes in tandem arrays, targeting a total of 17

arrays, including nine arrays not previously targeted, from the following families: AceTr (1 array),

Amino Acid/Auxin Permease (AAAP, 4 arrays), Cyclin M Mg^{2+} Exporter (CNNM, 1 array),

Equilibrative Nucleoside Transporter (ENT, 1 array), Major Facilitator (MFS, 2 arrays), Mitochondrial

Carrier (MC, 4 arrays), P-type ATPase (P-ATPase, 1 array), Voltage-gated Ion Channel (VIC, 2 arrays),

Zinc (Zn^{2+})-Iron (Fe^{2+}) Permease (ZIP, 1 array) (Supplementary Table 1). Upon analysis of the 13 drug

resistant mutants that survived the selection, only one AAAP array mutant (*LmxM.34.5350* and

LmxM.34.5360) was found to be null (Supplementary Table 1 and 3). For the *LmxM.18.1290* and

LmxM.18.1300 array where a null mutant was previously achieved (44), only double puromycin,

blasticidin resistant mutant populations retaining at least one copy of the targeted gene were recovered

(Supplementary Table 1 and 3, Supplementary Figure 2). As discussed previously (44), technical

challenges such as high sequence similarity and underestimated gene copy numbers can hinder the

isolation of null mutants from tandem arrays. For instance, at first, we failed in the isolation of a null

mutant for the glucose transporter array which harbours three genes (LmGT1-LmGT3), despite its

successful deletion in *L. mexicana* using a strategy of gene knockout by homologous recombination

(53,54). In this case, we repeated transfections using both puromycin and blasticidin selection, but only

double drug-resistant populations emerged that had retained at least some LmGT genes. It was only

after a third round of transfection and subsequent selection of clonal cell lines that we successfully

isolated three clones lacking the entire array (Supplementary Figure 5A-C). The doubling times of two

null mutant clones were measured and found to be significantly increased (8.73 and 8.24 h) compared

to that of the parental cells (5.14 h) (Supplementary Figure 5D-E). This suggests that although the GT-

array null mutants are viable, mutants that somehow retained one or several of the genes from the

targeted array may have a significant growth advantage in mixed populations. These data show that

while it is possible to achieve null array mutants, the technical challenges in identifying and isolating

array mutants precludes phenotype screens at scale with this bar-seq method.

Less than 30% of the Leishmania *transportome is essential for promastigote survival*

Consolidating data across the two independently generated libraries indicates that 225 (~71%)

transporter-encoding genes are dispensable for promastigote survival in standard *in vitro* laboratory

cultures (Figure 1; Supplementary Table 3; Supplementary Figure 1) (44). The successful deletion of

these genes is positive proof that they are not essential for cell proliferation under the tested conditions,

although they may still contribute to fitness. Conclusive statements cannot be made however about the

importance of genes where a deletion attempt failed. For the 91 genes refractory to deletion in two

independent screens (Supplementary Table 3), further attempts at gene deletion may yet prove

successful. Data released from the LeishGEM genome wide gene deletion screen

(https://browse.leishgem.org/) (55) already reports several transporter gene deletion mutants that were

not retrieved in the current screen. Examples include transporter-encoding genes located on

supernumerary chromosomes like the putative acidocalcisome inositol 1,4,5-triphosphate receptor/Ca^{2+}

release channel (LmxM.16.0280), one of the four amino acid transporters of the AAT1 locus, AAT1.4

(LmxM.30.0350), a glycosomal ABC transporter, GAT1 (LmxM.30.0540) and a major facilitator

(LmxM.30.0720); on diploid chromosomes a porphyrin transporter (LmxM.17.1430), the amino acid

transporter, AAT23.2 (LmxM.27.0680) and a mitochondrial carrier (LmxM.29.2240) (Supplementary

Table 3). Conditional gene knockout strategies would be required for conclusive functional validation

and stronger support for a claim of "essentiality" (56).

Prolonged culturing of transporter deletion mutants identifies novel growth fitness phenotypes in vitro

We next asked whether gene disruption had any effect on the relative growth rates of the surviving

promastigotes in standard laboratory cultures, over a one-week time course. To assess their relative

fitness, all 304 viable isolated barcoded transporter mutants and 13 array mutants, were combined into

a single masterpool. To this pool we added five barcoded parental control lines (SBL1-5), eight non-

transporter null mutants with previously characterised phenotypes [three independently barcoded

Δ*LPG1* (*LmxM.25.0010*, normal growth, important for sand fly colonisation), two independently

barcoded Δ*PF16* (*LmxM.20.1400*, normal growth, essential for sand fly colonisation), three

independently barcoded Δ*IFT88* (*LmxM.27.1130*, very slow growth, essential for sand fly colonisation)

(17), and one non-transporter null mutant Δ*LmxM.15.0240* (nonspecific lipid-transfer protein) (44)

(Supplementary Table 4). This masterpool was split into three separate flasks and grown in standard

M199 culture medium for seven days. Cultures were diluted into fresh medium twice during this period

(Figure 2A) to maintain the populations in the exponential phase of growth (Figure 2B). DNA was

sampled at baseline (0 hours), and after 24, 48, and 144 hours (Figure 2) and each mutant's relative

representation over time was assessed by measuring DNA barcode abundance at the sampled time points

and calculating the proportion of each barcode at a given time point relative to its representation in the

starting population (Supplementary Table 5). This showed that the parental control cells and a majority

of the mutants maintained a flat trajectory, indicating that they proliferated in the population at similar

rates. There were a small number of mutants with an upwards trajectory, indicating their proportion

within the pool increased over time, and a larger set where barcode proportions sharply decreased over

time (Figure 2C, Supplementary Table 5, Supplementary Figure 3,4). To quantify this, fitness scores

were calculated by comparing the change in barcode proportions for a given mutant to that of the mode

of the cell line change distribution (Supplementary Table 5). After 144 hours, seven mutants displayed

enhanced fitness in the promastigote *in vitro* pool (fitness score above 2 and $p < 0.05$, Figure 2D): two

mutants lacking a mitochondrial carrier (Δ*LmxM.15.0120* and Δ*LmxM.34.3330*), two ABC transporters

(Δ*LmxM.06.0100* and Δ*LmxM.11.1290*), two amino acid permeases (Δ*LmxM.30.1820* and

Δ*LmxM.27.0680*) and one hypothetical protein of the MSF family (Δ*LmxM.34.2810b*). In contrast, 107

transporter mutants exhibited significantly reduced fitness (score below 0.5 and $p < 0.05$, Figure 2D),

with two barcodes dropping below the detection limit at 48 h and eleven at 144 h (zero read counts in

all three replicates, Supplementary Table 5). Amongst the mutants disappearing rapidly from the

population was a confirmed deletion of ABCB3, which acts in heme and cytosolic iron/sulfur clusters

biogenesis and is required for *L. major* virulence (57). This severe loss of fitness in culture may explain

why previous attempts to generate null mutants for this transporter were unsuccessful (44,57). Still

detectable at the lowest levels were confirmed null mutants for a predicted sodium/hydrogen exchanger

of the CPA1 family (Δ*LmxM.14.0980* score 0.014, p=0.0001) and a putative calcium-transporting P-

ATPase (Δ*LmxM.32.1010*, score 0.009, p=0.004). Also significantly depleted were mutants for

predicted ADP/ATP carrier proteins: Δ*LmxM.07.0530* (MCP15, refractory to deletion, MCP

nomenclature taken from (58)) and the tandem array *LmxM.19.0200_LmxM.19.0210* (MCP5, refractory

to deletion). The deletion of the fourth predicted ADP/ATP carrier predicted in the *L. mexicana* genome

(LmxM.14.0990, MCP16) resulted in a less severe but also significant loss of fitness (score 0.17 and *p*

= 0.0016), indicating that these mitochondrial carrier proteins perform vital non-redundant functions in

promastigotes.

The culture conditions were designed to provide ample nutrients, a buffered environment and constant

temperature. However, over the 144 hours, cells experienced changes in population density and two

culture dilutions, thereby possibly being exposed to a variety of stresses, including microenvironmental

pH shifts, nutrient depletion, and waste metabolite accumulation. These data show that, although viable,

many of the null mutants were less fit than the parental cell lines and would be outcompeted over time

in mixed populations.

Loss and gain of fitness in transporter deletion mutants in vivo

Leishmania promastigotes naturally live in the alimentary tract of phlebotomine sand flies and we next

asked which mutants would be able to tolerate this more varied and harsher environment. To evaluate

the relative fitness of all viable transporter mutants *in vivo*, we created four sub-pools (named P1-P4)

which were added separately to blood feeds presented to female *Lutzomyia longipalpis* sand flies. Each

pool contained five barcoded parental control lines (SBL1-5), three non-transporter null controls with

established *in vitro* and *in vivo* promastigote phenotypes (ΔLPG1, ΔPF16, ΔIFT88), and on average 86

barcoded mutants per pool. We reasoned that this smaller pool size would reduce the chance of mutants

being lost at random considering the small volume ingested by the sand flies. Assuming that in

experimental conditions each *Lu. longipalpis* female feed 0.8-1 µl of blood(59), each fly would be

expected to ingest 16´000-20'000 if the blood-cell suspension was prepared at 2×10^7 parasites/ml,

guaranteeing an average of 186-233 parasites per mutant line, from a pool of 86 barcoded mutant lines

– a level within the range used in comparable bar-seq studies (16,32). DNA was collected from the

parasite-blood mixture at 0 hours (pre-infection) and from infected sand flies after 2 days (48 h) and 9

days (216 h) post blood meal (PBM) (Figure 3A). The barcode proportions for each mutant at each time

point were quantified by sequencing. In each cohort, the parental control lines remained stable over the

9-day infection (Supplementary Figure 4C-F), with many of the mutant barcodes following the same

trajectory as the parentals, indicating no fitness change. The trajectories of the control mutants Δ*IFT88*,

Δ*LPG1*, and Δ*PF16* indicated depletion of these mutants, as expected (16), albeit with some variation

between the different pools (Supplementary Figure 4C-F). To quantify these changes, fitness scores

were calculated to identify mutants that became significantly depleted or enriched in the flies

(Supplementary Table 5). The fitness scores of the mutants 9 days PBM in flies showed a positive

correlation (Pearson r = 0.6055) with the fitness scores of promastigotes measured after 144 h in culture

(Figure 3B). Across all sub-pools, 80 barcoded transporter mutants displayed significantly reduced

fitness (score < 0.5, $p < 0.05$), while nine mutants showed increased fitness (score > 2, $p < 0.05$) in the sand fly (Figure 3C-F).

There was a small number of mutants that showed a loss-of-fitness phenotype only in the fly (Figure 3B), including a UDP-Gal nucleotide sugar transporter (LmxM.22.1010 HUT1L), one amino acid transporter (LmxM.32.1420), two ABC transporters (LmxM.32.3260 ABCI4 and LmxM.32.1800 ABCD3, the ortholog of the *T. brucei* glycosomal transporter GAT2(60), one uncharacterised MSF transporter (LmxM.01.0440), a putative calcium motive p-type ATPase (LmxM.34.2080) and a subunit of the V-ATPAse (discussed below). While the substrates of most of these transporters remain to be defined, they may contribute to the parasite's metabolic adaptation to changing environments in the fly, or to the secretion of glycoconjugates (e.g. the HUT1L mutant). The importance of glycoconjugate secretion *in vivo* is supported by the phenotype of the control mutant *ΔLPG1*, which became strongly depleted in the flies. Loss of the Golgi GDP-Man transporter LPG2, which lacks a broader range of glycoconjugates including LPG (33,35,47,61), also resulted in mutants with low fitness scores in flies (16), as well as in *in vitro* cultures, although these did not pass the statistical significance test.

The gain-of-fitness mutants included three ABC transporters; ABCG3 (LmxM.06.0100), ABCH1 (LmxM.11.0040) and ABCA6 (LmxM.11.1290), one MFS protein (LmxM.34.2810), one UDP-galactose transporter (LmxM.24.0365), aquaglyceroporin 1 (AQP1, LmxM.30.0020), two folate-biopterin (FBT) transporters; FT1 (LmxM.10.0400) and LmxM.19.0920 and one voltage-gated calcium channels (VGCC) of the VIC family (LmxM.17.1440). The latter encodes one of two L-type VGCCs in *Leishmania*, previously shown to be sensitive to VGCC inhibitors (62). Interestingly, *ΔLmxM.17.1440* promastigotes also exhibited significantly increased fitness *in vitro* after 24h and 144 h (this study and (62)) while their abundance decreased in macrophages at 120 h (score = 0.5, $p < 0.05$ (62)). In contrast, the second VGCC (LmxM.33.0480) showed consistently reduced fitness both *in vitro* and *in vivo*, suggesting that these VGCCs have distinct roles in calcium homeostasis across life cycle stages. Similarly, stage-specific phenotypes were observed for the two predicted magnesium transporters of *L. mexicana*. Mutants where MGT2 (LmxM.25.1090) was targeted, but only double drug-resistant populations, refractory to gene deletion were isolated, resulted in decreased fitness in sand flies (score 0.01, p=0.008), while the MGT1 mutant (*ΔLmxM.15.1310*, null) was enriched in the flies 9 days PBM.

An MGT1 null mutant was previously found to have significantly reduced fitness in mice (44). Prior

work suggests AQP1 may function in volume regulation, osmotaxis and antimony [Sb(III)] uptake (63),

although its broader biological role remains unclear. While direct measurements sand fly gut or

macrophage phagolysosome osmolality are unavailable, estimates suggest values range from 85-448

mOsm/kg in a hematophagous insect gut (64) and 275-295 mOsm/kg in mammalian blood (65). The

phagolysosome is likely mildly hyperosmotic due to ion influx, digestion and acidification. How these

osmotic shifts across host environments are buffered in the absence of AQP1 remains to be investigated.

Similarly, the phenotypes of the FBT family mutants requires further study. Folate transporter FT1

(LmxM.10.0400) is known to be highly expressed in actively dividing promastigotes (66,67).

Leishmania, like many other eukaryotic cells, are folate (vitamin B9) auxotrophs, requiring external

folate uptake. Host folate availability can vary widely depending on nutritional status and microbiota,

implying that *Leishmania*'s 13 FBTs (Figure 1B) may be specialised for stage-specific roles. The mutant

for the FBT family biopterin transporter BT1 (Δ*LmxM.34.5150*) was significantly depleted from flies 9

days PBM (score 0.003, p=0.02) and completely lost from the promastigote *in vitro* culture.

Whether the higher fitness scores reflected more rapid proliferation or better survival or persistence in

the fly following excretion of the digested bloodmeal cannot be deduced from these data. Similarly, a

loss of barcodes from the population could indicate a higher death rate or slower proliferation. In the *in*

vitro assay, exponential growth of the population was precisely measured, showing a rate of 20.3

doublings over the 144 h time course (Supplementary Table 6). In the fly, the promastigotes normally

progress through a series replicative and non-replicative developmental stages. How many exact

doublings a wild type *L. mexicana* promastigote is expected to undergo during 9 days in *Lu. longipalpis*

is not precisely determined, but it is likely to be fewer than in the constant environment of a culture flask

(12). Previous reports, suggest that promastigotes may take ~ 29 h (Supplementary Table 6) to double

in the digestive tract of a female sand fly (12). However, this is a very rough estimate, since factors like

parasite death or loss from the fly during defecation of bloodmeal remnants, are likely the dominant

reasons for severe dropouts in the sand fly. Furthermore, the used standard M199 culture medium in this

study, is a high glucose medium routinely used for *in vitro* growth to maximise the growth rate and

density of *Leishmania* promastigotes, likely differing significantly from the natural sand fly gut

environment and its energy sources. Thus, small differences in growth rate would be hard to detect in the fly, but easily detectable in culture (12). Likely as a result of that, we observed a greater number of mutants with small but significant changes in fitness *in vitro* (107/304) compared to *in vivo* in sand flies (80/304) (Figure 3B, Supplementary Table 5). We can speculate that this reflects differences between the assays, where the *in vitro* populations underwent a larger number of doublings, combined with the technical advantage of sampling larger amounts of DNA from the *in vitro* cultured cells, allowing for the reliable detection of small differences in replication rate, which may not be detectable in the fly assay.

The V-ATPase of L. mexicana *is critical for survival and parasite differentiation in the sand fly gut*

Vacuolar proton ATPase (V-ATPase) pumps are multi-subunit protein complexes that acidify intracellular organelles by translocating protons across membranes. This complex was already shown to be important for the regulation of endocytosis in *T. brucei* bloodstream forms (68). We have recently demonstrated, that V-ATPases in *Leishmania* localise to a crescent-shaped region near the flagellar pocket (44) and while the loss of V-ATPase subunits had little effect on the growth of promastigotes in standard culture medium at neutral pH, it proved detrimental to promastigotes in acidified culture medium, and caused significant loss of fitness in macrophages and mice (44).

Here, the barcode trajectories of mutants lacking various V-ATPase subunits indicated a moderate decline after 48 hours of promastigote *in vitro* growth (Figure 4A) when the cells had reached a density of $>1 \times 10^7$ cells ml^{-1} (Figure 2B); at that density, dilution into fresh medium was required to maintain the population in log phase. Despite the decline in abundance, the barcodes of V-ATPase mutants were still represented within the pool after 144 h of continued exponential growth. In the sand flies, the decline was more pronounced within just 48 h PBM (Figure 4B), when the infected bloodmeal was still surrounded by peritrophic matrix, and declined further at 9 days PBM, resulting in lower fitness scores than observed *in vitro* (Supplementary Table 5).

To investigate this phenotype further, we introduced an ectopic copy of the V-ATPase subunit E (V$_1$E, Figure 4C) into the V$_1$E null mutant cell line (Δ*LmxM.36.*3100, KO) to generate an addback control

(AB) and compared the growth profiles of the three cell lines *in vitro* and *in vivo* (Figure 4D). The

growth rates of the KO, AB and parental cell lines were measured over 6 days, with one dilution at the

end of log phase on day 3. All three cell lines grew at a comparable rate for the first three days; after

dilution the null mutant cell line continued exponential growth but with a slightly longer doubling time.

The growth of the AB cell line was indistinguishable from that of the parental control (Figure 4D). We

next infected female *Lu. longipalpis* flies with the KO, AB or the parental *L. mex* Cas9 T7 cell line to

measure parasite abundance, location, and developmental stages over time (Figure 4 E-G). At 48 h PBM,

100% of flies infected with parental and AB lines showed heavy infections (> 1000 promastigotes/gut,

Figure 4E) and parasites were located within the endoperitrophic space surrounded by the peritrophic

matrix (Figure 4F). In contrast, the KO persisted in only 40 % of flies, and infections were mostly

moderate (100-1000 promastigotes/gut). By day 9 PBM, most flies infected with parental and AB lines

developed heavy infections with colonization of the stomodeal valve in 97 % of cases. In contrast, the

KO only established moderate or light infections, which were confined to the abdominal or proximal

thoracic midgut, and did not reach the cardia (Figure 4E,F).

Quantitative analysis of promastigotes at 9 days PBM showed significant differences in the distribution

of parasite morphotypes between KO, AB and parental cell lines. Elongated cells morphometrically

classified as "nectomonads" predominated in the KO, whereas leptomonads were less abundant, and

metacyclic promastigotes were completely absent when compared to AB and parental lines (Figure 4G,

Supplementary Table 7). Additionally, the KO exhibited a significantly longer body and flagella (16

µm, p=0.000; 16.48 µm, p=0.005) compared to the parental cells (10.67 µm; 14.91 µm) (Supplementary

Table 7). Expression of an episomal copy of the deleted gene in the KO, significantly restored body

length (11.57 µm, p=0.000), but not flagellum length (16.11 µm, p=0.747).

In laboratory cultures, promastigotes in stationary-phase culture may encounter metabolic stress as

cultures grow dense, including waste accumulation, nutrient depletion, and cell crowding, which may

cause the longer doubling times of the V-ATPase mutants (Figure 4A,D). In the sand fly, these

challenges are intensified by competition with the host and its microbiota for limited resources and

change of the external pH in the sand fly gut, which shifts from ~6 in unfed or sugar-fed insects to ~8.15

following a blood meal (69). Blood ingestion also triggers diuresis, a process in which hematophagous

insects, including sand flies, expel excess water and concentrate ingested blood, facilitating more blood uptake. This process is mediated by water absorption in the midgut and urine production by the malpighian tubules (69,70). Our results suggest that the V-ATPase may contribute to parasite fitness under these different stresses, beyond adaptation to a low pH environment. This could occur by regulating endocytosis, endo-/lysosomal trafficking, vesicle fusion, protein degradation, and autophagy (71–73) a pathway that has been shown to be important for the differentiation to amastigotes and metacyclic promastigotes *in vitro* (74). Here we show that the differentiation of V-ATPase mutants was delayed or prevented in the insect vector. Taken together, the mutant phenotypes *in vitro*, in the insect vector and in a mammalian host (44) identify the V-ATPase as being key to the parasite's ability to adapt to changing environments at every stage in its life cycle.

Materials and Methods

Leishmania parasites

Promastigote forms of the *L. mexicana* cell line *L. mex* Cas9 T7 (52) and all generated mutants in this study were either grown in T25 cm^2 flasks at 27 °C or flat bottom well plates at 27 °C + 5 % CO$_2$ in filter-sterilised M199 medium (Life Technologies) supplemented with 2.2 g/L NaHCO$_3$, 0.005% hemin, 40 mM 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES) pH 7.4 and 10 % FCS [referred in the text as "standard M199"]. 50 μg/ml Nourseothricin Sulphate and 32 μg/ml Hygromycin B were added to the medium for the maintenance of the *sp*Cas9 and T7 RNA polymerase transgenes (52).

Phlebotomine sand flies

A laboratory colony of *Lutzomyia longipalpis* (originating from Jacobina, Brazil) was maintained in the insectary of the Charles University (Prague, Czechia) under standard conditions (at 26 °C fed on 50% sucrose solution with a 14 h light/10 h dark photoperiod) as described previously (75). The use of laboratory mice for sand fly breeding has been approved by the Ministry of Education, Youth and Sports number MSMT-25062/2023-6. Mice were kept in the animal facility of Charles University in Prague in

accordance with institutional guidelines and Czech legislation (Act No. 246/1992 and 359/2012 coll. on

protection of animals against cruelty in present statutes at large), which complies with all relevant

European Union and international guidelines for experimental animals.

Revised transportome of Leishmania mexicana

This study included the genes that were previously reported in the 'TransLeish' mutant screen (44), plus

four newly identified putative membrane transporter genes: LmxM.03.0370 and LmxM.03.0390,

encode for proteins of the The Acetate Uptake Transporter (AceTr) Transporter Classification Data Base

(TCDB) family (51), LmxM.28.2380, belonging to the Selenoprotein P Receptor (SelP-Receptor)

family and LmxM.28.2410, encoding a protein of the Multidrug/Oligosaccharidyl-lipid/Polysaccharide

(MOP) Flippase TCDB family, thus updating the current size of the *L. mexicana* transportome to a total

of 316 putative members.

CRISPR-Cas9 gene knockouts

Gene deletions were done using the CRISPR-Cas9 barcoding method previously described (52).

Diagnostic PCRs for the validation of gene deletions was done as reported in (44) using ORF_Fw and

ORF_Rv primers (Supplementary Table 2). In addition to targeting each gene individually, a total of 17

tandem arrays were targeted and 8 non-transporter null mutant control cell lines were produced [three

independently barcoded Δ*LPG1*, two independently barcoded Δ*PF16*, and three independently barcoded

Δ*IFT88* (Supplementary Table 1). Fitness screens were done with populations for which gene deletions

were assessed by diagnostic PCR, without further subcloning. Exceptionally, for the deletion of the

glucose transporter array (LmGT1-GT3), new primers were designed that captured the dissimilar UTR

regions flanking the GT array. Drug resistant mutants were cloned by limiting dilution and ORF

verification primers for validation of resulting null mutant clones were also redesigned, so that all three

copies could be recognised (Supplementary Figure 5).

Generation of V-ATPase subunit E add-back cell line

The LmxM.36.3100 gene was cloned into the pT-add plasmid via restriction digestion cloning

(Supplementary File 1). Restriction sites *Spe*I and *EcoR*I were inserted into the 5' and 3' end of the

gene, respectively, with the following primers:

SubE_forward_SpeI 5' TCAAGACTAGTATGAGCGAGGCACGCCAAAT 3'

SubE_reverse_EcoRI 5' AATACAGAATTCTTACAGTGGCGCCTCGGTGT 3'

The ΔLmxM.36.3100 cell line was transfected with 5 µg of the newly generated pTadd-LmxM.36.3100

plasmid as described elsewhere (76). After approximately 15 hours following transfection, drug resistant

cells were selected by addition of phleomycin at a final concentration of 25 µg/ml and cells were kept

in presence of drug for the following 3 passages.

Pooling of cells for bar-seq experiments

The barcoded mutant and parental cell lines were combined in mixed pools, adding similar numbers of

each individual cell line. For the *in vitro* screen, a total of 290 individually targeted transporter mutants,

13 array transporter mutants, 5 barcoded parental lines (SBL1-5; barcodes introduced into the SSU locus

(16), and 9 non-transporter knock-out mutants, of which 8 acted as controls; 3 Δ*IFT88* (LmxM.27.1130),

3 Δ*LPG1* (LmxM.25.0010) and 2 Δ*PF16* (LmxM.20.1400) different barcoded mutants, were combined

into a pool of $1x10^5$ cells/ml (Supplementary Table 4), which was split into three aliquots for replicate

measurements of *in vitro* growth.

For sand fly infections, four separate experiments were conducted using distinct pools (Supplementary

Table 4, Pool membership). Pool 1 contained 75 individually targeted transporter mutant and one array

mutant, Pool 2 contained 71 individual transporter mutants and three array mutants, Pool 3 contained,

77 individual transporter mutants and 5 array mutants and Pool 4 contained 76 individual transporter

mutants and 4 array mutants. Each pool also contained five barcoded parental lines (SBL1-5) and three

non-transporter knock-out control mutants (Δ*IFT88, ΔLPG1, ΔPF16*). Each of the four pools was split

into three equal aliquots (replicates) in preparation for the infection of female *Lutzomyia longipalpis*

sand flies.

Growth curves in vitro

For the *in vitro* growth assay, the mixed pool was split into 3 equal aliquots (replicates), which were left to grow in separate flasks for 48 h, diluted to 1×10^6 cells/ml, grown for an additional 24 h, diluted to 1×10^5 cells/ml and grown for an additional 72 h (total 144 hours) at $27\ ^\circ C + 5\ \%$ CO_2 in standard M199. For the growth curves of individual cell lines, log phase promastigotes (between 2 and 4×10^6 parasites/ml) of V-ATPase Subunit E (LmxM.36.3100) knockout, add-back (AB) and *L. mex* Cas9 T7 (C9T7) cell lines were seeded in standard M199 at a density of 1×10^5 cells/ml. Parasites were left to grow at $27\ ^\circ C$ for 3 consecutive days and on day 3, were diluted back to 1×10^5 cells/ml density and left to grow at $27\ ^\circ C$. Growth was assessed by counting the cells every 24 hours with a CASY® cell counter (Cambridge Bioscience) using a 60 μm capillary and measurement range set between 2 and 15 μm. For each condition measurements from three replicate flasks were recorded.

Sand fly infections

For infections with pooled barcoded mutant populations, each pool was seeded at a density of 2×10^6 parasites/ml and grown for 24 h at $26\ ^\circ C$ in standard M199 with 250 μg/ml of Amikacin (Amikin). On the day of infection, a total of either 3×10^7 (Pools 1-3), or 1.8×10^7 (Pool 4) logarithmic growing parasites were washed three times with sterile 0.9% NaCl saline solution (Braun) and then resuspended in 300 μl of saline, which were then mixed with 2.7 ml of ram's defibrinated blood (LabMedia), previously heat inactivated at $56\ ^\circ C$ for 35 minutes. For each separate pool, three groups of 120-180 female sand flies, 4-5 days old, were allowed to feed on the parasite-blood mixture, through a skin membrane from a 1-day old chick, as previously described (62). Fully engorged females were separated and maintained at $26\ ^\circ C$ with free access to 40% sucrose solution. Infected sand flies were dissected at days 2 (48 h) and 9 (216 h) post blood-meal (PBM) (Supplementary Table 8). At day 2 PBM, a total of 3 to 9 female sand fly guts were checked to qualitatively assess the progress, localisation and intensity of infection by light microscopy. Parasite abundance was graded into three qualitative categories: negative, light (<100 parasites/gut), moderate (100-1000 parasites/gut) and heavy (>1000 parasites/gut), as described elsewhere (59). For infections with individual promastigote cell lines; female sand flies (5–

9 days old) were infected by feeding through a chick-skin membrane parasites from log-phase cultures

(3–4 day cultures), washed twice in sterile saline solution and resuspended in heat-inactivated ram blood

at concentration of 1 x 10^6 promastigotes/ml. Engorged sand flies were maintained as described above.

Female flies were dissected at days 2 and 9 PBM and the abundance and localisation of *Leishmania*

promastigotes in the sand fly digestive tract was examined as described above. Experiments were

performed in duplicates.

Sampling of DNA for sequencing

For the promastigote *in vitro* cultures, gDNA was extracted at 0 h, 24 h, 48 h and 144 h of growth from

approximately $1x10^7$ cells from each replicate culture, using the Wizard® SV Genomic DNA

Purification System (Promega) according to the manufacturer's instructions, eluting in 40 μl of bi-

distilled water (Ambion). For the *in vivo* experiments, gDNA was extracted from the pool after mixing

in standard M199 and from the parasite-blood mixture used for the infection (time point 0). For

extraction of genomic DNA from whole infected sand flies, a total of 17 to 82 specimens were collected

from each batch at 2- and 9-days post blood meal (PBM) (Supplementary Table 5). For both cells and

tissues, the High Pure PCR Template Preparation Kit (Roche) was used and all samples were eluted in

100 μl of VWR Life Sciences PCR grade water as previously described (16).

Bar-seq library preparation and sequencing

The preparation of bar-seq amplicon libraries was done as previously reported (44), with minor changes.

For the initial bar-seq amplicon PCR, 100 ng of gDNA isolated from promastigote cultures, 250 ng of

gDNA isolated from *Leishmania*-blood meal mix and 500 ng gDNA isolated from whole *Leishmania*-

infected female sand flies, were used. To account for the excess of host gDNA present in blood and

sand fly derived samples, the number of cycles for the same PCR was increased from 31 for *Leishmania*

culture derived samples to 35 for blood and sand fly samples. Raw sequencing files (fastq) for all

samples generated during this study were deposited in the European Nucleotide Archive (ENA) study

accession PRJEB90861 (ERP173867).

Quantification of barcoded cell line fitness

For each sample, we grouped and counted raw sequencing reads containing barcode sequences, normalised this to the sample total reads and calculated the change in the proportion of reads of each cell line in each sample compared to the zero time point, as previously reported (45). The majority of cell lines in a pool do not exhibit a fitness change, therefore we identified the mode (peak) of the distribution of changes in proportion as the reference to compare to in each replicate. For each time point and cell line we assigned fitness scores by dividing the median of the cell line's change in proportion over all replicates with the median over all modes. A fitness score above one indicates that proportion of barcodes from a particular cell line have increased faster relative to the bulk of the pooled cell lines, corresponding to faster growth and/or better survival than the bulk of the pooled cell lines from the start of the assay up to that time point. A fitness score below one indicates the inverse. P-values were calculated using a paired t-test of the log-transformed cell line changes in proportions against the corresponding reference values, testing the null hypothesis that the cell line change in proportion from all replicates of a particular cell line in a given time point cannot be distinguished from the change of the bulk of the pooled cell lines in all replicates of the same time point. Cell lines were labelled as having a strong fitness phenotype in a given time point if their p-value was below 0.05 and their fitness score was either below 0.5 (deleterious phenotype) or above 2 (beneficial phenotype).

In vivo *parasite morphometry*

Midgut smears of infected sand flies were fixed with methanol, stained with Giemsa, examined by light microscopy with an oil immersion objective and photographed (Olympus DP70) (Supplementary Figure 6, Supplementary Table 7). Body length, flagellar length and body width of 200 randomly selected promastigotes were measured on day 9 PBM using Fiji (65). Four morphological forms were distinguished, based on criteria previously described (10,28). Briefly; elongated nectomonads (EN), body length $\geqslant$14 µm; leptomonads (LE) body length <14 m and flagellar length $\leqslant$2 times body length and metacyclic promastigotes (MP), flagellar length >2 times body length and body length < 14 µm.

Haptomonads were not distinguished in this study as they are often remaining attached to the gut and can be underrepresented on gut smears. Differences in proportion of morphological forms and measurements were compared by a Tukey's HSD (honestly significant difference) test, using the software SPSS version 27.

Acknowledgements

We like to thank Pamela Nicholson and Daniela Steiner (NGS Facility, University of Bern) for help with Illumina sequencing, Vít Dvořák for maintaining the colony of *Lutzomyia longipalpis*, Kristýna Srstková for technical support during sand fly experiments and all past and current members of the Gluenz and Volf labs for helpful discussions.

Funding statement

AAW was the recipient of a Marie Skłodowska-Curie Individual Fellowship (trans-LEISHion-EU FP7, No. 798736) and is supported by a Marie Skłodowska-Curie Global Fellowship (LeishBlock-Horizon, No. 101148623). RJW is supported by a Wellcome Trust Henry Dale Fellowship (211075/Z/18/Z). This work was supported by a UKRI Medical Research Council grant (MR/V000446/1; This UK funded award was part of the EDCTP2 programme supported by the European Union), the Wellcome Trust (221944/A/20/Z, 200807/Z/16/Z, 104627/Z/14/Z) and the Wellcome Centre for Integrative Parasitology (WCIP) core Wellcome Centre Award (104111/Z/14/Z) and a project grant from the Swiss National Science Foundation (310030_220011).

Data availability statement

All data supporting the findings of this study are available within the article and its supplementary materials, which have been deposited to Figshare (https://figshare.com/) under Doi: 10.6084/m9.figshare.29481254 (Supplementary Figure 6 and Supplementary Table 7).

References

1. Pace D. Leishmaniasis. J Infect. 2014 Nov;69(S1):S10–8.

2. Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L. Phlebotomine sandflies and the spreading of leishmaniases and other diseases of public health concern. Med Vet Entomol. 2013 Jun;27(2):123–47.

3. Dostálová A, Volf P. *Leishmania* development in sand flies: parasite-vector interactions overview. Parasit Vectors. 2012 Dec;5(1):276.

4. Catta-Preta CMC, Ghosh K, Sacks DL, Ferreira TR. Single-cell atlas of *Leishmania* development in sandflies reveals the heterogeneity of transmitted parasites and their role in infection. Proc Natl Acad Sci. 2024 Dec 24;121(52):e2406776121.

5. Hart DT, Coombs GH. *Leishmania mexicana*: Energy metabolism of amastigotes and promastigotes. Exp Parasitol. 1982 Dec;54(3):397–409.

6. McConville MJ, Saunders EC, Kloehn J, Dagley MJ. *Leishmania* carbon metabolism in the macrophage phagolysosome- feast or famine? F1000Research. 2015 Oct 1;4:938.

7. Rosenzweig D, Smith D, Opperdoes F, Stern S, Olafson RW, Zilberstein D. Retooling *Leishmania* metabolism: from sand fly gut to human macrophage. FASEB J. 2008 Feb;22(2):590–602.

8. Saunders EC, Ng WW, Kloehn J, Chambers JM, Ng M, McConville MJ. Induction of a Stringent Metabolic Response in Intracellular Stages of *Leishmania mexicana* Leads to Increased Dependence on Mitochondrial Metabolism. Wilson ME, editor. PLoS Pathog. 2014 Jan 23;10(1):e1003888.

9. Gossage SM, Rogers ME, Bates PA. Two separate growth phases during the development of *Leishmania* in sand flies: implications for understanding the life cycle. Int J Parasitol. 2003 Sep;33(10):1027–34.

10. Pruzinova K, Sadlova J, Myskova J, Lestinova T, Janda J, Volf P. *Leishmania* mortality in sand fly blood meal is not species-specific and does not result from direct effect of proteinases. Parasit Vectors. 2018 Dec;11(1):37.

11.   Sadlova J, Homola M, Myskova J, Jancarova M, Volf P. Refractoriness of *Sergentomyia*

schwetzi to *Leishmania* spp. is mediated by the peritrophic matrix. Traub-Csekö YM, editor. PLoS Negl

Trop Dis. 2018 Apr 4;12(4):e0006382.

12.   Rogers ME, Chance ML, Bates PA. The role of promastigote secretory gel in the origin and

transmission of the infective stage of *Leishmania mexicana* by the sandfly *Lutzomyia longipalpis*.

Parasitology. 2002 May;124(5):495–507.

13.   Sádlová J, Volf P. Peritrophic matrix of *Phlebotomus duboscqi* and its kinetics during

Leishmania major development. Cell Tissue Res. 2009 Aug;337(2):313–25.

14.   Myšková J, Dostálová A, Pěničková L, Halada P, Bates PA, Volf P. Characterization of a

midgut mucin-like glycoconjugate of *Lutzomyia longipalpis* with a potential role in *Leishmania*

attachment. Parasit Vectors. 2016 Dec;9(1):413.

15.   Jecna L, Dostalova A, Wilson R, Seblova V, Chang KP, Bates PA, et al. The role of surface

glycoconjugates in *Leishmania* midgut attachment examined by competitive binding assays and

experimental development in sand flies. Parasitology. 2013 Jul;140(8):1026–32.

16.   Beneke T, Demay F, Hookway E, Ashman N, Jeffery H, Smith J, et al. Genetic dissection of a

Leishmania flagellar proteome demonstrates requirement for directional motility in sand fly infections.

PLoS Pathog. 2019 Jun;15(6).

17.   Cuvillier A, Miranda JC, Ambit A, Barral A, Merlin G. Abortive infection of *Lutzomyia*

longipalpis insect vectors by aflagellated LdARL-3A-Q70L overexpressing *Leishmania amazonensis*

parasites. Cell Microbiol. 2003 Oct;5(10):717–28.

18.   Bates PA. Transmission of *Leishmania* metacyclic promastigotes by phlebotomine sand flies.

Int J Parasitol. 2007 Aug;37(10):1097–106.

19.   Yanase R, Moreira-Leite F, Rea E, Wilburn L, Sádlová J, Vojtkova B, et al. Formation and

three-dimensional architecture of *Leishmania* adhesion in the sand fly vector. eLife [Internet]. 2023 May

10 [cited 2025 Jul 7];12. Available from: https://elifesciences.org/articles/84552

20.   Rogers M, Kropf P, Choi BS, Dillon R, Podinovskaia M, Bates P, et al. Proteophosophoglycans

Regurgitated by *Leishmania*-Infected Sand Flies Target the L-Arginine Metabolism of Host

Macrophages to Promote Parasite Survival. Ribeiro JMC, editor. PLoS Pathog. 2009 Aug

21;5(8):e1000555.

21. Ilg T, Stierhof YD, Wiese M, McConville MJ, Overath P. Characterization of phosphoglycan-

containing secretory products of *Leishmania*. Parasitology. 1994 Mar;108(S1):S63–71.

22. Stierhof YD, Bates PA, Jacobson RL, Rogers ME, Schlein Y, Handman E, et al. Filamentous

proteophosphoglycan secreted by *Leishmania* promastigotes forms gel-like three-dimensional networks

that obstruct the digestive tract of infected sandfly vectors. Eur J Cell Biol. 1999 Oct;78(10):675–89.

23. Volf P, Hajmova M, Sadlova J, Votypka J. Blocked stomodeal valve of the insect vector: similar

mechanism of transmission in two trypanosomatid models. Int J Parasitol. 2004 Oct;34(11):1221–7.

24. Yanase R, Pruzinova K, Owino BO, Rea E, Moreira-Leite F, Taniguchi A, et al. Discovery of

essential kinetoplastid-insect adhesion proteins and their function in *Leishmania*-sand fly interactions.

Nat Commun. 2024 Aug 13;15(1):6960.

25. Rogers ME, Hajmová M, Joshi MB, Sadlova J, Dwyer DM, Volf P, et al. *Leishmania* chitinase

facilitates colonization of sand fly vectors and enhances transmission to mice. Cell Microbiol. 2008

Jun;10(6):1363–72.

26. Schlein Y, Jacobson RL, Messer G. *Leishmania* infections damage the feeding mechanism of

the sandfly vector and implement parasite transmission by bite. Proc Natl Acad Sci. 1992 Oct

15;89(20):9944–8.

27. Ilg T, Stierhof YD, Craik D, Simpson R, Handman E, Bacic A. Purification and Structural

Characterization of a Filamentous, Mucin-like Proteophosphoglycan Secreted by *Leishmania* Parasites.

J Biol Chem. 1996 Aug;271(35):21583–96.

28. Rogers ME, Ilg T, Nikolaev AV, Ferguson MAJ, Bates PA. Transmission of cutaneous

leishmaniasis by sand flies is enhanced by regurgitation of fPPG. Nature. 2004 Jul;430(6998):463–7.

29. Rogers ME. The Role of *Leishmania* Proteophosphoglycans in Sand Fly Transmission and

Infection of the Mammalian Host. Front Microbiol [Internet]. 2012 [cited 2025 Apr 10];3. Available

from: http://journal.frontiersin.org/article/10.3389/fmicb.2012.00223/abstract

30. Kamhawi S, Ramalho-Ortigao M, Van M. Pham, Kumar S, Lawyer PG, Turco SJ, et al. A Role

for Insect Galectins in Parasite Survival. Cell. 2004 Oct;119(3):329–41.

31.   Sacks DL, Modi G, Rowton E, Späth G, Epstein L, Turco SJ, et al. The role of phosphoglycans

in *Leishmania* –sand fly interactions. Proc Natl Acad Sci. 2000 Jan 4;97(1):406–11.

32.   Baker N, Catta-Preta CMC, Neish R, Sadlova J, Powell B, Alves-Ferreira EVC, et al. Systematic

functional analysis of *Leishmania* protein kinases identifies regulators of differentiation or survival. Nat

Commun. 2021 Feb 23;12(1):1244.

33.   Späth GF, Lye LF, Segawa H, Sacks DL, Turco SJ, Beverley SM. Persistence without pathology

in phosphoglycan-deficient *Leishmania major*. Science. 2003 Aug;301(5637):1241–3.

34.   Sádlová J, Price HP, Smith BA, Votýpka J, Volf P, Smith DF. The stage-regulated HASPB and

SHERP proteins are essential for differentiation of the protozoan parasite *Leishmania major* in its sand

fly vector, *Phlebotomus papatasi: Leishmania* metacyclogenesis in the sand fly. Cell Microbiol. 2010

Dec;12(12):1765–79.

35.   Secundino N, Kimblin N, Peters NC, Lawyer P, Capul AA, Beverley SM, et al.

Proteophosphoglycan confers resistance of *Leishmania major* to midgut digestive enzymes induced by

blood feeding in vector sand flies: Proteophosphoglycan protects *L. major*. Cell Microbiol. 2010 Jan

20;12(7):906–18.

36.   Gazanion E, Seblova V, Votypka J, Vergnes B, Garcia D, Volf P, et al. *Leishmania infantum*

nicotinamidase is required for late-stage development in its natural sand fly vector, *Phlebotomus*

perniciosus. Int J Parasitol. 2012 Apr;42(4):323–7.

37.   Jecna L, Dostalova A, Wilson R, Seblova V, Chang KP, Bates PA, et al. The role of surface

glycoconjugates in *Leishmania* midgut attachment examined by competitive binding assays and

experimental development in sand flies. Parasitology. 2013 Jul;140(8):1026–32.

38.   Doehl JSP, Sádlová J, Aslan H, Pružinová K, Metangmo S, Votýpka J, et al. *Leishmania* HASP

and SHERP Genes Are Required for In Vivo Differentiation, Parasite Transmission and Virulence

Attenuation in the Host. Beverley SM, editor. PLOS Pathog. 2017 Jan 17;13(1):e1006130.

39.   Sunter JD, Yanase R, Wang Z, Catta-Preta CMC, Moreira-Leite F, Myskova J, et al. *Leishmania*

flagellum attachment zone is critical for flagellar pocket shape, development in the sand fly, and

pathogenicity in the host. Proc Natl Acad Sci. 2019 Mar 26;116(13):6351–60.

40.   Halliday C, Yanase R, Catta-Preta CMC, Moreira-Leite F, Myskova J, Pruzinova K, et al. Role

for the flagellum attachment zone in *Leishmania* anterior cell tip morphogenesis. Hill KL, editor. PLOS

Pathog. 2020 Oct 22;16(10):e1008494.

41.   Corrales RM, Vaselek S, Neish R, Berry L, Brunet CD, Crobu L, et al. The kinesin of the

flagellum attachment zone in *Leishmania* is required for cell morphogenesis, cell division and virulence

in the mammalian host. Hill KL, editor. PLOS Pathog. 2021 Jun 18;17(6):e1009666.

42.   Alcoforado Diniz J, Chaves MM, Vaselek S, Miserani Magalhães RD, Ricci-Azevedo R, De

Carvalho RVH, et al. Protein methyltransferase 7 deficiency in *Leishmania major* increases neutrophil

associated pathology in murine model. De Oliveira CI, editor. PLoS Negl Trop Dis. 2021 Mar

2;15(3):e0009230.

43.   Beneke T, Gluenz E. Bar-seq strategies for the LeishGEdit toolbox. Mol Biochem Parasitol.

2020 Sep;239:111295.

44.   Albuquerque-Wendt A, McCoy C, Neish R, Dobramysl U, Alagoz C, Beneke T, et al.

TransLeish: Identification of membrane transporters essential for survival of intracellular *Leishmania*

parasites in a systematic gene deletion screen. Nat Commun. 2025 Jan 2;16(1):299.

45.   Bengs F, Scholz A, Kuhn D, Wiese M. LmxMPK9, a mitogen-activated protein kinase

homologue affects flagellar length in *Leishmania mexicana*. Mol Microbiol. 2005 Mar;55(5):1606–15.

46.   Louradour I, Ferreira TR, Duge E, Karunaweera N, Paun A, Sacks D. Stress conditions promote

Leishmania hybridization in vitro marked by expression of the ancestral gamete fusogen HAP2 as

revealed by single-cell RNA-seq. eLife. 2022 Jan 7;11:e73488.

47.   Svárovská A, Ant TH, Seblová V, Jecná L, Beverley SM, Volf P. *Leishmania major*

Glycosylation Mutants Require Phosphoglycans (lpg2−) but Not Lipophosphoglycan (lpg1−) for

Survival in Permissive Sand Fly Vectors. Traub-Cseko YM, editor. PLoS Negl Trop Dis. 2010 Jan

12;4(1):e580.

48.   Kamhawi S. Phlebotomine sand flies and *Leishmania* parasites: friends or foes? Vol. 22, Trends

in Parasitology. 2006. p. 439–45.

49.   Volf P, Myskova J. Sand flies and *Leishmania*: specific versus permissive vectors. Trends

Parasitol. 2007 Mar;23(3):91–2.

50.   Myskova J, Svobodova M, Beverley SM, Volf P. A lipophosphoglycan-independent

development of *Leishmania* in permissive sand flies. Microbes Infect. 2007 Mar;9(3):317–24.

51.   Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, et al. The

Transporter Classification Database (TCDB): 2021 update. Nucleic Acids Res. 2021 Jan

8;49(D1):D461–7.

52.   Beneke T, Madden R, Makin L, Valli J, Sunter J, Gluenz E. A CRISPR Cas9 high-throughput

genome editing toolkit for kinetoplastids. R Soc Open Sci. 2017 May;4(5):1–16.

53.   Burchmore RJS, Rodriguez-Contreras D, Mcbride K, Barrett MP, Modi G, Sacks D, et al.

Genetic characterization of glucose transporter function in *Leishmania mexicana* [Internet]. Available

from: www.pnas.orgcgidoi10.1073pnas.0630165100

54.   Rodriguez-Contreras D, Feng X, Keeney KM, Bouwer HGA, Landfear SM. Phenotypic

characterization of a glucose transporter null mutant in *Leishmania mexicana*. Mol Biochem Parasitol.

2007 May;153(1):9–18.

55.   Aellig S, Billington K, Damasceno JD, Davidson L, Dobramysl U, Etzensperger R, et al.

LeishGEM: genome-wide deletion mutant fitness and protein localisations in *Leishmania*. Trends

Parasitol. 2024 Aug;40(8):675–8.

56.   Jones NG, Catta-Preta CMC, Lima APCA, Mottram JC. Genetically Validated Drug Targets in

Leishmania: Current Knowledge and Future Prospects. Vol. 4, ACS Infectious Diseases. American

Chemical Society; 2018. p. 467–77.

57.   Martínez-García M, Campos-Salinas J, Cabello-Donayre M, Pineda-Molina E, Gálvez FJ,

Orrego LM, et al. LmABCB3, an atypical mitochondrial ABC transporter essential for *Leishmania*

major virulence, acts in heme and cytosolic iron/sulfur clusters biogenesis. Parasit Vectors. 2016 Jan

5;9:7.

58.   Colasante C, Diaz PP, Clayton C, Voncken F. Mitochondrial carrier family inventory of

Trypanosoma brucei brucei: Identification, expression and subcellular localisation. Mol Biochem

Parasitol. 2009 Oct;167(2):104–17.

59.   Volfová V, Jančářová M, Volf P. Sand fly blood meal volumes and their relation to female body

weight under experimental conditions. Parasit Vectors. 2024 Aug 23;17(1):360.

60.   Yernaux C, Fransen M, Brees C, Lorenzen S, Michels PAM. *Trypanosoma brucei* glycosomal

ABC transporters: Identification and membrane targeting. Mol Membr Biol. 2006;23(2):157–72.

61.   Descoteaux A, Mengeling BJ, Beverley SM, Turco SJ. *Leishmania donovani* has distinct

mannosylphosphoryltransferases for the initiation and elongation phases of lipophosphoglycan

repeating unit biosynthesis. Mol Biochem Parasitol. 1998 Jul 1;94(1):27–40.

62.   Benaim G, García-Marchán Y, Reyes C, Uzcanga G, Figarella K. Identification of a

sphingosine-sensitive Ca2+ channel in the plasma membrane of *Leishmania mexicana*. Biochem

Biophys Res Commun. 2013 Jan;430(3):1091–6.

63.   Plourde M, Ubeda JM, Mandal G, Monte-Neto RLD, Mukhopadhyay R, Ouellette M.

Generation of an aquaglyceroporin AQP1 null mutant in *Leishmania major*. Mol Biochem Parasitol.

2015 Aug;201(2):108–11.

64.   Beyenbach KW. The plasticity of extracellular fluid homeostasis in insects. J Exp Biol. 2016

Sep 1;219(17):2596–607.

65.   Najem O, Shah MM, Zubair M, De Jesus O. Serum Osmolality. In: StatPearls [Internet].

Treasure Island (FL): StatPearls Publishing; 2025 [cited 2025 May 21]. Available from:

http://www.ncbi.nlm.nih.gov/books/NBK567764/

66.   Richard D, Kündig C, Ouellette M. A new type of high affinity folic acid transporter in the

protozoan parasite *Leishmania* and deletion of its gene in methotrexate-resistant cells. J Biol Chem.

2002 Aug 16;277(33):29460–7.

67.   Ouameur AA, Girard I, Légaré D, Ouellette M. Functional analysis and complex gene

rearrangements of the folate/biopterin transporter (FBT) gene family in the protozoan parasite

Leishmania. Mol Biochem Parasitol. 2008 Dec;162(2):155–64.

68.   Baker N, Hamilton G, Wilkes JM, Hutchinson S, Barrett MP, Horn D. Vacuolar ATPase

depletion affects mitochondrial ATPase function, kinetoplast dependency, and drug sensitivity in

trypanosomes. Proc Natl Acad Sci U S A. 2015 Jul;112(29):9112–7.

69.   Santos VC, Nunes CA, Pereira MH, Gontijo NF. Mechanisms of pH control in the midgut of

Lutzomyia longipalpis: roles for ingested molecules and hormones. J Exp Biol. 2011 May

1;214(9):1411–8.

70.   Te Brugge V, Ianowski JP, Orchard I. Biological activity of diuretic factors on the anterior

midgut of the blood-feeding bug, Rhodnius prolixus. Gen Comp Endocrinol. 2009 May;162(1):105–12.

71.   He C. Balancing nutrient and energy demand and supply via autophagy. Curr Biol CB. 2022

Jun 20;32(12):R684–96.

72.   Finbow ME, Harrison MA. The vacuolar H+-ATPase: a universal proton pump of eukaryotes.

Biochem J. 1997 Jun 15;324 (Pt 3)(Pt 3):697–712.

73.   Maxson ME, Grinstein S. The vacuolar-type H+-ATPase at a glance – more than a proton pump.

J Cell Sci. 2014 Dec 1;127(23):4987–93.

74.   Williams RA, Tetley L, Mottram JC, Coombs GH. Cysteine peptidases CPA and CPB are vital

for autophagy and differentiation in *Leishmania mexicana*. Mol Microbiol. 2006 Aug;61(3):655–74.

75.   Volf P, Volfova V. Establishment and maintenance of sand fly colonies. J Vector Ecol. 2011

Mar;36(SUPPL.1).

76.   Beneke T, Gluenz E. LeishGEdit: A Method for Rapid Gene Knockout and Tagging Using

CRISPR-Cas9. In: Methods in Molecular Biology. Humana Press Inc.; 2019. p. 189–210.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Figure Legends

Figure 1. Consolidated summary of gene deletion results for the transportome of *Leishmania mexicana*.

(A) Top, pie-charts showing the numbers of successful gene deletions (cyan) and non-successful deletion attempts (magenta), across two independent screens (44 and this study). Bottom, break-down of non-successful deletions attempts into two sub-categories: (i) Double drug-resistant populations where ORF is still detected (or PCR inconclusive) (yellow); (ii) Attempts where no drug resistant populations were ever recovered, or populations where resistant cells could only be recovered with single drug selection and ORF was still detected (dark pink). (B) Summary of gene deletion results separated into TCDB families (Supplementary Table 3); colours as for A.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Figure 2. Fitness of promastigote mutants *in vitro*.

(A) Overview of the experiment timeline for *in vitro* growth of promastigotes in culture. DNA sampling time-points are indicated by yellow arrows. Dark pink arrows denote dilutions, to (a) 1×10^6 parasites/ml and (b) 1×10^5 parasites/ml. (B) Growth profile of the masterpool of promastigote mutants over time. Data points are the average of three measurements; yellow dots indicate where gDNA was sampled. The dotted line indicates dilution of the cultures. (C) Trajectories of the average of normalised reads of the promastigote masterpool, relative to time-point "0 hours" (T_0). Red dotted line highlight relative barcode abundance of 1. Controls are shown in colour: dark blue, SBL1-5 parental cell lines; Cyan, ΔLPG1; Magenta, ΔIFT88; Yellow, ΔPF16. Grey, all other barcoded cell lines. (D) Volcano plot showing fitness scores against p-values of mutants from the promastigote masterpool after 144 hours of growth. Dashed lines demarcate fitness score thresholds of < 0.5 and > 2, and a significance threshold of $p < 0.05$. Barcodes meeting both threshold criteria are coloured black, non-significant values are grey. Controls are coloured as in D.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Figure 3. Fitness of promastigote mutants *in vivo*.

(A) Overview of the experiment timeline for sand fly infections. DNA sampling time-points are indicated by yellow arrows. (B) Fitness scores from the promastigote masterpool after 144 hours *in vitro* growth plotted against the fitness scores from all mutants 216 hours after infection of sand flies. Black dashed lines mark fitness score thresholds of < 0.5 and > 2. (C-F) Volcano plots showing fitness scores against p-values after 216 h (9 days PBM) in sand flies, separated by sub-pools (P1-P4). Dashed lines demarcate fitness score thresholds of < 0.5 and > 2, and a significance threshold of p < 0.05. Barcodes meeting both threshold criteria are colored black, non-significant are grey. Controls are shown in colour: Dark blue dots, SBL1-5 parental cell lines; Cyan, ΔLPG1; Magenta, ΔIFT88; Yellow, ΔPF16. Black numbers denote the abbreviated GeneIDs (LmxM.xx.xxxx) of selected mutants with the highest fitness changes.

"Identification of transporters essential for survival of *Leishmania* promastigotes in the digestive tract of sand flies"

Figure 4. V-ATPase is required for differentiation and colonisation in the sand fly vector.

(A) Trajectories of the average of normalised reads for V-ATPase deletion mutants from the promastigote masterpool (grey lines), relative to T_0. Dark blue lines, SBL1-5 parental controls. (B) Barcode trajectories for V-ATPase deletion mutants in sand flies, normalised to the start of the experiment. Colour code as in A. (C) Schematic of V-ATPase pump with each subunit labelled. Subunit E, for which the null mutant was individually characterised, is highlighted in magenta. (D) Growth of parental (PAR, dark blue), V-ATPase V_1E null mutant (KO, magenta) and V-ATPase V_1E add-back (AB, grey) mutant promastigotes *in vitro*. After 3 days of continuous growth, cultures were diluted back to 1 x 10^5 parasites/ml and monitored for an additional 3 days. (E) Parasite abundance in the digestive tract of dissected sand flies infected with PAR, AB or KO parasite lines, assessed at 2 days PBM (left) and 9 days PBM (right). (F) Location of PAR, AB or KO parasite lines at 2 days PBM (left) and 9 days PBM (right). NI, non-infected; ES, endoperitrophic space; AMG, abdominal midgut; TMG, thoracic midgut; CAR, cardia; SV, stomodeal valve. (G) Promastigote morphotypes of PAR, AB or KO parasite lines observed in infected gut smears after 9 days PBM.

Figure 4

Figure 1

Figure 2

Figure 3