Title: A nucleation distance sets the cell size-dependent P-body assembly

Authors: Xuefang Gu[1], Dexin Shen[1], Xing Quan[1], Tatsuhisa Tsuboi[1,2],*

Affiliations:

[1] Institute of Biopharmaceutical and Health Engineering, Tsinghua Shenzhen International Graduate School, Shenzhen 518055, China

[2] Tsinghua-SIGS & Jilin Fuyuan Guan Food Group Joint Research Center, Tsinghua Shenzhen International Graduate School, Shenzhen 518055, China

*To whom correspondence should be addressed:

Tatsuhisa Tsuboi

Institute of Biopharmaceutical and Health Engineering, Tsinghua Shenzhen International Graduate School, Shenzhen 518055, China

Tel: +86-152-1679-5216

E-mail address: ttsuboi@sz.tsinghua.edu.cn

Abstract:

Cells rely on various intracellular structures to regulate biological processes, including phase-separated granules. P-bodies are a type of membrane-less condensate found in eukaryotic cytoplasm that mediate mRNA degradation and storage. While the molecular composition of P-bodies is well studied, the relationship between their spatial organization and cell geometry, particularly cell size and organelle distribution, remains unclear. Here, we developed a conceptual framework to model phase-separation nucleation and validate the nucleation distance of a P-body in living yeast cells. Using microfluidic cell sorting and quantitative fluorescence microscopy, we found that smaller cells contain fewer, larger P-bodies, whereas larger cells form more, smaller granules. Monte Carlo simulations of P-body nucleation accurately recapitulate these cell size-dependent patterns, revealing a nucleation distance of approximately 2.2 μm. Furthermore, we demonstrate that increasing vacuolar volume reduces the effective cytoplasmic space and elevates the number of P-bodies, although cell size remains the dominant determinant. Our findings establish nucleation distance as a key parameter that links cell geometry to the biophysical regulation of condensate formation.

Keywords

P-body, cell size, phase separation, modeling, microfluidics

INTRODUCTION

Cellular structures act in concert to regulate essential processes such as intracellular transport, energy conversion, and gene expression [1–3]. Among these structures, membrane-less organelles formed through liquid-liquid phase separation (LLPS) play a critical role in gene expression regulation [4–6]. Cytoplasmic Processing bodies (P-bodies) are prototypical LLPS condensates in eukaryotic cells [3]. They sequester translationally repressed messenger RNAs (mRNAs) together with the decapping and decay enzymes [7,8], thereby orchestrating mRNA degradation, storage, and recycling [8–11]. P-body assembly involves several partially redundant scaffold proteins, including Edc3[12,13], Dcp2[14], Dhh1 [15], Pat1 [15,16], and Lsm4 [12,17], such that deletion of any one protein reduces P-body number and size and impairs the recruitment of client RNAs and proteins [18,19].

Phase separation begins with nucleation, the moment a small molecular cluster becomes large enough to overcome the energetic cost of forming a new interface. In supersaturated solutions or vapours, only molecules that diffuse within a reaction radius can bind productively; beyond this distance, collisions are too infrequent to surpass the nucleation barrier [20]. Once a condensate forms, it recruits nearby molecules, creating a depletion zone that reduces local supersaturation. New droplets can only nucleate beyond this depleted region. Together with the natural tendency of neighboring droplets to fuse and minimize surface energy, this phenomenon defines a characteristic nucleation distance (d_n), the typical spacing at which independent condensates can stably coexist [21]. Inside cells, molecular crowding and sub-diffusive dynamics further limit droplet fusion, allowing many condensates to persist at approximately d_n apart rather than merging into a single large phase[22]. Simulation using a coarse-grained protein model revealed that the stability of biomolecular condensates depends largely on the number of attractive interactions between

molecules[23]. By modulating concentration, crowding, or binding affinity, cells can tune d_n to

control both the number and spatial distribution of condensates.

Despite the complexity of the intracellular environment, nucleation in living cells still adheres to

fundamental physics constraints, the existence of an energy barrier, and the requirement for a

critical molecular cluster, with cellular components often acting as seeds to facilitate this process

[20]. The concept of a nucleation distance thus bridges physical theory and cell biology, as it arises

from the necessity of accumulating sufficient molecules within a reaction radius and is reinforced

by molecular depletion effects and coalescence dynamics, ultimately leading to a regulated, non-

random spatial distribution of condensates within the cell. These insights suggest that P-body

assembly is governed not only by protein composition and interactions but also by the physical

dimensions and spatial organization of the cell.

Cellular organization is determined by the geometry of intracellular compartments and directly

influences the scale and efficiency of biosynthetic processes [24–26]. As cells grow larger, organelles

generally scale proportionally [27–29]. Although the total protein concentration is maintained, the

concentration of individual proteins may increase or decrease with cell size [30–32]. Interestingly, P-

bodies are known to appear during G1 and expand in size through G2, without a corresponding

change in number [33], suggesting that spatial constraints, rather than protein abundance, may play

a key role in phase separation. Yet, the precise influence of cell size and spatial parameters on P-

body formation and dynamics remains largely unexplored.

Here, we extend this conceptual framework to quantitatively model condensate nucleation and

validate the presence of a characteristic nucleation distance in living cells. By combining

quantitative microscopy of size-sorted cells using microfluidic devices with Monte Carlo

simulations for nucleation, we find that larger cells tend to form more but smaller P-bodies,

whereas smaller cells generate fewer but larger P-bodies. The simulations identify a specific

nucleation distance (d_n) that aligns closely with our experimental observations. Moreover, in *sch9Δ*

and *vps1Δ* mutants, where vacuolar volume is enlarged and the effective cytoplasmic space

reduced, we observe a significant increase in P-body number, reinforcing the importance of spatial

restriction in Phase separation. Together, these results enable us to investigate how cells regulate

the nucleation distance to influence the formation, size, and distribution of phase-separated

compartments.

Results

P-bodies number increases with increasing yeast cell size

To examine the relationship between cell size and P-body formation under physiological

conditions, we set out to quantify how the spatial organization and cell geometry of a yeast cell

influence the initiation and abundance of P-bodies in vivo while avoiding genetic manipulations

that can secondarily perturb LLPS [34,35]. We developed a microfluidic device based on cell trapping

for long-period microscopy [36,37], machined with one hundred parallel trap lines whose constriction

widths gradually increase from 5.1 µm to 10 µm, enabling continuous perfusion of logarithmically

growing W303 cells and deterministic physical sorting by diameter (Fig. 1A, B, Supplementary

Fig. 1). This design provided a non-invasive and high-throughput method for examining a wide

range of cell sizes in a single experiment, while minimizing batch-to-batch variability

(Supplementary Fig. 1C). This platform preserved nutrient availability while imaging under an

identical glucose-starvation regime (0 % glucose, synthetic complete medium), a known

physiological condition that robustly induces P-body formation. The use of physical sorting rather

than genetic perturbation was crucial in isolating the impact of cell size per se, thereby avoiding

confounding effects such as stress responses, altered expression of phase-separation components,

or cell-cycle defects.

We tagged Dcp2, a core scaffold component of P-bodies, with a novel green fluorescent protein

variant, mAbbsinthe, which exhibits superior photostability and reduced aggregation relative to

commonly used GFPs (Supplementary Fig. 2A, B)[38]. Importantly, we confirmed that

Dcp2_mAbbsinthe retained P-body formation responsive to stress, validating it as a functional

reporter (Supplementary Fig. 2C). In addition, this choice was critical for long-term imaging of

condensate dynamics, especially under stress-induced conditions such as glucose starvation. Using

spinning-disk confocal microscopy, we captured three-dimensional image stacks of P-body

distribution in yeast cells of varying diameters (Fig. 1C). Quantitative analysis revealed a clear

positive correlation between cell size and the number of P-bodies (Fig. 1D). The mean P-body

number rose from 2.7 in 3.5–4.5 µm cells, 4.4 in 4.5-5.5 µm cells, to 6.2 µm in >5.5 µm cells.

These findings were corroborated using Edc3-GFP, an alternative P-body marker, validating that

the phenomenon is not tag-specific. Together, our results provide direct experimental evidence

that cell size is a determinant of P-body number, suggesting that physical cellular geometry plays

a role in influencing phase separation behavior.

A Monte-Carlo simulation based on nucleation identifies a characteristic nucleation distance

To test whether simple geometric constraints could explain the empirically observed scaling, we

implemented a Monte-Carlo simulation based on nucleation. In this model, we approximated the

yeast mother cell as a sphere and distributed virtual Dcp2 molecules randomly within its volume

(Fig. 2A). This abstraction was motivated by the physical properties of phase-separated

condensates, which behave as mesoscale entities with restricted mobility and volume-excluding

effects. Collisions resulting in center-to-center separations below a tunable threshold (nucleation

distance, d_n) trigger irreversible coalescence, conserving total mass. These fusion events were

iteratively calculated until all remaining molecular pairs were spaced beyond this threshold. We

initialized the model with 200 seed spheres per cell, each with radii ranging from 2.0 to 3.0 µm.

Across this range, a best-fit d_n of 2.2 µm simultaneously reproduced the monotonic increase in

punctum number with cell size (Fig. 1D, 2B, Supplementary Figure 3A). We then analyzed the

size of P-bodies. Since P-bodies were visualized through microscopy imaging, which potentially

results in deviations from their actual biological sizes. To ensure consistency between the

microscopy experiment and simulated data, we analyzed P-bodies size by calculating the

proportion of puncta in each size category. (Supplementary Figure 3B). We first matched the

category proportion between the two experiments for a cell size of 5.0 µm. With this size

classification, the simulation for the other two different size cells, 4.0 µm and 6.0 µm, matches

surprisingly well with the empirically observed triphasic size distribution of P-bodies (Fig. 2C -

G). Smaller cells tend to form larger P-bodies, whereas larger cells predominantly contain smaller

P-bodies. This agreement suggests that the physical spacing between successful nucleation events,

a feature not encoded in protein sequence or interaction affinity, acts as a central organizing

principle. To probe robustness, we performed sensitivity analyses varying initial molecular

numbers and cytoplasmic viscosity (Supplementary Figure 3C, D). The final puncta number was

largely invariant to particle count above ~200 and only weakly dependent on viscosity (within

biologically plausible ranges), indicating that d_n is the dominant determinant of condensate

distribution. These simulations highlight that spatial constraints and collision thresholds alone are

sufficient to explain the observed morphological transitions, without invoking complex

biochemical regulation.

Organelle occupancy amplifies nucleation frequency by restricting effective cytoplasmic

volume

While total cell volume increases during growth, the effective cytoplasmic space available for P-

body formation may be limited by the presence of large organelles such as the vacuole and nucleus.

To explore the impact of such intracellular crowding, we extended our simulation to incorporate

internal organelles occupying specific volumetric fractions. We implemented three distinct models

to simulate molecule fusion paths in the presence of a central spherical nucleus and vacuole,

representing non-permissive organelle space: (1) path-restricted fusion where molecular lines

crossing the nucleus were invalid, (2) path-shifted fusion where granules repositioned to avoid

organelle overlap, and (3) path-redirected fusion where particles navigated around obstacles using

a calculated shortest path (Fig. 3A–C).

Across all models, increasing organelle volume fractions consistently led to elevated final P-body

counts, even under a constant nucleation distance (Fig. 3D–G). This effect arose because internal

obstructions constrained the diffusion space, increasing local molecule concentrations and

reducing the effective reaction volume. These simulations suggest that intracellular crowding can

enhance condensate nucleation independent of changes in protein concentration or binding

affinity, highlighting a potential mechanism for tuning granule formation through organelle scaling

or spatial reorganization.

Vacuole mutants validate the influence of spatial confinement on P-body assembly

To test the simulation-derived hypothesis that intracellular crowding promotes condensate

nucleation, we examined two yeast mutants known to affect vacuolar morphology: *sch9Δ* and

vps1Δ. Both mutations are known to increase vacuole-to-cell volume ratios through distinct

mechanisms: *vps1Δ* via impaired membrane fission and *sch9Δ* via defects in nutrient sensing and

vacuolar growth regulation[27,39]. Quantification of vacuole-to-cell volume ratios confirmed

significantly increased organelle occupancy in both mutants compared to wild-type (Fig. 4A, B).

Fluorescence intensity analysis showed that Dcp2_mabbsinthe expression remained unchanged

(Fig. 4C), ensuring that any observed effects on P-bodies were not due to altered protein abundance.

As predicted by the simulation, *vps1Δ* cells exhibited significantly higher P-body counts across all

size bins (Fig. 4D), indicating that reduced cytoplasmic space can elevate nucleation rates.

Surprisingly, *sch9Δ* cells, despite also showing increased vacuole volume, did not exhibit a similar

rise in P-body number. Instead, both *sch9Δ* and *vps1Δ* displayed a marked shift toward smaller P-

body sizes (Fig. 4E, F), especially in larger cells.

This discrepancy can be reconciled by considering the nonlinear relationship between volume and

linear distance: a 10% reduction in cytoplasmic volume translates to only a ~0.17 μm. For details,

a 10% reduction in volume leads to only about 3.45% shrinkage in radius, which, for typical yeast

cytoplasmic dimensions (~5 μm), results in a 0.17 μm decrease in distance. Reduction in effective

radius, which is minor compared to the ~2.2 μm nucleation distance inferred from simulations.

Thus, small changes in cytoplasmic diameter may be insufficient to affect nucleation probability

but may still impair granule coalescence, favoring fragmentation in *sch9Δ* cells. In addition, the

simulated puncta numbers correspond to when the central sphere is calculated with less than 1%

of whole cell volume, using a d_n of 2.2 μm, which is far below the actual occupation by the

organelles (Fig. 3D-F). This suggests that effective intervention by the cellular organization must

be much smaller, possibly due to the condensate itself, which diffuses within the cell. These

findings experimentally confirm that intracellular crowding affects P-body morphology and

support the existence of a spatially constrained fusion threshold that governs the organization of

condensates.

205 Larger P-bodies exhibit delayed disassembly upon nutrient repletion

Having established that spatial parameters regulate condensate number and size, we next asked

whether these morphological features influence functional stability. In classical phase-separation

systems, larger droplets are thermodynamically favored and dissolve more slowly due to reduced

surface-to-volume ratios and kinetic buffering [40]. We hypothesized that large P-bodies would

persist longer upon glucose reintroduction, providing a functional correlate to their size. We

performed time-lapse confocal imaging of glucose-starved cells subjected to nutrient repletion and

tracked P-body disassembly over a 10-minute window. As expected, small and medium P-bodies

disappeared rapidly, with >80% disassembling within 4 minutes (Fig. 5A–B). In contrast, large P-

bodies were highly stable, often persisting until the final time point. This size-dependent

disassembly rate suggests that larger condensates have altered material properties, such as

increased viscosity or reduced surface curvature, that slow their dissolution. These results suggest

that condensate size is not merely a geometric consequence of fusion events but has functional

implications for granule stability and cellular memory. Larger P-bodies may act as more robust

repositories for mRNA and protein complexes, potentially influencing translational reactivation or

RNA decay following stress relief. This observation supports a model in which P-body size

encodes distinct biophysical states, governed by nucleation and fusion history, and modulated by

spatial constraints during formation.

Discussion

Our quantitative imaging combined with microfluidic cell size sorting and Monte Carlo modeling

shows that phase-separated P-bodies follow a characteristic nucleation distance (~2.2 μm), which

affects their number based on the overall cell diameter (Fig. 1, 2). By demonstrating that a single

geometric parameter is sufficient to recapitulate the full empirical relationship between cell size

and condensate number, we provide, to our knowledge, the first *in vivo* measurement of a

nucleation distance for any biomolecular condensate. This finding confirms long-standing

predictions from classical nucleation theory [20–23] and extends them to the crowded, heterogeneous

cytoplasm. Previous work has shown that many organelles scale with growth [24–29], but whether

equivalent principles govern LLPS compartments has been unclear. Our microfluidic approach

allowed us to modulate cell diameter without genetic perturbation or cell cycle arrest, cleanly

separating geometrical from biochemical variables. The experiments showed larger cells

consistently accumulated more, but smaller, P-bodies (Fig. 1D). The simulation showed that this

pattern naturally forms when nucleation events are prevented within d_n of an existing droplet: as

the cytoplasmic volume increases, the number of non-overlapping spheres of radius d_n roughly

increases with volume (Fig. 2A-C). These data argue that cells can modulate condensate

abundance passively, simply by growing. Our minimal model deliberately omits molecular

interactions, active transport, or cytoskeletal barriers, yet captures the essential statistics of P-body

distribution. This parsimonious view suggests that d_n signifies a fundamental effective reaction

radius that incorporates all microscopic determinants (multivalent interactions, crowding,

diffusivity) into a single mesoscale parameter. Future work can refine this framework by allowing

d_n to vary in response to environmental cues, post-translational modifications, or ATP levels,

ultimately enabling the quantitative prediction of condensate patterns from first principles.

Extending the model to include a central vacuole and nucleus showed that increasing internal

occupancy raises condensate number even at constant cell size (Fig. 3). The effect is conceptually

similar to the soft wall confinement used to promote vapour nucleation in microfluidic cavitation

chambers: reducing accessible volume raises the local supersaturation of phase separating

components and shortens the mean free path required to exceed the nucleation barrier. The vacuole

enlarged mutant *vps1Δ* validated this prediction in vivo, exhibiting higher P-body counts across

all size bins (Fig. 4D). By contrast, *sch9Δ* increased vacuole volume but did not change granule

number, highlighting that modest reductions in linear distance (< 0.1 μm) are negligible relative to

d_n, yet still fragment larger condensates and bias the population towards smaller droplets (Fig. 4E–

F). Together, these observations position spatial confinement as a tunable knob for condensate

biogenesis, complementing known biochemical regulators such as post translational modifications

or RNA scaffolding [9–12,14–18,41].

The strikingly slower dissolution of large P-bodies after glucose re-addition (Fig. 5) mirrors classic

Ostwald ripening behaviour, in which surface curvature drives exchange with the soluble pool.

Our data, therefore, suggest that the history of nucleation and fusion events leaves a lasting

physical imprint that regulates the lifetime of stress-induced mRNA repositories. A speculative

but attractive model is that large, stable granules favour selective retention of particular mRNAs

or protein complexes, endowing cells with a tunable memory of past stress episodes. P-bodies are

known to appear in G1 and grow through G2 without changing in number [33]. We reconcile this

observation with our findings by noting that budding yeast double their diameter only late in the

cell cycle; before this size threshold is crossed, the cytoplasm remains too small to permit

additional nucleation sites. More broadly, any physiological or pathological state that alters cell

volume or organelle allotment, such as hypertrophic growth, vacuolar fragmentation in aging, or

lysosomal enlargement in neurodegeneration, could reshuffle the cellular condensate landscape.

Indeed, impaired clearance of RNA granules is implicated in ALS and related disorders [6]. We

propose that disrupted spatial scaling, rather than intrinsic changes in phase-separating proteins,

may contribute to the aberrant accumulation of condensates in these contexts.

We imposed glucose starvation to synchronize P-body formation; whether d_n is conserved for other

stresses or constitutive granules, such as stress granules, remains to be tested. Additionally, the

model treats cells as perfect spheres and neglects the geometry of buds or cytosolic viscosity

gradients. High-speed light sheet imaging of nascent granules could capture the accurate

nucleation kinetics and refine these assumptions. Furthermore, our static snapshots cannot

distinguish between *de novo* nucleation and the fission of pre-existing droplets; combining

selective photoconversion with long-term tracking would resolve this issue. Finally, although we

focused on scaffold protein concentration, RNA itself can nucleate P-bodies [41]; incorporating

measured RNA fluxes may sharpen model accuracy. By quantitatively bridging cell-scale

geometry with mesoscale phase behaviour, our study uncovers a simple spatial rule that governs

how many condensates a cell can host. We anticipate that nucleation distance will prove to be a

unifying concept across diverse LLPS assemblies, offering a geometric lens through which to

interpret the dynamic self-organisation of the cytoplasm.

Methods

Yeast strains and plasmids

The yeast strains and plasmids used are listed in Supplementary File 1, and the oligonucleotides

used for plasmid construction and gene modification are listed in Supplementary File 2. To reduce

variability among the constructed yeast strains, the strains were created either through the

integration of a linear PCR product or a plasmid linearized through restriction digestion. The

variants of yeast expression plasmids (TTP257, PXQP017, PXQP032) for fluorescent protein were

constructed by swapping GFP of pFA6a-link-yoGFP-SpHis5 [42]with mAbb0.5 and mAbbsinthe[38]

through the combination of Gibson assembly and PCR. Fluorescent protein tagging for Dcp2 and

Edc3 with mAbb0.5, mAbbsinthe, and yoGFP was performed with PCR-mediated homologous

recombination using pFA6a-link-mAbb0.5-SpHis5, pFA6a-link- mAbbsinthe-SpHis5, and

pFA6a-link-yoGFP-SpHis5, and integrations were confirmed by microscopy. Su9-mCherry was

expressed by the TDH3 promoter from the integrated plasmid TTP076, and the strains with three

copies of the integrated plasmids in a cell were screened through microscopy and used for further

experiments. Deletion mutant strains were constructed with PCR-mediated homologous

recombination using pFA6a-hphMX6, and the integration of the hphMX6 cassette was confirmed

by PCR followed by electrophoresis.

Microfluidic device fabrication

The design and fabrication of the microfluidic device for yeast replicative aging were based on previously published work [43,44]. To enable the capture of yeast cells with different diameters, the initial 6 µm cell trap design was adjusted to include trap diameters ranging from 5.1 to 10 µm.

Setting up a microfluidic experiment

Before the microfluidics experiment, the device was placed under vacuum for 20 minutes to remove air from the channels. The device was then placed on the stage of an inverted spinning disk confocal microscope. Media ports were connected to plastic tubing, which in turn was connected to 20 mL syringes filled with fresh synthetic complete medium (SD) (CSM powder from Sunrise Science, #1001-100, supplemented with 2% glucose and 0.04% Tween-20). These syringes were positioned approximately 24 inches above the microscope stage to drive media flow via gravity. The waste port was connected to tubing positioned at stage height to maintain a stable flow of waste.

Yeast cells were grown in 3 mL of SC medium and cultured overnight at 30°C to an OD_{600} of more than 1.0. For loading, cells were diluted to an OD_{600} of 0.1 and cultured for 4 or 5 hours to an OD_{600} of 0.5-0.7. The media syringe was replaced with the syringe containing the yeast culture temporarily for loading cells. Most traps were filled within 1 to 2 minutes. Medium flow into the device was gravity-driven, allowing cells to enter and occupy the traps. Most traps were filled within 1 to 2 minutes. After loading, the yeast culture syringe was replaced with the media supply syringe, and all tubing heights were adjusted to maintain a height differential of approximately 60 inches.

Microscopy

A P-body visualization experiment in a 96-well plate was performed as follows: Yeast cells were grown in SC medium with 2% glucose in a 12 mL tube at 30°C with a rotator speed of 70 rpm. 100μl of mid-log phase yeast cells (OD600 of 0.4 to 0.7) were harvested and placed into a 96-well Glass Bottom Plate (Cellvis LLC) coated with 0.1 mg/mL concanavalin A (Sigma-Aldrich C2010). After 15 minutes of incubation, the well was washed twice with 100 μL of SC media without glucose and then filled with 100 μL of SC media without glucose.

Image data for visualizing P-bodies in a microfluidic device were collected as follows: Yeast cells were grown in SC medium using a 15ml glass tube at 30°C with a rotator speed of 70 rpm until they reached an OD_{600} of approximately 1.0. The following day, yeast cells were diluted to an OD_{600} of 0.1 and cultured for an additional 4 to 5 hours until they reached an OD_{600} of approximately 0.6, before being used in the experiment. To capture the cells in the microfluidic device, the culture was diluted 4-fold and transferred into a 20 mL syringe prefilled with fresh SC medium containing 0.04% Tween-20. After loading into microfluidics chips and 30 minutes of glucose starvation, an image was collected.

Microscopy: Cells were imaged at 23°C by an Eclipse Ti2-E Spinning Disk Confocal with Yokogawa CSU-X1 (Yokogawa) with 50 μm pinholes, located at the Tsinghua-SIGS Tsuboi Laboratory. Imaging was performed using a CFI Plan Apochromat Lambda 100x 1.49 NA oil objective. Z-stacks (300nm steps) were acquired by a Prime 95B sCMOS camera (Photometrics) (pixel size 0.11μm). Imaging was controlled using NIS-Elements software (Nikon). Images of the microfluidic device were acquired using the large-image acquisition mode. The scanning area was set to 0.1 × 1.5 mm with a 10% overlap between adjacent fields.

The measurement of the photobleach resistance

To measure the photo bleach resistance of GFPs, a 488 nm laser, and a 510–520 nm emission filter were used. The cells were imaged with 50% of pre-set laser power 20 times. 25 flames of z-section with 100 milliseconds were acquired in steps of 300nm. After background subtraction, the total fluorescence intensity was divided by the cell occupation area in the image. The fluorescence intensity was normalized to the initial intensity to obtain Relative Fluorescence Strength (RFS). The average RFS of the 30 cells was calculated, and the RFS curve was created by MATLAB (version 2021a) using polynomial curve fitting.

Molecular nucleation model for P-body formation

The model was based on Monte Carlo simulation approach. Since the yeast cell is approximately spherical, it was modeled as a spherical region in the simulation. P-bodies are randomly distributed in the cytoplasm and are also roughly spherical; therefore, in the simulation, they were also defined as spherical molecules. Details were as followed:

First, the initial number of molecules per cell and the number of simulated cells were defined. Cell diameter was used to represent different cell sizes. In each simulation, the initial position and volume of each molecule were assigned using a random number generator, and their positions were marked in a three-dimensional Cartesian coordinate system. The following procedure was employed to generate the three-dimensional coordinates for each molecule:

(1) Random angle generation: Two angles, θ and φ, representing the azimuthal and polar angles in spherical coordinates, were generated using uniform random distributions.

379   (2) Radial distribution: A radial coordinate was calculated based on the predefined cell radius, with

380   a cube root distribution applied to ensure uniform volume distribution.

381   (3) Position conversion: Using the standard spherical-to-Cartesian coordinate transformation, the

382   position in spherical coordinates was converted to three-dimensional Cartesian coordinates

383   according to the following equations:

384
$$x = r * cos(\theta) * sin(\varphi)$$

385
$$y = r * sin(\theta) * sin(\varphi)$$

386
$$z = r * cos(\varphi)$$

Since this study induced P-body formation by applying glucose starvation stress, the mobility of

P-bodies in this condition is very slow and can be neglected [45]; therefore, molecules were

considered to be in a static state. When the distance between two molecules fell below a critical

threshold (a predefined nucleation distance), these molecules were assumed to aggregate to

become one molecule.

The aggregation process was implemented as follows:

(1) Distance calculation and collision detection: To efficiently identify the closest pair of

molecules, an auxiliary function, findclosestpoint, was developed. Different models use different

findclosestpoint functions, as mentioned in Figure 3.

(2) Molecule merging: Upon collision, the volumes and positions of the two molecules were

merged. The resulting molecule had an increased volume, and its position was determined by

findclosestpoint.

(3) State update: The number and positions of molecules in the cell were updated accordingly after

each aggregation event.

The code updated the distribution of molecules in the cell based on the current positions and

volumes at each simulation step, progressively approaching the final aggregation state. The

simulation loop was terminated once the distances between all molecules exceeded the predefined

nucleation distance.

Quantification and statistical analysis

Detailed statistics, including the number of cells analyzed, mean value, standard deviation, and

standard error of the mean, are indicated in each figure legend. The Wilcoxon rank-sum test was

performed using GraphPad Prism (GraphPad software).

Data and materials availability

All data supporting the findings of this study are available within the paper and its supplementary

information. Yeast strains and plasmids used in the study are provided in Supplementary File 1.

The oligonucleotides used are listed in Supplementary File 2. Further information and requests for

raw data and reagents should be directed to and will be fulfilled by the lead contact, T.T.

(ttsuboi@sz.tsinghua.edu.cn).

Code Availability

The   Pytorch   codes   for   p-body   drug   screening   can   be   found   at

https://github.com/xuexuegu/nucleation.git.

Supplemental information

Supplementary File 1 | List of oligonucleotides used for yeast construction

Acknowledgments

We thank Nathan Shaner and Gerard Lambert for sharing mAbb0.5 and mAbbsinthe plasmids. We

thank Abdul Khan for his helpful discussions and feedback on the paper. This work was supported

in part by the Key Research and Development Program of the Ministry of Science and Technology

2024YFE0102700, 2023YFA0914303, Shenzhen Science and Technology Innovation

Commission WDZC20220811144737001, the Jilin Fuyuan Guan Food Group Co., Ltd, startup

fund OD2021031C, Interdisciplinary Research and Innovation Fund JC2022008, Overseas

Research Cooperation Fund HW2024009 from Tsinghua SIGS (to T.T.),

Author contributions

X.G. and T.T. conceived and designed the project. X.G. and X.Q. performed wet experiments and

analyzed image data. X.G. and D.S. performed image quantification and computational analysis.

X.G. and T.T. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1. van Bergeijk, P., Hoogenraad, C.C., and Kapitein, L.C. (2016). Right Time, Right Place: Probing the Functions of Organelle Positioning. Trends Cell Biol *26*, 121–134. https://doi.org/10.1016/j.tcb.2015.10.001.

2. Teti, A. (1992). Regulation of cellular functions by extracellular matrix. Journal of the American Society of Nephrology *2*, S83. https://doi.org/10.1681/ASN.V210s83.

3. Luo, Y., Na, Z., and Slavoff, S.A. (2018). P-Bodies: Composition, Properties, and Functions. Biochemistry *57*, 2424–2431. https://doi.org/10.1021/acs.biochem.7b01162.

4. West, A.P., and Shadel, G.S. (2017). Mitochondrial DNA in innate immune responses and inflammatory pathology. Nat Rev Immunol *17*, 363–375. https://doi.org/10.1038/nri.2017.21.

5. Ow, Y.-L.P., Green, D.R., Hao, Z., and Mak, T.W. (2008). Cytochrome c: functions beyond respiration. Nat Rev Mol Cell Biol *9*, 532–542. https://doi.org/10.1038/nrm2434.

6. Wang, B., Zhang, L., Dai, T., Qin, Z., Lu, H., Zhang, L., and Zhou, F. (2021). Liquid–liquid phase separation in human health and diseases. Signal Transduct Target Ther *6*, 290. https://doi.org/10.1038/s41392-021-00678-1.

7. Choi, Y.-J., and Song, K. (2022). Nst1, Densely Associated to P-Body in the Post-Exponential Phases of Saccharomyces cerevisiae, Shows an Intrinsic Potential of Producing Liquid-Like Condensates of P-Body Components in Cells. Int J Mol Sci *23*, 2501. https://doi.org/10.3390/ijms23052501.

8. Noda, N.N., Wang, Z., and Zhang, H. (2020). Liquid–liquid phase separation in autophagy. Journal of Cell Biology *219*. https://doi.org/10.1083/jcb.202004062.

9. Parker, R., and Sheth, U. (2007). P Bodies and the Control of mRNA Translation and Degradation. Mol Cell *25*, 635–646. https://doi.org/10.1016/j.molcel.2007.02.011.

10. Eulalio, A., Behm-Ansmant, I., and Izaurralde, E. (2007). P bodies: at the crossroads of post-transcriptional pathways. Nat Rev Mol Cell Biol *8*, 9–22. https://doi.org/10.1038/nrm2080.

11. Decker, C.J., and Parker, R. (2012). P-Bodies and Stress Granules: Possible Roles in the Control of Translation and mRNA Degradation. Cold Spring Harb Perspect Biol *4*, a012286–a012286. https://doi.org/10.1101/cshperspect.a012286.

12. Decker, C.J., Teixeira, D., and Parker, R. (2007). Edc3p and a glutamine/asparagine-rich domain of Lsm4p function in processing body assembly in *Saccharomyces cerevisiae*. J Cell Biol *179*, 437–449. https://doi.org/10.1083/jcb.200704147.

13. Xing, W., Muhlrad, D., Parker, R., and Rosen, M.K. (2020). A quantitative inventory of yeast P body proteins reveals principles of composition and specificity. Elife *9*. https://doi.org/10.7554/eLife.56525.

14. van Dijk, E. (2002). Human Dcp2: a catalytically active mRNA decapping enzyme located in specific cytoplasmic structures. EMBO J *21*, 6915–6924. https://doi.org/10.1093/emboj/cdf678.

15. Hurto, R.L., and Hopper, A.K. (2011). P-body components, Dhh1 and Pat1, are involved in tRNA nuclear-cytoplasmic dynamics. RNA *17*, 912–924. https://doi.org/10.1261/rna.2558511.

16.   Sachdev, R., Hondele, M., Linsenmeier, M., Vallotton, P., Mugler, C.F., Arosio, P., and Weis, K.
(2019). Pat1 promotes processing body assembly by enhancing the phase separation of the DEAD-
box ATPase Dhh1 and RNA. Elife *8*. https://doi.org/10.7554/eLife.41415.

17.   INGELFINGER, D., ARNDT-JOVIN, D.J., LÜHRMANN, R., and ACHSEL, T. (2002). The
human LSm1-7 proteins colocalize with the mRNA-degrading enzymes Dcp1/2 and Xrn1 in
distinct cytoplasmic foci. RNA *8*, 1489–1501. https://doi.org/10.1017/S1355838202021726.

18.   Rao, B.S., and Parker, R. (2017). Numerous interactions act redundantly to assemble a tunable size
of P bodies in *Saccharomyces cerevisiae*. Proceedings of the National Academy of Sciences *114*.
https://doi.org/10.1073/pnas.1712396114.

19.   Teixeira, D., and Parker, R. (2007). Analysis of P-Body Assembly in *Saccharomyces cerevisiae*.
Mol Biol Cell *18*, 2274–2287. https://doi.org/10.1091/mbc.e07-03-0199.

20.   Shimobayashi, S.F., Ronceray, P., Sanders, D.W., Haataja, M.P., and Brangwynne, C.P. (2021).
Nucleation landscape of biomolecular condensates. Nature *599*. https://doi.org/10.1038/s41586-
021-03905-5.

21.   Shin, Y., Chang, Y.C., Lee, D.S.W., Berry, J., Sanders, D.W., Ronceray, P., Wingreen, N.S.,
Haataja, M., and Brangwynne, C.P. (2018). Liquid Nuclear Condensates Mechanically Sense and
Restructure the Genome. Cell *175*. https://doi.org/10.1016/j.cell.2018.10.057.

22.   Lee, D.S.W., Choi, C.H., Sanders, D.W., Beckers, L., Riback, J.A., Brangwynne, C.P., and
Wingreen, N.S. (2023). Size distributions of intracellular condensates reflect competition between
coalescence and nucleation. Nat Phys *19*. https://doi.org/10.1038/s41567-022-01917-0.

23.   Espinosa, J.R., Joseph, J.A., Sanchez-Burgos, I., Garaizar, A., Frenkel, D., and Collepardo-
Guevara, R. (2020). Liquid network connectivity regulates the stability and composition of
biomolecular condensates with many components. Proc Natl Acad Sci U S A *117*.
https://doi.org/10.1073/pnas.1917569117.

24.   Schmoller, K.M., and Skotheim, J.M. (2015). The Biosynthetic Basis of Cell Size Control. Trends
Cell Biol *25*, 793–802. https://doi.org/10.1016/j.tcb.2015.10.006.

25.   Goehring, N.W., and Hyman, A.A. (2012). Organelle Growth Control through Limiting Pools of
Cytoplasmic Components. Current Biology *22*, R330–R339.
https://doi.org/10.1016/j.cub.2012.03.046.

26.   Chan, Y.-H.M., and Marshall, W.F. (2010). Scaling properties of cell and organelle size.
Organogenesis *6*, 88–96. https://doi.org/10.4161/org.6.2.11464.

27.   Chan, Y.-H.M., and Marshall, W.F. (2014). Organelle Size Scaling of the Budding Yeast Vacuole
Is Tuned by Membrane Trafficking Rates. Biophys J *106*, 1986–1996.
https://doi.org/10.1016/j.bpj.2014.03.014.

28.   Rafelski, S.M., Viana, M.P., Zhang, Y., Chan, Y.-H.M., Thorn, K.S., Yam, P., Fung, J.C., Li, H.,
Costa, L. da F., and Marshall, W.F. (2012). Mitochondrial Network Size Scaling in Budding
Yeast. Science (1979) *338*, 822–824. https://doi.org/10.1126/science.1225720.

29. Jorgensen, P., Edgington, N.P., Schneider, B.L., Rupeš, I., Tyers, M., and Futcher, B. (2007). The
Size of the Nucleus Increases as Yeast Cells Grow. Mol Biol Cell *18*, 3523–3532.
https://doi.org/10.1091/mbc.e06-10-0973.

30. Lanz, M.C., Zhang, S., Swaffer, M.P., Ziv, I., Götz, L.H., Kim, J., McCarthy, F., Jarosz, D.F.,
Elias, J.E., and Skotheim, J.M. (2024). Genome dilution by cell growth drives starvation-like
proteome remodeling in mammalian and yeast cells. Nat Struct Mol Biol *31*, 1859–1871.
https://doi.org/10.1038/s41594-024-01353-z.

31. Zhurinsky, J., Leonhard, K., Watt, S., Marguerat, S., Bähler, J., and Nurse, P. (2010). A
Coordinated Global Control over Cellular Transcription. Current Biology *20*, 2010–2015.
https://doi.org/10.1016/j.cub.2010.10.002.

32. Newman, J.R.S., Ghaemmaghami, S., Ihmels, J., Breslow, D.K., Noble, M., DeRisi, J.L., and
Weissman, J.S. (2006). Single-cell proteomic analysis of S. cerevisiae reveals the architecture of
biological noise. Nature *441*, 840–846. https://doi.org/10.1038/nature04785.

33. Safieddine, A., Benassy, M.-N., Bonte, T., Slimani, F., Pourcelot, O., Kress, M., Ernoult-Lange,
M., Courel, M., Coleno, E., Imbert, A., et al. (2024). Cell-cycle-dependent mRNA localization in
P-bodies. Mol Cell *84*, 4191-4208.e7. https://doi.org/10.1016/j.molcel.2024.09.011.

34. Harvey, S.L., Enciso, G., Dephoure, N., Gygi, S.P., Gunawardena, J., and Kellogg, D.R. (2011). A
phosphatase threshold sets the level of Cdk1 activity in early mitosis in budding yeast. Mol Biol
Cell *22*, 3595–3608. https://doi.org/10.1091/mbc.e11-04-0340.

35. Jorgensen, P., Nishikawa, J.L., Breitkreutz, B.-J., and Tyers, M. (2002). Systematic Identification
of Pathways That Couple Cell Growth and Division in Yeast. Science (1979) *297*, 395–400.
https://doi.org/10.1126/science.1070850.

36. Li, Y., Jiang, Y., Paxman, J., O'Laughlin, R., Klepin, S., Zhu, Y., Pillus, L., Tsimring, L.S., Hasty,
J., and Hao, N. (2020). A programmable fate decision landscape underlies single-cell aging in
yeast. Science (1979) *369*, 325–329. https://doi.org/10.1126/SCIENCE.AAX9552.

37. Zhou, Z., Liu, Y., Feng, Y., Klepin, S., Tsimring, L.S., Pillus, L., Hasty, J., and Hao, N. (2023).
Engineering longevity - design of a synthetic gene oscillator to slow cellular aging. Science (1979)
380. https://doi.org/10.1126/science.add7631.

38. Lambert, G.G., Depernet, H., Gotthard, G., Schultz, D.T., Navizet, I., Lambert, T., Adams, S.R.,
Torreblanca-Zanca, A., Chu, M., Bindels, D.S., et al. (2020). Aequorea's secrets revealed: New
fluorescent proteins with unique properties for bioimaging and biosensing. PLoS Biol *18*,
e3000936. https://doi.org/10.1371/journal.pbio.3000936.

39. Tsuboi, T., Viana, M.P., Xu, F., Yu, J., Chanchani, R., Arceo, X.G., Tutucci, E., Choi, J., Chen,
Y.S., Singer, R.H., et al. (2020). Mitochondrial volume fraction and translation duration impact
mitochondrial mRNA localization and protein synthesis. Elife *9*.
https://doi.org/10.7554/eLife.57814.

40. Ostwald ripening (2014). In The IUPAC Compendium of Chemical Terminology (International
Union of Pure and Applied Chemistry (IUPAC)). https://doi.org/10.1351/goldbook.O04348.

41.   Xing, W., Muhlrad, D., Parker, R., and Rosen, M.K. (2020). A quantitative inventory of yeast P
558   body proteins reveals principles of composition and specificity. Elife *9*.
559   https://doi.org/10.7554/eLife.56525.

560   42.   Lee, S., Lim, W.A., and Thorn, K.S. (2013). Improved Blue, Green, and Red Fluorescent Protein
561   Tagging Vectors for S. cerevisiae. PLoS One *8*, e67902.
562   https://doi.org/10.1371/journal.pone.0067902.

563   43.   Li, Y., Jiang, Y., Paxman, J., O'laughlin, R., Klepin, S., Zhu, Y., Pillus, L., Tsimring, L.S., Hasty,
564   J., and Hao, N. A programmable fate decision landscape underlies single-cell aging in yeast.

565   44.   Paxman, J., Zhou, Z., O'Laughlin, R., Liu, Y., Li, Y., Tian, W., Su, H., Jiang, Y., Holness, S.E.,
566   Stasiowski, E., et al. (2022). Age-dependent aggregation of ribosomal RNA-binding proteins links
567   deterioration in chromatin stability with challenges to proteostasis. Elife *11*.
568   https://doi.org/10.7554/eLife.75978.

569   45.   Joyner, R.P., Tang, J.H., Helenius, J., Dultz, E., Brune, C., Holt, L.J., Huet, S., Müller, D.J., and
570   Weis, K. (2016). A glucose-starvation response regulates the diffusion of macromolecules. Elife *5*.
571   https://doi.org/10.7554/eLife.09376.

Figure 1 | P-bodies number increases with increasing yeast cell size

(A) Different sizes of the cells are captured in the microfluidic device based on the well size

written below.

(B) The size of the cells in 96-well plates and microfluidic chips. Each dot corresponds to one

cell (n>90). Statistical significance was assessed by the Unpaired T-test (**** P < 0.0001).

(C) P-bodies in varying sizes of yeast cells were labeled with Dcp2_mAbbsinthe and analyzed 30

minutes after glucose starvation. Yellow arrows indicate P-bodies in mother cells. Scale bar, 2

μm.

(D) P-body number per cell across different cell sizes. Yeast cells expressing Dcp2_mAbbsinthe

or Edc3_mAbbsinthe were categorized into three size ranges based on cell diameter. Statistical

significance was assessed by Unpaired T-test (ns P > 0.05, n>30).

Figure 2 | A Monte-Carlo simulation based on nucleation identifies a characteristic nucleation distance

(A) Particle nucleation model based on Monte Carlo simulation. 1. Generate particles: Initial distribution of molecules in 3D space before simulation, each blue dot represents a molecule. 2. Analyze the closest distance: Each molecule recognizes the closest one and then fuses. 3. No more

merge: Result of the merge model simulation, showing puncta left after fusion, each granule

represents a P-body.

(B) Nucleation distance correlates with the final puncta number. Based on the nucleation distance,

final puncta numbers were simulated for each cell size. (n = 200 per condition). Horizontal dashed

lines indicate the mean P-body number per cell of different cell size groups in Figure 1D.

(C) Size fraction of P-body in the 4.5-5.5μm cell size by the manual classification of microscope

image.

(D) The definition of the simulated size of the P-body. The boundary for different sizes of P-bodies

in the simulation is based on the fraction in C.

(E) Size fraction of P-body in the different cell sizes by the manual classification of microscope

image data.

(F) Size of P-bodies in different cell sizes at the d_n = 2.2 μm in the simulation.

(G) Fractions of the simulated P-body size in the different cell sizes based on the definition in D.

Figure 3 | Organelle occupancy amplifies nucleation frequency by restricting effective cytoplasmic volume

(A-C) Schematic diagrams of three different molecular nucleation models in a cell with a central occupation by an organelle (gray circle).

(A) Model 1: Molecules fuse with their closest neighbor only if the connecting line does not intersect the central organelle.

(B) Model 2: Molecules fuse with their closest neighbor even if the direct path crosses the nucleus, but the merged granule is repositioned to avoid overlap with the central organelle.

(C) Model 3: Molecules fuse with their closest neighbor based on the true shortest path around the nucleus, allowing nucleus-aware trajectories.

619   (D-F) The final puncta number under different nucleation distances and varying organelle

620   fractions. The color represents the different organelle fractions relative to the cell.

Figure 4 | Vacuole mutants validate the influence of spatial confinement on P-body assembly

(A) Visualization of P-bodies in WT, *sch9Δ*, and *vps1Δ* yeast strains expressing Dcp2_mAbbsinthe under glucose starvation for 30 minutes. Cells are grouped by cell diameter (3.5-4.5, 4.5-5.5, >5.5), and P-bodies are indicated by arrows. Images are z-projections of 42 slices. Scale bar, 2 μm.

(B) Quantification of vacuole-to-cell volume ratio in WT, *sch9Δ*, and *vps1Δ* strains. Each dot

represents an individual experiment (n = 3 biological replicates). Statistical analysis was performed

using one-way ANOVA with multiple comparisons (**$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$).

(C) Intensity of Dcp2_mAbbsinthe fluorescence in WT, *sch9Δ*, and *vps1Δ* strains. No significant

differences were observed. Each dot represents an individual experiment.

(D) P-body number per cell across different cell diameter groups (3.5-4.5 μm, 4.5-5.5 μm, >5.5

μm) in WT, *sch9Δ*, and *vps1Δ* strains. Statistical significance was determined by two-way

ANOVA with multiple comparisons (ns $P > 0.05$, ****$P < 0.0001$).

(E) Proportion of large, middle, and small P-bodies in *sch9Δ* cells across the cell size groups.

(F) Proportion of large, middle, and small P-bodies in *vps1Δ* cells across the cell size groups.

Figure 5 | Larger P-bodies exhibit delayed disassembly upon nutrient repletion

(A) Time-lapse images of P-bodies in yeast cells after glucose addition, following prior glucose starvation of 30 minutes. Images were taken at 1, 2, 4, 8, and 10 minutes after the addition of glucose. Yellow arrows indicate large P-bodies, white arrows indicate small P-bodies. Images are z-projections of 42 slices. Scale bar, 2 μm.

(B) Quantification of the number of P-bodies per cell over time after glucose addition. Small P-bodies rapidly disaggregate following glucose treatment, while large P-bodies are more stable. P-

647   body size is primarily determined at the initial time point of formation. n> 45 cells within three

648   independent experiments.

649   (C) A schematic model description of P-body formation in yeast cells. Upon stress induction,

650   molecular components (blue dots) and mRNA (orange lines) condense into P-bodies via nucleation

651   within a spatial nucleation distance (~2.2 µm).

A

B

C

654 **Supplementary Figure 1 | Design of microfluidics for different sizes of the cells**

655 (A) Design of microfluidics for different sizes of the cells.

656 (B) Example of a microfluidic device with a well size of from 5.1 μm to 10 μm.

657 (C) The microfluidics can capture different size of the cells consistently (n ≥ 30 cells per repeat).

Supplementary Figure 2 | Dcp2 labeled with mAbbsinthe exhibits similar condensation and

greater resistance to photobleaching compared to GFP

(A) Z-stack images of Dcp2 labeled with mAbb0.5, mAbbsinthe, and YoGFP. Cells were

visualized upon glucose (G) starvation for 0, 10, 20, and 30 minutes. The scale bar indicates 2 μm.

(B) Various Dcp2-GFPs were photobleached under a 10-Watt laser for a 50-ms exposure for each

time point. The fluorescence recovery was monitored over 20 seconds (n = 30 cells), and the

relative fluorescence strength was plotted.

(C) Number of P-bodies per cell for mAbbsinthe and YoGFP-tagged Dcp2 under glucose

starvation for 30 min. Statistical significance was assessed by the Unpaired T-test (ns P > 0.05,

n>30).

Supplementary Figure 3 | Effects of nucleation distance and molecular number on P-body

formation simulation.

(A) Relationship between the final puncta number across different nucleation distances with

different sizes of the cell.

(B) Example of the different P-body sizes. Yellow boxes highlight the representative P-bodies in

the different size groups. 1 - large, 2 - middle, and 3 - small groups, respectively.

(C) The final puncta number is independent of molecular concentration in the nucleation

simulation. Bar graph shows the final number of simulated P-bodies when varying the initial

number of molecules (200, 500, 1000). Statistical significance was assessed by the Unpaired T-

test (ns P > 0.05, n = 200 simulations per group).

(D) Punta size distribution does not change upon molecular number.