Brain-wide connectivity patterns of feedforward and feedback cortico-cortical neurons in

the mouse secondary visual cortex

Richard G. Dickson[1], Matthew W. Jacobs[1], John M. Ratliff[1], Alec L.R. Soronow[1], Faye An[1],

Walid A. Yuqob[1], Euiseok J. Kim[1,2,3]*

[1]Department of Molecular, Cell, and Developmental Biology, University of California, Santa

Cruz, CA 95064, USA

[2]Institute for the Biology of Stem Cells, University of California, Santa Cruz, CA 95064, USA

[3]Lead Contact

Running title: Brain-wide circuits for FF/FB neurons

*Corresponding Author: Euiseok J. Kim

Department of Molecular, Cell, and Developmental Biology, University of California, Santa

Cruz, 1156 High Street, Santa Cruz, CA 95064, USA

Phone: 831-502-7233

Fax: 831-459-3139

ekim62@ucsc.edu

ACKNOWLEDGEMENTS

We thank Bin Chen, Bradley Colquitt, and David Feldheim for reading the manuscript and

Efrain Hernandez Alvarez for mouse husbandry and histology. We thank undergraduate

volunteers Eugene Kim, Jordan Chong, Hannah Tao, Lily Taylor-Hunt, Paul Pham, and Julian

Swartz for tissue processing and histology. We acknowledge technical support from Benjamin

Abrams, UCSC Life Sciences Microscopy Center, RRID: SCR_021135. Purchase of the Zeiss

880 confocal microscope used in this research was made possible through the National Institutes

of Health s10 Grant 1S10OD23528-01. We acknowledge support from the UCSC start-up fund,

the Whitehall Foundation, the Hellman Fellows Program, the E. Matilda Ziegler Foundation for

the Blind, BRAIN Initiative at the National Institutes of Health RF1MH132591 and the National

Institute of Neurological Disorders and Stroke at the National Institutes of Health,

R01NS128771 (E.J.K.), the Koret scholar program (A.L.R.S.), and a National Institutes of

Health IRACDA Postdoctoral Training Grant, K12GM139185 (R.G.D.).

ABSTRACT

Feedforward and feedback cortico-cortical neurons are distinct yet spatially intermingled

subtypes distributed across cortical layers, playing specialized roles in sensory and cognitive

processing. However, whether their presynaptic inputs differ to support these functions remains

unknown. Using projection- and layer-specific monosynaptic rabies tracing, we mapped brain-

wide long-distance inputs to multiple feedforward and feedback neuron types in VISl (also

known as LM), the mouse secondary visual cortex. Overall, long-distance input patterns for these

feedforward and feedback neurons were largely similar, as all received the majority of their

inputs from VISp, the primary visual cortex, along with substantial inputs from various other

cortical and visual thalamic regions. Despite their similarities, these feedforward and feedback

types differed in the proportion of long-distance cortical inputs originating from specific visual,

retrosplenial, and auditory cortices. These findings reveal the input connectivity patterns of

cortico-cortical neurons based on feedforward and feedback projections, providing an anatomical

framework for future studies on their functions and circuit integration.

KEYWORDS

cortical hierarchy, cortico-cortical neurons, feedforward and feedback neurons, mouse visual

cortex, rabies tracing

KEY POINTS

1) Feedforward and feedback cortico-cortical projection neurons in VISl form largely distinct

populations that are distributed across multiple cortical layers.

2) Four different feedforward and feedback cortico-cortical projection neuron types receive

broadly similar patterns of long-distance input across the brain.

3) Despite overall similarities, feedforward and feedback neurons differ in the relative

proportions of inputs from specific areas, including visual, retrosplenial, and auditory cortices.

INTRODUCTION

The mammalian cortex is responsible for high-level perception and cognition and is composed of numerous interconnected cortical areas (Felleman and Van Essen, 1991; Oh et al., 2014; Gămănuț et al., 2018). These areas communicate through long-distance inter-areal connections mediated by excitatory cortico-cortical projection neurons (CCPNs). Two key organizational features of the mammalian cortex are its six-layered laminar structure and its hierarchical organization (Felleman and Van Essen, 1991; Cajal, 1995). Each cortical layer is considered as a different information-processing unit (Harris and Shepherd, 2015). CCPNs in a specific layer of each cortical area can be further divided into feedforward projection neurons, which send axons to higher-order cortical areas, and feedback projection neurons, which project to lower-order cortical areas. These two neuronal types are thought to process distinct types of information, feedforward neurons primarily transmitting sensory input from the external world and feedback neurons modulating contextual factors such as attention, brain states, and background information (Hupé et al., 1998; Nassi et al., 2013; Keller et al., 2020; Debes and Dragoi, 2023). Despite their distinct roles, these feedforward and feedback neurons are spatially intermingled within cortical layers (Berezovskii et al., 2011). Recent advances have elucidated their functional roles (Keller et al., 2020; Fişek et al., 2023), however, it remains unclear whether feedforward and feedback projection neurons within a specific layer of a given cortical area receive distinct or similar brain-wide presynaptic inputs. Uncovering this would provide insight into their proposed functional differences.

The mouse visual cortex serves as an ideal model for investigating the brain-wide connectivity of layer- and projection-specific cell types at a cellular level. This is due to the availability of viral genetic tools that enable precise labeling of neurons based on their axonal projections (e.g., retrograde tracer such as AAVretro) and laminar identity (e.g., layer-specific Cre-expressing

transgenic mouse lines) (Gerfen et al., 2013; Tervo et al., 2016; Chatterjee et al., 2018; Matho et

al., 2021). Additionally, the hierarchical organization of mouse visual cortical areas has been

well characterized both anatomically and functionally (Harris et al., 2019; Siegle et al., 2021;

D'Souza et al., 2022). The mouse visual cortex comprises the primary visual cortex (VISp) and

at least ten different higher visual areas (HVAs), including the lateromedial (VISl) and

posteromedial (VISpm) visual cortices (Wang and Burkhalter, 2007). In the cortical hierarchy,

VISp represents the lowest level, VISl occupies an intermediate position, and VISpm is at a

higher level (Harris et al., 2019; D'Souza et al., 2022). Within this framework, CCPNs in VISl

that project to VISpm are defined as feedforward neurons, whereas those projecting to VISp are

defined as feedback neurons (D'Souza et al., 2022). This makes VISl particularly suitable for

studying spatially intermingled feedforward and feedback CCPNs.

Previous studies have mapped the monosynaptic inputs of layer-specific excitatory neuronal

types (Vélez-Fort et al., 2014; DeNardo et al., 2015; Kim et al., 2015, 2020; Yao et al., 2023a).

These studies suggest that long-distance input connectivity to specific layer-defined neuronal

classes can be either distinct or largely comparable and that local connectivity varies (DeNardo

et al., 2015; Yao et al., 2023a). However, it remains unclear whether feedforward and feedback

CCPNs within the same cortical layer receive distinct brain-wide monosynaptic inputs,

highlighting a critical gap in our understanding of cortical circuit organization.

To investigate the brain-wide monosynaptic input connectivity of layer-specific feedforward and

feedback CCPNs in the mouse visual cortex, we employed the intersectional G-deleted rabies

tracing method - cTRIO (cell-type specific inputs and outputs relationship) (Schwarz et al.,

2015). Using an intersectional approach that combined layer-specific Cre recombinase with

projection-specific tTA trans-activator expression, we defined distinct subtypes of feedforward

and feedback CCPNs in layer 2/3 (L2/3) and 5 (L5) of VISl. We then mapped their long-distance

monosynaptic inputs using rabies tracing, leveraging two avian receptors, TVA and TVA[66T], for

avian sarcoma leukosis virus glycoprotein EnvA (Wickersham et al., 2007; Miyamichi et al.,

2013). Thus we are able to trace the monosynaptic inputs to neuronal subtypes which were

previously inaccessible to brainwide mapping efforts. Our findings reveal that while the overall

input patterns to L2/3 and L5 feedforward and feedback CCPNs in VISl are largely comparable,

there are notable differences in the proportions of long-distance inputs from the visual, auditory,

and retrosplenial cortices. The experimental design presented here provides a framework for

investigating hierarchical brain-wide networks across different species and brain structures

beyond the mouse visual cortex. This study also provides a comprehensive presynaptic input

connectivity map that will serve as a foundation for future experimental and theoretical

investigations into cortico-cortical neuronal functions.

MATERIALS AND METHODS

Resource and materials availability

Further information and requests for resources and materials should be directed to and will be

fulfilled by the lead contact, Euiseok J. Kim (ekim62@ucsc.edu). Microscopy and any additional

data reported in this paper will be shared by the lead contact upon reasonable request.

Experimental animals

GENSAT BAC transgenic *SepW1-Cre* NP39 and *Tlx3-Cre* PL56 mouse lines have been

previously described (Gerfen et al., 2013). The transgenic mice were maintained on C57BL/6J

backgrounds. C57BL/6J mice were used as wild-type. Both male and female mice were used. All

mice were housed with a 12-hour light and 12-hour dark cycle and *ad libitum* access to food and

water. All animal procedures were performed in accordance with the University of California,

Santa Cruz animal care and use committee (IACUC)'s regulations.

Virus preparation

All adeno-associated viruses (AAVs) and SAD B19 EnvA-pseudotyped glycoprotein (G)-deleted

rabies virus (EnvA+RVdG) were produced by the Salk Viral Core GT3: scAAVretro-hSyn-BFP

(1.26×10^{12} GC/ml), scAAVretro-hSyn-H2B-eGFP (1.16×10^{13} GC/ml), scAAVretro-hSyn-H2B-

mCherry (1.21×10^{13} GC/ml), AAVretro-nEF-lox66/71-tTA (4.8×10^{13} GC/ml), AAV8-TRE-

DIO-oG (5.92×10^{12} GC/ml), AAV8-TRE-DIO-eGFP-T2A-TVA (7.00×10^{13} GC/ml), AAV8-

TRE-DIO-TVA^{66T}eGFP-P2A-oG (7.40×10^{13} GC/ml), and EnvA+RVdG-mCherry (7.71×10^{8}-

1.43×10^{10}) Infectious Unit (IU)/ml.

Animal surgery for virus injection

For all surgeries anesthesia induction was with 100 mg/kg of ketamine and 10 mg/kg of xylazine

cocktail via intraperitoneal injections, and anesthesia was maintained as needed with ~1%

isoflurane. Mice were mounted in a stereotax (RWD instruments) for surgery and stereotaxic

injections. For rabies tracing experiments, *SepW1-Cre* NP39 or *Tlx3-Cre* PL56 mice received

AAV helper injections at postnatal day (P)63-P125. 50-200 nl of AAVretro-nEF-lox66/71-tTA

was injected into the center of the medial secondary visual cortex (VISpm), primary visual

cortex (VISp), or anterior cingulate area (ACA) using the following coordinates: 1.6 mm rostral,

1.5 mm lateral relative to lambda and 0.4-0.6 mm ventral from the pia for VISpm or 1.1 mm

rostral, 2.6 mm lateral to lambda and 0.45-0.67 mm ventral from the pia for VISp, 0.4 mm

rostral, 0.4 mm lateral to bregma and 0.7-0.8 mm ventral from the pia for ACA. A 50 nl mixture

of 9:1 AAV8-TRE-DIO-oG and AAV8-TRE-DIO-eGFP-T2A-TVA, a 50 nl mixture of 2:7:1

AAV8-TRE-DIO-oG, 1x PBS and AAV8-TRE-DIO-eGFP-T2A-TVA, or 50 nl of AAV8-TRE-

DIO-TVA^{66T}eGFP-P2A-oG was injected into the center of the lateral secondary visual cortex

(VISl), using the following coordinates: 0.7 mm rostral, 3.65 mm lateral relative to lambda and

0.4-0.6 mm ventral from the pia. We injected AAVs using air pressure using a 1 ml syringe with

18G tubing adaptor and tubing. To prevent virus backflow, the pipette was left in the brain for 5-

10 minutes after the completion of the injection. After recovery, mice were given water with

ibuprofen (30 mg/kg) and subcutaneous carprofen (5 mg/kg). Two weeks after AAV helper

injection, 100 nl of EnvA+RVdG-mCherry was injected into the same site in VISl using a 1 ml

syringe-mediated air pressure. After recovery, mice were again given water with ibuprofen (30

mg/kg) and subcutaneous carprofen (5 mg/kg) and housed for seven days to allow for trans-

synaptic rabies spread and fluorescent protein expression.

For retrograde colocalization studies, mice were prepared and head-fixed as previously with the

same coordinates for VISpm, VISp, and ACA as above. For triple AAV colocalization

experiments, combination of 50-100 nl of scAAVretro-hSyn-BFP, scAAVretro-hSyn-H2B-

mCherry, scAAVretro-hSyn-H2B-eGFP were injected into one of VISp, VISpm, and ACA.

AAV was left to express for 18-21 days before the tissue harvest and histology. For

VISpm/VISp/ACA chemical tracer colocalization experiments, 50-100 nl of cholera toxin

subunit B (CTB) conjugated to Alexa fluor 647 (CTB-647, 1 mg/ml, Invitrogen C34778), Alexa

fluor 555 (CTB-555, 1 mg/ml, Invitrogen C22843) and/or Alexa fluor 488 (CTB-488, 1 mg/ml,

Invitrogen C34775) was injected into VISpm, VISp, and/or ACA. Three to six days later, the

tissue was harvested for histology. Different fluorophore colors were rotated between the three

injection sites across different animals. Detailed information of injections on individual mice can

be found in Table 1.

Histology and image analysis

Brains were harvested after trans-cardiac perfusion using phosphate-buffered saline (PBS)

followed by 4% paraformaldehyde (PFA). Brains were dissected out from skulls and post-fixed

with 2% PFA and 15% sucrose in PBS at 4°C overnight and then immersed in 30% sucrose in

PBS at 4°C before sectioning. Using a freezing microtome, 50-100 μm coronal brain sections

were cut and stored in cryogen (30% ethylene glycol, 30% glycerol, 30% ddH$_2$0, 10% 10x PBS)

at 4°C, every other section was stained and mounted for data analysis and quantification. Before

staining, sections were washed 3x 10 minutes in PBS and permeabilized and blocked in PBS/5%

normal donkey serum (Millipore S30-100ML)/1% Triton-X 100 (Sigma-Aldrich X100-100ML)

for two hours at room temperature. To enhance eGFP and mCherry signals, free-floating sections

were incubated at 4°C overnight with goat anti-GFP (1:1000; Rockland 600-101-215,

RRID:AB_218182) and rabbit anti-dsRed (1:500; Clontech 632496, RRID:AB_10013483)

primary antibodies in PBS/0.5% normal donkey serum/0.1% Triton-X 100, followed by 3x 10

minutes in PBS wash and incubation with the appropriate secondary antibodies conjugated with

Alexa 488 or 568 (1:500; Invitrogen A-11055 (RRID:AB_2534102) or A-10042

(RRID:AB_2534017), respectively). Sections were counterstained with 10 μM DAPI (4',6-

diamidino-2-phenylindole) in PBS for 30 minutes to visualize cell nuclei, followed by 3x 10

minutes in PBS wash. Immunostained tissue sections were mounted on slides with a polyvinyl

alcohol mounting medium containing DABCO (PVA-DABCO) and allowed to air-dry overnight.

All sections were scanned with a 10x/0.45 NA (wd-2.0 mm) objective on a Zeiss AxioImager Z2

Widefield Microscope using tile scan and Z-stack (every 6-8 μm). Representative images were

taken with the Zeiss 880 Confocal Microscope with 10x/0.45 NA (wd-2.0 mm) or 40x/ 0.95 NA

corr (wd-0.25 mm) objectives. Scanned images were first processed in Zeiss ZEN 3.11 with

DAPI images for alignment and fluorescent images for eGFP+ or mCherry+ cell detections

exported for processing. Subsequent image files were processed and analyzed using Bell Jar for

long-distance input mapping (Soronow et al., 2025). For alignment, the center Z-stack DAPI

images were used to predict the section's xy coordinates and cut angle for alignment to the Allen

Brain Common Coordinate Framework reference atlas (Wang et al., 2020). Alignments were

then fine-tuned in Bell Jar to ensure fit and minimize counting rabies-labeled cells resulting from

minor TVA expression leakage in non-Cre-expressing neurons (Wall et al., 2010). Layers for

VISpm and VISp were also manually fine-tuned to ensure accuracy. The exported Zeiss ZEN

files with mCherry+ signals were used for automated cell counting. Using Bell Jar, max

projection images were produced from the Z-stacks, and these max projections were then

sharpened in the program before detection. Detection thresholds were kept stable unless manual

counting validation showed less than 90% accuracy of cell labeling, in which case the confidence

interval was adjusted (three out of 15 brains) to ensure accurate counting. Cell detection and

brain alignment were then integrated, and the long-distance ipsi-hemispheric cortical input

fractions were calculated and graphed using GraphPad Prism 10.1.1. For thalamic input

proportions, areas and cells were drawn and counted manually on Zeiss ZEN 3.11 using the

Allen Brain Common Coordinate Framework reference atlas (Wang et al., 2020).

For VISpm/VISp/ACA CTB colocalization experiments, all CTB labeled cells were counted

manually due to weaker fluorescent signals that could not reliably be detected using Bell Jar's

automated cell detection. For each animal, three sections corresponding to the same three VISl

containing plates from the Allen CCFv3 were selected for counting. Using Fiji, VISl and laminar

borders were drawn over DAPI images, cells were labelled individually and then overlayed

labels were checked for co-localization.

Quantification and Statistical analysis

The values of n and what n represents are reported in results and figure legends. For statistical

significance, two-way ANOVA with Tukey's multiple comparisons test was performed with

GraphPad Prism 10.1.1. A one-way ANOVA with the Fisher's Least Significant Difference

(LSD) post hoc for multiple comparisons test was performed using IBM SPSS 30. Unless

otherwise noted, statistical significance is indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p <$

0.001; ****$p < 0.0001$. In Fig. 4C, significance is denoted differently: *(black) $p < 0.05$ and

*(magenta) $p < 0.01$.

RESULTS

Feedforward and feedback cortico-cortical projection neurons of VISl are largely non-

overlapping and distributed across layers

We first investigated the long-distance projection targets of CCPNs in VISl using published data

from the Allen Mouse Brain Connectivity Atlas (Harris et al., 2019). This dataset provides high-

resolution maps of eGFP+ axonal projections in adult mouse brains, generated by injecting

AAV1-CAG-FLEx-eGFP into Cre-expressing transgenic mice to label specific neuronal

populations in different brain regions. To examine CCPNs in VISl, we analyzed data from *Cux2-*

IRES-Cre and *Tlx3-Cre* mice, which label layers 2-4 (L2-4) and L5 CCPNs, respectively. Whole-

brain projection maps revealed that both L2-4 and L5 CCPNs extend axons to more than 13

cortical areas, including visual areas as VISp, VISpm, and VISpor and non-visual areas such as

the temporal association cortex (TEa) and the anterior cingulate area, dorsal part (ACAd) (Fig.

1A-B). Quantitative analysis of the axonal projection volumes across these regions showed that

L2-4 and L5 CCPNs exhibit similar projection patterns, with the highest projection volume in

VISp ($39.43 \pm 3.20\%$ in *Cux2-IRES-Cre* and $29.75 \pm 3.80\%$ in *Tlx3-Cre*, mean $\pm$ SEM),

followed by VISpm, VISpor, VISli, and VISrl (Fig. 1B). Notably, L2-4 CCPNs exhibit sparse

axonal projections to ACAs (1.10 ± 0.28%), whereas 3.55 ± 0.95% of the total axonal

projections from L5 CCPNs target ACAd (Fig. 1B).

Next, we investigated whether CCPNs in VISl that project to three major cortical target areas

(VISp, ACA/MOs (secondary motor cortex), and VISpm) represent distinct or overlapping cell

types at the single-cell level. To retrogradely label these projection neurons with three different

fluorescent markers, we stereotaxically injected AAVretro-hSyn-H2B-mCherry into VISp,

AAVretro-hSyn-BFP into ACA/MOs, and AAVretro-hSyn-H2B-eGFP into VISpm in wild-type

adult C57BL/6J mice (Fig. 1C). Thus, VISl neurons projecting to VISpm, VISp, and ACA/MOs,

designated as VISl→VISpm, VISl→VISp, and VISl→ACA, respectively. In VISl of these mice,

labeled CCPNs were distributed across layers 2-6. The majority of neurons were labeled by a

single fluorescent protein (VISl→VISpm: 90.97 ± 2.08%; VISl→VISp: 92.25 ± 5.04%;

VISl→ACA: 87.94 ± 4.60%, n = 2 mice) with relatively few exhibiting double or triple labeling

(Fig. 1C). These findings indicate that CCPNs projecting to VISpm or ACA/MOs (feedforward,

FF) and VISp (feedback, FB) do not extensively share axonal collaterals consistent with a

previous study (Berezovskii et al., 2011).

To evaluate whether this observation is consistent when using chemical tracers that avoid

potential viral co-transduction effects of AAVretros, we stereotaxically injected 50 nl of cholera

toxin subunit B (CTB) conjugated to Alexa Fluor 647, 555, or 488 into the ACA, VISp, and

VISpm, respectively, alternating fluorophores across animals. In VISl of these triple CTB

injected mice, most neurons were labeled by a single fluorescent CTB (VISl→ACA: 75.11 ±

10.04%; VISl→VISp: 95.93 ± 1.06%; VISl→VISpm: 85.00 ± 12.92%, n= 4 mice), with

relatively few neurons exhibiting double or triple labeling (Fig. 1E, F). In a separate set of

experiments, we injected 100 nl of CTB conjugated to Alexa Fluor 647 or 555 into VISp or

VISpm, respectively, to maximize labeling of neurons projecting to these areas while minimizing

unintended spread to adjacent regions. Even under these double injection conditions, the majority

of VISl neurons were labeled by a single fluorescent CTB (VISl→VISp: 94.40 ± 1.05%;

VISl→VISpm: 94.95 ± 0.87%, n=3 mice) (Fig. 1E). Therefore, our results indicate that

VISl→VISpm, VISl→VISp, and VISl→ACA CCPNs are largely distinct neuronal

subpopulations.

CCPNs are distributed across all cortical layers from L2/3 to L6. To determine the laminar

distribution of the three VISl CCPN types projecting to ACA, VISp, and VISpm, we quantified

their soma locations as percentages across layers (Fig. 1G). All three types were present

throughout L2-6, with the highest proportions in L5 (VISl→ACA: 41.50 ± 8.62%; VISl→VISp:

31.82 ± 3.97%; VISl→VISpm: 45.07 ± 7.29%; n = 3 mice, Fig. 1G). Notably, VISl→ACA

CCPNs were more abundant in the upper division of L2/3 (L2/3^{upper}) compared to the lower

division (L2/3^{lower}) (28.56 ± 3.91% vs. 9.08 ± 5.48%, Fig. 1G). VISl→VISpm CCPNs showed a

similar trend, with higher proportions in L2/3^{upper} than L2/3^{lower} (12.29 ± 1.77% vs. 7.93 ±

2.87%, Fig. 1G). In contrast, VISl→VISp CCPNs were more abundant in L2/3^{lower} compared to

L2/3^{upper} (14.66 ± 2.67% vs. 9.58 ± 3.40%, Fig. 1G). Despite these laminar differences, all three

CCPN types displayed substantial overlap across the full depth of L2/3 and within L5.

Brain-wide monosynaptic input mapping of feedforward and feedback CCPNs in VISl

We examined whether long-distance inputs to feedforward and feedback neurons align with one

of three possible organizational models (Fig. 2A). In Model 1, these neurons receive inputs from

the same areas they project to, establishing a reciprocal connectivity pattern that facilitates

looped computations. In Model 2, feedforward and feedback neurons differ in the type of

information they relay; feedforward neurons preferentially receive inputs from lower sensory

areas such as VISp and the dorsal lateral geniculate nucleus of the thalamus (LGd), while

feedback neurons receive inputs from higher cortical areas that convey contextual information

such as higher cortical areas and the lateral posterior thalamic nucleus (LP). In Model 3, both

feedforward and feedback neurons receive inputs from the same brain regions, making them

anatomically indistinguishable. In this scenario, despite similar long-distance anatomical inputs,

feedforward and feedback neurons may exhibit distinct functional roles due to differences in

local circuit computations. Alternatively, the long-distance inputs themselves may differ

functionally, even if their anatomical connectivity proportion from each brain region appears

similar.

To differentiate between these three models, we mapped and characterized the brain-wide long-

distance inputs to distinct feedforward and feedback CCPNs in VISl. We implemented an

intersectional rabies tracing strategy called cTRIO to map monosynaptic inputs in a projection-

and layer-specific manner. We first injected a helper virus mixture of AAV8-TRE-DIO-oG

(rabies glycoprotein) (Kim et al., 2016) and AAV8-TRE-DIO-eGFP-T2A-TVA into VISl, along

with AAVretro-nEF-lox66/71-tTA (hereafter AAVretro-tTA) into projection targets, including

VISpm, VISp, and ACA, in L2/3-specific *SepW1-Cre* or L5 CCPN-specific *Tlx3-Cre* transgenic

mice (Gerfen et al., 2013). This ensured Cre- and tTA-dependent expression of the avian TVA

receptor and rabies glycoprotein (oG). This approach enabled selective labeling of feedforward

and feedback neurons in L2/3 or L5 (Fig. 2D). 14 days later, EnvA-pseudotyped glycoprotein

(G)-deleted rabies virus (EnvA+RVdG)-mCherry was injected into the same VISl location to

enable monosynaptic input tracing from the starter neurons. After seven days, brains were

harvested, sectioned, immunostained with goat anti-eGFP (Rockland 600-101-215,

RRID:AB_218182) and rabbit anti-dsRed (Clontech 632496, RRID:AB_10013483) antibodies,

and imaged to quantify brain-wide long-distance inputs.

Control experiments to optimize layer- and projection-specific monosynaptic rabies tracing

We conducted a series of experiments to optimize our monosynaptic rabies tracing approach. Our goal was to minimize false-positive long-distance labeling and prevent contamination of the starter neuron population. Such issues could arise from 'leaky' expression of transgenes such as TVA and oG independent of Cre and tTA activity (Wall et al., 2010; Lavin et al., 2020), as well as from unintended damage to long-distance axonal tracts in the white matter (Watakabe and Hirokawa, 2018) (Fig. 3).

To examine the absence of false-positive long-distance rabies labeling, we first injected AAV8-TRE-DIO-eGFP-T2A-TVA (without oG) into VISl and AAVretro-tTA into VISp of *Tlx3-Cre* mice (Fig. 3A). 14 days later, we administered EnvA+RVdG-mCherry into VISl and harvested the brains seven days post-injection (Fig. 3A). As expected, in the absence of oG, no mCherry+ rabies-labeled neurons were detected outside of the injection site (VISl) (Fig. 3B-B').

In additional controls, we injected either a mixture of AAV8-TRE-DIO-eGFP-T2A-TVA and AAV8-TRE-DIO-oG (Fig. 3C), or AAV8-TRE-DIO-TVA^{66T}eGFP-P2A-oG alone (Fig. 3E) into VISl or VISp, but without AAVretro-tTA injections. In both cases, following EnvA+RVdG-mCherry and a seven-day incubation period, no mCherry+ rabies-labeled neurons outside of the injection site (Fig. 3C-D', E-F')

Next, to assess the purity of starter neurons and verify whether L2/3 or L5 VISl→VISp CCPNs contained only VISl→VISp neurons, we injected AAV8-TRE-DIO-TVA^{66T}eGFP-P2A-oG into VISl and a mixture of AAVretro-tTA and Alexa Fluor 647-conjugated cholera toxin subunit B (CTB) into VISp of *Tlx3-Cre* mice (Fig. 3G-H), and collected brains three weeks later. Confocal imaging showed that TVA^{66T}eGFP-P2A-oG labeled L5 neurons in VISl were also labeled with CTB, showing that they were VISl→VISp neurons (Fig. 3H).

To determine whether VISl→VISpm neurons were included in the L2/3 VISl→VISp starter

population due to passing axons in the grey matter of lower cortical layers (Watakabe and

Hirokawa, 2018), we injected AAV8-TRE-DIO-TVA^{66T}eGFP-P2A-oG into VISl, AAVretro-tTA

into VISp, and CTB into VISpm of *SepW1-Cre* mice (Fig. 3I-J). Confocal images of brain

sections harvested three weeks later showed that TVA^{66T}eGFP-P2A-oG expressing neurons were

not labeled with CTB, indicating that virus-labeled starter neurons did not include VISl→VISpm

neurons (Fig. 3J). These experiments validate our approach, ensuring specificity in identifying

monosynaptic inputs while preventing false positives and potential starter population

contamination.

Long-distance cortical inputs onto four different CCPN types in VISl

We examined whether the four types of feedforward and feedback CCPNs (L2/3 VISl→VISpm,

L2/3 VISl→VISp, L5 VISl→VISpm, L5 VISl→VISp) in VISl receive similar or distinct

monosynaptic inputs from other cortical areas within the ipsilateral hemisphere (Fig. 4 and Table

2). Rabies-labeled long-distance input neurons to those four types of CCPNs were distributed

throughout both visual and non-visual cortical areas (Fig. 4A), with the highest proportion

originating from VISp (L2/3 VISl→VISpm 0.409 ± 0.046, L2/3 VISl→VISp 0.380 ± 0.030, L5

L5 VISl→VISpm 0.492 ± 0.016, L5 VISl→VISp 0.582 ± 0.023, Fig. 4B). We quantified and

ranked the top 20 cortical regions contributing inputs along the cortical hierarchy (Harris et al.,

2019). While overall input patterns among the four CCPN types were largely similar, we

identified key differences. L5 VISl→VISp feedback neurons received a greater proportion of

long-distance inputs from VISp compared to L5 VISl→VISpm feedforward neurons (Fig. 4B).

Similarly, L2/3 VISl→VISpm feedforward neurons received a significantly greater proportion of

their inputs from VISpm than L2/3 VISl→VISp feedback neurons (Fig. 4C).

Further statistical analysis using one-way ANOVA revealed that at least one of the four CCPN

types exhibited a significantly different proportional input than other CCPNs in eight out of the

20 cortical areas analyzed (Fig. 4C): VISpm, VISpl, AUDd, AUDp, RSPv, RSPagl, RSPd, and

MOs. Overall, these findings suggest that while the four CCPN types receive broadly

comparable long-distance cortical inputs, they exhibit distinct preferences for specific cortical

regions, reflecting their functional roles in cortical processing. Notably, L5 VISl→VISpm

CCPNs receive proportionally more retrosplenial cortex input, suggesting a greater role in visuo-

spatial memory, and L2/3 CCPNs regardless of projection receive proportionally more auditory

cortex input suggesting a greater role for L2/3 CCPNs in audio-visual integration (Fig. 4B-C).

We next investigated the laminar distribution of long-distance presynaptic inputs from VISp and

VISpm to four distinct types of feedforward and feedback CCPNs in VISl. Although prior

studies have reported preferential long-distance connectivity between superficial layers and

between deep layers (DeNardo et al., 2015; Kim et al., 2015), it remains unclear whether CCPNs

in specific layers of a given cortical area receive layer-specific long-distance inputs, depending

on their feedforward or feedback projections. To address this, we quantified the proportion of

input neurons across each layer of VISp and VISpm (Fig. 5A-B) for four distinct CCPN types in

VISl: L2/3 VISl→VISpm, L2/3 VISl→VISp, L5 VISl→VISpm, and L5 VISl→VISp. With a

few exceptions, the laminar distribution of input neurons in VISp and VISpm did not reveal

statistically significant differences across these neuronal groups (Fig. 5A-B).

We investigated whether feedforward neurons in VISl that project to cortical areas with higher

hierarchy scores than VISpm exhibit more distinct input connectivity patterns from those of

VISl→VISp compared to VISl→VISpm neurons. Previous studies have shown that the cortical

hierarchy in rodents is relatively shallow compared to that of primates (Harris et al., 2019;

D'Souza et al., 2022). The hierarchy scores based on layer-termination patterns of cortico-

cortical connection (Harris et al., 2019) for select cortical areas are as follows: VISp: -0.239,

VISl: -0.148, VISpm: 0.074, ACAd: -0.060, ACAv: 0.404, and MO: 0.282., To determine if the

shallow hierarchy in the mouse visual cortex was masking hierarchy dependent input differences,

we injected AAVretro-tTA into ACA/MOs and a mixture of AAV8-TRE-DIO-oG and AAV8-

TRE-DIO-eGFP-T2A-TVA into VISl of *Tlx3-Cre* mice to label VISl→ACA/MOs (hereafter,

VISl→ACA) neurons and traced their monosynaptic inputs across the cortex. We then compared

these results with monosynaptic input tracing of L5 VISl→VISpm feedforward neurons and L5

VISl→VISp feedback neurons (Fig. 5C). We found broadly similar input connectivity profiles

across these three L5 CCPN subtypes. Thus, the input connectivity of L5 VISl→ACA CCPNs is

comparable to that of VISl→VISpm CCPNs and does not exhibit greater differences from

VISl→VISp CCPNs than from VISl→VISpm CCPNs. Notably, the proportion of inputs from

VISp to VISl→VISp neurons was higher than that to VISl→VISpm and VISl→ACA neurons

(Fig. 5C).

Long-distance thalamic input neuron profiles onto four different CCPN types in VISl

Next, we examined whether the four types of feedforward and feedback CCPNs in VISl receive

distinct presynaptic inputs from visual thalamic regions. In the mouse visual system, LGd

projects to VISl and other higher visual cortical areas in addition to VISp, while LP and the

laterodorsal nucleus of the thalamus (LD) also provide input to VISl and other visual cortical

regions (Bienkowski et al., 2019).

In rabies-traced brain sections, we identified three thalamic nuclei, LGd, LD, and LP, and

quantified the proportion of rabies-labeled neurons among all these input areas (Fig. 6A). Across

all four CCPN types, LP provided the majority of thalamic inputs, followed by LGd and LD,

consistent with a previous study (Yao et al., 2023a) (Fig. 6B). While the overall proportions of

thalamic inputs were similar, we observed that L2/3 VISl→VISpm neurons received a lower

proportion of inputs from LGd compared to L5 VISl→VISpm neurons (L2/3 VISl→VISpm:

0.102 ± 0.008 and L5 VISl→VISpm 0.323 ± 0.027, p=0.0030, two-way ANOVA with Tukey's

multiple comparisons test) (Fig. 6B). Conversely, L2/3 VISl→VISpm neurons received a higher

proportion of inputs from LP than L5 VISl→VISpm neurons (L2/3 VISl→VISpm 0.887 ± 0.004,

L5 VISl→VISpm 0.657 ± 0.024, p=0.0021, two-way ANOVA with Tukey's multiple

comparisons test) (Fig. 6B).

DISCUSSION

Using a layer- and projection-specific monosynaptic rabies tracing approach, we mapped brain-

wide inputs to five distinct types of feedforward and feedback CCPNs in VISl. This study

provides the first comprehensive connectivity maps to investigate the organizational principles

governing feedforward and feedback neurons in relation to their laminar positions and the

hierarchical positions of their target areas. Our findings indicate that long distance input

connectivity patterns are largely comparable across CCPN subtypes, supporting Model 3 (Fig.

2A), but notable differences emerge based on their laminar and projection identities. First, L2/3

VISl→VISpm CCPNs received a higher proportion of inputs from their own projection target,

VISpm, compared to L2/3 VISl→VISp and other L5 CCPNs. Similarly, L5 VISl→VISp CCPNs

received a higher proportion of inputs from VISp, their projection target, compared to L5

VISl→VISpm or VISl→ACA CCPNs. These results are in accordance with Model 1 (Fig. 2A),

in which monosynaptic inputs preferentially originate from the neurons' axonal targets (Kim et

al., 2020; Siu et al., 2021). These insights provide a foundation for future functional and

theoretical studies aimed at understanding brain-wide cortical computations at the cellular level.

444 The input-output organization of neural networks provides crucial insights into the information

445 they receive and the signals they transmit for functional processing. Therefore, understanding

446 neuronal connectivity patterns is essential for a comprehensive understanding of neural circuit

447 function and computation (Zeng and Sanes, 2017). Previous studies have demonstrated that

448 distinct neuronal cell types in the mammalian brain exhibit characteristic input connectivity

449 patterns. Even within broader neuronal classes, spatially intermingled subtypes display subtype-

450 specific input connectivity (e.g., dopaminergic D1 receptor versus D2 receptor expressing spiny

451 projection neurons in the striatum or serotonergic neuron subtypes in the dorsal raphe; see also

452 whole-brain rabies tracing data) (Wall et al., 2016; Ren et al., 2018). A recent study found that

453 functionally distinct neuronal types in layer 2/3 of the mouse somatosensory cortex exhibit

454 distinct brain-wide input connectivity patterns (Inácio et al., 2025). For cortico-cortical

455 connectivity in the visual cortices, a previous study found that VISp CCPNs projecting to VISpm

456 or VISal received inputs from the same respective areas, suggesting a like-to-like connectivity

457 pattern (Kim et al., 2020). A previous study observed laminar differences in like-to-like

458 connectivity using Channelrhodopsin-2-assisted circuit mapping, where there is stronger like-to-

459 like connectivity in L5 compared to L2/3 neurons (Young et al., 2021). This observation of like-

460 to-like connectivity is further supported by a study in macaques, where V1 neurons projecting to

461 V2 receive monosynaptic feedback inputs from V2 but not from other V1-projecting areas (Siu

462 et al., 2021). In the mouse VISl, the secondary visual cortex, projection-specific input

463 connectivity is observed, though it appears less pronounced than in CCPNs of VISp, the primary

464 visual cortex. One possible explanation is that VISl→VISpm and VISl→VISp neurons share

465 similar medial axonal projection trajectories, whereas VISp→VISpm and VISp→VISal (VISp

466 CCPNs projecting to anterolateral, AL, visual cortex) neurons exhibit medial versus lateral

467 projection patterns. Another, not mutually exclusive, possibility is that the cortical hierarchy in

the mouse visual system is relatively shallow, making it more challenging to detect strong

hierarchical connectivity features.

Although little is known about LP and LGd projections to VISl, our observed LP:LGd input

ratios are largely consistent with previous rabies-tracing data examining thalamic inputs to layer-

specific VISl populations (Yao et al., 2023a). Notably, we found that VISl L5 FF neurons

projecting to VISpm receive proportionally more LGd input compared to their L2/3 counterparts,

suggesting a Model 2 input organization in which FF neurons receive stronger input from

feedback-associated areas such as LGd. However, this pattern is not observed in L2/3 FF

neurons, indicating a non-symmetrical input organization between FF and FB neurons across

layers. Importantly, our data describe input proportions rather than functional strengths; future

quantitative and functional studies, such as electrophysiological recordings or other synaptic

connectivity measures, will be necessary to clarify the strength and dynamics of these

thalamocortical connections.

To validate the projection identities of our starter neurons, we performed CTB injections into

both target and non-target projection areas (Fig. 3G-J). However, given the collateralization

properties of some CCPNs (Han et al., 2018; Peng et al., 2021), it is possible that a subset of

starter neurons also projects to additional areas beyond the intended targets. Notably, axonal

reconstructions and RNA barcode-based mapping (MAPseq) by Han et al. demonstrated that

approximately 88% of VISp CCPNs project to only one or two of the six higher visual areas.

Similarly, Berezovskii et al. (2011) using dextran amine tracers reported that FF (to VISp) and

FB (to VISal) CCPNs in VISl are largely non-overlapping, with less than 5% overlap. Our

observations of VISl axonal collateralization are consistent with these findings, while

acknowledging potential differences between VISl and VISp as well as variations in

experimental approaches. Nonetheless, we note that potential collateral projections of VISl

CCPNs to areas beyond ACA, VISp, and VISpm could introduce variability in rabies tracing

results and complicate efforts to restrict analyses exclusively to single projection-defined

populations.

Monosynaptic rabies tracing is a powerful approach for mapping anatomical connections across

the brain; however, this method does not provide information about synaptic strength or other

functional properties of these connections (Callaway and Luo, 2015). For example, although

feedforward and feedback CCPNs may receive similar proportions of inputs from a given area, it

remains unclear whether these input neurons are functionally distinct—for example, whether

they exhibit different receptive field properties or belong to different functional subtypes. To

gain additional insight into the functional differences and similarities between feedforward and

feedback CCPNs, future studies could complement rabies tracing with electrophysiological or

imaging approaches. Additionally, advances in high-throughput molecular profiling, including

single-cell RNA sequencing, have revealed that cortical neurons within a single layer can be

further classified into distinct subtypes based on unsupervised clustering (Tasic et al., 2018; Yao

et al., 2023b). Beyond their broad excitatory versus inhibitory or laminar identities, it would be

valuable to examine whether long-distance input neurons exhibit specific connectivity patterns

among these molecularly defined subtypes of CCPNs (Condylis et al., 2022).

It is also possible that, despite receiving similar long-distance inputs, feedforward and feedback

neurons differ in their local circuit architecture. To further test Model 3 (Fig. 2A) and uncover

additional anatomical distinctions, future studies using TVA^{66T}-dependent rabies tracing for local

inputs, combined with molecular characterization of input neurons and electrophysiological

profiling would be informative.

In this study, we used feedforward (to VISpm or ACA) and feedback (to VISp) CCPNs in VISl

as a model to investigate connectivity differences. Given that mammalian visual cortical areas

are organized into dorsal and ventral streams with strong intra-stream connectivity, mapping

inputs to feedforward and feedback neurons targeting areas within a stream, such as POR, a

higher-order ventral stream area, could offer valuable insights (D'Souza et al. 2022; Harris et al.

2019). Although POR represents an informative target for examining feedforward and feedback

connectivity, precise stereotaxic targeting remains technically challenging. Future studies

including POR could further clarify ventral stream-specific connectivity. Another key question

for future research is whether similar connectivity profiles would be observed in other cortical

regions, such as secondary somatosensory cortex projecting to primary somatosensory cortex as

feedback CCPNs and secondary somatosensory cortex projecting to primary and secondary

motor cortices as feedforward CCPNs (Yamashita et al., 2018). Given that the cortical hierarchy

in mice is relatively shallow compared to that of higher mammals (Harris et al., 2019; D'Souza

et al., 2022), it would also be interesting to explore these questions in species with a more robust

hierarchical organization, such as marmosets or macaques.

Declaration of Interests

The authors declare no competing interests that could have influenced the contents of this

publication.

Author Contributions

Richard G. Dickson performed investigations, supervised students, collected and curated data,

analyzed data, prepared visualizations, and contributed to writing the original draft and revisions.

Matthew W. Jacobs performed investigations, supervised students, and collected data. John M.

Ratliff performed investigations, collected and analyzed data, and contributed to writing and

revisions. Alec L. R. Soronow performed investigations. Faye An and Walid A. Yuqob collected

and analyzed data. Euiseok J. Kim conceived and designed the study, supervised the project,

analyzed and visualized data, contributed resources, acquired funding, and contributed to writing

the original draft and revisions. All authors had full access to the data, take responsibility for its

integrity, and approved the final manuscript.

FIGURE LEGENDS

Figure 1. Whole-brain axonal projections and largely segregated feedforward and feedback CCPNs in VISl. (A) Top-down views of whole-brain 3D reconstructions of axonal projections from VISl. (Top) L2-4 CCPNs in a *Cux2-IRES-Cre* brain (Experiment: 287495026). (Bottom) L5 CCPNs in a *Tlx3-Cre* brain (Experiment: 520750912). Data adapted from the Allen Brain Connectivity Atlas (https://connectivity.brain-map.org/projection). **(B)** Percentage of total axon projection volume across 13 cortical areas from L2-4 VISl CCPNs in *Cux2-IRES-Cre* (magenta, n = 2 mice) and L5 VISl CCPNs in *Tlx3-Cre* (cyan, n = 3 mice). **(C)** Confocal microscope images showing the injection sites of AAVretro-H2B-eGFP, AAVretro-H2B-mCherry, and AAVretro-BFP targeting ACAd/MO, VISp, and VISpm, respectively. Allen Reference Atlas (ARA) section numbers are in the top left corner. **(D)** Representative confocal microscope images showing VISl CCPNs retrogradely labeled based on their axonal projections to ACAd/MO, VISp, or VISpm. Arrows indicate neurons that project to more than one of these target areas. **(E)** Proportions of VISl neurons projecting to multiple target areas. Triple CTB injection experiments show the fraction of VISl neurons projecting to single, double, or all three target areas (ACA, VISp, VISpm). Counts per CCPN type across all animals were: 94 cells for VISl→ACA, 523 cells for VISl→VISp, and 573 cells for VISl→VISpm (n =4 mice). Double CTB injections show the fractions of VISl neurons projecting to one or both target areas (VISp or VISpm). Counts per CCPN type across all animals were: 601 cells for VISl→VISp and 763 cells for VISl→VISpm (n = 3 mice). **(F)** Representative confocal images of CTB-labeled VISl neurons. Top row: Injection sites of CTB tracers conjugated to three fluorophores targeting ACA (yellow), VISp (magenta), and VISpm (cyan); ARA section numbers are indicated above each panel. Middle and bottom rows: VISl CCPNs retrogradely labeled according to their projections. Yellow arrows highlight neurons projecting to ACA, and white arrows indicate neurons projecting to ACA, VISp, and VISpm. **(G)** Percentage distribution of the somata of three VISl CCPN types across cortical layers. A total of 1190 retrograde labeled neurons were analyzed in 6 mice. Data are represented as mean ± SEM.

Figure 2. Long-distance connectivity models and projection- and layer-specific rabies tracing strategy. (A) Three proposed models of long-distance monosynaptic input connectivity to feedforward (FF) and feedback (FB) neurons. **(B)** Schematic of the intersectional rabies tracing strategy for projection- and layer-specific long-distance input tracing to VISl neurons. **(C)** Representative scanning microscope images overlaid with the Allen Brain Common Coordinate Framework reference atlas (Wang et al., 2020), showing AAVretro-tTA targeting in three different areas: VISpm, VISp, and ACA. **(D)** Representative confocal microscope images showing eGFP+mCherry+ starter neurons (arrowheads) in VISl projecting to VISp, VISpm, and ACA in *SepW1-Cre* or *Tlx3-Cre* mice.

Figure 3. Control brains. (A) Schematic of the viral genetic tracing strategy for control injections without oG (rabies glycoprotein). **(B)** Fluorescence microscope images showing the absence of labeled cells outside the injection sites **(B')**. **(C,E)** Schematic of the viral genetic tracing strategy for control injections without tTA. **(D-D', F-F')** Fluorescence microscope images showing the absence of labeled cells outside the injection sites **(D', F')**. **(G, I)** Schematic of the viral genetic tracing strategy used to verify the axonal projection targets of labeled starter neurons (TVA^{66T}eGFP-P2A-oG+) by CTB injections into either the expected projection target, VISp **(G-H)**, or a non-target area, VISpm **(I-J)**. **(H)** Yellow arrowheads indicate starter neurons co-labeled with eGFP (cyan) and CTB (magenta), indicating their axonal projections to VISp.

Figure 4. Long-distance input mapping of layer- and projection-specific CCPNs in VISl. (A) Representative confocal images showing long-distance mCherry+ input neurons in LP, VISp, VISpm, and RSP for four distinct feedforward and feedback CCPN types in VISl of *SepW1-Cre* or *Tlx3-Cre* mice. **(B)** The ipsilateral cortical proportion of input neurons across 20 cortical areas, sorted by cortical hierarchy (Harris et al., 2019). The red arrowhead indicates the rank of VISl along the cortical hierarchy. n = 3 mice per each group. Two-way ANOVA with Tukey's multiple comparisons test, * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001. **(C)** Results of one-way ANOVA with Fisher's Least Significant Difference

(LSD) post-hoc test for eight selected input regions. *(black) p<0.05, *(magenta) p<0.01. Data are
represented as mean ± SEM.

Figure 5. Further analyses of long-distance input neurons to CCPNs in VISl. (A) The distributions of
presynaptic input neurons across VISp layers to four distinct types of CCPNs in VISl. **(B)** The
distribution of presynaptic input neurons across VISpm layers to four distinct types of CCPNs in VISl.
(C) The proportion of long-distance input neurons located in each of the 20 cortical regions for the three
L5 CCPN types in VISl. n = 3 mice per each group for each figure. Two-way ANOVA with Tukey's
multiple comparisons test, * p<0.05, ** p<0.01, **** p<0.0001. Data are represented as mean ± SEM.

Figure 6. Long-distance thalamic input mapping of layer- and projection-specific CCPNs in VISl.
(A) Representative confocal microscope images showing rabies-labeled mCherry+ input neurons in LGd
or LP. **(B)** Proportions of long-distance inputs from each visual thalamic region to four distinct types of
feedforward and feedback starter neurons in VISl. n = 3 mice per each group. Two-way ANOVA with
Tukey's multiple comparisons test, * p<0.05. Data are represented as mean ± SEM.

Table 1. Animal and experimental condition summary

Table 2. Quantification of rabies tracing in individual mouse brains

Abbreviations: AAV, Adeno associated virus; ACA, Anterior cingulate area; ACAd, Anterior cingulate
area dorsal; ACAv, Anterior cingulate area ventral; AIp, Agranular insular area, posterior part; AUDd,
Dorsal auditory area; AUDp, Primary auditory area; AUDpo, Posterior auditory area; AUDv, Ventral
auditory area; CCPN, Cortico-cortical projection neurons; CTB, Cholera toxin subunit B; cTRIO, Cell
specific tracing the relationship between input and output; ECT, Ectorhinal area; ENTl, Entorhinal area,
lateral part; ENTm, Entorhinal area, medial part, dorsal zone; EnvA, Envelope protein of avian leukosis
virus; FB, feedback neurons; FF, feedforward neurons; GU, Gustatory areas; HVA, Higher visual area;
LD, Lateral dorsal nucleus of thalamus; LGd, Lateral geniculate nucleus dorsal; LP, Lateral posterior
nucleus of the thalamus; Mop, Primary motor area; Mos, Secondary motor cortex; oG, optimized rabies
glycoprotein; ORBl, Orbital area, lateral part; PBS, Phosphate buffered saline; PERI, Perirhinal area;
PFA, paraformaldehyde; RSP, Retrosplenial cortex; RSPagl, Retrosplenial cortex agranular; RSPd,
Retrosplenial cortex dorsal; RSPv, Retrosplenial cortex ventral; SSp-bfd, Primary somatosensory area,
barrel field; SSp-ll, Primary somatosensory area, lower limb; SSp-m, Primary somatosensory area,
mouth; SSp-n, Primary somatosensory area, nose; SSp-ul, Primary somatosensory area, upper limb; SSp-
un, Primary somatosensory area, unassigned; SSp-tr, Primary somatosensory area, trunk; SSs,
Supplemental somatosensory area; TEa, Temporal association area; tTa, Tetracycline trans activator;
TVA, Receptor for avian leukosis virus; VISa, Anterior area secondary visual cortex; VISal,
Anterolateral secondary visual cortex; VISam, Anteromedial secondary visual cortex; VISC, Visceral
area; VISl (LM), Lateral secondary visual cortex; VISli, Latero intermediate secondary visual cortex;
VISp, Primary visual cortex; VISpl, Posterolateral secondary visual cortex; VISpm, Posteromedial
secondary visual cortex; VISpor, Postrhinal secondary visual cortex; VISrl, Rostrolateral secondary visual
cortex.

REFERENCES

Berezovskii VK, Nassi JJ, Born RT. 2011. Segregation of feedforward and feedback projections in mouse
visual cortex. J Comp Neurol [Internet] 519:3672–3683. Available from:
http://dx.doi.org/10.1002/cne.22675

Bienkowski MS, Benavidez NL, Wu K, Gou L, Becerra M, Dong H-W. 2019. Extrastriate connectivity of
the mouse dorsal lateral geniculate thalamic nucleus. J Comp Neurol [Internet] 527:1419–1442.
Available from: http://dx.doi.org/10.1002/cne.24627

Cajal SR y. 1995. Cajal's histology of the nervous system of man and vertebrates. New York, NY:
Oxford University Press.

Callaway EM, Luo L. 2015. Monosynaptic circuit tracing with glycoprotein-deleted rabies viruses. J
Neurosci [Internet] 35:8979–8985. Available from: http://dx.doi.org/10.1523/JNEUROSCI.0409-
15.2015

Chatterjee S, Sullivan HA, MacLennan BJ, Xu R, Hou Y, Lavin TK, Lea NE, Michalski JE, Babcock KR,
Dietrich S, Matthews GA, Beyeler A, Calhoon GG, Glober G, Whitesell JD, Yao S, Cetin A, Harris
JA, Zeng H, Tye KM, Reid RC, Wickersham IR. 2018. Nontoxic, double-deletion-mutant rabies
viral vectors for retrograde targeting of projection neurons. Nat Neurosci [Internet] 21:638–646.
Available from: https://doi.org/10.1038/s41593-018-0091-7

Condylis C, Ghanbari A, Manjrekar N, Bistrong K, Yao S, Yao Z, Nguyen TN, Zeng H, Tasic B, Chen
JL. 2022. Dense functional and molecular readout of a circuit hub in sensory cortex. Science
[Internet] 375:eabl5981. Available from: http://dx.doi.org/10.1126/science.abl5981

Debes SR, Dragoi V. 2023. Suppressing feedback signals to visual cortex abolishes attentional
modulation. Science [Internet] 379:468–473. Available from:
http://dx.doi.org/10.1126/science.ade1855

DeNardo LA, Berns DS, DeLoach K, Luo L. 2015. Connectivity of mouse somatosensory and prefrontal
cortex examined with trans-synaptic tracing. Nat Neurosci [Internet] 18:1687–1697. Available from:
https://www.nature.com/articles/nn.4131

D'Souza RD, Wang Q, Ji W, Meier AM, Kennedy H, Knoblauch K, Burkhalter A. 2022. Hierarchical and
nonhierarchical features of the mouse visual cortical network. Nat Commun [Internet] 13:503.
Available from: http://dx.doi.org/10.1038/s41467-022-28035-y

Felleman DJ, Van Essen DC. 1991. Distributed hierarchical processing in the primate cerebral cortex.
Cereb Cortex [Internet] 1:1–47. Available from: http://dx.doi.org/10.1093/cercor/1.1.1-a

Fişek M, Herrmann D, Egea-Weiss A, Cloves M, Bauer L, Lee T-Y, Russell LE, Häusser M. 2023.
Cortico-cortical feedback engages active dendrites in visual cortex. Nature [Internet] 617:769–776.
Available from: http://dx.doi.org/10.1038/s41586-023-06007-6

Gămănuţ R, Kennedy H, Toroczkai Z, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Burkhalter A.
2018. The mouse cortical connectome, characterized by an ultra-dense cortical graph, maintains
specificity by distinct connectivity profiles. Neuron [Internet] 97:698–715.e10. Available from:
http://dx.doi.org/10.1016/j.neuron.2017.12.037

Gerfen CR, Paletzki R, Heintz N. 2013. GENSAT BAC cre-recombinase driver lines to study the
functional organization of cerebral cortical and basal ganglia circuits. Neuron [Internet] 80:1368–
1383. Available from: http://dx.doi.org/10.1016/j.neuron.2013.10.016

Han Y, Kebschull JM, Campbell RAA, Cowan D, Imhof F, Zador AM, Mrsic-Flogel TD. 2018. The logic
of single-cell projections from visual cortex. Nature [Internet] 556:51–56. Available from:
https://pubmed.ncbi.nlm.nih.gov/29590093/

Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, Bernard A, Bohn P, Caldejon S, Casal L, Cho
A, Feiner A, Feng D, Gaudreault N, Gerfen CR, Graddis N, Groblewski PA, Henry AM, Ho A,
Howard R, Knox JE, Kuan L, Kuang X, Lecoq J, Lesnar P, Li Y, Luviano J, McConoughey S,
Mortrud MT, Naeemi M, Ng L, Oh SW, Ouellette B, Shen E, Sorensen SA, Wakeman W, Wang Q,
Wang Y, Williford A, Phillips JW, Jones AR, Koch C, Zeng H. 2019. Hierarchical organization of
cortical and thalamic connectivity. Nature [Internet] 575:195–202. Available from:
http://dx.doi.org/10.1038/s41586-019-1716-z

Harris KD, Shepherd GMG. 2015. The neocortical circuit: themes and variations. Nat Neurosci [Internet]
18:170–181. Available from: http://dx.doi.org/10.1038/nn.3917

Hupé JM, James AC, Payne BR, Lomber SG, Girard P, Bullier J. 1998. Cortical feedback improves
discrimination between figure and background by V1, V2 and V3 neurons. Nature [Internet]
394:784–787. Available from: http://dx.doi.org/10.1038/29537

Inácio AR, Lam KC, Zhao Y, Pereira F, Gerfen CR, Lee S. 2025. Brain-wide presynaptic networks of
functionally distinct cortical neurons. Nature [Internet]:1–11. Available from:
https://www.nature.com/articles/s41586-025-08631-w

Keller AJ, Roth MM, Scanziani M. 2020. Feedback generates a second receptive field in neurons of the
visual cortex. Nature [Internet] 582:545–549. Available from: http://dx.doi.org/10.1038/s41586-020-
2319-4

Kim EJ, Jacobs MW, Ito-Cole T, Callaway EM. 2016. Improved monosynaptic neural circuit tracing
using engineered rabies virus glycoproteins. Cell Rep [Internet] 15:692–699. Available from:
http://dx.doi.org/10.1016/j.celrep.2016.03.067

Kim EJ, Juavinett AL, Kyubwa EM, Jacobs MW, Callaway EM. 2015. Three types of cortical layer 5
neurons that differ in brain-wide connectivity and function. Neuron [Internet] 88:1253–1267.
Available from: http://dx.doi.org/10.1016/j.neuron.2015.11.002

Kim EJ, Zhang Z, Huang L, Ito-Cole T, Jacobs MW, Juavinett AL, Senturk G, Hu M, Ku M, Ecker JR,
Callaway EM. 2020. Extraction of Distinct Neuronal Cell Types from within a Genetically
Continuous Population. Neuron [Internet] 107:274–282.e6. Available from:
http://dx.doi.org/10.1016/j.neuron.2020.04.018

Lavin TK, Jin L, Lea NE, Wickersham IR. 2020. Monosynaptic tracing success depends critically on
helper virus concentrations. Front Synaptic Neurosci [Internet] 12:6. Available from:
http://dx.doi.org/10.3389/fnsyn.2020.00006

Matho KS, Huilgol D, Galbavy W, He M, Kim G, An X, Lu J, Wu P, Di Bella DJ, Shetty AS,
Palaniswamy R, Hatfield J, Raudales R, Narasimhan A, Gamache E, Levine JM, Tucciarone J,
Szelenyi E, Harris JA, Mitra PP, Osten P, Arlotta P, Huang ZJ. 2021. Genetic dissection of the
glutamatergic neuron system in cerebral cortex. Nature [Internet] 598:182–187. Available from:
http://dx.doi.org/10.1038/s41586-021-03955-9

Miyamichi K, Shlomai-Fuchs Y, Shu M, Weissbourd BC, Luo L, Mizrahi A. 2013. Dissecting local
circuits: parvalbumin interneurons underlie broad feedback control of olfactory bulb output. Neuron
[Internet] 80:1232–1245. Available from: http://dx.doi.org/10.1016/j.neuron.2013.08.027

Nassi JJ, Lomber SG, Born RT. 2013. Corticocortical feedback contributes to surround suppression in V1
of the alert primate. J Neurosci [Internet] 33:8504–8517. Available from:
http://dx.doi.org/10.1523/JNEUROSCI.5124-12.2013

Oh SW, Harris JA, Ng L, Winslow B, Cain N, Mihalas S, Wang Q, Lau C, Kuan L, Henry AM, Mortrud
MT, Ouellette B, Nguyen TN, Sorensen SA, Slaughterbeck CR, Wakeman W, Li Y, Feng D, Ho A,
Nicholas E, Hirokawa KE, Bohn P, Joines KM, Peng H, Hawrylycz MJ, Phillips JW, Hohmann JG,
Wohnoutka P, Gerfen CR, Koch C, Bernard A, Dang C, Jones AR, Zeng H. 2014. A mesoscale
connectome of the mouse brain. Nature [Internet] 508:207–214. Available from:
http://dx.doi.org/10.1038/nature13186

Peng H, Xie P, Liu L, Kuang X, Wang Y, Qu L, Gong H, Jiang S, Li A, Ruan Z, Ding L, Yao Z, Chen C,
Chen M, Daigle TL, Dalley R, Ding Z, Duan Y, Feiner A, He P, Hill C, Hirokawa KE, Hong G,
Huang L, Kebede S, Kuo H-C, Larsen R, Lesnar P, Li L, Li Q, Li X, Li Y, Li Y, Liu A, Lu D, Mok
S, Ng L, Nguyen TN, Ouyang Q, Pan J, Shen E, Song Y, Sunkin SM, Tasic B, Veldman MB,
Wakeman W, Wan W, Wang P, Wang Q, Wang T, Wang Y, Xiong F, Xiong W, Xu W, Ye M, Yin
L, Yu Y, Yuan J, Yuan J, Yun Z, Zeng S, Zhang S, Zhao S, Zhao Z, Zhou Z, Huang ZJ, Esposito L,
Hawrylycz MJ, Sorensen SA, Yang XW, Zheng Y, Gu Z, Xie W, Koch C, Luo Q, Harris JA, Wang
Y, Zeng H. 2021. Morphological diversity of single neurons in molecularly defined cell types.
Nature [Internet] 598:174–181. Available from: https://doi.org/10.1038/s41586-021-03941-1

Ren J, Friedmann D, Xiong J, Liu CD, Ferguson BR, Weerakkody T, DeLoach KE, Ran C, Pun A, Sun
Y, Weissbourd B, Neve RL, Huguenard J, Horowitz MA, Luo L. 2018. Anatomically defined and
functionally distinct dorsal raphe serotonin sub-systems. Cell [Internet] 175:472–487.e20. Available
from: http://dx.doi.org/10.1016/j.cell.2018.07.043

Schwarz LA, Miyamichi K, Gao XJ, Beier KT, Weissbourd B, DeLoach KE, Ren J, Ibanes S, Malenka
RC, Kremer EJ, Luo L. 2015. Viral-genetic tracing of the input-output organization of a central
noradrenaline circuit. Nature [Internet] 524:88–92. Available from:
https://www.nature.com/articles/nature14600

Siegle JH, Jia X, Durand S, Gale S, Bennett C, Graddis N, Heller G, Ramirez TK, Choi H, Luviano JA,
Groblewski PA, Ahmed R, Arkhipov A, Bernard A, Billeh YN, Brown D, Buice MA, Cain N,
Caldejon S, Casal L, Cho A, Chvilicek M, Cox TC, Dai K, Denman DJ, de Vries SEJ, Dietzman R,
Esposito L, Farrell C, Feng D, Galbraith J, Garrett M, Gelfand EC, Hancock N, Harris JA, Howard
R, Hu B, Hytnen R, Iyer R, Jessett E, Johnson K, Kato I, Kiggins J, Lambert S, Lecoq J,
Ledochowitsch P, Lee JH, Leon A, Li Y, Liang E, Long F, Mace K, Melchior J, Millman D,
Mollenkopf T, Nayan C, Ng L, Ngo K, Nguyen T, Nicovich PR, North K, Ocker GK, Ollerenshaw
D, Oliver M, Pachitariu M, Perkins J, Reding M, Reid D, Robertson M, Ronellenfitch K, Seid S,
Slaughterbeck C, Stoecklin M, Sullivan D, Sutton B, Swapp J, Thompson C, Turner K, Wakeman
W, Whitesell JD, Williams D, Williford A, Young R, Zeng H, Naylor S, Phillips JW, Reid RC,
Mihalas S, Olsen SR, Koch C. 2021. Survey of spiking in the mouse visual system reveals functional
hierarchy. Nature [Internet] 592:86–92. Available from: http://dx.doi.org/10.1038/s41586-020-
03171-x

Siu C, Balsor J, Merlin S, Federer F, Angelucci A. 2021. A direct interareal feedback-to-feedforward
circuit in primate visual cortex. Nat Commun [Internet] 12:4911. Available from:
http://dx.doi.org/10.1038/s41467-021-24928-6

Soronow ALR, Jacobs MW, Dickson RG, Kim EJ. 2025. Bell Jar: A semiautomated registration and cell
counting tool for mouse neurohistology analysis. eNeuro [Internet] 12. Available from:
http://dx.doi.org/10.1523/ENEURO.0036-23.2025

Experiment	Mouse ID	Sex	Strain	From	Projection	Mouse age (P) at virus #1/#2 injection	Virus #1a (oG) and #1b (TVA)	Virus #1a (oG) titer	Virus #1b (TVA) titer	Virus #2 AAVretro	Virus #2 titer	Mouse age (P) at virus #3 injection	Virus #3 (rabies)	Virus #3 titer	Total virus/tracer incubation time (day)
FF/FB identity: Triple AAV	414	F	C57	VISl	ACA, VISp, VISpm	89	50 nl of scAAVretro-hSyn-H2B-eGFP into ACA, 50 nl of scAAVretro-hSyn-H2B-mCherry into VISp, 50 nl of AAVretro-BFP into VISpm	NA	NA	NA	NA	NA	NA	NA	21
FF/FB identity: Triple AAV	415	F	C57	VISl	ACA, VISp, VISpm	89	50 nl of scAAVretro-hSyn-H2B-eGFP into VISp, 50 nl of scAAVretro-hSyn-H2B-mCherry into VISpm, 50 nl of AAVretro-BFP into ACA	NA	NA	NA	NA	NA	NA	NA	21
FF/FB Identity: Double CTB	124	F	Tlx3-Cre	VISl	VISpm, VISp	104	50 nl of CTB-555 in VISpm, 50 nl of CTB-647 in VISp	NA	NA	NA	NA	NA	NA	NA	3
FF/FB Identity: Triple CTB	353	M	SepW1-Cre	VISl	ACA, VISp, VISpm	72	50 nl of CTB-647 in VISpm, 50 nl of CTB-555 in VISp, 50 nl of CTB-488 in ACA	NA	NA	NA	NA	NA	NA	NA	4
FF/FB Identity: Triple CTB	354	M	SepW1-Cre	VISl	ACA, VISp, VISpm	72	50 nl of CTB-555 in VISpm, 50 nl of CTB-488 in VISp, 50 nl of CTB-647 in ACA	NA	NA	NA	NA	NA	NA	NA	4
FF/FB Identity: Triple CTB	JR0845	M	C57	VISl	ACA, VISp, VISpm	64	50 nl of CTB-555 in VISpm, 50 nl of CTB-488 in VISp, 50 nl of CTB-647 in ACA	NA	NA	NA	NA	NA	NA	NA	6
FF/FB Identity: Triple CTB	JR0846	M	C57	VISl	ACA, VISp, VISpm	61	50 nl of CTB-647 in VISpm, 50 nl of CTB-555 in VISp, 50 nl of CTB-488 in ACA	NA	NA	NA	NA	NA	NA	NA	6
FF/FB Identity: Triple CTB	JR0847	M	C57	VISl	ACA, VISp, VISpm	61	50 nl of CTB-488 in VISpm, 50 nl of CTB-647 in VISp, 50 nl of CTB-555 in ACA	NA	NA	NA	NA	NA	NA	NA	6
FF/FB Identity: Triple CTB	JR0848	M	C57	VISl	ACA, VISp, VISpm	61	50 nl of CTB-555 in VISpm, 50 nl of CTB-488 in VISp, 50 nl of CTB-647 in ACA	NA	NA	NA	NA	NA	NA	NA	6
FF/FB Identity: Triple CTB	JR0849	M	C57	VISl	ACA, VISp, VISpm	61	50 nl of CTB-647 in VISpm, 50 nl of CTB-555 in VISp, 50 nl of CTB-488 in ACA	NA	NA	NA	NA	NA	NA	NA	6
FF/FB Identity: Triple CTB	JR0850	M	C57	VISl	ACA, VISp, VISpm	61	50 nl of CTB-488 in VISpm, 50 nl of CTB-647 in VISp, 50 nl of CTB-555 in ACA	NA	NA	NA	NA	NA	NA	NA	6
Control: No G-protein test	392	M	Tlx3-Cre	VISl	VISp	91	50 nl mixture of PBS and AAV8-TRE-DIO-eGFP-T2A-TVA into VISl	NA	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	105	100 nl of EnvA+RVdG-mCherry into VISl	1.43E+10	21
Control: No G-protein test	406	M	Tlx3-Cre	VISl	VISp	139	50 nl mixture of PBS and AAV8-TRE-DIO-eGFP-T2A-TVA into VISl	NA	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	153	100 nl of EnvA+RVdG-mCherry into VISl	1.43E+10	21
Control: No G-protein test	405	M	Tlx3-Cre	VISl	VISp	139	50 nl mixture of PBS and AAV8-TRE-DIO-eGFP-T2A-TVA into VISl	NA	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	153	100 nl of EnvA+RVdG-mCherry into VISl	1.43E+10	21
Control: Colocalization AAV retro with CTB	386	F	Tlx3-Cre	VISl	VISp	126	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl. 50 nl of CTB-647 in VISpm, 50 nl of CTB-555 in VISp	7.44E+13	NA	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	126	NA	NA	14
Control: Colocalization AAV retro with CTB	403	F	SepW1-Cre	VISl	VISp	131	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl. 50 nl of CTB-647 in VISp, 50 nl of CTB-555 in VISpm	7.44E+13	NA	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	NA	NA	NA	14
Control: Colocalization AAV retro with CTB	343	F	Tlx3-Cre	VISl	VISpm	149	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl. 10 nl of CTB-647 into VISpm	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	163	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Control: Colocalization AAV retro with CTB	344	F	Tlx3-Cre	VISl	VISpm	140	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl. 10 nl of CTB-647 into VISpm	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	154	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Control: No TTA test	94	M	Tlx3-Cre	VISl	NA	337	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	NA	NA	NA	NA	NA	NA	21
Control: No TTA test	425	M	SepW1-Cre	VISl	NA	74	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	7.44E+13	NA	NA	84	100 nl of EnvA+RVdG-mCherry into VISl	7.11E+09	21
Control: No TTA test	427	F	SepW1-Cre	VISl	NA	63	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA	1.41E+14	7.00E+12	NA	NA	77	100 nl of EnvA+RVdG-mCherry into VISl	7.11E+09	21
Control: No TTA test	428	F	SepW1-Cre	VISl/VISp	NA	63	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA	1.41E+14	7.00E+12	NA	NA	77	100 nl of EnvA+RVdG-mCherry into VISl	7.11E+09	21
Control: No TTA test	429	F	Tlx3-Cre	VISl	NA	51	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	7.44E+13	NA	NA	65	100 nl of EnvA+RVdG-mCherry into VISl	7.11E+09	21
Control: No TTA test	432	F	Tlx3-Cre	VISl	NA	49	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	7.44E+13	NA	NA	63	100 nl of EnvA+RVdG-mCherry into VISl	7.11E+09	21
Rabies: L2/3 VISl-VISpm	326	M	SepW1-Cre	VISl	VISpm	73	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	87	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L2/3 VISl-VISpm	327	M	SepW1-Cre	VISl	VISpm	73	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	87	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L2/3 VISl-VISpm	328	M	SepW1-Cre	VISl	VISpm	73	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	87	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L2/3 VISl-VISp	329	M	SepW1-Cre	VISl	VISp	68	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	82	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L2/3 VISl-VISp	161	F	SepW1-Cre	VISl	VISp	67	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA	1.41E+14	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	81	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L2/3 VISl-VISp	60	F	Tlx3-Cre	VISl	VISp	65	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	NA	100 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	79	100 nl of EnvA+RVdG-mCherry into VISl	1.78E+09	21
Rabies: L5 VISl-VISpm	67	F	Tlx3-Cre	VISl	VISpm	69	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	NA	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	83	100 nl of EnvA+RVdG-mCherry into VISl	1.07E+09	21
Rabies: L5 VISl-VISpm	143	M	Tlx3-Cre	VISl	VISpm	65	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	1.41E+14	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	79	100 nl of EnvA+RVdG-mCherry into VISl	1.07E+09	21
Rabies: L5 VISl-VISpm	118	F	Tlx3-Cre	VISl	VISpm	106	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	1.41E+14	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	120	100 nl of EnvA+RVdG-mCherry into VISl	1.07E+09	21
Rabies: L5 VISl-ACA	325	M	Tlx3-Cre	VISl	ACA	92	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	106	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L5 VISl-ACA	356	F	Tlx3-Cre	VISl	ACA	74	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	2.36E+13	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	89	100 nl of EnvA+RVdG-mCherry into VISl	7.71E+08	21
Rabies: L5 VISl-ACA	396	F	Tlx3-Cre	VISl	ACA	125	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	1.41E+14	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	139	100 nl of EnvA+RVdG-mCherry into VISl	1.43E+10	21
Rabies: L5 VISl-VISp	122	F	Tlx3-Cre	VISl	VISp	97	50 nl mixture of AAV8-TRE-DIO-oG and AAV8-TRE-DIO-GFP-T2A-TVA into VISl	1.41E+14	7.00E+12	50 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	111	100 nl of EnvA+RVdG-mCherry into VISl	1.07E+09	21
Rabies: L5 VISl-VISp	63	M	Tlx3-Cre	VISl	VISp	70	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	NA	200 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	84	100 nl of EnvA+RVdG-mCherry into VISl	1.78E+09	21
Rabies: L5 VISl-VISp	64	F	Tlx3-Cre	VISl	VISp	71	50 nl of AAV8-TRE-DIO-TVA66TeGFP-P2A-oG into VISl	7.44E+13	NA	100 nl of AAVretro-nEF-lox66/71-tTA	4.86E+13	85	100 nl of EnvA+RVdG-mCherry into VISl	1.07E+09	21

Raw counts in each brain area | Proportion of ipsilateral cortical inputs | nts in each thala / Proportion of thalamic inputs

Experiment	Mouse ID	VISp	AUDp	SSp-n	AUDv	SSp-ll	AUDd	SSp-un	SSp-ul	SSp-m	SSp-bfd	SSs	VISrl	RSPd	MOp	VISpl	RSPagl	RSPv	VISal	VISpm	SSp-tr	ORBI	AUDpo	VISpor	ACAv	VISam	VISa	ECT	MOs	PERI	TEa	Alp	GU	VISC	ACAd	ENTl	ENTm	LGd	LD	LP	VISp	AUDp	SSp-n	AUDv	SSp-ll	AUDd	SSp-un	SSp-ul	SSp-m	SSp-bfd	SSs	VISrl	RSPd	MOp	VISpl	RSPagl	RSPv	VISal	VISpm	SSp-tr	ORBI	AUDpo	VISpor	ACAv	VISam	VISa	ECT	MOs	PERI	TEa	Alp	GU	VISC	ACAd	ENTl	ENTm	LGd	LD	LP
L2/3 VISl-VISpm	326	660	129	1	32	12	151	16	13	0	144	103	69	16	11	3	14	5	40	161	102	3	78	31	2	23	183	2	17	0	18	0	2	4	15	18	3	4	1	41	31.75%	6.20%	0.05%	1.54%	0.58%	7.26%	0.77%	0.63%	0.00%	6.93%	4.95%	3.32%	0.77%	0.53%	0.14%	0.67%	0.24%	1.92%	7.74%	4.91%	0.14%	3.75%	1.49%	0.10%	1.11%	8.80%	0.08%	0.82%	0.00%	0.87%	0.00%	0.10%	0.19%	0.72%	0.87%	0.14%	8.70%	2.17%	89.13%
L2/3 VISl-VISpm	327	1038	184	1	21	1	55	4	3	2	98	26	140	50	2	14	30	7	72	128	7	1	64	61	0	42	55	4	17	0	77	2	0	1	7	5	15	9	1	73	46.55%	8.25%	0.04%	0.94%	0.04%	2.47%	0.18%	0.13%	0.09%	4.39%	1.17%	6.28%	2.24%	0.09%	0.63%	1.35%	0.31%	3.23%	5.74%	0.31%	0.04%	2.87%	2.74%	0.00%	1.88%	2.47%	0.14%	0.76%	0.00%	3.45%	0.09%	0.00%	0.04%	0.31%	0.22%	0.67%	10.84%	1.20%	87.95%
L2/3 VISl-VISpm	328	193	15	0	0	0	13	0	0	0	17	6	46	1	0	8	5	5	33	32	1	0	6	3	1	15	4	1	9	2	19	1	0	0	2	1	0	1	0	8	44.57%	3.46%	0.00%	0.00%	0.00%	3.00%	0.00%	0.00%	0.00%	3.93%	1.39%	10.62%	0.23%	0.00%	1.85%	1.15%	7.62%	7.39%	0.23%	0.00%	1.39%	0.69%	0.23%	3.46%	0.92%	0.14%	2.08%	0.46%	4.39%	0.23%	0.00%	0.00%	0.46%	0.23%	0.00%		11.11%	0.00%	88.89%
L2/3 VISl-VISp	60	94	23	0	3	0	14	0	0	0	14	2	14	1	0	1	2	0	11	11	1	0	15	4	0	8	12	1	0	0	10	0	0	0	1	6	5	1	0	5	37.30%	9.13%	0.00%	1.19%	0.00%	5.56%	0.00%	0.00%	0.00%	5.56%	0.79%	5.56%	0.40%	0.00%	0.40%	0.79%	0.00%	4.37%	4.37%	0.40%	0.00%	5.95%	1.59%	0.00%	3.17%	4.76%	0.19%	0.00%	0.00%	3.97%	0.00%	0.00%	0.00%	0.40%	2.38%	1.98%	16.67%	0.00%	83.33%
L2/3 VISl-VISp	161	403	54	0	18	0	27	0	1	0	30	11	32	31	0	22	26	14	36	32	2	0	16	27	0	12	15	8	0	1	64	3	0	12	0	22	3	30	2	58	44.09%	5.91%	0.00%	1.97%	0.00%	2.95%	0.00%	0.11%	0.00%	3.28%	1.20%	3.50%	3.39%	0.00%	2.41%	2.84%	1.53%	3.94%	3.50%	0.22%	0.00%	1.75%	2.95%	0.00%	1.31%	1.64%	0.43%	0.00%	0.11%	7.00%	0.33%	0.00%	1.31%	0.00%	2.41%	0.33%	33.33%	2.22%	64.44%
L2/3 VISl-VISp	329	1191	238	0	41	5	161	0	7	0	256	74	307	93	9	31	58	49	19	175	58	9	60	18	2	264	299	3	49	1	36	2	1	1	49	5	11	17	12	137	33.28%	6.65%	0.00%	1.15%	0.14%	4.50%	0.00%	0.20%	0.00%	7.15%	2.07%	8.58%	2.60%	0.25%	0.87%	1.62%	1.37%	0.53%	4.89%	1.62%	0.25%	1.68%	0.50%	0.06%	7.38%	8.35%	0.07%	1.37%	0.03%	1.01%	0.06%	0.03%	0.03%	1.37%	0.14%	0.31%	10.24%	7.23%	82.53%
L5 VISl-VISpm	67	76	0	1	0	1	0	0	0	0	0	5	7	10	0	0	5	5	7	2	0	0	0	2	0	2	0	1	1	0	1	2	0	3	0	10	2	3	0	6	53.52%	0.00%	0.70%	0.00%	0.70%	0.00%	0.00%	0.00%	0.00%	0.00%	3.52%	4.93%	7.04%	0.00%	0.00%	3.52%	3.52%	4.93%	1.41%	0.00%	0.00%	0.00%	1.41%	0.00%	1.41%	0.00%	0.70%	0.70%	0.00%	0.70%	1.41%	0.00%	2.11%	0.00%	7.04%	1.41%	33.33%	0.00%	66.67%
L5 VISl-VISpm	118	290	7	0	2	1	14	1	2	0	18	0	27	29	0	3	45	10	18	21	2	0	3	60	0	27	4	0	0	29	2	0	0	0	0	0	1	23	3	59	47.08%	1.14%	0.00%	0.32%	0.16%	2.27%	0.16%	0.32%	0.00%	2.92%	0.00%	4.38%	4.71%	0.00%	0.49%	7.31%	1.62%	2.92%	3.41%	0.32%	0.00%	0.49%	9.74%	0.00%	4.38%	0.65%	0.00%	0.00%	4.71%	0.32%	0.00%	0.00%	0.00%	0.00%	0.00%	0.16%	27.06%	3.53%	69.41%
L5 VISl-VISpm	143	601	12	0	10	0	10	3	1	0	28	22	42	60	0	16	54	63	61	25	7	0	10	44	0	24	55	0	0	86	1	0	3	0	4	4		46	3	77	48.23%	0.96%	0.00%	0.80%	0.00%	0.80%	0.24%	0.08%	0.00%	2.25%	1.77%	3.37%	4.82%	0.00%	1.28%	4.33%	5.06%	4.90%	2.01%	0.56%	0.00%	0.80%	3.53%	0.00%	1.93%	4.41%	0.00%	0.00%	6.90%	0.08%	0.00%	0.24%	0.00%	0.32%	0.32%		36.51%	2.38%	61.11%
L5 VISl-VISp	63	226	4	7	8	4	3	0	0	0	13	9	22	20	0	4	10	6	30	14	5	0	1	9	2	6	2	0	0	0	2	1	2	4	0			5	1	21	54.59%	0.97%	1.69%	1.93%	0.97%	0.72%	0.00%	0.00%	0.00%	3.14%	2.17%	5.31%	4.83%	0.00%	0.97%	2.42%	1.45%	7.25%	3.38%	1.21%	0.00%	0.24%	2.17%	0.48%	1.45%	0.48%	0.00%	0.00%	0.00%	0.48%	0.24%	0.48%	0.97%	0.00%			18.52%	3.70%	77.78%
L5 VISl-VISp	64	646	32	0	10	0	2	0	0	0	17	4	40	21	0	22	5	10	30	23	3	0	57	29	0	15	7	3	0	1	63	0	0	0	2	2		17	3	101	62.06%	3.07%	0.00%	0.96%	0.00%	0.19%	0.00%	0.00%	0.00%	1.63%	0.38%	3.84%	2.02%	0.00%	2.11%	0.48%	0.96%	2.88%	2.21%	0.29%	0.00%	5.48%	2.79%	0.00%	1.44%	0.67%	0.13%	0.00%	0.10%	6.05%	0.00%	0.00%	0.00%	0.19%	0.19%		14.05%	2.48%	83.47%
L5 VISl-VISp	122	308	12	1	2	3	2	0	0	0	9	2	34	7	0	32	7	6	22	15	1	0	10	39	0	1	0	0	0	10	0	0	1	3	0			13	0	23	58.44%	2.28%	0.19%	0.38%	0.57%	0.38%	0.00%	0.00%	0.00%	1.71%	0.38%	6.45%	1.33%	0.00%	6.07%	1.33%	1.14%	4.17%	2.85%	0.19%	0.00%	0.57%	7.40%	0.00%	0.19%	0.00%	0.00%	0.00%	1.90%	0.00%	0.00%	0.00%	0.19%	0.57%			36.11%	0.00%	63.89%
L5 VISl-ACA	325	320	20	0	0	0	14	0	0	0	24	16	42	34	0	4	41	3	12	25	4	0	2	14	0	12	14	0	0	0	11	0	0	0	3	2		NA	NA	NA	51.86%	3.24%	0.00%	0.00%	0.00%	2.27%	0.00%	0.00%	0.00%	3.89%	2.59%	6.81%	5.51%	0.00%	0.65%	6.65%	0.49%	1.94%	4.05%	0.65%	0.00%	0.32%	2.27%	0.00%	1.94%	2.27%	0.00%	0.00%	0.00%	1.78%	0.00%	0.00%	0.00%	0.49%	0.32%		NA	NA	NA
L5 VISl-ACA	356	103	7	0	3	0	1	1	0	0	5	1	9	2	0	0	1	0	41	2	0	0	4	2	0	0	0	1	0	19	0	0	1	0	1			NA	NA	NA	50.49%	3.43%	0.00%	1.47%	0.00%	0.49%	0.49%	0.00%	0.00%	2.45%	0.49%	4.41%	0.98%	0.00%	0.49%	0.00%	20.10%	0.98%	0.00%	0.00%	1.96%	0.98%	0.00%	0.00%	0.00%	0.49%	0.00%	9.31%	0.00%	0.00%	0.49%	0.00%	0.00%	0.49%			NA	NA	NA
L5 VISl-ACA	396	164	35	0	67	0	18	0	0	0	12	23	21	11	0	0	4	9	0	6	0	0	14	17	0	16	15	38	0	14	79	0	0	0	12	5		NA	NA	NA	30.26%	6.46%	0.00%	12.36%	0.00%	3.32%	0.00%	0.00%	0.00%	2.21%	4.24%	3.87%	2.03%	0.00%	0.00%	0.74%	1.66%	0.00%	1.11%	0.00%	0.00%	2.58%	3.14%	0.00%	2.95%	2.77%	7.01%	0.00%	2.58%	14.58%	0.00%	0.00%	0.00%	2.21%	0.92%		NA	NA	NA

Projection- and layer- specific long-range input tracing
SepW1-Cre (L2/3) or Tlx3-Cre (L5 CCPNs)

Axonal target-
dependent
VISpm or ACA input

Relaying information-
dependent
context inputs
(VISpm, HCtx, LP)

Inputs are common

VISl

VS.

VS.

VISp input

sensory inputs
(VISp, LGd)

Common Inputs

Model 1

Model 2

Model 3

Figure 1

Figure 2

Figure 3

Figure 4

A L2/3: SepW1-Cre VISl→VISpm (FF)   L2/3: SepW1-Cre VISl→VISp (FB)

LGd   LGd   LP   LP

DAPI
RV*dG*-mCherry

100μm

B