A periplasmic protein complex mediates cell envelope integrity and intrinsic multidrug
resistance in *Mycobacterium tuberculosis*

Thaís Klevorn[1,2], Christopher Brown[3*], Christine D. Hardy[4*], Bonnie J. Cuthbert[4*], Amanda
Spencer[5], Adrián Jinich[5], Luming Chan[1], Jessica Mendoza[4], Rodger de Miranda[4], Heather Kim[1],
Dirk Schnappinger[1], Kyu Rhee[2,3], Celia Goulding[4,6], Sabine Ehrt[1,2+]

[1]Department of Microbiology and Immunology, Weill Cornell Medical College, New York, NY, USA

[2]Immunology and Microbial Pathogenesis Graduate Program, Weill Cornell Graduate School of
Medical Sciences, Cornell University, New York, New York, USA

[3]Department of Medicine, Weill Cornell Medical College, New York, New York, USA

[4]Department of Molecular Biology & Biochemistry, University of California Irvine, Irvine, California,
USA

[5]Department of Chemistry & Biochemistry, Skaggs School of Pharmacy & Pharmaceutical
Sciences, University of California San Diego, San Diego, California, USA

[6]Department of Pharmaceutical Sciences, University of California Irvine, Irvine, California, USA

*These authors contributed equally.

[+]Corresponding author.

Abstract

The intrinsic drug resistance of *Mycobacterium tuberculosis* (Mtb) is a major barrier to effective
tuberculosis (TB) treatment and is largely attributed to its complex, impermeable cell envelope.
We identified a periplasmic protein complex comprising FecB and Rv3035 that is essential for
maintaining cell envelope integrity and mediating intrinsic multidrug resistance in Mtb. FecB
interacts directly with Rv3035, forming a stable heterodimer that also associates with the cell
envelope biosynthesis proteins AftB and Rv0227c. We report the structures of Rv3035 alone and
in complex with FecB, and identify critical residues required for complex formation and function.
Co-essentiality analysis and genetic interaction experiments further supported a functional link
between FecB, Rv3035 and AftB, the enzyme responsible for the last step in the biosynthesis of
the cell envelope components arabinogalactan and lipoarabinomannan. Mtb mutants of *fecB*,
rv3035 and *aftB* accumulated trehalose monomycolates in the outer membrane, consistent with
a perturbation of membrane assembly and/or maintenance. Importantly, we show that FecB is
required for Mtb virulence in mice, underscoring its physiological relevance. Altogether, our
findings uncover a previously uncharacterized periplasmic protein complex that supports cell
envelope integrity and intrinsic drug resistance in Mtb, highlighting FecB and Rv3035 as promising
targets for therapeutic intervention.

Introduction

Tuberculosis (TB) remains the world's leading cause of death from a single infectious agent and is responsible for over 1 million deaths anually[1]. Unlike most bacterial infections, successful TB treatment requires multiple antibiotics taken for several months, even in drug-susceptible cases. This prolonged therapy is in part due to the intrinsic drug resistance of *Mycobacterium tuberculosis* (Mtb), the causative agent of TB[2]. A key contributor to this resistance is Mtb's complex, multi-layered cell envelope, which acts as a protective barrier against antibiotics and host immune defenses[3]. Despite its central role in Mtb's pathogenicity and drug resistance, the molecular mechanisms governing the structure, assembly, and function of the mycobacterial cell envelope remain incompletely understood[4].

To identify novel targets that could potentiate existing TB therapies and shorten treatment duration, we previously performed a genome-wide chemical-genetic screen using transposon mutagenesis and sequencing[5]. This identified *fecB* (Rv3044) as a gene whose disruption sensitized Mtb to multiple antibiotics that have various mechanisms of action. These included rifampicin and ethambutol, which are part of the standard treatment regimen for drug susceptible TB, and pretomanid which is part of the BPaL (bedaquiline, pretomanid, linezolid) regimen used to treat drug-resistant TB[5]. Functional studies of a Mtb *fecB* knockout strain (Δ*fecB*) revealed that *fecB* is essential for maintaining the impermeability of the Mtb cell envelope [5].

In Mtb, FecB is annotated as a periplasmic substrate-binding protein homologous to FecB from *Escherichia coli*, which mediates ferric citrate transport[6]. However, Mtb lacks several canonical components of the ferric citrate transport system. Instead, Mtb FecB binds heme and the siderophore carboxymycobactin and interacts with components of the mycobacterial siderophore transport machinery[7]. Despite these biochemical insights, the physiological role of FecB in Mtb is not fully defined. Here, we investigated how FecB contributes to Mtb's intrinsic antibiotic resistance and uncovered a critical functional interaction with Rv3035, a previously uncharacterized periplasmic protein. Rv3035 not only interacts with FecB, but also with AftB (Rv3805c), likely bridging these two proteins to form a larger complex that is important for cell envelope biogenesis.

Given the urgent need for shorter and more effective TB treatments, elucidating mechanisms that underlie Mtb's intrinsic drug resistance is of high therapeutic relevance[2,3]. By characterizing the function of FecB and Rv3035 in Mtb, this study advances our understanding of periplasmic proteins in mycobacteria and highlights these proteins as promising targets for combination therapies aimed at enhancing the efficacy of existing anti-TB drugs.

Results

Mtb Δ*fecB* displays a transcriptional iron deprivation signature and accumulates siderophores

Transcriptome analysis can provide functional insights by identifying pathways that are differentially regulated in a mutant strain compared to a wild type (WT) strain. These changes

may reflect the mutant strain's need to adapt in the absence of the gene of interest[8]. To gain
insight into the function of FecB, we performed transcriptomic analysis by RNA sequencing of Mtb
WT, Δ*fecB* and a complemented strain (Δ*fecB::fecB,* containing a chromosomally integrated
plasmid expressing *fecB* from the *hsp60* promoter). We decided to compare the transcriptional
profile of the strains in regular growth conditions with detergent (0.05% Tween80) and without
detergent, because the presence of Tween80, although helpful to prevent excessive clumping of
mycobacterial cultures, also strips the mycobacterial capsule and some surface exposed lipids[9],
making it a less physiologically relevant condition.

Compared to WT and the complemented strain, Δ*fecB* significantly upregulated the *iniBAC*
operon, which is known to be induced in response to cell wall stress[10], both in the presence and
absence of detergent in the medium (Fig. 1A and data S1). Furthermore, in detergent-free
conditions, 61% of the upregulated (22 of 36) and 13% of downregulated (3 of 23) differentially
expressed genes (DEGs) were related to iron utilization. The upregulated DEGs included all 14
siderophore biosynthesis genes (*mbtA-mbtN*), the genes that encode the IrtA/B transporter for
siderophore uptake (*irtA*, *irtB*) and the ESX-3 secretion system substrates (*esxG*, *esxH*), which
are required for mycobactin-mediated iron acquisition. The downregulated DEGs included *bfrB*,
encoding the iron storage protein bacterioferritin (Fig. 1, A and B). When grown in detergent-free
media, Δ*fecB* thus displays a transcriptional iron deprivation signature even though the media
contains a high concentration of iron. This iron deprivation signature is conditional, and it was not
present when Tween80 was included in the medium (Fig. 1A). One potential explanation for this
difference in transcriptional response is that Tween80's stripping of the capsule and some outer
membrane lipids might facilitate uptake of iron in Δ*fecB* during growth in regular media.

Further supporting the transcriptomics data, the iron starvation signature in Δ*fecB* in detergent-
free medium was accompanied by an accumulation of cell-associated siderophores as identified
by mass-spectrometry analysis of lipid extracts (Fig. 1C). Taken together, these data suggest that
Δ*fecB* may be less efficient in acquiring iron in detergent-free media due to interrupted
siderophore export through the periplasm, mycobacterial outer membrane and/or capsule.

FecB is non-critical for mycobactin-mediated iron acquisition in Mtb

To further probe FecB's potential role in iron acquisition, we analyzed the growth of Δ*fecB*, WT
and Δ*fecB*::*fecB* in iron-limiting conditions (Fig. 1D). We included a siderophore mutant,
MtbΔ*mbtK*, to ensure that the growth medium was iron limiting. After starving the strains for iron
to deplete internal iron stores and then growing them in either high or low iron conditions (iron-
chelated medium with 160 μM or 1 μM ammonium ferric citrate, respectively), we observed only
a minor growth defect of Δ*fecB* compared to the control strains, both in the presence (Fig. 1D) or
absence of detergent (fig. S1A). Upon passaging the strains growing in low iron into both high
and low iron conditions, the growth defect was still subtle and much weaker than the strong growth
defect of Δ*mbtK* (Fig. 1D).

FecB can bind to ferric- and apo-carboxymycobactin with high affinity *in vitro*, and this binding is
disrupted by mutating key residues E339, R240 and Y242[7]. To test whether binding to
siderophores is required for complementing Δ*fecB*'s iron-related phenotypes, we cloned and

expressed FecB variants harboring the mutations E339S together with R240S or Y242S in $\Delta fecB$.
The FecB variants with significantly reduced binding affinity to ferric-carboxymycobactin
(E339S_R240S and E339S_Y242S) fully reversed the iron deprivation transcriptional signature
(examined by representative gene expression of *mbtB*, *ppe37* and *irtA*) in detergent-free media
(Fig. 1E), and the subtle growth defect in low iron (fig. S1B). This complementation was
particularly remarkable given that the variant proteins were not clearly detectable in whole cell
lysates via western blot (fig. S1C). However, we note that the FecB E339S_R240S and
E339S_Y242S double mutants bind apo-carboxymycobactin with similar (or slightly higher)
affinity than WT FecB (fig. S1D). Thus, we cannot rule out that binding apo-siderophores is an
important function of FecB.

Taken together, these results suggest that FecB plays a role in iron acquisition in Mtb, however,
the subtle growth defect of $\Delta fecB$ in iron-limiting conditions points to a condition-dependent non-
critical function that is likely compensated for by other gene products required for iron acquisition.

FecB's role in intrinsic multidrug resistance is distinct from its activity in iron homeostasis

FecB is critical for Mtb's cell envelope impermeability and resistance to multiple antibiotics[5]. To
test whether FecB's activity in iron homeostasis is related to its activity in mediating cell envelope
integrity and multidrug resistance in Mtb, we tested whether the FecB variants with reduced
binding to ferric-carboxymycobactin could rescue the antibiotic hypersusceptibility of $\Delta fecB$.
Vancomycin and rifampicin were chosen for minimal inhibitory concentration (MIC) assays
because $\Delta fecB$ is significantly more susceptible to these drugs compared to WT[5].
Complementation of $\Delta fecB$ with the FecB double mutants (E339S_R240S and E339S_Y242S)
fully rescued $\Delta fecB$'s hypersusceptibility to vancomycin and rifampicin (Fig. 2A) indicating that
ferric-carboxymycobactin binding is not required for FecB's function in mediating antibiotic
resistance.

To explore whether siderophore accumulation in $\Delta fecB$ (Fig. 1C) leads to antibiotic
hypersusceptibility, we interrupted siderophore biosynthesis via CRISPR interference (CRISPRi)
knock down (KD) of the mycobactin biosynthesis gene *mbtI* in $\Delta fecB$. If siderophore accumulation
is directly related to increased antibiotic susceptibility in $\Delta fecB$, reducing siderophore production
by knocking down *mbtI* should relieve this phenotype. We validated the KD of *mbtI* in WT and
$\Delta fecB$ backgrounds by growing the strains in iron-limiting conditions, in the presence or absence
of anhydrotetracycline (ATc) to activate CRISPRi (fig. S2A). In the presence of ATc, the *mbtI* KD
strains failed to grow with ammonium ferric citrate concentrations lower than 1 μM, similarly to
$\Delta mbtK$, indicating that siderophore production is significantly impaired through silencing of *mbtI*.
We next measured the MICs of vancomycin and rifampicin against the *mbtI* KD strain, in the
presence or absence of ATc (Fig. 2B). Silencing *mbtI* with ATc did not alter the MICs, indicating
that siderophore accumulation is likely not the cause of antibiotic hypersusceptibility in $\Delta fecB$.

To further test whether siderophores are involved in the antibiotic hypersusceptibility of $\Delta fecB$, we
tested whether induction of siderophore biosynthesis would enhance $\Delta fecB$'s susceptibility to
vancomycin and rifampicin. We performed MIC assays in low iron medium, where siderophore
synthesis is induced, and in regular medium, where siderophore synthesis is suppressed (Fig. 2C

and fig. S2B). Additionally, since an iron-starvation transcriptional signature, which includes the
overexpression of siderophore biosynthesis genes, was observed in Δ*fecB* only in the absence of
detergent (Fig. 1A), we also tested whether the mutant's antibiotic hypersusceptibility would
increase in detergent-free medium compared to medium containing 0.05% Tween80 (fig. S2C).
Contrary to these hypotheses, the differences of the vancomycin and rifampicin MICs against
Δ*fecB* compared to WT and Δ*fecB*::*fecB* decreased in low iron medium compared to regular iron
rich medium (Fig. 2C). Similarly, a reduced shift in MICs was observed in detergent-free conditions
(fig. S2C). These results do not support a direct involvement of siderophores in the antibiotic
hypersusceptibility phenotype of Δ*fecB.* The reduced differences in antibiotic MICs might be due
to slowed growth rates of the strains in low iron medium and increased clumping in detergent-free
medium, which hinder accurate optical density measurements. In summary, these experiments
dissociate FecB's role in iron homeostasis from that in antibiotic resistance.

FecB and Rv3035 interact in Mtb and likely function in the same cellular pathway

FecB is classified as a periplasmic substrate-binding protein that has been shown to interact with
siderophore-mediated iron transport proteins[7]. Since our data point to an alternative role for FecB
in mediating intrinsic drug resistance in Mtb, we set out to identify additional physical interactors
of FecB that could shed light on other pathways in which FecB might participate. Pull-down and
mass spectrometry analysis of FLAG-tagged FecB was performed and analyzed compared to the
pull-down of an WT lysate control to eliminate non-specific interactions.

The most highly enriched interactor in our dataset was the uncharacterized protein Rv3035 (Fig.
3A). Rv3035 is annotated as an essential hypothetical protein containing WD40 repeats that
together are predicted to form an eight-bladed β-propeller structure. These domains can serve as
scaffolds in protein-protein interactions and the formation of multiprotein complexes and play
important roles in prokaryotes and eukaryotes[11]. We confirmed binding of FecB to Rv3035 by co-
immunoprecipitation of the tagged proteins (fig. S3) and then set out to identify Rv3035's physical
interactors by using Rv3035 as the bait in a second pull-down experiment. Satisfyingly, FecB was
the most enriched protein in the pull-down of Rv3035-3xFLAG, compared to the WT lysate control
(Fig. 3B). In addition, AftB and Rv0227c, which are the other two most enriched interacting
partners of FecB, were also among the most enriched partners of Rv3035 (Fig. 3, A and B, and
data S2).

To further inform our proteomics findings, we investigated the genetic interactions of *fecB* via co-
essentiality/co-fitness analysis. This method allows the detection of gene pairs that share similar
fitness profiles across forward genetic screens, and thus, can help identify genes that encode
proteins with similar or related function[12]. Using a publicly available CRISPRi chemical-genetic
interaction dataset which profiled gene essentiality across 80 drug-treatment and CRISPRi pre-
depletion conditions[13], we performed co-essentiality analysis using generalized least squares
(GLS) regression of *fecB*, *rv3035*, and the genes encoding their most enriched common protein
interactors, *aftB* and *rv0227c*. While *rv0227c* did not share an essentiality profile with the others,
fecB, *rv3035* and *aftB* were all hits in each other's co-essential gene clusters (Fig. 3, C and D).

This analysis identified 13 genes co-essential with *fecB*, 19 with *rv3035* and 22 with *aftB*, among
which 4 were shared by all three (Fig. 3D and data S3).

FecB forms a complex with Rv3035 *in vitro*

To confirm the direct binding of Rv3035 and FecB *in vitro*, we cloned His-tagged Rv3035 from
Mycobacterium thermoresistible (Mth). Mth Rv3035 shares 67% sequence identity with Mtb
Rv3035 (fig. S4A) and was used because full-length or N-terminal truncation constructs of His-
tagged Mtb Rv3035 failed to yield soluble protein. Rv3035 is predicted to have an N-terminal
signal peptide by SignalP 6.0[14] and to be a lipoprotein, where the N-terminal cysteine residue is
predicted to form a thioether linkage to diacylglycerol[15,16]. Here, we have designated this N-
terminal cysteine as the first amino acid of the mature form of Rv3035 (fig. S4B). Our Mth Rv3035
protein construct begins at residue 2 of the predicted mature protein and was expressed with a
cleavable, N-terminal His-tag. Mth Rv3035-His was purified to homogeneity and the His-tag
cleaved, with size-exclusion chromatography (SEC) of the resulting protein suggesting that Mth
Rv3035 is monomeric (fig. S5A).

Mth Rv3035-His was also co-expressed with untagged Mtb FecB and found to co-elute by Ni-
affinity chromatography (fig. S5B). The Mth Rv3035 and Mtb FecB complex was stable during
SEC and displayed a molecular weight of 79.1 kDa, consistent with a heterodimer of Rv3035 and
FecB (fig. S5A). The affinity of Mth Rv3035 for Mtb FecB was quantified using isothermal titration
calorimetry (ITC). This analysis revealed tight 1:1 binding of Mth Rv3035 and Mtb FecB, with a
K_D of 27.5 ± 4.6 nM (Fig. 4A).

Structure determination of Rv3035 and the FecB-Rv3035 complex

We next solved the structure of Mth Rv3035 by X-ray crystallography to 1.9 Å (table S1). Mth
Rv3035 crystallized with two subunits in the asymmetric unit. In both subunits, there is clear
electron density for all but the N-terminal seven residues of Rv3035. The two subunits are virtually
identical with a root mean square deviation (rmsd) of 0.46 Å over 392 residues.

The structure of Rv3035 adopts a β-propeller fold, composed of eight 4-stranded β-sheets, where
each β-sheet is referred to as a blade (Fig. 4B and fig. S4A). At the N-terminus of the protein,
preceding the β-propeller fold, are ~40 residues that form an ordered loop region, which occupies
the bottom half of the central cavity formed by the β-propeller. A small portion of this N-terminal
region forms a β-strand (β0, residues 27-32) that forms a β-sheet with the interior β-strand (β26)
of blade H (Fig. 4B and fig. S4A). The top half of the cavity is occupied by a second ordered loop
region formed by residues 171-197 containing a small α-helix (fig. S4A), which appears to act as
a cap (α-cap) for the central pore (Fig. 4B).

Mth Rv3035 contains two disulfide bonds (Fig. 4B, fig. S4A, and fig. S6A). The first is formed
between C71 and C87 and connects the exterior β-strand (β5) of blade B to the loop region on
the α-cap side of the β-propeller, and the second is formed between C187 and C194 connecting
the α-cap to the ordered loop region following it, perhaps stabilizing the cap region. A structural

homology search of Mth Rv3035 utilizing the DALI server[17] (table S2) demonstrated that the
closest hits have extremely high structural homology with Rv3035 and the majority are accessory
proteins that assist in remodeling protein assemblies; for example, two of the top hits are BamB
from Gram-negative bacteria.

To determine the protein-protein interacting residues of the FecB-Rv3035 complex, we solved the
X-ray crystal structure of the FecB-Rv3035 complex to a resolution of 3.35 Å (table S1). There
are minimal structural differences between FecB alone[7] and FecB in complex with Rv3035 (fig.
S6, C and D). One notable difference is the presence of a disulfide bond between C56 and C104
(fig. S6D), which is absent in the previous FecB crystal structure. Likewise, there are little
differences between the structures of Rv3035 alone and in complex with FecB (fig. S6, E and F).

The FecB-Rv3035 complex interface is small and involves 729 Å^2 (5.6%) of the FecB surface,
and 692 Å^2 (4.3%) of Rv3035 (Fig. 4C). Although the interface is small, there are multiple
interactions between the two proteins including 19 hydrogen bonds and salt bridges that are $\leq$
3.5 Å and six interactions that are < 3.0 Å (Fig. 4D and table S3). The interface involves residues
from Rv3035 blade B (β4-β5) and residues from the loop that precedes blade B (between β1 and
β2) (Fig. 4D and fig. S4A). The interface also involves the backbone nitrogen atom from disulfide
bond forming C71 (Fig. 4D and fig. S6B). Strikingly, only nine Rv3035 residues are responsible
for the polar interactions at the interface, with six residues forming multiple contacts with FecB:
R43, K46, E77, R83, Q84, and R85, where R83 forms four polar contacts at the interface. FecB
residues participating in the polar contacts at the interface are localized to the second lobe of
FecB and are located in α7 (D223, H226), the loop between α8 and α9 (D258), the loop between
α10B and β11 (A291, D292), and α13 (S320, R323, D324)[7]. Similar to Rv3035, several FecB
residues are responsible for multiple contacts at the interface: H226, D292, S320, R323 and
D324; where D292 forms five and D324 forms four polar contacts at the interface. The FecB-
Rv3035 interface leaves the ligand binding site of FecB accessible to solvent, and thus still able
to accommodate a ligand (Fig. 4C).

Key FecB residues in binding Rv3035

To test the functional importance of FecB residues involved in binding Rv3035, we mutated FecB
D292 to alanine alone or H226, D292 and D324 to alanine together, and tested binding of these
proteins to Rv3035 by protein interaction pull-down assays and by ITC. After determining that the
mutants were properly folded by circular dichroism (fig. S7A), we utilized FecB-His and untagged
Rv3035 in a nickel-affinity pull-down assay (fig. S7B). While WT FecB-His copurified with Rv3035,
FecB D292A alone and the FecB triple mutant H226A_D292A_D324A failed to pull down Rv3035.
In addition, no binding to Rv3035 was observed for the single and triple FecB mutants by ITC
analysis (fig. S7C). Thus, the nanomolar affinity (K_D of ~30 nM) interaction between FecB and
Rv3035 can be completely disrupted by mutating one FecB residue, D292.

Rv3035 mediates intrinsic multidrug resistance in Mtb

Having demonstrated that FecB and Rv3035 interact, we next set out to probe the cellular function
of Rv3035. In TnSeq screens, *rv3035* mutants are usually annotated as having a "growth defect"[18]
and are often absent from TnSeq screen input libraries. That is likely the reason why *rv3035* was
not a hit in our TnSeq chemical-genetic interaction screen that identified *fecB* as important for
intrinsic antibiotic resistance in Mtb[5]. Because *fecB* and *rv3035* display similar growth defects by
CRISPRi[19], we reasoned that it would be possible to generate a *rv3035* knockout strain (Δ*rv3035*)
to test directly whether Rv3035 mediates intrinsic resistance to different antibiotics. Although
Δ*rv3035* colonies took longer to grow on agar plates than WT Mtb, the mutant strain exhibited
similar growth kinetics as WT and Δ*fecB* in liquid media (fig. S8A). Additionally, we generated a
Δ*fecB* Δ*rv3035* double knockout as well as the respective complemented mutants.

We performed MIC assays using these strains for several antibiotics having different mechanisms
of action to which Δ*fecB* was hypersusceptible (Fig. 5A). Isoniazid was included as a control
because its MIC is not affected by the absence of *fecB*. The single Δ*rv3035* and double
Δ*fecB*Δ*rv3035* mutants showed very similar antibiotic hypersusceptibility patterns as Δ*fecB*, with
reduced MICs compared to WT. As the basis for Δ*fecB*'s hypersusceptibility to antibiotics is likely
a more permeable cell envelope compared to WT Mtb[5], we also tested whether Δ*rv3035* and
Δ*fecB*Δ*rv3035* are more permeable to fluorescently labelled vancomycin and the reporter dye
calcein-AM. Both mutants phenocopied Δ*fecB* with increased uptake of BODIPY-vancomycin (Fig.
5B) and both were hyperpermeable to calcein-AM similarly to Δ*fecB* (Fig. 5C).

Δ*fecB* also displayed altered colony morphology and a growth defect when growing as single
colonies on agar plates, appearing smooth on high density spots and often failing to grow in lower
spot dilutions (Fig. 5D). This makes experiments that rely on the quantification of Δ*fecB* colony
forming unit (CFU) titers unreliable. Although we do not know the basis of this phenotype, it could
be related to cell envelope alterations. Δ*rv3035* and Δ*fecB*Δ*rv3035* displayed the same defect
(Fig. 5D). Taken together, these results demonstrate that Rv3035 is important for intrinsic
multidrug resistance, mediates cell envelope impermeability and is required for regular growth on
solid media. Importantly, deleting *fecB* and *rv3035* in the same strain did not result in enhanced
phenotypes (Fig. 5, A-D), a result that it is expected for genes functioning in the same pathway.

To test whether binding between FecB and Rv3035 is required for maintaining cell envelope
integrity, we complemented Δ*fecB* with an integrated plasmid expressing the FecB triple mutant
H226A_D292A_D324A which fails to bind Rv3035 *in vitro* (fig. S7C). First, we confirmed that the
FecB triple mutant is stably expressed *in vivo* (fig. S8B). Then, we tested whether Δ*fecB*
complemented with the triple mutant (Δ*fecB*::*fecB*-H226A_D292A_D324A) can rescue Δ*fecB*'s
hypersusceptibility to vancomycin and rifampicin. The MICs of these drugs were unchanged in
Δ*fecB*::*fecB*-H226A_D292A_D324A compared to Δ*fecB* (Fig. 5E). Similarly, the FecB triple mutant
failed to rescue Δ*fecB*'s growth defect in solid media (fig. S8C). These results indicate that FecB's
binding to Rv3035 is critical to mediate intrinsic drug resistance in Mtb.

FecB and Rv3035 interact with cell envelope biosynthesis proteins

Our pull-down data revealed that FecB and Rv3035 also interact with the cell envelope
biosynthesis proteins AftB and Rv0227c (Fig. 3, A and B), which was confirmed by co-

immunoprecipitation assays (fig. S9 and S10). Moreover, *fecB* and *rv3035* are co-essential with
aftB (Fig. 3C), which suggests they may perform similar functions and could be part of a protein
complex[12]. AftB catalyzes the last step in the biosynthesis of the arabinan domain of
arabinogalactan (AG) and lipoarabinomannan (LAM), adding the terminal arabinofuranose (Ara*f*)
residue to those molecules[20]. In AG, the β-(1 → 2)-linked terminal-Ara*f* residues provide mycolic
acid (MA) attachment sites which are critical for the formation of the mycolyl–arabinogalactan–
peptidoglycan (mAGP) complex[21,22]. Rv0227c (named polar growth factor A, or PgfA, in
Mycobacterium smegmatis) is an inner membrane protein that interacts with MmpL3, the
transporter for the cell envelope component trehalose monomycolate (TMM), and with a TMM
analog, suggesting that it participates in TMM transport across the plasma membrane[23].

In the periplasm, TMMs may donate a mycoloyl residue to AG to form the mAGP complex or to
another TMM molecule to form trehalose dimycolate (TDM or cord factor)[24]. While MAs attached
to AG (mAG) are the major components of the inner leaflet of the mycobacterial outer membrane
(MOM), TMMs, TDMs and free MAs are part of the outer leaflet of the MOM[3]. These lipids are
critical determinants of Mtb's cell envelope impermeability[3]. Since AftB and Rv0227c are involved
in the biosynthesis of mAG and the MOM, the interaction of FecB and Rv3035 with these proteins
suggests that they participate in the assembly or maintenance of mAG and/or the MOM.

We used AlphaFold 3[25] to predict the potential protein complexes formed by FecB, Rv3035, AftB
and/or Rv0227c, modeling different combinations of these proteins (fig. S11 and S12). The FecB
and Rv3035 complex showed a very high per-residue model confidence score (pIDDT, fig. S11),
and has the same interface and orientation observed in the experimental structure of FecB-
Rv3035, superimposing with a rmsd of 0.91 Å over 696 residues (Fig. 4C). The complexes
containing AftB (with or without Rv0227c) showed a higher confidence score when modeled with
Rv3035 than with FecB alone (fig. S12, B and C). These structural models crossed the interface
predicted template modeling (ipTM) score threshold for high-quality prediction of 0.8 when AftB
was modelled with FecB and Rv3035, with Rv3035 and Rv0227c, or when all 4 proteins were
modeled together (fig. S12). This suggests that Rv3035 bridges the interaction between FecB and
AftB, and between FecB and Rv0227c.

AftB is more vulnerable to knockdown in Δ*fecB* than in WT

Because *aftB* shares an essentiality profile with *fecB* and *rv3035* (Fig. 3C), we decided to further
explore the genetic interaction between *fecB* and *aftB*. We depleted *aftB* with a weak CRISPRi
single guide RNA in both WT and Δ*fecB*. While this weak guide caused a slight growth defect in
WT upon induction of the CRISPRi system, knockdown of *aftB* by the same weak guide in Δ*fecB*
completely abrogated growth (Fig. 6A). This indicates that *aftB* is more vulnerable to knockdown
in Δ*fecB* than in WT as a moderate depletion of *aftB* leads to a profound loss of fitness in the
absence of *fecB*.

FecB is required to maintain normal levels of MAs, TMMs and TDMs in Mtb

To test whether FecB is required for proper assembly of the MOM, we performed lipidomics
analysis of total extractable lipids from WT, Δ*fecB* and Δ*fecB*::*fecB* strains (Fig. 6, B and E, and
data S4). Contrary to expectations that mycolate levels would be reduced in Δ*fecB* due to their
role in mediating envelope impermeability, the mutant displayed a higher abundance of TMMs
and free MAs compared to WT and Δ*fecB*::*fecB*.

Since TMMs accumulate in the plasma membrane of TMM transport mutants[23,26–28], we next
assessed if lipid trafficking was disrupted in Δ*fecB*. We treated bacterial pellets with the anionic
surfactant dioctylsulfosuccinate sodium (AOT) dissolved in heptane to strip non-covalently bound
lipids off the MOM[29]. In *Corynebacterium glutamicum*, AOT in heptane forms reverse micelles
which solubilize and capture OM surface lipids but does not penetrate other layers of the cell
envelope[29]. We validated this approach by analyzing known lipids extracted from the supernatants
and pellets of AOT treated samples, corresponding to the outer leaflet of the MOM (labelled OM)
and all other extractable lipids (labelled inner membrane or IM), respectively (fig. S13, A-D, and
data S4). Δ*fecB* displayed a higher abundance of TMMs and free MAs in both IM and OM
fractions, compared to WT and Δ*fecB*::*fecB* (Fig. 6, C-D and F-G, and data S4). We used the OM
fraction to visualize and quantify TDMs via thin-layer chromatography (Fig. 6H and fig. S13E).
Although TDM levels are inversely correlated with TMMs in mutants with defective TMM
transport[23,26–28], Δ*fecB* also displayed increased TDM levels compared to controls.

To validate the lipidomics data with an alternative method, we used the solvatochromic trehalose
probe DMN-Tre to track MA incorporation into TMMs via flow cytometry[30]. Upon entering the cell,
DMN-Tre is mycolylated to form DMN-Tre monomycolates, which are subsequently inserted into
the MOM, where the hydrophobic environment activates the probe's fluorescence[30].
Corroborating the lipid analysis, Δ*fecB* incorporated more DMN-Tre in its cell envelope than WT,
shown as increased mean fluorescence intensity compared to WT and Δ*fecB*::*fecB* (Fig. 6I), a
phenotype also shared with Δ*rv3035* (fig S13F).

Together, these results suggest that TMM transport occurs normally in Δ*fecB*, and the
accumulation of TMMs, TDMs and free MAs in the mutant is caused either by increased mycolate
lipid biosynthesis or blockage of a downstream process that uses these species as precursors,
such as incorporation of mycolic acids into mAG. Our RNA-seq data did not show differences in
transcript levels for MA/TMM biosynthesis genes in Δ*fecB* (data S1), thus favoring the second
hypothesis.

The *aftB* KD strains also displayed increased incorporation of DMN-Tre upon CRISPRi activation
(Fig. 6J). Since *aftB* directly controls the addition of mycolylation sites on AG, it is probable that
the *aftB* KD strain accumulates mycolates that do not attach to the cell wall[20,31] and the increased
incorporation of DMN-Tre reflects the redirection of the mycolate pool towards the MOM.

FecB is essential for virulence of Mtb in mice

The mycobacterial cell envelope is a formidable barrier that protects Mtb not only from antibiotics
but also from host immune responses[3]. Thus, we hypothesized that *fecB* might also be required
for Mtb to establish a successful infection in mice. The unreliability of Δ*fecB* to grow as single

colonies on agar plates (Fig. 5D), however, impairs the quantification of the bacterial burden in
infected mice. To circumvent this problem, we implemented a genetic switch to activate *fecB*'s
expression in the presence of a tetracycline inducer (TetON)[32] and allow quantification of CFUs
from mice infected with Δ*fecB* harboring the TetON switch (FecB-TetON). The conditional
expression of *fecB* was validated via MIC assays with vancomycin and rifampicin, which showed
that addition of ATc rescued the antibiotic hypersusceptibility of FecB-TetON (Fig. 7A).

Mice were infected via the aerosol route with FecB-TetON and received either regular or
doxycycline (doxy) chow throughout the course of the experiment (Fig. 7, B and C). Lung and
spleen homogenates from both groups were cultured on agar containing ATc to activate *fecB*
expression *ex vivo*. FecB-TetON grew and established a high titer chronic infection in the lungs
of mice fed with doxy-containing chow (Fig. 7B). In contrast, in mice receiving regular chow, FecB-
TetON grew during the first 14 days post-infection (p.i.), after which CFU counts remained
relatively stable for another 8 weeks p.i. Dissemination to the spleen was delayed, and similarly
to the lung, spleen bacterial burden was significantly lower in mice receiving regular versus doxy
chow (Fig. 7C). Thus, we conclude that *fecB* is required for Mtb virulence and plays a critical role
during growth in the acute phase of infection but is not required for Mtb's persistence in the host.

Discussion

The mycobacterial cell wall, comprised of peptidoglycan, arabinogalactan and mycolic acids, has
been a key target of several anti-TB drugs[33]. Two of the four first-line TB drugs, isoniazid and
ethambutol, target mycolic acid and arabinogalactan biosynthesis, for instance. Although these
drugs were developed more than 60 years ago, they remain part of the standard TB chemotherapy
due to their success in treating drug-susceptible Mtb infections, which validates efforts to explore
novel inhibitors that target the cell wall of Mtb[10]. Despite being an important target for drug
development and playing a central role in Mtb pathogenesis and intrinsic drug resistance, many
aspects of the Mtb cell wall, and more broadly, cell envelope, such as assembly and maintenance
remain unknown. In the current study, we describe two periplasmic proteins, FecB and Rv3035,
that work together to mediate cell envelope impermeability and intrinsic multidrug resistance in
Mtb.

Our transcriptomic data indicate that Mtb lacking *fecB* upregulates the *iniBAC* operon (**iso**n**i**azid
induced genes B, A and C), which is named for being induced by isoniazid, ethambutol, and other
inhibitors of cell wall biosynthesis[10]. Upregulation of the *iniBAC* genes in Δ*fecB* occurred
independently of the presence of detergent in the media and most likely reflects a constitutive
defect in cell wall integrity, which results in Δ*fecB's* increased membrane permeability[5].

The iron-deprivation transcriptional signature and accumulation of siderophores in Δ*fecB* grown
in detergent-free media points to *fecB* having an activity in iron homeostasis. However, whether
this is a direct or indirect effect remains unclear. While FecB can bind ferric-carboxymycobactin
with high affinity *in vitro*[7], experiments probing the requirement of *fecB* for iron acquisition *in vivo*
showed only minor growth phenotypes (Fig. 1D and fig. S1B). FecB variants with lower binding
affinity to ferric-carboxymycobactin (FecB E339S_R240S and E339S_Y242S) rescued the iron-
deprivation transcriptional signature and slower growth of Δ*fecB* in iron-limiting conditions,

suggesting that binding to iron-loaded siderophores is dispensable in FecB's ability to reverse
those phenotypes. On the other hand, FecB E339S_R240S and E339S_Y242S have slightly
higher binding affinity to apo-carboxymycobactin and thus we could not test the *in vivo*
requirement of FecB's binding to iron-free apo-siderophores using these variants. Importantly,
using FecB E339S_R240S and E339S_Y242S, and other genetic and microbiological
approaches, we showed that FecB's role in iron homeostasis is likely not connected to its role in
mediating antibiotic resistance in Mtb.

Specific evidence for FecB's function in cell envelope integrity came to light upon identification of
FecB's most enriched physical interactors: Rv3035, AftB and Rv0227c. In *Mycobacterium*
marinum, FecB was also found to interact with AftB and the Rv3035 homologue MMAR_1667[34].
Although AftB's function has been well described in the literature[20,35,36], only a few studies explore
the function of Rv0227c[23,37] and Rv3035[34] homologues in mycobacteria. Here we report on the
structure of Rv3035 alone and the FecB-Rv3035 complex, which identified key FecB residues
involved in binding Rv3035 that we were able to validate. Moreover, we show that FecB's binding
to Rv3035 is required for its function in mediating antibiotic resistance in Mtb. MtbΔ*rv3035* and
the double knockout Δ*fecB*Δ*rv3035* displayed the same phenotypes indicative of perturbed cell
envelope integrity as Δ*fecB*. This indicates that the absence of either protein equally impacts the
function of the complex and thus both are required to fulfill their function.

Structural homology analysis for Rv3035 revealed two classifications of structural homologs (table
S2). One class represents assembly factors that contribute to scaffolding in larger complexes,
facilitating the coordinated movement of multiple protein components. This includes BamB of the
prokaryotic β-barrel assembly machinery (BAM) complex, and the eukaryotic ODA16, an
assembly factor involved in outer dynein arm intraflagellar transport[38]. When comparing these
structural homologs to Rv3035, both the N-terminal propeller cavity plug region and the α-cap
present in Rv3035 are absent in these assembly factors. The second class of Rv3035 structural
homologs have enzymatic activity. Strikingly, these structural homologs - *Acidithiobacillus*
ferrooxidans tetrathionate hydrolase[39], *Methylacidiphilum fumariolicum* SolV methanol
dehydrogenase[40], and the *Devosia albogilva* quinone-dependent dehydrogenase[41] - all have
secondary structure elements similar to the Rv3035 α-cap and bear an N-terminal propeller
central cavity plug. However, in these enzymatic structural homologs, the propeller fold is more
extensively decorated than in Rv3035. Indeed, Rv3035—due to the presence of N-terminal
propeller cavity plug and the propeller decorations—has greater similarity with its enzymatic
structural homologs than the assembly factor homologs. Although our studies show that Rv3035
acts as a scaffolding protein in its interactions with FecB, its resemblance to the enzymatic
homologs raises the possibility that Rv3035 may possess an as-yet unidentified enzymatic
activity.

What is the function of FecB and Rv3035? And how do these proteins contribute to intrinsic
multidrug resistance in Mtb? Our initial attempts at answering these questions point to FecB's and
Rv3035's interaction with AftB. Corroborating our proteomics dataset, the genes encoding FecB,
Rv3035 and AftB share CRISPRi essentiality profiles, and AlphaFold 3 predicts a very high
confidence model of the complex formed by the three proteins. In this model, Rv3035 bridges the
interaction of FecB and AftB, and the two periplasmic proteins are positioned in close proximity to

AftB's active site cavity[42]. The proposed orientation suggests that the FecB-Rv3035 complex
could serve as scaffold for AftB's attachment to its substrate AG or play a role in controlling AftB's
arabinofuranosyltransferase activity. If one of these hypotheses is correct, one would expect
reduced terminal arabinofuranose residues (mycolylation sites) on AG and, consequently, lower
levels of MA covalently attached to AG which would compromise the structure and integrity of the
cell wall[35]. A decreased ratio of arabinose to galactose in AG was observed in a *M. marinum fecB*
mutant, supporting the hypothesis that FecB influences AftB's function[34]. Additionally, an
impairment of AftB's function leads to production of a truncated LAM that is pro-inflammatory[20]
and can also influence cell envelope integrity[34,43].

Our lipidomics dataset further supports the hypothesis that FecB participates in AftB's function as
the increased levels of TMMs, TDMs and free MAs observed in Δ*fecB* could be a direct
consequence of reduced attachment of MAs to the cell wall (fig. S14). This phenotype is observed
in Mtb treated with ethambutol[44] (inhibitor of AG biosynthesis), and in deletion mutants of *aftB*[20]
and *aftC*[31] (another arabinofuranosyltransferase) in *C. glutamicum* and *M. smegmatis*,
respectively, which all show increased levels of trehalose mycolates. Finally, while Δ*fecB*, Δ*rv3035*
and *aftB* KD strains showed similar increased incorporation of DMN-Tre into the MOM, this
phenotype was enhanced when *aftB* was selectively depleted in Δ*fecB* (Fig. 6J). Moreover, such
depletion also completely inhibited the growth of Δ*fecB* (Fig. 6A)*.* This indicates that residual
AftB's activity remains in Δ*fecB* and further disruption of mycolate turnover is possible by targeting
aftB in the absence of *fecB*. Based on this, we posit that FecB, and by extent Rv3035, support
but are not essential for AftB's arabinofuranosyltransferase activity *in vivo*. This new
understanding of the requirements for cell envelope integrity in Mtb can be used to inform future
TB drug discovery efforts.

Materials and Methods

Bacterial strains and culture conditions

Mtb strains used in this study are listed in data S5. For most experiments, the strains were cultured
in Middlebrook 7H9 medium (BD, #271310) supplemented with 0.2% glycerol, 0.5% bovine serum
albumin fraction V, 0.2% dextrose, 0.085% NaCl and 0.05% Tween80 (Sigma, # P8074), this is
referred to as "regular" medium. For detergent-free conditions, Tween80 was omitted from the
media recipe. For growth in iron-limiting conditions, self-made 7H9 medium without ammonium
ferric citrate (AFC) was prepared and chelated for 2 days with 20g/liter of BioRad Chelex 100
resin. After filter-sterilization, metals other than iron were added according to their concentrations
in Middlebrook 7H9. To prepare Low/High Iron media, 1µM or 160µM of AFC was added to iron-
chelated medium, respectively. For reference, Middlebrook 7H9 medium contains 150µM AFC.
Where required, antibiotics or small molecules were used at the following concentrations:
hygromycin (50µg·ml^{-1}), kanamycin (25µg·ml^{-1}), zeocin (25µg·ml^{-1}), streptomycin (25µg·ml^{-1}),
apramycin (60µg·ml^{-1}) and ATc (200ng·ml^{-1} for CRISPRi strains and 1000ng·ml^{-1} for FecB-
TetON). The strains were grown at 37°C and 5% CO_2 in a standing incubator, unless otherwise
indicated.

Plasmids construction

Plasmids used in Mtb strains are listed under the strain "full name" after the double colon in data S5. Plasmids starting with pGM were constructed using Invitrogen Gateway Cloning system. Constructs harboring the gene of interest were synthesized by GenScript. CRISPRi constructs were generated as previously described[19].

RNA extraction

Total RNA was extracted from 10mL of mid-log phase bacterial cultures by first adding 1:1 guanidinium thiocyanate buffer followed by centrifugation for 10 minutes at 4°C. Bacterial pellets were bead-beaten into 0.9mL Trizol, 0.2mL chloroform was added to each sample and RNA in the aqueous phase was purified using the RNA Clean and Concentrator-25 kit (Zymo Research #R1017). After elution, samples were incubated with TURBO DNase (Ambion, AM2238) at 37°C for 1h to remove residual genomic DNA. Samples were then repurified using the RNA Clean and Concentrator-25 kit or RNA Clean and Concentrator-5 kit (Zymo Research #R1015).

RNA-sequencing (RNA-seq) and analysis

Ribosomal RNA (rRNA) was removed from total RNA as previously described[45]. The resulting rRNA-depleted RNA was then concentrated with the RNA Clean and Concentrator-5 kit (Zymo Research #R1015). RNA-seq libraries were created from these samples using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New England Biolabs, E7760L) along with NEBNext Multiplex Oligos for indexing (New England Biolabs). The libraries were purified using Dynabeads MyOne Streptavidin C1 (Thermo Fisher #65002), assessed for quality with a Bioanalyzer, and pooled based on their unique indexes. Sequencing was carried out on an Illumina NextSeq2000 platform at the Genomics Resources Core Facility of Weill Cornell Medicine.

RNA-seq data analysis was conducted using R. Sequencing reads underwent trimming and filtering with Rfastp, followed by alignment to the H37Rv reference genome (GenBank AL123456.3; RefSeq NC_000962.3; GCF_000195955.2; Genome assembly ASM19595v2) using the Rsubread package[46,47]. BAM alignment files were generated, sorted, and indexed using Rsamtools. Read quantification was carried out with featureCounts, and differential gene expression analysis was performed using DESeq2[48,49].

Lipid extraction

Mtb cultures were grown in detergent-free 7H9 containing fatty acid-free BSA until mid-log phase. Bacterial pellets were transferred to sterile filters[50] which were added directly to 16 x 100mm glass tubes containing 2:1 chloroform: methanol or to 2mL screw cap tubes containing 1mL of 10mM

AOT dissolved in heptane or PBS as a control. For AOT treatment, pellets in AOT or PBS were
agitated in a bead-beater at low speed (2800rpm) for 10s to knock the cells off the filters followed
by removal of filters and incubation in a cooling rack for 1h, with mixing by inversion of the tubes
every 5 minutes. 500µL of AOT supernatant was added to 16 x 100mm glass tubes containing
2:1 chloroform: methanol. AOT and PBS pellets were resuspended in 600µL of 2:1 chloroform:
methanol and added to 16 x 100mm glass tubes containing 2:1 chloroform: methanol.

Liquid chromatography-mass spectrometry (LC-MS) analysis

Lipid extracts were centrifuged at 4000rpm for 15 minutes at 4°C and the supernatant decanted
into new 16 x 100mm borosilicate tubes. The tubes were dried using a GeneVac evaporator under
vacuum with gentle heating to 40°C. The residual residue was transferred to pre-weighed 4mL
amber vials using 2mL 1:1 chloroform: methanol. The solution was again dried using the GeneVac
evaporator. The vials were re-weighed to obtain the dry weight of lipid extract, and this mass was
used to prepare a 1mg/mL solution in 1:1 chloroform: methanol. Lipids were stored at -20°C as a
stock solution. A 150µL aliquot of stock solution was transferred to a glass insert tube (250µL,
Agilent Technologies) nested within a microfuge tube. The aliquot was evaporated to dryness
using an argon air stream then resuspended in 150µL of a solution of 70% hexanes and 30%
isopropyl alcohol. After briefly vortexing to resuspend, the samples were centrifuged at 4000rpm
for 15 minutes at 4°C. A volume of 100µL of the supernatant was transferred to a 2mL amber vial
(Agilent) equipped with 250µL glass insert tubes and capped with 9mm septum-embedded screw
caps (Agilent). A 10µL injection of this sample was analyzed using liquid chromatography mass
spectrometry (LC-MS) on an Agilent 6520 ESI-QTOF. Separation of lipids using a 3µm diol column
(2.1 x 150mm) and gradient elution from solvent A (70% hexane, 30% isopropanol, .02% formic
acid, .01% ammonium hydroxide) to solvent B (70% isopropanol, 30% methanol, .02% formic
acid, .01% ammonium hydroxide) as previously described[51]. Following chromatography, the
sample underwent electrospray ionization at 325°C with 5 L/min drying gas flow at 5500 V and
30 psig nebulizer pressure and ions were detected by time-of-flight in both positive and negative
modes (m/z range: 100 – 3000). Internal calibrants (922.009798 positive mode, 983.0325
negative mode) were infused continuously.

Lipids were identified using unique mass-retention time identifiers for masses exhibiting the
expected isotopic distributions as previously reported[51]. Integrations of ion abundance were
determined using Agilent Profinder 8.0 and Qualitative Analysis 6.0.

Growth curves

Mtb strains were inoculated from frozen glycerol stocks and grown with appropriate selection
antibiotics in 7H9 medium until mid-log phase at 37°C.

For pre-depletion of iron internal stores, bacteria were washed in PBS or iron-chelated 7H9 and
diluted to an OD_{580nm} of 0.05 in iron-chelated 7H9. After 4-5 generations, bacteria were spun down
at 800rpm for 8 minutes to generate a single cell suspension. Bacteria were diluted to an OD_{580nm}

of 0.025 in the indicated media and added to flasks which were incubated standing at 37°C. For
serial passaging, when cultures in high iron media reached stationary phase, low iron cultures
were spun down at 800rpm for 8 minutes to generate a single cell suspension and bacteria were
diluted to an OD_{580nm} of 0.025 in both high/low iron media. OD measurements from cultures in
flasks were performed in a spectrophotometer. For growth curves in low iron media without
detergent, iron pre-depleted cultures were washed in PBS or detergent-free iron-chelated 7H9,
diluted to an OD_{580nm} of 0.05 and 100μL of bacteria were added to a 96-well plate in triplicates.
OD measurements from plates were performed in a plate reader. Cultures grown in 96-well plates
were resuspended prior to OD_{580nm} readings.

For CRISPRi strains, cultures were split into two flasks and 200ng/mL of ATc added to one of the
flasks 2-3 days prior to setting up the growth curve to pre-deplete the gene of interest. Fresh ATc
(200ng/mL) was again added to the media (+ATc conditions) before distributing bacteria into
96/384-well plates in triplicates.

Reverse transcription quantitative polymerase chain reaction (RT qPCR)

Primers and probes for RT qPCR are listed in table S4. Primers and probes were ordered from
TIB MOLBIO, LLC. cDNA was generated using MluV reverse transcriptase (New England Biolabs)
and approximately 500ng of total RNA. Parallel no-reverse transcriptase reactions were prepared
as a control for residual genomic DNA. qPCR was performed in the LightCycler® 480 System
(Roche). The housekeeping gene *sigA* (Rv2703) was used as an internal control. *sigA* probes
were labeled with 5' LC760/3' BBQ. Probes targeting genes of interest were all labeled with 5'
6FAM/3' BBQ or 3' BHQ.

Minimal inhibitory concentrations (MIC) assays

Mtb strains were inoculated from frozen glycerol stocks and grown with appropriate selection
antibiotics in 7H9 medium until mid-log phase at 37°C. Bacteria were diluted to an OD_{580nm} of 0.01
in 7H9 and 50μL of bacteria were added to a 384-well plate in triplicates per antibiotic
concentration, or to triplicate control wells without antibiotics. The HP D300e digital dispenser was
used to deliver antibiotics to plate wells. Bacteria were grown in standing cultures at 37°C and
OD_{580nm} was measured in a plate reader between days 10-21 following treatment (time point is
indicated in each figure legend). For MICs in low iron or detergent-free conditions, bacterial
cultures were first washed in PBS once or twice prior to dilution into the respective media of the
MIC assay. For CRISPRi strains, cultures were split into two flasks and 200ng/mL of ATc added
to one of the flasks 2-3 days prior to setting up the MIC assay to pre-deplete the gene of interest.
Fresh ATc (200ng/mL) was again added to the media of the assay (+ATc conditions) before
distributing bacteria into plates.

Pull-down tandem mass spectrometry

Sample preparation

Whole-cell lysates were collected from log-phase cultures grown in the shaking incubator,
incubated with 1% n-dodecyl -D-maltoside for 2h on a rotator at 4°C, followed by incubation with
anti-DYKDDDDK (FLAG) magnetic agarose (Pierce #A36797). Captured proteins were
dissociated from magnetic agarose by addition of 3x DYKDDDDK (3xFLAG) peptide (Pierce
#A36805), according to the manufacturer's instructions. Sodium Dodecyl Sulfate (SDS) was
added to the peptide eluate at a final concentration of 2% and samples were boiled at 95°C for
10 minutes. Boiled samples were transferred to protein lo-bind tubes. FLAG- (or 3xFLAG-) tagged
bait proteins were detected in eluates on an SDS-Polyacrylamide Gel Electrophoresis (PAGE) gel
prior to mass spectrometry.

23µg of proteins per sample were flash frozen on dry ice and subsequently dried using a Speed-
Vac. Next, 23µl of lysis buffer (5% SDS, 50mM TEAB) was added to initiate the S-Trap™ Micro
Spin digestion protocol (PROTIFI). The proteins underwent reduction by 4mM dithiothreitol (DTT)
and alkylation by 20mM iodoacetamide. Following an overnight digestion with trypsin at 37 °C,
peptides were eluted first by addition of 40µL of 50mM triethylammonium bicarbonate (TEAB) and
spun at 4000 rpm for 1 min. This was followed by a 1-minute spin with 40µL of 0.2% formic acid
(FA) and, lastly, 40µL of 50% acetonitrile (ACN). The three elution were then pooled together,
dried using a Speed-Vac, and reconstituted in 20µL of 0.1% FA in 5% ACN. After reconstitution,
the samples were centrifuged at 16,000 rpm for 16 min and an 18µL aliquot of the supernatant
was transferred to a non-binding HPLC vials.

LC-MS/MS and data analysis workflows

Data from the first experiment (pull-down of FecB-FLAG) was acquired on a TimsTOF Pro2
(Bruker) mass spectrometer, which was coupled to a nanoElute LC system from Bruker. 5µL of
peptides were loaded and separated on an in-house 75 µm I.D. fused silica analytical column
packed with 25-cm ReproSil-Pur C18-AQ (Dr. Maisch, GmbH, 120 Å, 3 µm) particles to a gravity-
pulled tip, using an aqueous mobile phase (A) of water and 0.1% FA and an organic mobile phase
(B) of ACN and 0.1% FA. A 30-minute gradient was employed for DDA-PASEF with a flow rate
set at 500 nL/min. The captive nano-electrospray voltage was maintained at 1600V, using one
column configuration (no trap). The DDA-PASEF method consists of 10 MS/MS PASEF scans per
topN acquisition cycle, with ramp and accumulation times of 100 ms, covering an m/z range from
100 to 1700 and an ion mobility range (1/K0) from 0.70 to 1.30 V s/cm^2. Collision energy settings
followed a linear function of ion mobility, ranging from 20 eV at 0.6 V s/cm^2 to 59 eV at 1.6 V
s/cm^2, using default parameters. Calibration of the instrument was performed using three ions
from the ESI-L Tuning Mix (Agilent) (m/z 622, 922, 1222).

The custom workflow 'LFQ-MBR' from FragPipe version 21.1 was used, including database
search with MSFragger (version 4.0), deep-learning prediction rescoring with MSBooster,
Percolator and ProteinProphet (Philosopher version 5.1.0) for PSM validation and protein
inference. The raw files from TimsTOF Pro2 were searched against the database
"Mycobacterium_tuberculosis_H37Rv_v4modified" appended with common contaminant
proteins. Decoy reversed sequences were appended to the search database. The default

MSFragger search parameters were used, except precursor and fragment mass tolerances were
set to 20 ppm; enzyme: trypsin (not cutting before P); peptide length 7-30. Variable modifications
were set to oxidation of Met and acetylation of protein N-t, and fixed modifications to
carbamidomethylation of Cys. MSBooster, Percolator and ProteinProphet default options were
used, and results were filtered by 1% FDR at protein level. The searched results were then
imported to Scaffold 5 Q+S (Proteome Software Inc., v5.3) for data visualization.

Data from the second experiment (pull-down of Rv3035-3xFLAG) was acquired using a
NanoAcquity UPLC (Waters Corporation, Milford, MA) coupled to an Orbitrap Fusion Lumos
Tribrid (Thermo Fisher Scientific, Waltham, MA) mass spectrometer. Peptides were trapped and
separated using an in-house packed precolumn packed with 2cm ReproSil-Pur C18-AQ (Dr.
Maisch, GmbH, 120 Å, 5 µm) particles plus and an in-house 75 µm I.D. fused silica analytical
column (as previously described) and the same solvent composition. 3.8µL sample was injected,
and the peptides were trapped at a flow rate of 4µL/min with 5% mobile phase B for 4 min, followed
by a gradient elution with 5-35% B at a flow rate of 300nL/min over 120 min (total run time 145
min). Mass spectra were acquired over m/z 375-1500 Da with a resolution of 120,000 (m/z 200).
Tandem mass spectra were acquired using data-dependent acquisition with an isolation width of
1.6 Da, HCD collision energy of 30%, resolution of 15,000 (m/z 200), maximum injection time of
50 ms, and an AGC target of 50,000.

Raw data files from the Orbitrap Fusion Lumos Tribrid were peak processed with Proteome
Discoverer (version 2.5, Thermo Scientific) followed by identification using Mascot Server (Matrix
Science, v2.6.2) against the same database used in the previous experiment. Search parameters
were as follows: tryptic digestion with up to 2 missed cleavages; peptide N-terminal acetylation,
methionine oxidation, and N-terminal glutamine to pyroglutamate conversion were specified as
variable modifications. Carbamidomethylation of cysteines was set as static modification.
Assignments were made using a 10-ppm mass tolerance for the precursor and 0.6 Da mass
tolerance for the fragment ions. All non-filtered search results were processed by Scaffold 5 Q+S
at 1.0% false-discovery rate (FDR) for peptides and 99% threshold for proteins (2 peptides
minimum).

For each experiment, Total Spectrum Count (TSC) was exported from Scaffold 5 Q+S into a
Microsoft Excel spreadsheet. Zeros were replaced with ones in WT control samples to allow fold-
change (FC) calculation of every protein detected in the samples containing the bait protein. For
normalization, we summed the TSC values for all proteins in each replicate, calculated the
average of the sum of the three replicates and divided the average by the sum value of each
replicate (this is the normalization factor). Then, we multiplied each protein TSC by the
normalization factor of that replicate sample, calculated the median, log2FC of tagged group over
WT control, and p-value by performing T-tests of the normalized values in each row. P-values
were transformed into -log10(p-value) for visualization of the data as a volcano plot.

Co-essentiality analysis

We performed co-essentiality analysis on CRISPR interference (CRISPRi) data from Li, Poulton
et al.[13], which profiled Mtb fitness under 80 experimental conditions. These conditions span 9

antibiotics (BDQ = bedaquiline; CLR = clarithromycin; EMB = ethambutol; INH = isoniazid;
LVX = levofloxacin; LZD = linezolid; RIF = rifampicin, STR = streptomycin; VAN = vancomycin),
each tested under 9 screening conditions defined by a combination of pre-depletion time (1, 5, or
10 days of CRISPRi induction before drug exposure) and drug concentration ("Low," "Medium,"
or "High" partially inhibitory doses). We used only the $\log_2$ fold change (LFC) values from negative
selection screens, which report the fitness effects of gene knockdown.

Each gene was thus represented by an 80-dimensional LFC vector capturing its essentiality
profile across the drug/time/concentration matrix. To account for covariance across screens—
such as batch effects—we computed an 80 × 80 empirical covariance matrix (Σ) from the LFC
data. We then applied generalized least squares (GLS), following the approach of Wainberg *et*
al.[52] to compute pairwise co-essentiality relationships among 3,979 genes. This yielded a 3,979
× 3,979 matrix of GLS coefficients and corresponding p-values.

We corrected for multiple hypothesis testing using the Benjamini–Hochberg procedure
(fdrcorrection in the Python statsmodels package). Gene pairs with a false discovery rate (FDR)-
adjusted p-value < 0.001 were considered significantly co-essential, resulting in approximately
16,000 high-confidence gene pairs.

Design of expression constructs for *M. thermoresistibile* (Mth) Rv3035

Initial attempts to express Mtb Rv3035 in *E. coli* using N- or C-terminal 6xHis-tag (His) constructs
with full-length and various N-termini truncations failed to produce soluble Rv3035. To improve
solubility, we turned to a thermostable homolog of Rv3035, Mth Rv3035, Accession no.
WP_040546537, and constructed two *E. coli* codon-optimized constructs cloned into pET28a with
a TEV-cleavable N-terminal His-tag (GenScript, Piscataway, NJ). The Mth Rv3035 construct that
encodes for the mature form of Rv3035 without its predicted signal peptide (SignalP 6.0[14]) and
starts at residue Gly2 (fig. S4B), produced soluble Rv3035 only when expressed in *E. coli*
T7SHuffle cells (New England Biolabs, Ipswich, MA). Notably, T7SHuffle cells promote disulfide
bond formation[53], and Rv3035 is predicted to harbor two disulfide bonds.

Generation of FecB mutants for *in vitro* studies

A pET28a FecB construct encoding Mtb FecB (without the signal peptide, mature FecB starting
at residue 39) with a thrombin-cleavable N-terminal 6xHis-tag[7] was used as the template for *in*
vitro site-directed mutagenesis using Pfu Ultra Fusion HS DNA polymerase (Agilent) with the
primers listed in table S5 and then confirmed by DNA sequencing (GeneWiz from Azenta Life
Sciences). The FecB construct encoding no His-tag was generated by using the pET28a FecB
plasmid and excising the encoded *fecB* gene with *Nde*I and *BamH*I and ligating it into a pET22b
vector.

Expression and purification of Mth Rv3035 and the Mth Rv3035-Mtb FecB complex

Mth Rv3035-His was expressed in *E. coli* T7SHuffle cells in Terrific Broth (TB) media containing
0.4% glycerol and 50µg/mL kanamycin (Kan). To express the Mth Rv3035-Mtb FecB complex,
which we refer to as the FecB-Rv3035 complex throughout the manuscript, both pET28a Mth
Rv3035-His and pET22b Mtb FecB plasmids were co-transformed into T7SHuffle cells with
selection on LB + 50µg /mL Kan and 100µg/mL ampicillin (Amp) agar plates. A fresh transformant
was used to inoculate TB containing 0.4% glycerol, 50µg/mL Kan and 100µg/mL Amp. Cells were
grown at 37°C to an OD_{600nm} of ~ 0.6, before protein expression was induced by the addition of
0.5mM isopropyl β-D-1-thiogalactopyranoside (IPTG). The cells grown overnight at 20°C and
harvested by centrifugation.

Both Mth Rv3035-His and the FecB-Rv3035-His complex were purified using the same protocol.
Cell pellets were resuspended in 10x vol(mL)/w(g)of cells in Buffer A (150mM NaCl, 50mM Tris,
pH 7.8) with 0.25mM phenylmethylsulfonyl fluoride (PMSF) with stirring at 4°C for 30 minutes.
Cells were lysed by sonication and cell debris was pelleted by centrifugation at 14,000rpm for 1h
at 4°C. The cell lysate was passed through a 1µm syringe filter to remove insoluble debris and
applied to a His-trap column (Cytiva, Marlborough, MA) pre-equilibrated with Buffer A using an
AKTA Start FPLC instrument (Cytiva, Marlborough, MA). Following a 10x column volume (CV)
wash step with Buffer A, a gradient of Buffer A from 0 to 500mM imidazole over 20 CVs elute His-
tagged proteins. Fractions with containing Mth Rv3035-His or the FecB-Rv3035-His complex
were concentrated using an Amicon concentrator (MilliporeSigma, 30kDa cut-off) with buffer
exchange into Buffer A.

Expression and purification of Mtb FecB and mutants

Purification of WT FecB and FecB mutant proteins were carried out in a similar manner as
previously described[7]. FecB WT or mutants were expressed in BL21-(DE3) cells and cells were
grown in TB + 0.4% glycerol + 50µg/mL kanamycin at 37°C to an OD_{600nm} of ~0.6, before protein
expression was induced with IPTG (0.5mM). Cell growth was continued overnight for 20h at 20°C,
before cells were harvested by centrifugation. Cell pellets were resuspended in 10x vol(mL)/w(g)
Buffer B (350 mM NaCl, 50 mM Tris, pH 7.8, 10% glycerol) with 0.25mM PMSF with stirring at
4°C for 30 minutes. Cells were lysed by sonication and cell debris was pelleted by centrifugation
at 14,000rpm for 1h at 4°C. The cell lysate was filtered and applied to a Ni-NTA resin pre-
equilibrated with Buffer B using gravity flow. The column was washed with Buffer B and eluted
with a stepwise elution gradient of imidazole going up to 600mM imidazole in Buffer B. Fractions
found to contain FecB by SDS-PAGE analysis (expected MW ~37 kDa) were pooled and dialyzed
against Buffer B overnight at 4°C.

Removal of N-terminal His-tag from Rv3035-His and FecB-His

For cleavage of His-tags from Rv3035-His and from the FecB-Rv3035-His complex, TEV was
added to approximately 30-100mg of protein at a final concentration ~0.1mg/ml TEV and
incubated overnight at 4°C. The cleavage reaction was passed over a Ni-NTA column pre-
equilibrated with Buffer A + 10mM imidazole and chased with Buffer A + 10mM imidazole. Flow-

through fractions containing Rv3035 or FecB-Rv3035 complex were concentrated and buffer
exchanged into Buffer A.

For the isolation of tag-free WT FecB, which was used for some ITC experiments, a similar
procedure was used but with thrombin at ~30U thrombin:1mg FecB-His. The cleavage reaction
was passed over a Ni-NTA column pre-equilibrated with Buffer B + 10mM imidazole and chased
with Buffer B + 10mM imidazole. Flow-through fractions containing WT FecB were concentrated
and buffer exchanged into Buffer B.

Isothermal titration calorimetry (ITC) experiments

ITC experiments were carried out using a Malvern MicroCal PEAQ-ITC instrument (Malvern,
United Kingdom). Prior to the experiments, proteins were dialyzed into Buffer B and the same
dialysis buffer was used for protein dilutions. 19-point titrations were carried out using 200µM Mth
R3035 in the syringe and 20µM FecB in the cell, whereas 13-point titrations were carried out using
300µM Rv3035 and 30µM FecB-His mutant protein, or 200µM Rv3035-His and 20µM FecB-His.
Notably, the His-tag on FecB did not appreciably change the affinity of FecB for Rv3035. Curve
fitting was carried out using the Malvern instrument analysis software.

Structure determination of Mth Rv3035 and the FecB-Rv3035 complex

Initial screening of Mth Rv3035 (15mg/mL) and FecB-Rv3035 complex (38mg/mL) were carried
out in 96-well trays using a Mosquito liquid handler (SPT Labtech, Cambridge, MA) and
commercially available crystallization screens by the hanging drop vapor diffusion method.

Mth Rv3035 alone

Diffraction-quality Mth Rv3035 crystals were grown with a crystallization reservoir of 0.1M sodium
chloride; 0.1M HEPES, pH 7.5; 1.6M ammonium sulfate with Rv3035 at 15 mg/mL and a reservoir
to protein ratio of 1:1 at room temperature. The crystals were cryocooled in mother liquor
supplemented with 25% glycerol. Data was collected at SSRL 12-2 at the wavelength 1 Å. Data
was processed in xds[54] to a resolution of 1.9 Å (table S1). A model of Mth Rv3035 was generated
on the AlphaFold server[25] and was used for molecular replacement (MR) in phaser[55] in phenix.
The MR solution placed two subunits in the asymmetric unit (ASU) and was improved by autobuild
in phenix[56], and refined in phenix.refine[57]. Following autobuild, the N-terminal residues 10-37 were
built *de novo* in coot[58]. The N-terminal 9 residues of Rv3035 could not be modeled for both
subunits, and for subunit B, the last C-terminal residue also could not be modeled due to
unresolved electron density. The model was further refined in phenix.refine and coot to a final
R_{work}/R_{free} of 16.1 / 20.25 (table S1).

FecB-Rv3035 complex

Diffraction-quality crystals of the FecB-Rv3035 complex were obtained with a crystallization
reservoir of 3M sodium chloride; 0.1M Bis-Tris, pH 7.0 with a reservoir to protein complex ratio of
1: 1. To further improve diffraction quality of the FecB-Rv3035 complex, an additive screen was
performed (Hampton Research, Aliso Viejo, CA), with the final conditions used being a
crystallization reservoir of 3M sodium chloride; 0.1M Bis-Tris, pH 7.0; 3% trehalose, with the
Rv3035-FecB complex at 38mg/ml and a reservoir to protein ratio of 1:1. Crystals were
cryoprotected in reservoir solution with 25% glycerol and flash frozen with liquid nitrogen. Data
was collected at SSRL 12-2 at the wavelength 1 Å.

Data was processed in XDS[54] to a resolution of 3.3 Å (table S1). The phases were solved by MR
in phaser[55] in phenix, placing a single subunit of the deposited FecB structure (PDB ID 7UQ0,[7])
in the ASU in complex with a single subunit of the Mth Rv3035 structure solved above. The MR
solution was improved in autobuild and refined as described above in phenix.refine[57] and coot[58]
to a final R_{work}/R_{free} of 25.2/26.3 (table S1).

Knockout construction

Δ*rv3035* and Δ*fecB*Δ*rv3035* mutants were generated using a suicide plasmid approach as
previously described[59] with a few modifications. Constructs containing 500bp fragments
corresponding to regions upstream and downstream of *rv3035* were ordered from GenScript and
cloned into a temperature-sensitive deletion vector pDE43-XSTS to flank the zeocin resistance
cassette and generate pKO-XSTS-Zeo-*rv3035* via Gateway Cloning. Mtb WT and Δ*fecB* were
transformed with both the pKO-XSTS-Zeo-*rv3035* and a *rv3035* complementation plasmid
(pGMCS-T02-P*hsp60*-SD1-*rv3035*_A-HA) to generate a merodiploid strain since initial attempts
of generating the knockout with pKO-XSTS-Zeo-*rv3035* alone had failed. Transformants were
plated on Middlebrook 7H10 agar (BD #262710) containing 0.5% glycerol, 10% OADC (BD
#212351) and 25µg/mL Zeo (for WT) or 50µg/mL Hyg and 25µg/mL Zeo (for Δ*fecB*) and incubated
at 37°C until colonies were approx. 1mm in diameter. Colonies were tested for reporter gene XylE
activity directly on plates by addition of a drop of catechol solution (1g in 20mL water). After 30
minutes, candidate colonies that turned yellow were selected and grown in regular 7H9 with
appropriate antibiotics until they reached an OD_{580nm} of 1-1.5. Candidates were then grown on
7H10 agar with 10% sucrose at 37°C until colonies were approx. 1mm in diameter. Colonies were
test for reporter gene activity as above, but this time white colonies (that had successfully lost the
suicide plasmid) were selected. Genomic DNA (gDNA) was extracted from a few mutant
candidates and allelic exchange was confirmed by PCR using primers listed in table S6. To obtain
the final knockout mutants, we replaced the *rv3035* complementation plasmid integrated into attL5
with an empty plasmid containing an apramycin resistance cassette (pGMCA-OXOX).
Transformants were plated on 7H10 containing Zeo (or Zeo and Hyg) and 60µg/mL apramycin.
Δ*rv3035* and Δ*fecB*Δ*rv3035* mutants took about 6-8 weeks to form visible colonies. Candidate
mutants were selected and tested for streptomycin sensitivity in liquid culture as an indication that
the copy of *rv3035* (carrying streptomycin resistance) was successfully replaced in attL5. Finally,
gDNA was extracted for whole genome sequencing to confirm deletion of *rv3035* and insertion of
zeocin and apramycin cassettes.

Cell envelope permeability assays

BODIPY Fluorescent (FL) vancomycin (Invitrogen #V34850) uptake was performed as previously reported[5] with a few modifications. Briefly, mid-log phase Mtb cultures were washed once and adjusted to an OD_{580nm} of 2 in PBS and incubated with 2µg/mL of BODIPY FL vancomycin. 350µL sample aliquots were taken at the 30 minutes incubation time point, washed twice in PBS 0.05% Tween80, and resuspended in 350µL PBS. Negative control samples were treated the same way but did not receive BODIPY FL vancomycin. 100µL of each sample was distributed in triplicate wells in a black-sided 96-well plate. Fluorescence was measured (excitation wavelength of 485nm and emission wavelength of 538nm) and normalized to the OD_{580nm} of the final suspension to account for cell loss during washing steps.

For the calcein-AM (Invitrogen #C3099) uptake assay, mid-log phase cultures were adjusted to an OD_{580nm} of 0.8 in regular 7H9 and 100µL aliquots were added to the wells of a black-sided 96-well plate in triplicates. 100µL of calcein-AM (1µg/mL in 7H9 0.4% glucose) was added to the wells resulting in a final concentration of 0.5µg/mL calcein-AM. Fluorescence was measured (excitation wavelength of 495nm and emission wavelength of 520nm) at 2 minutes intervals over a course of 60 minutes. Fluorescence was normalized to OD_{580nm} measured at 60 minutes.

Spot assay

Mig-log phase Mtb cultures were adjusted to an OD_{580nm} of 0.3 and 10-fold serial dilutions in PBS 0.05% Tween80 were spotted on 7H10 agar with 5µL per spot. Plates were incubated at 37°C and imaged after 2-3 weeks. To confirm colony morphology alteration and growth defect of Δ*fecB*, Δ*rv3035* and Δ*fecB* Δ*rv3035* mutants, plates were incubated for 4-5 additional weeks, after which the phenotypes were unchanged.

DMN-Tre labeling and flow cytometry

Mid-log phase Mtb cultures were adjusted to OD_{580nm} of 0.5 in regular 7H9. For CRISPRi strains, cultures were split into two flasks and 200ng/mL of ATc was added to one of the flasks 2-3 days prior to labeling with DMN-Tre to pre-deplete the gene of interest. DMN-Tre probe[30] was added to 500µL of culture at a final concentration of 100µM. Samples were incubated at 37°C overnight for approximately 20h. On the next day, cultures were spun down at 10,000rpm for 3 minutes, and the supernatant containing DMN-Tre was discarded. Samples were fixed with 4% PFA (Biolegend #420801) for 4h at room temperature. After two washes in PBS 0.05% Tween80, bacteria were resuspended in PBS. Data collection was performed on a FACSymphony A5 Cell Analyzer (BD Biosciences) at the Weill Cornell Medicine Flow Cytometry Core Facility using the 405nm violet laser and the 405 [425/50] detector. Fluorescence data were obtained for 200,000 events per sample and analyzed in FlowJo software (BD) by first gating on the bacterial population in the side scatter versus forward scatter plot followed by calculation of the mean fluorescence intensity.

Mouse infection

Sixty-four female, eight-week-old *Mus musculus* C57BL/6 (Jackson Laboratories) were infected with ~100 colony forming units (CFU)/mouse using an Inhalation Exposure System (Glas-Col). Mice in the doxycycline group started to receive doxycycline chow 2 days before aerosol infection. The FecB-TetON strain was grown to mid-log phase in the presence of 1µg/mL ATc, a single cell suspension was prepared in PBS 0.05% Tween80 and then resuspended in PBS for a final OD_{580nm} of 0.2. Lungs and spleen were homogenized in PBS and plated on 7H10 agar containing 1µg/mL ATc to determine CFU/organ at the indicated time points. This experiment was performed in accordance with the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health, with approval from the Institutional Animal Care and Use Committee of Weill Cornell Medicine.

Quantification and statistical analysis

Data analysis and generation of graphs were performed in Prism version 10.5 (GraphPad).

Acknowledgements

We thank Sae Woong Park (Weill Cornell Medicine) for sharing the Δ*mbtK* and Δ*mbtI* strains, Carolina Trujillo (Weill Cornell Medicine) for assistance with the animal experiment, and Allison Fay (Memorial Sloan Kettering Cancer Center) for her advice on the thin layer chromatography experiment. We thank the staff of the University of Massachusetts Chan Medical School Mass Spectrometry Facility and Kadamba Papavinasasundaram for their help with the pull-down tandem mass spectrometry experiments. We thank the Kamariza Lab at the University of California, Los Angeles for gifting an aliquot of the DMN-Tre probe. We thank the staff of the Weill Cornell Medicine Genomics Resources Core Facility and Flow Cytometry Core Facility for their contribution and advice on the RNA-seq and flow cytometry experiments, respectively. We thank the staff of the Advanced Light Source at Berkeley National Laboratories and the Stanford Synchrotron Radiation Lightsource for their invaluable help in data collection.

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. 2139291 and the Potts Memorial Foundation Pre-doctoral Fellowship.

References

1.   2024 Global tuberculosis report.

2.   Poulton, N. C. & Rock, J. M. Unraveling the mechanisms of intrinsic drug resistance in Mycobacterium tuberculosis. *Front Cell Infect Microbiol* **12**, 997283 (2022).

3.   Dulberger, C. L., Rubin, E. J. & Boutte, C. C. The mycobacterial cell envelope - a moving
target. *Nat Rev Microbiol* **18**, 47–59 (2020).

4.   Jackson, M., Stevens, C. M., Zhang, L., Zgurskaya, H. I. & Niederweis, M. Transporters
Involved in the Biogenesis and Functionalization of the Mycobacterial Cell Envelope.
(2020) doi:10.1021/acs.chemrev.0c00869.

5.   Xu, W. *et al.* Chemical Genetic Interaction Profiling Reveals Determinants of Intrinsic
Antibiotic Resistance in Mycobacterium tuberculosis. (2017) doi:10.1128/AAC.01334-17.

6.   Staudenmaier, H., Van Hove, B., Yaraghi, Z. & Braun, V. Nucleotide sequences of the
fecBCDE genes and locations of the proteins suggest a periplasmic-binding-protein-
dependent transport mechanism for iron(III) dicitrate in Escherichia coli. *J Bacteriol* **171**,
2626–2633 (1989).

7.   de Miranda, R. *et al.* Differentiating the roles of Mycobacterium tuberculosis substrate
binding proteins, FecB and FecB2, in iron uptake. *PLoS Pathog* **19**, (2023).

8.   Wang, L. *et al.* Multiple genetic paths including massive gene amplification allow
Mycobacterium tuberculosis to overcome loss of ESX-3 secretion system substrates.
Proc Natl Acad Sci U S A **119**, e2112608119 (2022).

9.   Leisching, G., Pietersen, R. D., Wiid, I. & Baker, B. Virulence, biochemistry, morphology
and host-interacting properties of detergent-free cultured mycobacteria: An update.
Tuberculosis **100**, 53–60 (2016).

10.   Alland, D., Steyn, A. J., Weisbrod, T., Aldrich, K. & Jacobs, W. R. Characterization of the
Mycobacterium tuberculosis iniBAC promoter, a promoter that responds to cell wall
biosynthesis inhibition. *J Bacteriol* **182**, 1802–1811 (2000).

11.   Schapira, M., Tyers, M., Torrent, M. & Arrowsmith, C. H. WD40 repeat domain proteins: a
novel target class? *Nature Reviews Drug Discovery 2017 16:11* **16**, 773–786 (2017).

12.   Jinich, A. *et al.* Genome-wide co-essentiality analysis in Mycobacterium tuberculosis
reveals an itaconate defense enzyme module. *bioRxiv* 2022.09.27.509804 (2022)
doi:10.1101/2022.09.27.509804.

13.   Li, S. *et al.* CRISPRi chemical genetics and comparative genomics identify genes
mediating drug potency in Mycobacterium tuberculosis. *Nature Microbiology 2022 7:6* **7**,
766–779 (2022).

14.   Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using protein
language models. *Nature Biotechnology 2022 40:7* **40**, 1023–1025 (2022).

15.   Smithers, L., Olatunji, S. & Caffrey, M. Bacterial Lipoprotein Posttranslational
Modifications. New Insights and Opportunities for Antibiotic and Vaccine Development.
Front Microbiol **12**, 788445 (2021).

16. Drage, M. G. *et al.* Mycobacterium tuberculosis lipoprotein LprG (Rv1411c) binds
triacylated glycolipid agonists of Toll-like receptor 2. *Nat Struct Mol Biol* **17**, 1088–1095
(2010).

17. Holm, L. & Rosenström, P. Dali server: conservation mapping in 3D. *Nucleic Acids Res*
38, W545–W549 (2010).

18. Dejesus, M. A. *et al.* Comprehensive essentiality analysis of the Mycobacterium
tuberculosis genome via saturating transposon mutagenesis. *mBio* **8**, (2017).

19. Bosch, B. *et al.* Genome-wide gene expression tuning reveals diverse vulnerabilities of
M. tuberculosis. *Cell* **184**, 4579-4592.e24 (2021).

20. Seidel, M. *et al.* Identification of a Novel Arabinofuranosyltransferase AftB Involved in a
Terminal Step of Cell Wall Arabinan Biosynthesis in Corynebacterianeae, such as
Corynebacterium glutamicum and Mycobacterium tuberculosis. *Journal of Biological*
Chemistry **282**, 14729–14740 (2007).

21. Liu, Y. *et al.* Structural insights into terminal arabinosylation of mycobacterial cell wall
arabinan. *Nature Communications 2025 16:1* **16**, 1–16 (2025).

22. Poulton, N. C. & Rock, J. M. Unraveling the mechanisms of intrinsic drug resistance in
Mycobacterium tuberculosis. *Front Cell Infect Microbiol* **12**, 997283 (2022).

23. Gupta, K. R. *et al.* An essential periplasmic protein coordinates lipid trafficking and is
required for asymmetric polar growth in mycobacteria. *Elife* **11**, (2022).

24. Daffé, M. The mycobacterial antigens 85 complex – from structure to function and
beyond. *Trends Microbiol* **8**, 438–440 (2000).

25. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500 (2024).

26. Degiacomi, G. *et al.* Essentiality of mmpL3 and impact of its silencing on Mycobacterium
tuberculosis gene expression. *Sci Rep* **7**, 1–8 (2017).

27. Fay, A. *et al.* Two accessory proteins govern MmpL3 mycolic acid transport in
mycobacteria. *mBio* **10**, (2019).

28. Varela, C. *et al.* MmpL Genes Are Associated with Mycolic Acid Metabolism in
Mycobacteria and Corynebacteria. *Chem Biol* **19**, 498–506 (2012).

29. Bansal-Mutalik, R. & Nikaido, H. Quantitative lipid composition of cell envelopes of
Corynebacterium glutamicum elucidated through reverse micelle extraction. *Proc Natl*
Acad Sci U S A **108**, 15360–15365 (2011).

30. Kamariza, M. *et al.* Rapid detection of Mycobacterium tuberculosis in sputum with a
solvatochromic trehalose probe. *Sci Transl Med* **10**, (2018).

31. Birch, H. L. *et al.* Biosynthesis of mycobacterial arabinogalactan: identification of a novel
α(1→3) arabinofuranosyltransferase. *Mol Microbiol* **69**, 1191 (2008).

32.   Gandotra, S., Schnappinger, D., Monteleone, M., Hillen, W. & Ehrt, S. In vivo gene
silencing identifies the Mycobacterium tuberculosis proteasome as essential for the
bacteria to persist in mice. *Nat Med* **13**, 1515–1520 (2007).

33.   Abrahams, K. A. & Besra, G. S. Mycobacterial cell wall biosynthesis: a multifaceted
antibiotic target. *Parasitology* **145**, 116–133 (2018).

34.   Lissner, R. *et al.* A new role for lipoproteins LpqZ and FecB in orchestrating mycobacterial
cell envelope biogenesis. *bioRxiv* 2025.06.19.660376 (2025)
doi:10.1101/2025.06.19.660376.

35.   Jankute, M. *et al.* Disruption of mycobacterial aftB results in complete loss of terminal β(1
→ 2) arabinofuranose residues of Lipoarabinomannan. *ACS Chem Biol* **12**, 183–190
(2017).

36.   Liu, Y. *et al.* Structural insights into terminal arabinosylation of mycobacterial cell wall
arabinan. *Nature Communications 2025 16:1* **16**, 1–16 (2025).

37.   Patel, O. *et al.* Crystal structure of the putative cell-wall lipoglycan biosynthesis protein
LmcA from Mycobacterium smegmatis. *Acta Crystallogr D Struct Biol* **78**, 494–508
(2022).

38.   Taschner, M., Mourão, A., Awasthi, M., Basquin, J. & Lorentzen, E. Structural basis of
outer dynein arm intraflagellar transport by the transport adaptor protein ODA16 and the
intraflagellar transport protein IFT46. *Journal of Biological Chemistry* **292**, 7462–7473
(2017).

39.   Kanao, T. *et al.* Reaction mechanism of tetrathionate hydrolysis based on the crystal
structure of tetrathionate hydrolase from Acidithiobacillus ferrooxidans. *Protein Science*
30, 328–338 (2021).

40.   Jahn, B. *et al.* Similar but Not the Same: First Kinetic and Structural Analyses of a
Methanol Dehydrogenase Containing a Europium Ion in the Active Site. *ChemBioChem*
19, 1147–1153 (2018).

41.   Yang, H. *et al.* Structure-Function Analysis of a Quinone-Dependent Dehydrogenase
Capable of Deoxynivalenol Detoxification. *J Agric Food Chem* **70**, 6764–6774 (2022).

42.   Jackson, M. *et al.* Inactivation of the antigen 85C gene profoundly affects the mycolate
content and alters the permeability of the Mycobacterium tuberculosis cell envelope. *Mol
Microbiol* **31**, 1573–1587 (1999).

43.   Sparks, I. L. *et al.* Lipoarabinomannan mediates localized cell wall integrity during division
in mycobacteria. *Nature Communications 2024 15:1* **15**, 1–17 (2024).

44.   Kilburn, J. O. & Takayama, K. Effects of ethambutol on accumulation and secretion of
trehalose mycolates and free mycolic acid in Mycobacterium smegmatis. *Antimicrob
Agents Chemother* **20**, 401–404 (1981).

45. Culviner, P. H., Guegler, C. K. & Laub, M. T. A simple, cost-effective, and robust method
for rrna depletion in rna-sequencing studies. *mBio* **11**, (2020).

46. Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster, cheaper and
better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res* **47**,
(2019).

47. Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

48. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

49. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).

50. Nandakumar, M., Prosser, G. A., De Carvalho, L. P. S. & Rhee, K. Metabolomics of
mycobacterium tuberculosis. *Methods in Molecular Biology* **1285**, 105–115 (2015).

51. Layre, E. *et al.* A Comparative Lipidomics Platform for Chemotaxonomic Analysis of
Mycobacterium tuberculosis. *Chem Biol* **18**, 1537–1549 (2011).

52. Wainberg, M. *et al.* A genome-wide atlas of co-essential modules assigns function to
uncharacterized genes. *Nat Genet* **53**, 638–649 (2021).

53. Lobstein, J. *et al.* SHuffle, a novel Escherichia coli protein expression strain capable of
correctly folding disulfide bonded proteins in its cytoplasm. *Microb Cell Fact* **11**, 1–16
(2012).

54. Kabsch, W. XDS. *Acta Crystallogr D Biol Crystallogr* **66**, 125–132 (2010).

55. McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658–674
(2007).

56. Terwilliger, T. C. *et al.* Iterative model building, structure refinement and density
modification with the PHENIX AutoBuild wizard. *Acta Crystallogr D Biol Crystallogr* **64**,
61–69 (2007).

57. Afonine, P. V. *et al.* Towards automated crystallographic structure refinement with
phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352–367 (2012).

58. Emsley, P. & Cowtan, K. Coot: Model-building tools for molecular graphics. *Acta
Crystallogr D Biol Crystallogr* **60**, 2126–2132 (2004).

59. Ganapathy, U. *et al.* Two enzymes with redundant fructose bisphosphatase activity
sustain gluconeogenesis and virulence in Mycobacterium tuberculosis. *Nature
Communications 2015 6:1* **6**, 1–12 (2015).

1086

1087

Main Figures

 Figure 1. FecB plays a non-critical role in iron homeostasis in Mtb. (A) Volcano plots of
 transcriptome changes in Δ*fecB* compared to the complemented strain (Δ*fecB*::*fecB*) during log-
 phase growth in regular media without detergent or with 0.05% Tween80; highlighted genes are
 also differentially expressed in Δ*fecB* vs WT. (B) Heatmap of iron-related, differentially expressed
 genes (DEG) in strains grown in detergent-free media. DEGs were selected by padj ≤ 0.05 and
 |log2FC| ≥ 0.5; common hits in Δ*fecB* vs Δ*fecB*::*fecB* and Δ*fecB* vs WT are shown. (C) Relative
 abundance of representative mycobactin and carboxymycobactin species detected as [M+Fe-

2H]+ from lipid extracts analyzed by liquid chromatography-mass spectrometry (LC-MS). (D)
Growth curves of iron pre-depleted strains in iron-chelated media with 160μM (high iron) or 1μM
(low iron) ammonium ferric citrate (AFC). Cultures in low iron were grown until day 17 (Passage
1) and diluted to starting OD=0.025 in high/low iron media (Passage 2). Δ*mbtK* (mycobactin
biosynthesis mutant) was used as a positive control that cannot grow in iron-limiting conditions.
(E) Differential expression by reverse transcription quantitative PCR (RT qPCR) of representative
iron-related genes (*mbtB*, *ppe37* and *irtA*) and *fecB* in strains grown in detergent-free media.
Statistical significance in (C) and (E) was determined by one-way ANOVA and Tukey post-hoc
test. Error bars: ±SD. N independent experiments = 2 with a total of five replicate cultures (A-B),
2 (C), 5 (D) and 1 (E) each performed with triplicate cultures.

1108

1109

Figure 2. FecB's roles in iron homeostasis and multidrug resistance are not functionally
related. MIC assays for vancomycin and rifampicin, percent growth calculated from no drug
control wells. (A) Δ*fecB* complemented with FecB variants with lower binding affinity to ferric
carboxymycobactin (Δ*fecB*::*fecB*-E339S_R240S and Δ*fecB*::*fecB*-E339S_Y242S) and control
strains (day 14). (B) CRISPRi KD of *mbtI* in WT/Δ*fecB* in the absence or presence of ATc inducer
(day 14). (C) WT, Δ*fecB* and Δ*fecB*::*fecB* strains in low iron (iron-chelated media with 1μM AFC)
or regular media (day 21). Error bars: ±SD. N independent experiments = 2 (A, C), 1 (B), each
performed with triplicate cultures.

1118

1119

Figure 3. FecB and Rv3035 interact in Mtb and share essentiality profiles. Protein interaction
profiling of (A) FecB and (B) Rv3035. FecB-FLAG and Rv3035-3xFLAG were immunoprecipitated
from live Mtb and interacting proteins were identified by mass spectrometry. WT lysate was used
a control for unspecific interactions. (C) Heatmap of co-essentiality Generalized Least Squares
(GLS) coefficients for selected gene pairs. The genes shown were selected based on three "query
genes": *fecB*, *rv3035*, and *aftB*. We included all gene pairs involving at least one of these queries
that met a significance threshold of FDR-adjusted p-value < 0.001. Each square's color represents
the GLS coefficient (i.e., the strength of co-essentiality), while size corresponds to statistical
significance ($-\log_{10}$ of the FDR-adjusted p-value). Diagonal elements represent self-comparisons
(each gene's $\log_2$ fold change profile against itself); while biologically uninformative, they are
displayed at full size as a visual benchmark for perfect correlation. Some gene-gene combinations
did not pass the significance threshold and are shown with smaller squares. (D) Venn diagram
showing genes that are co-essential with *fecB* (red circle), *rv3035* (yellow circle) and *aftB* (blue
circle), and their overlap. N independent experiments = 1 (A-B), each performed with triplicate
cultures.

1135

1136

Figure 4. Binding affinity of FecB and Rv3035 and X-ray crystal structures of Mth Rv3035
and the FecB-Rv3035 complex. (A) A 19-point ITC experiment was carried out by titrating 200μM
Mth Rv3035 into 20μM Mtb FecB. The resulting thermogram is shown (left graph), resulting in the
following calculated parameters: $K_D = 27.5 \pm 4.6$ nM; N(sites) = 1.03 $\pm$ 0.004 (right graph). (B)
Cartoon depiction of the structure of Mth Rv3035. The β-propeller fold is colored in light purple for
β-strands and white for loops, each blade was assigned a letter, and the additional secondary
structure elements are colored in pink. The disulfide bonds are shown in stick with yellow sulfur
atoms. (C) Cartoon depiction of the structure of the FecB-Rv3035 complex. Rv3035 is colored as
in (B). FecB secondary structure elements are colored in light green and loops in wheat. (D) A

 zoom in of the interface of the FecB-Rv3035 complex with interacting residues shown in stick,
1147 and hydrogen bond and salt bridge interactions are represented in black dashed lines.

1148

1149

Figure 5. FecB and Rv3035 function in the same pathway that mediates cell envelope
impermeability and antibiotic resistance in Mtb. (A) MIC assays for vancomycin, bedaquiline,
rifampicin, pretomanid, SQ109 and isoniazid, percent growth calculated from no drug control wells
(day 10). (B) BODIPY-vancomycin uptake after 30min of incubation, fluorescence intensity
normalized by OD_{580nm}. Statistical significance was determined by one-way ANOVA and Tukey
post-hoc test. (C) Calcein-AM uptake, fluorescence intensity normalized by OD_{580nm} at 60min. (D)
Spot Assay. (E) MIC assay for vancomycin and rifampicin, percent growth calculated from no drug
control wells. Δ*fecB* complemented with FecB triple mutant that does not bind Rv3035
(Δ*fecB*::*fecB*-H226A_D292A_D324A) and control strains (day 14), #1-2 indicate individual
colonies (biological replicates). Error bars: ±SD. N independent experiments = 2 (A-D), 1 (E),
each performed with triplicate cultures.

1161

1162

Figure 6. Enhanced vulnerability of *aftB* and accumulation of mycolates in Mtb Δ*fecB*. (A)
Growth curve of CRISPRi KD of *aftB* in WT/Δ*fecB* in the absence or presence of ATc inducer;
strains were grown in 384-well plate in standing conditions. Lipidomics analysis by LC-MS of total
lipids (B, E) and fractions from AOT treatment (C-D, F-G) identifying (B-D) trehalose
monomycolate (TMM) species and (E-G) free-mycolic acid (MA) species. (H) Trehalose
dimycolate (TDM) spot intensity quantified from fig. S12E by densitometry analysis using ImageJ
software. (I-J) Flow cytometry analysis of DMN-Tre labelled cells using 405nm laser and 525/50
channel. Representative histograms (left) and quantification (right) of mean fluorescence intensity
of DMN-Tre probe. Unstained (Unst) WT was used as a control for autofluorescence. Statistical
significance was determined by two-way (B-G) or one-way (H-J) ANOVA and Tukey post-hoc test.

1173 Error bars: ±SD. N independent experiments = 2 (A, I-J), 1 (B-H), each performed with triplicate
1174 cultures.

1175

1176

Figure 7. FecB is required for Mtb virulence. (A) MIC assays for vancomycin and rifampicin, in
the absence or presence of ATc to induce *fecB* expression in the FecB-TetON strain (day 14).
Percent growth calculated from no drug control wells. (B-C) Growth and persistence of FecB-
TetON in the (B) lungs and (C) spleens of mice receiving doxy or regular chow. Data are average
CFU counts from four mice per time point. Dashed black lines represent lower limit of detection.
Statistical significance was determined by unpaired T-tests for each time point, p-value is shown
above each pair analyzed. Error bars: ±SD. N independent experiments = 1 performed with
triplicate cultures (A), 1 (B-C).

1185