Running Title: *CCL22* upregulation in cancer cells via paracrine signaling

Title: Tumor cells upregulate *CCL22* in a STING-dependent manner in response to paracrine

factors released by STING-activated myeloid cells and type I interferons

Elmira M Lomashvili[1], Jihyun Kim[2], Lingwei Kong[1], Pamela R. Cook[1]*

[1]Department of Biomedical Sciences, Mercer University School of Medicine, Macon, GA, USA

[2]Department of Biomedical Sciences, Mercer University School of Medicine, Columbus, GA,

USA

*Corresponding: Pamela R. Cook cook_p@mercer.edu

ABSTRACT

Immunosuppressive elements within the tumor microenvironment include both regulatory T cells

(Tregs) and M2 macrophages. A well-described mechanism of Treg recruitment occurs via the

chemokine CCL22, and CCL22 has also recently been implicated in the polarization of tumor-

associated macrophages to the M2a subtype. Our lab and others have shown that CCL22 is

upregulated in cancer cells following activation of the STimulator of INterferon Genes (STING).

STING triggers immune responses against pathogenic and self-DNA mislocalized to the

cytoplasm, which can accumulate in cancer cells due to chromosomal instability, damaged

mitochondria, and increased expression of LINE-1 retrotransposons. STING activation has been

associated with both anti- and pro-tumor immune responses, and a potential mechanism of

STING-mediated immune evasion is through CCL22 upregulation. CCL22 was first

characterized in macrophages, and here we investigate the effects of STING activation on

CCL22 expression in macrophages and monocytes. We report that human macrophages and

monocytes are resistant to *CCL22* upregulation by STING, but that STING-activated

macrophages and monocytes release unidentified paracrine factor(s) that dramatically increase

CCL22 upregulation in cancer cells in a manner that remains STING-dependent, as evidenced by

the inability of STING knockout cells to upregulate *CCL22* in response to these factors. We

further found that exogenous type I interferons (IFNs), a major downstream product of STING

activation, also upregulate *CCL22* in cancer cells via a STING-dependent mechanism and that

exogenous IFNβ can directly activate STING.

INTRODUCTION

The STimulator of INterferon Genes (STING) is an effector protein that transmits signals

from various cytosolic DNA sensors to transcription factors, including interferon regulatory

factor 3 (IRF3), which leads to the expression of type I interferon (*IFN*) genes and downstream

immune responses (1-3). Healthy, non-mitotic eukaryotic cells confine DNA to the nucleus and

mitochondria; thus, the presence of cytosolic DNA often signals infection by microbes, making

STING an important sentinel in innate immune surveillance. Cancer cells are often characterized

by self-DNA mislocalized to the cytoplasm due to chromosomal instability, damaged

mitochondria, and reactivated LINE-1 retrotransposons (4, 5). LINE-1-mediated activation of

STING is thought to contribute to inflammation in multiple disease states, including age-related

neuropathologies and cancer (5-8). In the context of cancer, LINE-1 is highly expressed in many

epithelial tumors (9-11), and its activation of STING in those cells may have important clinical

implications on host anti-tumor immune responses. STING activation in cancer can boost anti-

tumor immunity due to the release of type I IFNs, and indeed STING agonists represent

promising immunotherapy agents (12, 13). However, the roles of STING in carcinogenesis,

tumor progression, and metastasis are complex, and in some cases STING activation has been

associated with impaired anti-tumor immunity (14-16).

Our lab and others have demonstrated a link between STING activation and increased

expression of a chemokine associated with immunosuppression in the tumor microenvironment,

the C-C-motif ligand 22 (*CCL22*) (17-19). Correlations between *CCL22* expression and adverse

outcomes across various cancer types have been reported in multiple studies (18, 20-24),

although CCL22 has also been associated with increased patient survival in some cancers,

particularly for colon and head and neck squamous cell carcinomas (19 and references therein).

CCL22 recruits regulatory T-cells (Tregs) via its interaction with the CCR4 receptor, which may

reduce tumor immune responses and promote immunosuppression in some cancers (25, 26).

Recently, CCL22 has also been reported to have a role in polarizing tumor-associated

macrophages to the M2a pro-tumor subtype (27).

CCL22, also known as macrophage-derived chemokine (MDC), was originally

discovered in macrophages (28, 29). Macrophages and other myeloid cells are major targets of

STING agonists in the tumor microenvironment (27). Here we sought to investigate the

relationship between STING activation and *CCL22* upregulation in macrophages and monocytes.

Although we found that STING activation did not increase *CCL22* expression in macrophages or

monocytes, we found that factor(s) released from these myeloid cells dramatically upregulated

CCL22 in cancer cells in a paracrine fashion. Moreover, we determined that this paracrine effect

remains dependent on STING in the cancer cells, based on results from our STING knockout cell

line. We previously showed that *CCL22* upregulation in response to STING activation occurs

predominantly via the STING-IRF3 axis (19), the same axis that triggers robust upregulation of

type I IFNs. We thus hypothesized that type I IFNs could be the relevant paracrine factor(s)

released by myeloid cells that upregulate *CCL22* in cancer cells. Although we found that

exogenous type I IFNs did indeed increase *CCL22* expression in cancer cells, the myeloid-

derived paracrine factors tended to have a larger effect, particularly in HeLa cells. Surprisingly,

we also found that upregulation of *CCL22* in response to type I IFNs remained dependent on

STING, and that IFNβ itself could activate STING.

MATERIALS AND METHODS

Cells and cell culture

Cell lines used in the study include MCF7 (human mammary gland adenocarcinoma), THP-1

(human monocytic leukemia), both purchased from ATCC, human HeLa cervical

adenocarcinoma cells, a kind gift from Dr. Anthony Furano from the National Institute of

Diabetes and Digestive and Kidney Diseases as described in (19), and HeLa$^{-/-TMEM173}$ STING

(TMEM173) knockout cells, engineered from the aforementioned wild type HeLa with the

CRISPR-Cas 9 system, also described in (19). All cells were cultured in DMEM high glucose

with GlutaMAX and pyruvate (Gibco, cat. 10569010) supplemented with 10% fetal bovine

serum (Gibco OneShot, non-heat-inactivated, cat. A3160401) and 1x antibiotic-antimycotic (100

units/mL penicillin, 100 ug/mL streptomycin, 0.25 ug/mL Amphotericin B; Gibco, cat.

15240096) in a humidified incubator at 37°C with 5% CO_2. THP-1 culture media was

additionally supplemented with 1x 2-mercaptoethanol (Gibco, cat. 21985023).

Cell culture treatments (STING agonists, IFNα, IFNβ, and MMHAR-2 block)

Four commercially available STING agonists were used in the study: a stabilized analog of the

cyclic dinucleotide canonical STING agonist 2'3'-cGAMP (2'3'-cGAM(PS)2(Rp/Sp)), hereafter

referred to as cGAM(PS)2 (Invivogen, cat. tlrl-nacga2srs), a stabilized cyclic adenine

monophosphate-inosine monophosphate (cAIM(PS)2 Difluor (Rp/Sp), hereinafter referred to as

cAIM(PS)2 (Invivogen, cat. tlrl-nacairs-2), E7766, a macrocycle-bridged diammonium salt

(Chemietek, cat. CT-E7766), and the diamidobenzimidazole diABZI (Selleckchem, cat.

S8796). All agonists except diABZI were water soluble and reconstituted using a companion

vial of sterile, endotoxin-free LAL water; diABZI was reconstituted in dimethyl sulfoxide

(DMSO). All agonists were diluted to a final concentration of 30 uM in complete cell culture

media and applied to cells as specified in the figures. Human recombinant IFNα stock (pbl assay

science, cat. 11101-1) was diluted to concentrations indicated in the figure legends in complete

cell culture media. Lyophilized recombinant human IFNb (PeproTech, cat. 300-02) was initially

reconstituted in sterile, endotoxin-free LAL water and further diluted into sterile 0.1% bovine

serum albumin (BSA) in 1x phosphate-buffered saline (PBS) to 100 ng/uL and further diluted to

concentrations indicated in the figure legends in complete cell culture media before adding to

cells. MMHAR-2 blockade was performed by pretreating cells with 8.3 µg/mL of the MMHAR-

2 antibody, then adding indicated treatments diluted in cell culture media, reducing the final

concentration of MMHAR-2 to 5 µg/mL. Cells were then incubated for approximately 30 hours

prior to harvesting for either Western blots or RTqPCR.

Macrophage differentiation and generation of conditioned media

THP-1 cells were differentiated into macrophages by plating cells in complete culture media

supplemented with 150 nM phorbol 12-myristate 13-acetate (PMA) and incubating for 96 hours,

with one media change performed after 48 hours to replenish PMA. PMA-containing media was

then removed, and cells were allowed to rest for 24 hours prior to treatment with cGAM(PS)2.

Conditioned media from both macrophages and undifferentiated monocytes was obtained by

treating cells with fresh complete media, with or without the addition of 30 uM cGAM(PS)2,

incubating for 24 hours, then gently removing media, centrifuging at 300 x g, and transferring to

cancer cells.

Cell imaging

The morphological changes during THP-1 differentiation were imaged using an EVOS FLoid

system.

RNA purification and RTqPCR

Cells were harvested using TRIzol reagent (Invitrogen, cat. 15596026) followed by column RNA

purification and on-column DNase digestion according to the TRIzol two-step protocol from the

Monarch total RNA miniprep kit (NEB, cat. T2010S). Purity and concentration of the RNA were

assessed using spectrophotometry. First-strand cDNA synthesis was performed using the

LunaScript RT SuperMix Kit (NEB, cat. E3010S) or LunaScript (-) RT SuperMix control with

added M-MuLV Reverse Transcriptase (NEB). qPCR was performed on a QuantStudio 3 real-

time PCR machine using TaqMan Fast Advanced Master Mix (Applied Biosystems, cat.

A44359). Singleplexed and duplexed 5' endonuclease assays were performed using the

following TaqMan Assays: *CCL22* (Hs00171080); IFN-β (Hs01077958_s1; beta actin

(Hs99999903); and 18s (Hs99999901). For duplexed assays, a beta-actin primer-limited assay

(IDT) was used. Samples for qPCR were assayed in technical triplicates, and data were analyzed

with the delta-delta Ct method; when multiple housekeeping genes were measured, their

combined geometric mean was used for analysis.

Cell lysis and Western blotting

Cells were lysed for immunoblotting in 3% SDS, 25 mM Tris-HCl pH 7.4, and 0.5 mM EDTA

supplemented with 3X HALT protease and phosphatase inhibitor cocktail (ThermoFisher, cat.

78440). Lysate homogenization was achieved with QIAshredder columns (Qiagen, cat. 79656),

after which protein concentration was determined using the BioRad DC Protein Assay (cat.

5000112). Equal amounts of lysates were resolved with SDS-PAGE. Transfers were performed

using a wet tank system overnight at 28 V or the BioRad Trans-Blot Turbo Transfer system.

Blots were imaged with an infrared Li-Cor Odyssey CLx Imager and processed using Image

Studio (Li-Cor). Positive controls for phospho-STING (S366) were prepared as commercially

described by transfecting THP-1 differentiated macrophages with dsDNA (pcDNA3.1 (+) puro),

(3.33 ug/mL) using Opti-MEM (Gibco, cat. 31-985-062) as the DNA diluent and TransIT-LT1

(Mirus, cat. MIR 2300) at a ratio of 1 ug DNA: 2 uL TransIT-LT1; cells were harvested 4 hours

after transfection with the lysis buffer described above.

Antibodies

Antibodies for Western blots included: anti-phospho(S366)-STING (Cell Signaling Technology,

cat. 19781S); anti-phospho(Y701)-Stat1 (Cell Signaling Technology, cat. 9167); anti-IFNAR2

(Cell Signaling Technology, cat. 53883); and anti-beta-tubulin (Abcam, cat. ab6046). The

antibody used for IFNAR blockade was anti-human interferon alpha/beta receptor chain 2, clone

MMHAR-2 (pbl assay science, cat. 21385-1).

Statistical analyses

Statistical analyses were performed using GraphPad Prism 9. Tests for significance are described

in the figure legends, and statistical significance was defined as p-value ≤ 0.05.

RESULTS

STING-activated macrophages and monocytes do not upregulate *CCL22*. Macrophages are a

major source of CCL22, reflected in the historical name of CCL22, macrophage-derived

chemokine (MDC). Since one of the targets of STING agonists are tumor-associated

macrophages, we sought to determine the effect of STING activation on *CCL22* in these cells.

Using the well-established THP-1 monocyte-to-macrophage differentiation model (Fig. 1A), we

incubated THP-1 monocytes with phorbol myristate acetate (PMA) for four days to establish

adherent, differentiated macrophages. Following differentiation, we treated the THP-1-derived

macrophages with the canonical human STING agonist cGAM(PS)2 for 24 hours.

Unexpectedly, we found that cGAM(PS)2 did not increase *CCL22* expression but rather reduced

it (Fig. 1B). To ensure that we had not missed an increase in *CCL22* at an earlier time point, we

monitored *CCL22* expression at 0, 6, 12, 24, and 48 hours, which revealed no significant increase

at any time point (Fig. S1). Rather, the time course in Fig. S1 shows a downward trend in *CCL22*

expression, although these decreases did not reach statistical significance in the context of a

Kruskal-Wallis multiple comparisons test, which generally has less power to identify differences

in means compared with the *t* test used to analyze data in Figure 1B. Indeed, when analyzing

only the 24-hour time-course data with an unpaired, two-tailed *t* test, the decrease in *CCL22* was

again significant (p = 0.0290). To determine whether failure to activate *CCL22* in macrophages

was due to a lack of STING activation, we confirmed that cGAM(PS)2 induced STING

phosphorylation in macrophages at both 6 and 24 hour time points (Fig. 1C). Consistent with

results from macrophages, undifferentiated THP-1 monocytes also failed to upregulate *CCL22* in

response to cGAM(PS)2 (Fig. 1D).

STING-activated macrophages and monocytes release factor(s) that upregulate *CCL22* in cancer cells. Macrophages have important roles in anti-tumor immunity but can also contribute to immunosuppressive environments (30). We thus sought to determine whether STING-activated macrophages might release factors that could act via paracrine signaling to alter *CCL22* expression in cancer cells. *CCL22* has been reported to play a role in breast (20, 22, 24) and cervical (18, 23) cancers, and we used human breast (MCF7) and cervical (HeLa) cancer cells in these experiments. These cell lines were also used in our previous study, where we showed that both CCL22 mRNA and protein were increased upon direct STING activation with cytosolic DNA or cGAM(PS)2 (19). To investigate potential paracrine activation of *CCL22* in these cells, we first treated macrophages with cGAM(PS)2 or a mock control for 24 hours, then removed the media and applied it to cancer cells for 24 hours. Both MCF7 (Fig. 2A) and HeLa cells (Fig. 2B) exposed to conditioned media from macrophages treated with cGAM(PS)2 exhibited a robust increase in *CCL22* compared to media from untreated macrophages or treatment with cGAM(PS)2 alone, although conditioned media from untreated macrophages also increased *CCL22* in HeLa cells, albeit to a lower extent. Monocytes also released factors that increased *CCL22* above levels induced by direct STING activation in MCF7 (Fig. 2C) and HeLa (Fig. 2D) cells.

Paracrine activation of *CCL22* in cancer cells activates STING and is dependent on STING. Our previous study showed that *CCL22* upregulation in response to cytosolic dsDNA and cGAM(PS)2 in HeLa cells was dependent on STING. However, in the current experiments, it was possible that paracrine-mediated activation of *CCL22* in these cells was independent of

STING. To investigate this, we first determined whether conditioned media from STING-

activated macrophages had any effect on STING activation in HeLa cells, as determined by

analysis of STING phosphorylation on serine 366 (31, 32). Figure 3A shows that STING in

HeLa cells was indeed activated by media obtained from STING-activated macrophages, and

this activation was greater than when HeLa cells were treated directly with cGAM(PS)2.

Importantly, in our previous study, we confirmed that free-floating DNA, which could possibly

have been present in the conditioned media, did not activate STING in HeLa cells (19). Notably,

the degree of STING activation by each treatment in Figure 3A generally corresponds to relative

levels of *CCL22* upregulation in Figure 2B, including the slight increases observed in response to

media from untreated macrophages. Similar results for STING phosphorylation were observed

when treating HeLa cells with conditioned media from monocytes (Fig. S2A), which again

parallels increases in *CCL22* in Figure 2D. Note that we did not examine STING

phosphorylation on S366 in MCF7 cells due to our previous finding that STING activation in

MCF7 cells with cytosolic dsDNA or pharmacological agonists occurred in the absence of

phosphorylation on S366 for unknown reasons, which are further discussed in that study (19). To

determine whether the observed STING activation was *required* for paracrine-mediated

upregulation of *CCL22*, we used a STING (gene name *TMEM173*) knockout cell line, HeLa$^{-/-}$

^{TMEM173}, which we previously established and characterized from the parental HeLa cells used

here (19). Treatment of the HeLa$^{-/-TMEM173}$ STING knockout cells with media from STING-

activated macrophages showed no statistically significant increase in *CCL22* expression,

indicating that intact STING signaling in the cancer cells was still required for upregulated

CCL22 (Fig. 3B). Similar results were obtained using media from STING-activated monocytes

(Fig. S2B).

Type I IFNs have modest effects on *CCL22* in HeLa cells that remain STING-dependent.

We previously found that STING-mediated *CCL22* upregulation in HeLa cells is mediated

predominantly via the STING-IRF3 axis, which triggers robust expression of type I IFNs (19).

We therefore hypothesized that the relevant paracrine factor(s) leading to the increase in *CCL22*

might be type I IFNs. Overall however, IFNα failed to elicit strong *CCL22* expression in either

cell type, and effects of IFNβ plateaued in dose-response curves, indicating that the relevant

paracrine factor was unlikely a higher concentration of type I IFNs.

In HeLa cells, IFNβ did not significantly upregulate *CCL22* at any dose at 24 hours (Fig.

S3A), and subsequent time-course analyses indicated the maximum response to both IFNα and

IFNβ occurred at 48 hours. However, even at 48 hours, IFNα failed to significantly upregulate

CCL22 in HeLa cells except at the highest dose (100 ng/mL), and then only to an average of 15

fold (Fig. 4A). IFNβ increased *CCL22* at the lowest dose (1 ng/mL) in HeLa cells, but only to

approximately 10 fold, and higher doses of IFNβ failed to elicit any additional increase (Fig. 4B).

MCF7 cells upregulated CCL22 in response to type I IFNs more rapidly than HeLa cells,

showing higher levels at 24 hours compared to 48 hours (Fig. S3B). At the highest dose, IFNα

increased *CCL22* by an average of only 6.7 fold in 24 hours (Fig. 4C). Paralleling the relative

effects of IFNα and IFNβ in HeLa cells, IFNβ also increased *CCL22* in MCF7 cells more than

IFNα, with an average of 57 fold change in 24 hours at the highest dose (Fig. 4D). However, the

effect of IFNβ was still below that observed from paracrine activation of MCF7 cells by

conditioned media from STING-activated monocytes (76.5 fold, Fig. 2C), though it was slightly

above levels induced by conditioned media from STING-activated macrophages (46.3 fold, Fig.

2A).

We next tested the effect of exogenous IFNα and IFNβ on *CCL22* upregulation in the

context of STING knockout cells, HeLa$^{-/-TMEM173}$. Neither IFNα (Fig. 4E) nor IFNβ (Fig. 4F)

significantly affected *CCL22* expression in STING knockout cells. These data were unexpected

and indicate that the effect of exogenous IFNα and IFNβ on *CCL22* is dependent on STING.

Although STING is well-known to trigger expression of type I *IFNs* downstream of IRF3, to our

knowledge, a requirement of STING for the effects of exogenous type I IFNs has not been

reported. To determine whether STING was activated by type I IFN signaling, we treated wild-

type HeLa cells with IFNβ and then performed a Western blot to assess STING phosphorylation,

and indeed found that exogenous IFNβ resulted in activated STING (Fig. 4G). We did not test

the effect of IFNα due to the comparatively minimal effect we observed from IFNα on *CCL22*

relative to IFNβ. To rule out the possibility that type I IFNs failed to upregulate *CCL22* in the

STING KO cells due to an inadvertent disruption of the interferon receptor (IFNAR) and

downstream Stat1 signaling, we confirmed that our STING KO cells exhibited the same degree

of Stat1 phosphorylation in response to IFNα and IFNβ as wild-type cells (Fig. 4H). As

expected, we found that cGAM(PS)2, which activates STING and STING-mediated type I IFNs,

also slightly induced Stat1 in the wild-type cells, but this activation was blocked in the STING

KO cells, reflecting the lack of STING-mediated upregulation of type I IFNs available to bind

and activate the IFNAR-Stat1 pathway in those cells (Fig. 4H). Confirming intact pStat1

signaling in the STING KO cells is consistent with our hypothesis that *CCL22* upregulation in

response to exogenous type I IFN is mediated through STING, not Stat1. Given that Stat1

phosphorylation was equivalently induced in wild-type and STING knockout cells by type I

IFNs, it was also unlikely that IFNAR expression would have been reduced in the STING

knockout cells, but to confirm this, we also show equivalent levels of IFNAR in both cell lines

(S3C).

The inability of type I IFNs to activate *CCL22* expression in our STING KO cells,

combined with confirmation of intact IFNAR expression and signaling in STING KO cells,

indicated that *CCL22* upregulation in wild-type cells by type I IFNs occurs via STING. To

further examine the role of IFNAR signaling on *CCL22* activation in the context of type I IFNs,

direct STING activation with cGAM(PS)2, or paracrine activation using conditioned media, we

performed an IFNAR blockade using a well-characterized neutralizing monoclonal antibody

from PBL Assay Science, anti-human IFN-alpha/beta receptor chain 2, clone MMHAR-2. To

first confirm that MMHAR-2 exerted a neutralizing effect on IFNAR signaling, we analyzed the

ability of this antibody to reduce Stat1 phosphorylation in response to IFNα and IFNβ, as well as

cGAM(PS)2. MMHAR-2 somewhat reduced levels of phosphorylated Stat1 in response to

IFNα, but not IFNβ (Fig. S3D). This difference might be attributable to the reported higher

affinity of IFNβ for IFNARs compared to IFNα (33, 34). Given that cGAM(PS)2 upregulates

type I IFNs through its activation of STING, we also observed as expected that the MMHAR-2

blockade reduced Stat1 phosphorylation following treatment with cGAM(PS)2 (Fig. S3D). We

also confirmed that MMHAR-2 exerted no activating effect on Stat1 (Fig. S3E). After

establishing the neutralizing potential of MMHAR-2 in our system, we sought to determine the

contribution of IFNAR signaling to *CCL22* upregulation in response to type I IFNs,

cGAM(PS)2, and paracrine factors from STING-activated macrophages. Figure 4I shows the

relative degree of inhibition induced by MMHAR-2 for each treatment. Although not statistically

significant, MMHAR-2 produced an observable downward trend in *CCL22* expression for all

treatments except cGAM(PS)2. The neutralizing effect of MMHAR-2 was most noticeable in

cells treated with IFNα, where the p-value is approaching significance at 0.0659, and these

results parallel those observed for MMHAR-2 inhibition of IFNα-mediated Stat1

phosphorylation in Figure S3D. Also in Figure S3D, the very slight increase in Stat1

phosphorylation from cGAM(PS)2 treatment, as well as its return to baseline with MMHAR-2

pretreatment, most likely reflects the activity of cGAM(PS)2-induced type I IFNs downstream of

STING. That the IFNAR block had the least effect on *CCL22* upregulation by cGAM(PS)2 is

consistent with our other data indicating that *CCL22* upregulation downstream of STING in

HeLa cells is not mediated primarily or solely by type I IFNs.

STING agonists differentially upregulate *CCL22* and *IFNβ*

We were curious, given the differential effects of type I IFNs and cGAM(PS)2 on *CCL22,*

whether different STING agonists would have distinctive effects on *CCL22* compared to *IFNβ*.

We tested a panel of four STING agonists that included two cyclic and two and non-cyclic

dinucleotides. The cyclic dinucleotides included cGAM(PS)2 and the stabilized analog of

adenosine-inosine monophosphate (cAIMP(PS)2) (35, 36). The two non-cyclic dinucleotides

were diamidobenzimidazole (diABZI) (37-39) and E7766, a locked, macrocycle-bridged agonist

(38, 40, 41). In HeLa cells, all agonists increased *CCL22* but had no significant effect on *IFNβ*

expression (Fig. 5A-5D). Three of the STING agonists increased *IFNβ* in MCF7 cells to

statistically significant levels, but *CCL22* was increased in response to all agonists, including

diABZI, the agonist that did not statistically increase *IFNβ* (Fig. 5E-5H). These results are

consistent with a mechanism of *CCL22* induction via STING that does not rely exclusively on

type I IFNs.

DISCUSSION

STING activation in tumors can promote anti-tumor immune responses, and STING

agonists represent an emerging class of immunotherapy agents (13, 42, 43). However, multiple

reports also indicate that STING can induce immunosuppressive tumor environments (reviewed

in 44). Several pro-tumor mechanisms of STING have been described, including upregulation of

programmed death ligand (44) and indoleamine 2,3-dioxygenase (45), radiation-induced

recruitment of myeloid suppressor cells (46), and DMBA exposure (47). One example of

immune equilibrium mediated by STING is the polarization of macrophages, where STING has

been reported to induce both anti-tumor M1 as well as pro-tumor M2 subtypes (44, 48, 49).

Our lab and others have found that STING activation can also increase expression of the

chemokine CCL22 in cancer cells (17-19). CCL22 is implicated in tumor immune evasion

through its recruitment of regulatory T cells (25, 26) and, more recently, by promoting

polarization of the M2 phenotype in tumor-associated macrophages (27). We previously showed

that STING activation by cytosolic dsDNA and the STING agonist cGAM(PS)2 can upregulate

CCL22 expression in epithelial cancer cells via the STING-IRF3 axis (19). In the present study,

we show that macrophages and monocytes do not themselves upregulate *CCL22* in response to

STING activation but rather release unidentified factor(s) that in turn increase *CCL22* expression

in a paracrine fashion in cancer cells. STING-activated myeloid cells that can induce neighboring

cancer cells to upregulate CCL22 via paracrine signaling may thus contribute to

immunosuppressive tumor environments by favoring the recruitment of Tregs and polarization of

M2a macrophages.

Based on our prior finding that *CCL22* upregulation was downstream of the STING-IRF3

axis, we hypothesized that type I IFNs could be the relevant paracrine factor(s). However,

treatment of cancer cells with exogenous IFNα or IFNβ did not, in most cases, achieve the same

level of *CCL22* upregulation as conditioned media from STING-activated myeloid cells. For

example, in HeLa cells, treatment with type I IFNs increased *CCL22* by 10-20 fold, while

treatment with conditioned media from STING-activated macrophages increased *CCL22* by up

to 8,000 fold. In MCF7 cells, conditioned media from monocytes, but not macrophages, also

increased *CCL22* above levels induced by type I IFNs, although the effect was more modest.

354   Surprisingly, we found that *CCL22* upregulation by exogenous type I IFNs, as well as

paracrine factors, remained dependent on STING, as evidenced by the lack of *CCL22*

upregulation in STING knockout cells. Although STING-dependent expression of type I IFNs is

well-described, a requirement for STING for a functional effect of exogenous type I IFNs has, to

our knowledge, not been previously described. One possible explanation could be our finding

that IFNβ itself induces activating phosphorylation of STING, which also has not been

previously reported. However, other studies have indicated connections between type I IFNs and

the transcriptional regulation of *cGAS* and the gene that encodes STING, *TMEM173* (50, 51).

These findings raise important questions about feedback mechanisms between type I IFNs and

STING. Regarding such feedback, LINE-1 retrotransposons, which are reported to activate

cGAS-STING (5), were recently shown to be upregulated in both monocytes and neutrophils

exposed to IFNα (52). Many solid tumors upregulate LINE-1 expression (9, 53), making LINE-1

a possible source of CCL22 in cancer cells via cGAS-STING activation. The potential for

additional LINE-1 upregulation in myeloid cells by type I IFNs may further amplify STING

signaling in those cells, increasing the release of paracrine factors affecting *CCL22*.

369   More studies are needed to identify the paracrine factor(s) released from STING-

activated myeloid cells that induce *CCL22*. Multiple reports show that a variety of factors can

influence CCL22 regulation, including but not limited to IL-1, IL-4, IL-5, IFN-gamma (IFN-γ),

and tumor necrosis factor alpha (TNF-α), with some factors having different effects on CCL22

based on cell type (e.g. 28, 29, 54-59). A study in 2011 by Faget et al. showed that peripheral

blood mononuclear cells promoted increased secretion of CCL22 from breast cancer cells and

identified roles for monocyte-derived TNF-α and IL-1β, as well as exogenous IFN-γ (60).

Whether any of these factors, or combinations thereof, are among the paracrine factors released

by STING-activated myeloid cells in our study remains to be determined. However, one

important potential candidate, especially given the STING-dependency we observed for *CCL22*

upregulation in cancer cells, could be cGAMP itself. Intercellular transfer of cGAMP has been

shown to occur via several mechanisms, including cell-cell contacts such as phagocytosis and

gap junctions, as well as the release and uptake of cGAMP from the extracellular space through

various transport channels in the cell membrane (4, 61-65). Although these studies have largely

focused on cGAMP released by tumor cells that is taken up by myeloid cells, the converse may

also be true.

The findings presented here offer new insights into possible mechanisms of STING-

mediated immunosuppression in cancer. Future studies will be needed to determine clinical

associations between STING activation and increased intratumoral CCL22, and whether

increased levels of CCL22 in this context correlate with immunosuppression. In addition,

identification of the paracrine factor(s) released from STING-activated myeloid cells that can

increase CCL22 in cancer cells could reveal therapeutic strategies targeting Treg recruitment and

M2 polarization in solid tumors.

REFERENCES

1. Zhang, B., P. Xu, and A. Ablasser. 2025. Regulation of the cGAS-STING Pathway. *Annu Rev Immunol* 43: 667-692.

2. Motwani, M., S. Pesiridis, and K. A. Fitzgerald. 2019. DNA sensing by the cGAS-STING pathway in health and disease. *Nat Rev Genet* 20: 657-674.

3. Hopfner, K. P., and V. Hornung. 2020. Molecular mechanisms and cellular functions of cGAS-STING signalling. *Nat Rev Mol Cell Biol* 21: 501-521.

4. Dvorkin, S., S. Cambier, H. E. Volkman, and D. B. Stetson. 2024. New frontiers in the cGAS-STING intracellular DNA-sensing pathway. *Immunity* 57: 718-730.

5. Mathavarajah, S., and G. Dellaire. 2024. LINE-1: an emerging initiator of cGAS-STING signalling and inflammation that is dysregulated in disease. *Biochem Cell Biol* 102: 38-46.

6. Gamdzyk, M., D. M. Doycheva, C. Araujo, U. Ocak, Y. Luo, J. Tang, and J. H. Zhang. 2020. cGAS/STING Pathway Activation Contributes to Delayed Neurodegeneration in Neonatal Hypoxia-Ischemia Rat Model: Possible Involvement of LINE-1. *Mol Neurobiol* 57: 2600-2619.

7. Gorbunova, V., A. Seluanov, P. Mita, W. McKerrow, D. Fenyo, J. D. Boeke, S. B. Linker, F. H. Gage, J. A. Kreiling, A. P. Petrashen, T. A. Woodham, J. R. Taylor, S. L. Helfand, and J. M. Sedivy. 2021. The role of retrotransposable elements in ageing and age-associated diseases. *Nature* 596: 43-53.

8. De Cecco, M., T. Ito, A. P. Petrashen, A. E. Elias, N. J. Skvir, S. W. Criscione, A. Caligiana, G. Brocculi, E. M. Adney, J. D. Boeke, O. Le, C. Beauséjour, J. Ambati, K. Ambati, M. Simon, A. Seluanov, V. Gorbunova, P. E. Slagboom, S. L. Helfand, N. Neretti, and J. M. Sedivy. 2019. L1 drives IFN in senescent cells and promotes age-associated inflammation. *Nature* 566: 73-78.

9. Rodic, N., R. Sharma, R. Sharma, J. Zampella, L. Dai, M. S. Taylor, R. H. Hruban, C. A. Iacobuzio-Donahue, A. Maitra, M. S. Torbenson, M. Goggins, M. Shih Ie, A. S. Duffield, E. A. Montgomery, E. Gabrielson, G. J. Netto, T. L. Lotan, A. M. De Marzo, W. Westra, Z. A. Binder, B. A. Orr, G. L. Gallia, C. G. Eberhart, J. D. Boeke, C. R. Harris, and K. H. Burns. 2014. Long interspersed element-1 protein expression is a hallmark of many human cancers. *Am J Pathol* 184: 1280-1286.

10. Burns, K. H. 2017. Transposable elements in cancer. *Nat Rev Cancer* 17: 415-424.

11. McKerrow, W., N. Doudican, N. Frazzette, S. A. Evans, A. Rocha, L. Kagermazova, J. M. Sedivy, N. Neretti, J. Carucci, J. D. Boeke, and D. Fenyö. 2022. LINE-1 Retrotransposon expression in cancerous, epithelial and neuronal cells revealed by 5' single-cell RNA-Seq. *bioRxiv*: 2021.2001.2019.427347.

12. Gan, Y., X. Li, S. Han, Q. Liang, X. Ma, P. Rong, W. Wang, and W. Li. 2021. The cGAS/STING Pathway: A Novel Target for Cancer Therapy. *Front Immunol* 12: 795401.

13. Xu, Y., and Y. Xiong. 2024. Targeting STING signaling for the optimal cancer immunotherapy. *Front Immunol* 15: 1482738.

14.   Zhou, Z., S. Huang, F. Fan, Y. Xu, C. Moore, S. Li, and C. Han. 2024. The multiple faces
of cGAS-STING in antitumor immunity: prospects and challenges. *Med Rev (2021)* 4:
173-191.

15.   Samson, N., and A. Ablasser. 2022. The cGAS-STING pathway and cancer. *Nat Cancer*
3: 1452-1463.

16.   Yue, B., W. Gao, J. F. Lovell, H. Jin, and J. Huang. 2025. The cGAS-STING pathway in
cancer immunity: dual roles, therapeutic strategies, and clinical challenges. *Essays*
Biochem 69.

17.   Liang, D., H. Xiao-Feng, D. Guan-Jun, H. Er-Ling, C. Sheng, W. Ting-Ting, H. Qin-
Gang, N. Yan-Hong, and H. Ya-Yi. 2015. Activated STING enhances Tregs infiltration
in the HPV-related carcinogenesis of tongue squamous cells via the c-jun/CCL22 signal.
Biochim Biophys Acta 1852: 2494-2503.

18.   Ni, H., H. Zhang, L. Li, H. Huang, H. Guo, L. Zhang, C. Li, J. X. Xu, C. P. Nie, K. Li, X.
Zhang, X. Xia, and J. Li. 2022. T cell-intrinsic STING signaling promotes regulatory T
cell induction and immunosuppression by upregulating FOXP3 transcription in cervical
cancer. *J Immunother Cancer* 10.

19.   Kim, J., J. V. Pena, H. P. McQueen, L. Kong, D. Michael, E. M. Lomashvili, and P. R.
Cook. 2024. Downstream STING pathways IRF3 and NF-kappaB differentially regulate
CCL22 in response to cytosolic dsDNA. *Cancer Gene Ther* 31: 28-42.

20.   Gobert, M., I. Treilleux, N. Bendriss-Vermare, T. Bachelot, S. Goddard-Leon, V. Arfi, C.
Biota, A. C. Doffin, I. Durand, D. Olive, S. Perez, N. Pasqual, C. Faure, I. Ray-Coquard,
A. Puisieux, C. Caux, J. Y. Blay, and C. Ménétrier-Caux. 2009. Regulatory T cells
recruited through CCL22/CCR4 are selectively activated in lymphoid infiltrates
surrounding primary breast tumors and lead to an adverse clinical outcome. *Cancer Res*
69: 2000-2009.

21.   Lecoq, I., K. L. Kopp, M. Chapellier, P. Mantas, E. Martinenaite, M. Perez-Penco, L.
Ronn Olsen, M. B. Zocca, A. Wakatsuki Pedersen, and M. H. Andersen. 2022. CCL22-
based peptide vaccines induce anti-cancer immunity by modulating tumor
microenvironment. *Oncoimmunology* 11: 2115655.

22.   Li, Y. Q., F. F. Liu, X. M. Zhang, X. J. Guo, M. J. Ren, and L. Fu. 2013. Tumor secretion
of CCL22 activates intratumoral Treg infiltration and is independent prognostic predictor
of breast cancer. *PLoS One* 8: e76379.

23.   Wang, Q., E. Schmoeckel, B. P. Kost, C. Kuhn, A. Vattai, T. Vilsmaier, S. Mahner, D.
Mayr, U. Jeschke, and H. H. Heidegger. 2019. Higher CCL22+ Cell Infiltration is
Associated with Poor Prognosis in Cervical Cancer Patients. *Cancers (Basel)* 11.

24.   Wiedemann, G. M., M. M. Knott, V. K. Vetter, M. Rapp, S. Haubner, J. Fesseler, B.
Kuhnemuth, P. Layritz, R. Thaler, S. Kruger, S. Ormanns, D. Mayr, S. Endres, and D.
Anz. 2016. Cancer cell-derived IL-1alpha induces CCL22 and the recruitment of
regulatory T cells. *Oncoimmunology* 5: e1175794.

25.   Anz, D., M. Rapp, S. Eiber, V. H. Koelzer, R. Thaler, S. Haubner, M. Knott, S. Nagel, M.
Golic, G. M. Wiedemann, F. Bauernfeind, C. Wurzenberger, V. Hornung, C. Scholz, D.
Mayr, S. Rothenfusser, S. Endres, and C. Bourquin. 2015. Suppression of intratumoral

CCL22 by type i interferon inhibits migration of regulatory T cells and blocks cancer
progression. *Cancer Res* 75: 4483-4493.

26. Sugiyama, D., H. Nishikawa, Y. Maeda, M. Nishioka, A. Tanemura, I. Katayama, S.
Ezoe, Y. Kanakura, E. Sato, Y. Fukumori, J. Karbach, E. Jager, and S. Sakaguchi. 2013.
Anti-CCR4 mAb selectively depletes effector-type FoxP3+CD4+ regulatory T cells,
evoking antitumor immune responses in humans. *Proc Natl Acad Sci U S A* 110: 17945-
17950.

27. Wang, Q., K. Sudan, E. Schmoeckel, B. P. Kost, C. Kuhn, A. Vattai, T. Vilsmaier, S.
Mahner, U. Jeschke, and H. H. Heidegger. 2022. CCL22-Polarized TAMs to M2a
Macrophages in Cervical Cancer In Vitro Model. *Cells* 11.

28. Mantovani, A., P. A. Gray, J. Van Damme, and S. Sozzani. 2000. Macrophage-derived
chemokine (MDC). *J Leukoc Biol* 68: 400-404.

29. Yamashita, U., and E. Kuroda. 2002. Regulation of macrophage-derived chemokine
(MDC, CCL22) production. *Crit Rev Immunol* 22: 105-114.

30. Jung, M., and B. Bonavida. 2024. Immune Evasion in Cancer Is Regulated by Tumor-
Asociated Macrophages (TAMs): Targeting TAMs. *Crit Rev Oncog* 29: 1-17.

31. Liu, S., X. Cai, J. Wu, Q. Cong, X. Chen, T. Li, F. Du, J. Ren, Y. T. Wu, N. V. Grishin,
and Z. J. Chen. 2015. Phosphorylation of innate immune adaptor proteins MAVS,
STING, and TRIF induces IRF3 activation. *Science* 347: aaa2630.

32. Zhang, C., G. Shang, X. Gui, X. Zhang, X. C. Bai, and Z. J. Chen. 2019. Structural basis
of STING binding with and phosphorylation by TBK1. *Nature* 567: 394-398.

33. Lavoie, T. B., E. Kalie, S. Crisafulli-Cabatu, R. Abramovich, G. DiGioia, K. Moolchan,
S. Pestka, and G. Schreiber. 2011. Binding and activity of all human alpha interferon
subtypes. *Cytokine* 56: 282-289.

34. Schreiber, G. 2017. The molecular basis for differential type I interferon signaling. *The
Journal of Biological Chemistry* 292: 7285-7294.

35. Andersen, A. H. F., R. Olesen, K. L. Jonsson, J. F. Hojen, C. Krapp, K. Mack, M. K.
Thomsen, L. Ostergaard, M. Tolstrup, F. Dagnaes-Hansen, M. R. Jakobsen, and P. W.
Denton. 2019. cAIMP administration in humanized mice induces a chimerization-level-
dependent STING response. *Immunology* 157: 163-172.

36. Skouboe, M. K., A. Knudsen, L. S. Reinert, C. Boularan, T. Lioux, E. Perouzel, M. K.
Thomsen, and S. R. Paludan. 2018. STING agonists enable antiviral cross-talk between
human cells and confer protection against genital herpes in mice. *PLoS Pathog* 14:
e1006976.

37. Guerini, D. 2022. STING Agonists/Antagonists: Their Potential as Therapeutics and
Future Developments. *Cells* 11.

38. Motedayen Aval, L., J. E. Pease, R. Sharma, and D. J. Pinato. 2020. Challenges and
Opportunities in the Clinical Development of STING Agonists for Cancer
Immunotherapy. *J Clin Med* 9.

39. Su, T., Y. Zhang, K. Valerie, X. Y. Wang, S. Lin, and G. Zhu. 2019. STING activation in
cancer immunotherapy. *Theranostics* 9: 7759-7771.

40. Amouzegar, A., M. Chelvanambi, J. N. Filderman, W. J. Storkus, and J. J. Luke. 2021.
STING Agonists as Cancer Therapeutics. *Cancers (Basel)* 13.

41. Le Naour, J., L. Zitvogel, L. Galluzzi, E. Vacchelli, and G. Kroemer. 2020. Trial watch:
STING agonists in cancer therapy. *Oncoimmunology* 9: 1777624.

42. Wang, Y., Y. Zhu, Y. Cao, Y. Li, Z. Zhang, J. S. Fleishman, S. Cheng, J. Chen, and M.
Ding. 2025. The activation of cGAS-STING pathway offers novel therapeutic
opportunities in cancers. *Front Immunol* 16: 1579832.

43. Hines, J. B., A. J. Kacew, and R. F. Sweis. 2023. The Development of STING Agonists
and Emerging Results as a Cancer Immunotherapy. *Curr Oncol Rep* 25: 189-199.

44. Zhang, Y., Y. Wang, P. Mu, X. Zhu, and Y. Dong. 2024. Bidirectional regulation of the
cGAS-STING pathway in the immunosuppressive tumor microenvironment and its
association with immunotherapy. *Front Immunol* 15: 1470468.

45. Lemos, H., E. Mohamed, L. Huang, R. Ou, G. Pacholczyk, A. S. Arbab, D. Munn, and A.
L. Mellor. 2016. STING Promotes the Growth of Tumors Characterized by Low
Antigenicity via IDO Activation. *Cancer Res* 76: 2076-2081.

46. Liang, H., L. Deng, Y. Hou, X. Meng, X. Huang, E. Rao, W. Zheng, H. Mauceri, M.
Mack, M. Xu, Y. X. Fu, and R. R. Weichselbaum. 2017. Host STING-dependent MDSC
mobilization drives extrinsic radiation resistance. *Nat Commun* 8: 1736.

47. Ahn, J., T. Xia, H. Konno, K. Konno, P. Ruiz, and G. N. Barber. 2014. Inflammation-
driven carcinogenesis is mediated through STING. *Nat Commun* 5: 5166.

48. Ma, R., T. Ji, D. Chen, W. Dong, H. Zhang, X. Yin, J. Ma, X. Liang, Y. Zhang, G. Shen,
X. Qin, and B. Huang. 2016. Tumor cell-derived microparticles polarize M2 tumor-
associated macrophages for tumor progression. *Oncoimmunology* 5: e1118599.

49. Nakajima, S., K. Mimura, A. Kaneta, K. Saito, M. Katagata, H. Okayama, M. Saito, Z.
Saze, Y. Watanabe, H. Hanayama, T. Tada, W. Sakamoto, T. Momma, H. Ohira, and K.
Kono. 2023. Radiation-Induced Remodeling of the Tumor Microenvironment Through
Tumor Cell-Intrinsic Expression of cGAS-STING in Esophageal Squamous Cell
Carcinoma. *Int J Radiat Oncol Biol Phys* 115: 957-971.

50. Ma, F., B. Li, S. Y. Liu, S. S. Iyer, Y. Yu, A. Wu, and G. Cheng. 2015. Positive feedback
regulation of type I IFN production by the IFN-inducible DNA sensor cGAS. *J Immunol*
194: 1545-1554.

51. Ma, F., B. Li, Y. Yu, S. S. Iyer, M. Sun, and G. Cheng. 2015. Positive feedback
regulation of type I interferon by the interferon-stimulated gene STING. *EMBO Rep* 16:
202-212.

52. Le, E., F. Moadab, X. Wang, R. Najjar, S. J. Van den Bogaerde, A. Bays, J. LaCava, and
T. Mustelin. 2025. Interferons and Cytokines Induce Transcriptional Activation of the
Long-Interspersed Element-1 in Myeloid Cells from Autoimmune Patients. *Eur J*
Immunol 55: e2451351.

53.   Ardeljan, D., M. S. Taylor, D. T. Ting, and K. H. Burns. 2017. The Human Long
554   Interspersed Element-1 Retrotransposon: An Emerging Biomarker of Neoplasia. *Clin*
555   *Chem* 63: 816-822.

556   54.   Faffe, D. S., T. Whitehead, P. E. Moore, S. Baraldo, L. Flynt, K. Bourgeois, R. A.
557   Panettieri, and S. A. Shore. 2003. IL-13 and IL-4 promote TARC release in human
558   airway smooth muscle cells: role of IL-4 receptor genotype. *Am J Physiol Lung Cell Mol*
559   *Physiol* 285: L907-914.

560   55.   Fujii-Maeda, S., K. Kajiwara, K. Ikizawa, M. Shinazawa, B. Yu, T. Koga, M. Furue, and
561   Y. Yanagihara. 2004. Reciprocal regulation of thymus and activation-regulated
562   chemokine/macrophage-derived chemokine production by interleukin (IL)-4/IL-13 and
563   interferon-gamma in HaCaT keratinocytes is mediated by alternations in E-cadherin
564   distribution. *J Invest Dermatol* 122: 20-28.

565   56.   Fukuda, K., Y. Fujitsu, K. Seki, N. Kumagai, and T. Nishida. 2003. Differential
566   expression of thymus- and activation-regulated chemokine (CCL17) and macrophage-
567   derived chemokine (CCL22) by human fibroblasts from cornea, skin, and lung. *J Allergy*
568   *Clin Immunol* 111: 520-526.

569   57.   Horikawa, T., T. Nakayama, I. Hikita, H. Yamada, R. Fujisawa, T. Bito, S. Harada, A.
570   Fukunaga, D. Chantry, P. W. Gray, A. Morita, R. Suzuki, T. Tezuka, M. Ichihashi, and
571   O. Yoshie. 2002. IFN-gamma-inducible expression of thymus and activation-regulated
572   chemokine/CCL17 and macrophage-derived chemokine/CCL22 in epidermal
573   keratinocytes and their roles in atopic dermatitis. *Int Immunol* 14: 767-773.

574   58.   Bonecchi, R., S. Sozzani, J. T. Stine, W. Luini, G. D'Amico, P. Allavena, D. Chantry, and
575   A. Mantovani. 1998. Divergent effects of interleukin-4 and interferon-gamma on
576   macrophage-derived chemokine production: an amplification circuit of polarized T helper
577   2 responses. *Blood* 92: 2668-2671.

578   59.   Xiao, T., S. Kagami, H. Saeki, M. Sugaya, T. Kakinuma, H. Fujita, S. Yano, H. Mitsui,
579   H. Torii, M. Komine, A. Asahina, K. Nakamura, and K. Tamaki. 2003. Both IL-4 and IL-
580   13 inhibit the TNF-alpha and IFN-gamma enhanced MDC production in a human
581   keratinocyte cell line, HaCaT cells. *J Dermatol Sci* 31: 111-117.

582   60.   Faget, J., C. Biota, T. Bachelot, M. Gobert, I. Treilleux, N. Goutagny, I. Durand, S. Léon-
583   Goddard, J. Y. Blay, C. Caux, and C. Ménétrier-Caux. 2011. Early detection of tumor
584   cells by innate immune cells leads to T(reg) recruitment through CCL22 production by
585   tumor cells. *Cancer Res* 71: 6143-6152.

586   61.   Blest, H. T. W., and L. Chauveau. 2023. cGAMP the travelling messenger. *Front*
587   *Immunol* 14: 1150705.

588   62.   Carozza, J. A., V. Bohnert, K. C. Nguyen, G. Skariah, K. E. Shaw, J. A. Brown, M.
589   Rafat, R. von Eyben, E. E. Graves, J. S. Glenn, M. Smith, and L. Li. 2020. Extracellular
590   cGAMP is a cancer cell-produced immunotransmitter involved in radiation-induced anti-
591   cancer immunity. *Nat Cancer* 1: 184-196.

592   63.   Mekers, V. E., V. M. Kho, M. Ansems, and G. J. Adema. 2022. cGAS/cGAMP/STING
593   signal propagation in the tumor microenvironment: Key role for myeloid cells in
594   antitumor immunity. *Radiother Oncol* 174: 158-167.

595   64.   Ritchie, C., J. A. Carozza, and L. Li. 2022. Biochemistry, Cell Biology, and
596   Pathophysiology of the Innate Immune cGAS-cGAMP-STING Pathway. *Annu Rev*
597   *Biochem* 91: 599-628.

598   65.   Wang, M., P. Xu, and Q. Wu. 2024. Cell-to-cell communications of cGAS-STING
599   pathway in tumor immune microenvironment. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 53:
600   15-24.

     FIGURE LEGENDS

Figure 1. STING-activated macrophages and monocytes do not upregulate *CCL22*. A)

Microscopic images (20x brightfield) of THP-1 monocytes (upper left) and fully differentiated

THP-1-derived macrophages (lower right) are shown. **B)** THP-1-derived macrophages were

treated with cGAM(PS)2 (30 μM) for 24 hours, then harvested in TRIzol for RNA purification

and RTqPCR. Relative *CCL22* gene expression is shown. **C)** THP-1-derived macrophages were

treated with cGAM(PS)2 (30 μM) for 6 and 24 hours, then whole cell lysates (20 μg) and

antibody positive control (7 μg) were resolved with SDS-PAGE, transferred to nitrocellulose,

and probed for phosphorylated STING and beta-tubulin. **D)** THP-1 monocytes were treated with

cGAM(PS)2 (30 μM) and harvested for RTqPCR and *CCL22* gene expression as described in *B*.

Error bars represent standard deviations of independent biological replicates. Statistical analyses

were performed with unpaired, two-tailed *t* test (*B*) or non-parametric Wilcoxon signed-rank test

(*D*) due to non-Gaussian distribution of the treated sample per Shapiro-Wilk test.

Figure 2. STING-activated macrophages and monocytes release factor(s) that upregulate

***CCL22* in cancer cells. A)** MCF7 and **B)** HeLa cells were treated for 24 hours with cGAM(PS)2

(30 μM) or conditioned media (CM) from macrophages that were either untreated or treated with

cGAM(PS)2 (30 μM) for 24 hours prior to media collection. Cells were harvested with TRIzol

for RNA purification and RTqPCR to assess relative expression of *CCL22*. **C)** MCF7 and **D)**

HeLa cells were treated for 24 hours with cGAM(PS)2 (30 μM) or conditioned media (CM) from

monocytes that were either untreated or treated with cGAM(PS)2 (30 μM) for 24 hours prior to

media collection. Cells were harvested with TRIzol for RNA purification and RTqPCR to

determine expression of *CCL22* as above. For all figures, error bars represent standard deviations

of independent biological replicates. Statistical analyses were performed with one-way ANOVAs

on log-transformed data and Tukey's pairwise comparisons (*A*, *C*, and *D*) or with Brown-

Forsythe and Welch ANOVA (*B*) due to significant differences among standard deviations.

Figure 3. Paracrine activation of *CCL22* in cancer cells activates STING and is dependent

on STING. *A)* HeLa cells were treated for 24 hours with cGAM(PS)2 (30 μM) or conditioned

media (CM) from macrophages that were either untreated or treated with cGAM(PS)2 (30 μM)

for 24 hours. Whole cell lysates (25 μg) and antibody positive control (5 μg) were resolved with

SDS-PAGE, transferred to nitrocellulose and probed for phosphorylated STING and beta-

tubulin. The image shown is representative of at least three independent experiments. *B)* HeLa$^{-/-}$

^{TMEM173} STING knockout cells were treated for 24 hours with cGAM(PS)2 (30 μM) or

conditioned media (CM) from macrophages that were either untreated or treated with

cGAM(PS)2 (30 μM) for 24 hours prior to media collection. Cells were harvested with TRIzol

for RNA purification and RTqPCR to determine relative expression of *CCL22*. Error bars

represent standard deviations of independent biological replicates. Statistical analysis was

performed using one-way ANOVA and Tukey's pairwise comparisons.

Figure 4. Type I IFNs have modest effects on *CCL22* in HeLa cells that remain STING-

dependent. HeLa *(A-B)* and MCF7 cells *(C- D)* were treated with exogenous type I IFNs at

indicated concentrations for 48 (HeLa) or 24 hours (MCF7), then harvested with TRIzol for

RNA purification and RTqPCR to determine relative gene expression of *CCL22*. *E-F)* HeLa $^{-/-}$

^{TMEM173} STING knockout cells were treated with exogenous type I IFNs at indicated

concentrations for 48 hours, then harvested with TRIzol for RNA purification and RTqPCR

analysis of *CCL22*. Error bars represent standard deviations of independent biological replicates.

Statistical analyses were performed using one-way ANOVA with Dunnett's multiple

comparisons test (*A, C, E*), one-way ANOVA on log-transformed data with Tukey's pairwise

comparisons (*B, D*), and unpaired, two-tailed t test (*F*). *G)* HeLa wild type cells were treated

with IFNβ (100 ng/mL) for 48 hours, and whole cell lysates (60 μg) and antibody positive

control (5 μg) were resolved with SDS-PAGE, transferred to nitrocellulose and probed for

phosphorylated STING and beta-tubulin. *H)* HeLa wild type (WT) and HeLa $^{-/-TMEM173}$ STING

knockout (KO) cells were treated as indicated for approximately 30 hours prior to harvesting;

whole cell lysates (15 μg) were resolved with SDS-PAGE, transferred to nitrocellulose, and

probed for phosphorylated Stat1 and beta-tubulin. *I)* HeLa cells were pretreated for 3 hours with

MMHAR-2 neutralizing antibody, then indicated treatments were added, including conditioned

media (CM) from macrophages treated with cGAM(PS)2 (30 μM), and cells were incubated for

approximately 48 hours prior to harvest in TRIzol for RNA purification and RTqPCR to assess

relative changes in *CCL22* expression. Statistical analyses were performed using one-way

ANOVA with Šídák's multiple comparisons test.

Figure 5. STING agonists differentially upregulate *CCL22* and *IFNβ*. HeLa *(A-D)* and

MCF7 cells *(E-H)* were treated with 30 μM of each indicated STING agonist or DMSO vehicle

control (for diABZI) for 24 hours, then harvested in TRIzol for RNA purification and RTqPCR

using TaqMan assays for *CCL22* or *IFNb*. Error bars represent standard deviations of

independent biological replicates. Statistical analyses were performed on log-transformed data

using unpaired, two-tailed t tests except for C and G, which were analyzed with a one-way

ANOVA and Šídák's multiple comparisons test.

Fig. 1

A

monocytes

Macrophage
differentiation
with PMA

macrophages

B

CCL22

Relative expression mRNA

cGAM(PS)2 μM   0   30

**

C

cGAM(PS)2 30 μM

—   24 hr   6 hr   Ab + control

37 kDa — .... pSTING

50 kDa — .... β-tubulin

D

CCL22

Relative expression mRNA

cGAM(PS)2 μM   0   30

ns

Fig. 2

Fig. 3

A

cGAM(PS)2 (30 µM): − + CM − + Ab + control

37 kDa —   pSTING

50 kDa —   β-tubulin

B

CCL22

Relative expression mRNA

ns

cGAM(PS)2 (30 µM)	−	+	−	−
CM macrophages, untreated	−	−	+	−
CM macrophages, treated	−	−	−	+

Fig. 4

Fig. 5

Fig. S1

Supplementary Figure 1. THP-1-derived macrophages were untreated (UN) or treated with cGAM(PS)2 (30 μM) for indicated times, then harvested in TRIzol for RNA purification and RTqPCR. Relative gene expression of *CCL22* is shown. Error bars represent standard deviations of independent biological replicates. Statistical significnace was determined uisng Kruskal-Wallis multiple comparisons test.

Fig. S2

Supplementary Figure 2. *A)* HeLa cells (wild type) were untreated or treated for 24 hours with either cGAM(PS)2 (30 μM) or conditioned media (CM) from untreated monocytes or monocytes treated with cGAM(PS)2 (30 μM) for 24 hours. Whole cell lysates (60 μg) and antibody positive control (15 μg) were resolved with SDS-PAGE, transferred to nitrocellulose and probed for phosphorylated STING and beta-tubulin. *B)* HeLa [-/- TMEM173] (STING KO) cells were untreated or treated with either cGAM(PS)2 (30 μM) or conditioned media (CM) from untreated monocytes or monocytes treated with cGAM(PS)2 (30 μM) for 24 hours. Cells were harvested in TRIzol for RNA purification and RTqPCR to determine relative levels of *CCL22*.

Fig. S3

Supplementary Figure 3. *A)* HeLa cells were treated for 24 hours with exogenous IFNβ at indicated concentrations, then harvested in TRIzol for RNA purification and RTqPCR to assess relative fold changes in *CCL22* expression. Error bars represent standard deviations of independent biological replicates, and statistical analysis was performed with one-way ANOVA and Dunnett's pairwise comparsions. *B)* MCF7 cells were treated for 24 or 48 hours with 10 ng/mL of IFNα or IFNβ, then harvested for RTqPCR as in *A*. *C)* Whole cell lysates (40 μg) from HeLa wild type (WT) and HeLa [-/-TMEM173] STING knock out (KO) cells were resolved with SDS-PAGE, transferred to nitrocellulose, and probed for IFNAR2 and beta-tubulin. *D-E)* HeLa cells were pretreated for 3 hours with MMHAR-2 neutralizing antibody, then indicated treatments were added and cells incubated for approximately 48 hours, at which time whole cell lysates (5 μg) were resolved with SDS-PAGE, transfered to nitrocellulose and probed with anti-pStat1 and β-tubulin.