Dynamic Modulation of Beta-Band Oscillations in the LGN

and Their Role in Visual Processing

Henry J. Alitto[1,2+], Alyssa N. Sanchez[1,3,4], Prescot C. Alexander[1,4], and W. Martin Usrey[1,2]

[1] Center for Neuroscience, University of California, Davis, Davis, CA, 95618, USA

[2] Department of Neurobiology, Physiology, and Behavior, University of California, Davis,

Davis, CA, 95618, USA

[3] Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA, 94720

[4] Department of Neuroscience, University of California, Berkeley, Berkeley, CA, 94720

[3]Information Systems and Modeling Group (A-1), Los Alamos National Laboratory, Los

Alamos, NM, 87545, USA

[+] hjalitto@ucdavis.edu

Running title: Thalamocortical LGN beta oscillations

Key words: thalamus, cortex, contrast gain control, arousal, spatial attention

Number of pages: 14

Number of figures: 10

Number of words for abstract: 218

Number of words for introduction: 645

Number of word discussion: 1306

Acknowledgments: We thank K.E. Neverkovec, Jeffrey S. Johnson, D.J. Sperka, and R. Oates
for expert technical assistance. We also thank Rober E. Boshra and Nancy Kopell for their
comments on the manuscript.

31
Conflict of Interest: Authors report no conflict of interest

Funding sources: This work was supported by NIH grants EY012576, EY036242, and
P50MH132642

<u>Abstract</u>

Neuronal oscillations are a ubiquitous feature of thalamocortical networks and can be

dynamically modulated across processing states, enabling thalamocortical communication to

flexibly adapt to varying environmental and behavioral demands. The lateral geniculate nucleus

(LGN), like all thalamic nuclei, engages in reciprocal synaptic interactions with the cortex,

relaying retinal information to and receiving feedback input from primary visual cortex (V1).

While retinal excitation is the primary driver of LGN activity, retinal synapses represent a

minority of the total synaptic input onto LGN neurons, allowing for both retinogeniculate and

geniculocortical signals to be influenced by nonretinal sources. To gain a holistic view of

network processing in the geniculocortical pathway, we performed simultaneous extracellular

recordings from the LGN and V1 of behaving macaque monkeys, measuring local field

potentials (LFPs) and spiking activity. These recordings revealed prominent beta-band

oscillations coherent between the LGN and V1 that influenced spike timing in the LGN and were

statistically consistent with a feedforward process from the LGN to V1. These thalamocortical

oscillations were suppressed by visual stimulation, spatial attention, and behavioral arousal,

strongly suggesting that these oscillations are not a feature of active visual processing. Instead,

they appear analogous to occipital lobe, alpha oscillations recorded in humans and may

represent a signature of signal suppression that occurs during periods of low engagement or

active distractor suppression.

Significance Statement

Oscillations within thalamocortical networks in the awake state are generally believed to enhance communication between the thalamus and cortex, allowing circuits to flexibly respond to changes in sensory, behavioral, and cognitive demands. Here, we show that oscillations within and between the LGN and V1 are suppressed by increases in visual stimulation, increases in behavioral arousal, and shifts in covert spatial attention. We therefore conclude that these oscillations are not a mechanism to enhance the transmission of retinal information through the LGN to V1. Instead, we propose that they are a signature of signal suppression that occurs when network engagement is low or during active distractor suppression.

Introduction

The thalamus is the inextricable partner of the cortex, serving as a critical hub for sensory and cognitive information flow from the periphery to the cortex, as well as providing a transthalamic pathway for communication between cortical areas (Jones, 1985; Sherman and Guillery, 2013; Usrey and Sherman, 2023). Indeed, all thalamic relay nuclei have reciprocal connections with the cortex, and all cortical areas have reciprocal connections with the thalamus. Neuronal oscillations are a ubiquitous feature of thalamic network activity, varying with sensory, behavioral, and cognitive demands (Bazhenov et al., 2000; Steriade, 2006; Saalmann et al., 2012; Zhou et al., 2016; Bazinet et al., 2021; Jayachandran et al., 2023). For higher order thalamus, including the pulvinar, oscillations are thought to modulate thalamocortical communication, allowing networks to adapt to changing demands (Saalmann et al., 2012; Sweeney-Reed et al., 2014; Helfrich and Knight, 2016; Halassa and Kastner, 2017; Hwang and Shine, 2023). Whether oscillations serve similar roles in first-order thalamus remains unclear.

In the visual system, the lateral geniculate nucleus (LGN) of the dorsal thalamus is a first order nucleus and the primary source of retinal information to the cortex, receiving monosynaptic excitation from the retina and sending monosynaptic excitation to primary visual cortex (V1; Usrey and Alitto, 2015). While the LGN's primary role is relaying retinal signals to V1, this transmission is not absolute; instead, retinal signals are selectively gated according to visual processing demands and behavioral state (Usrey et al., 1998; O'Connor et al., 2002; McAlonan et al., 2008; Rathbun et al., 2010; Alitto et al., 2011, 2019; Wimmer et al., 2015; Fisher et al., 2017; Crombie et al., 2024). This gating is influenced by non-retinal inputs, which comprise the majority of LGN synapses, including cortical feedback from V1, GABAergic

inhibition from local interneurons and the thalamic reticular nucleus, and cholinergic input from

the brainstem (Harting and Guillery, 1976; Steriade et al., 1988; Guillery and Harting, 2003;

Wang et al., 2006; Usrey and Alitto, 2015; Krueger and Disney, 2019).

Previous work identified coherent beta-band oscillations between the LGN and V1 in

behaving macaques (Bastos et al., 2014). While these geniculocortical oscillations exhibit

temporal features consistent with feedforward transmission, it remains unknown whether they

are modulated by network demands. To address this, we quantified how geniculocortical

oscillation magnitude changes as the LGN becomes more engaged in visual processing due to

increased visual stimulation or behavioral demands. We hypothesize that if these oscillations

actively facilitate the transmission of retinal information through the LGN to V1, then they should

increase in magnitude as the strength of the visual stimulus increases and the behavioral

significance of the visual stimulus increases.

To gain a holistic understanding of oscillations in the geniculocortical pathway, we

recorded neuronal activity from behaving macaques and monitored local in the LGN and V1

across behavioral and visual conditions. Consistent with previous reports (Bastos et al., 2014),

we found robust beta-band oscillations within the LGN that were coherent with oscillations in

retinotopically neighboring locations in V1 and were consistent with a feedforward

thalamocortical process. We further found that these beta-band oscillations influenced spike

timing within the LGN, providing a mechanism for these oscillations to be communicated from

the LGN to V1. In contrast to the hypothesis that these oscillations facilitate visual processing,

we observed that they were suppressed during periods of increased LGN processing demands.

In particular, the geniculocortical oscillations were suppressed by increased visual stimulation,

heightened arousal, and shifts of spatial attention toward the LGN receptive field center. These

dynamics resemble those of occipital alpha oscillations in human EEG, which may represent a

signature of signal suppression that occurs when processing demands are low or during active

distractor suppression. A possible source for these oscillations is the thalamic reticular nucleus,

which provides GABAergic inhibition to the LGN and is modulated by behavioral arousal and

attention (Guillery and Harting, 2003; McAlonan et al., 2008; Wimmer et al., 2015).

MATERIALS AND METHODS

Six adult rhesus monkeys (*Macaca mulatta*; four female and two male) were used for this study. All experimental procedures conformed to National Institutes of Health and United States Department of Agriculture guidelines and were approved by the Institutional Animal Care and Use Committee at the University of California, Davis. Under full surgical anesthesia, the animals received a cranial implant consisting of a head post for head stabilization and two recording cylinders that allowed access to the LGN and V1.

Electrophysiological recordings. Extracellular recordings from the LGN and V1 were made with platinum-glass electrodes (1–2 MV; Alpha Omega) using a microdrive (40 mm electrode travel; Thomas Motorized Electrode Manipulator, Thomas RECORDING) mounted on the recording chamber. Continuous voltage signals were amplified (A-M Systems), filtered (0.1–5 kHz), and recorded using a Micro1401 data acquisition system (28 kHz) and Spike2 software (Cambridge Electronic Design, CED). To extract local-field potentials (LFPs), extracellular recordings were low-pass filtered (stopband edge frequency = 250 Hz) with a 6th degree, Chebyshev filter (Matlab: cheby2, filtfilt). LFP signals were subsequently subsampled to 500 Hz and notch filtered at 60 Hz to remove mains noise (Matlab, custom software).

Visual stimulation and receptive field mapping. Visual stimuli were generated with a ViSaGe (Cambridge Research Systems) and presented on a gamma-corrected CRT monitor (Sony or Mitsubishi) positioned ~80 cm in front of the animal. The display had a resolution of 1024x768, a refresh rate of 100-140Hz, and a mean luminance of ~38 cd/m2. Receptive fields of LGN and V1 neurons were manually mapped using small (<0.5°), user-controlled, computer-generated visual stimuli (custom software with Spike2 interface).

Behavioral training and performance. Data was collected while the animals performed either a passive fixation or a spatial attention task. For both tasks, eye position and pupil size were monitored using an Eye-Trac6 infrared, eye tracker (Applied Science Laboratories) with a custom Spike2 interface. For passive fixation, the animals fixated on a small (~0.2° diameter) fixation dot, using a fixation window of 0.8°-1.2° diameter. During passive fixation, sine wave gratings varying in stimulus contrast, spatial frequency, and size were presented centered over the retinotopic coordinates of the recorded neuron's receptive field.

Spatial attention. Two animals were trained to perform a spatial attention task. For this task, the animals initiated a trial by fixating a central fixation dot similar to the fixation dot used for passive fixation. When the animal obtained fixation, two sine wave gratings (temporal frequency

= 5 Hz, spatial frequency = 1.0 cycles/°) appeared at equidistant locations from the fovea,

typically at an eccentricity of 4-8°. Each sine wave grating was surrounded by a colored circle

(one green, one red) and the color of the circle at each location was constant throughout the

task. The color of the fixation point (either red or green) indicated to the animal which grating

was 95% likely to change contrast (valid trials) after a delay with an exponential hazard function

of 0.6 – 1.8s. On 5% of the trials (invalid trials), the uncued grating increased in contrast (same

hazard function). The base contrast of the grating was set to be near the C50 (contrast to evoke

a half-maximum response) of the recorded LGN neuron, as estimated through hand mapping,

and the contrast change was titrated so that the animal performed ~ 85% correctly. After the

contrast change, the animal was given 1s to respond by making a saccade to the location of the

change, and the animal was rewarded for correct responses with a small juice reward.

Psychophysical performance (accuracy and reaction time) for valid and invalid trials was

compared to ensure that the animals were applying spatial attention as anticipated.

Analysis

Oscillatory LFP activity. LFP power spectra from LGN and V1 recordings were computed using

a Fourier transform with a Hanning window (custom Matlab script). To identify significant

oscillatory activity, we first estimated the mean and standard deviation of the non-oscillatory,

fractal component via a bootstrap analysis of the LFP power spectra. Specifically, we iteratively

sampled the power spectrum with replacement 1000 times, drawing the same number of trials

as in the original dataset for each iteration, and fitting each to a linear, log-log relationship

between power and frequency:

171
$$power_{fractal} = m * \frac{1}{\log_{10} f} + b.$$

172 Here, m and b were optimized to capture the linear relationship between log(power) and

173 log(frequency), respectively. From this, the residual oscillatory power was defined as

174
$$power_{oscillatory} = power_{total} - mean(power_{fractal})$$

175 Similarly, the Z-score of the oscillatory power was defined as

176
$$Z_{oscillatory} = (power_{oscillatory} - mean(power_{fractal}) \, / \, std(power_{fractal})$$

177 Z-scores greater than 5.0 were taken as evidence of statistically significant oscillatory activity at

178 a given frequency.

179   *LGN-V1 phase synchrony:* Frequency-specific LGN-V1 field-field phase synchrony was

180   estimated using pairwise phase consistency (ppc, Vinck et al., 2010):

181
$$ppc = \sum_i \sum_j \cos \Phi_i * \cos \Phi_j + \sin \Phi_i * \sin \Phi_j$$

Here, Φ_i and Φ_j are the phase angles between the LGN and V1 on trials i and j, respectively.

Low ppc values near 0 indicate no consistent relationship between LGN and V1 LFP phase,

while ppc values near 1.0 indicate a constant phase relationship between LGN and V1.

To determine the significance of PPC values, we estimated the expected standard deviation of

PPC under the null hypothesis of no consistent phase relationship. We modeled the null as a

random walk, where each step is taken from the distribution of $\cos(\Phi_i - \Phi_j)$ where Φ_i and Φ_j

are independently drawn from a uniform distribution over [-π, π]. In this case, $\cos(\Phi_i - \Phi_j)$

follows an arcsine distribution on [-1, 1], which has a variance of 0.5. If we assume that the

expected step size is 0 (Vinck et al., 2010) and the variance of the step size is 0.5, then the total

displacement after N steps:

192
$$S_N = \sum_{i=1}^{N} x_i$$

193   This is a sum of independent and identically distributed variables, so the variance of the total
194   displacement is:

195
$$var(S_N) = N * var(x_i) = N * (0.5) = N/2$$
196

197   Since PPC considers normalized displacement,

198
$$PPC \sim \bar{S}_N = \frac{S_N}{N}$$

199   This is the mean of N zero-mean random steps. Thus,

200
$$var(\bar{S}_N) = var\left(\frac{S_N}{N}\right) = \frac{var(S_N)}{N^2} = \frac{\frac{N}{2}}{N^2} = \frac{1}{2N}$$

201
$$std(\bar{S}_N) = 1/\sqrt{2N}$$

Here, N is the number of pairwise comparisons. PPC values were then z-scored based on this

analytically defined N-dependent standard deviation. The validity of this estimation of the null

distribution was supported by simulation of PPC values based on random phases using a range

of N values (Figure S1). Values exceeding 5.0 standard deviations were considered evidence of

significant field-field phase synchrony.

Directionality of LGN-V1 communication. To determine if LGN-V1 phase synchrony resulted

from either a feedforward or feedback process, we calculated the directionality of LGN-V1 LFP

interactions using phase slope index (psi, (Nolte et al., 2008b). Positive and negative psi values

are statistically consistent with feedforward and feedback processes, respectively. The

significance of psi values was not estimated for individual recordings but instead was estimated

for the sample distribution as a whole using a standard t-test to determine statistically significant

differences from 0.

Spike-field phase synchrony. LGN spike-field phase synchrony was estimated with ppc, using

the phase of 0.4s LFP snippets centered on spike times. Because recordings were made using

single, high-impedance electrodes, the LFP data was contaminated with residual spike

waveforms that could not be removed by lowpass filtering. These spike waveform artifacts

potentially interfered with the ability to detect spike-field phase synchrony, particularly in the

beta band, where we saw the best evidence for significant oscillatory activity in the LGN. To

deal with this issue, we reasoned that the strength of frequency-specific spike-field synchrony

would depend upon the presence of strong oscillatory activity in the LGN, while the influence of

the spike artifact would be independent of the underlying oscillatory activity. Therefore, we

calculated ppc values for epochs of strong and weak beta band oscillatory activity and defined

the uncontaminated estimate of LGN spike-field phase synchrony as the difference between

these two values.

Matching Pursuit. To visualize spike-related transients in the local field potential, we used

matching pursuit (mp), a greedy algorithm that decomposes signals into linear combinations of

time–frequency localized basis functions ("atoms") drawn from an overcomplete dictionary. This

approach adaptively selects atoms that best match the signal structure, allowing high-resolution

representations of both rhythmic and transient features. We implemented the mp algorithm in

Matlab following the procedure described by (Chandran et al., 2016), using Gabor atoms

spanning a range of scales. Matching pursuit was used solely for visualization and was not

involved in any quantitative analyses of spike–LFP phase relationships.

RESULTS

The lateral geniculate nucleus (LGN) is the primary source of retinal information to the cerebral cortex, receiving monosynaptic excitation from retinal ganglion cells and sending monosynaptic excitation to primary visual cortex (V1). The primary goal of this study was to gain a holistic view of network dynamics in the LGN by recording local field potentials (LFPs, n = 1,117) from behaving macaque monkeys (n = 6) under various behavioral conditions and visual stimulation paradigms. Single-unit spiking activity was also recorded. In a subset of recordings, we simultaneously recorded from both the LGN and V1 at similar retinotopic locations (n = 227). Unless otherwise stated, recordings were made while the monkeys passively fixated on a central location with visual stimuli centered over the LGN receptive field.

As will be shown below, our results from the LGN recordings show a prominent oscillation in the beta-frequency band (low beta = 11-23 Hz, high beta = 24 – 35 Hz), that influenced spike timing within the LGN, was coherent with oscillations in V1, and was statistically consistent with a feedforward, thalamocortical process from the LGN to V1. These thalamocortical oscillations were suppressed by visual stimulation, behavioral arousal, and spatial attention directed toward the LGN receptive field.

Beta-band oscillations in the LGN

Our first objective was to measure LGN power spectra with the goal of determining whether there are significant neuronal oscillations in the local network. Towards this goal, extracellular recordings were made from six passively fixating macaque monkeys, totaling 888 recordings from both parvocellular and magnocellular layers (Figure 1, column 1). To determine the presence of significant neuronal oscillations, each power spectrum was fit to a 1/f equation on a log-log scale (Figure 1, column 2). This allowed us to estimate and remove the non-oscillatory, fractal component (Figure 1, column 3), leaving an estimate of the oscillatory power. A bootstrap analysis was used to estimate the significance of the residual oscillatory power at each frequency (Figure 1, column 4).

Neuronal oscillations were most commonly observed in the LGN LFP in the range of 11-35 Hz (Figure 2A and B). This roughly corresponds to the lower (11-23 Hz) and higher (24-35 Hz) segments of the beta-frequency band (Kopell et al., 2000; Jensen et al., 2005; Cohen et al., 2007). Indeed, the distribution of significant frequencies has two clear peaks, one centered over the low-beta range and one centered over the high-beta range (Figure 2B). Across our six animals, 81% of recordings showed significant beta-band oscillations in either the low- or high-

beta band (75.9% low-beta, 30.4% high-beta), with a range of 58.8% - 96.3% across animals. In

5 of 6 animals, low-beta oscillations were the dominant neural oscillation (Figure 2A-F, H), with

one animal having a near-even distribution of low-beta and high-beta oscillations (Figure 2G). In

general, we detected few functional differences between low- and high-beta oscillations. In the

sections below, we identify any significant differences between low-beta and high-beta

oscillatory properties, using the term beta oscillation when no differences were observed. Less

commonly, significant neural oscillations were also seen in other frequency bands. In particular,

we observed significant alpha-band oscillations (6-10 Hz), gamma-band oscillations (40-70 Hz),

and broad-band, high-frequency oscillations (>70 Hz) in a minority of recordings (17.9%, 10%,

23%, respectively). These oscillations will not be emphasized here but may be of interest for

future studies.

279 Coherent thalamocortical beta-band oscillations

Given that the primary role of the LGN is to relay retinal information to V1, we next sought to

understand the relationship between beta-band oscillations in the LGN and V1. Towards this

goal, in a subset of our recordings, simultaneous recordings were made from regions in the

LGN and V1 that had a maximum retinotopic displacement of 3.0° visual degrees (minimum =

0.1°, median = 1.4°). From these recordings, pairwise phase consistency (ppc) was used to

calculate field-field phase synchrony between the LGN and V1 (see Materials and Methods).

Similar to the distribution of significant neural oscillations within the LGN, significant

LGN-V1 field-field synchrony was detected in the beta-band frequency range, as can be clearly

seen in the four example LGN-V1 pairs (Figure 3A-D) and the sample average (Figure 3E1).

Across all four monkeys, 80.6% and 63.9% of recordings had significant LGN-V1 phase

synchrony in the low-beta and high-beta bands, respectively. As expected, the probability of

finding significant beta-band LGN-V1 phase synchrony was dependent upon the presence of

significant beta-band oscillatory power in the LGN ($Z_{ßlow}$ = 2.17, p = 0.0147; $Z_{ßhigh}$ = 4.9, p < 10^{-}

^{5}). Similar relationships were seen for the alpha band (Z_{alpha} = 7.0, p < 10^{-5}) and the high-

frequency bands (Z_{alpha} = 4.6, p = 10^{-5}), but not the gamma band (Z_{gamma} = 0.034, p = 0.4866).

Further, the magnitude of LGN-V1 beta-band phase synchrony was highly dependent upon the

strength of the underlying LGN oscillation (Figure 3E2-3), being significantly stronger during

periods of high oscillatory strength compared to periods of low oscillatory strength for both low-

beta (ppc_{weak} = 0.10±0.01, ppc_{strong} = 0.17±0.02, p < 10^{-5}) and high-beta oscillations (ppc_{weak} =

0.11±0.01, ppc_{strong} = 0.20±0.02, p < 10^{-5}) (Figure 3E2-3). This indicates that beta-band

coherence is not simply a reflection of recording from two synaptically connected areas. Instead,

the dependence of the LGN-V1 phase synchrony on the strength of the underlying LGN

oscillation indicates stronger communication between the LGN and V1 when beta oscillations

are present. Finally, using phase slope index (psi) as a measure of the directionality of LGN-V1

communication, we determined that beta oscillations were statistically consistent with a

feedforward, thalamocortical process from the LGN to V1 (Figure 3F).

Spike-field phase synchrony in the LGN

We next sought to determine the functional influence of LGN beta-band oscillations on LGN

spiking activity. Before describing these interactions, it is useful to consider the involvement of

artifactual contributions that can affect the analyses performed and a means to avoid them. The

data analyzed for this study were collected with single, high-impedance electrodes, typically with

well-isolated single units with large signal-to-noise ratios (Figure 4, column 1). Previous studies

have demonstrated that when the spiking activity and LFP are recorded from the same

electrode, the residual spike waveform, which cannot be completely removed by high-pass

filtering, will cause artifactual interactions between the LFP and spiking activity (Ray et al., 2008;

Ray, 2015). Indeed, the influence of residual spike waveforms is evident in the spike-triggered

average LFP waveform (Figure 4, column 2), and the residual spike waveform underlies an

artificial, broad-band, high-frequency spike-field interaction that dominates the average LGN ppc

curves (Figure 4, column 3). The relationship between the spike waveform and the high-

frequency coherence can be further visualized using matching pursuit (see Materials and

Methods), a spectral analytic technique that allows one to achieve temporal resolution that

exceeds the theoretical limits of traditional Fourier analysis. As can be seen from the average

matching pursuit across the sample of LGN recordings (Figure 4C), the broadband, high-

frequency, spike-field synchrony is restricted to a narrow temporal window (~8ms) surrounding

the time of LGN action potentials. This is more consistent with a spike artifact than an ongoing

oscillation. Moreover, we found that the strength of the broadband, high-frequency, spike-field

phase synchrony was highly correlated with the signal-to-noise of the underlying spike

waveform (Figure 4D, r = 0.56, p < 10^{-5}). Thus, the residual spike waveform would interfere with

the ability to measure spike-field interactions using traditional approaches.

To quantify spike-field phase synchrony and avoid the influence of the spike artifact, we

applied the following reasoning. The strength of spike-field synchrony should be proportional to

the underlying oscillatory power. Thus, there should be a higher degree of synchrony during

periods when the oscillatory power is high compared to periods where it does not reach

statistical significance. By comparison, the influence of the spike artifact should not depend

upon the strength of the underlying neural oscillation. We therefore calculated spike-field ppc for

periods of both strong and weak oscillatory power (Figure 5A-C, red and blue curves,

respectively) and subtracted these two curves to obtain a spike artifact-free estimate of spike-

field phase synchrony (Figure 5D and E). Across the sample of LGN recordings, the majority of

cells had weak, statistically nonsignificant, spike-field phase synchrony, even when the analysis

was restricted to epochs of strong beta-band activity (Figure 5F, see example C; ppc_{strong} =

0.001±5.3*10^{-4}, ppc_{weak} = 0.0006±1.7*10^{-4}). However, a minority of LGN recordings exhibited

statistically significant beta-band, spike-field synchrony (Figure 5, see examples A and B), and

the percentage of cells increased during epochs of strong beta-band activity (97/397, 24.4%,

ppc_{strong} = 0.016±0.003) relative to epochs of weak beta-band activity (33/397, 8.3%, ppc_{weak} =

0.012±0.003, p < 10^{-5}) (Figure 5E-G). Thus, if beta oscillations are indeed a feed-forward,

thalamocortical signal from the LGN to V1, then the signal is likely conveyed by a minority of the

LGN neurons.

Thalamocortical beta-band oscillations are suppressed by active visual processing

If beta-band thalamocortical oscillations are a mechanism that facilitates the transmission of

retinal information to the cortex, then it seems reasonable to expect that both LGN oscillatory

strength and LGN-V1 field-field phase synchrony would increase during periods when (1) visual

stimulation is high, (2) spatial attention is directed toward the recorded retinotopic location, and

(3) behavioral arousal is high. Each of these is examined below, and the results do not support

the conclusion that thalamocortical beta-band oscillations are a mechanism underlying active

visual processing. Instead, these oscillations are more consistent with a signature of

suppression that is active when visual and behavioral processing demands are low or the

demands for distractor suppression are high.

To quantify the influence of visual stimulation on thalamocortical oscillations, we first

investigated the influence of stimulus strength on beta-band oscillations in the geniculocortical

pathway. For this analysis, epochs of visual stimulation with sine wave gratings were divided

into strong and weak classifications based on the stimulus contrast and size. Gratings that were

larger than 1.5° (as large as 10°) and 50% contrast (up to 100% contrast) were considered

"strong stimuli", while gratings smaller than 1.0° (as small as 0.3°) and less than 25% contrast

(as low as 5% contrast) were considered "weak stimuli". As can be seen from both the

examples and the sample average, both LGN beta-band oscillatory power (Figure 6A-E,

$index_{ßlow}$ = -0.05±0.006, p < 10^{-5}; $index_{ßhigh}$ = -0.05±0.01, p < 10^{-4}) and LGN-V1 beta-band phase

synchrony (Figure 7A-E; $index_{ßlow}$ = -0.08±0.02, p < 0.001; $index_{ßhigh}$ = -0.12±0.02, p < 10^{-5})

decreased during periods of strong visual stimulation compared to periods of weak visual

stimulation. Importantly, the decrease in both measures was restricted to the beta-frequency

band and was not observed in other frequency bands (Figure 6F and Figure 7F). Additionally,

when the analysis was restricted to recordings with significant alpha-band LGN-V1 phase

synchrony (Figures 6F and 7F), there was a significant increase in alpha-band oscillatory

strength ($\text{index}_\alpha = 0.07\pm0.03$, $p = 0.011$) and LGN-V1 phase synchrony ($\text{index}_\alpha = 0.17\pm0.05$, $p =$

0.0017) during the presentation of strong visual stimuli.

Oscillatory activity throughout the brain is modulated by changes in behavioral state.

This includes changes in behavioral arousal as well as changes in cognitive demands. Previous

studies have demonstrated that the LGN is sensitive to levels of behavioral arousal (Livingstone

and Hubel, 1981; Crombie et al., 2024). Thus, we next investigated the influence of arousal on

thalamocortical beta-band oscillations (Figure 8). Towards this goal, we monitored pupil size in

a subset of recordings for two monkeys as an indirect yet reliable physiological measure of

overall behavioral arousal (Reimer et al., 2014; McGinley et al., 2015; Vinck et al., 2015;

Crombie et al., 2024). From these measurements, we divided trials into low- and high-arousal

states, representing the bottom and top 25^{th} percentiles, respectively.

Increases in behavioral arousal, as inferred through increases in pupil size, were

associated with decreases in thalamocortical oscillations. We found a decrease in the high beta-

band LGN oscillatory power (Figure 8A-E; $\text{index}_{\beta high} = -0.19\pm0.02$, $p < 10^{-5}$ $\text{index}_{\beta low} =$

0.005 ± 0.01, $p = 0.58$) as well as a decrease in the low and high beta-band, LGN-V1phase

synchrony (Figure 9A-E; $\text{index}_{\beta low} = -0.12\pm0.04$, $p = 0.001$; $\text{index}_{\beta high} = -0.13\pm0.02$, $p < 10^{-5}$).

Importantly, these decreases were specific to the beta-frequency band and were not

representative of an overall decrease in network activity (Figure 8F and Figure 9F).

Finally, we recorded from two macaque monkeys while they performed a spatial

attention task where their attention was either directed toward or away from the retinotopic

position of the LGN receptive field (see Materials and Methods). Importantly, the animals

displayed psychophysical behavioral signatures of performing the task as anticipated.

Specifically, they were faster and more accurate at reporting changes at the attended location

(valid cue trials) compared to changes at the unattended location (invalid cue trials) (Alitto et al.,

2025). For these experiments, recordings were only made from the LGN, so results are limited

to the influence of spatial attention on LGN oscillatory strength.

Shifting spatial attention toward the retinotopic position of the LGN receptive field caused

a small, but significant, decrease in thalamocortical oscillations as assessed via LGN beta-band

oscillatory strength (Figure 10; $\text{index}_{\text{ßlow}}$ = -0.02±0.006, p = 0.002; $\text{index}_{\text{ßhigh}}$ = -0.04±0.02, p =

0.007). Importantly, the changes to LGN oscillatory power were specific to the beta-frequency

band and did not represent an overall decrease in network activity (Figure 10F3). In contrast, we

saw a small but significant increase in the firing rate of LGN neurons when attention was shifted

toward the LGN receptive field, relative to when attention was shifted away from the LGN

receptive field (Alitto et al., 2025). We speculate that the difference in firing rate may be

associated with the change in beta-band oscillatory power, which may serve to suppress the

transmission of retinal signals that encode visual distractors, such as the unattended drifting

grating.

In summary, this study revealed prominent beta-band oscillations in the LGN that are

coherent with V1 and are statistically consistent with a feedforward, thalamocortical process

from the LGN to V1. These oscillations are suppressed by visual stimulation, behavioral arousal,

and shifts in spatial attention toward the LGN receptive field, indicating they do not represent a

mechanism underlying active visual processing.

DISCUSSION

Consistent with Bastos et al. (2014), we observed robust beta-band oscillations in the lateral geniculate nucleus (LGN) that were coherent with activity in primary visual cortex (V1) in a manner that was statistically consistent with a feedforward thalamocortical signal. These oscillations influenced spike timing in a subset of LGN neurons, suggesting a plausible mechanism for transmitting rhythmic activity to the cortex. Critically, however, we found that LGN-V1 beta-band oscillations were suppressed, rather than enhanced, during periods of increased network engagement. Specifically, this suppression occurred during strong visual stimulation, elevated arousal, and covert shifts of spatial attention toward the LGN receptive field. This inverse relationship between beta-band power and sensory-cognitive demands strongly suggests that thalamocortical oscillations do not enhance information transmission between the LGN and V1. Instead, our findings are more consistent with the robust literature on cortical alpha-band rhythms, which are understood to reflect active suppression of task-irrelevant input and are modulated by attention, arousal, and perceptual load. In the sections below, we expand on this proposed functional analogy between these thalamocortical beta oscillations and cortical alpha oscillations, discuss their shared sensitivity to behavioral context, and explore potential circuit mechanisms that may support this dynamic form of suppression.

Oscillatory mechanisms of signal suppression in the visual system

EEG and LFP recordings from the awake brain, in both humans and animals, are often dominated by cortical alpha-band oscillations (Lopes da Silva et al., 1973; Başar et al., 2001; Palva and Palva, 2007; Rihs et al., 2007; Bollimunta et al., 2011; Mo et al., 2011; Haegens et al., 2015). One of the earliest observations was that alpha power increases when subjects close their eyes during quiet wakefulness, a state long associated with reduced sensory input and diminished visual processing (reviewed in Shomer, 2017). While alpha-band oscillations were once viewed as a signature of cortical idling, they are now widely believed to represent an active inhibitory mechanism that limits interference from task-irrelevant inputs and modulates cortical excitability in accordance with behavioral demands (Kelly et al., 2006; Klimesch et al., 2007; Rihs et al., 2007). This suppressive function has been observed across a range of experimental paradigms. Alpha-band power increases over task-irrelevant visual regions during spatial attention tasks (Gutteling et al., 2022; Yang et al., 2024), across unattended sensory modalities during modality-specific attention (Foxe et al., 1998), and in motor cortex during the inhibition of learned responses (Hummel et al., 2002). In working memory tasks, elevated alpha-band power

suppresses interference from new visual input during the maintenance phase (Herrmann et al.,
2004; Tuladhar et al., 2007; Jensen and Mazaheri, 2010). Importantly, alpha-band activity is not
merely present during suppression, but actively tracks the demands of suppression. For
example, in spatial attention tasks, alpha-band power increases more strongly over regions
representing distractors when those distractors are physically similar to targets (Gutteling et al.,
2022). This observation aligns with perceptual load theory, which posits that the degree of
distractor suppression increases with task difficulty. It is also now well established that alpha-
band activity influences perceptual awareness. Trial-to-trial variability in alpha-band power
predicts whether near-threshold visual stimuli are consciously detected (Di Gregorio et al.,
2022).

Based on the modulation of beta-band oscillations in our dataset by both arousal and
spatial attention, we propose that these oscillations in the macaque geniculocortical pathway
are functionally analogous to the alpha-band oscillations commonly observed in human cortical
EEG. Similar to cortical alpha, LGN-V1 beta-band oscillations were strongest during states of
low engagement and were reduced when sensory or attentional demands increased. This
pattern was evident across multiple behavioral contexts. First, during passive fixation and the
presentation of behaviorally irrelevant visual stimuli, beta-band power increased during periods
of low arousal, as indexed by pupil size. While fixation on a 1° target requires minimal but
sustained engagement, we interpret these oscillations as consistent with quiet wakefulness or
partial disengagement. Second, during a spatial attention task that required significantly more
engagement, we observed that beta-band oscillations were stronger when the distractor, rather
than the target, overlapped the LGN receptive field. This spatial selectivity closely mirrors the
retinotopically specific increases in alpha-band power observed in visual cortex when attention
is directed away from a given location. Together, these findings suggest that LGN-V1 beta-band
oscillations serve as a dynamic index of functional inhibition, suppressing inputs that are
behaviorally irrelevant or unattended. We emphasize that our data demonstrate a correlational
relationship consistent with this suppressive function. Future experiments involving causal
perturbation will be necessary to establish whether beta-band oscillations in the LGN actively
gate information flow to the cortex and/or modulate perceptual sensitivity.

Although we conclude that the LGN-V1 beta-band oscillations observed in macaques
are functionally analogous to cortical alpha-band oscillations, we refrain from labeling them as
"alpha" due to frequency range considerations. Specifically, these oscillations often extended
above 30 Hz, and there is no precedent in the literature for referring to such high-frequency

activity as alpha. To maintain consistency with existing nomenclature, we instead adopt a

conventional labeling scheme: low-beta (11–24 Hz) and high-beta (25–35 Hz) (HajiHosseini et

al., 2012; Zhang et al., 2019). An alternative approach would have been to label the low-

frequency component as alpha and the high-frequency component as beta, as has been done in

some prior studies (e.g., Schmiedt et al., 2014). However, this binary distinction would imply a

functional dissociation between the two sub-bands—an interpretation not supported by our data.

Importantly, while the oscillations we report occupy the beta-frequency range, their peak

frequencies varied across animals and sometimes overlapped with traditional alpha-band

frequencies (e.g., 10–14 Hz). This observation is consistent with prior findings showing

substantial inter-individual variability in alpha frequency (Benwell et al., 2019) and even trial-by-

trial variability within individuals (Minami et al., 2020). Moreover, recent work suggests that

functionally analogous rhythms may appear at lower frequencies in other species, such as 3–5

Hz rhythms in rodent visual thalamus (Nestvogel and McCormick, 2022). Taken together, these

findings highlight the view that the functional role of an oscillation cannot be fully determined by

its frequency alone. Ultimately, while frequency-based labels offer a useful shorthand, they are

only meaningful to the extent that they reflect shared mechanisms and functions across

experimental contexts.

The neural origin of LGN-V1 oscillations

The LGN and V1 are interconnected not only via feedforward excitatory projections from the

LGN to V1 but also through corticothalamic feedback projections originating from layer 6 of V1.

This feedback provides both direct excitation to LGN relay neurons and indirect inhibition via the

thalamic reticular nucleus (TRN) and local interneurons(Sherman and Guillery, 2006). Given this

bidirectional architecture, distinguishing whether oscillations observed in the LGN-V1 circuit

arise through feedforward or feedback (or both) processes is nontrivial.

To probe directionality, we applied a phase slope index (psi, which revealed that LGN-

V1 beta-band synchrony was statistically consistent with a feedforward signal from LGN to V1.

While psi is less susceptible to spurious inferences than other metrics like Granger causality

(Nolte et al., 2008), it is ultimately a statistical inference based on relative phase delays and

cannot alone establish the origin of the observed oscillations. Indeed, evidence from the broader

literature suggests that alpha-band oscillations—functionally similar to the LGN-V1 beta-band

oscillations described here—can arise from both thalamic and cortical sources. In thalamus,

rhythmic inhibition from the TRN and high-threshold burst firing in specialized relay neurons

have been implicated in alpha-frequency generation (Hughes and Crunelli, 2005). In cortex, infragranular activity, particularly in cortical layer 5, has been proposed as a source of alpha-band oscillatory activity, especially during attentionally relevant processing (Bollimunta et al., 2008; Spaak et al., 2012).

We therefore propose that LGN-V1 beta-band oscillations arise from both thalamic and cortical generators, which are then synchronized via recurrent thalamocortical loops. Such hybrid origins have been described for alpha-band rhythms and may also account for the behavioral sensitivity and feedforward dynamics of the beta-band oscillations found in this study.

REFERENCES

Alitto H, Rathbun DL, Vandeleest JJ, Alexander PC, Usrey WM (2019) The augmentation of retinogeniculate communication during thalamic burst mode. J Neurosci 39:5697–5710.

Alitto HJ, Johnson JS, Usrey WM (2025) Spatial Attention Weakly Modulates Visual Responses in the Lateral Geniculate Nucleus. bioRxiv:2025.05.16.652890 Available at: http://biorxiv.org/content/early/2025/05/19/2025.05.16.652890.abstract.

Alitto HJ, Moore BD, Rathbun DL, Usrey WM (2011) A comparison of visual responses in the lateral geniculate nucleus of alert and anaesthetized macaque monkeys. J Physiol 589:87–99.

Başar E, Başar-Eroglu C, Karakaş S, Schürmann M (2001) Gamma, alpha, delta, and theta oscillations govern cognitive processes. Int J Psychophysiol 39:241–248 Available at: https://www.sciencedirect.com/science/article/pii/S0167876000001458.

Bastos AM, Briggs F, Alitto HJ, Mangun GR, Martin Usrey W (2014) Simultaneous recordings from the primary visual cortex and lateral geniculate nucleus reveal rhythmic interactions and a cortical source for gamma-band oscillations. J Neurosci 34:7639–7644.

Bazhenov M, Timofeev I, Steriade M, Sejnowski T (2000) Spiking-bursting activity in the thalamic reticular nucleus initiates sequences of spindle oscillations in thalamic networks. J Neurophysiol 84:1076–1087.

Bazinet V, Vos de Wael R, Hagmann P, Bernhardt BC, Misic B (2021) Multiscale communication in cortico-cortical networks. Neuroimage 243:1–18.

Benwell CSY, London RE, Tagliabue CF, Veniero D, Gross J, Keitel C, Thut G (2019) Frequency and power of human alpha oscillations drift systematically with time-on-task. Neuroimage 192:101–114.

Bollimunta A, Chen Y, Schroeder CE, Ding M (2008) Neuronal mechanisms of cortical alpha oscillations in awake-behaving macaques. J Neurosci 28:9976–9988.

Bollimunta A, Mo J, Schroeder CE, Ding M (2011) Neuronal mechanisms and attentional modulation of corticothalamic alpha oscillations. J Neurosci 31:4935–4943.

Chandran KSS, Mishra A, Shirhatti V, Ray S (2016) Comparison of matching pursuit algorithm with other signal processing techniques for computation of the time-frequency power spectrum of brain signals. J Neurosci 36:3399–3408.

Cohen MX, Elger CE, Ranganath C (2007) Reward expectation modulates feedback-related negativity and EEG spectra. Neuroimage 35:968–978 Available at: http://dx.doi.org/10.1016/j.neuroimage.2006.11.056.

Crombie D, Spacek MA, Leibold C, Busse L (2024) Spiking activity in the visual thalamus is coupled to pupil dynamics across temporal scales. PLoS Biol 22:e3002614.

Di Gregorio F, Trajkovic J, Roperti C, Marcantoni E, Di Luzio P, Avenanti A, Thut G, Romei V (2022) Tuning alpha rhythms to shape conscious visual perception. Curr Biol 32:988-998.e6 Available at: https://www.sciencedirect.com/science/article/pii/S0960982222000148.

Fisher TG, Alitto HJ, Usrey WM (2017) Retinal and nonretinal contributions to extraclassical surround suppression in the lateral geniculate nucleus. J Neurosci 37.

Foxe JJ, Simpson G V, Ahlfors SP (1998) Parieto-occipital approximately 10 Hz activity reflects anticipatory state of visual attention mechanisms. Neuroreport 9:3929–3933.

Guillery RW, Harting JK (2003) Structure and connections of the thalamic reticular nucleus: Advancing views over half a century. J Comp Neurol 463:360–371.

Gutteling TP, Sillekens L, Lavie N, Jensen O (2022) Alpha oscillations reflect suppression of distractors with increased perceptual load. Prog Neurobiol 214:102285 Available at: https://doi.org/10.1016/j.pneurobio.2022.102285.

Haegens S, Barczak A, Musacchia G, Lipton ML, Mehta AD, Lakatos P, Schroeder CE (2015) Laminar profile and physiology of the α rhythm in primary visual, auditory, and somatosensory regions of neocortex. J Neurosci 35:14341–14352.

HajiHosseini A, Rodríguez-Fornells A, Marco-Pallarés J (2012) The role of beta-gamma oscillations in unexpected rewards processing. Neuroimage 60:1678–1685 Available at: https://www.sciencedirect.com/science/article/pii/S1053811912001528.

Halassa MM, Kastner S (2017) Thalamic functions in distributed cognitive control. Nat Neurosci 20:1669–1679.

Harting JK, Guillery RW (1976) Organization of retinocollicular pathways in the cat. J Comp Neurol 166:133–144.

Helfrich RF, Knight RT (2016) Oscillatory Dynamics of Prefrontal Cognitive Control. Trends Cogn Sci 20:916–930.

Herrmann CS, Senkowski D, Röttger S (2004) Phase-locking and amplitude modulations of EEG alpha: Two measures reflect different cognitive processes in a working memory task. Exp Psychol 51:311–318.

Hughes SW, Crunelli V (2005) Thalamic mechanisms of EEG alpha rhythms and their pathological implications. Neuroscientist 11:357–372.

Hummel F, Andres F, Altenmüller E, Dichgans J, Gerloff C (2002) Inhibitory control of acquired

motor programmes in the human brain. Brain 125:404–420.

Hwang K, Shine JM (2023) Thalamic Contribution to Cortical Network Functions. In: The
Cerebral Cortex and Thalamus (Usrey WM, Sherman SM, eds), pp 494–502. Oxford
University Press.

Jayachandran M, Viena TD, Garcia A, Veliz AV, Leyva S, Roldan V, Vertes RP, Allen TA (2023)
Nucleus reuniens transiently synchronizes memory networks at beta frequencies. Nat
Commun 14:1–15.

Jensen O, Goel P, Kopell N, Pohja M, Hari R, Ermentrout B (2005) On the human sensorimotor-
cortex beta rhythm: sources and modeling. Neuroimage 26:347–355.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: gating
by inhibition. Front Hum Neurosci 4:186.

Jones EG (1985) The thalamus / Edward G. Jones. New York: Plenum Press.

Kelly SP, Lalor EC, Reilly RB, Foxe JJ (2006) Increases in alpha oscillatory power reflect an
active retinotopic mechanism for distracter suppression during sustained visuospatial
attention. J Neurophysiol 95:3844–3851.

Klimesch W, Sauseng P, Hanslmayr S (2007) EEG alpha oscillations: The inhibition-timing
hypothesis. Brain Res Rev 53:63–88.

Kopell N, Ermentrout GB, Whittington MA, Traub RD (2000) Gamma rhythms and beta rhythms
have different synchronization properties. Proc Natl Acad Sci U S A 97:1867–1872.

Krueger J, Disney AA (2019) Structure and function of dual-source cholinergic modulation in
early vision. J Comp Neurol 527:738–750.

Livingstone MS, Hubel DH (1981) Effects of sleep and arousal on the processing of visual
information in the cat. Nature 291:554–561.

Lopes da Silva FH, van Lierop THM., Schrijer CF, Storm van Leeuwen W (1973) Organization
of thalamic and cortical alpha rhythms: Spectra and coherences. Electroencephalogr Clin
Neurophysiol 35:627–639 Available at:
https://www.sciencedirect.com/science/article/pii/0013469473902162.

McAlonan K, Cavanaugh J, Wurtz RH (2008) Guarding the gateway to cortex with attention in
visual thalamus. Nature 456:391–394.

McGinley MJ, David S V, McCormick DA (2015) Cortical membrane potential signature of
optimal states for sensory signal detection. Neuron 87:179–192.

Minami S, Oishi H, Takemura H, Amano K (2020) Inter-individual differences in occipital alpha
oscillations correlate with white matter tissue properties of the optic radiation. eNeuro 7.

Mo J, Schroeder CE, Ding M (2011) Attentional modulation of alpha oscillations in macaque
inferotemporal cortex. J Neurosci 31:878–882.

Nestvogel DB, McCormick DA (2022) Visual thalamocortical mechanisms of waking state-
dependent activity and alpha oscillations. Neuron 110:120-138.e4 Available at:
https://www.sciencedirect.com/science/article/pii/S089662732100773X.

Nolte G, Ziehe A, Kramer N, Popescu F, Muller K-R (2008a) Comparison of Granger Causality
and Phase Slope Index. JMLR Workshop Conf Proc 6:267–276.

Nolte G, Ziehe A, Nikulin V V., Schlögl A, Krämer N, Brismar T, Müller KR (2008b) Robustly
estimating the flow direction of information in complex physical systems. Phys Rev Lett
100:1–4.

O'Connor DH, Fukui MM, Pinsk MA, Kastner S (2002) Attention modulates responses in the
human lateral geniculate nucleus. Nat Neurosci 5:1203–1209.

Palva S, Palva JM (2007) New vistas for α-frequency band oscillations. Trends Neurosci
30:150–158 Available at:
https://www.sciencedirect.com/science/article/pii/S0166223607000264.

Rathbun DL, Warland DK, Usrey WM (2010) Spike timing and information transmission at
retinogeniculate synapses. J Neurosci Off J Soc Neurosci 30:13558–13566.

Ray S (2015) Challenges in the quantification and interpretation of spike-LFP relationships. Curr
Opin Neurobiol 31:111–118.

Ray S, Hsiao SS, Crone NE, Franaszczuk PJ, Niebur E (2008) Effect of stimulus intensity on
the spike-local field potential relationship in the secondary somatosensory cortex. J
Neurosci 28:7334–7343.

Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS (2014) Pupil
fluctuations track fast switching of cortical states during quiet wakefulness. Neuron
84:355–362.

Rihs TA, Michel CM, Thut G (2007) Mechanisms of selective inhibition in visual spatial attention
are indexed by alpha-band EEG synchronization. Eur J Neurosci 25:603–610.

Saalmann YB, Pinsk MA, Wang L, Li X, Kastner S (2012) The pulvinar regulates information
transmission between cortical areas based on attention demands. Science (80-) 337:753–
756.

Sherman SM, Guillery RW (2006) Exploring the Thalamus and Its Role in Cortical Function, 2nd
ed. Cambridge, MA: MIT Press.

Sherman SM, Guillery RW (2013) Functional Connections of Cortical Areas: A New View from
the Thalamus. Available at: https://doi.org/10.7551/mitpress/9780262019309.001.0001.

Shomer DL (2017) Niedermeyer's electroencephalography: basic principles, clinical
applications, and related fields (Schomer DL, Lopes da Silva FH, eds). Oxford University
Press. Available at: https://doi.org/10.1093/med/9780190228484.001.0001.

Spaak E, Bonnefond M, Maier A, Leopold DA, Jensen O (2012) Layer-specific entrainment of
gamma-band neural activity by the alpha rhythm in monkey visual cortex. Curr Biol
22:2313–2318.

Steriade M (2006) Grouping of brain rhythms in corticothalamic systems. Neuroscience
137:1087–1106.

Steriade M, Paré D, Parent A, Smith Y (1988) Projections of cholinergic and non-cholinergic
neurons of the brainstem core to relay and associational thalamic nuclei in the cat and
macaque monkey. Neuroscience 25:47–67.

Sweeney-Reed CM, Zaehle T, Voges J, Schmitt FC, Buentjen L, Kopitzki K, Esslinger C,
Hinrichs H, Heinze H-J, Knight RT, Richardson-Klavehn A (2014) Corticothalamic phase
synchrony and cross-frequency coupling predict human memory formation. Elife 3:e05352.

Tuladhar AM, ter Huurne N, Schoffelen J-M, Maris E, Oostenveld R, Jensen O (2007) Parieto-
occipital sources account for the increase in alpha activity with working memory load. Hum
Brain Mapp 28:785–792.

Usrey WM, Alitto HJ (2015) Visual Functions of the Thalamus. Annu Rev Vis Sci 1:351–371.

Usrey WM, Reppas JB, Reid RC (1998) Paired-spike interactions and synaptic efficacy of retinal
inputs to the thalamus. Nature 395:384–387 Available at: https://doi.org/10.1038/26487.

Usrey WM, Sherman SM eds. (2023) The cerebral cortex and thalamus. Oxford University
Press. Available at: https://doi.org/10.1093/med/9780197676158.001.0001.

Vinck M, Batista-Brito R, Knoblich U, Cardin JA (2015) Arousal and locomotion make distinct
contributions to cortical activity patterns and visual encoding. Neuron 86:740–754 Available
at: http://dx.doi.org/10.1016/j.neuron.2015.03.028.

Vinck M, van Wingerden M, Womelsdorf T, Fries P, Pennartz CMA (2010) The pairwise phase
consistency: A bias-free measure of rhythmic neuronal synchronization. Neuroimage
51:112–122 Available at: http://dx.doi.org/10.1016/j.neuroimage.2010.01.073.

Wang W, Jones HE, Andolina IM, Salt TE, Sillito AM (2006) Functional alignment of feedback
effects from visual cortex to thalamus. Nat Neurosci 9:1330–1336.

Wimmer RD, Schmitt LI, Davidson TJ, Nakajima M, Deisseroth K, Halassa MM (2015) Thalamic
control of sensory selection in divided attention. Nature 526:705–709.

Yang X, Fiebelkorn IC, Jensen O, Knight RT, Kastner S (2024) Differential neural mechanisms
underlie cortical gating of visual spatial attention mediated by alpha-band oscillations. Proc
Natl Acad Sci U S A 121:e2313304121.

Zhang R V., Featherstone RE, Melynchenko O, Gifford R, Weger R, Liang Y, Siegel SJ (2019)
High-beta/low-gamma frequency activity reflects top-down predictive coding during a
spatial working memory test. Exp Brain Res 237:1881–1888 Available at:
https://doi.org/10.1007/s00221-019-05558-3.

Zhou H, Schafer RJ, Desimone R (2016) Pulvinar-Cortex interactions in vision and attention.
Neuron 89:209–220 Available at: http://dx.doi.org/10.1016/j.neuron.2015.11.034.

 FIGURE LEGENDS

Figure 1: LGN local field potential. (A1) Snippet of an LGN extracellular recording, lowpass

filtered (<250 Hz) to extract LFP activity (M1 = animal ID). (A2) Power spectrum from the full,

representative recording. Dashed gray line = 1/f fit. (A3) Oscillatory power, defined as the

residual power after subtracting the 1/f fit. (A4) Z-score of the oscillatory power. Power above

the gray shaded box was considered evidence of significant oscillatory activity (B-F) Additional

examples, one from each monkey included in this study.

Figure 2: Oscillatory activity in LGN LFP. (A) Distribution of significant residual oscillatory

activity in the LGN across all 6 animals. (B). Percentage of LGN recordings with significant

oscillatory activity in 5 frequency bands: alpha 7-10 Hz, low-beta 11-23 Hz, high-beta 24-35 Hz,

gamma 40-70 Hz, and high-frequency >70 Hz. (C1) Average LGN power spectrum for monkey

M1 (n = # of recordings). (C2) Average LGN oscillatory activity for monkey M1. (C3).

Percentage of recordings with significant oscillatory activity at each frequency for monkey M1.

(D-H) Same as C, one row for each animal included in this study.

Figure 3: LGN-V1 field-field phase synchrony. (A1) Example of simultaneously recorded LGN

and V1 power spectra (LGN = blue, V1 = orange). (A2) Example field-field ppc for paired

recording shown in A. (B-D) Additional examples, same as A. (E1) Average LGN-V1 field-field

ppc across our sample. (E2) PPC separated into trials with weak or strong beta oscillations

(blue trace and red trace, respectively) for recordings with significant low beta-band LGN-

V1phase synchrony. (E3) Same as E2, but for recordings with significant high beta-band LGN-

V1 phase synchrony. (F1) Average LGN-V1phase slope index (psi) for recording with significant

low beta-band LGN-V1 phase synchrony. Positive and negative values are indicative of

feedforward (LGN→V1) and feedback processes (LGN←V1), respectively. (F2) Same as F1,

but for recordings with significant high beta-band LGN-V1 phase synchrony. (F3) Sample

distribution for frequency band specific psi values (* = p< 0.001, ** = p<10^{-5}).

Figure 4: Residual spike waveforms in LFP. (A1) Example snippet of raw continuous voltage

before either high-pass filtering (to extract action potentials) or low-pass filtering (to extract

LFP). (A2) Spike-triggered LFP waveform (after low-pass filtering) from the example recording

shown in A1. (A3) Pairwise phase consistency for the current example. (B) Additional example,

same as A. (C) Average spectrogram, calculated via matching pursuit, across all LGN

recordings. (D) Broadband, high-frequency spike-field phase synchrony was strongly related to

the underlying signal-to-noise ratio of the LGN spike waveform.

Figure 5: Beta-band spike-field phase synchrony in the LGN. (A1) Example snippets of weak

(top trace, blue) and strong (bottom trace, red) beta band activity in LGN LFP. (A2) Example

power spectra for periods of weak and strong beta band activity (blue and red traces,

respectively). (A3) Example spike-triggered average LFP waveforms from same periods of weak

and strong beta-band activity. (A4) Example PPC curve for the same periods of weak and

strong beta-band activity. (B-C) Additional examples, same as A. (D) Average ppc curves

across all LGN recordings, separated into periods of weak and strong beta-band activity. (E)

Difference PPC curve: strong – weak. (F) Distribution of PPC z-scores for periods of weak and

strong beta band activity. Values outside of the gray shaded region were considered significant.

(G) Scatter plot of beta band PPC values for periods of weak and strong beta-band activity. For

visualization purposes, values were capped at 0.06.

Figure 6: Visual stimulation suppresses beta-band oscillatory strength in the LGN (A1) Example

LGN power spectra for weak and strong visual stimuli, gray trace and black trace respectively.

(A2) LGN oscillatory power for current example. (B-D) additional examples, same as A. (E1)

Sample distribution for LGN beta-band oscillatory power: weak vs strong visual stimulation (pink

= low beta band; yellow = high beta band). (E2) Sample distribution of stimulus strength index

for LGN beta-band oscillatory power (index = (strong – weak)/(strong+weak)). (F1) Average

LGN oscillatory power curves (blue = weak stim, red = strong stim) for LGN recordings with

significant low beta-band oscillatory strength. Frequency is defined relative to the peak low-beta

band LGN oscillatory power (separately for each curve). (F2) Same as F1 for LGN recordings

with significant high beta-band oscillatory strength (F3) Frequency band specific index values (**

= p< 0.01).

Figure 7: Visual stimulation suppresses beta band LGN-V1 phase synchrony. (A1) Example

LGN and V1 power spectra during weak visual stimulation, blue and orange curves respectively.

(A2) Example LGN-V1 field-field PPC curves: weak stimulation = gray trace, strong stimulation

= black trace. (B-D) Additional examples, same as A. (E1) population distribution for LGN-V1

beta-band PPC: weak vs strong visual stimulation (pink dots = high beta-band, yellow dots =

high beta-band). (E2) Population distribution of stimulus strength index for LGN-V1 beta-band

PPC (index = (strong – weak)/(strong-weak)). (F1) Average LGN-V1 PPC (blue = weak stim, red

= strong stim) for LGN recordings with significant low beta-band LGN-V1 phase synchrony.

Frequency is defined relative to the peak low beta-band LGN oscillatory power (separately for

each curve). (F2) Same as F1 for paired recordings with significant high beta-band LGN-V1

phase synchrony (F3) Frequency band specific index values (* = p<0.05, ** = p< 0.01, *** = p<

10^{-5}).

Figure 8: Behavioral arousal suppresses beta-band oscillations in the LGN. (A1) Example LGN

power spectra for low and high arousal trials, gray trace and black trace respectively. (A2) LGN

oscillatory power for current example. (B-D) additional examples, same as A. (E1) Sample

distribution for LGN beta-band oscillatory power: low vs high arousal (pink = low beta band;

yellow = high beta band). (E2) Sample distribution of arousal index for LGN beta-band

oscillatory power (index = (high – low)/(high+low)). (F1) Average LGN oscillatory power curves

(blue = low arousal, red = high arousal) for LGN recordings with significant low beta-band

oscillatory strength. Frequency is defined relative to the peak low beta-band LGN oscillatory

power (separately for each curve). (F2) Same as F1 for LGN recordings with significant high

beta-band oscillatory strength (F3) Frequency band specific index values (*** = p< 10^{-5}).

Figure 9: Behavioral arousal suppresses beta band LGN-V1 phase synchrony. (A1) Example

LGN and V1 power spectra during low behavioral arousal, blue and orange curves respectively.

(A2) Example LGN-V1 field-field PPC curves: low arousal = gray trace, high arousal = black

trace. (B-D) Additional examples, same as A. (E1) population distribution for LGN-V1 beta band

PPC: low vs high behavioral arousal (pink dots = high beta-band, yellow dots = high beta-band).

(E2) Population distribution of arousal index for LGN-V1 beta-band PPC (index = (high –

low)/(high-low)). (F1) Average LGN-V1 PPC (blue = low arousal, red = high arousal) for LGN

recordings with significant low beta-band LGN-V1 phase synchrony. Frequency is defined

relative to the peak low beta=band LGN oscillatory power (separately for each curve). (F2)

Same as F1 for paired recordings with significant high beta-band LGN-V1 phase synchrony (F3)

Frequency band specific index values (* = p<0.05, ** = p< 0.01, *** = p< 10^{-5}).

Figure 10: Shifts in spatial attention modulate beta-band activity in the LGN. (A1) Example LGN

power spectra when spatial attention was directed away from and toward an LGN receptive

field, gray trace and black trace, respectively. (A2) Example LGN oscillatory power. (B-D)

additional examples, same as A. (E1) Sample distribution for LGN beta-band oscillatory power:

away vs toward the LGN receptive field power (pink = low beta band; yellow = high beta band)

(E2) Population distribution of spatial attention index for LGN beta-band oscillatory power (index

= (toward – away)/(toward+high)). (F1) Average LGN oscillatory power curves (blue = away, red

= toward) for LGN recordings with significant low beta-band oscillatory strength. Frequency is

defined relative to the peak low beta-band LGN oscillatory power (separately for each curve).

(F2) Same as F1 for LGN recordings with significant high beta-band oscillatory strength (F3)

Frequency band specific index values (* = p< 0.05).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10