Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Martina Pandinelli[1,2], Mohsen Parto-Dezfouli[1,3], Eleni Psarou[1,2], Iris Grothe[1,4], Pascal Fries[1,2,3]

[1]Ernst Strüngmann Institute (ESI) for Neuroscience in Cooperation with Max Planck Society, 60528 Frankfurt am Main, Germany

[2]Donders Institute for Brain, Cognition and Behavior, Radboud University Nijmegen, 6525, Nijmegen, Netherlands

[3]Max Planck Institute for Biological Cybernetics, 72076, Tübingen, Germany

[4]Brain Research Institute, Cognitive Neurophysiology Group, 28359 Bremen, Germany

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Significance

Natural visual experience often entails repetitions of the same stimulus. This allows optimization of the brain's processing of those stimuli. Indeed, repeated stimuli typically induce decreasing firing rates and increasing gamma-band synchronization in many visual areas. However, in this study in awake macaque primary visual cortex, we report surprising opposite effects: under some conditions, stimulus repetition can result in neuronal firing rate increases and gamma-band synchronization decreases. We argue that these effects might relate to the fact that the repeated stimulus was not predictive of a reward. In fact, when we isolated the responses to a second stimulus that was added later in each trial and that was predictive of reward, the effects reverted again to the typical pattern.

Abstract

Stimulus repetition is abundant, because the environment is redundant and/or because it is redundantly sampled. This offers an opportunity to optimize the processing of repeated stimuli. Indeed, stimulus repetition leads to classically described neuronal response decreases, and to more recently described neuronal gamma synchronization increases (sometimes preceded by decreases for a few trials). Here, we used a full-screen colored background (FSCB) and a flashed black bar, while recording multi-unit activity (MUA) and local field potentials (LFP) from area V1 of an awake macaque monkey. We found that the FSCB repetition induced neuronal response increases (sometimes preceded by decreases for a few trials) and gamma synchronization decreases (preceded by increases for a few trials). These effects are largely opposite to the dominant previous findings. Intriguingly, these surprising effects reversed when we isolated the responses to the flashed black bar. We discuss these findings, considering differences to previous studies with regards to the subject of the study, the stimuli and the task. We notice that in studies reporting classical results for gamma, sometimes in combination with firing rates, the stimuli were typically (partly) predictive of the reward. Here, we found non-classical results for the FSCB that was not reward predictive, and classical results for the black bar that was reward predictive. Whether this has revealed a general effect of reward predictive versus non-predictive stimuli will require further investigation with stimuli and task designs tailored specifically for this question.

Keywords: stimulus repetition; primary visual cortex (V1); multi-unit activity (MUA); gamma-band oscillations; reward prediction

Introduction

Repetition is a key feature of visual experience on several levels. On a behavioral level, the eyes of a subject tend to perform return saccades, as a strategy to optimize foraging (Psarou et al., 2025; Wilming et al., 2013), and thereby revisit the same objects. This brings the same stimuli repeatedly into the receptive fields (RFs) of a given neuron or neuronal group in retinotopically organized visual cortical areas, like primary visual cortex, V1. On a stimulus

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

level, a natural visual scene can contain dynamic stimulus repetitions, when e.g. image
elements move back and forth and thereby in and out of the mentioned RFs. On a combined
stimulus and behavioral level, stimuli can contain spatial repetitions (Torralba & Oliva, 2003),
like e.g. trees in a forest, or faces in a crowd, and the behavioral sampling of these repeated
items can stimulate those RFs repeatedly with similar stimuli. All these types of stimulus
repetition offer the brain the opportunity to optimize its computation (Olshausen & Field,
1996; Schwartz et al., 2007).

Repetition of visual stimuli in experimental contexts has generally led to decreases in
neuronal firing rates (Desimone, 1996; Li et al., 1993) and in the fMRI BOLD (functional
magnetic imaging blood oxygenation level dependent) response (Grill-Spector et al., 2006).
This effect, classically described as repetition suppression, has no detrimental impact on
behavioral performance (Grill-Spector et al., 2006). Maintained performance might be
achieved by neuronal synchronization (Gotts, 2003; Gotts et al., 2012) which might increase
the impact of the synchronized neurons by means of feedforward coincidence detection
(Salinas & Sejnowski, 2001) or entrainment (Fries, 2015). Indeed, increased neuronal
gamma-band synchronization within and between awake macaque areas V1 and V4 has
been reported as a result of stimulus repetition (N. M. Brunet et al., 2014). This repetition-
related gamma increase has been confirmed by studies that additionally demonstrated
stimulus specificity of the repetition effect, both in monkeys (Peter et al., 2021; Psarou et al.,
2025) and humans (Stauch et al., 2021).

In the present study, we report surprising findings obtained serendipitously while analyzing
the effect of stimulus repetition in a dataset that had originally been collected to further
investigate a different previously reported phenomenon: visually induced gamma-band
activity rhythmically modulates the gain of the spike response to a stimulus change (Ni et al.,
2016). In the present experiment, we recorded multi-unit activity (MUA) and local field
potentials (LFPs) from awake macaque area V1, while inducing sustained gamma-band
activity with a full-screen-colored background (FSCB) (Peter et al., 2019; Shirhatti & Ray,
2018). On top of the FSCB and at an unpredictable time, we then presented in the RFs of
the recorded V1 neurons, a black bar that evoked neuronal spike responses.

This paradigm differed in several aspects from previous studies that had described
repetition-related changes in gamma-band activity. These previous studies had used
sustained stimulation with gratings or natural stimuli (N. M. Brunet et al., 2014; Peter et al.,
2021; Psarou et al., 2025; Stauch et al., 2021), whereas here we used the FSCB and a
flashed bar. Furthermore, in those previous studies, the stimuli that induced the investigated
neuronal activity underwent changes or offsets that temporally predicted the reward. In the
present experiment, reward was tightly predicted by the black bar, whereas the FSCB did
not add to reward prediction.

This paradigm produced repetition-related changes in neuronal activity that were surprising,
as they differed drastically from the results of those previous studies. The previous studies
had confirmed numerous reports that stimulus repetition leads to MUA decreases (N. Brunet
et al., 2014; Peter et al., 2021; Psarou et al., 2025). Furthermore, they revealed that stimulus
repetition can lead to initial gamma decreases for a few (typically 4-10) repetitions (Peter et

al., 2021; Stauch et al., 2021), always followed by gamma increases for the remaining
repetitions (N. M. Brunet et al., 2014; Peter et al., 2021; Psarou et al., 2025; Stauch et al.,
2021). By contrast, the present study revealed that the repetition of the FSCB induced MUA
increases, preceded by MUA decreases for some conditions and/or task period. The
repetition of the FSCB consistently induced gamma increases for some initial trials followed
by gamma decreases for the remaining trials. Even more surprisingly, the repetitions of the
black bar induced MUA decreases and initial gamma decreases followed by gamma
increases, largely following results from previous studies.

Results

Stimuli and task are illustrated in Fig. 1A. After an inter-trial interval with a randomly colored
noise mask, each trial started with the presentation of a gray FSCB and a central fixation
point (we refer to gray as a color, simply for ease of use). As soon as the monkey attained
fixation, the FSCB color remained gray in 10% of the trials (gray-condition), or turned red in
90% of the trials (red-condition). The sequence of gray- and red-condition trials was
pseudorandom. After 1.4-1.7 s, while the FSCB remained unchanged, a black bar (width:
0.1 dva (degrees of visual angle), length: 0.3 dva) was flashed for 0.2 s into the previously
mapped RF, while the monkey was required to keep fixation. In different trials, the bar was
presented at different contrast levels, pseudorandomly chosen from 5%, 10%, 20%, 40%,
80%, or 100% in each trial of the red condition, and always at full contrast (100%) in the
gray-condition (see Methods for definition of contrast). Thus, the trials of the red-condition
contained approximately equal numbers of trials of the 6 distinct contrast conditions. If the
monkey kept fixation, the bar was followed by a gray FSCB and reward.

This paradigm offered an interesting opportunity to investigate repetition-related changes in
neuronal activity, because it differed from previous studies as outlined in the introduction.
We studied neuronal activity in response to the repetition of the FSCB and the repetition of
the FSCB with the black bar superimposed. Moreover, we studied the neuronal response to
the repetition of the black bar in isolation, by subtracting the response to the FSCB.

To analyze neuronal activity, we defined three periods indicated by the blue-shaded
rectangles in Fig. 1B, C. The "FSCB-transient" period started with the fixation onset, which
also triggered the red FSCB onset in the red condition; the FSCB-transient period was
200 ms long. The "FSCB-sustained" period corresponded to the last 200 ms end-aligned to
the onset of the black bar. Finally, the "bar-transient" period started with the onset of the
black bar and was 200 ms long. As expected, the LFP (Fig. 1B) and MUA (Fig. 1C)
averaged over all trials and recording sites showed phasic responses during the FSCB-
transient period in the red condition, much smaller phasic responses during this period in the
gray condition (as there was no color change, just fixation onset), and contrast-dependent
phasic responses in the bar-transient period.

For each of the conditions and periods, we first computed the raw LFP power spectra
averaged over recording sites and all trials (Fig. S1). The raw LFP power spectrum is not
very sensitive for showing rhythmical neuronal synchronization (Vinck et al., 2013).

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Nevertheless, these raw power spectra revealed gamma peaks or at least gamma bumps for
all combinations of conditions and task periods. Note that clear gamma peaks were visible
also for the FSCB-transient and FSCB-sustained periods of the gray condition. This was
unexpected, and potentially due to a color-aftereffect, as few gray-condition trials were
interspersed between many red-condition trials. We nevertheless opted to use the gray
condition as a baseline condition for the normalization of the red condition. This is mainly
useful for the investigation of power spectra, which are otherwise dominated by a 1/f
component. Importantly, when we analyzed the raw LFP power spectra, all results remained
qualitatively identical.

For the normalization, we expressed the power during the red condition as percent change
relative to the gray condition (from the corresponding task period). For the bar-transient
period, this was done separately per contrast condition. This was first done per recording
site, and the resulting LFP power change spectra were then averaged over recording sites.
They showed prominent beta and gamma peaks for the FSCB-transient period (Fig. 1D),
and they were dominated by a large gamma peak for both the FSCB-sustained period
(Fig. 1E) and the bar-transient period (Fig. 1F). For the bar-transient period, they differed
across contrast conditions, mainly in the alpha-beta range (Fig. 1F). For further analysis of
the gamma-band power, we defined the gamma band, separately per task period, as the
peak frequency in the average power change spectrum plus/minus 10% of that frequency,
resulting in 55±5.5 Hz for the FSCB-transient period, 48±4.8 Hz for the FSCB-sustained
period, and 47±4.7 Hz for the bar-transient period (as indicated by the thin vertical lines and
gray rectangles in Fig. 1 D-F).

Repetition-related changes in MUA

We investigated repetition-related changes in MUA across trials. We first normalized all MUA
values per recording site and per task period, by expressing MUA as percent of the MUA in
the gray condition, averaged over the respective task period. These normalized MUA values
were subsequently averaged over recording sites and plotted per trial (Fig. 2). We separately
considered all combinations of color conditions (Fig. 2 rows) and task periods (Fig. 2
columns). Note that the bar-transient period during the red condition contained six different
contrast conditions. Therefore, we also compiled a plot of repetition-related MUA changes,
for which we averaged the data first per contrast condition and then over contrast conditions
(Fig. 2D).

For several of the combinations of color condition and task period, MUA showed an initial
repetition-related decrease followed by an increase. Similar breakpoints had been previously
observed by us in several other datasets, both for MUA and gamma in awake macaque V1
(Peter et al., 2021; Psarou et al., 2025) and for gamma in human MEG (Stauch et al., 2021).
In these datasets, the breakpoints had been determined by eye and occurred at trial 4-10 for
gamma and 4-20 for MUA. In the current dataset, visual inspection of the MUA results
suggested that breakpoints could occur around trial 20 when we considered the gray
condition (Fig. 2E-G), and around trial 120 when we considered the red condition (Fig. 2A).
We noticed that the red condition contained 6 different contrast conditions for the bar that

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

was flashed into the RF, whereas the gray condition contained only 1 contrast condition. We
further noticed that the breakpoint appeared around trial number 20 times the number of
contrast conditions. We hypothesize that this might be related to the fact that the bar was
temporally predictive of the reward (see Discussion for details). With this in mind, we opted
for a segmented regression, where we fixed the number of segments to two, and the
breakpoint to occur between trial 5-20 times the number of contrast conditions. This indeed
provided excellent fits. For consistency, we applied this approach also to the combinations of
color condition and task period, for which visual inspection did not reveal a breakpoint
(Fig. 2B-D).

Most importantly, this analysis revealed that after the initial MUA decrease, which occurred
for some conditions and task periods, all conditions and task periods showed repetition-
related MUA increases until the end of the session. This was opposite to many previous
reports, including our own, of repetition-related MUA decreases (N. M. Brunet et al., 2014;
Peter et al., 2021; Psarou et al., 2025; Stauch et al., 2021).

We also investigated those repetition-related MUA changes as a function of time within the
three task periods. For this, we focused on the trials after the breakpoint, and we pooled
over red and gray conditions. After pooling, per task period, we determined the breakpoint
(Fig. S2) and divided the remaining trials into three equal-sized trial bins, the early, middle
and late trials (Fig. 3). Within each trial bin, we averaged the normalized MUA (blue, gray
and red lines in Fig. 3A-C), and we statistically compared the late to the early trials (Fig. 3D-
F). This showed that repetition-related MUA increases occurred throughout all three task
periods.

Repetition-related changes in LFP gamma power

We next investigated repetition-related changes in LFP gamma-band power, using the same
approaches as used for MUA. Gamma showed initial repetition-related increases followed by
decreases, consistently for all combinations of color condition and task period (Fig. 4A-F)
except for the bar-transient period in the gray condition (Fig. 4G). The segmented regression
located the breakpoint to trial number 8-16 for the gray condition with one bar-contrast
condition (Fig. 4E-G), at trial number 7 when we averaged over bar-contrast conditions in the
red condition (Fig. 4D), and at trial number 30-50 for the red condition, which corresponds to
trial number 5-8.3 after dividing by the 6 bar-contrast conditions (see last paragraph for the
rationale of this). These breakpoint positions were consistent with the hypothesis that the
breakpoint occurs at trial number 5-15 times the number of bar-contrast conditions.

Most importantly, the observation of repetition-related gamma decreases preceded by initial
increases is opposite to our previous studies using other stimuli and task paradigms. In
those previous studies, we had consistently found repetition-related gamma increases,
preceded in some cases by initial decreases for 4-10 trials.

We also investigated those repetition-related LFP-power changes as a function of frequency
within the three task periods, following the same approach as for MUA (for trial-pooled
gamma results and breakpoints see Fig. S3). This revealed, for the FSCB-transient period,

repetition-related LFP power increases in the alpha band and decreases in both the beta
and gamma bands (Fig. 5A, D). For the FSCB-sustained and the bar-transient period,
repetition-related changes were dominated by gamma decreases (Fig. 5B, C, E, F). Note
that both the FSCB-transient and the bar-transient period most likely contained event-related
potentials (ERPs), both for the red and the gray conditions. As we used the gray condition
for normalization, the ERP-related spectral perturbations might have influenced the
normalized spectra. Therefore, we also considered an alternative normalization, in which the
different task periods of the red conditions were always normalized by the FSCB-sustained
period of the gray condition. This left the results essentially unchanged (Fig. S4).

Repetition-related changes in MUA-LFP gamma synchronization

We also investigated MUA-LFP synchronization, quantified by the MUA-LFP pairwise phase
consistency metric (Vinck et al., 2010). As the PPC is poorly defined for single trials, we
used the same trial-binning approach as in Figs. 3 and 5.

During the FSCB-transient (Fig. 6A, D) and the bar-transient periods (Fig. 6C, F), MUA-LFP
PPC spectra were dominated by low-frequency synchronization, most likely due to the
transient neuronal responses jointly evoked in MUA and LFP by the FSCB and bar onset,
respectively. Those low-frequency components were the only that showed repetition-related
changes, namely an increase during the FSCB-transient period and a decrease during the
bar-transient period.

During the FSCB-sustained period, the MUA-LFP PPC spectrum showed a low-frequency
and a gamma component. Both showed repetition-related decreases. The MUA-LFP PPC
gamma component and its repetition-related decrease corresponds closely to the gamma
component in the LFP power spectra, suggesting that the LFP power spectra reflect local
neuronal synchronization.

Isolating responses to the bar and their repetition-related changes

As mentioned above, the repetition-related changes observed for the current stimuli and task
were largely opposite to the changes observed previously (N. M. Brunet et al., 2014; Psarou
et al., 2025; Stauch et al., 2021). Those previous studies had used gratings or photos of
objects to induce neuronal activity, and those stimuli underwent changes or offsets that
temporally predicted the reward. In the present study, the stimulus that temporally predicted
reward was the bar. We have already analyzed the neuronal responses to the bar on top of
the FSCB. Yet, this response to the compound FSCB + bar stimulus might be dominated by
the response to the FSCB, and we therefore set out to isolate the response to the bar by
subtracting the response to the FSCB alone. As a good estimate for this isolated FSCB
response, we used the FSCB-sustained period. We subtracted the FSCB-sustained results
from the bar-transient results, and refer to the difference as bar-only results (Fig. 7). We
analyzed this separately for all red-condition trials pooled (Fig. 7A, D), for the red-condition
trials averaged per bar-contrast condition (Fig. 7B, E), and for the gray trials (Fig. 7C, F). To
our great surprise, those bar-only results showed again the same repetition-related changes

that we had found in previous studies, namely for MUA a monotonic decrease (see
Fig. 7A-C for significances), and for gamma an initial decrease followed by an increase for
the rest of the trials (Fig. 7D-F, both highly significant for most conditions).

254 Correlations between repetition-related and stimulus-driven changes

Finally, we tested whether repetition-related changes in neuronal activity were correlated
with stimulus-driven changes in neuronal activity. MUA in V1 can be enhanced by many
stimuli, and the repetition of those stimuli typically causes MUA response reductions. By
contrast, we found, for all conditions except the bar-only condition, repetition-related MUA
response increases. Intriguingly, the MUA response in awake macaque V1 to large uniform
red surfaces had previously been reported to be a reduction below baseline (Peter et al.,
2019). If this stimulus-induced MUA reduction were itself reduced by stimulus repetition, just
like stimulus repetition typically reduces stimulus-induced MUA enhancements, then this
might explain our surprising observation of repetition-related MUA increases. If so, the
repetition-related MUA increases should be negatively correlated with stimulus-induced MUA
changes: the larger the stimulus-induced MUA decrease, the larger the repetition-related
MUA increase. We quantified stimulus-induced MUA changes as the ratio of the red
condition over the gray condition, and refer to this as the stimulus-related response ratio
(SR). We quantified repetition-related MUA changes as the ratio of the last 30 trials over the
first 30 trials, and refer to this as the repetition-related response ratio (RR). RR values were
positively correlated to SR values during all task periods (Fig. 8A-C). This analysis also
revealed that the stimuli in this dataset mostly did not suppress but enhance MUA, leading to
positive SR values. Both of these observations do not support the possibility that the
repetition-related MUA increases were due to repetition-related reductions in stimulus-
induced response reductions.

We repeated the same analysis for gamma-band power. For gamma, RR values were not
correlated to SR values during the FSCB-transient period (Fig. 8D), and they were
negatively correlated during the FSCB-sustained and bar-transient period (Fig. 8E, F).

278 Discussion

279 Summary

We report surprising serendipitous findings from investigating repetition-related changes in
MUA and neuronal gamma-band activity in a new dataset. Such repetition-related changes
have been reported in several previous publications, whose overall results can be
summarized as follows: When a given stimulus is repeatedly presented, (1) MUA responses
decrease monotonically with stimulus repetition, with a slope that is steep for the initial ~20
trials and then flattens, (2) neuronal gamma-band responses increase with stimulus
repetition, and this increase can be preceded by a decrease for the initial 4-10 trials. These
repetition-related effects were largely reversed in the present dataset: (1) MUA responses
decreased for the initial trials in some of the conditions or task periods, but subsequently,

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

MUA responses increased for the remaining trials; (2) neuronal gamma-band responses
consistently increased for the initial trials in both the red and the gray conditions and all task
periods, and subsequently, neuronal gamma-band responses decreased for the remaining
trials for all red-condition task periods and showed the same trend for the gray-condition task
periods. However, when we isolated the bar-only response by analyzing the bar-transient
minus the FSCB-sustained period, this largely reverted the effects back to those found
previously.

Limitations of the study

These surprising findings have been obtained from an experiment that investigated a
different effect, namely gamma-rhythmic gain modulation (Ni et al., 2016), and this latter
purpose determined stimuli and task. This experimental design leaves several possible
explanations for the surprising findings, and it will require future experiments to arbitrate
between them. Yet, the observations reported here and in our previous studies on repetition-
related gamma changes do allow to argue against some and in favor of some other,
intriguing, explanations, and we will spend most of this discussion on this exercise.

Note that also our original report of repetition-related gamma increases was based on a
serendipitous finding in a dataset that had been obtained to study selective visual attention
(N. M. Brunet et al., 2014).

The surprising findings are most likely due to experimental differences compared to previous
studies. The main differences that appear plausible are: (1) Differences between animals, (2)
differences between stimuli, (3) differences between tasks.

An effect of the specific animal?

The animal used in the present study is different from the ones used in (N. M. Brunet et al.,
2014) and (Peter et al., 2021), but the same as used in one previous study . This latter study
with this same animal showed repetition-related changes in MUA and gamma-band
responses that were highly similar to the ones reported in (N. M. Brunet et al., 2014) and
(Peter et al., 2021). Thus, the specific animal used in the present study is likely not the
reason for the surprising findings. Yet, this one previous study with this same animal also
used similar stimuli and tasks as parts of (Peter et al., 2021) and (Stauch et al., 2021),
leaving open the possibility that differences in the present study in either one of those
aspects were relevant.

An effect of the specific stimuli?

An effect of using uniform, colored stimuli?

322 The previous studies reporting repetition-related gamma increases had used gratings (N. M.
323 Brunet et al., 2014; Peter et al., 2021; Stauch et al., 2021), and one of them had additionally
324 used photos of natural objects (Peter et al., 2021). None of them had used uniform, colored
325 surfaces. Thus, this specific choice of stimuli might appear like a plausible candidate to

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

326 explain the surprising findings. However, we had six recording sessions from the same
327 animal, in which we used uniform colored discs in a task that was highly similar to (Psarou et
328 al., 2025). While this was a very limited dataset, the results were fully in line with our
329 previously published reports (Fig. S5) and do not support an explanation of the surprising
330 findings as a result of using these stimuli.

An effect of using full-screen stimuli?

332 The previous studies reporting repetition-related gamma increases had mostly used stimuli
333 that did not cover the full screen. However, the original report demonstrated the effect first
334 and very clearly for full-screen gratings (Fig. 1A-C of (N. M. Brunet et al., 2014)). Thus, the
335 use of full-screen stimuli could clearly be eliminated as explanation for the surprising
336 findings.

An effect of the specific task?

This leaves the specific task as an intriguing candidate explanation. To explore this, we first
review the main aspects of the tasks that have induced repetition-related gamma increases
in previous studies.

In the original report of repetition-related gamma increases (N. M. Brunet et al., 2014),
several different datasets were used. Most of the data were recorded while macaques
performed selective attention tasks, during which two patches of drifting grating were
presented and one was cued as behaviorally relevant. One of the patches was inside the RF
of the recorded neurons, the other one outside, and attention was randomly cued per trial to
the stimulus inside or outside. The cued stimulus had to be monitored for a randomly timed
change in shape (for the ECoG data in that study) or color (for the V4 microelectrode data in
that study), which the animal needed to report by releasing a bar. The behavioral response
prompted the disappearance of both the attended and the non-attended stimulus, and their
disappearance immediately preceded reward delivery. This same study also reported
repetition-related gamma increases for another ECoG dataset. For this dataset, a monkey
monitored the fixation point for a color change, while the neuronal activity was induced by a
full-field patch of drifting grating. Similar to the attention tasks, the behavioral report of the
fixation-point color change prompted the disappearance of the stimulus, which immediately
preceded reward delivery.

In the subsequent study (Peter et al., 2021), macaques monitored stimuli for randomly timed
changes, which were small localized contrast decrements randomly positioned on the
stimuli. Those stimuli were either natural stimuli (color photos of isolated leaves, flowers,
sweets, fruits, vegetables) or grating patches. Change detection was again reported
manually, which prompted stimulus disappearance and subsequent reward delivery.

In the next study (Stauch et al., 2021), human subjects monitored a large grating patch for a
suprathreshold (50%) contrast decrement of the entire grating combined with a near-
threshold orientation change. The contrast change prompted the behavioral report of the
orientation change. The behavioral report in turn prompted stimulus disappearance and

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

subsequent presentation of a smiley, which provided positive feedback irrespective of
behavioral response correctness.

In the most recent study (Psarou et al., 2025), macaques fixated while patches of grating
were presented in the RFs of the recorded neurons. The duration of stimulus presentation
was either 0.8-0.9 s for variable-stimulus blocks, or 1-1.3 s for fixed-stimulus blocks (see ref
for details), and the stimulus offset was immediately followed by reward delivery.

In summary, in the previous datasets for which a repetition-related gamma increase has
been reported (N. M. Brunet et al., 2014; Peter et al., 2021; Stauch et al., 2021), the stimulus
that activated the recorded neurons was behaviorally relevant for obtaining the reward,
and/or stimulus changes or stimulus offsets were predictive of the upcoming reward.

In the present study, as soon as the animal acquired fixation, the FSCB appeared and
stayed on throughout the trial. After 1400 to 1700 ms, randomly timed, the bar was
presented inside the RF of the recorded neurons for 200 ms. After the bar disappeared, the
FSCB turned gray, which was a change for the red-condition trials and no change for the
gray-condition trials, and this was followed by reward delivery after a constant interval of 100
ms. Thereby, the bar was predictive of the upcoming reward. Note that when we isolated the
neuronal responses to the bar by subtracting the responses to the FSCB, we found similar
repetition-related changes as in previous studies. In comparison, the FSCB was less
predictive of the reward, as it was presented for a variable amount of time before reward
delivery, and it also did not contain any temporal structure predictive of the upcoming bar
and/or reward. For the FSCB, the main findings were opposite to the previous studies.
Based on these considerations, we would like to speculate as follows.

We will first focus on the changes that we have observed for many repetitions. In the
previous studies, that was the repetition-related firing-rate decrease and gamma increase;
we will later discuss the changes observed in some datasets (including the present one) for
the initial repetitions. Those many-repetition-related changes might be explained by a
statistical dependence of the reward on the stimuli. When a stimulus repeatedly predicted a
reward, we observed that the stimulus-induced neuronal responses showed firing-rate
decreases and gamma increases. By contrast, and this is the main novel insight from the
present study, when a stimulus was repeatedly presented but not reward predictive, or at
least there was another stimulus that was much more tightly reward-predictive, then the
stimulus-induced neuronal responses showed firing-rate increases and gamma decreases.

We next turn to the changes that we have observed for the initial repetitions in some of the
previous studies. In those previous studies this was mainly a firing-rate decrease that was
steeper than for the later repetitions, and a gamma decrease for the initial 4-10 repetitions
preceding the later gamma increase. This gamma decrease was observed for non-
overtrained natural stimuli in macaques and for non-overtrained gratings in human subjects
(Peter et al., 2021; Stauch et al., 2021), and it was not observed for highly-overtrained
gratings in macaques (N. M. Brunet et al., 2014; Peter et al., 2021; Psarou et al., 2025).
Therefore, we have previously suggested that the initial presentations of non-overtrained
stimuli induce an arousal reaction, which leads to increased gamma, and that this arousal

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

and its enhancing effect on gamma decline over the course of few trials. In the present
study, we find similar initial gamma decreases when we isolate the response to the bar, and
we find opposite changes, i.e., initial gamma increases when we analyze the response to the
FSCB.

In addition, we here make one intriguing observation, which holds both for the responses to
the FSCB and the responses to the black bar: the number of those initial trials was similar to
previous studies, i.e., in the range of 7 to 20 when we analyzed the gray condition, for which
only one bar contrast was used, and when we analyzed the red condition after averaging
over the different bar conditions. However, when we concatenated all trials of the red
condition, the number of those initial trials was larger, in the range of 30 to 120. When this
larger number was divided by 6, i.e., the number of bar-contrast conditions for the red
condition, then we ended up with similar initial-trial numbers as in the other cases. This
suggests that the changes observed for the initial trials "count" the number of repetitions of a
particular black-bar. This might be related to the abovementioned, speculative, mechanism
that results from the pairing of a particular stimulus with reward. Yet note, that this putative
explanation departs from our previous interpretation of the initial-trial related changes as an
arousal-related effect. The arousal in response to the FSCB is not expected to be modulated
by the pairing of the preceding black bar with reward.

We emphasize that these latter paragraphs offer speculations intended to explain all
observations as parsimoniously as possible, namely as the learning of a stimulus' reward
predictability. Yet, those observations have partly been made serendipitously – this applies
to the original study (N. M. Brunet et al., 2014) and the present one – and therefore the
stimuli and tasks were not optimized to investigate the observed phenomena, to minimize
confounds, and to dissect different possibilities of explanation. At the same time, the
observed effects are so striking and appear so fundamental, and the datasets from invasive
recordings in awake non-human primates are so precious, that we think those serendipitous
findings deserve to be reported. That said, the speculations offered as putative
interpretations definitively require rigorous testing in other datasets and experiments that
allow the dissection of different possibilities, and we anticipate that this will be a highly fruitful
avenue for future research.

Acknowledgments

This work was supported by DFG (FR2557/1-1, FR2557/2-1, FR2557/5-1, FR2557/7-1), EU (HEALTH-F2-2008-200728-BrainSynch, FP7-604102-HBP), NIH (1U54MH091657-WU-Minn-Consortium-HCP). We thank Dr. Alina Peter for the most useful scientific input, and Rasmus Roese for the invaluable day to day assistance and supervision in the laboratory work.

Author contributions

Conceptualization: M.P., E.P., P.F.; investigation: M.P., E.P., I.G.; methodology: M.P., E.P., I.G., P.F.; software: M.P., M.P.D., E.P.; formal analysis: M.P., M.P.D., E.P., P.F.; writing – original draft; M.P., P.F. writing – review & editing, all authors; visualization: M.P., M.P.D., P.F.; supervision: P.F.; funding acquisition, P.F.

Declaration of interests

P.F. has a patent on thin-film electrodes and is a member of the Advisory Board of CorTec GmbH (Freiburg, Germany).

Supplemental information

Document S1. Figures S1☐S5.

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Figure legends

Fig. 1. Stimuli, paradigm, task periods, and stimulus evoked and induced MUA and LFP responses. (*A*) Schematic illustration of stimuli and task periods for the gray condition (upper row) and red condition (lower row). (*B*) LFP averaged over trials and recording sites, separately for the gray and the red conditions, shown in the corresponding colors. (*C*) Same as *B*, but for MUA. (*D*) LFP power spectra during the FSCB-transient period for the red condition, averaged over recording sites and trials. Power is expressed as percentage change relative to the corresponding task period of the gray condition. (*E*) Same as *D*, but for FSCB-sustained period. (*F*) Same as D, but for bar-transient period. For the bar-transient period (B, C, F), the different bar-contrast conditions are color coded according to the legend to the right of panel *C*.

Fig. 2. Trial-by-trial analysis of repetition-related changes in MUA responses. (*A*) MUA responses for the FSCB-transient period of the red condition, expressed as percent change to the corresponding gray condition, averaged over recording sites and sessions, separately per trial. Repetition-related changes were fitted with a segmented linear regression, whose results are shown as yellow regression lines, separated by a vertical line at the break point. The text above the plot provides the trial number of the break point (BP), and the slope and p-value for the first regression segment (S1, p1) and the second segment (S2, p2). The respective data points were colored magenta and blue. (*B*) Same as *A*, but for the FSCB-sustained period. (C) Same as *A*, but for the probe-transient period. (*D*) Same as *A*, but for the probe-transient period after averaging separately per bar-contrast condition. (*E*) Same as *A*, but for the FSCB-transient period of the gray condition. (*F*) Same as *A*, but for the FSCB-sustained period of the gray condition. (*G*) Same as *A*, but for the bar-transient period of the gray condition.

Fig. 3. Time-course analysis of repetition-related changes in MUA responses. (*A*) MUA responses, expressed as percent change relative to the average FSCB-sustained period of the gray condition, for the FSCB-transient period of both the red and gray condition combined, separately for the early, middle and late trial group as indicated by the color legend right of panel *C*. The trial groups were defined after excluding the trials before the break point identified by the segmented linear regression. (*B*) Same as *A*, but for the FSCB-sustained period. (*C*) Same as *A*, but for the bar-transient period. (*D*) The black line shows the difference of the late minus the early trial group in *A*. The gray-shaded region indicates the range of differences expected by chance. All observed differences were significant. (*E*) Same as *D*, but corresponding to panel *B* for the FSCB-sustained period. (*F*) Same as *D*, but corresponding to panel *C* for the bar-transient period.

Fig. 4. Trial-by-trial analysis of repetition-related changes in LFP gamma-band responses. Same as Fig. 2, but for LFP gamma-band power instead of MUA responses.

Fig. 5. Spectral analysis of repetition-related changes in LFP power. Same as Fig. 3, but for LFP power changes during the different task periods, expressed as percent change relative to the corresponding task periods of the gray condition.

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Fig. 6. Spectral analysis of repetition-related changes in MUA-LFP PPC. Same as Fig. 3, but
for MUA-LFP PPC spectra, which are intrinsically normalized.

Fig. 7. Isolation of MUA (upper row) and LFP gamma power (lower row) responses during
the bar-transient period after subtracting the responses to the FSCB-sustained period. (*A*)
Same as Fig. 2C, but after subtracting the data from the FSCB-sustained period. (*B*) Same
as Fig. 2D, but after subtracting the data from the FSCB-sustained period. (*C*) Same as
Fig. 2G, but after subtracting the data from the FSCB-sustained period. (*D*) Same as
Fig. 4C, but after subtracting the data from the FSCB-sustained period. (*E*) Same as Fig. 4D,
but after subtracting the data from the FSCB-sustained period. (*F*) Same as Fig. 4G, but
after subtracting the data from the FSCB-sustained period.

Fig. 8. Analysis of correlation between repetition-related and stimulus related modulation. In
all plots, each dot represents one recording site, the x-axis value represents the site's
stimulus-related modulation (stimulation-over-baseline ratio), and the y-axis value represents
the site's repetition-related modulation (late-over-early ratio). The upper row shows the MUA
data, the lower row shows the LFP gamma power data. The three columns show the results
separately for the three task periods, as indicated on the top. The insets in each plot provide
the respective Pearson correlation coefficients and the corresponding p-values.

Fig. S1. Raw LFP power spectra, without normalization. Each plot shows the raw LFP power
spectrum, averaged over trials and recording sites, separately for the red and gray condition
as indicated by the inset legends. The different plots show the data for the different task
periods. For the bar-transient period, the data were either averaged over all contrast
conditions (*C*), or separately per contrast condition (*D*), as indicated by the color legend to
the right.

Fig. S2. Same as Fig. 2, but for red- and gray-condition trials combined. As this combines
trials of different conditions, those trials do not constitute repetitions of the same stimulus,
and correspondingly, the x-axis is not labeled "Repetition number" but "Trial number".

Fig. S3. Same as Fig. 4, but for red- and gray-condition trials combined. As this combines
trials of different conditions, those trials do not constitute repetitions of the same stimulus,
and correspondingly, the x-axis is not labeled "Repetition number" but "Trial number".

Fig. S4. Same as Fig. 5, but always using the FSCB-sustained period for normalization.

Fig. S6. Classical results obtained in the same animal for different stimuli and task. (*A*)
Illustration of task paradigm and stimuli. (*B*) Trial-by-trial analysis of repetition-related
changes in MUA responses to the red-color disk stimuli. MUA responses for the stimulation
period, averaged over recording sites and sessions, separately per trial. The yellow line
shows the linear regression, whose slope and p-value are given as inset. (*C*) Same as *B*, but
for grating patches. (*D*) Same as *B*, but for LFP gamma power. (*E*) Same as *C*, but for LFP
gamma power.

*Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-
band synchronization in awake macaque V1*

Materials and Methods

Surgical procedures

All experiments on NHP were performed in compliance with the German and European ethical guidelines and regulations for the protection of animals. The experimental procedures were reviewed and approved by the German regional authority, the Regierungspräsidium Darmstadt.

One adult male macaque monkey (*Macaca Mulatta*), 19 years of age, weighing 13 kg, was used in this study. We used one rather than the more typical two or three animals, because we recently showed that the inference with the addition of a second or third animal is still limited to the sample (Fries & Maris, 2022; Laurens, 2022; Psarou et al., 2024). Thus, doubling or tripling the number of used animals would not change the quality of the inference, and ethical considerations mandate the use of one animal. All surgical procedures performed on the animal were carried out under general anesthesia, and analgesic and antibiotic treatments were administered in the postoperative time period.

The procedure for head-post implantation followed a two-step approach, described in detail in (Psarou et al., 2023). Subsequently, a modular custom-made titanium recording chamber for acute recordings was implanted. The chamber was fixed over the occipital area of the skull corresponding to visual area V1, on the left hemisphere. AAV virus injections into V1, used for other experiments, were performed at locations remote from the recording locations used here.

Visual Stimulation

Visual stimuli were presented on an LG 32GK850G-B monitor (LG Electronics Inc., Seoul, South Korea), and gamma correction was applied in all tasks (gamma: 1.57). The monkey's eyes were at a distance of 83.5 cm from the screen. The precise times of black-bar presentation were recorded by simultaneously presenting a small stimulus in the corner of the monitor, occluded from the monkey's view, and recording that small stimulus with a custom-made photo-diode signal.

Receptive field (RF) estimation

The RF position and orientation preference were determined after averaging neuronal responses over all electrode contacts on the laminar probe that were deemed to be located inside the cortex, as is explained in the corresponding section below. The electrode contacts were densely spaced, such that RF positions and orientation preferences were expected to be similar across simultaneously recorded contacts.

RF position was determined at the beginning of each recording session by using the back-projection method described in (Fiorani et al., 2014). Per trial, one bar moving in one of eight possible directions across the entire monitor was presented while the monkey kept fixation

on a central fixation spot. The moving bar was 0.1 deg wide and had a speed of 8 deg/s. Ten
correct trials for each direction were collected, for a total of 80 correct trials.

The MUA spike density functions were accumulated over time to reconstruct a spatial
response map that indicated the position of the receptive field, after correcting for the
response latency which maximized the back-projection response. The MUA's response-
based receptive field position was inferred by smoothing MUA responses with a Gaussian
function for each direction of movement, and taking the 10th and 90th percentile and fitting
an ellipse to the resulting data points.

Orientation tuning task

The orientation preference of the located RF was inferred with a dedicated orientation tuning
task. Per trial, a 0.1 degrees of visual angle (dva) wide and 0.3 dva long bar of one of six
possible orientations, was presented at the location of the RF, while the monkey was
required to keep fixation on the central fixation spot. A trial was considered correctly
performed if fixation was maintained. Ten correct trials for each orientation were collected for
a total of 60 trials. The preferred orientation was inferred by directly comparing the MUA (or
LFP in case of low MUA signal) response amplitude across the 6 orientations and choosing
the orientation with the highest related MUA response. The screen background during the
orientation-tuning task was gray (RGB 128, 128, 128).

Behavioral task

During training and recordings, the monkey sat in a primate chair located inside a fully
shielded recording booth, with dim illumination. All recordings were performed under head
fixation. The behavioral paradigm was created using ARCADE (Dowdall et al., 2018), a
custom software built in house using MATLAB.

The monkey performed a fixation task. The animal acquired fixation on a central black
fixation dot, 0.3 dva diameter, positioned in the center of the gray screen, and maintained
fixation in a tolerance radius of 0.5 dva from the fixation spot. A trial was considered
correctly performed when the monkey kept fixation for the whole duration of a trial (1650 to
1950 ms), and was rewarded with the delivery of grape juice.

Each trial consisted of the presentation of a sequence of visual stimuli on the monitor. Each
trial began by showing a gray background screen (RGB coordinates [128, 128, 128])
together with the central fixation dot. The monkey was allowed a time period of maximally 10
seconds to acquire fixation, yet they typically attained fixation much quicker.

As soon as the animal started fixating, a full-screen colored background (FSCB) was shown.
The FSCB was red colored (RGB coordinates [255,0,0]) in 90% of the trials (red-condition),
while in 10% of the trials, the screen maintained the same gray color as the initial
background (gray-condition). The gray trials were taken as a baseline reference for the red
trials. The FSCB time period had a constant duration of 1450 ms, plus a time jitter ranging
from 0 to 250 ms to precede the following section of the trial.

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

After the FSCB time period, the FSCB, red or gray, was kept in place, and an additional
visual stimulus in the shape of a black bar was presented inside the previously determined
RF for 200 ms. The black bar had a length of 0.3 dva, a thickness of 0.1 dva and was
oriented according to the preference of the recorded neurons. In the red condition, the black
bar stimulus was presented at 6 contrast levels (see below), while in the gray condition, the
black bar stimulus was always shown at full contrast. After bar offset, a waiting period of
100 ms elapsed after which the animal was rewarded. Fixation breaks led to immediate
termination of the trial, and no reward was given.

During each inter-trial interval (ITI, 700 ms), a colorful noise mask was presented on the
screen. The noise mask was composed of squares of height and width ranging from 0.1 to
0.5 dva in steps of 0.01 dva, and each square had a color randomly selected from the
possible RGB color coordinates. This mask was intended to counteract color adaptation
effects.

Color contrast and normalized color space

618 In the red condition, the black bar stimulus was presented at six contrast levels: 5%, 10%,
619 20%, 40%, 80%, 100%. The contrast was calculated based on the Weber contrast formula:

$$\frac{I - Ib}{Ib}$$

620 , where I indicates the luminance of the stimulus, and I_b indicates the luminance of the
621 background. In order to define contrast levels appropriately, color coordinates for the red
622 background and for the black bar foreground stimulus were chosen from the CIELAB
623 standardized color space (Gravesen, 2015), then translated to RGB coordinates.

Color patches and oriented gratings task

Results shown in Fig. S5 were obtained by analyzing data collected with a dedicated task.
Every trial of the task started with an inter-trial interval (ITI) of 500 ms, presenting a grey
screen (RGB [128,128,128]). An acquire-fixation time window started right after the ITI. A
central fixation dot of 0.5 dva, dark gray in color (RGB [204,204,204]), was presented on the
screen. In this time window, the monkey was allowed a maximum of 10000 ms to acquire
fixation. After fixation was acquired, the monkey was required to maintain fixation for the
duration of a baseline time window of 1000 – 1300 ms, during which the screen kept the
initial grey color. At the end of the baseline window, the visual stimulus was presented during
a 1300 – 1800 ms long time window. The stimulus consisted in a circular patch of uniform
color or of grating. The circular-colored patch had a 7 dva diameter, covering RFs, and could
assume one of the following colors: RGB [149,112,198], [188,107,180], [191,115,129],
[160,130,60], [103,143,6], [44,147,59], [37,140,128], [90,126,179]. The grating stimulus was
also circular, had a 3.5 dva diameter, a spatial period of 0.5 dva, and was approximately
centered on the RF. The grating could assume one of four possible orientations: 0, 45, 90 or
135 degrees. If the monkey maintained fixation for the whole trial duration, the trial was
considered correct, and the monkey was rewarded with delivery of grape juice.

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Eye position

Eye position was recorded at 1000 Hz using an Eyelink 1000 system (Eyelink Inc., SR research Ottawa, Canada). The system was calibrated on the left eye at the beginning of every recording session, and the left-eye signal was recorded.

Recordings

In each recording session, one laminar probe was introduced through the intact dura into the cortex. Laminar probes featured 16 channels in one column, spaced by 50 micron (linear V-probes from Plexon Inc., Dallas, Texas, USA). The probe was guided by a guide tube that rested on the dura. For the recordings, the guide tube served as ground and the shank of the probe as reference. Recordings were performed in a fully shielded booth acting as a Faraday cage that prevented disturbance of the electrophysiological signal by line noise. Correspondingly, line-noise removal was not necessary and not performed.

Electrophysiological data were acquired using Tucker Davis Technologies (TDT, Alachua, Florida, USA) systems. Raw data were recorded and digitized by a TDT PZ2 pre-amplifier at 24.4140625 kHz.

Data Analysis

All analyses were carried out in MATLAB and using the FieldTrip toolbox (Oostenveld et al., 2011).

Preprocessing

Raw data were downsampled to 1000 Hz. For the LFP, this entailed upsampling to 3,125,000 Hz, downsampling to 25,000 Hz, filtering with a stop-band at 500 Hz (IIR filter) and downsampling to 1000 Hz. For the MUA, this entailed upsampling to 3,125,000 Hz, band-pass filtering at 300-12000 Hz, rectification, stop-band filtering at 500 Hz (IIR filter) and downsampling to 1000 Hz.

Recording site selection

During each recording session, we performed one penetration with one laminar probe. The probe was positioned under online visual inspection of the neuronal signals and their general responsiveness to visual stimulation, such that typically all recording sites provided good MUA and LFP responses. Yet, some recording sites did not provide clear RFs, and those recording sites were identified by visual inspection and not used for further analyses. The remaining recordings sites will be referred to as "selected recording sites", or "selected sites".

Spectral analysis

For the spectral analysis, the LFP in the 200 ms long FSCB-transient, FSCB-sustained or bar-transient periods was multiplied with a Hann taper, zero-padded to 1s, and Fourier transformed using the FieldTrip function ft_freqanalysis.

Baseline normalization

All analyses except Fig. 1B, C and Fig. S4 use the gray condition as baseline, separately per
task period. For the MUA, separately per recording site, each task period provided one value
for the gray condition, averaged over time points and trials. This gray-condition value was
used to express the corresponding red-condition MUA as percent change relative to the gray
condition. The same was done for the LFP, except that the LFP was first transformed to a
spectrum, and the averaging and normalization was done per frequency, providing a
frequency-wise LFP power change spectrum during the red condition relative to the gray
condition.

For Fig. S4, the same procedure was followed, except that the normalization was not by the
corresponding task period of the gray condition, but always by the FSCB-sustained period of
the gray condition.

For Fig. 1B, C the signals, both the raw LFP and the MUA, were expressed as percent
change relative to the average respective values during the gray-condition FSCB-sustained
period. This was done per recording site, with subsequent averaging over sites.

Estimation of gamma peak frequency and gamma band

The LFP power change spectra as compiled for the main analyses were used to determine
the gamma peak frequency. The gamma peak frequency was estimated after averaging over
all recording sites from all sessions, yet separately for the FSCB-transient, FSCB-sustained
and bar-transient. The frequency associated with the maximum power change value in the
30-100 Hz range was defined as the gamma peak frequency, and a band of +/- 10% around
the peak was defined as the gamma band.

Analysis of repetition-related changes across trials

To visualize and quantify repetition-related changes across trials, LFP gamma power and
MUA were averaged over recording sites and sessions, separately per trial. For this, only the
trials present in all sessions were used. In other words, the session with the smallest number
of trials for the respective stimulation condition determined the number of trials that was
used from all other sessions. When all trials of both the red and the gray condition were
considered together, this resulted in 813 trials (Fig. S2 and S3). When all trials of the red
condition were considered, this resulted in 750 trials (Fig. 2A, B, C, Fig. 4A, B, C, Fig. 7A,
D). When all trials of the red condition were considered, yet averaged separately per bar-
contrast condition, this resulted in 120 trials (Fig. 2D, Fig. 4D, Fig.7B, E). When all trials of
the gray condition were considered, this resulted in 63 trials (Fig. 2E, F, G, Fig. 4E, F, G, Fig.
7C, F).

Visual inspection of the resulting trial-by-trial MUA and gamma values suggested that they
could generally be described as a linear increase, or a linear decrease, or a combination of
those two with a relatively well-defined break point in between. Furthermore, as discussed in
the results and discussion sections, visual inspection suggested that the break point
occurred at approximately trial number five to 20, times the number of bar-contrast
conditions that was relevant for the respective analysis. Finally, we intended to treat all data

*Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-
band synchronization in awake macaque V1*

as uniformly as possible. Therefore, we fitted the data with two linear regression lines,
separated at a break point that was constrained to lie at trial five to 20, times the number of
bar-contrast conditions that was relevant for the respective analysis. The number of bar
contrast conditions that was relevant for the respective analysis was seven for the all-trials
condition (Fig. S2 and S3), six for the red condition (Fig. 2A, B, C; Fig. 4A, B, C; Fig. 7A, D),
one for the gray condition (Fig. 2E, F, G; Fig. 4E, F, G; Fig. 7C, F), and also one for the red
condition after averaging over bar-contrast conditions (Fig. 2D; Fig. 4D; Fig. 7B, E). The
decision about the precise trial constituting the break point was made objectively by using
the approach described in (Bogartz, 1968). This approach is one version of a segmented
linear regression. The algorithm iteratively considers each trial as a potential break point,
and subsequently selects that trial as break point that minimizes the sum of the squared
errors of the two regression lines.

PPC analysis

Local neuronal synchronization was assessed by calculating the Pairwise Phase
Consistency (PPC) (Vinck et al., 2010) between MUA and the LFP (Fig. 6), as implemented
in the FieldTrip toolbox (Oostenveld et al., 2011). For each session, respectively for each
laminar-probe penetration, we calculated PPC between all combinations of MUA and LFP
from all selected recording sites, except between MUA and LFP from the same recording
site. This latter exclusion was made to minimize spectral bleeding of spikes into the LFP. We
used the "unipolar" LFP, i.e., the LFP referenced against the laminar-probe shaft.
Subsequently, the MUA-LFP PPC was averaged over site-pair combinations. PPC was
based on Fourier transforms of the MUA and the LFP during the task period and trial groups
as described in the results section.

Statistical analysis

Statistical testing was performed (1) for the segmented linear regressions quantifying the
trial-by-trial MUA or gamma responses (Fig. 2, 4, 7, S2, S3), (2) for the simple linear
regression quantifying the trial-by-trial MUA or gamma responses in the very few trials with
color or grating patches (Fig. S5), (3) for the comparison of MUA change time courses, LFP
change spectra, or MUA-LFP PPC spectra between late and early trial groups, including
correction for multiple comparisons across time points or frequencies (Fig. 3, 5, 6, S4).

Regarding the statistical tests for case (1): The segmented linear regression was performed
on the observed trial-by-trial data, giving two observed regression slopes. Subsequently, the
trial numbers were randomized, and the segmented linear regression repeated, and this
randomization was performed 1000 times. The slopes obtained across randomizations were
collected in two randomization distributions, one for each segment. Per segment, we first
determined whether the observed slope was positive or negative and then determined the
number of randomization slopes with the same sign that exceeded the observed slope. This
number was divided by the total number of randomizations and multiplied by two, and the
resulting value was taken as p-value of a two-sided test.

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Regarding the statistical tests for case (2): This followed the same approach as case (1),
except that the regression was not segmented, but performed across all of the few available
trials.

Regarding the statistical tests for case (3): We first obtained the observed difference time
course or observed difference spectra of the late minus the early trial group. Subsequently,
we randomly exchanged trials between the late and the early trial groups and recalculated
the differences, and this was performed 1000 times. Importantly, for each randomization,
only the minimal and the maximal difference – across all time points or frequencies - was
retained and placed into the minimal randomization distribution and the maximal
randomization distribution, respectively. The 2.5th percentile of the minimal randomization
distribution and the 97.5th percentile of the maximal randomization distribution were taken as
thresholds for statistical significance at a false-positive level of 5%, including correction for
the multiple comparisons across time points or frequencies, respectively (Nichols & Holmes,
2002).

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

References

772 Bogartz, R. S. (1968). A least squares method for fitting intercepting line segments to a set
773   of data points. *Psychological Bulletin, 70*(6), 749-755.
774   https://doi.org/10.1037/h0026713

775
776 Brunet, N., Vinck, M., Bosman, C. A., Singer, W., & Fries, P. (2014, Oct). Gamma or no
777   gamma, that is the question. *Trends Cogn Sci, 18*(10), 507-509.
778   https://doi.org/10.1016/j.tics.2014.08.006

779
780 Brunet, N. M., Bosman, C. A., Vinck, M., Roberts, M., Oostenveld, R., Desimone, R., De
781   Weerd, P., & Fries, P. (2014, Mar 4). Stimulus repetition modulates gamma-band
782   synchronization in primate visual cortex [Research Support, Non-U.S. Gov't]. *Proc
783   Natl Acad Sci U S A, 111*(9), 3626-3631. https://doi.org/10.1073/pnas.1309714111

784
785 Desimone, R. (1996, Nov 26). Neural mechanisms for visual memory and their role in
786   attention [Review]. *Proc Natl Acad Sci U S A, 93*(24), 13494-13499.
787   http://www.ncbi.nlm.nih.gov/pubmed/8942962

788
789 Dowdall, J., Schmiedt, J., Stephan, M., & Fries, P. (2018). Arcade: a modular multithreaded
790   stimulus presentation software for the Real-Time control of stimuli, actions and
791   reward during behavioral experiments. *SfN, 2018*(254.18), 2018.

792
793 Fiorani, M., Azzi, J. C., Soares, J. G., & Gattass, R. (2014, Jan 15). Automatic mapping of
794   visual cortex receptive fields: a fast and precise algorithm. *J Neurosci Methods, 221*,
795   112-126. https://doi.org/10.1016/j.jneumeth.2013.09.012

796
797 Fries, P. (2015, Oct 7). Rhythms for Cognition: Communication through Coherence [Review].
798   *Neuron, 88*(1), 220-235. https://doi.org/10.1016/j.neuron.2015.09.034

799
800 Fries, P., & Maris, E. (2022, Jun 2). What to Do If N Is Two? *J Cogn Neurosci, 34*(7), 1114-
801   1118. https://doi.org/10.1162/jocn_a_01857

802
803 Gotts, S. J. (2003). *Mechanisms underlying enhanced processing efficiency in neural
804   systems* Carnegie Mellon University]. Pittsburgh, PA.

805
806 Gotts, S. J., Chow, C. C., & Martin, A. (2012). Repetition priming and repetition suppression:
807   A case for enhanced efficiency through neural synchronization. *Cogn Neurosci, 3*(3-
808   4), 227-237. https://doi.org/10.1080/17588928.2012.670617

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

810 Gravesen, J. (2015, 2015/11/01/). The metric of colour space. *Graphical Models, 82*, 77-86.
811   https://doi.org/https://doi.org/10.1016/j.gmod.2015.06.005

812
813 Grill-Spector, K., Henson, R., & Martin, A. (2006, Jan). Repetition and the brain: neural
814   models of stimulus-specific effects. *Trends Cogn Sci, 10*(1), 14-23.
815   https://doi.org/10.1016/j.tics.2005.11.006

816
817 Laurens, J. (2022). The statistical power of three monkeys. *bioRxiv*,
818   2022.2005.2010.491373. https://doi.org/10.1101/2022.05.10.491373

819
820 Li, L., Miller, E. K., & Desimone, R. (1993, Jun). The representation of stimulus familiarity in
821   anterior inferior temporal cortex [Research Support, Non-U.S. Gov't]. *J Neurophysiol,*
822   *69*(6), 1918-1929. http://www.ncbi.nlm.nih.gov/pubmed/8350131

823
824 Ni, J., Wunderle, T., Lewis, C. M., Desimone, R., Diester, I., & Fries, P. (2016, Oct 5).
825   Gamma-Rhythmic Gain Modulation. *Neuron, 92*(1), 240-251.
826   https://doi.org/10.1016/j.neuron.2016.09.003

827
828 Nichols, T. E., & Holmes, A. P. (2002, Jan). Nonparametric permutation tests for functional
829   neuroimaging: a primer with examples [Research Support, Non-U.S. Gov't]. *Hum*
830   *Brain Mapp, 15*(1), 1-25. http://www.ncbi.nlm.nih.gov/pubmed/11747097

831
832 Olshausen, B. A., & Field, D. J. (1996, Jun 13). Emergence of simple-cell receptive field
833   properties by learning a sparse code for natural images [Research Support, U.S.
834   Gov't, P.H.S.]. *Nature, 381*(6583), 607-609. https://doi.org/10.1038/381607a0

835
836 Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source
837   software for advanced analysis of MEG, EEG, and invasive electrophysiological data
838   [Research Support, Non-U.S. Gov't]. *Comput Intell Neurosci, 2011*, 156869.
839   https://doi.org/10.1155/2011/156869

840
841 Peter, A., Stauch, B. J., Shapcott, K. A., Kouroupaki, K., Schmiedt, J. T., Klein, L., Klon-
842   Lipok, J., Dowdall, J. R., Schölvinck, M. L., Vinck, M., Schmid, M. C., & Fries, P.
843   (2021). Stimulus-specific plasticity of macaque V1 spike rates and gamma. *Cell*
844   *Reports, 37*(10), 110086. https://doi.org/doi.org/10.1016/j.celrep.2021.110086

845
846 Peter, A., Uran, C., Klon-Lipok, J., Roese, R., van Stijn, S., Barnes, W., Dowdall, J. R.,
847   Singer, W., Fries, P., & Vinck, M. (2019, Feb 4). Surface color and predictability
848   determine contextual modulation of V1 firing and gamma oscillations. *Elife, 8*.
849   https://doi.org/10.7554/eLife.42101

850
851 Psarou, E., Katsanevaki, C., Maris, E., & Fries, P. (2024). Would you agree if N is three? On
852   statistical inference for small N. *bioRxiv*, 2024.2008. 2026.609821.

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Psarou, E., Parto-Dezfouli, M., Grothe, I., Peter, A., Roese, R., & Fries, P. (2025).
Repetition-related gamma plasticity in macaque V1 and V2 is highly stimulus specific
and robust to stimulus set size. *bioRxiv*, 2025.2006. 2020.660699.

Psarou, E., Vezoli, J., Schölvinck, M. L., Ferracci, P. A., Zhang, Y., Grothe, I., Roese, R., &
Fries, P. (2023, Jun 1). Modular, cement-free, customized headpost and connector-
chamber implants for macaques. *J Neurosci Methods, 393*, 109899.
https://doi.org/10.1016/j.jneumeth.2023.109899

Salinas, E., & Sejnowski, T. J. (2001, Aug). Correlated neuronal activity and the flow of
neural information [Research Support, U.S. Gov't, Non-P.H.S.

Review]. *Nat Rev Neurosci, 2*(8), 539-550. https://doi.org/10.1038/35086012

Schwartz, O., Hsu, A., & Dayan, P. (2007, Jul). Space and time in visual context. *Nat Rev
Neurosci, 8*(7), 522-535. https://doi.org/10.1038/nrn2155

Shirhatti, V., & Ray, S. (2018, Apr 24). Long-wavelength (reddish) hues induce unusually
large gamma oscillations in the primate primary visual cortex [Research Support,
Non-U.S. Gov't]. *Proc Natl Acad Sci U S A, 115*(17), 4489-4494.
https://doi.org/10.1073/pnas.1717334115

Stauch, B. J., Peter, A., Schuler, H., & Fries, P. (2021, Aug 24). Stimulus-specific plasticity in
human visual gamma-band activity and functional connectivity. *Elife, 10*, e68240.
https://doi.org/10.7554/eLife.68240

Torralba, A., & Oliva, A. (2003, Aug). Statistics of natural image categories. *Network, 14*(3),
391-412. https://www.ncbi.nlm.nih.gov/pubmed/12938764

Vinck, M., van Wingerden, M., Womelsdorf, T., Fries, P., & Pennartz, C. M. (2010, May 15).
The pairwise phase consistency: A bias-free measure of rhythmic neuronal
synchronization. *Neuroimage, 51*(1), 112-122.
https://doi.org/10.1016/j.neuroimage.2010.01.073

Vinck, M., Womelsdorf, T., Buffalo, E. A., Desimone, R., & Fries, P. (2013, Nov 20).
Attentional modulation of cell-class-specific gamma-band synchronization in awake
monkey area V4. *Neuron, 80*(4), 1077-1089.
https://doi.org/10.1016/j.neuron.2013.08.019

Wilming, N., Harst, S., Schmidt, N., & König, P. (2013). Saccadic momentum and facilitation
of return saccades contribute to an optimal foraging strategy. *PLoS Comput Biol,
9*(1), e1002871. https://doi.org/10.1371/journal.pcbi.1002871

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Pandinelli et al., Surprising effects of stimulus repetition on neuronal firing rates and gamma-band synchronization in awake macaque V1

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5